Single-cell genomics of uncultured bacteria reveals dietary fiber

responders in the mouse gut microbiota

Rieka Chijiiwa[1,2#], Masahito Hosokawa[3,4,5,#,*], Masato Kogawa[1,2], Yohei Nishikawa[1,2], Keigo

Ide[1,2], Chikako Sakanashi[3], Kai Takahashi[1], Haruko Takeyama[1,2,3,4*].

[1]Department of Life Science and Medical Bioscience, Graduate School of Advanced Science

and Engineering, Waseda University, 2-2 Wakamatsucho, Shinjuku-ku, Tokyo 162-8480,

Japan.

[2] Computational Bio Big-Data Open Innovation Laboratory (CBBD-OIL), National Institute

of Advanced Industrial Science and Technology, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555,

Japan.

[3]Research Organization for Nano and Life Innovation, Waseda University, 513

Wasedatsurumaki-cho, Shinjuku-ku, Tokyo 162-0041, Japan.

[4] Institute for Advanced Research of Biosystem Dynamics, Waseda Research Institute for

Science and Engineering, Graduate School of Advanced Science and Engineering,

Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan.

[5]PRESTO, Japan Science and Technology Agency (JST), 5-3 Yonban-cho, Chiyoda-ku,

Tokyo, 102-0075, Japan

# These authors equally contributed to this work.

* Correspondence should be addressed to M.H. (m.hosokawa@aoni.waseda.jp) and H.T.

(haruko-takeyama@waseda.jp)

Abstract

Background

The gut microbiota can have dramatic effects on host metabolism; however, current genomic

strategies for uncultured bacteria have several limitations that hinder their ability to identify

responders to metabolic changes in the microbiota. In this study, we describe a novel single-

cell genomic sequencing technique that can identify metabolic responders at the species level

without the need for reference genomes, and apply this method to identify bacterial

responders to an inulin-based diet in the mouse gut microbiota.

Results

Inulin feeding changed the mouse fecal microbiome composition to increase *Bacteroides* spp.,

resulting in the production of abundant succinate in the mouse intestine. Using our massively

parallel single-cell genome sequencing technique, named SAG-gel platform, we obtained 346

single-amplified genomes (SAGs) from mouse gut microbes before and after dietary inulin

supplementation. After quality control, the SAGs were classified as 267 bacteria, spanning

two phyla, four classes, seven orders, and 14 families, and 31 different strains of SAGs were

graded as high- and medium-quality draft genomes. From these, we have successfully

obtained the genomes of the dominant inulin-responders, *Bacteroides* spp., and identified

their polysaccharide utilization loci and their specific metabolic pathways for succinate

production.

Conclusions

Our single-cell genomics approach generated a massive amount of SAGs, enabling a

functional analysis of uncultured bacteria in the intestinal microbiome. This enabled us to

estimate metabolic lineages involved in the bacterial fermentation of dietary fiber and

metabolic outcomes such as short-chain fatty acid production in the intestinal environment

based on the fibers ingested. The technique allows the in-depth isolation and characterization

of uncultured bacteria with specific functions in the microbiota and could be exploited to

improve human and animal health.

Keywords

Single-cell genomics, gut microbiome, uncultured bacteria, mouse, inulin, *Bacteroides*.

Background

The gut microbiota plays a crucial role in the control of host physiology and

metabolism through the release and transformation of metabolites like short-chain fatty acids

(SCFAs) [1, 2]. Many studies have demonstrated that dietary fiber supplementation can

modulate gut microbiota composition and promote SCFA production [3-5]. This dietary fiber

metabolism is particularly important for non-digestible inulin-type fructans classified as

prebiotics, which are substrates that promote the growth of beneficial microorganisms in the

gut [6-12]. Previous reports indicated that several bacterial taxa in the intestine, including

Firmicutes, *Bacteroides*, and *Bifidobacterium*, utilize fructan, and that dietary fructan can

result in the growth of *Actinobacteria*, *Firmicutes*, or *Bacteroides*. However, our

understanding of nutrient sensing and utilization by gut microbiota is limited, and there is a

lack of predictability of the response of the microbiota to such dietary interventions. Thus, to

enhance the benefits of dietary fiber to the host, it is important to identify inulin-responders

in the complex microbiome community of the host intestine and characterize their

mechanisms of inulin fermentation and potential for SCFA production.

Recently developed metagenomic approaches can provide comprehensive microbial

profiling and putative functions using 16S rRNA gene amplicon sequencing and shotgun

metagenome sequencing. These methods have enabled the exploration of complex

communities in the gut environment without the need for *in vitro* cultivation of the intestinal

microbiome. However, 16S rRNA gene sequencing provides snapshots of microbial

composition, but does not provide information on microbial metabolic functions [13].

Shotgun metagenomics provides the complete genomic content of the microbial community,

but it is generally difficult to link taxonomic and functional information for individual

microbial strains [14]. Importantly, these techniques rely on the availability of reference

microbial genomes for accurate functional assignments of specific organisms and precise

taxonomic classification in the microbiome [15-17]. However, an estimated half of human

gut species lack a reference genome. Thus, a new culture-independent genomic approach is

required to enhance the reference genomes along with conventional metagenomic approaches

and unveil the taxonomy and functions of the yet uncultured microbes at the strain level.

In this study, we used single-cell genome sequencing to identify microbial responders

of dietary fiber, inulin, in mouse gut microbiota. Our massively parallel single bacterial

genome sequencing technique, named single-cell amplified genomes in gel beads sequencing

(SAG-gel), utilizes a gel matrix that enables a serial enzymatic reaction for tiny genome

amplification and provides a support for the isolation of the amplified genomes. We targeted

the family *Bacteroidaceae* from massively produced single-cell amplified genomes and

characterized their polysaccharide utilization loci (PUL) and SCFA production pathways [7,

8, 11].

Near-complete genomes obtained via single-cell genome sequencing provided us new

insights into the mechanisms of microbial proliferation and SCFA production in the intestine.

The resulting genome set can serve as the basis for strain-specific comparative genomics to

understand uncultured organisms in the gut microbiome. These findings will help us to

predict the metabolic fermentation of dietary fibers based on the presence and ability of the

specific responders.

Results

Effects of dietary fiber inulin on gut microbiota composition

First, we assessed changes in mouse gut microbiota composition before and after

inulin supplementation. To account for diurnal oscillations in gut microbiota composition [18,

19], feces were sampled in the morning and evening after two weeks of continuous inulin

feeding. Cellulose, which is generally poorly fermented by gut microbes, was used as a

control diet fiber. As shown in Figure 1a, the mouse fecal microbiota mainly consisted of

three classes, *Bacteroidales*, *Lactobacillales*, and *Clostridiales*, regardless of feeding

conditions and sampling time. *Lachnospiraceae* and *Streptococcaceae* were dominant (25-

30% and 17-45%, respectively) in the mice feces collected before fiber feeding and from the

cellulose-fed mice, while in the feces from the inulin-fed mice, the abundance of

Bacteroidaceae increased relative to other bacteria (17% in morning and 41% in evening),

while that of *Lachnospiraceae* decreased (20% in morning and 12% in evening). These

relative abundance profiles indicate a microbial composition increment of *Bacteroidaceae* in

the feces of inulin-fed mice beyond typical diurnal oscillations in comparison to that in the

cellulose-fed mice (p-values <0.005, Tukey's HSD test).

To identify the specific bacterial taxa associated with inulin fermentation, we

compared the fecal microbiota of mice before fiber feeding and inulin-fed mice using the

LDA effect size (LEfSe) analysis [20]. Differentially abundant taxa in the fecal microbiota

and their predominant bacteria are shown in Figures 1b and 1c. These results primarily

indicated the diurnal differences, but revealed that *Bacteroidales* were abundant (LDA score

>2, p-value <0.05) in inulin-fed mice at both time points.

To confirm this effect, we monitored time-dependent microbiota composition changes

in inulin-fed mice from another animal lot. The feces of inulin-fed mice were sampled in the

morning for two weeks. As shown in Figure 1d, *Bacteroidaceae* rapidly increased on day 1,

though there were individual differences in microbial composition. On days 3 to 5,

Bacteroidaceae were most abundant in all mice, which were housed in three different cages.

After this rapid increase, *Bacteroidaceae* gradually decreased but maintained their higher

percentage in all mice except those in cage 1. In this lot of mice, we observed different

microbiota compositional changes compared to the previous experiment, such as an increase

in *Bifidobacteriaceae* in all cages, and in *Akkermansiaceae* in cage 1 during days 6 to 14.

Despite differences in microbial composition between cages and animal lots, the results

strongly suggest that *Bacteroidaceae* is a pivotal responder in the intestinal

microenvironment of inulin-fed mice. In addition, we found two main *Bacteroidaceae spp.*

(OTU2 and OTU4) in all mice, regardless of the fiber feeding condition. As shown in

Additional file 2: Figure S1, OTU2 were rapidly increased after starting inulin feeding and

showed significant compositional increment after 2-weeks of continuous inulin feeding (p-

values < 4e-02, Tukey's HSD test). In contrast, OTU4 showed a constant composition during

inulin feeding, while they showed significant increment compared to that in the cellulose-fed

mice in the evening.

We also evaluated cecum SCFAs to observe the effects of inulin on gut microbiome

metabolic function. Figures 1e and 1f show the concentrations of butyrate and succinate in

the cecum contents of mice used for 16S rRNA gene sequencing, measured by gas

chromatography (GC). Although the total SCFA concentration depended on the diet and

sampling time points in mice before fiber feeding and in the cellulose-fed mice (data not

shown), samples from inulin-fed mice consistently contained higher butyrate and succinate

concentrations.

Taken together, microbial composition and metabolome analyses suggest that inulin

feeding changes the microbiome composition to increase *Bacteroides* spp. and microbial

metabolic functions, resulting in increased succinate and butyrate concentrations in the

mouse intestine.

Single-cell resolution genome analysis of the fecal microbiome of inulin-fed mice

Using the single-cell sequencing based on the SAG-gel platform (Figure 2a), mouse

intestinal microbes were captured in agarose gel beads by microfluidic droplet generator. Cell

lysis with an enzyme cocktail, genomic DNA purification [21], and subsequent whole-

genome amplification [22, 23] were all performed in the agarose gel matrix. After sorting the

single beads, which contain single-cell amplified DNA, into standard PCR plates using a cell

sorter, the DNA samples were re-amplified as a single-cell amplified genome (SAG) library.

From 464 single-cell captured beads derived from mice fecal microbiota samples before (180

SAGs) and after inulin or cellulose feeding (166 SAGs), a total of 346 SAGs, named

IMSAG_001 to 346, were sequenced from the SAG library (Figure 2b). According to the

standards developed by the Genomic Standards Consortium [24], we recovered one high-

quality draft genome, 43 medium-quality draft genomes, 258 low-quality draft genomes, 38

short genome fragments (< 100 kb or 0% completeness), and six contaminated genomes

(Figure 2b and Additional file 1: Table S1). The average completeness and contamination of

the draft genomes were 31.8% and 0.6%, respectively, with an average of 296 contigs (Figure

2c). Notably, only 15 of the 346 (4.3%) SAGs had >5% contamination. Approximately 15%

of SAGs were classified as high- or medium-quality draft genomes, ranging from 1.3-3.6 Mb

in total contig length. The high- and medium-quality draft genomes had an average N50 of

16.4 kb, while the low-quality draft genomes had an average of 10.4 kb (Figure 2d). An

average of 17 tRNAs were detected in the high- and medium-quality draft genomes (Figure

2e). Of the 302 SAG draft genomes, 267 (88%) recovered the 16S rRNA gene, 71% were

>1,400 bp in size (Figure 2f), and 25% contained an rRNA operon that encoded the 16S, 23S,

and 5S rRNA genes in single contigs (Additional file 1: Table S1). When we conducted

nucleotide BLAST searches against a database of 16S rRNA gene sequences, 284 SAGs were

similar to intestinal microbe 16S rRNA gene sequences with >80% identity, whereas 216

SAGs showed <97% identity with the top search results, suggesting that many SAGs were

obtained from uncultured bacteria that have not yet been registered in the database

(Additional file 1: Table S2).

9   Qualified SAGs were classified as 267 bacteria, spanning two phyla, four classes,

seven orders, and 14 families with GTDB-Tk [25]. Of all samples, 123 (46%) were classified

as *Lachnospiraceae*, followed by *Bacteroidaceae* (65 (24%)), and *Muribaculaceae* (27

(10%); Additional file 1: Table S2). The phylogenetic distribution of these SAGs was slightly

different from that of 16S rDNA genes acquired from a metagenomic sample, and the SAGs

covered 62 of 298 operational taxonomic units (OTUs; 21%; Additional file 1: Table S3).

The SAGs covered seven major bacterial families in the metagenomic analysis (>1% of the

total OTUs in the inulin and cellulose samples) but missed the genomes of some gram-

positive bacteria, such as *Streptococcaceae*, *Enterococcaceae*, and *Staphylococcaceae*.

Moreover, some SAGs had strain marker gene identities ≥99%, and average nucleotide

identities (ANIs) ≥95% were co-assembled to improve their completeness as single strain

genomes, which were named IMSAGC_001 to 028 (Additional file 1: Table S4). Finally, we

obtained 31 different SAG strains (22 composite and 9 raw) that were graded as high- (7) or

medium-quality (24) draft genomes among seven families (Table 1 and Additional file 1:

Table S5). These included two *Bacteroidaceae* spp. (IMSAGC_001 and IMSAGC_004) that

had identical 16S rRNA genes to *Bacteroidaceae* (OTU2 and OTU4, respectively), which

were potential inulin-responders in inulin-fed mice feces (Additional file 1: Tables S3 and S5,

and Additional file 2: Figure S1).

Genomic features of *Bacteroides* spp. draft genomes

Bacteroides spp. are capable of metabolizing fructans, a class of plant and microbial

polysaccharides [7, 9, 11]. The completeness of the inulin responder *Bacteroides* draft

genomes IMSAGC_001 and IMSAGC_004 were sufficiently high (≥97%) for comparison

with publicly available genomes of *Bacteroides* spp. known to be fructan utilizers or showing

similarity to these genomes (Additional file 1: Table S6 and Figure 3a). Both had >4.2 Mb

genome sizes, 43% GC content, and >3,500 coding sequences (CDSs), similar to the

genomes of *Bacteroides* spp. including *B. thetaiotaomicron* (Bt), *B. ovatus* (Bo), *B. caccae*

(Bc), *B. vulgatus* (Bv), *B. uniformis* (Bu), *B. fragilis* (Bf), and *B. acidifaciens* (Ba; Figure 3a

and Table S6). Genes in these *Bacteroides* groups were clustered into 5,013 different

orthogroups. The orthogroup distributions between different *Bacteroides* group members are

shown in Figure 3b; of these orthogroups, 1,150 were defined as core orthogroups.

IMSAGC_001 and IMSAGC_004 shared a number of different orthogroups with specific

strains, such as Ba (196 orthogroups) and Bv (99 orthogroups), consistent with the

phylogenetic analysis (Figure 3a). Although the vast majority of the orthogroups (151 and 57

in IMSAGC_001 and IMSAGC_004, respectively) were classified as hypothetical proteins of

unknown function, most orthogroups shared between IMSAGC_001 and Ba belonged to

category O (carbohydrate transport and metabolism; nine orthogroups), followed by

categories T and P (signal transduction mechanisms and inorganic ion transport and

metabolism, respectively; six orthogroups each). In addition, IMSAGC_001 and IMSAGC

004 shared 47 unique orthogroups (37 with unknown functions).

Identification of putative polysaccharide utilization loci from *Bacteroides* spp. draft genomes

Most of the glycan degrading and import machinery within *Bacteroides* genomes are encoded within clusters of coregulated genes known as polysaccharide utilization loci (PUL). The defining characteristic of a PUL is the presence of a pair of genes homologous to *susD* and *susC*, which encode outer membrane proteins that bind inulin and import its digestion products [8, 11, 26]. The *susC* and *susD* homologs are usually associated with genes encoding the machinery necessary to convert extracellular polysaccharides into intracellular monosaccharides, such as glycoside hydrolases (GHs) or polysaccharide lyases (PLs). In addition, most PULs contain or are closely linked to genes encoding an inner membrane-associated sensor-regulator system, including the hybrid two-component system. Notably, some *Bacteroides* species, such as Bc, Bo, Bt, and Bu, can utilize inulin with similar efficiency to glucose [7, 8, 11], while Bv lacks *susC*- and *susD*-like genes within its fructan PUL and cannot use inulin. Bt and Bo encode a number of predicted signal peptidase (SP)II-containing GHs/PLs and act as public goods producers in various networks of polysaccharide utilizers [7]. These producer-derived GHs are extracellularly transported to create polysaccharide degradation products that can be utilized for growth of concomitant recipients, such as Bv, at considerable distance from the producer.

We detected PULs containing inulinases in two *Bacteroides* draft genomes (IMSAGC_001 and IMSAGC_004; Figures 4a and 4b). We also identified other *Bacteroidales*, such as IMSAGC_014 and IMSAGC_006, that had no PULs encoding inulinases. At the whole genome level, IMSAGC_001 showed similarity to inulin utilizers such as Bo, Bc, and Bt, while IMSAGC_004 was similar to the inulin non-utilizer Bv (Figure 3a). In IMSAGC_001, the sequences of GH genes, including inulinase, showed high similarity to those of Bc and Bo, which have superior ability to use inulin [7, 11] and had

highest similarity to Ba (Figure 4c). IMSAGC_004 also contained a GH gene similar to that of known inulin utilizers. The GHs of IMSAGC_001 and IMSAGC_004 are predicted to be periplasmic, as they contain SPI cleavage sites. In addition, IMSAGC_001, IMSAGC_004, and Ba have no predicted SPII-containing GHs and PLs in their inulin utilization loci (Figure 4b). Since *Bacteroides* spp. show differences in numbers of GH/PL-encoding genes and uniformity among strains of a given species [7], we compared the number of SPII-containing GH/PLs among *Bacteroides* genomes. IMSAGC_001 had a medium level of SPII-containing GH/PLs (56, Additional file 1: Table S7), while IMSAGC_004 and Ba had extremely low levels of SPII-containing GH/PLs (26 and 32, respectively). This result indicated that IMSAGC_001 and IMSAGC_004 might contain different types of machinery for polysaccharide utilization from Bt and Bo, which have several SPII-containing GH/PLs, which act as producers for a larger repertoire of polysaccharides.

For *susC/D* homologs, IMSAGC_001 and IMSAGC_004 clustered with Ba and apart from other inulin utilizers (Bc, Bo, Bu, and Bt; Figure 4d). Homologs of the SusC and SusD outer membrane proteins are the defining features of the Sus-system, and genes encoding these proteins are found as a pair in most *Bacteroidetes* genomes. Inulin chains that may or may not be extracellularly cleaved are threaded through the SusC porin, and all degradation occurs in the periplasm. In Bo, deletion of the *susC* and *susD* orthologs results in significant growth impairment on inulin, and the orthologs are predicted to function in inulin binding and import without prior extracellular digestion [9, 26]. As our data suggested that IMSAGC_001 and IMSAGC_004 had low similarity to the Bo *susC/D* homologs, we next compared genome sequence similarities through structure prediction-based analysis. Based on structure prediction, the susC/D homologs in IMSAGC_001 and IMSAGC_004 could form complex structures composed of homodimers of an extracellular SusD-like lipoprotein and an integral membrane SusC-like transporter (Additional file 2: Figure S2). These

predicted structures showed high similarity to the Ba complex at the amino acid level (Additional file 2: Figure S2).

In summary, *Bacteroides* IMSAGC_001, which were significantly increased in inulin-fed mouse feces, and IMSAGC_04, which showed no apparent changes, possess putative PULs that include *susC/D* homolog pairs, a putative outer membrane lipoprotein, a putative inner membrane monosaccharide importer, a putative fructokinase, and a putative inulin-specific GH. No other SAGs had higher similarity to *Bacteroides* PUL. We predict that these bacteria import inulin directly via SusC and SusD for degradation to monosaccharides in the periplasm without prior extracellular degradation.

Identification of metabolic characters in inulin-utilizing strains and other bacterial SAGs

Having identified key responder bacterial strains for inulin metabolism, we next examined their fundamental metabolic functions in the mouse gut microbiome. Additional file 1: Tables S8 and S9 summarize the metabolic functions of all high- and medium-quality genomes obtained in this study (IMSAGC series with >80% completeness). Notably, inulin-fed mice displayed significant increases in *Bacteroides* in their microbial composition and succinate and butyrate concentrations in their SCFAs. For succinate, we focused on the reductive tricarboxylic acid (TCA) cycle, as it produces the major end product of energy metabolism in anaerobic microorganisms. Many bacteria have incomplete TCA cycles and are commensal obligate anaerobes, including *Bacteroides* spp., which use a branched TCA cycle to support fumarate respiration [27, 28]. The end product of fumarate reduction, succinate, is secreted into the extracellular environment. In IMSAGC_001 and IMSAGC_004, we detected reductive and incomplete TCA cycles from oxaloacetate to succinate (Additional

file 2: Figure S3 and Additional file 1: Table S8), while no other predominant SAG contained

both an inulin PUL and a succinate pathway.

In single-cell genome sequencing, the profiling of SAG metabolic pathways provides

various insights into the traits of diverse microbiota at once, because each metabolic gene is

specifically assigned to specific bacterial strains (Additional file 1: Table S9). As an

interesting functional signature, we detected the presence of antibiotic resistance modules,

such as vancomycin resistance, D-Ala-D-Ser type, in specific *Lachnospiraceae* strains

(IMSAGC_003 and IMSAGC_007). We observed differences in conserved biosynthetic

pathways in cofactor and vitamin metabolism, such as biotin and cobalamin synthesis, among

different intestinal microbes. Comparing IMSAGC_001, IMSAGC_004, and other

Bacteroides spp., IMSAGC_004 lacks proline and tryptophan biosynthesis pathways, as well

as the first carbon oxidation pathway in the TCA cycle, but has a conserved cobalamin

biosynthetic pathway similar to those of Bf, Bu, and Bv. Both IMSAGC_001 and

IMSAGC_004 lack glycine betaine/proline transport systems, similar to Ba and Bu. The

inulin nonresponsive *Bacteroidales* spp. (IMSAGC_014 and IMSAGC_006) lacked some

carbohydrate metabolism pathways and saccharide transport systems compared to known

polysaccharide utilizers. These results suggest that inulin-responsive *Bacteroides*

IMSAGC_001, IMSAGC_004, and inulin non-responder *Bacteroidales* spp. play different

metabolic roles in the mouse gut microbiota.

Discussion

Inulin (β2-1 fructan) is resistant to host-mediated digestion, resulting in remarkable

fecal microbiome changes and cecum SCFA concentrations through intestinal microbial

fermentation. In this study, to elucidate the inulin-responders in mouse intestine microbiota,

we conducted 16S rRNA gene sequencing for the assessment of changes in the inulin-

dependent microbiota composition, metabolome analysis for the evaluation of SCFA production, and single-cell genome sequencing for obtaining the draft genomes of inulin-responders. The microbiome composition and SCFA amounts displayed typical diurnal oscillations, and inulin intake resulted in significant changes beyond these. *Bacteroides* spp. are known to utilize levan and inulin (Figures 1 and 3, Additional file 1: Table S6). Microbiome composition monitoring demonstrated rapid responses by *Bacteroides* IMSAGC_001 to inulin supplementation and dynamic daily changes. However, reference *Bacteroides* strains have different origins and uncultured strains present in individual hosts could have different metabolic features. *Bacteroides* members have differing abilities to use these various plant polysaccharides. Species that utilize a particular polysaccharide (producers), such as Bo, liberate polysaccharide degradation products that are consumed by other recipients unable to grow using the intact polysaccharide [7, 8].

Functional characterization of bacteria using 16S rRNA gene sequencing databases may lead to misinterpretations. From a partial 16S rRNA gene sequence, responder bacteria can only be identified at the family level, *Bacteroidaceae*, and insights on the inulin utilization abilities of specific bacteria in the diverse microbial community cannot be obtained (Figure 1 and Additional file 1: Table S3). Therefore, single-cell sequencing was used to analyze closely related species and compare uncultured bacterial genomes to identify inulin-responders in complex microbial communities.

The advantage of single-cell sequencing with the SAG-gel platform is that it can select specific sequencing samples after generating large numbers of SAGs. Since we can adjust the total sequencing effort according to the number of SAGs required, it is not necessary to produce a large number of reads to cover target bacteria with the same depth as in shotgun metagenomic sequencing. Since each SAG is composed of tens of thousands of reads on average, limiting the analysis to specific strains enables extremely cost-efficient

whole-genome sequencing of target cells. Indeed, approximately half of the SAGs were

obtained from up to 96-plex indexes in single sequence runs by MiSeq (total ~3Gb). This

prescreening can utilize marker genes, such as 16S rRNA genes, to select *Bacteroidaceae* and

other abundant strains based on initial 16S rRNA gene sequencing. Even if the target bacteria

were selected from the results of initial 16S rRNA gene sequencing, it is challenging to

demonstrate direct agreement between the MAGs and the target bacteria as most MAGs lack

the rRNA and tRNA sequences in the contigs [15-17]. Thus, SAGs with functional and

phylogenetic information are useful in combination with stepwise multilayer analyses,

including screening through 16S rRNA gene sequencing, to achieve the strategic genomic

analysis of target bacteria.

Conversely, nearly 90% of the SAGs obtained in this study contained 16S rRNA gene

sequences, 70% of which were full-length, and many existed in operons with the 5S and 23S

rRNA genes. The SAGs provide the genetic information of dominant microbes by covering

dominant OTUs existing in >1% of the fecal microbiome community. However, there is still

room for improvement. For example, modifying the cell lysis procedure could enhance the

variety of SAGs detected, including gram-positive bacteria. In 16S rRNA gene homology

analysis, a considerable number of SAGs showed low similarity to reference genomes,

suggesting that these SAGs contain the genomes of species currently undescribed in public

databases. Furthermore, by acquiring duplicate SAG data from identical strains, we could

assemble the draft genomes using a cleaning and co-assembly strategy [22, 23]. For example,

IMSAGC_001 and IMSAGC_004 were duplicated 41 and 12 times, respectively, allowing

the construction of composite draft genomes, which are very advantageous for obtaining

genomes of medium or high quality. These two main Bacteroides *spp.* were present in all

mice, regardless of inulin feeding (Table S3). Thus, we utilized raw SAGs for assembly

improvement from all mice regardless of fiber feeding conditions. According to the definition

used in Almeida et al. [16], 40% of the draft genomes of quality higher than medium were

qualified as near-complete (>90% completeness and < 5% contamination, Additional file

1:Table S5). Hence, the SAG-gel platform can provide increased numbers of draft genomes

from single microbial cells, with higher completeness than other single-cell genome

sequencing techniques [22, 23, 29-32]. The ratio of high- and medium-quality draft genomes

was comparable with recent metagenomic assembly and binning strategies for obtaining

MAGs as draft genomes [15-17].

Pan-genome analysis within the same environment reveals what bacteria are present

and which harbor specific genes (Additional file 1:Table S9). Our results revealed PUL and

succinate synthesis pathways in inulin responder *Bacteroidaceae*. In addition, comparative

SAG analysis allowed us to understand the conservation of various pathways, such as those

involved in antibiotic resistance and vitamin synthesis, in each bacterial strain in the

intestinal microbiome. Single strain-resolution fingerprinting of the microbiome enables the

identification of bacteria or genes responsible for specific metabolic function from a complex

microbial community. In this study, two responder *Bacteroides* genomes had inulin-utilizing

PUL clusters that showed similarity to known inulin utilizers. Our genome analyses

suggested that *Bacteroides* IMSAGC_001 and IMSAGC_004, which can take up and degrade

inulin, grow predominantly in the intestines of inulin-fed mice, and produce succinate

through the glycolytic system and a reductive partial TCA cycle for release into the intestinal

environment. We have identified the potential inulin utilizers *Bacteroides* IMSAGC_001 and

IMSAGC_004 in uncultured fecal microbiomes as a new subgroup with Ba. Ba was

originally isolated from mouse cecum as two strains that differed in their starch hydrolysis

functions [33]. Their major end products of glucose fermentation are acetate and succinate,

similar to those predicted for *Bacteroides* IMSAGC_001 and IMSAGC_004. Our data

suggest that *Bacteroides* IMSAGC_001 and IMSAGC_004 utilize inulin for their metabolism

and lack the ability to liberate the fructose for other species in the same manner as Bf and Ba

[7]. The significant increase in *Bacteroides* IMSAGC_001 over IMSAGC_004 (Additional

file: Figure S1) may be due to differences in the cell growth rate, as well as inulin uptake and

degradation rates, given the differences in GHs and other PUL genes.

Conclusion

In this study, single-cell genomics enabled detailed microbiome analysis and

identified inulin-responders with putative roles in dietary fiber metabolism. The technique

generates a massive preparation of SAGs, enabling the functional analysis of uncultured

bacteria in the intestinal microbiome and the consideration of host responses. The analysis

makes it possible to estimate metabolic lineages in bacterial PUL and metabolic outcomes in

the intestinal environment, such as SCFA production, based on ingested fibers. Genetic and

functional differences between intestinal bacterial species are predictive of *in vivo*

competitiveness in the presence of dietary fibers. Taxonomic and functional genes regulating

bacterial metabolic capacity can serve as potential biomarkers in microbiome datasets and

can be sequenced to assess microbiota control via diet. In the future, it may be possible to

estimate responders in advance based on the presence of these microorganisms and genes.

Methods

Mice

Wild-type six-week-old male mice from the BALB/c background (Tokyo Laboratory

Animals Science Co., Japan) were used. All mice were housed in a 12:12 light:darkness cycle

(Zeitgeber time (ZT) 0 = 8 AM, ZT 12 = 8 PM). Before feeding experiments, the mice were

fed AIN93M (Oriental Yeast Co., Japan) for one week. After that, the mice were fed inulin-

or cellulose-supplemented diets for two weeks in the morning or evening. To prepare the

fiber-supplemented feed, AIN93M powder was mixed with 5% inulin or 5% cellulose, and

then pellets were formed with tap water. Fecal samples were collected for 16S rRNA gene

sequencing and single-cell sequencing, and cecum samples were harvested before and after

the two-week fiber feeding for metabolomic analysis. All samples were stored at -80 °C.

DNA extraction from fecal samples

Fecal samples (200 mg) were homogenized with 20 mL phosphate-buffered saline

(PBS) with a Pellet Pestle™ Cordless Motor (Fisher Scientific). The fecal suspensions were

then filtered through 100-µm nylon mesh to remove eukaryotic cells and other debris, and the

flow-through suspensions were collected as the bacterial cell fraction. Filtered fecal

suspensions were then centrifuged at $9,000 \times g$ for 10 min at 4 °C, and the bacterial pellets

were resuspended in 800 µL Tris-ethylenediaminetetraacetic acid (TE) 10 buffer (10 mM

Tris-HCl/10 mM EDTA). Bacterial pellets were serially incubated with 100 µL lysozyme

(150 mg/mL) at 37 °C for 1 h, 20 µL achromopeptidase (100 U/µL) at 37 °C for 30 min, and

50 µL proteinase K (20 mg/mL) with 50 µL 20% sodium dodecyl sulfate (SDS) at 55 °C for 1

h. The lysates were then treated with equal volumes of phenol:chloroform:isoamyl alcohol

and centrifuged at $6,000 \times g$ for 10 min. After centrifugation, DNA in the upper aqueous

layer was transferred to another tube and then pelleted by adding 3M sodium acetate solution

and isopropanol, followed by centrifugation at $6,000 \times g$ for 10 min. DNA pellets were rinsed

with 70% ethanol, dried, and dissolved in TE buffer (10 mM Tris-HCl/1 mM EDTA). The

collected DNA samples were purified with 10 µg/mL RNase and incubated at 37 °C for 30

min and precipitated by adding equal volumes of 20% polyethylene glycol solution

(PEG6000). After centrifugation at $10,000 \times g$ for 10 min at 4 °C. DNA pellets were rinsed

with 70% ethanol, dried, and dissolved in TE buffer. The extracted DNA samples were stored

at -20 °C.

16S rRNA gene sequencing and analysis

The V3–V4 hypervariable regions of 16S rRNA genes were analyzed according to the

Illumina protocol for 16S Metagenomic Sequencing Library Preparation. Barcoded

amplicons, amplified with 341F and 806R primers (Forward, 5′-

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG-3′;

reverse, 5′-

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATC

C-3′) were sequenced using the Illumina MiSeq 2×300 bp platform with MiSeq Reagent Kit

v3 (Illumina Co.) according to the manufacturer's instructions. Raw reads were processed

using QIIME2 v.2019.1 [34] using the DADA2 [35] plugin to denoise quality filter reads, call

amplicon sequence variants (ASVs), and generate a feature table of ASV counts and host

metadata. In the quality filtering step, the datasets were truncated to read length of 270 to 250

base pairs for the forward and reverse reads (all other parameters were set to default values).

After quality filtering, bacterial taxonomies were assigned to the ASV feature table using the

Naïve Bayesian Q2 feature classifier as implemented in QIIME2. We compared the data

against a SILVA reference database [36] trained on the V3-V4 region of the 16S rRNA gene.

Merged paired-end sequence reads of each sample were trimmed from the primer

sequences, and low-quality reads (base error rate $> 1\%$) and chimeric sequences were

removed using USEARCH [37]. Filtered sequence reads were assigned to operational

taxonomic units (OTUs) by open-reference OTU picking at 97% identity using UPARSE

algorithm [38] against a SILVA reference database [36]. Taxonomic assignment of the

reference sequence was used as the taxonomy for each OTU, and taxonomy table at the

family level was generated.

LEfSe analysis was performed to identify taxa displaying the largest differences in

microbiota abundance between groups [20]. Only taxa with LDA scores >2.0 and $p < 0.05$,

as determined by Wilcoxon signed-rank test, are shown. All data analyses were performed

using R software v3.5.3.

Metabolome analysis from cecum samples

SCFA concentrations of cecum contents were determined by GC-mass spectrometry.

In brief, 50 mg cecum contents were homogenized with ether. After vigorous shaking using a

Shake Master Neo (Bio-Medical Science) at 1,500 rpm for 10 min, homogenates were

centrifuged at $1,000 \times g$ for 10 min, then the top ether layers of each sample were collected

and transferred to new glass vials. Aliquots of the ether extracts (80 μL) were mixed with 16

mL N-tert-butyldimethylsilyl-N- methyltrifluoroacetamide. The vials were sealed tightly and

heated at 80 °C for 30 min in a water bath. Derivatized samples were loaded onto a 6890N

Network GC System (Agilent Technologies) equipped with an HP-5ms column (0.25 mm ×

30 m × 0.25 mm) and a 5973 Network Mass Selective Detector (Agilent Technologies). The

temperature program used was 60 °C for 3 min, 5 °C/min increases up to 120 °C, then

20 °C/min increases up to 300 °C. Each sample (1.0 mL) was injected with a run time of 30

min. SCFA concentrations were quantified by comparing their peak areas with standards.

SAG-gel processing step 1: single bacterial cell isolation with gel beads

Microfluidic droplet generators were fabricated and used for in-bead bacterial genome

amplification sequencing. After homogenization of mouse feces in PBS, the supernatant was

recovered by centrifugation at $2,000 \times g$ for 2 s, followed by $15,000 \times g$ for 3 min. The

resulting cell pellets were suspended in PBS, filtered through 35-μm nylon mesh, washed

twice, and resuspended in 1.5% agarose in Dulbecco's PBS (DPBS; Thermo Fisher

Scientific).

Prior to single-cell encapsulation, cell suspensions were adjusted to 0.1 cells/bead to

prevent the encapsulation of multiple cells in single beads. Using the droplet generator, single

microbial cells were encapsulated in droplets and collected in a 1.5 mL tube, which was

chilled on ice for 15 min to form the gel matrix. After solidification, collected droplets were

broken with 1H,1H,2H,2H-perfluoro-1-octanol to collect the beads. Then, the gel beads were

washed with 500 µL acetone (Sigma), and the solution was mixed vigorously and centrifuged.

The acetone supernatant was removed, and 500 µL isopropanol (Sigma) was added, and the

solution was mixed vigorously and centrifuged. The isopropanol supernatant was removed,

and the gel beads were washed three times with 500 µL DPBS.

SAG-gel processing step 2: massively parallel single-cell genome amplification

Next, individual cells in beads were lysed by submerging the gel beads in lysis

solutions: first, 50 U/µL Ready-lyse Lysozyme Solution (Epicentre), 2 U/mL Zymolyase

(Zymo research), 22 U/mL lysostaphin (MERCK), and 250 U/mL mutanolysin (MERCK) in

DPBS at 37 °C overnight; second, 0.5 mg/mL achromopeptidase (MERCK) in DPBS at

37 °C for 8 h; and third, 1 mg/mL Proteinase K (Promega) with 0.5% SDS in DPBS at 40 °C

overnight. At each reagent replacement step, the gel beads were washed three times with

DPBS and then resuspended in the next solution. After lysis, the gel beads were washed with

DPBS five times and the supernatant was removed. Then, the beads were suspended in

Buffer D2 and subjected to multiple displacement amplification using a REPLI-g Single Cell

Kit (QIAGEN).

After WGA at 30 °C for 2 h and 65 °C for 3 min, the gel beads were washed three

times with 500 µL DPBS. Then, the beads were stained with 1× SYBR Green in DPBS. After

confirming DNA amplification by the presence of green fluorescence in the gel, fluorescence-positive beads were sorted into 0.8 µL DPBS in 96-well plates by a FACSMelody cell sorter (BD Bioscience) equipped with a 488 nm excitation laser. Green fluorescence intensities were detected through a 550-nm long-pass dichroic filter and a 525-nm band-pass filter using fluorescence detector 1 (FL1) with a PMT voltage of 650 and a logarithmic gain. The sample flow rate was adjusted to approximately 200 events/s. At least 5,000 particles were analyzed for each histogram. After droplet sorting, 96-well plates proceeded through a second round of WGA or were stored at -30 °C.

SAG-gel processing step 3: multiplex single-cell genome sequencing

After gel bead collection in 96-well plates, second-round MDA was performed with the REPLI-g Single Cell Kit. Buffer D2 (0.6 µL) was added to each well and incubated at 65 °C for 10 min. Then, 8.6 µL of MDA mixture was added and incubated at 30 °C for 120 min. The MDA reaction was terminated by heating at 65 °C for 3 min. After second-round amplification, master library plates of single amplified genomes (SAGs) were prepared. For quality control, aliquots of the SAGs were transferred to replica plates, which were used for DNA yield quantification using a Qubit dsDNA High Sensitivity Assay Kit (Thermo Fisher Scientific) and 16S rRNA gene Sanger sequencing (FASMAC) with the V3-V4 primers described above. Based on the DNA amount and 16S rRNA gene sequence, after removing the contaminated samples, to perform whole-genome sequencing, we screened the SAG samples that had enough DNA for library preparation with qualified 16S rRNA sequences obtained by Sanger sequencing (more than 400 bp and HQ 40% were assessed using the Geneious software (Biomatters, Ltd.)). For sequencing analysis, Illumina libraries were prepared using amplicons from the second-round MDA product, using the Nextera XT DNA Library Prep Kit (Illumina) according to the manufacturer's instructions. Each SAG library

was sequenced using an Illumina MiSeq 2 × 75bp or HiSeq 2 × 150bp configuration

(GENEWIZ).

Single-cell genome sequencing analysis

The sequence reads were assembled *de novo* using SPAdes 3.9.0 (options: --sc --

careful --disable-rr) [39], and the contigs were qualified by QUAST 4.5 [40]. CheckM 1.0.6

[41] with the "lineage_wf" command was used to assess the completeness and the

contamination read rate after the contigs smaller than 1,000 bp were removed. Composite

draft genomes were constructed by co-assembly of SAG groups with ≥99% identical marker

genes detected by CheckM and ≥95% ANI. When an assembled composite draft genome

showed ≥10% completeness, the composite draft was reassembled from the same SAG group

by ccSAG [23]. CDS, rRNAs, and tRNAs were extracted from SAG contigs or composite

draft genomes by prokka 1.13 [42], and the metabolic functions were estimated by

Genomaple 2.3.2 [43]. Taxonomic annotation of draft genomes was performed by GTDB-Tk

0.3.0, [25], and *Bacteroides* phylogenetic trees were constructed by the neighbor-joining

method using concatenated marker genes detected by CheckM. PUL clusters of *Bacteroides*

composite genomes were detected by BLASTP [44] searches against the Bo cluster with

default parameters, and phylogenetic trees were constructed from nucleotide sequences of the

PUL clusters in the same manner. Orthologous groups were inferred by OrthoFinder 2.3.3

[45] for additional evaluation of genomic similarities among *Bacteroides* strains.

Structure prediction

Homology modeling of proteins encoded by each gene included in the *Bacteroides*

PUL cluster was performed using SWISS-MODEL [46] in automated mode. Based on the

predicted protein structure, structural alignment was performed using MATRAS [47]. SusC-

SusD complex structure construction was performed by MODELLER 9.20 [48] using

template search and construction scripts created by HOMCOS [49] and using the crystal

structure of BT1762-1763 (PDB_5T4Y) as a template.

Additional files

Additional file 1: Table S1. Statistics for 346 SAGs obtained from the mouse fecal

microbiome **Table S2.** Taxonomic annotation for 346 SAGs and 28 SAGCs obtained from

the mouse fecal microbiome **Table S3.** Comparison of OTU detection between 16S rRNA

gene amplicon sequencing and single-cell sequencing **Table S4.** Statistics for 28 composite

SAGs (IMSAGCs) obtained from the mouse fecal microbiome. **Table S5.** Detailed statistics

for medium and high-quality draft genomes obtained from the mouse fecal microbiome

Table S6. Statistics for *Bacteroides* genomes used in comparative genome analysis **Table S7.**

Numbers of signal peptide-containing predicted glycoside hydrolases/polysaccharide lyases

Table S8. Statistics for high completeness SAGs (> 80%) associated with inulin uptake and

TCA cycle-mediated metabolism **Table S9.** Statistics for conserved metabolic modules in

qualified draft genomes and known *Bacteroides* spp. reference genomes

Additional file 2: Figure S1. Comparison of abundance profiles of IMSAGC_001 (OTU2)

and IMSAGC_004 (OTU4) in 16S rRNA gene sequencing (a), (b) Violin plots indicate

abundance profiles of IMSAGC_001 (OTU2) (a) and IMSAGC_004 (OTU4) (b) before and

after two weeks of inulin or cellulose feeding in Lot 1 mice (n=5, Tukey's HSD test). (c)

Time-dependent changes in abundance profiles of IMSAGC_001 (OTU2) and IMSAGC_004

during 2-weeks inulin feeding in Lot 2 mice (n=9). **Figure S2** Predicted structures of sus

homologs in IMSAGC_001 and IMSAGC_004 (a) Representation of the predicted SusC-

SusD complex protein structure in IMSAGC_001. Units are shown in different colors. The

crystal structure of BT1762-1763 (PDB_5T4Y) in *Bacteroides thetaiotaomicron* was used as

template. (b), (c) Trees based on structural alignments of SusC (b) and SusD (c) among

fructan-utilizing *Bacteroides* strains. **Figure S3.** Predicted partial reverse TCA cycles in

IMSAGC 001 and IMSAGC 004 Succinate is formed by the reversal of partial TCA cycle

reactions. Pyruvate is carboxylated to form oxaloacetate, which is then reduced to malate,

fumarate, and succinate.

Declarations

Ethics approval and consent to participate

The protocols of all animal studies were approved by the Committee for Animal

Experimentation of the School of Science and Engineering, Waseda University (No. 2017-

A057 and 2018-A068)

Consent for publication

Not applicable.

Availability of data and materials

Sequence raw data obtained from mouse gut microbes and high- and medium quality draft

genomes are listed in Table 1 and Additional file1: Data from Table S5 were deposited in the

DNA Data Bank of Japan (DDBJ) under the accession number PRJDB8805.

Competing interests

M.H. and H.T. are shareholders in bitBiome, to which the patents pertaining to the SAG-gel

workflow were transferred.

Funding

This work was supported by JST-PRESTO grant number JPMJPR15FA, MEXT KAKENHI grant numbers 18H01801 and 17H06158, and Cabinet Office, Government of Japan, Cross-ministerial Strategic Innovation Promotion Program (SIP), "Technologies for creating next-generation agriculture, forestry and fisheries"(funding agency: Bio-oriented Technology Research Advancement Institution, NARO).

Authors' contributions

M.H., R.C., and H.T. conceived and designed the experiments. M.H., Y.N., and K.M. developed the SAG-gel workflow. R.C. conducted animal experiments, metagenomic analysis, and metabolomic analysis. M.H., Y.N. M.K. C.S., and K.T. conducted single-cell genomics experiments and collected the data. M.H., R.C., M.K., and K.I conducted bioinformatic analysis of the metagenomic data and single-cell genomic data. M.H., R.C., M.K., K.I., and H.T. wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgments

References

1.   Neis EP, Dejong CH, Rensen SS: The role of microbial amino acid metabolism in host metabolism. Nutrients. 2015;7**:**2930-46.

2.   Human Microbiome Project C: Structure, function and diversity of the healthy human microbiome. Nature. 2012;486**:**207-14.

3.   Macia L, Tan J, Vieira AT, Leach K, Stanley D, Luong S, Maruya M, Ian McKenzie

C, Hijikata A, Wong C, et al: Metabolite-sensing receptors GPR43 and GPR109A

facilitate dietary fibre-induced gut homeostasis through regulation of the

inflammasome. Nat Commun. 2015;6**:**6734.

4.   Poeker SA, Geirnaert A, Berchtold L, Greppi A, Krych L, Steinert RE, de Wouters T,

Lacroix C: Understanding the prebiotic potential of different dietary fibers using an in

vitro continuous adult fermentation model (PolyFermS). Sci Rep. 2018;8**:**4318.

5.   Kovatcheva-Datchary P, Shoaie S, Lee S, Wahlstrom A, Nookaew I, Hallen A,

Perkins R, Nielsen J, Backhed F: Simplified Intestinal Microbiota to Study Microbe-

Diet-Host Interactions in a Mouse Model. Cell Rep. 2019;26**:**3772-83 e6.

6.   Schroeder BO, Birchenough GMH, Stahlman M, Arike L, Johansson MEV, Hansson

GC, Backhed F: Bifidobacteria or Fiber Protects against Diet-Induced Microbiota-

Mediated Colonic Mucus Deterioration. Cell Host Microbe. 2018;23**:**27-40 e7.

7.   Rakoff-Nahoum S, Coyne MJ, Comstock LE: An ecological network of

polysaccharide utilization among human intestinal symbionts. Curr Biol. 2014;24**:**40-

9.

8.   Rakoff-Nahoum S, Foster KR, Comstock LE: The evolution of cooperation within the

gut microbiota. Nature. 2016;533**:**255-9.

9.   Bolam DN, Sonnenburg JL: Mechanistic insight into polysaccharide use within the

intestinal microbiota. Gut Microbes. 2011;2**:**86-90.

10. Vandeputte D, Falony G, Vieira-Silva S, Wang J, Sailer M, Theis S, Verbeke K, Raes

J: Prebiotic inulin-type fructans induce specific changes in the human gut microbiota.

Gut. 2017;66**:**1968-74.

11.   Sonnenburg ED, Zheng H, Joglekar P, Higginbottom SK, Firbank SJ, Bolam DN,

2   Sonnenburg JL: Specificity of polysaccharide use in intestinal bacteroides species

3   determines diet-induced microbiota alterations. Cell. 2010;141**:**1241-52.

4   12.   Catry E, Bindels LB, Tailleux A, Lestavel S, Neyrinck AM, Goossens JF, Lobysheva

5   I, Plovier H, Essaghir A, Demoulin JB, et al: Targeting the gut microbiota with inulin-

6   type fructans: preclinical demonstration of a novel approach in the management of

7   endothelial dysfunction. Gut. 2018;67**:**271-83.

8   13.   Langille MG, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA,

9   Clemente JC, Burkepile DE, Vega Thurber RL, Knight R, et al: Predictive functional

10   profiling of microbial communities using 16S rRNA marker gene sequences. Nat

11   Biotechnol. 2013;31**:**814-21.

12   14.   Scholz M, Ward DV, Pasolli E, Tolio T, Zolfo M, Asnicar F, Truong DT, Tett A,

13   Morrow AL, Segata N: Strain-level microbial epidemiology and population genomics

14   from shotgun metagenomics. Nat Methods. 2016;13**:**435-8.

15   15.   Pasolli E, Asnicar F, Manara S, Zolfo M, Karcher N, Armanini F, Beghini F, Manghi

16   P, Tett A, Ghensi P, et al: Extensive Unexplored Human Microbiome Diversity

17   Revealed by Over 150,000 Genomes from Metagenomes Spanning Age, Geography,

18   and Lifestyle. Cell. 2019;176**:**649-62 e20.

19   16.   Almeida A, Mitchell AL, Boland M, Forster SC, Gloor GB, Tarkowska A, Lawley

20   TD, Finn RD: A new genomic blueprint of the human gut microbiota. Nature.

21   2019;568**:**499-504.

22   17.   Nayfach S, Shi ZJ, Seshadri R, Pollard KS, Kyrpides NC: New insights from

23   uncultivated genomes of the global human gut microbiome. Nature. 2019;568**:**505-10.

24   18.   Thaiss CA, Zeevi D, Levy M, Zilberman-Schapira G, Suez J, Tengeler AC,

25   Abramson L, Katz MN, Korem T, Zmora N, et al: Transkingdom control of

microbiota diurnal oscillations promotes metabolic homeostasis. Cell. 2014;159**:**514-29.

19.   Thaiss CA, Levy M, Korem T, Dohnalova L, Shapiro H, Jaitin DA, David E, Winter DR, Gury-BenAri M, Tatirovsky E, et al: Microbiota Diurnal Rhythmicity Programs Host Transcriptome Oscillations. Cell. 2016;167**:**1495-510 e12.

20.   Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C: Metagenomic biomarker discovery and explanation. Genome Biol. 2011;12**:**R60.

21.   Zhang M, Zhang Y, Scheuring CF, Wu CC, Dong JJ, Zhang HB: Preparation of megabase-sized DNA from a variety of organisms using the nuclei method for advanced genomics research. Nat Protoc. 2012;7**:**467-78.

22.   Hosokawa M, Nishikawa Y, Kogawa M, Takeyama H: Massively parallel whole genome amplification for single-cell sequencing using droplet microfluidics. Sci Rep. 2017;7**:**5199.

23.   Kogawa M, Hosokawa M, Nishikawa Y, Mori K, Takeyama H: Obtaining high-quality draft genomes from uncultured microbes by cleaning and co-assembly of single-cell amplified genomes. Sci Rep. 2018;8**:**2059.

24.   Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK, Schulz F, Jarett J, Rivers AR, Eloe-Fadrosh EA, et al: Minimum information about a single amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea. Nat Biotechnol. 2017;35**:**725-31.

25.   Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA, Hugenholtz P: A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. Nat Biotechnol. 2018;36**:**996-1004.

26.   Glenwright AJ, Pothula KR, Bhamidimarri SP, Chorev DS, Basle A, Firbank SJ, Zheng H, Robinson CV, Winterhalter M, Kleinekathofer U, et al: Structural basis for

nutrient acquisition by dominant members of the human gut microbiota. Nature. 2017;541**:**407-11.

27. Spiga L, Winter MG, Furtado de Carvalho T, Zhu W, Hughes ER, Gillis CC, Behrendt CL, Kim J, Chessa D, Andrews-Polymenis HL, et al: An Oxidative Central Metabolism Enables Salmonella to Utilize Microbiota-Derived Succinate. Cell Host Microbe. 2017;22**:**291-301 e6.

28. Connors J, Dawe N, Van Limbergen J: The Role of Succinate in the Regulation of Intestinal Inflammation. Nutrients. 2018;11.

29. Lan F, Demaree B, Ahmed N, Abate AR: Single-cell genome sequencing at ultra-high-throughput with microfluidic droplet barcoding. Nat Biotechnol. 2017;35**:**640-6.

30. Xu L, Brito IL, Alm EJ, Blainey PC: Virtual microfluidics for digital quantification and single-cell sequencing. Nat Methods. 2016;13**:**759-62.

31. Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ, Cheng JF, Darling A, Malfatti S, Swan BK, Gies EA, et al: Insights into the phylogeny and coding potential of microbial dark matter. Nature. 2013;499**:**431-7.

32. Rinke C, Lee J, Nath N, Goudeau D, Thompson B, Poulton N, Dmitrieff E, Malmstrom R, Stepanauskas R, Woyke T: Obtaining genomes from uncultivated environmental microorganisms using FACS-based single-cell genomics. Nat Protoc. 2014;9**:**1038-48.

33. Miyamoto Y, Itoh K: Bacteroides acidifaciens sp. nov., isolated from the caecum of mice. Int J Syst Evol Microbiol. 2000;50 Pt 1**:**145-8.

34. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ, Arumugam M, Asnicar F, et al: Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol. 2019;37**:**852-7.

35.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP: DADA2:

High-resolution sample inference from Illumina amplicon data. Nat Methods.

2016;13**:**581-3.

36.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO:

The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. Nucleic Acids Res. 2013;41**:**D590-6.

37.   Edgar RC: Search and clustering orders of magnitude faster than BLAST.

Bioinformatics. 2010;26**:**2460-1.

38.   Edgar RC: UPARSE: highly accurate OTU sequences from microbial amplicon reads.

Nat Methods. 2013;10**:**996-8.

39.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, et al: SPAdes: a new genome assembly

algorithm and its applications to single-cell sequencing. J Comput Biol. 2012;19**:**455-

77.

40.   Gurevich A, Saveliev V, Vyahhi N, Tesler G: QUAST: quality assessment tool for

genome assemblies. Bioinformatics. 2013;29**:**1072-5.

41.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW: CheckM: assessing

the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. Genome Res. 2015;25**:**1043-55.

42.   Seemann T: Prokka: rapid prokaryotic genome annotation. Bioinformatics.

2014;30**:**2068-9.

43.   Arai W, Taniguchi T, Goto S, Moriya Y, Uehara H, Takemoto K, Ogata H, Takami

H: MAPLE 2.3.0: an improved system for evaluating the functionomes of genomes

and metagenomes. Biosci Biotechnol Biochem. 2018;82**:**1515-7.

44.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: Basic local alignment search

2   tool. J Mol Biol. 1990;215**:**403-10.

3   45.   Emms DM, Kelly S: OrthoFinder: solving fundamental biases in whole genome

4   comparisons dramatically improves orthogroup inference accuracy. Genome Biol.

5   2015;16**:**157.

6   46.   Schwede T, Kopp J, Guex N, Peitsch MC: SWISS-MODEL: An automated protein

7   homology-modeling server. Nucleic Acids Res. 2003;31**:**3381-5.

8   47.   Kawabata T: MATRAS: A program for protein 3D structure comparison. Nucleic

9   Acids Res. 2003;31**:**3367-9.

48.   Marti-Renom MA, Stuart AC, Fiser A, Sanchez R, Melo F, Sali A: Comparative

protein structure modeling of genes and genomes. Annu Rev Biophys Biomol Struct.

2000;29**:**291-325.

49.   Fukuhara N, Kawabata T: HOMCOS: a server to predict interacting protein pairs and

interacting sites by homology modeling of complex structures. Nucleic Acids Res.

2008;36**:**W185-9.

Figure legends

Figure 1. Fecal microbial composition and cecum metabolites in inulin-fed mice

a. Relative family-level abundance profiles of mouse fecal microbiomes in the morning and

evening before and after two weeks of inulin or cellulose feeding. Box plots indicate the

phylogenetic composition of the fecal microbiota samples obtained from each mouse by 16S

rRNA gene sequencing (n = 5).

b, c. Linear discriminant analysis (LDA) scores computed for differentially abundant taxa in

the fecal microbiomes of mice before fiber feeding, cellulose-, and inulin-fed mice in the

morning (**b**) or evening (**c**) obtained from each mouse by 16S rRNA gene sequencing. The

length indicates the effect size associated with each taxon. *P* <0.05 by Wilcoxon signed-rank

test; LDA score >2. **d.** Time-dependent changes in relative family-level abundance profiles of

the microbiota in each mouse by 16S rRNA gene sequencing. Nine mice in three different

cages from a different animal lot from (**a**) were fed inulin for two weeks, and fecal samples

were collected in the morning for microbiome composition analysis. Box plots indicate the

phylogenetic compositions of fecal microbiota samples from each mouse (n = 3, one from

each cage). **e, f.** Concentrations of butyrate (**e**) and succinate (**f**) in the cecum of mice before

and after two weeks of inulin or cellulose feeding in the morning or evening. The mice are

the same as those shown in (**a**). Violin plots indicate the SCFA concentration in each mouse

(n = 5, Tukey's HSD test).

Figure 2. Single-cell genome sequencing of fecal microbiomes in inulin-fed mice

a. Workflow for SAG-gel-based single-cell genome sequencing of bacterial cells in a

complex microbial community. Individual bacterial cells are randomly captured in picoliter-

sized gel beads and processed by in-gel lysis and whole-genome amplification (WGA).

Single-cell amplified genomes (SAGs) in the gel are fluorescently detected and sorted into

well plates as a SAG library. The SAGs are re-amplified for further analysis by NGS, amplicon sequencing, and storage. **b-f** Assembly qualities of 324 SAGs, excluding short fragments (length < 100 kb) and contaminated samples (contamination > 10%). **b.** Completeness and contamination statistics for 302 SAGs (IMSAGs) and 22 composite SAGs (IMSAGCs). **c-f.** The number of contigs, N50 values, number of tRNAs, and 16S rRNA gene lengths for 324 SAGs.

Figure 3. Comparative analysis of the inulin responder *Bacteroides* genomes IMSAGC_001 and IMSAGC_004

a. Phylogenetic tree of inulin-associated *Bacteroides* spp. found in this study and reference *Bacteroides* strains. The tree is based on the alignment of concatenated amino acid sequences of collocated sets of ubiquitous, single-copy genes within a phylogenetic lineage, as used in CheckM analysis. IMSAGC_001 and IMSAGC_004 are indicated by black circles.

b. UpSet plot comparing shared orthogroup counts between reference *Bacteroides* strains. Orthogroups shared between IMSAGC_001 and other *Bacteroides* group members are colored in orange, whereas orthogroups shared between IMSAGC_004 and other *Bacteroides* group members are colored in blue. Orthogroups shared between IMSAGC_001, IMSAGC_004, and other members are colored in green. Bt, *B. thetaiotaomicron*; Bo, *B. ovatus*; Bc, *B. caccae*; Bv, *B. vulgatus*; Bu, *B. uniformis*; Bf, *B. fragilis*; Ba, *B. acidifaciens*.

Figure 4. Identification of putative inulin utilization loci in the inulin responder *Bacteroides* genomes IMSAGC_001 and IMSAGC_004

a. Polysaccharide utilization loci (PUL) from reference *Bacteroides* species, IMSAGC_001, and IMSAGC_004. Common predicted functions are color-coded while intervening unrelated genes are in gray. Signal peptide, SP; glycoside hydrolase, GH; polysaccharide lyase, PL. **b.**

1   Heat map of the identities of amino acid sequences in PUL genes between inulin responder

2   (001 and 004) and non-responder (006 and 014) *Bacteroidales* and the known inulin utilizer

3   *B. ovatus*. **c, d.** Phylogenetic analysis of glycoside hydrolase (**c**) and a concatenated sequence

4   containing *SusC* and *SusD* (**d**) between reference *Bacteroides* species, IMSAGC_001, and

5   IMSAGC_004. IMSAGC_001 and IMSAGC_004 are indicated with black circles.

Table 1 Statistics for high and medium quality draft genomes

SAG ID	Lineage[a]	# contigs	Total length (Mb)	Largest contig (bp)	N50 (bp)	GC (%)	# CDS	Completeness (%)[b]	Contamination (%)[b]	SAG quality[c]
IMSAGC_001	Bacteroidota;Bacteroidia;Bacteroidales;Bacteroidaceae;Bacteroides;Bacteroides acidifaciens	369	4.93	125378	29923	43.1	4165	97.27	1.12	High
IMSAGC_002	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;28-4	709	5.08	70754	13000	44.6	4725	93.87	1.78	Medium
IMSAGC_003	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;CAG-65	308	4.62	140873	48295	50.7	4203	98.85	3.97	Medium
IMSAGC_004	Bacteroidota;Bacteroidia;Bacteroidales;Bacteroidaceae;Bacteroides_B	305	4.22	106819	34968	43.1	3598	98.62	0.8	Medium
IMSAGC_005	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;CAG-95	317	4.23	113574	39279	43.6	4110	96.45	1.45	Medium
IMSAGC_006	Bacteroidota;Bacteroidia;Bacteroidales;Muribaculaceae;UBA3263;GCA_001689615.1	265	2.42	113083	24935	50.4	2284	85.47	0.89	Medium
IMSAGC_007	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae	496	5.06	122936	25158	47.1	4957	95.89	3.59	Medium
IMSAGC_008	Bacteroidota;Bacteroidia;Bacteroidales;Muribaculaceae;CAG-873	380	2.66	59381	16733	52.1	2436	78.2	1.58	Medium
IMSAGC_009	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;CAG-95	597	5.02	113931	23340	44.4	4633	95.33	2.87	Medium
IMSAGC_010	Firmicutes;Bacilli;Lactobacillales;Lactobacillaceae;Lactobacillus;Lactobacillus johnsonii	54	1.91	214598	104736	34.3	1789	99.22	0.78	High
IMSAGC_011	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;COE1	431	4.25	143151	27757	37.9	3664	92.66	2.59	High
IMSAGC_012	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;Eubacterium_J	419	3.92	117373	25561	45.0	3748	92.92	4.75	Medium
IMSAGC_013	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;14-2	884	5.14	67964	11980	44.6	4945	85.42	0.95	Medium
IMSAGC_014	Bacteroidota;Bacteroidia;Bacteroidales;Bacteroidaceae;F0040	273	2.64	87270	20006	47.1	2340	95.18	1.58	Medium
IMSAGC_015	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;Dorea	388	2.55	63422	12891	45.3	2429	67.29	2.53	Medium
IMSAGC_016	Bacteroidota;Bacteroidia;Bacteroidales;Muribaculaceae;CAG-485	385	1.94	50916	8558	46.4	1735	56.91	0.5	Medium
IMSAGC_017	Firmicutes;Bacilli;Erysipelotrichales;Erysipelatoclostridiaceae;Erysipelatoclostridium;Erysipelatoclostridium cocleatum	299	2.69	71470	19801	28.9	2384	95.28	0.94	Medium
IMSAGC_019	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;Dorea	311	2.42	198629	20896	44.1	2370	62.98	0	Medium
IMSAGC_020	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;CAG-95	756	4.13	136678	11653	45.8	4150	57.89	2.63	Medium
IMSAGC_021	Bacteria;Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;Dorea	642	3.03	69412	9299	45.8	2954	60.15	2.58	Medium
IMSAGC_027	Bacteroidota;Bacteroidia;Bacteroidales;Muribaculaceae;CAG-1031;GCA_001689585.1	314	2.00	63820	15566	50.5	1738	59.25	0	Medium
IMSAGC_028	Bacteroidota;Bacteroidia;Bacteroidales;Rikenellaceae;Alistipes	265	1.73	77838	22105	52.5	1581	56.76	1.92	Medium
IMSAG_013	Firmicutes_A;Clostridia;Oscillospirales;CAG-272	229	1.72	93908	16943	45.4	1622	59.65	0	Medium
IMSAG_025	Bacteroidota;Bacteroidia;Bacteroidales;Muribaculaceae;CAG-485	667	2.51	52448	6077	45.3	2558	50	7.76	Medium
IMSAG_044	Firmicutes;Bacilli;Lactobacillales;Lactobacillaceae;Lactobacillus_H;Lactobacillus_H reuteri_A	200	1.82	108153	17958	38.3	1735	86.54	0.54	Medium
IMSAG_049	Firmicutes_A;Clostridia;Lachnospirales	317	1.74	50322	11294	39.6	1762	54.88	1.34	Medium
IMSAG_117	Firmicutes;Bacilli;Lactobacillales;Lactobacillaceae;Lactobacillus_B;Lactobacillus_B murinus	314	1.86	38237	10806	40.5	1824	78.75	1.31	Medium
IMSAG_185	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;CAG-65	317	2.25	78895	17758	49.9	2117	54.01	1.45	Medium
IMSAG_192	Bacteroidota;Bacteroidia;Bacteroidales;Muribaculaceae;Muribaculum;Muribaculum intestinale	342	1.89	59599	11500	49.3	1711	57.55	0	Medium
IMSAG_249	Firmicutes_A;Clostridia;Lachnospirales;Lachnospiraceae;CAG-95	428	2.83	108242	17278	44.4	2659	55.18	0.91	Medium
IMSAG_250	Firmicutes_A;Clostridia;Oscillospirales;DTU089;Eubacterium_R	147	1.35	87509	28237	37.8	1309	68.68	0	Medium

[a] Assigned using GTDB-Tk.

[b] Estimated using CheckM.

[c] Classified according to GSC MISAG.

a

Before | Cellulose | Inulin

Abundance (%)

Morning Evening | Morning Evening | Morning Evening

Remainder
Streptococcaceae
Lachnospiraceae
Ruminococcaceae
Lactobacillaceae
Muribaculaceae
Enterococcaceae
Rikenellaceae
Staphylococcaceae
Erysipelotrichaceae
Bacteroidaceae

b

Morning

Before | Cellulose | Inulin

P_Bacteroidetes
O_Bacteroidales
C_Bacteroidia
F_Bacteroidaceae
O_Verrucomicrobiales
F_Akkermansiaceae
P_Verrucomicrobia
C_Verrucomicrobiae
F_Ruminococcaceae
C_Erysipelotrichia
O_Erysipelotrichales
F_Erysipelotrichaceae
F_Rikenellaceae
P_Proteobacteria
O_Enterobacteriales
C_Gammaproteobacteria
F_Enterobacteriaceae
C_Bacilli
O_Lactobacillales
P_Firmicutes
F_Enterococcaceae
F_Staphylococcaceae
O_Bacillales
F_Streptococcaceae

LDA SCORE (log 10)

c

Evening

Before | Cellulose | Inulin

F_Bacteroidaceae
O_Bacteroidales
P_Bacteroidetes
C_Bacteroidia
F_Enterococcaceae
P_Firmicutes
C_Clostridia
O_Clostridiales
F_Streptococcaceae
F_Lachnospiraceae
O_Bacillales
F_Staphylococcaceae
F_Ruminococcaceae
F_Lactobacillaceae
F_Rikenellaceae

LDA SCORE (log 10)

d

Day 0 | Day 1 | Day 3 | Day 4 | Day 5 | Day 6 | Day 7 | Day 10 | Day 14

Abundance (%)

Cage No. 1 2 3

Remainder
Rikenellaceae
Prevotellaceae
Akkermansiaceae
Muribaculaceae
Bifidobacteriaceae
Lactobacillaceae
Streptococcaceae
Ruminococcaceae
Lachnospiraceae
Bacteroidaceae

e

Butyrate

μmol/g cecal contents

0.005
0.0009
0.003

Before Cellulose Inulin | Before Cellulose Inulin
Morning | Evening

f

Succinate

μmol/g cecal contents

1e-07
< 1e-07
3e-07
2e-07

Before Cellulose Inulin | Before Cellulose Inulin
Morning | Evening