Dissecting the structural and functional roles of a vicinal iron-binding site in encapsulated ferritins

Cecilia Piergentili[1,♣], Didi He[2,#,♣], Jennifer Ross[3,♣], Will A. Stanley[1], Laurène Adam[1], C. Logan Mackay[3], Kevin J. Waldron[4], David J. Clarke[3*], Jon Marles-Wright[1*]

[1]School of Natural and Environmental Sciences, Newcastle University. Newcastle upon Tyne, NE1 7RU; [2]Institute of Quantitative Biology, Biochemistry and Biotechnology, School of Biological Sciences, The University of Edinburgh, Max Born Crescent, Edinburgh, EH9 3BF; [3]EaStCHEM School of Chemistry, The University of Edinburgh, Joseph Black Building, David Brewster Road, Edinburgh, Scotland, EH9 3FJ; [4]Institute for Cell and Molecular Biosciences, Newcastle University, Newcastle upon Tyne, NE2 4HH.

[#]*Current address: Structural Genomics Consortium, University of Oxford, Old Road Campus Research Building, Roosevelt Drive, Oxford, OX3 7DQ*

[♣] These authors contributed equally to this work

[*] Corresponding authors

Contact details:

Email: Dave.clarke@ed.ac.uk

Tel: +44(0)131 650 4808

Email: Jon.marles-wright1@ncl.ac.uk

Tel: +44(0)191 2084855

Abstract

The Encapsulated ferritin-like proteins belong to the universally distributed ferritin superfamily, which function as iron detoxification and storage systems. The encapsulated ferritins have a distinct annular structure and must associate with an encapsulin nanocage to form a competent iron store capable of holding significantly more iron than classical ferritins. The catalytic mechanism of iron oxidation in the ferritin family is still an open question due to differences in organisation of the ferroxidase catalytic site and secondary metal binding sites vicinal to this. We have previously identified a putative metal binding site on the inner surface of *Rhodospirillum rubrum* encapsulated ferritin at the interface between the two-helix subunits and proximal to the ferroxidase centre. Here we present a comprehensive structural and functional study to investigate the functional relevance of this putative iron entry site by means of enzymatic assays, mass-spectrometry, and X-ray crystallography. We show that catalysis occurs in the ferroxidase centre and suggest a dual role for the secondary site as an electrostatic trap guiding ferrous ions toward the ferroxidase centre and, at the same time, acting as a barrier protecting the ferroxidase site against non-cognate inhibiting species. Moreover, confinement of encapsulated ferritins within the encapsulin nanocage, while enhancing the ferritin ability to undergo catalysis, does not influence the specific function of the secondary site.

Introduction

The ferritin superfamily consists of a number of structurally and functionally distinct members, each of which is built around a four-helix bundle scaffold for an iron-binding ferroxidase active site (Andrews, 2010). The quaternary structure of these protein families show a high degree of plasticity: from the monomeric rubrerythrin proteins (Dillard et al., 2011; Cardenas et al., 2016); dimeric IMEF proteins (Giessen et al. 2019); dodecameric DNA protection in Starved cells (DPS) (Grant et al., 1998; Pesek et al., 2011; Martinez et al., 1997); the 24-meric classical (Ftn) and bacterioferritins (Bfr) (Andrews, 1998, Pfaffen et al., 2013; Frolow et al., 1994; Harrison et al., 1996); and the decameric encapsulated ferritins (EncFtn) (He et al. 2016, 2019). The rubrerythrin and DPS proteins play key roles in the protection of cells from oxidative stress (Ratnayake et al., 2000; Sztukowska et al., 2002; Almirón et al., 1992; Martinez and Kolter, 1997; Choi et al., 2000; Nair et al., 2004; Chiancone and Ceci, 2010; De Martino et al., 2016); while DPS and the other members of the ferritin superfamily are vital iron stores. These latter proteins oxidise and convert free Fe(II) to inert Fe(III) oxyhydroxide or Fe(III) phosphate mineral forms, to provide a reservoir of bioavailable iron (Chasteen and Harrison, 1999). The nanocages formed by the quaternary structure of DPS, Ftn, and Bfr, allow iron oxidation and mineralisation to occur within a biochemically privileged environment within the cell, which protecting the host cell from iron-induced oxidative stress. Interestingly, the recently described IMEF and EncFtn proteins have ferroxidase activity, but they do not form nanocage structures on their own, instead they must be localised to the lumen of encapsulin nanocages to form a competent iron-store (Giessen et al., 2019; He et al., 2016, 2019).

The catalytic route to iron mineralisation in the different ferritin iron-stores is still a subject of vigorous debate (Xu and Chasteen, 1991; Proulx-Curry and Chasteen, 1995; Yang et al., 1998; Chasteen and Harrison, 1999; Baaghil et al., 2003; Ebrahimi et al., 2009) and given the lack of absolute conservation in the ferroxidase centre, it is likely that there is no single conserved mechanism for their activity ((Xu and Chasteen, 1991; Proulx-Curry and Chasteen, 1995; Yang et al., 1998; Chasteen and Harrison, 1999; Baaghil et al., 2003; Ebrahimi et al., 2009). All of the ferritin family proteins that have been structurally characterised have additional iron-binding sites to the ferroxidase centre. In the case of sites proximal to the ferroxidase centre in DPS and ferritins, these have been suggested to be iron-entry, transit, and exit points from the ferroxidase centre (Theil et al., 2013; Masuda et al., 2010; Behera et al., 2015), or the sites of electron transfer from Fe(II) substrates to the FOC (Yao et al., 2012). The subunit boundaries in the DPS and ferritin nanocages form a variety of symmetrical pores, which have variously been suggested and shown to be entry and exit sites to the lumen for iron, water and phosphate substrates (Treffry and Harrison, 1978; Andrews, 1998; Liu et al., 2007; Bellapadrona et al., 2009; Giessen et al., 2019). Whatever their function, these secondary metal sites are essential to the correct functioning of the ferritin iron-stores (Treffry et al., 1998; Ebrahimi et al., 2016; Hagen et al., 20refs).

In our previous work on the EncFtn family we identified a number of conserved secondary metal-binding sites in their structures (He et al., 2016; 2019). A partially occupied 'weak' site on the outer surface of the EncFtn decamer (Figure 1A in He et al., 2016) and a highly occupied 'entry' site on the inner face, 17; which is proximal to the ferroxidase centre (Figure 1B in He et al., 2016). Both of these sites are found at dimer interfaces between the two-helix EncFtn subunits and are formed when the four-helical ferritin fold is reconstituted by the iron-mediated assembly of the EncFtn ferroxidase centre (Ross et al., *submitted*). In our structure of the *Rhodospirillum rubrum* EncFtn protein, the entry site coordinates a calcium ion, from the crystallisation solution, in a symmetrical arrangement of glutamic acid residues (E31/E34) and a tryptophan residue (W38) contributed by each chain from the ferroxidase centre dimer (Ross et al., *submitted*) (Figure 1). The metal coordination distances between the side-chain carboxylic acid oxygens and calcium ion are 2.1 Å Ca – Glu31-Ox, 2.7 Å Glu34-Ox, which is similar to the Fe-Glu-Ox coordination distances seen in the FOC of 2.2-2.6 Å.

We hypothesise that this site acts as an entry site for iron ions approaching the ferroxidase centre, and acts as a filter to control the access of metal ions to the active site of the protein. The glutamic acid residues in this site are arranged in such a way as to capture and coordinate an Fe(II) ion in an octahedral geometry before passing it to the FOC. The positioning of W38 adjacent to this site suggests that the indole ring sterically constrains the position of the E31 side chain to an optimal position for metal coordination. We have considered several hypotheses regarding the functional role of this additional metal-binding site, including a gate prior to the channel leading to FOC; an electrostatic trap to direct metal ions to the FOC; a discriminatory filter to avoid mismetallation and consequent inhibition of the active site (Figure 2); or for protein stability and assembly. These functions may overlap, and the site may play a number of key roles in the stability and activity of the Encapsulated ferritins.

In order to understand the role of this site a number of single-point variants of the *R. rubrum* EncFtn protein were produced by site-directed mutagenesis to replace the residue of interest with non-metal binding residues. These variants were produced in the isolated *R. rubrum* EncFtn protein and an *R. rubrum* Encapsulin:EncFtn (Enc:EncFtn) co-expression system. We produced and purified recombinant E31A, E34A and W38G/A variants of EncFtn and the Enc:EncFtn complex and analysed the solution and gas phase behaviour of these proteins to understand the influence of these variants on the stability of the EncFtn complex. We obtained X-ray crystal structures of the E31A and E34A variants of the EncFtn protein and their structures display altered entry site geometry, with the abrogation of metal binding in this site in the E34A variant. The functional role of the secondary metal binding site was explored biochemically, we show that changes of metal coordination in this site enhances the ferroxidase activity of the EncFtn protein, while increasing its susceptibility to inhibition by competing zinc ions. Taken together, these results indicate that this metal binding site plays a minor role in stabilising the oligomeric state of the EncFtn decamer, and primarily acts as an electrostatic filter to control metal access to the ferroxidase centre.

Methods
Cloning
Expression constructs for the *R. rubrum* EncFtn protein (pACYCDuet-1-Rru_EncFtn) and Enc:EncFtn protein complex (pACYCDuet-1-Enc:EncFtn) were produced in the previous work (He et al., 2016). Site-directed mutagenesis was performed using Stratagene QuikChange kit following manufacturer instructions. Protein nomenclature used throughout the text is shown in Supplementary Table S1. Primers used in this work for this purpose are listed in Supplementary Table S2.

Protein production and purification
Expression plasmids for the *R. rubrum* EncFtn protein and Enc:EncFtn complex were transformed into *Escherichia coli* BL21(DE3) cells and grown in LB medium and expressed as described previously (He et al., 2016; 2019).

EncFtn and its variants, were purified as described in He et al., 2019; concisely, the proteins were purified by combining anion exchange chromatography (AXC) and Size exclusion chromatography (SEC). Purity of protein samples was analysed by SDA-PAGE.

Enc:EncFtn wild-type and variant complex cell pellets were resuspended in 10-times (v/w) of Lysis Buffer (20 mM HEPES pH 8.0, 2 mM $MgCl_2$, 1 mg/ml lysozyme, 20 units/ml benzonase). Sonication on ice (5 minutes at 30 seconds on/off cycles, 35 %) of cell sample was followed by incubation for 30 minutes at 37 °C using a water bath. Cell lysate was clarified by centrifugation (35,000 x g, 30 minutes, 4 °C) and the supernatant was collected and incubated at 80 °C for 10 minutes. After cooling the sample was centrifuged again (30 minutes, 35,000 x g, 4 °C), and the supernatant was collected and filtered using a 0.22 um syringe filter (Millipore, UK). Enc:EncFtn protein complex sample was loaded onto a 20 ml anion exchange column (Hi-Trap Q-sepharose FF, GE Healthcare) pre-

equilibrated with Buffer QA (50 mM Tris, pH 8.0). Un-bound protein was washed with Buffer QA.
A step gradient elution was performed by mixing Buffers QA and QB (50 mM Tris, pH 8.0, 1 M
NaCl). Since Enc:EncFtn does not bind the Q-sepharose column, proteins of interest were found in
flow-through sample, as confirmed by 15 % (w/v) SDS-PAGE. Protein was concentrated using a
Vivaspin centrifugal concentrator (MWCO 10 kDa) following the manufacturer's instructions.
Concentrated protein was then applied to a Size exclusion column (HiLoad 16/600 Superdex 200, GE
Healthcare) pre-equilibrated with Buffer GF (50 mM Tris, pH 8.0, 150 mM NaCl). Fractions of
interest were run on a 15% (w/v) SDS-PAGE to assess protein purity and oligomerisation state, and
further concentrated using a Vivaspin centrifugal concentrator as before. After concentration, protein
samples were analysed by 10% Tricine-SDS-PAGE (Schägger, 2006) in order to resolve bands
around 30 kDa. Elution volumes of EnFtn and Enc:EncFtn complexes from the size exclusion column
are listed in Supplementary Table S3.
162
His-tagged protein production and purification
Hexahistidine tagged EncFtn E31A and E34A variants (EncFtn_sH) were produced in *E. coli*
BL21(DE3) as described above and purified following the same protocols as previously described
(He et al., 2016). Briefly, a 5 ml HisTrap column (GE Healthcare, UK) was used for immobilised
metal affinity chromatography (IMAC), followed by size exclusion chromatography (HiLoad 16/600
Superdex 200, GE Healthcare) using the protocol described above.
169
Protein quantification
Both untagged and His-tagged EncFtn proteins were quantified using the Pierce BCA Assay kit
(Thermo Fisher Scientific) following the manufacturer's instructions for the Test-tube procedure.
Concentration of Enc:EncFtn proteins was determined using the Beer-Lambert equation, measuring
absorbance at 280 nm with a Nanodrop Spectrophotometer (Thermo Scientific) and using an
extinction coefficient calculated by ProtParam tool on the Expasy platform entering a sequence
composed of 3 x Enc and 2 x Enc-Ftn sequences. This ratio was determined by examination of SDS-
PAGE gels of Enc:EncFtn proteins and densitometry of bands.
178
Transmission electron microscopy
TEM imaging was performed as previously described (He et al., 2016) on purified Enc:EncFtn
proteins. Proteins were diluted in Buffer GF to a final concentration of 0.1 mg/ml before being spotted
on glow-discharged 300 mesh carbon-coated copper grids. Excess liquid was removed by blotting
with filter paper (Whatman, UK) before sample was stained with 0.2 % uranyl acetate, blotted again,
and air-dried. Grids were imaged using a Hitachi HT7800 transmission electron microscope (LaB6
filament) and images were collected with an EMSIS Xarosa camera (Radius software). A
representative set of images are presented here.
187
Mass spectrometry analysis
Sample preparation, instrumentation, native MS conditions, ion mobility-MS conditions and the
method of collision cross section (CCS) determination have all been reported in previous works by
our group (He et al., 2016; 2019). For collision induced unfolding (CIU) experiments, protein samples
were prepared and the Synapt G2 was set up as for IM-MS. Samples were analysed with incremental
increasing trap voltages between 10 V to 60 V, using 2 V increments. Data was processed using
Spyder CIUSuite (Dixit et al., 2018).
195
Protein crystallization and X-ray data collection
EncFtn_sH variants (E31A or E34A) were concentrated to 10 mg/ml (based on extinction coefficient
calculation) and subjected to crystallisation under similar conditions to the wild-type protein
(WT)（Supplementary Table S4）．Crystallization drops were set up in 24−well Linbro plates using
the hanging drop vapor diffusion method at 292 K. Glass coverslips were set up with 1–2 µl protein
mixed with 1 µl well solution and sealed over 1 ml of well solution. Crystals appeared after one week

to two months and were mounted using a LithoLoop (Molecular Dimensions Limited, UK),
transferred briefly to a cryoprotection solution containing well solution supplemented with 1 mM
$FeSO_4$ (in 0.1% (v/v) HCl) and 20% (v/v) PEG 200, which were subsequently flash cooled in liquid
nitrogen.
206
All crystallographic datasets were collected at 10–100 eV above the experimentally determined Fe-
$K\alpha$ edge on the macromolecular crystallography beamlines at Diamond Light Source (Didcot, UK)
at 100 K using Pilatus 6M detectors. Diffraction data were integrated and scaled using XDS (Kabsch,
2010) and symmetry related reflections were merged with Aimless (Evans, 2011). Data collection
statistics are shown in Table 3. The resolution cut-off used for structure determination and refinement
was determined based on the $CC_{1/2}$ criterion proposed by Karplus and Diederichs, 2012.
The structures of the EncFtn_sH variants were determined by molecular replacement using decameric
wild-type protein (PDB ID: 5DA5) as the search model (He et al., 2016). A single solution comprising
three decamers in the asymmetric unit was found by molecular replacement using Phaser (McCoy et
al., 2007). The initial model was rebuilt using Phenix.autobuild (Adams et al., 2010) followed by
cycles of refinement with Phenix.refine (Afonine et al., 2012), with manual rebuilding and model
inspection in Coot (Emsley et al., 2010). The final model was refined with isotropic B-factors,
torsional NCS restraints, and with anomalous group refinement to correctly model the iron ions in
the ferroxidase centre. The model was validated using MolProbity (Chen et al., 2010). Structural
superimpositions were calculated using Coot (Emsley et al., 2010). Crystallographic figures were
generated with PyMOL. Model refinement statistics are shown in Table 3. The final models and
experimental data are deposited in the PDB.
224
Ferroxidase assay of EncFtn proteins
Enzymatic activity of EncFtn proteins was tested by ferroxidase assay, as previously described (He
et al., 2019). Fe(II) aliquots were prepared under anaerobic conditions by dissolving FeSO4·7H2O in
0.1% (v/v) HCl. Purified protein was diluted anaerobically in Buffer H (10 mM HEPES, pH 8.0, 150
mM NaCl), previously purged with gaseous nitrogen, to a final concentration of 10 µM (EncFtn).
Protein and ferrous ion samples were added to a quartz cuvette (Hellma) under aerobic conditions at
a final concentration of 10 µM and 50 µM, respectively, corresponding to a FOC:Fe ratio of 1:10.
Absorbance at 315 nm was recorded every second for 1500 s by a UV-visible spectrophotometer
(PerkinElmer Lambda 35), using the TimeDrive software. The same experiment was performed in
the absence of the enzyme to determine the oxidation of ferrous salts by atmospheric oxygen.
The same setup was adopted for recording the activity of EncFtn_E34A and EncFtn_E31A/E34A in
a separate assay at higher concentration (20 µM EncFtn and 100 µM Fe(II)) for comparison purposes.
Data presented here are the mean of three technical replicates of time zero-subtracted progress curves
with standard deviations calculated from the mean.
Calculation of the EncFtn enzymatic reaction initial rate (v_0) was made by applying the Linear
Regression tool in GraphPad (Prism8) on the absorbance at 315 nm measured for the first 10 s or 200
s, when curves are still linear and following an order zero kinetics (Figure 10, Supplementary Tables
S5, S6, S8). Slopes obtained from these curves correspond to initial reaction rate. $v_0^{variant}$/ $v_0^{wild-type}$
factors were calculated by dividing the variants initial rates by the initial rate of the wild-type protein
(Supplementary Tables S7).
245
Ferroxidase assay of Enc:EncFtn protein complexes
Ferroxidase assay of Enc:EncFtn protein complexes was performed in the same conditions as those
used for EncFtn, but with different protein and iron salt concentration. Enc:EncFtn was diluted to a
final concentration of 25 µM which corresponds to 10 µM EncFtn based on a 3:2 Enc:EncFtn ratio,
as previously observed by SDS-PAGE by our group (Figure 1A in He et al., 2016). Fe(II) sample was
diluted to 50 µM in the reaction system to maintain a final ratio of 1:10 FOC:Fe(II).
Calculation of the Enc-EncFtn enzymatic reaction initial rate (v_0) and $v_0^{variant}$/ $v_0^{wild-type}$ factors was
carried out as described in the above section (Figure 10, Supplementary Tables S5, S6, S7, S8).

Zinc inhibition of ferroxidase activity of EncFtn proteins

Ferroxidase assay was performed as above with EncFtn (10 µM) and Fe(II) samples (50 µM) in the presence of 34 µM $ZnSO_4 \cdot 7H_2O$. The chosen concentration corresponds to the IC50 value previously determined for EncFtn_StrepII, under the same experimental conditions, by our group (He et al., 2019). Oxygen-free metal samples were added to the quartz cuvette under aerobic conditions, followed by the protein sample. Data were replicated three times and mean and standard deviation of time zero-subtracted progress curves were calculated. Negative control was performed by monitoring A_{315} of Zn(II) and Fe(II) salts mixed in the absence of enzyme.

Accession codes and datasets

Data sets supporting this paper have been deposited in appropriate public data repositories. Please see figure legends and tables for links to these.

Results

Purification of encapsulated ferritins from R. rubrum

In order to explore the functional roles of the residues in the putative entry site of EncFtn, a number of single-point variants were produced by site-directed mutagenesis, selectively substituting each metal-binding residue in the site with alanine or glycine. The resulting set of variants were also generated in association with the encapsulin component to produce Enc:EncFtn complexes. All the proteins were produced by heterologous expression in *Escherichia coli.*

To investigate the influence of these changes to the proteins in solution, all proteins were subjected to size-exclusion chromatography using an S200 column calibrated with standards of known molecular mass (Figure 3A/B). The resulting fractions were analysed by SDS-PAGE (Figure S1A) and the average molecular masses of the EncFtn proteins were confirmed by LC-MS (Table 1). Each of the variant EncFtn proteins eluted with a similar profile, with a major peak at retention volume around 60 ml, a second peak at retention volume of 75 ml, and the final peak at 82 ml (Figure 3A, Supplementary Table S3). The E34A variant had significantly more protein in the 82 ml peak when compared to the wild type and E31A variant, suggesting that this change affects the stability of the protein complex.

Based on our prior work on the assembly pathway of the *Haliangium ochraceum* EncFtn protein (Ross et al., *submitted*) and native mass spectrometry of other representative members of the EncFtn family (He et al., 2019) these peaks would correspond to decamer, tetramer, and dimer species. The anomalous size estimation based on calibration standards can be attributed to the fact that the EncFtn proteins used in this study were full length proteins with the flexible localisation sequence, which would increase the hydrodynamic radius of the oligomers considerably.

The Enc:EncFtn protein complexes all eluted as single peaks starting in the void volume of the S200 column, indicating the formation of large protein complexes not able to enter the matrix pores (Figure 3B, Supplementary Table S3). SDS-PAGE analysis of peak fractions confirmed the presence of both the EncFtn and encapsulin proteins (Figure S1) To determine if these complexes had formed correctly assembled encapsulin nanocages we performed transmission electron microscopy on uranyl acetate stained samples (Figure 4). The encapsulins purified with wild type and variant EncFtn all showed characteristic 25 nm encapsulin cages with visible internal density for the EncFtn proteins, consistent with what previously observed for Rru_Enc-EncFtn wild-type (He et al., 2016).

Native mass spectrometry of EncFtn proteins

To interrogate the stability and oligomeric states of the EncFtn variants we performed native MS and ion mobility MS experiments on the samples purified by size-exclusion chromatography (Figure 5A) (He et al., 2019). In native MS analysis, EncFtn_E31A behaves similarly to EncFtn and adopts a decameric charge state distribution (Figure 5A, pink circles). In contrast, under the same instrumental

conditions, EncFtn_E34A and EncFtn_W38G display significant differences – with both displaying
decameric and monomeric charge state distributions (Figure 5, stressed with pink circles and blue
triangles, respectively). The presence of monomeric with relatively high charge states (8+ to 16+),
suggests a degree of gas phase dissociation from the decameric complex during native MS analysis.
Even with careful control of instrument conditions, this monomeric dissociation product could not be
completely eliminated, suggesting that the EncFtn_E34A and EncFtn_W38G variants are less stable
than the wild type protein. The difference in gas phase stability of the EncFtn variants was further
highlighted when increasing the sampling cone voltage and/or activating the trap voltage of the mass
spectrometer, which activates the complex and causes dissociation (Figure S2). EncFtn_W38G fully
dissociates into its monomer charge state distribution at lower activation conditions than the other
variants, further highlighting its instability (Figure S2D).
To further investigate the gas phase stability of the variants, ion mobility collision induced unfolding
(CIU) experiments were performed (Eschweiler et al. 2015). The complexes were unfolded by
gradually increasing the trap voltage prior to the gas phase conformation of the native protein ions is
monitored by ion mobility mass spectrometry. The resulting ion mobility profiles for an individual
protein ion is then plotted as a function of activation voltage to produce a heat map to reveal discrete
conformations and gas phase unfolding transitions (Figure 6). It is clear from these experiments, that
two discrete conformations of each EncFtn variant are observed during CIU; a major conformation
with a compact structure and a minor conformation with a more extended, unfolded structure (Figure
6, Figure S3). At minimal activation, EncFtn_WT, EncFtn_E34A and EncFtn_E31A all exist in a
single, compact decameric conformation with a drift time of between 10 and 11 ms, suggesting the
same overall structure. As activation is increased all three variants undergo a discrete transition to a
more extended conformation (characterised by a drift time of 12.5 ms) at around 30 V activation,
followed by a more complex transition to higher drift time conformations with activation above ca.
40V. The close similarity of these CIU profiles strongly suggest have similar gas phase structures and
stability (Figure 6, Figure S3). In contrast, EncFtn_W38G occurs in a more extended conformation
(12.5 ms) throughout the CIU experiment, with the compact conformation only observed as a minor
species at low activations. This is in accord with the native MS studies and suggests that the W38G
substitution has a significant structural effect and destabilises the decameric EncFtn assembly.
335
By calibration of the ion mobility data, the collision cross sections (CCS) of both the discrete
conformations of the EncFtn decamer were determined (Table 2). The compact conformation of
EncFtn_WT has a CCS of 71.40 nm^2; whereas the more extended conformation, observed at higher
activation has a CCS of 80.62 nm^2 (Table 2). We note that the CCs of the compact conformer is
significantly larger than the CCS calculated from the EncFtn_ sH is crystal structure of 58.7 nm^2 and
58.2 nm^2 observed in previously reported IM-MS experiments (He et al., 2016). However, prior
studies were performed on truncated EncFtn_WT which has the C-terminal localisation sequence
region removed. The higher observed CCS for EncFtn_WT suggests that the localisation sequence is
on the surface of the decamer as it has an increased cross-sectional area. This is in accord with our
observations for the size-exclusion experiments, where the EncFtn variants elute with an apparently
higher hydrodynamic radius than the expected oligomeric states.
347
X-ray crystal structures of R. rubrum EncFtn variants
Given the observation E31A and E34A variants do not display any significant differences in terms of
oligomeric state and stability both in solution and the gas phase, we subjected truncated his-tagged
versions of these proteins to crystallisation screening to determine any changes at the metal binding
sites. We obtained crystals in similar conditions to the wild type sequence protein. We were unable
to obtain crystals of the W38G, or W38A variants of the EncFtn protein, indicating that the
conformational differences seen in the mass spectrometry experiments destabilise the quaternary
structure of the protein sufficiently to present a barrier to stable crystallisation.
356

The structures of the *R. rubrum* EncFtn_sH E31A and E34A variants were determined by X-ray crystallography and confirmed to adopt the same overall quaternary fold as the wild type sequence EncFtn, with RMSD Cα differences of 0.2 A^2 over 91 Cα and 0.17 Å^2 over 90 Cα for the E31A and E34A variants respectively, and an RMSD Ca of 0.23 Å^2 over 90 Cα between the two variants. This indicates that these amino acid substitutions do not significantly alter the structure of the protein in the E31A and E34A variants.

Inspection of anomalous difference maps calculated for the X-ray data collected near to the Fe-*Kα* edge indicate the presence of anomalously scattering metal ions in the ferroxidase centre for both the E31A and E34A variants. We have assigned these as iron ions based on the size of the peaks and the nature of this site (Figure 7A/B, 8A/B). The ferroxidase centre of the E31A variant has an additional peak of electron density assigned as a water molecule, bridging the di-iron centre.

The entry site of the E31A variant appears to coordinate a metal ion via the Glu34 side-chain, inspection of anomalous maps allowed us to assign this as a calcium ion; however, the coordination distances are much longer than the wild type, at 3.8 Å compared to 2.8 Å (Figure 7C.). Loss of the coordinating Glu31 side chain ligand does not appear to cause gross structural rearrangements in this site, and the change in metal ion binding can be ascribed to the change in the coordination environment around this site.

The entry site of the E34A variant does not appear to coordinate any metal ions and the side chain of Glu31 is flipped away from this site towards Arg42 (Figure 8C), the Trp38 indole ring is also moved away from the site when compared to the wild type protein. These changes greatly disrupt the metal coordination in this site.

Ferroxidase activity of EncFtn and Enc:EncFtn proteins
While the MS and crystallographic data were in agreement with respect to the quaternary structure of the E31A and E34A variants, our inability to produce crystals of the W38G protein and reduced stability in our MS experiments highlight the importance of this residue for the oligomerisation of the EncFtn decamer. The changes in metal ion coordination seen in the E31A and E34A crystal structures suggest these may play a role in controlling entry of iron ions to the ferroxidase centre. To test this hypothesis, we investigated the activity of all of the variants produced, both in isolation, and in complex with the encapsulin protein. The rationale behind producing single point mutants was to discern between the different actions that the E31-E34-W38 residues could exert on the enzyme activity. At the start of our work we delineated a number of hypothesis involving this site, including it being an access gate at the entry of the channel, and providing attractive forces which, as a consequence, guide the metal ions to the FOC, buried at the dimer interface.

The ferroxidase activity of the wild-type EncFtn protein was confirmed to be comparable to that previously determined for the EncFtn_StrepII variant (He et al., 2019). Both the W38A and E31A mutations clearly affect the activity of the protein, as oxidation of Fe(II) to Fe(III) occurs faster and both curves present a distinct shape to the wild-type protein (Figure 9). The EncFtn_E34A variant shows a smaller increase in catalytic activity when compared to the other variants (Figure 9).

Considering these differences in activity in the context of the X-ray crystal structures, it is clear that disruption of the metal coordination in this entry site has a significant impact on the catalytic activity of the EncFtn protein. The EncFtn_E31A structure shows that the gating provided by this mutant is partially diminished, as is its binding ability. However, the residual glutamic acid, Glu34, is still able to coordinate metal ions, albeit less tightly than the wild type given the increased distance between these residues and lack of additional coordination from Glu31 (Figure 7). This variant possesses the highest ferroxidase activity, along with EncFtn_W38A.

The crystal structure of the EncFtn_E34A variant shows that the mutation of Glu34 into Ala results
in a reorganisation of the site, with the side chain of Glu3 flipped by 117°, and is no longer correctly
positioned for metal coordination, essentially removing both the electrostatic attraction and gating of
this site (Figure 8). A ferroxidase assay carried out with an EncFtn_E31A/E34Avariant shows that
its activity is comparable to EncFtn_E34A (Figure 11). This data corroborates our hypothesis that in
EncFtn_E34A, the gating function is completely abolished, and no attraction is conveyed by the
residual Glu31. Hence, in Rru_EncFtn_E34A metal ions slowly reach the FOC by diffusion only. As
no gating is provided at this level, we would expect to observe a higher activity, however the absence
of attracting forces results in modest increase when compared to wild type. The difference in activity
between EncFtn_E31A and EncFtn_E34A shows how critical Glu34 is and more broadly the
importance of this electrostatic attraction for the activity of the protein.

As we were unable to produce crystals of either the W38G or W38A variants, we could not determine
the effect of removing this tryptophan residue on the structure of the entry site. However, analysis of
the crystal structure of the wild type EncFtn shows that the tryptophan indole ring appears to sterically
constrain the Glu31 side chain in the correct position for metal-coordination in WT (Figure 1, He et
al., 2016), we propose that the removal of Trp38 results in the Glu31 side chain moving away from
the site. Our ferroxidase assay data presents this variant as having a comparable activity profile to the
E31A variant, we therefore propose that the two variants possess similar entry site structures. The
broader destabilising effects of the loss of the tryptophan side chain are harder to model, but the loss
of a side chain of the size of the tryptophan indole may lead to further structural rearrangements
around the internal dimer interfaces engaged by this residue.

The same ferroxidase assay was performed with the Enc:EncFtn complex, with wild-type and variants
carrying the same mutations introduced in Rru_EncFtn protein (Figure 9). These deviations from
wild-type behaviour were compared to what observed in the absence of the encapsulin
nanocompartment (Figure 9A). Enzymatic reaction initial rates (v_0) were calculated for each of the
variants (in the first 200 s of the assay) and then divided by the initial rate of the corresponding wild-
type protein (EncFtn or Enc:EncFtn) (Supplementary Table S7). This allowed comparison across the
two experimental setups (EncFtn versus Enc:EncFtn proteins). Data show that E31A and W38X
(where X is Gly or Ala) mutations have a greater effect on activity within both systems (about 4 folds
faster when compared to wild-type initial rate) while removal of Glu34 results in a 1.5-2-folds
increase in the initial velocity v_0 (Figure 10, Supplementary Table S7). It appears that the EncFtn
variants and wild-type retain the same relative differences in their enzymatic activities, whether they
are associated to the encapsulin shell or not. This outcome suggests that the gating and the attraction
properties are actually exerted by the entry site instead of the pore level.

However, if we examine Enc:EncFtn_E31A curve in more details, we notice that it is not completely
linear in the first 200 s of reaction (Figure 10B). Hence, we chose to directly compare the initial
velocity (v_0) of EncFtn_E31A and Enc:EncFtn_E31A within this shorter period of time (Figure S5).
Although EncFtn_E31A data are far too noisy to allow an acceptable fit of the curve resulting from
the Linear Regression tool (Supplementary Table S8), a qualitative comparison can still be made
(Figure S5). When encapsulated, EncFtn_E31A appears to catalyse the ferroxidase reaction much
faster than other variants, including EncFtn_E31A when not compartmentalised within A0974
(Supplementary Table S8). After the first 20-30 s the reaction starts to slow down, reaching a rate
comparable to Enc:EncFtn_W38G. It is plausible to believe that the nanocompartment improves the
stabilisation of the inner protein, resulting in a different reaction profile.

Zinc inhibition of ferroxidase activity of Rru_EncFtn
In order to test the hypothesis that the entry site acts as a selectivity filter, and is able to discriminate
between metal ions approaching the FOC, ferroxidase assays were performed with the addition of
zinc as a competitive inhibitor of ferroxidase activity. The activity of EncFtn wild-type and variant

proteins (10 μM, monomer) with 50 μM Fe(II) was assayed in the presence of 34 μM Zn(II) (Figure
12A). The selected zinc concentration was previously determined to be the IC50 for wild type
EncFtn_StrepII (He et al., 2019). This concentration was chosen to allow identification of the impact
of inhibition across the set of proteins. In line with our previous results the EncFtn wild type sequence
protein showed a 54% inhibition (He et al., 2019) (Figures 12A/B). The activity of the three variant
proteins (E31, E34, W38) showed a markedly increased relative inhibition when compared to their
enhanced activity in the absence of zinc (66%, 68%, and 88%, respectively) (Figures 12A/B).

The increased inhibition seen in the variants can be explained by the changes in metal coordination
at this site. With all the variants, the change in metal coordination at the putative entry site permits
greater access to the FOC for competing metal ions, as they are no longer effectively captured in the
entry site. Of note is the fact that in the presence of zinc, the activity of each variant is the same as
the wild type protein.

We have shown that substitution of residues in the entry site triad (E31, E34, W38) with non-metal
binding residues results in the enhancement of enzymatic activity (Figure 9A). Our crystal structures
support the conclusion that this is a consequence of the creation of a wider and more loosely
coordinating site, which allows the faster passage of metal ions to the FOC due to a reduced energy
barrier at this site (Figure 7, Figure 8). Furthermore, when these key residues are removed the protein
is more susceptible to inhibition by Zn(II). As wild type EncFtn presents an extra metal binding site
with which competing metals would interact, this presents a barrier to the FOC, which would trap
metal ligands before they are able to reach the FOC itself. In the absence of this site both iron and
competing metals can approach the FOC, either productively in the case of iron, or unproductively in
the case of zinc.

Discussion
Our work presents a comprehensive structure/function study investigating the role of the putative
metal ion entry site proximal to the ferroxidase centre FOC in encapsulated ferritin-like proteins. To
investigate the role of this site we produced single point mutations in residues in this site to replace
metal coordinating residues in the *R. rubrum* EncFtn protein with alanine (E31A, E34A) and to
remove the large tryptophan side chain proximal to this site (W38G/A).

Solution and gas-phase analyses of variant proteins showed that removal of the metal coordinating
ligands did not have a significant effect on protein oligomerisation, however the removal of the Trp38
side chain destabilised the protein considerably. X-ray crystal structures of the E31A and E34A
variants highlighted changes in metal coordination at this site.

The effect of these changes on the activity of the protein was tested using ferroxidase assays and zinc
inhibition experiments. Removal of the Glu31 and Trp38 side chains significantly enhanced the
activity of the enzyme. While removal of metal coordinating residues increased the susceptibility of
the protein to inhibition with zinc.

Taken together these results support the hypothesis that this metal ion binding site has a dual function
in modulating the rate of ferroxidase activity of the EncFtn protein and protecting the ferroxidase
centre from competing metal ions by acting as an electrostatic trap prior to the active site. These
results also confirm that the ferroxidase centre is indeed the active site of this enzyme, as the loss of
the metal ion entry site enhances enzyme activity rather than abrogates it. Why the EncFtn protein
has evolved a mechanism to effectively slow its activity is still an open question. There may be a
trade-off between enzyme activity and the availability of a sink for the electrons liberated from the
iron in the ferroxidase centre. The catalytic mechanism and electron sink for ferritins is still a subject
of vigorous debate and the EncFtn family presents a unique structural arrangement of the active site,
which will no doubt present a distinct mechanism when probed further. The mineralisation step is

also likely to be a rate limiting step and the entry site may provide a brake to ensure productive
coupling of iron oxidation and mineralisation.
515
Encapsulation of the EncFtn protein within the encapsulin nanocage enhances the activity of all of
the EncFtn variants but does not change the way in which the variants affect the activity, indicating
that the function of the entry site is not changed through association with the encapsulin shell. These
results suggest that the encapsulin shell acts to guide productive iron ion interactions with the EncFtn
protein, indeed, our previous work showed that the encapsulin shell interacts with a significant
amount of iron in the absence of the EncFtn protein. This warrants future investigation into the ability
of the encapsulin shell to conduct metal ions to the EncFtn active site and the role of the encapsulin
shell pores in the access of substrates to the encapsulated enzyme.

532
533
534 **Figure 1. Putative iron ion entry site in *Rhodospirillum rubrum* EncFtn.**
535 Coordination of calcium (yellow sphere) within the dimer interface by Glu31 (purple) and Glu34
536 (pink) from the two different subunits. Iron ions coordinated in the ferroxidase centre (FOC) are
537 depicted as red spheres. Trp38 side chains (orange) from both chains are also shown.
538

540
541
Figure 2. Illustration of the hypotheses for the functional role of the putative metal ion entry
site in *Rhodospirillum rubrum* EncFtn.
A, B, Schematic view of a putative pathway for incoming ferrous ions moving towards the active
site (FOC). EncFtn is represented by a grey oval, where the putative entry sites and ferroxidase
centres are shown in red and yellow, respectively. A portion of the encapsulin is displayed in pale
blue, with emphasis given to the 5-fold pore used by ferrous ions to reach the compartmentalised
ferritin protein. Ferrous ions (blue circles) are shown to be converted into ferric ions (yellow
circles) in the ferroxidase centre with the concomitant reduction of molecular oxygen (*B*). Several
hypotheses (gating (*C*), selectivity filter (*D*), and attraction (*E*)) on the role played by the residues
contributing to the putative entry site are shown in *C-E*. Ferrous ions are depicted as blue circles,
while non-cognate species are in dark green.
553

Figure 3. Purification of recombinant *R. rubrum* EncFtn and Enc:EncFtn protein complexes.
Recombinant EncFtn (***A***) and Enc:EncFtn (***B***) proteins were purified by anion exchange column
(Hi-Trap Q-sepharose FF, GE Healthcare) and then subjected to analytical size-exclusion
chromatography using a Superdex 200 16/60 column (GE Healthcare) previously equilibrated with
50 mM Tris-HCl, pH 8.0, 150 mM NaCl. Investigation of the elution profiles in Figure 3A reveal
that all EncFtn proteins show the main peak at around 60 ml, diagnostic of oligomerisation states
close to 20-mer, and smaller peaks at ~ 76 ml and 82 ml (indicative of smaller assembly states,
Supplementary Table S3). Enc:EncFtn proteins elute in a single peak around 46 ml, suggesting that
EncFtn is compartmentalised within the Enc shell. doi. 10.6084/m9.figshare.9885557

564
565

566
567
568
Figure 4. Transmission electron micrographs of *Rhodospirillum rubrum* Enc:EncFtn
complexes.
Negative stain TEM images of *R. rubrum* Enc:EncFtn nanocompartments stained with uranyl
acetate. Enc:EncFtn_WT (***A***), Enc:EncFtn_E34A (***B***), Enc:EncFtn_W38G (***C***), and
Enc:EncFtn_E31A (***D***). Samples were imaged at a magnification of 100,000, scale bars correspond
to 50 nm. Proteins were diluted in Buffer GF to 0.1 mg/ml.
575

Figure 5. Native MS Analysis of *Rhodospirillum rubrum* EncFtn variants.
All spectra were acquired in 100 mM ammonium acetate, pH 8.0. nESI spectra of EncFtn variants
displaying decamer (pink circles) and monomer (blue triangle) charge state distributions.

586
587
Figure 6. Collision Induced unfolding (CIU) of the *Rhodospirillum rubrum* EncFtn variants determined by ion mobility mass spectrometry.
The ion mobility drift time of the 26+ charge state of each EncFtn variant is displayed as a function of increasing collision voltage. All variants display similar CIU profiles, with a compact structural conformation (10.5 ms) present at low activation which transitions to a more extended conformation (drift time 12.5-13.5 ms) as activation increases. The onset of this transition occurs at around 32 V in WT, E31A, and E34A; However, the transition occurs at significantly lower activation energy in W38G and the extended structural conformation (12.5 ms) dominates the profile, even at low activation voltages.
597
598

599
Figure 7. Comparison of metal binding sites in EncFtn_sH wild type and E31A structures.
Metal binding sites in the EncFtn E31A variant and WT are shown as stick representations with Cα
coloured in marine and white, respectively. The final experimental 2mFo-DFc density map is
shown contoured at 1.5 σ around the metal binding sites of the E31A variant as a blue mesh. An
NCS-averaged anomalous difference map is superimposed on the FOC of E31A variant as an
orange mesh and contoured at 20 σ. Residues from dimers are differentiated with a prime symbol in
their labels. The distance between metal ions and coordinating residues is marked by dashed lines
with distances shown in Å. *A*, Aligned FOCs with di-iron bound as orange spheres in E31A mutant
and red spheres in WT. A water molecule shown as a red cross is coordinated above the di-iron of
E31A mutant. *B*, Top view of *A* with residues labelled. *C*, Metal ion entry site comparison between
E31A mutant and WT with both Ca(II) bound in grey and brown spheres, respectively. In the E31A
mutant the Ca(II) moves further from FOC and is coordinated by E34 from two chains in a distance
of 3.4 Å/3.8 Å by contrast to the 2.8 Å/2.9 Å in the WT.
613
614

615
Figure 8. Comparison of metal binding sites in the E34A and WT *Rhodospirillum rubrum*
EncFtn_sH structures.
Metal binding sites in the EncFtn E34A mutant and WT are shown as stick representations with Cα
coloured in forest and white, respectively. The final calculated 2mFo-DFc density map is shown
contoured at 1.5 σ around the metal binding sites of the E34A variant as a blue mesh. An NCS-
averaged anomalous map is superimposed on FOC of E34A mutant as an orange mesh and
contoured at 20 σ. Residues from dimers are differentiated with a prime symbol in their labels. The
distance between metal ions and coordinating residues is marked by dashed lines and labelled with
numbers nearby and Å as the unit. *A*, Aligned FOCs with di-iron ions bound as orange spheres in
E34A variant and red spheres in WT. *B*, Top view of *A* with residues labelled. *C*, Metal ion entry
site comparison between E34A variant and WT. No metal is coordinated in this site of E34A mutant
while WT binds a Ca(II) shown in grey sphere. The E31 in E34A mutant moves away by 117° and
would be 6 Å away from the position of the Ca(II) of the WT. The indole ring of W38 rotates
towards the central Ca(II) site of the WT by 38°.
630

631
Figure 9. Ferroxidase activity of *Rhodospirillum rubrum* EncFtn and Enc:EncFtn complexes.
A, EncFtn wild-type (green line), EncFtn_E34A (pink line), EncFtn_E31A (blue line), and
EncFtn_W38A (orange line), (10 μM, monomer) were incubated with 50 μM FeSO$_4$.7H$_2$O (10
times molar equivalent Fe(II) per FOC) and progress curves of the oxidation of Fe(II) to Fe(III) was
monitored at 315 nm at room temperature. The background oxidation of iron at 50 μM in enzyme-
free control is shown for reference (dotted grey line). Solid lines represent the average (n = 3) of
technical replicates, shaded areas represent standard deviation from the mean. Protein and iron
samples were prepared anaerobically in Buffer H (10 mM HEPES pH 8.0, 150 mM NaCl), and 0.1
% (v/v) HCl, respectively. *B*, Enc:EncFtn wild-type (dark green line), Enc:EncFtn_E34A (dark pink
line), Enc:EncFtn_E31A (dark blue line), and Enc:EncFtn_W38G (chestnut line) (25 μM
Enc:EncFtn, corresponding to 10 μM EncFtn monomer) were incubated with 50 μM FeSO$_4$.7H$_2$O
and progress curves of the oxidation of Fe(II) to Fe(III) was monitored at 315 nm at room
temperature. The background oxidation of iron at 50 μM in enzyme-free control is shown for
reference (dotted grey line). Solid lines represent the average (n = 3) of technical replicates, shaded
areas represent standard deviation from the mean. Protein and iron samples were prepared
anaerobically in Buffer H (10 mM HEPES pH 8.0, 150 mM NaCl), and 0.1 % (v/v) HCl,
respectively. doi.10.6084/m9.figshare.9885575
649
650

Figure 10. Linear Regression on first 200s of ferroxidase assays with EncFtn and Enc:EncFtn complexes.

A, Data from first 200 s of the assays shown in Figure 9*A* were processed using the Linear Regression tool in GraphPad (Prism8) to determine v_0, initial enzymatic rate, from slope of each of the linear curves (see Table S5). *B*, Data from first 200 s of the assays shown in Figure 9*B* were processed using the Linear Regression tool in GraphPad (Prism8) to determine v_0, initial enzymatic rate, from slope of each of the linear curves (see Table S6). *C*, Variants initial enzymatic rate ($v_0^{variant}$) calculated from linear regression curves in the first 200 s of ferroxidase assay experiments were divided by the slope calculated for the corresponding wild-type species ($v_0^{wild-type}$). Evaluation of the resulting $v_0^{variant}/v_0^{wild-type}$ factors allow comparison between enzymatic kinetics of EncFtn (solid coloured columns) and Enc:EncFtn (striped columns) proteins. Colour coding is consistent with Figures 9*A,B* and Figures 10*A,B* and columns are labelled with corresponding mutation. Horizontal green line represents wild type's $v_0^{variant}/v_0^{wild-type}$ ratio = 1. Representations of EncFtn or Enc:EncFtn systems (non-descriptive of true size ratios) are shown above each column for clarity. doi.10.6084/m9.figshare.9885575

Figure 11. Comparison of ferroxidase activities of *Rhodospirillum rubrum*_EncFtn_E34A and *Rhodospirillum rubrum* EncFtn_E31A/E34A.

EncFtn_E34A (pink line) and EncFtn-E31A/E34A (dark purple line), (20 μM, monomer) were incubated with 100 μM FeSO$_4$.7H$_2$O (10 times molar equivalent Fe(II) per FOC) and progress curves of the oxidation of Fe(II) to Fe(III) was monitored at 315 nm at room temperature. The background oxidation of iron at 100 μM in enzyme-free control is shown for reference (dotted grey line). Solid lines represent the average (n = 3) of technical replicates, shaded areas represent standard deviation from the mean. Protein and iron samples were prepared anaerobically in Buffer H (10 mM HEPES pH 8.0, 150 mM NaCl), and 0.1 % (v/v) HCl, respectively.

doi.10.6084/m9.figshare.9885575

Table 1. Average molecular masses of encapsulated ferritins obtained by LC-MS.

Protein	Observed Mass (Da)	Assignment	Theoretical Mass (Da)
Rru_EncFtn_WT	15060.8 Da	monomer without Met	15061.4 Da
		dimer without Met	
Rru_EncFtn_E31A	15003.6 Da	monomer without Met	15003.4 Da
Rru_EncFtn_E34A	15003.3 Da	monomer without Met	15003.4 Da
Rru_EncFtn_W38G	14932.3 Da	monomer without Met	14932.2 Da

The starting Methionine residue is not always retained, and this has been indicated in "Assignment".

Table 2. Collision cross sections (CCS) of the compact and extended conformations of the Rru-EncFtn variants

Protein	Compact CCS (nm^2)	Extended CCS (nm^2)
Rru_EncFtn_ WT	71.40	80.62
Rru_EncFtn_E31A	69.50	82.65
Rru_EncFtn_E34A	71.30	81.68
Rru_EncFtn_W38G	71.17	81.41

CCS values were calculated using Driftscope v2.5.

709 **Table 3. Data collection and refinement statistics.**

	*E31A**	*E34A**
Data collection		
Wavelength	1.74	1.74
Resolution range	49.40 - 2.66 (2.75 - 2.66)	48.73 - 2.19 (2.27 - 2.19)
Space group	P 1 2_1 1	P 1 2_1 1
Unit cell (Å) a b c	97.89 120.16 139.37	97.79 120.14 139.34
β (°)	95.33	95.20
Total reflections	310,119 (30,187)	906,663 (86,277)
Unique reflections	170,103 (16,471)	31,4834 (30,221)
Multiplicity	1.8 (1.8)	2.9 (2.9)
Completeness (%)	93.30 (90.35)	96.37 (92.66)
Mean I/sigma(I)	5.45 (1.40)	8.04 (1.43)
Wilson B-factor ($Å^2$)	39.14	33.05
R_{merge}	0.136 (0.657)	0.094 (0.688)
R_{meas}	0.184 (0.878)	0.116 (0.842)
R_{pim}	0.123 (0.578)	0.065 (0.478)
$CC_{1/2}$	0.978 (0.517)	0.995 (0.579)
CC*	0.994 (0.826)	0.999 (0.857)
Refinement		
Reflections used in refinement	170,031 (16,463)	314,739 (30,212)
Reflections used for R_{free}	8,346 (794)	15,590 (1548)
R_{work}	0.195 (0.297)	0.184 (0.282)
R_{free}	0.236 (0.327)	0.212 (0.315)
CC_{work}	0.946 (0.748)	0.959 (0.768)
CC_{free}	0.914 (0.684)	0.952 (0.689)
Model		
Number of non-hydrogen atoms	22,382	22,760
macromolecules	21,988	22,072
ligands	65	48
solvent	329	640
Protein residues	2680	2696
RMS(bonds) (Å)	0.007	0.005
RMS(angles) (°)	1.03	0.87
Ramachandran		
favoured (%)	99.69	99.77
allowed (%)	0.31	0.23
outliers (%)	0.00	0.00
Rotamer outliers (%)	0.26	1.07
Clashscore	4.79	3.01
Average B-factor ($Å^2$)	41.64	40.53
macromolecules	41.66	40.52
ligands	55.57	46.17
solvent	37.08	40.62
PDB ID	6SUW	6SV1

710 Statistics for the highest-resolution shell are shown in parentheses. *Friedel pairs are treated as
711 separate reflections for data analysis and refinement.
712

References

Adams PD, Afonine PV, Bunkóczi G, Chen VB, Davis IW, Echols N, Headd JJ, Hung L-W, Kapral GJ, Grosse-Kunstleve RW, McCoy AJ, Moriarty NW, Oeffner R, Read RJ, Richardson DC, Richardson JS, Terwilliger TC, Zwart PH. 2010. *PHENIX*: a comprehensive Python-based system for macromolecular structure solution. *Acta Cryst*. **D66**:213-221. doi:10.1107/S0907444909052925

Afonine PV, Grosse-Kunstleve RW, Echols N, Headd JJ, Moriarty NW, Mustyakimov M, Terwilliger TC, Urzhumtsev a, Zwart PH, Adams PD. 2012. Towards automated crystallographic structure refinement with phenix.refine. *Acta Cryst*. **D68(4)**:352-67. doi: 10.1107/S0907444912001308

Almirón M, Link AJ, Furlong D, Kolter R. 1992. A novel DNA-binding protein with regulatory and protective roles in starved Escherichia coli. *Genes Dev* **6**:2646–2654. doi:10.1101/gad.6.12b.2646

Andrews SC. 1998. Iron storage in bacteria. *Adv Microb Physiol* **40**:281-351

Andrews SC. 2010. The ferritin-like superfamily:evolution of the biological iron storeman from a rubrerythrin like ancestor. *Biochim Biophys Acta – General Subjects* **1800**:691-705. https://doi.org/10.1016/j.bbagen.2010.05.010

Baaghil S, Lewin A, Moore GR, Le Brun NE. 2003. Core formation in Escherichia coli bacterioferritin requires a functional ferroxidase center. *Biochem* **42(47)**:14047-56. doi:10.1021/bi035253u

Behera RK, Torres R, Tosha T, Bradley JM, Goulding CW, Theil EC. 2015. Fe(II) substrate transport through ferritin protein cage ion channels influences enzyme activity and biomineralisation. *J Biol Inorg Chem*. **20(6)**:957-969. doi:10.1007/s00775-015-1279-x

Bellapadrona G, Stefanini S, Zamparelli C, Theil EC, Chiancone E. 2009. Iron translocation into and out of Listeria innocua Dps and size distribution of the protein-enclosed nanomineral are modulated by the electrostatic gradient at the 3-fold "ferritin-like" pores. *J Biol Chem*. **284(28)**:19101-9. doi: 10.1074/jbc.M109.014670

Cardenas JP, Quatrini R, Holmes DS. 2016 Aerobic lineage of the oxidative stress response protein rubrerythrin emerged in an ancient microaerobic, (hyper)thermophilic environment. *Front Microbiol* **7**:1822 doi:10.3389/fmicb.2016.01822

Chasteen ND, Harrison PM. 1999. Mineralization in ferritin: an efficient means of iron storage. *J Struct Biol*. **126**:182-194. doi:10.1006/jsbi.1999.4118

Chen VB, Arendall WB, Headd JJ, Keedy DA, Immormino RM, Kapral GJ, Murray LW, Richardson JS, Richardson DC. 2010. *MolProbity*: all-atom structure validation for macromolecular crystallography. *Acta Cryst*. **D66(1)**:12-21 doi: 10.1107/S0907444909042073

Chiancone E, Ceci P. 2010. The multifaceted capacity of Dps proteins to combat bacterial stress conditions: Detoxification of iron and hydrogen peroxide and DNA binding. *Biochim Biophys Acta*. **1800(8)**:798-805. doi: 10.1016/j.bbagen.2010.01.013

Choi SH, Baumler DJ, Kaspar CW. 2000. Contribution of *dps* to acid stress tolerance and oxidative stress tolerance in Escherichia coli O157:H7. *Appl Environ Microbiol* **66**:3911–3916. doi:10.1128/AEM.66.9.3911-3916.2000

Crow A, Lawson TL, Lewin A, Moore GR, Le Brun NE. 2009. Structural basis for iron mineralization by bacterioferritin. *J Am Chem Soc* **131**:6808–6813 doi: 10.1021/ja8093444

De Martino M, Ershov D, van den Berg PJ, Tans SJ, Meyer AS. 2016. Single-cell analysis of the Dps Response to oxidative stress. *J Bacteriol* **198(11)**:1662-1674. doi: 10.1128/JB.00239-16.

Dillard BD, Demick JM, Adams MW, Lanzilotta WN. 2011. A cryo-crystallographic time course for peroxide reduction by rubrerythrin from Pyrococcus furiosus. *J Biol Inorg Chem.* **16(6)**:949-59. doi:10.1007/s00775-011-0795-6

Dixit SM, Polasky DA, Ruotolo BT. 2018. Collision induced unfolding of isolated proteins in the gas phase: past, present, and future. *Curr Opin Chem Biol.* **42**:93-100. doi: 10.1016/j.cbpa.2017.11.010

Ebrahimi HK, Hagedoorn PL, Jongejan JA, Hagen WR. 2009. Catalysis of iron core formation in Pyrococcus furiosus ferritin. *J Biol Inorg Chem* **14(8)**:1265-74 doi:10.1007/s00775-009-0571-z

Ebrahimi KH, Bill E, Hagedoorn P-L, Hagen W. 2016. Spectroscopic evidence for the role of a site of the di-iron catalytic center of ferritins in tuning the kinetics of Fe(II) oxidation. *Mol BioSyst.* **12**:3576-3588. doi: 10.1039/C6MB00235H

Emsley P, Lohkamp B, Scott WG, Cowtan K. 2010. Features and development of Coot. *Acta Cryst* **D66**:486-501. doi:10.1107/S0907444910007493

Evans PR. 2011. An introduction to data resolution: space-group determination, scaling and intensity statistics. *Acta Cryst* **D76(4)**:282-92. doi:10.1107/S090744491003982X

Frolow F, Kalb AJ, Yariv J. 1994. Structure of a unique twofold symmetric haem-binding site. *Nat Struct Biol* 453-460.

Giessen TW, Orlando BJ, Verdegaal AA, Chambers MG, Gardener J, Bell DC, Birrane G, Liao M, Silver PA, 2019. Large protein organelles form a new iron sequestration system with high storage capacity. *eLife.* **8**:e46070 doi: 10.7554/eLife.46070

Grant RA, Filman DJ, Finkel SE, Kolter R, Hogle JM. 1998. The crystal structure of Dps, a ferritin homolog that binds and protects DNA. *Nat Struct Biol.* **5**:294-303.

Hagen WR, Hagedoorn P-L, Ebrahimi KH. 2017. The working of ferritin: a crossroad of options. *Metallomics.* **9**:595-605.

Harrison PM, Arosio P. 1996. The ferritins: molecular properties, iron storage function and cellular regulation. *Biochim Biophys Acta* **1275**:161-203.

He D and Marles-Wright J. 2015. Ferritin family proteins and their use in biotechnology. *New Biotech* **32**:651-657 doi:org/10.1016/j.nbt.2014.12.006

He D, Hughes S, Vanden-Hehir A, Georgiev K, Altenbach K, Tarrant E, Mackay CL, Waldron KJ, Clarke DJ, Marles-Wright J. 2016. Structural characterisation of encapsulated ferritin provides insight into iron storage in bacterial nanocompartments. *eLife* **5**. doi:10.7554/eLife.18972

816 He D, Piergentili C, Ross J, Tarrant E, Tuck LR, Mackay CL, McIver Z, Waldron KJ, Clarke DJ,
817 Marles-Wright J. 2019. Conservation of the structural and functional architecture of encapsulated
818 ferritins in bacteria and archaea. *Biochem J* **476(6)**:975-989. doi:10.1042/BCJ20180922
819
820 Kabsch W. 2010. Integration, scaling, space-group assignment and post-refinement. *Acta Cryst.*
821 **D66(2)**:133-44. doi:10.1107/S0907444909047374
822
823 Karlplus PA, Diederichs K. 2012. Linking crystallographic model and data quality. *Science*
824 **336(6084)**:1030-33. doi:10.1126/science.1218231
825
826 Liu XS, Patterson LD, Miller MJ, Theil EC. 2007. Peptides selected for
827 the protein nanocage pores change the rate of iron recovery from the ferritin mineral. *J Biol*
828 *Chem.* **282(44)**:31821-5. doi:10.1074/jbc.C700153200
829
830 Martinez A, Kolter R. 1997. Protection of DNA during oxidative stress by the nonspecific DNA-
831 binding protein Dps. *J Bacteriol.* **179**:5188-5194. doi:10.1128/jb.179.16.5188-5194.1997
832
833 Masuda T, Goto F, Yoshihara T, Mikami B. 2010. Crystal structure of plant ferritin reveals a
834 novel metal binding site that functions as a transit site for metal transfer in ferritin. *J Biol Chem*
835 **285**:4049-4059. doi:10.1074/jbc.M109.059790
836
837 McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MS, Storoni LC, Read RJ. 2007. *Phaser*
838 crystallographic software. *J Appl Cryst.* **40**:658-674. doi:10.1107/S0021889807021206
839
840 Nair S, Finkel SE. 2004. Dps protects cells against multiple stresses during stationary phase. *J*
841 *Bacteriol* **186**:4192–4198. doi:10.1128/JB.186.13.4192-4198.2004
842
843 Pesek J, Büchler R, Albrecht R, Boland W, Zeth K. 2011. Structure and mechanism of iron
844 translocation by a Dps protein from Microbacterium arborescens. *J Biol Inorg Chem.* **286**:34872-
845 34882. doi:10.1074/jbc.M111.246108
846
847 Pfaffen S, Abdulqadir R, Le Brun NE, Murphy MEP. 2013. Mechanism of ferrous iron binding and
848 oxidation by ferritin from a pennate diatom. *J Biol Inorg Chem*. **288**:14917-14925.
849 doi:0.1074/jbc.M113.454496
850
851 Proulx-Curry PM, Chasteen ND. 1995. Molecular aspects of iron uptake and storage in ferritin.
852 *Coord Chem Rev* **144**:347–368. doi:10.1016/0010-8545(95)01148-I
853
854 Ratnayake DB, Wai SN, Shi Y, Amako K, Nakayama H, Nakayama K. 2000. Ferritin from the
855 obligate anaerobe *Porphyromonas gingivalis*: purification, gene cloning and mutant studies.
856 *Microbiol* **146**:1119-1127. doi:10.1099/00221287-146-5-1119
857
858 Ross J, Lambert T, Piergentili C, He D, Waldron KJ, Mackay CL, Marles-Wright J, Clarke DJ.
859 2019. Mass spectrometry reveals the assembly pathway of encapsulated ferritins and highlights a
860 dynamic ferroxidase interface. *Chem Comm* (submitted for review) doi xxxx
861
862 Schägger H. 2006. Tricine-SDS-PAGE. *Nat Protoc*. **1(1)**:16-22. doi:10.1038/nprot.2006.4
863
864 Sztukowska M, Bugno M, Potempa J, Travis J, Kurtz DM Jr. 2002. Role of rubrerythrin in
865 the oxidative stress response of Porphyromonas gingivalis. *Mol Microbiol*. **44(2)**:479-88.
866 doi:10.1046/j.1365-2958.2002.02892.x
867

Theil EC, Behera RK, Tosha T. 2013. Ferritins for chemistry and for life. *Coord Chem Rev*.
257(2):579-586. doi:10.1016/j.ccr.2012.05.013

Treffry A, Harrison PM. 1978. Incorporation and release of inorganic phosphate in horse spleen
ferritin. *Biochem J*. **171**:313–320.

Treffry A, Zhao Z, Quail MA, Guest JR, Harrison PM. 1998. How the presence of three iron
binding sites affects the iron storage function of the ferritin (EcFtnA) of *Escherichia coli*. *FEBS*
432:1873-3468

Xu B, Chasteen ND. 1991. Iron oxidation chemistry in ferritin. Increasing Fe/O2 stoichiometry
during core formation. *J Biol Chem.* **266(30)**:19965-70.

Yang X, Chen-Barrett Y, Arosio P, Chasteen ND. 1998. Reaction paths of iron oxidation and
hydrolysis in horse spleen and recombinant human ferritins. *Biochem* **37(27)**:9743-50.

Yao H, Wang Y, Lovell S, Kumar R, Ruvinsky AM, Battaile KP, et al. 2012. The structure of the
BfrB-Bfd complex reveals protein-protein interactions enabling iron release from bacterioferritin.
J Am Chem Soc. **134**:13470-13481. doi:10.1021/ja305180n

Zhao G, Bou-Abdallah F, Arosio P, Levi S, Janus-Chandler C, Chasteen ND. 2003. Multiple
pathways for mineral core formation in mammalian apoferritin. The role of hydrogen peroxide.
Biochem **42(10)**:3142-50. doi:10.1021/bi027357v

Zheng H, Cooper DR, Porebski PJ, Shabalin IG, Handing KB, Minor W. 2017. CheckMyMetal: a
macromolecular metal-binding validation tool. *Acta Crystallogr D Struct Biol*. **73(3)**:223–233.
doi:10.1107/S2059798317001061
895

Competing interests

The authors declare that no competing interests exist.

Author contributions

CP, Conception and design, Acquisition of data, Analysis and interpretation of data, Drafting or revising the article.

DH, Conception and design, Acquisition of data, Analysis and interpretation of data.

JR, Conception and design, Acquisition of data, Analysis and interpretation of data. Drafting or revising the article.

WAS, Acquisition of data.

LA, Acquisition of data.

CLM, Acquisition of data.

KJW, Acquisition of data, Interpretation of data.

DJC, Conception and design, Acquisition of data, Analysis and interpretation of data, Drafting or revising the article.

JMW, Conception and design, Acquisition of data, Analysis and interpretation of data, Drafting or revising the article.

Funding

This work was supported a Royal Society Research Grant awarded to JMW [RG130585] and a BBSRC New Investigator Grant to JMW and DJC [BB/N005570/1]. CP and WAS are funded by the BBSRC New Investigator Grant [BB/N005570/1]. JMW is funded by Newcastle University. DJC and JR are funded by the University of Edinburgh. JR is funded by a BBSRC EastBio DTP studentship [BB/M010996/1]. DH was funded by the China Scholarship Council. KJW was funded by the Wellcome Trust and Royal Society through a Sir Henry Dale Fellowship awarded to KJW [098375/Z/12/Z].

Abbreviations

Enc, encapsulin; EncFtn, encapsulated ferritin; FOC, ferroxidase centre; MS, mass spectrometry; nESI, Native nanoelectrospray ionization; CIU, collision induced unfolding.