Season-specific carry-over of early-life associations in a monogamous bird species

Ralf H.J.M. Kurvers[1,2,*], Lea Prox[3,4], Damien R. Farine[4,5,6], Coretta Jongeling[2] & Lysanne Snijders[7]

[1] Center for Adaptive Rationality, Max Planck Institute for Human Development, Lentzeallee 94, 14195 Berlin, Germany

[2] Resource Ecology Group, Wageningen University and Research, Droevendaalsesteeg 3a, 6708 PB Wageningen, the Netherlands

[3] Behavioral Ecology & Sociobiology Unit, German Primate Center, Kellnerweg 4, 37077 Göttingen, Germany

[4] Department of Biology, University of Konstanz, Universitätsstraße 10, 78464 Konstanz, Germany

[5] Department of Collective Behavior, Max Planck Institute of Animal Behavior, Universitätsstraße 10, 78464 Konstanz, Germany.

[6] Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Universitätsstraße 10, 78464 Konstanz, Germany

[7] Department of Evolutionary Ecology, Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Straße 17, 10315 Berlin, Germany

[*] Author for correspondence: kurvers@mpib-berlin.mpg.de

Abstract

1. Social bonds can have important fitness consequences. Although there is increasing evidence that social bonds carry over across contexts, few studies have investigated whether social bonds formed early in life are carried over to adulthood. For example, juveniles of monogamous species go through a major life-history stage transition—pair formation—during which the pair bond becomes a central unit of the social organization. At present, it remains unclear if pair members retain their early-life social bonds after pair formation and, if so, when they display these preferences most strongly.

2. Here we investigated whether bonds formed early in life carry over into adulthood and whether carry-over was dependent on season, in a monogamous species. Moreover, we investigated the role of familiarity, genetic relatedness and aggression on the perseverance of social bonds.

3. We studied the social structure before and after pair formation in captive barnacle geese (*Branta leucopsis*), a highly social, long-lived, monogamous species. We constructed social networks of groups of geese before pair formation, during the subsequent breeding season, and in the following wintering season. From these networks, we identified the preferred associations among individuals, and how these are carried over during seasonal changes.

4. We found that early-life associations in females were lost during the breeding season, but resurfaced during the subsequent wintering season. In males, the early-life associations persisted across both seasons, and were mediated by genetic relatedness and familiarity. The high level of aggressiveness of males, but not females, in the breeding season suggests that males may have played a key role in shaping both their own social environment and that of their female partners.

5. By taking advantage of a captive population in which animals had no temporal or spatial

restrictions to re-associate after pairing and for which we had complete data on their

social history, we show that early-life social relationships can be maintained well into

later life. Such bonds can be sustained even if they are temporarily disrupted, for example

due to reproductive behaviour. Our findings therefore highlight that the early-life social

environment can have life-long consequences on individuals' social environment.

Key-words: aggression, early life, familiarity, genetic relatedness, monogamous, pair

formation, social bonds, social networks

Introduction

Consistency of social bonds affects the extent to which individuals can develop cooperative interactions, build up reputations and form stable dominance hierarchies (Trivers 1971; Axelrod & Hamilton 1981; Dugatkin 1997; Wiley, Steadman, Chadwick & Wollerman 1999; Forkman & Haskell 2004; Nowak 2006; Samuni *et al.* 2018; Carter *et al.* 2019) and can, ultimately, affect fitness (Beletsky & Orians 1989; Silk 2007; Cameron, Setsaas & Linklater 2009; Silk *et al.* 2009; Silk *et al.* 2010; Kohn 2017). There is increasing evidence showing that social bonds carry over across time, place, and context. Shizuka *et al.* (2014) showed that golden-crowned sparrows (*Zonotrichia atricapilla*) that flocked together in one winter flocked together in the subsequent winter more often than expected based on the degree of home range overlap. Roosting associations in Bechstein's bats (*Myotis bechsteinii*) and Natterer's bats (*Myotis nattereri*) were found to remain stable across several years despite high fission-fusion dynamics (Kerth, Perony & Schweitzer 2011; Zeus, Reusch & Kerth 2018). Stanley, Mettke-Hofmann, Hager and Shultz (2018) revealed that, despite seasonal fluctuations in gregariousness and overall weak social bonds, semi-feral ponies (*Equus caballus*) maintained stable association preferences over three years. Finally, Firth and Sheldon (2016) showed that great tits' (*Parus major*) winter social associations carried over into their subsequent breeding season, as individuals bred nearer to those they were most associated with during winter. Despite evidence that social associations between individuals can persist across time and context, it is still largely unknown when these relationships are formed in the first place.

Experiences in early life are known to regularly carry over to adulthood, influencing survival and reproductive performance (Lindström 1999). Early-life social conditions are generally known to have long-lasting effects (Stanton & Mann 2012; Leris & Reader 2016; Langenhof & Komdeur 2018). Despite the likely significance of the early-life period for the formation of social bonds, only a few studies have tracked the social bonds of the same

individuals from their early life into adulthood (e.g., Linklater & Cameron 2009; Mitani 2009;

Carter, Brand, Carter, Shorrocks & Goldizen 2013), and almost exclusively in mammals. As

they mature, individuals of many taxa go through major life-history transitions, such as

dispersal (Blumstein, Wey & Tang 2009; Linklater & Cameron 2009) and pair formation

(Kurvers *et al.* 2013). These transitions will strongly impact the structure of an individual's

social environment, but perturbations in the social environment do not necessarily impact the

stability of already-established bonds (e.g., Kerth, Perony & Schweitzer 2011). It remains an

open question if a life-history transition such as pair formation will cause preferred associations

formed in early life to disappear or whether these may persist into adulthood.

Here, we examined the social structure of barnacle geese, a highly social species, before

and after pair formation, in order to study the persistence (or disappearance) of early-life

associations after a major life-history transition: pair formation. Barnacle geese are long-lived

monogamous birds, which generally find a partner at 2–3 years of age (Choudhury & Black

1994). Barnacle geese are very selective in choosing a mate; they sample one to six potential

mates in so-called "trial liaisons" before settling with a permanent partner (Choudhury & Black

1993; van der Jeugd & Blaakmeer 2001). The pair bond is extremely strong and pair members

generally stay in close proximity to each other and remain together until one of them dies

(Owen, Black & Liber 1988; Black 2001). Pairs with a longer pair bond show higher

reproductive success than pairs with a shorter pair bond (Black 2001). Pair formation is thus a

crucial step in the life history of geese. Here we took advantage of controlled experiments to

study how pair formation impacted associations formed early in life, allowing us to circumvent

problems with missing individuals in naturalistic data. We followed a captive population of

barnacle geese over several years, quantifying their social network structure before and after

pair formation. To study mediating factors underlying the potential persistence of social bonds,

we furthermore quantified the degree of familiarity and genetic relatedness in the population,

as well as individuals' aggressive interactions.

In the wild, barnacle geese, like many migratory waterfowl, show a high site fidelity to their breeding, wintering and staging areas (Percival 1991; Robertson & Cooke 1999), creating opportunities for long-term maintenance of social bonds established early in life. The extent to which individuals maintain such non-sexual bonds across seasons is not well understood, partly because of the challenges of following individuals across space and time. van der Jeugd, van der Veen and Larsson (2002) showed that during breeding, female, but not male, barnacle geese nested closer to siblings than expected by chance. This occurred not only when the female siblings bred in the same island as their parents, but also when they nested on a different island. Moreover, this relationship was only observed for siblings born in the same year. Given these observations, we predicted that females would show a higher likelihood of maintaining their early-life bonds than males after mating. We constructed social networks from observations collected during the breeding and wintering seasons. Given the elevated levels of aggression and territoriality during the breeding season (Owen & Wells 1979), we expected that associations with early-life companions after pair formation would be stronger in the wintering than the breeding season. Finally, we expected that both genetic relatedness and familiarity would positively impact the strength of social bonds.

Materials and methods

Study subjects

In late 2007 we obtained two mixed-sex groups of barnacle geese (see Fig. 1 for study timeline, and Kurvers *et al.* 2013 for more details). The two groups consisted of 21 (13 female, 8 male) and 23 individuals (8 female, 15 male) respectively. The groups were housed in separate outdoor aviaries (12 x 15 m) at the Netherlands Institute of Ecology (NIOO) in Heteren, the

Netherlands. Groups were visually, but not acoustically, isolated from each other. The aviaries

consisted of bare soil with a large pond (6 x 1 m) with running water for drinking and bathing.

Birds received ad libitum food consisting of a mixture of grains and pellets occasionally

supplemented with grass. Most individuals (40 of 44) were born in 2007 and were thus

approximately 5 months old upon arrival. All birds were captive-born, wing-clipped and (upon

arrival) fitted with uniquely coded white leg rings for identification. Birds from the two groups

had different origins, implying that birds within (but not between) groups could have a high

genetic relatedness (see below). Birds within a group will henceforth be referred to as "familiar"

individuals, and birds between groups as "unfamiliar". Geese lived for approximately 1.5 years

in these familiarity groups before the start of the social network observations (see below).

Genetic relatedness

We determined the genetic relatedness between each pair of individuals using a high-resolution

374 single nucleotide polymorphism (SNP) marker set, developed for the barnacle goose

(Jonker *et al.* 2012). We took a small blood sample (± 1 ml) from each individual from the

brachial vein and preserved it in ethanol. Entire genomic DNA was isolated using the Gentra

Systems Puregene DNA purification kit. Genotyping was conducted with Vera Code assays on

an Illumina BeadXpress (described in Kraus *et al.* 2011). We calculated the pairwise genetic

relatedness (*r*) using the program Coancestry (version 1, Wang 2011). To determine which

relatedness estimator best fitted our data, we used the empirical SNP allele frequencies of our

population and simulated 500 dyads of geese of varying relatedness coefficients. Meeting real

conditions as closely as possible, we simulated mainly unrelated dyads ($n = 430$) but also dyads

of close familial relationships, namely 30 full sibs dyads ($r = 0.5$), 10 half sibs dyads ($r = 0.25$),

and 30 first cousins dyads ($r = 0.125$). Based on this simulation, we found that the maximum-

likelihood estimator of Milligan (2003) performed best and used it for the final estimation of *r*.

This produced a strong correlation with expected values of *r* ($r^2 = 0.9$; analysis of the simulated

data set carried out with default settings). Subsequently, all pairwise relatedness values of the

experimental geese were obtained from Coancestry with standard settings (see Kurvers *et al.*

2013 for more details).

Observations prior to pair formation: early life

After living for approximately 1.5 years in their respective familiarity groups—and prior to any

pair formation—we separated all geese into two single-sex groups (June 2009; Fig. 1) to study

the factors shaping intra-sexual social association preferences while avoiding any confounding

intersexual interactions. Social network observations were conducted in the home aviaries.

Foraging associations were studied on five grass patches (40 x 20 cm, 1.5m apart) which were

replaced twice a day to avoid depletion. Other food sources were removed during the

observations. The sex of one individual could not be reliably determined at the time of the

network observations. This individual was not included in the early-life observations, but joined

the flock after these observations finished. This individual was excluded from all analyses.

Each single-sex group was observed (9 a.m.–1 p.m.) for 15 days (females: 22 June–12

July 2009; males: 13–30 July 2009). The presence of all individuals on the patches was recorded

every 4 minutes. This interval was longer than the mixing interval among individuals, in order

to ensure independence of observations (Croft, James & Krause 2008). Group composition was

rarely the same in consecutive records (females: 5.9%; males: 7.9%). Observations were

occasionally interrupted for 10 minutes in the event of an external disturbance. Since patch size

(40 x 20 cm) and group size (mean females = 1.9, range 1–5; mean males = 2.0, range 1–5;

Suppl. Fig. S1a, b) were small, we assumed that animals grazing on the same patch during a

sampling period were associating (a.k.a. gambit of the group, Whitehead 2008; Franks, Ruxton

& James 2010).

The results of the network observations before pair formation are reported in Kurvers

et al. (2013). In brief, in both sexes, familiarity and genetic relatedness predicted association

strength, whereas boldness and dominance did not. Therefore, we focus here on the role of

familiarity and genetic relatedness in governing the stability of long-term associations.

Observations after pair formation: breeding and wintering

Geese remained in the single-sex groups for six months and were then placed together in one

group (January 2010; Fig. 1). By then, most geese were 2.5 years of age, approximating the

average age of final pair formation in barnacle geese (Choudhury & Black 1994; van der Jeugd

& Blaakmeer 2001). Pair formation started rapidly, and most geese quickly formed a stable pair

relationship. Before starting the first post-pair formation network observations, 37 (of 44) geese

formed stable pair relationships. In total, geese formed 12 pairs, four triplets (three geese

continuously moving together as one unit without any aggression; two triplets consisted of two

females and one male, and two triplets of one female and two males) and seven geese (3 males,

4 females) remained unpaired. The occurrence of triplets has also been observed in the wild,

with a third party joining a pair between 10 months and 4 years, however incidence in the field

was lower (Black, Choudhury & Owen 1996). After the geese formed pair relationships, and

during their first breeding season, we again conducted observations of social associations. We

placed 10 grass patches (40 x 40 cm) in the aviary. Observations were conducted for five weeks

(31 May–2 July 2010, 25 observation days) in two 2-hour blocks per day (8 a.m.–1 p.m.)

following the same observation protocol described above. During the breeding season, many of

the paired individuals started building nests and laying eggs. We regularly checked all nests

and removed any eggs to avoid the birth of goslings.

To study the stability of associations across seasons, we repeated the observations of

social associations six months later, during the wintering season, following the same protocol

(17 December 2010–4 February 2011, 13 observation days). Comparing the pair status of all

geese between the breeding and wintering seasons, we observed that all pair relationships but

one remained the same. This reflects the strong and long lasting pair bonds in barnacle geese,

which generally stay together until one of the pair members dies (Black 2001). One individual

died in between the breeding and wintering observations. To facilitate comparison across

seasons, we removed this individual from all analyses. The final sample size was thus 42

individuals (20 females, 22 males).

Agonistic interactions

During all three observation periods (early life, breeding, wintering), we collected data on

agonistic interactions to study the role of aggression on the stability of early-life bonds. In

between scoring the presence of individuals on the patches, we scored the winner and loser of

agonistic interactions, defined as a direct confrontation between two geese and ranging from

threats with lowered head and neck to active chases with flapping wings. In the early-life

observations, we identified a total of 1,429 interactions in the female group and 2,619

interactions in the male group. We then identified 3,411 and 786 interactions during the

subsequent breeding and wintering periods, respectively.

Statistical analyses

Stability of early-life connections

We generated undirected weighted networks for each of the three periods (early life, breeding

season, and wintering season) from the observation data. Edge weights in the networks were

calculated using the simple ratio index (SRI) as an association measure using the 'asnipe'

package (Farine 2013) in R (v. 3.4.4). The SRI indicates the probability of observing two

individuals in association with each other given that one was observed. Values range from 0

(two individuals were never observed together) to 1 (two individuals were always observed

together). The SRI is considered an effective measure of association strength provided there are

no large sampling biases (Ginsberg & Young 1992; Farine & Whitehead 2015; Hoppitt &

Farine 2018). Since all observations were performed within aviaries in which all individuals

feeding on all patches could be easily observed, we did not expect a strong sampling bias.

We first analysed whether early-life associations (within the same sex) were maintained

after pair formation, i.e. during the subsequent breeding and wintering season. Within each sex,

we performed Mantel tests using the built-in node permutation test from the 'vegan' library

Mantel test function—running 50,000 permutations and using the spearman correlation

(Oksanen *et al.* 2019)—to compare dyads' association strength between (i) early life and

breeding season, and (ii) early life and wintering season. We started with a separate matrix for

each sex for the early-life associations, one matrix (containing both sexes) for the breeding

season, and one matrix for the wintering season. To compare the matrix correlation for females

(/males), we therefore first subset all females (/males) from the breeding and wintering matrix,

constructing single-sex matrices. To confirm that our results were not mainly driven by the

associations of the unpaired individuals (as we were primarily interested in the possible

continuation of associations after pair formation), we repeated these analyses after further

excluding associations between the unpaired individuals.

To investigate which factors might underlie a possible continuation of associations after

pair formation, we studied the effect of genetic relatedness and familiarity on association

strength during the breeding and wintering season. For each sex, we created a genetic

relatedness matrix (containing the genetic relatedness value of each unique pair) and a

familiarity matrix (containing information on whether a pair belonged to the same familiarity

group, yes/no). We used Mantel tests (see above) to test the correlations of these matrices with

the association matrices during breeding and wintering, conducting separate tests for females

and males.

Patch visits and aggression

As we found sex differences in the continuation of early-life associations (see below), we

investigated the role of patch visitation rates and aggression as potential underlying

mechanisms. For each individual, for each period, we calculated the mean number of patch

visits per hour. We then used a Mann-Whitney U test to determine whether the sexes differed

in their patch visitation rates in each of the three seasons. Similarly, for each individual and for

each season, we determined the mean number of aggressive interactions initiated per hour, and

tested whether the sexes differed in their likelihood to display aggression in each of the three

seasons.

Ethical permission

The animal ethical committee of both the Royal Netherlands Academy of Arts and Sciences

(KNAW) and the Wageningen University approved all experiments [protocol numbers:

2010008.b (blood sampling)]. Geese were obtained from a waterfowl breeding farm (Kooy and

Sons, 't Zand, the Netherlands) and returned there after the experiments had finished.

Results

Sex- and season-dependent effects of persistence of early-life associations

We constructed social networks for each of the three observation periods (i.e., early life,

breeding season, and wintering season, see Fig. 2), and compared the stability of the social

associations that were formed early in life to those in subsequent seasons. We found that a

female's association strengths from early life (i.e. prior to pair formation) were not significantly

correlated with her association strengths to the same pool of individuals (i.e. other females) in

the subsequent breeding season ($r = 0.04$; $p = 0.32$; Fig. 3A). Her early-life association strengths

were, however, significantly correlated with her association strengths in the subsequent wintering season (r = 0.18; p = 0.019; Fig. 3B). A male's early-life association strengths were significantly correlated with his association strengths to other males in both the subsequent breeding (r = 0.26; p = 0.002; Fig. 3C) and the following wintering season (r = 0.31; p = 0.002; Fig. 3D).

When excluding the associations among individuals that remained unpaired (7 of 42; but maintaining the associations between paired and unpaired individuals), we obtained similar results (females: early life vs. breeding: r = 0.00, p = 0.49; early life vs. wintering: r = 0.19, p = 0.018; males: early life vs. breeding: r = 0.26, p = 0.003; early life vs. wintering: r = 0.31, p = 0.003).

Relatedness and familiarity drive associations in males, but not females

In females, there was no effect of genetic relatedness or familiarity on association strength in the breeding season (genetic relatedness: r = -0.06, p = 0.79; familiarity: r = 0.04, p = 0.29). During the wintering season, both factors became more positively—yet not significantly—correlated with association strength (genetic relatedness: r = 0.09, p = 0.12; familiarity: r = 0.10, p = 0.11). For males, we found a positive effect of genetic relatedness and familiarity on association strength during breeding (genetic relatedness: r = 0.11, p = 0.041; familiarity: r = 0.15, p = 0.027) and wintering season (genetic relatedness: r = 0.09, p = 0.070; familiarity: r = 0.23, p = 0.007).

Elevated aggression in males and during the breeding season

We investigated two mechanisms that might drive the disappearance of social associations in females—but not males—during the breeding season. First, females may simply visit the food patches less than males during the breeding season, which would lower their opportunities for maintaining bonds. Though females visited patches less than males in the single-sex groups

before pair-formation (W = 50, p < 0.001; Fig. 4A), females visited the patches equally often

as males in the breeding and wintering season (both p > 0.35; Fig. 4B, C), ruling out this

explanation.

A second mechanism might be that males, being generally the more dominant member

of a pair, play a stronger role in determining the association members of a pair than do females,

especially in the breeding season. Before pair formation, males showed a higher level of

aggression than females in the single-sex groups (W = 142, p = 0.049*; Fig. 4D). Likewise,

males showed a substantially higher level of aggression than females during the breeding (W =

8, p < 0.001; Fig. 4E) and wintering seasons (W = 39, p < 0.001; Fig. 4F). Males displayed

equal levels of aggression towards males and females in the breeding season (Wilcoxon signed-

rank test: V = 76, p = 0.10, Fig. 4E) but slightly higher levels of aggression towards males than

females in the wintering season (V = 50, p = 0.02; Fig. 4F). As expected, male geese displayed

higher levels of aggression in the breeding than in the wintering season (Wilcoxon signed-rank

test, males only: V = 240, p < 0.001; Fig. 4E, F). In accordance, the mean group size at patches

was higher during the wintering than during the breeding season (mean group size wintering:

2.4; breeding: 2.0; Suppl. Fig. S1c, d). Moreover, paired individuals were more tolerant to the

presence of other individuals (i.e., not belonging to the pair) at a patch during wintering season.

In the breeding season, in 80% of cases when pair members were observed together on a patch,

there were no other individuals present. In the wintering season, this percentage dropped to

60% (mean number of non-pair individuals at a patch with a pair during breeding: 0.26, during

wintering: 0.76; Suppl. Fig. S2c, d). This is also apparent in the network graphs showing more

edges during the wintering season (Fig. 2C, D).

Discussion

Maintaining stable social associations across time and contexts can have adaptive benefits

(Kohn 2017). Yet the importance of the early-life period for the formation of such long-term

stable bonds has so far received little attention. Here we show that early-life social associations

can persist after a major life-history transition—pair formation—in a monogamous and long-

lived bird species. However, which associations were maintained and when depended on sex

and season. Early-life associations in females were lost during the breeding season, but

resurfaced again during the subsequent wintering season. In males, the early-life associations

persisted across both seasons, and were mediated by genetic relatedness and familiarity.

Elevated male aggression likely influenced the limited number of contacts outside of the pair

bond during the breeding season—and thereby the extent to which early-life associations could

be maintained, especially by females during the breeding season. Our findings extend the

understanding of how social bonds develop and are maintained over different life-history

phases and how their importance to individuals may vary with season.

Across taxa, females are well-known for maintaining long-term social bonds (Silk,

Alberts & Altmann 2003; Cameron, Setsaas & Linklater 2009; Carter, Brand, Carter, Shorrocks

& Goldizen 2013; Ilany, Booms & Holekamp 2015), which is commonly attributed to their

reproductive strategies. In our study, males, not females, maintained their social preferences

throughout both the breeding and wintering season; these social bonds were mediated by

familiarity and relatedness. Similar examples have been found in red-winged blackbirds

(*Agelaius phoeniceus*), where males that bred close to familiar neighbours fledged more

offspring (Beletsky & Orians 1989) and Assamese macaques (*Macaca assamensis*), where

males with stronger social bonds sired more offspring (Schulke, Bhagavatula, Vigilant &

Ostner 2010). Males can clearly form, and benefit from, long-lasting social bonds.

Social bonds in males often take the form of coalitions or alliances (Schulke,

Bhagavatula, Vigilant & Ostner 2010; Gilby *et al.* 2013; Connor *et al.* 2017), benefiting

individuals by providing cooperation partners in agonistic interactions with conspecifics. But

reduced aggression, for instance, via dear-enemy effects (Temeles 1994), can likewise be an

important benefit of maintaining long-term social bonds, especially in territorial animals

(Jaeger 1981; Chuang, Kam & Bee 2017; Siracusa *et al.* 2019). Additionally, familiarity can

influence vigilance behaviour, with individuals showing reduced vigilance in the proximity of

well-known conspecifics (Kutsukake 2006; Carter, Macdonald, Thomson & Goldizen 2009).

In a monogamous prey species such as barnacle geese, in which the male spends much of his

time during the breeding season on vigilance at the expense of foraging (Forslund 1993), such

benefits can be especially substantial. Lastly, familiarity-based associations have been linked

to faster predator-evasion responses and increased foraging success (Griffiths, Brockmark,

Höjesjö & Johnsson 2004; Carter, Macdonald, Thomson & Goldizen 2009). The expected

benefits of maintaining long-term social bonds vary with ecological and social conditions

(Maher & Burger 2011; Kappeler, Barrett, Blumstein & Clutton-Brock 2013; Connor *et al.*

2017), and studying the social structure of both sexes in taxa with distinct space use (e.g., natal

philopatry), life history (e.g., long life-span), social organization (e.g., fission-fusion), and

mating system (e.g., monogamy) characteristics, offers us greater insight into the drivers and

constraints of maintaining long-term stable bonds in animal societies.

The apparent loss of female social preferences in the breeding season is surprising, but

adds to our understanding of social flexibility by showing that the display of social preferences

not only varies between species and individuals (Kappeler, Barrett, Blumstein & Clutton-Brock

2013) but can also have sex-specific effects across seasons. Female chacma baboons (*Papio*

hamadryas ursinus) were similarly found to vary in their social preferences depending on

ecological context (Henzi, Lusseau, Weingrill, van Schaik & Barrett 2009): Social preferences

were more pronounced when food resources were scarce, but females acted more

"gregariously" (i.e., without social preferences) when resources were plentiful. For geese, winter is a time when resources become scarce; barnacle geese, however, are highly gregarious in winter (Black, Prop & Larsson 2014), possibly to access social information on foraging opportunities (Drent & Swierstra 1977; Kurvers *et al.* 2009; Kurvers *et al.* 2010) (but see Kurvers *et al.* 2014). Alternatively, the fluctuations in female social preferences may have been the result of the male partner unselectively excluding his partner's preferred—but not his own—early-life companions during the breeding season. The high level of aggression displayed by the males during the breeding season compared to the wintering season supports this hypothesis.

The lack of social preferences displayed by females in the breeding season appears to be in contrast to previous findings. An earlier study on wild barnacle geese found that female, but not male, geese exhibited social preferences in terms of nesting proximity, with females nesting closer to familiar females (van der Jeugd, van der Veen & Larsson 2002). It is possible that nest choice offers female geese an alternative way to maintain early-life social bonds, circumventing the potentially controlling influence of their partner during foraging interactions. Possibly, breeding in proximity to familiar and/or related conspecifics provides context-specific benefits that foraging together does not. Intraspecific brood parasitism and adoption is common in waterfowl (Choudhury, Jones, Black & Prop 1993; Forslund & Larsson 1995; Anderholm *et al.* 2009a; Andersson, Åhlund & Waldeck 2019) and breeding close to related individuals may decrease the costs, through inclusive fitness benefits, of having to care for additional offspring. Moreover, breeding in proximity could enable siblings to defend each other's nests against unrelated brood parasites. Female siblings may also actively or passively support each other in the acquisition of high-quality nest locations. Indeed, in geese, social support from family members is known to give individuals an advantage in competitive interactions, starting from an early age (Black & Owen 1989; Raveling, Sedinger & Johnson 2000; Scheiber, Weiss, Frigerio & Kotrschal 2005; Scheiber, Kotrschal & Weiss 2009). Lastly, nesting closely to

familiars may facilitate cooperative nest defence (Olendorf, Getty & Scribner 2004;

Grabowska-Zhang, Sheldon & Hinde 2012). Nest predation is a major threat for geese (Drent

& Prop 2008) that exposes the females in particular to considerable predation risk (Samelius &

Alisauskas 2006).

Our findings suggest that different types of social bonds may be beneficial in different

contexts and that the early-life period can be a crucial time for the formation of these bonds.

The next step is to disentangle whether individuals actively choose to (re)associate with familiar

companions depending on season- and context-dependent benefits, or whether the observed

fluctuations in social preferences are an emergent property following relatively simple season-

dependent social processes, such as heightened aggression. The first scenario may have

important implications for our understanding of the cognitive abilities of animals. It implies

that they can keep track of multiple types of relationships in a large fission–fusion society,

despite extended breaks, and that they can re-evaluate the benefits of these relationships

depending on context. Given the strong evidence for birthplace-independent long-term kin

discrimination in both migrating and captive barnacle geese (van der Jeugd, van der Veen &

Larsson 2002; Anderholm *et al.* 2009b; Kurvers *et al.* 2013), this level of cognitive ability is

certainly feasible and makes the long-lived barnacle goose an interesting study system to further

examine such mechanisms. Complex social patterns can be driven by cognitive ability, but also

emerge from relatively simple processes (Kappeler 2019), such as site fidelity combined with

season- and sex-dependent aggression. Unravelling how these mechanisms underlie social

complexity in a diversity of social systems will be central to our understanding of the evolution

of animal societies.

Acknowledgements

We thank Bart van Lith from the Netherlands Institute of Ecology for the caretaking of the

birds, and Chantal Althuizen and Jan Baar for help during the social network observations. We

thank Herbert Prins, Ron Ydenberg and Sip van Wieren for their guidance on the experimental

design. We thank the Faunafonds and the Koninklijke Nederlands Jagers Vereniging (KNJV)

for financial support, and Bart Nolet and Marcel Klaassen for facilitating the animal holding

facilities. We thank Rudy Jonker for blood sampling; Henk van der Jeugd for sexing the geese;

and Robert Kraus, Richard Crooijmans, Martien Groenen, Qiong Zhang and Bert Dibbits for

the genetic lab work. Finally, we thank Deborah Ain for helpful suggestions on the manuscript.

D.R.F. was funded by the Max Planck Society and the DFG Centre of Excellence 2117 "Centre

for the Advanced Study of Collective Behaviour" (ID: 422037984). L.S. was funded by a

postdoc fellowship of the Alexander von Humboldt Stiftung (Ref 3.3 - NLD - 1192631 - HFST-

P).

Authors' contributions

R.H.J.M.K and C.J. designed the study and collected the behavioural data. R.H.J.M.K, L.P.,

and D.R.F. analysed the data. R.H.J.M.K and L.S. wrote the manuscript with substantial input

from all other authors.

Data Availability Statement

The raw social association observations, the matrices containing the SRI values per sex per

season, the matrices containing the familiarity and genetic relatedness data per sex, and the raw

aggression data are uploaded on the Open Science Framework for referee inspection (link). The

R code for the main analyses is also uploaded here. These files will be made public in case of

acceptance.

References

Anderholm, S., Marshall, R.C., van der Jeugd, H.P., Waldeck, P., Larsson, K. & Andersson, M. (2009a).

Nest parasitism in the barnacle goose: evidence from protein fingerprinting and microsatellites.

Animal Behaviour, 78, 167-174.

Anderholm, S., Waldeck, P., Van der Jeugd, H.P., Marshall, R.C., Larsson, K. & Andersson, M.

(2009b). Colony kin structure and host-parasite relatedness in the barnacle goose. Molecular

Ecology, 18, 4955-4963.

Andersson, M., Åhlund, M. & Waldeck, P. (2019). Brood parasitism, relatedness and sociality: a kinship

role in female reproductive tactics. Biological Reviews, 94, 307-327.

Axelrod, R. & Hamilton, W. (1981). The evolution of cooperation. Science, 211, 1390-1396.

Beletsky, L.D. & Orians, G.H. (1989). Familiar neighbors enhance breeding success in birds.

Proceedings of the National Academy of Sciences, 86, 7933-7936.

Black, J.M. (2001). Fitness consequences of long-term pair bonds in barnacle geese: monogamy in the

extreme. Behavioral Ecology, 12, 640-645.

Black, J.M., Choudhury, S. & Owen, M. (1996). Do barnacle geese benefit from lifelong monogamy?

Oxford Ornithology Series, 6, 91-117.

Black, J.M. & Owen, M. (1989). Parent Offspring Relationships in Wintering Barnacle Geese. Animal

Behaviour, 37, 187-198.

Black, J.M., Prop, J. & Larsson, K. (2014) The barnacle goose. Bloomsbury Publishing.

Blumstein, D.T., Wey, T.W. & Tang, K. (2009). A test of the social cohesion hypothesis: interactive

female marmots remain at home. Proceedings of the Royal Society B-Biological Sciences, 276,

3007-3012.

Cameron, E.Z., Setsaas, T.H. & Linklater, W.L. (2009). Social bonds between unrelated females

increase reproductive success in feral horses. Proceedings of the National Academy of Sciences

of the United States of America, 106, 13850-13853.

Carter, A.J., Macdonald, S.L., Thomson, V.A. & Goldizen, A.W. (2009). Structured association patterns

and their energetic benefits in female eastern grey kangaroos, Macropus giganteus. Animal

Behaviour, 77, 839-846.

Carter, G.G., Farine, D.R., Crisp, R.J., Vrtilek, J.K., Ripperger, S. & Page, R. (2019). Development of
new food-sharing relationships among nonkin vampire bats. bioRxiv, 534321.

Carter, K.D., Brand, R., Carter, J.K., Shorrocks, B. & Goldizen, A.W. (2013). Social networks, long-
term associations and age-related sociability of wild giraffes. Animal Behaviour, 86, 901-910.

Choudhury, S. & Black, J.M. (1993). Mate selection behavior and sampling strategies in geese. Animal
Behaviour, 46, 747-757.

Choudhury, S. & Black, J.M. (1994). Barnacle geese preferentially pair with familiar associates from
early-life. Animal Behaviour, 48, 81-88.

Choudhury, S., Jones, C.S., Black, J.M. & Prop, J. (1993). Adoption of young and intraspecific nest
parasitism in barnacle geese. The Condor, 95, 860-868.

Chuang, M.-F., Kam, Y.-C. & Bee, M.A. (2017). Territorial olive frogs display lower aggression
towards neighbours than strangers based on individual vocal signatures. Animal Behaviour,
123, 217-228.

Connor, R.C., Cioffi, W.R., Randic, S., Allen, S.J., Watson-Capps, J. & Krutzen, M. (2017). Male
alliance behaviour and mating access varies with habitat in a dolphin social network. Scientific
Reports, 7, 46354.

Croft, D.P., James, R. & Krause, J. (2008) Exploring animal social networks. Princeton University Press,
Princeton.

Drent, R.H. & Prop, J. (2008). Barnacle goose Branta leucopsis survey on Nordenskiöldkysten, west
Spitsbergen 1975–2007: breeding in relation to carrying capacity and predator impact.
Circumpolar Stud, 4, 59-83.

Drent, R.H. & Swierstra, P. (1977). Goose flocks and food finding: field experiments with Barnacle
Geese in winter. Wildfowl, 28, 15-20.

Dugatkin, L.A. (1997) Cooperation among animals: an evolutionary perspective. Oxford University
Press on Demand.

Farine, D.R. (2013). Animal social network inference and permutations for ecologists in R using asnipe.
Methods in Ecology and Evolution, 4, 1187-1194.

Farine, D.R. & Whitehead, H. (2015). Constructing, conducting and interpreting animal social network

analysis. Journal of Animal Ecology, 84, 1144-1163.

Firth, J.A. & Sheldon, B.C. (2016). Social carry-over effects underpin trans-seasonally linked structure

in a wild bird population. Ecology Letters, 19, 1324-1332.

Forkman, B. & Haskell, M. (2004). The maintenance of stable dominance hierarchies and the pattern of

aggression: support for the suppression hypothesis. Ethology, 110, 737-744.

Forslund, P. (1993). Vigilance in relation to brood size and predator abundance in the barnacle goose,

Branta leucopsis. Animal Behaviour, 45, 965-973.

Forslund, P. & Larsson, K. (1995). Intraspecific nest parasitism in the barnacle goose: behavioural

tactics of parasites and hosts. Animal Behaviour, 50, 509-517.

Franks, D.W., Ruxton, G.D. & James, R. (2010). Sampling animal association networks with the gambit

of the group. Behavioral Ecology and Sociobiology, 64, 493-503.

Gilby, I.C., Brent, L.J.N., Wroblewski, E.E., Rudicell, R.S., Hahn, B.H., Goodall, J. & Pusey, A.E.

(2013). Fitness benefits of coalitionary aggression in male chimpanzees. Behavioral Ecology

and Sociobiology, 67, 373-381.

Ginsberg, J.R. & Young, T.P. (1992). Measuring association between individuals or groups in

behavioral studies. Animal Behaviour, 44, 377-379.

Grabowska-Zhang, A., Sheldon, B. & Hinde, C. (2012). Long-term familiarity promotes joining in

neighbour nest defence. Biology Letters, 8, 544-546.

Griffiths, S.W., Brockmark, S., Höjesjö, J. & Johnsson, J. (2004). Coping with divided attention: the

advantage of familiarity. Proceedings of the Royal Society of London. Series B: Biological

Sciences, 271, 695-699.

Henzi, S.P., Lusseau, D., Weingrill, T., van Schaik, C.P. & Barrett, L. (2009). Cyclicity in the structure

of female baboon social networks. Behavioral Ecology and Sociobiology, 63, 1015-1021.

Hoppitt, W.J. & Farine, D.R. (2018). Association indices for quantifying social relationships: how to

deal with missing observations of individuals or groups. Animal Behaviour, 136, 227-238.

Ilany, A., Booms, A.S. & Holekamp, K.E. (2015). Topological effects of network structure on long-

term social network dynamics in a wild mammal. Ecology Letters, 18, 687-695.

Jaeger, R.G. (1981). Dear enemy recognition and the costs of aggression between salamanders. The

American Naturalist*, 117*, 962-974.

Jonker, R.M., Zhang, Q., Van Hooft, P., Loonen, M., Van der Jeugd, H.P., Crooijmans, R., . . . Kraus,

R.H.S. (2012). The development of a genome wide SNP set for the barnacle goose *Branta*

leucopsis. PLoS ONE*, 7*, e38412.

Kappeler, P.M. (2019). A framework for studying social complexity. Behavioral Ecology and

Sociobiology*, 73*, 13.

Kappeler, P.M., Barrett, L., Blumstein, D.T. & Clutton-Brock, T.H. (2013). Constraints and flexibility

in mammalian social behaviour: introduction and synthesis. Philosophical Transactions of the

Royal Society B: Biological Sciences*, 368*, 20120337.

Kerth, G., Perony, N. & Schweitzer, F. (2011). Bats are able to maintain long-term social relationships

despite the high fission-fusion dynamics of their groups. Proceedings of the Royal Society B*,

278*, 2761-2767.

Kohn, G.M. (2017). Friends give benefits: autumn social familiarity preferences predict reproductive

output. Animal Behaviour*, 132*, 201-208.

Kraus, R.H.S., Kerstens, H., Van Hooft, P., Crooijmans, R., Van Der Poel, J., Elmberg, J., . . . Groenen,

M. (2011). Genome wide SNP discovery, analysis and evaluation in mallard (*Anas*

platyrhynchos). BMC Genomics*, 12*, 150.

Kurvers, R.H., Straates, K., Ydenberg, R.C., van Wieren, S.E., Swierstra, P.S. & Prins, H.H. (2014).

Social information use by barnacle geese Branta leucopsis, an experiment revisited. Ardea*, 102*,

173-181.

Kurvers, R.H.J.M., Adamczyk, V.M.A.P., Kraus, R.H.S., Hoffman, J.I., van Wieren, S.E., van der

Jeugd, H.P., . . . Jonker, R.M. (2013). Contrasting context dependence of familiarity and kinship

in animal social networks. Animal Behaviour*, 86*, 993-1001.

Kurvers, R.H.J.M., Eijkelenkamp, B., van Oers, K., van Lith, B., van Wieren, S.E., Ydenberg, R.C. &

Prins, H.H.T. (2009). Personality differences explain leadership in barnacle geese. Animal

Behaviour*, 78*, 447-453.

Kurvers, R.H.J.M., Prins, H.H.T., van Wieren, S.E., van Oers, K., Nolet, B.A. & Ydenberg, R.C. (2010).

The effect of personality on social foraging: shy barnacle geese scrounge more. Proceedings of

the Royal Society B-Biological Sciences, 277, 601-608.

Kutsukake, N. (2006). The context and quality of social relationships affect vigilance behaviour in wild

chimpanzees. Ethology, 112, 581-591.

Langenhof, M.R. & Komdeur, J. (2018). Why and how the early-life environment affects development

of coping behaviours. Behavioral Ecology and Sociobiology, 72, 34.

Leris, I. & Reader, S.M. (2016). Age and early social environment influence guppy social learning

propensities. Animal Behaviour, 120, 11-19.

Lindström, J. (1999). Early development and fitness in birds and mammals. Trends in Ecology &

Evolution, 14, 343-348.

Linklater, W.L. & Cameron, E.Z. (2009). Social dispersal but with philopatry reveals incest avoidance

in a polygynous ungulate. Animal Behaviour, 77, 1085-1093.

Maher, C.R. & Burger, J.R. (2011). Intraspecific variation in space use, group size, and mating systems

of caviomorph rodents. Journal of Mammalogy, 92, 54-64.

Milligan, B.G. (2003). Maximum-likelihood estimation of relatedness. Genetics, 163, 1153-1167.

Mitani, J.C. (2009). Male chimpanzees form enduring and equitable social bonds. Animal Behaviour,

77, 633-640.

Nowak, M.A. (2006). Five rules for the evolution of cooperation. Science, 314, 1560-1563.

Oksanen, J., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., . . . Wagner, H. (2019)

vegan: Community Ecology Package. R package version 2.5-4. .

Olendorf, R., Getty, T. & Scribner, K. (2004). Cooperative nest defence in red–winged blackbirds:

reciprocal altruism, kinship or by–product mutualism? Proceedings of the Royal Society of

London. Series B: Biological Sciences, 271, 177-182.

Owen, M., Black, J.M. & Liber, H. (1988) Pair bond duration and timing of its formation in barnacle

geese (Branta leucopsis). *Waterfowl in Winter* (ed. M.W. Weller). University of Minnesota

Press, Minneapolis.

Owen, M. & Wells, R. (1979). Territorial behaviour in breeding geese--a re-examination of Ryder's

hypothesis. Wildfowl, 30, 20-26.

Percival, S.M. (1991). The Population-Structure of Greenland Barnacle Geese Branta-Leucopsis on the

Wintering Grounds on Islay. Ibis, 133, 357-364.

Raveling, D.G., Sedinger, J.S. & Johnson, D.S. (2000). Reproductive success and survival in relation to

experience during the first two years in Canada geese. The Condor, 102, 941-945.

Robertson, G.J. & Cooke, F. (1999). Winter philopatry in migratory waterfowl. The Auk, 116, 20-34.

Samelius, G. & Alisauskas, R.T. (2006). Sex-biased costs in nest defence behaviours by lesser snow

geese (Chen caerulescens): consequences of parental roles? Behavioral Ecology and

Sociobiology, 59, 805-810.

Samuni, L., Preis, A., Mielke, A., Deschner, T., Wittig, R.M. & Crockford, C. (2018). Social bonds

facilitate cooperative resource sharing in wild chimpanzees. Proceedings of the Royal Society

B, 285, 20181643.

Scheiber, I.B.R., Kotrschal, K. & Weiss, B.M. (2009). Benefits of family reunions: Social support in

secondary greylag goose families. Hormones and Behavior, 55, 133-138.

Scheiber, I.B.R., Weiss, B.M., Frigerio, D. & Kotrschal, K. (2005). Active and passive social support

in families of greylag geese (Anser anser). Behaviour, 142, 1535-1557.

Schulke, O., Bhagavatula, J., Vigilant, L. & Ostner, J. (2010). Social Bonds Enhance Reproductive

Success in Male Macaques. Current Biology, 20, 2207-2210.

Shizuka, D., Chaine, A.S., Anderson, J., Johnson, O., Laursen, I.M. & Lyon, B.E. (2014). Across-year

social stability shapes network structure in wintering migrant sparrows. Ecology Letters, 17,

998-1007.

Silk, J.B. (2007). Social components of fitness in primate groups. Science, 317, 1347-1351.

Silk, J.B., Alberts, S.C. & Altmann, J. (2003). Social bonds of female baboons enhance infant survival.

Science, 302, 1231-1234.

Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., . . . Cheney, D.L.

(2009). The benefits of social capital: close social bonds among female baboons enhance

offspring survival. Proceedings of the Royal Society B-Biological Sciences, 276, 3099-3104.

Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., . . . Cheney, D.L.

(2010). Strong and Consistent Social Bonds Enhance the Longevity of Female Baboons. Current

Biology, 20, 1359-1361.

Siracusa, E.R., Wilson, D.R., Studd, E.K., Boutin, S., Humphries, M.M., Dantzer, B., . . . McAdam,

A.G. (2019). North American red squirrels mitigate costs of territory defence through social

plasticity. Animal Behaviour, 151, 29-42.

Stanley, C.R., Mettke-Hofmann, C., Hager, R. & Shultz, S. (2018). Social stability in semiferal ponies:

networks show interannual stability alongside seasonal flexibility. Animal Behaviour, 136, 175-

184.

Stanton, M.A. & Mann, J. (2012). Early social networks predict survival in wild bottlenose dolphins.

PLoS ONE, 7, e47508.

Temeles, E.J. (1994). The role of neighbours in territorial systems: when are they 'dear enemies'? Animal

Behaviour, 47, 339-350.

Trivers, R.L. (1971). The Evolution of Reciprocal Altruism. The Quarterly Review of Biology, 46, 35-

57.

van der Jeugd, H.P. & Blaakmeer, K.B. (2001). Teenage love: the importance of trial liaisons, subadult

plumage and early pairing in barnacle geese. Animal Behaviour, 62, 1075-1083.

van der Jeugd, H.P., van der Veen, I.T. & Larsson, K. (2002). Kin clustering in barnacle geese:

familiarity or phenotype matching? Behavioral Ecology, 13, 786-790.

Wang, J.L. (2011). COANCESTRY: a program for simulating, estimating and analysing relatedness and

inbreeding coefficients. Molecular Ecology Resources, 11, 141-145.

Whitehead, H. (2008) Analyzing animal societies: Quantative methods for vertebrate social analysis.

University of Chicago Press, Chicago.

Wiley, R.H., Steadman, L., Chadwick, L. & Wollerman, L. (1999). Social inertia in white-throated

sparrows results from recognition of opponents. Animal Behaviour, 57, 453-463.

Zeus, V.M., Reusch, C. & Kerth, G. (2018). Long-term roosting data reveal a unimodular social network

in large fission-fusion society of the colony-living Natterer's bat (Myotis nattereri). Behavioral

Ecology and Sociobiology, 72, 99.

Figure 1

Figure 1. Schematic overview of the experimental procedures. Geese were procured in

November 2007—at which point almost all were juveniles (born in 2007)—and placed in one

of two familiarity groups. In July 2009, males and females of both groups were placed together

and we quantified the social network structure of both groups (to examine the role of familiarity

and genetic relatedness on social structure, Kurvers *et al.* 2013). In January 2010, all geese were

placed together in one group and individuals rapidly started to form pairs. In the subsequent

breeding season (June 2010) and wintering season (January 2011), we quantified the social

network structure of the entire group. Note that the sex of one individual could not be reliably

determined at the time of the first network observation and hence was not included in the early-

life observations. Another individual died in between the breeding and wintering season. Both

individuals were excluded from all analyses, resulting in a final sample size of 42 (20 females,

22 males).

Figure 2

Figure 2. The (A) female and (B) male network in single-sex groups prior to pair formation

and the network of all individuals after pair formation during the (C) breeding, and (D)

wintering season. Yellow/blue circles represent females/males respectively. For visualization,

we removed associations below SRI values of 0.005. Thin/thick lines represent SRI values

below/above 0.1. Coloured lines indicate a pair bond. Networks were created with the ggnet2

function in R using the Fruchterman-Reingold algorithm for node placement.

Figure 3

Figure 3. The relationship between the association strength of female dyads comparing (A) early life and breeding season, and (B) early life and wintering season, and the relationship between the association strength of male dyads comparing (C) early life and breeding season, and (D) early life and wintering season. Lines are linear regression lines, including 95% confidence bands.

Figure 4. The mean number of patches visited per hour for (A) females and males in the single-sex groups prior to pair formation, and for both sexes during the (B) breeding and (C) wintering seasons after pair formation. The mean number of aggressive events initiated per hour for (D) females and males in the single-sex groups prior to pair formation, and for both sexes during the (E) breeding and (F) wintering season after pair formation. (E, F) Colours indicate the sex of the individual receiving the aggression. Dots represent individuals. Boxplots show median, interquartile ranges and whiskers show the lowest/highest values within the 1.5 interquartile range.

707 # Supporting Information

709 Belonging to

711 **Season-specific carry-over of early-life associations in a monogamous bird**

712 **species**

714 Ralf H.J.M. Kurvers, Lea Prox, Damien R. Farine, Coretta Jongeling & Lysanne Snijders

Supplementary Figure S1. (A, B) The frequency of observed group sizes during the early-life social network observations for (A) females and (B) males. Red dashed lines represent mean group size (mean group size females: 1.92; males: 1.96). Males frequented patches at a higher rate (see also Fig. 4A). (C, D) The frequency of observed group sizes during the (C) breeding and (D) wintering season. Red dashed lines represent mean group size (mean group size breeding: 2.0; wintering: 2.4). Note that the overall higher frequency of group sizes in the breeding season is due to more observation days during breeding (n=25 days) than during wintering season (n=13 days). The mean patch visit rate was in fact slightly higher in wintering than in breeding season (see Fig. 4B, C).

Supplementary Figure S2. (A, B) The observed likelihood that when a paired individual was present on a patch that its partner was also present ('partner'), that it was together with another individual but not its partner ('social'), or that it was alone ('alone'), for the (A) breeding, and (B) wintering season. (C, D) The observed likelihood of the number of individuals not belonging to the pair, which were present at a patch in the presence of a pair during the (C) breeding, and (D) wintering season. To illustrate, in the breeding season, in 80% of cases when pair members were observed together on a patch, there would be no other individuals (i.e., not belonging to that pair) present. In the wintering season, this percentage was around 60%. Red dashed lines represent the average number of non-paired geese present with a pair (mean during breeding: 0.26, during wintering: 0.76).