The sensory representation of causally controlled objects

Kelly B. Clancy[1] and Thomas D. Mrsic-Flogel[1]

1. Sainsbury Wellcome Centre, UCL, 25 Howland Street London, W1T 4JG, UK;
previously at the Biozentrum, University of Basel

Correspondence: Kelly Clancy (k.clancy@ucl.ac.uk) and Thomas Mrsic-Flogel (t.mrsic-
flogel@ucl.ac.uk)

Summary
Intentional control over external objects is informed by our sensory experience of

them. To study how causal relationships are learned and effected, we devised a

brain machine interface (BMI) task utilising wide-field calcium signals. Mice

learned to entrain activity patterns in arbitrary pairs of cortical regions to guide a

visual cursor to a target location for reward. Brain areas that were normally

correlated could be rapidly reconfigured to exert control over the cursor in a

sensory feedback-dependent manner. Higher visual cortex was more engaged

when expert but not naïve animals controlled the cursor. Individual neurons in

higher visual cortex responded more strongly to the cursor when mice controlled

it than when they passively viewed it, with the greatest response boosting as the

cursor approached target location. Thus, representations of causally-controlled

objects are sensitive to intention and proximity to the subject's goal, potentially

strengthening sensory feedback to allow more fluent control.

27
Introduction

How does the brain infer a causal relationship between its activity and the sensed world,
and how does this affect the sensory encoding of controlled external objects? Actions
and perceptions reciprocally affect one another in a mutual dialog (Dewey, 1896). The
sense of causal control can be operationalised to inferring a causal relationship between
a subject's internally generated actions or activity, and their outcome in the external
world (Haggard, 2017). In motor learning, for example, the relationship between an
action and its outcome can be learned and re-learned throughout adulthood as animals
acquire new motor skills. Brain machine interfaces (BMI) are a method for investigating
how subjects learn arbitrary action-outcome relationships (Fetz, 1969; Bakay and
Kennedy, 1998; Nicolelis, 2001; Donoghue, 2002; Carmena et al., 2003; Sitaram et al.,
2017). When learning to control a BMI, rodents have been found to employ the same
mechanisms as implicated in motor learning (Koralek et al., 2012; Neely et al., 2018).
But unlike motor learning, wherein animals learn a task and researchers must search for
correlates of the behaviour in patterns of neural activity, BMIs allow the experimenter to
precisely control sensory feedback, as well as prescribe the requisite activity patterns
necessary for successful task execution, which can then be changed day to day. Thus,
animals learning neuroprosthetic control of external objects must engage in continuous
self-monitoring to assess the contingency between their neural activity and its outcome,
preventing them from executing a habitual or fixed motor pattern, and encouraging
animals to learn arbitrary new sensorimotor mappings on the fly.
50
A key aspect of this self-monitoring is the sensory feedback from the object being
controlled by the agent. Yet little is known about how causally controlled objects are
represented in the brain. Studies have implicated parietal cortex in intention, as well as
in the subjective assessment of agency over outcome. In human subjects, disrupting
activity in parietal cortex with TMS temporarily ablates self-reported agency (Chambon
et al., 2015). Parietal activity has been found to be involved in representing task rules,
the value of competing actions, and visually-guided real-time motor plan updating: both
in humans (Pisella et al., 2000; Kahnt et al., 2014; Wisniewski et al., 2015; Zapparoli et
al., 2020) and non-human primates (Andersen et al., 1997; Sugrue et al., 2004). Motor
plans can be decoded from parietal activity, and its responses are task-, expectation-,
and goal-dependent, in humans, (Rushworth et al., 2001; Desmurget et al., 2009; Aflalo
et al., 2015), non-human primates (Mountcastle et al., 1975; Gnadt and Andersen, 1988;
Lacquaniti et al., 1995; Johnson et al., 1996; Churchland et al., 2008), and rodents

(Licata et al., 2017; Pho et al., 2018; Mohan et al., 2018, 2019). All of this evidence
suggests that, across multiple species, parietal cortex plays a role in intentional, goal
directed behaviours (Rizzolatti et al., 1997; Andersen and Buneo, 2002; Andersen and
Cui, 2009). However, previous studies of the role of parietal cortex in intention have not
examined how causal control affects sensory representations across different
sensorimotor contingencies.
70
To address this, we devised a mouse model of adaptive causal control. Animals learned
to guide a visual feedback cursor to a target location in order to obtain a reward using
activity in experimenter-defined cortical areas, recorded with widefield imaging. This had
the added benefit of acting as an unbiased screen to identify dorsal cortical areas
involved in learning the task. We found that higher visual areas, including anteromedial
cortex, AM, were more engaged when expert animals controlled the BMI. These higher
areas are considered by some to be a putative homologue of parietal cortex in mice
(Harvey et al., 2012; Licata et al., 2017; Mohan et al., 2018; Pho et al., 2018; Lyamzin
and Benucci, 2019). In order to gain insight into what this task-related activity looks like
on an individual neuron level, we targeted single-cell recordings to the functionally-
identified task-relevant region AM, and found that the visual cursor elicited larger
responses when an animal was controlling it in a closed-loop configuration than when
passively viewing it in an open-loop configuration (Bagur et al., 2018). Responses were
highest when the cursor was closest to the target zone, and were sensitive to the
cursor's instantaneous trajectory: they were greater when the cursor was moving
towards the target than away from it. Thus, the sensory representation of the visual
object was sensitive to the subject's intention, as well as its perception of the object's
instantaneous trajectory with respect to its goal. Given that animals had to relearn a
changing sensorimotor contingency on the fly, we surmised that the heightened sensory
representation of the cursor at the target position might serve to strengthen the signal to
adjudicating areas for informing fluent control over external objects. Indeed, the neural
activity in AM during the task condition carried significantly more information about the
cursor identity than during the passive playback condition.
94
95
96 **Results**
97

Goal-directed control of a visual cursor using areal signals

In order to investigate how causal control over external objects is effected and encoded

in mammalian cortex, we trained mice to control a visual feedback cursor using real-time

calcium signals recorded with wide-field imaging (largely reflecting the summed spiking

activity of local cells (Makino et al., 2017; Clancy et al., 2019). We imaged the dorsal

cortex in transgenic mice expressing the calcium indicator GCamp6s in CaMKII+

pyramidal neurons (Wekselblatt et al., 2016), assigning two small frontal regions to

control a visual feedback cursor (Figure 1A-B, Supplemental Video 1-3), similar to a task

described previously (Clancy et al., 2014; Koralek et al., 2012). Animals were head-fixed

under the widefield microscope, and free to run on a styrofoam wheel. The animal's

goal was to bring a visual cursor (a copy of which was presented to both eyes on two

separate monitors flanking each side of the mouse) to a target position in the centre of

its visual field (Figure 1C). Animals could achieve this by increasing activity in control

region 1 (R1) relative to control region 2 (R2). If activity in R2 was greater than R1, the

cursor moved towards the back of the animal's visual field. Thus, in this design, animals

could not hit the target simply by generally increasing or decreasing activity across

cortex, but had to differentially modulate the activity of these experimenter-specified

control regions (Figure 1D-F; Supp. Figure 1A-B). These control regions were usually

placed over ipsilateral motor areas and were changed from day to day (Figure 1B,

Supplemental Table 1). Control regions were deliberately small ($\sim$0.1 mm^2) as we

reasoned that it would be easier for animals to control a smaller area, but this was not

systematically tested. We avoided using anterior lateral motor cortex (ALM) in the

control regions, to minimise the effect of licking.

Figure 1. A widefield-imaging based brain machine interface
a. Task schematic. Clockwise starting from cartoon mouse: widefield signals were imaged from head-fixed animals in real time and transmuted into the position of a visual cursor. Two small regions (R1 and R2) were used for controlling the cursor, and activity recorded from these areas was fed into a decoder such that their activity opposed one another. Example dF/F for the two regions is shown at top, with blue arrows denoting trial starts, and pink arrows denoting target hits. Activity averaged around hits for one example animal, one day, shown for R1 and R2. Animals had to increase activity in R1 relative to R2 to bring the cursor to a rewarded position at the centre of their visual field, at which point they could collect a reward after a 1 second delay. **B.** Positions of control ROIs (R1 in blue, R2 in green) for all 7 animals over the course of training (averaging 15 days each), superimposed on the Allen Brain Atlas (totalling 104 pairs). **C.** Feedback schematic: the cursor could take one of 8 potential positions on screen, with position 8, the target, rewarded. **D.** ΔF/F in control regions triggered around hits for three example animals on one day of training, indicating different strategies animals use to achieve reward. Pink line indicates the time of target hit, grey line indicates reward delivery. **E.** Activity in R1 subtracted by activity in R2, averaged around target hits for all mice on a day of training (n=7 mice, shading represents s.e.m.). **F.** Animals could not control cursor using lick alone. ΔF/F triggered around lick bouts in spontaneous activity for the same three example animals on one day of training, indicating animals could not achieve the differential activation of R1 and R2 using lick alone. Grey line indicates time of lick.

The visual feedback cursor could take one of eight positions on the monitors (Figure 1C), and the cursor had to be held at the target for 300 milliseconds to count as a hit. When animals succeeded in holding the cursor at the target position, a soya milk reward was delivered after a one second delay. If animals failed to bring the cursor to the target position within 30 seconds, the trial was considered a miss, and a white noise miss-cue was played, followed by a brief time out. Chance performance was assessed using

152 spontaneous activity recorded before the task began, and represents the estimated hit
153 rate that the animal would achieve using spontaneous fluctuations of neural signals
154 alone. Animals improved their performance over training (Figure 2A, n = 7 mice), and
155 took less time to reach a criterion performance of 50% hit trials over days (Figure 2B).
156
Animals could perform the task without overt movements, licking, or eye saccades (Supp.
Figure 1C-E). Several recent papers have shown that many motor activities can
influence cortical activity (Stringer et al., 2019; Musall et al., 2019; Orsolic et al., 2019;
Salkoff et al., 2020), raising the possibility that mice may adopt a motor strategy to
control the activation of cortical regions used for BMI. Our analyses show that animals
could perform the task without gross overt movements that were detected by monitoring
licking, eye saccades or running speed (Supp. Figure 1C-E), though it is possible that
subtler movement were not detected. This is a question, common to all BMI studies, that
is ultimately unanswerable without full recordings from every muscle in the body. Here
we did not take full body videos of the animals performing the task, but in future work
including video monitoring can offer insights on subtler motor outputs. Nevertheless,
animals in this study could discover arbitrary activation patterns of different brain areas
within and between training sessions, and use these activations to control a visual cursor
in a manner dependent on sensory feedback: that is, they could flexibly work out the
arbitrary coupling of action patterns (whether purely confined to the brain, or involving
peripheral muscle contractions) with sensory feedback in order to achieve a goal. The
flexibility with which the animals could rapidly readjust their control of arbitrary brain
areas suggests some degree of adaptability in decorrelating otherwise highly correlated
brain areas (see below).
176

Figure 2. Animals learn to control a visual cursor using areal neural activity
A. Behavioural performance (percentage of trials the animal successfully reached the target within the 30 second trial window) increased above chance over the course of days. Shaded region denotes chance s.e.m., assessed as how often spontaneous activity would have achieve hits, averaged across 7 mice. **B.** Animals achieve 50% performance faster over the course of days of training, calculated by taking a moving average of the number of times the animal successfully reached the target per trial. **C.** Average hits per minute (grey) increased over the course of a training session, and recovered within minutes when control regions were changed (n = 7 mice on day six of training, shading denotes 95% confidence interval, see Supplemental Table 1). Individual hit times, pooled for all animals, shown below rate curves. At chance baseline, animals should perform around 0.3 hits/minute (here, animals started at slightly worse than chance, likely due to the fact that some of the control regions positions had been changed from the previous day). After R1 and R2 were swapped mid-session, the estimate of how well the animals would have done using the original control regions is shown in teal: the previous ("ghost") decoder's performance starts near the animal's previous hit rate, suggesting they persist with their original strategy, but this drops off as animals discover the new rewarded contingency and recover their performance. **D.** Performance dropped to near chance when rewards were given randomly, but recovered when target and reward was again coupled on a subsequent day (n = 5 mice). Performance also dropped to chance when the visual feedback cursor was uncoupled

198 from neural activity and presented at random (n = 3 mice), even though animals could
199 still achieve reward using the previously learned activity pattern.
200
201
Control regions could be changed from day to day or within a session: hit rates improved
over time within a session, and recovered after control regions were changed, indicative
of learning (Figure 2C, Supp. Figure 2A-B). After recovery, the animals' hit rate before
and after this control region switch were not statistically different (Supp. Figure 2C). The
decoder transformation for both the pre- and post-switch conditions was calculated using
the same spontaneous baseline taken before the training session. To test if the
behaviour was goal-directed, we dissociated the reward from the target position. On day
8 of training, animals were allowed to perform the task as usual, but the training session
was constrained to 30 minutes. Thereafter, the visual feedback was coupled to the
animal's neural activity as before, but rewards were given randomly, at the same rate as
an expert animal engaged in the task (~1.3 hits/minute). During this random reward
condition, the animals' target hit rate dropped to chance, indicating that the behaviour
was goal-oriented (Figure 2D). The fluorescence signals representing the difference
between R1 and R2 increased over the normal training session, indicative of increased
efficacy of control, and decreased when reward was randomised, again suggesting that
animals were effecting the requisite neural patterns in a goal-directed manner (Supp.
Figure 2D-E). Animals were able to recover their performance following reinstatement of
the normal task on the next day of training (Figure 2D). For a subset of animals, the
visual feedback was then randomised (referred to as the random feedback condition):
the visual cursor was presented at random positions, though animals could still achieve
the target with the appropriate neural activity patterns independent of cursor position.
The animals' ability to bring the cursor to the target dropped to chance levels without
meaningful visual feedback (Figure 2D). Activity leading to hits did not resemble activity
induced by licking around reward collection, as evidenced in both the normal task and
random-reward conditions (Supp. Figure 2F). Altogether, this suggests that animals
performed the task in a flexible, goal-directed, visual feedback-dependent manner.
228

229 **Exploration and exploitation in neural activity space**

231 As control regions were changed day-to-day, the activity patterns necessary for
232 successful BMI control had to be re-learned each session. Example fluorescence traces
233 from control regions indicate the areas were initially highly spontaneously correlated

(Figure 3A, top trace; Supplemental Figure 1A-B; Supplemental Video 3). Early in the training session, hits were preceded by diverse activity patterns (Figure 3A, middle trace). By late in the training session, the activity patterns leading to hits were more consistent (Figure 3A, bottom trace). Animals found different ways to achieve this consistency, sometimes by sweeping activity through R1 towards R2, or by depressing R2 while activating R1 (Figure 1D, Supp. Videos 1-3).

In order to achieve the prescribed activity pattern, animals had to functionally decorrelate the two control regions, which were usually highly spontaneously correlated. Indeed, activity in dorsal cortex was globally correlated in animals pre-task, as indicated by correlation maps using R1 and R2 as seed pixels (Figure 3B; Supp. Figure 1B). Although activity in dorsal cortex was globally correlated in animals pre-task, correlations between these areas decreased during task performance (Figure 3B, Supp. Figure 3B), as did the correlations between the control regions and primary visual cortex, V1, (Supp. Figure 3C). Correlations between the control regions and primary somatosensory cortex, S1, as well as between S1 and V1 were unchanged between spontaneous activity ans the task (Supp. Figure 3C-D). Only periods of task performance, and not reward collection or inter-trial waiting periods, were included in these analyses. Animals could arbitrarily decorrelate different regions: in a separate cohort of animals trained using auditory feedback instead of the visual feedback cursor, animals could also decorrelate visual control areas (Figure 3B, rightmost panel; Supp. Figure 4). Interestingly, the task-induced correlation patterns were invariably bilateral, even when control regions were ipsilateral to each other. Not every training session resulted in the kind of decorrelated maps evident in Figure 3B; reflecting that some areas are harder for the animals to decorrelate than others: animals took longer to reach criterion 50% performance when control regions were spontaneously highly correlated (Figure 3C). The distance between control regions did not affect the time animals required to reach 50% performance (Supp. Figure 3A), suggesting that areas that were the most spontaneously correlated, irrespective of their distance from one another, were the hardest to use to perform the task.

Figure 3. Exploration and exploitation of neural activity patterns

A. Areal signals were highly spontaneously correlated before the training session (top trace). Animals explored different activation patterns early in the training session (middle trace) in order to discover and exploit successful patterns by the end of the session

(bottom trace). Pink arrows denote target hits. Pearson's correlation between R1 and R2
indicated on the right of each trace. Right: run, lick and saccade averaged around hits
for this training session, grey shading denotes 95% confidence interval around mean.
Pink arrow denotes target hits, black arrow denotes reward delivery. **B.** Correlation map
across cortex for 3 animals using activity in R1 (top row) and R2 (bottom row) as seeds,
during spontaneous activity and during the BMI task. Mouse 3 was part of a separate
cohort trained on an auditory task (see Methods). Animals could decorrelate normally
correlated areas for task performance.   **C.** Animals took longer to reach criterion
performance (50% hits/attempt) if control regions were highly spontaneously correlated
(data from 7 mice, 9 days of training starting on day 4). **D.** The average variance of
activity for R1 (top panel) and R2 (bottom panel) was greater around hits early in a
training session than late in the session (n = 7 mice, day 8 of training, shading indicates
95% confidence interval around mean) indicating that mice honed in on more reliable
and reproducible control strategies over the course of a single training session. **E.** Early
in the session, neural activity around the control regions had high spectral entropy (a
proxy of signal randomness) as animals used stochastic bursts of activity to explore the
neural patterns that would yield reward. By late in the session, animals had discovered a
successful activity pattern to exploit, and spectral entropy in control area activity
decreased. Shaded area indicates 95% confidence interval around mean.

The variance in R1 and R2 activity peaked around hits early in a training session as

animals explored strategies that would yield reward, as has been shown previously for

the activity of individual cells learning to control a BMI device using a fixed decoder

(Zacksenhouse et al., 2007; Athalye et al., 2017). This variance decreased later in the

session as animals discovered and exploited reliable, reproducible strategies (Figure

3D). Early in the session, the spectral entropy of activity (a measure of the spectral

power distribution of a signal, and a proxy for its complexity, see Methods) around the

control regions increased as animals explored activity patterns that would yield reward,

then dramatically dropped as they discovered successful patterns and reliably exploited

them (Figure 3E). At the start of a session, or when control regions were changed,

animals faced with uncertain task rules 'explored' activity space through stochastic

bursting, and gradually switched to effecting stereotyped patterns they could reliably

exploit after having probed the rules of their environment (Tervo et al., 2014).

Expert performance correlated with increased activity in higher visual areas

We sought to determine what cortical areas were most active during task performance,

and how this changed over learning. On the first day of training, primary visual cortex

was most active during the task, but as animals became expert in the task, higher visual

areas were recruited (Figure 4A-C): in particular, anteromedial cortex (AM),

posteriomedial cortex (PM), and rostrolateral visual cortex (RL), similar to previous studies of learning visually-guided tasks in mice (Wekselblatt et al., 2016; Orsolic et al., 2019). Areas AM and RL are considered by some to be parietal cortex homologues in mouse cortex, though there is no widespread agreement on this point (Harvey et al., 2012; Licata et al., 2017; Mohan et al., 2018; Pho et al., 2018; Lyamzin and Benucci, 2019). After the final day of training, animals were shown playback of the cursor positions using their previous task performance (hereafter referred to as the passive playback condition). Activity in these higher areas was not evident in animals passively watching the cursor, suggesting their recruitment was specific to goal-oriented task engagement (Figure 4). A separate cohort of animals trained using an auditory feedback cursor had variable task-active areas (Supp. Figure 4A, n = 4 mice), but, as with the visual task, higher activity was also evident in RL, which is a multimodal sensory associative area with neurons responsive to visual, touch and auditory stimuli (Mohan et al., 2018).

Figure 4: Higher visual areas were more active during expert task performance

A. Activation maps for individual animals on day 1 of training (top row), when animals were naïve, versus day 9 of training (bottom row), when animals performed the task expertly, calculated using the normalised activity difference for task-on minus task-off periods. Each map has been registered to the Allen Brain Atlas (overlaid) using stereotaxic marks. Control regions are shown as slightly larger than they actually were for better visibility. **B.** Activation maps during task performance on day 1, day 9, and during passive playback of a previous session (representing the normalised

activity difference for task-on minus task-off periods). Each map has been registered to the Allen Brain Atlas by stereotaxic marks and then averaged across 7 mice. **C.** The relative ratio of task-activation in higher visual areas versus V1 increased over training. When animals passively viewed playback of the same session, higher visual areas were less active. Red bars indicate mean ratios (n=7 mice, paired t-test, Bonferroni corrected).

Population tuning of neurons shifted towards target position

Having identified brain areas implicated in BMI control, we recorded spiking from

individual cells while animals performed the task, in order to investigate the task-

dependent increase in calcium signals with cellular resolution. We chose to record from

functionally identified area AM, due to its recruitment over learning, and used multi-

channel silicon probes to record spikes from individual neurons while simultaneously

imaging the rest of dorsal cortex (Figure 5A, see Methods, (Xiao et al., 2017; Clancy et

al., 2019; Barson et al., 2018; Peters et al., 2019)). We obtained 16-49 single units per

recording, spanning all cortical layers. Units could be classified as regular-spiking (RS)

or fast-spiking, putative interneurons (FS) depending on spike width (see Methods,

Figure 5A). We recorded from 131 units in 7 mice performing the task (example

recording shown in Figure 5B), and 128 units from the same animals passively viewing a

playback of the previous training session's cursor positions. Due to electrode shift over

the recording sessions, we do not have recordings of the same units across both

conditions, thus our analyses are limited to population response differences. During the

task, population firing was significantly increased for cursor positions closer to the target,

and this was true both of FS and RS units (Figure 5C-E, Supp. Figure 5A-C). The target-

dependent increased spiking could not be explained by reward expectation, as a subset

of animals were given reward at the target position during passive playback as well

(Supp. Figure 5C).

Figure 5. Cursor tuning shifts to target location
A. Electrophysiological recordings were targeted to AM while animals were performing the imaging-based BMI task, with control regions in anterior motor cortex. Inset shows example waveforms from three isolated units (fast spiking unit in red). **B.** Example spiking during three successful trials (trials denoted in blue, hits denoted with pink arrows). At top, traces, from top to bottom, are: pupil diameter, running velocity, and visual cursor position. **C.** Normalised tuning to cursor positions for all single units. Each row represents the normalised firing responses to each of eight cursor positions for every recorded unit in the task (top, N = 131 units) and playback (bottom, N = 128 units) conditions. Firing responses were taken as the average firing rate for a period from 80 to 200 ms from the onset of the cursor presentation. **D.** Average population firing rates for each cursor position during task performance (red) and passive playback (blue). Shaded regions indicate 95% confidence levels. **E.** (Left) Mean firing rate for regular spiking (RS) units to different cursor positions during task performance (95% confidence interval indicated by shading, n = 7 mice). (Right) Mean firing rate for fast spiking (FS) units to different cursor positions during task performance (95% confidence interval indicated by shading, n = 7 mice).

Responses were dependent on the preceding cursor position—firing to the cursor positions closest to the target was higher if the cursor swept towards the target, but lower if the cursor swept away from it. The opposite was true for the cursor positions

farther from the target: firing was higher if the cursor swept away from the target (Figure
6A-B). We surmised this might allow downstream areas to more effectively decode the
cursor identity for more effective behavioural performance, and so performed a classifier
analysis on the population neural responses (Meyers, 2013). Indeed, the cursor identity
could be much more effectively decoded from AM neural responses during task
performance vs. playback (Figure 6C). During the task, the classifier could perform
above chance (12.5%) even before the cursor was present, suggesting that the neural
responses also encode intention or expectation. In the passive playback condition, the
classifier only performed above chance after the cursor had been presented. However,
the classifier failed to decode the direction that the cursor was traveling in (towards or
away from the target) in either the task or playback conditions (Figure 6D).

Figure 6. Cursor identity was more decodable from neural activity during task performance

A. Mean firing rate to different cursor positions depended on whether the preceding cursor position was sweeping towards (red) or away (black) from the direction of the target during task (t-test, Bonferroni corrected, * indicates p<0.05, ** indicates p<0.01). **B.** Cursor sweep direction had little effect on firing rates when animals were passively viewing playback. **C.** Classification accuracy for cursor identity, from population responses trained on real (blue) and shuffled data (grey), for neural responses during task performance (left panel) and passive playback (right panel). Shaded regions denote 95% confidence interval. During task performance, the classifier could infer the upcoming cursor position even before presentation (chance level at 12.5%), and rose higher after presentation, suggesting that neural responses encode expectation. This was not true of the passive playback condition, in which case the classifier only performed above chance during the cursor presentation period. **D.** Classification

accuracy for cursor direction, from population responses trained on real (blue) and shuffled labels data (grey), for neural responses during task performance (left panel) and passive playback (right panel). Shaded regions denote 95% confidence interval. A trained classifier could not perform above chance (50%) in predicting, from firing alone, the direction that the visual cursor was moving, in either the task or playback conditions.

Pupil diameter and running velocity were significantly decorrelated during task

performance compared to playback (Supp. Figure 5F), suggesting distinct mechanisms

underlie pupil size modulation in the two conditions. At the population level, firing rates

were uncorrelated with both pupil diameter and pupil position during the task, while firing

rates were weakly correlated with pupil diameter during playback (Supp. Figure 5G-H).

The fluorescence activity of the control regions was also uncorrelated with running

velocity and pupil diameter (Supp. Figure 5I-J).

To understand the relationship between the firing rate of individual neurons and dorsal

cortex-wide activity, we correlated the spike trains of individual units with fluorescence

activity across the brain to build affiliation maps for each unit (Xiao et al., 2017; Clancy et

al., 2019; Barson et al., 2018; Peters et al., 2019). We aligned these maps to the

common coordinate framework of the Allen Brain Atlas using stereotaxic marks on the

skull, and sorted these maps by units' preference for different cursor locations. During

task performance, the cells most responsive to the target and target-adjacent cursor

position were significantly more correlated with activity across the dorsal cortex (Supp.

Figure 6A). This could mean that the boosting around the target might be the result of a

cortex-wide signal, or that units tuned to the target correlate more strongly with the rest

of dorsal cortex during task performance, but not playback (Supp. Figure 6B-C).

Discussion

The idea of representation is fundamental to the idea of computation, and the neocortex

appears to employ hierarchies of transformed representations. It has become

increasingly clear that cortical sensory representations are not strictly veridical

reproductions of the outside world, but are shaped by a subject's internal states and

goals. We sought to understand how having causal control over an external object

affects the cortical sensory representation of that object, given that fluent control must be
informed by a constant dialogue between action and sensation (Haggard, 2017).
Animals learned to causally control an external object using neural calcium signals
recorded by widefield imaging, and did so by discovering and exploiting experimentally-
defined mappings between their neural activity and the visual feedback that led to
reward. This technique enabled us to identify cortical areas involved in task performance
and then to target recordings from individual cells in these areas while animals were
engaged in the task. We found that higher visual cortical areas, including area AM, were
recruited during expert BMI control, and that single units in AM encoded the same visual
cursor differently depending on whether the animal was causally controlling it, or
passively viewing it. These results lends evidence to the idea that activity in AM, a
potential homolog of parietal cortex, encodes a subject's intention and self-monitoring of
sensorimotor transformations (Andersen and Buneo, 2002; Andersen and Cui, 2009;
Desmurget et al., 2009; Aflalo et al., 2015; Cui, 2016).
423
Previous work indicates that subjects can learn to control neuroprosthetic devices using
single cells or bulk electrophysiological signals (Fetz, 1969; Bakay and Kennedy, 1998;
Nicolelis, 2001; Serruya et al., 2002; Carmena et al., 2003; Weiskopf et al., 2003;
Sitaram et al., 2007; Koralek et al., 2012; Hochberg et al., 2012; Collinger et al., 2013;
Clancy et al., 2014; Sadtler et al., 2014; Prsa et al., 2017; Sitaram et al., 2017;
Trautmann et al., 2019), but this is the first work, to our knowledge, to employ control
using population calcium signals. This technique allowed us to monitor much of the
dorsal cortical network as animals learned neuroprosthetic control, whereas previous
BMI work has been limited to recording from neighbouring neurons (Koralek et al., 2012;
Clancy et al., 2014; Sadtler et al., 2014; Prsa et al., 2017). This lends evidence to the
idea that using aggregate population signals (for example, from infrared imaging
(Shevelev, 1998; Abdelnour and Huppert, 2009)), rather than electrophysiological
recordings from individual neurons to manipulate neuroprosthetic devices might afford
more stable and minimally invasive control, robust to losing signals from, or damage to,
individual control cells. We also found that the spontaneous activity correlations between
the control regions, and not their spatial distance, was predictive of the animals' fluency
of performance, instructive for BMI design considerations, and similar to what has been
found in other work assessing the limits of BMI learning (Sadtler et al., 2014; Clancy et
al., 2014; Oby et al., 2019).

In order to learn the arbitrary action-outcome relationships required to perform BMI
tasks, animals must match internally generated actions or activity with their sensory
consequences. To probe how animals learned these contingencies, we changed the
regions that controlled the BMI between and within training sessions: meaning that
animals could not rely on a habitual activity pattern or strategy, but had to continually
explore different neural patterns to achieve reward on different training days. Animals did
so by 'exploring' with highly variable neural activity patterns early in a training session,
until they discovered a successful activity pattern they could reliably exploit (Figure 3).
Target hit rates dropped when reward was dispensed randomly, unlinked to the target
zone, indicating the animals' task performance was goal-directed, and not habitual.

Animals could decorrelate normally correlated brain areas during task execution, even
though spontaneous activity was widely correlated across cortex in mice not performing
the task, in agreement with previous work (Ledochowitsch et al., 2013). We found this to
be true of both anterior and posterior cortical areas—in mice trained to control an
auditory cursor, for example, posterior visual control areas could also be decorrelated
during the task (Figure 3B). However, animals performed better when control regions
were less spontaneously correlated (Figure 3C). Pupil and locomotion also became
significantly decorrelated during task performance, indicating that task-engagement and
locomotion may engage distinct arousal mechanisms (Supp. Figure 5, Vinck et al., 2015;
Reimer et al., 2016; Clancy et al., 2019).

By imaging dorsal cortex as animals performed this task, we were able to screen for
cortical areas engaged during expert BMI control. On the first day of training, V1 was
most active during the task, but as animals learned the task over days, higher visual
areas, including AM, PM and RL, became more active as animals controlled the cursor.
When the same cursors were played back to animals in an open-loop fashion (e.g. not
controlled by animals), activation was again mainly evident in V1, suggesting these
higher areas were involved in the goal-directed aspect of task performance. AM and RL
are considered by some to be rodent homologues of parietal cortex (Glickfeld and Olsen,
2017; Wang et al., 2011), which has been shown in humans to be involved in intention
and monitoring the mapping between action and outcome (Andersen and Buneo, 2002;
Desmurget et al., 2009; Andersen and Cui, 2009; Aflalo et al., 2015; Cui, 2016).
However, it was unclear whether the recruitment of these higher visual areas over

learning was related to the fact that they are involved with sensorimotor transformations generally, or because they're involved more specifically in planning and intentional control.

In order to clarify the role of these higher visual areas in causal control, we targeted extracellular recordings in one of the functionally identified task-active areas, AM, to probe task-related changes in the spiking of single units. We found that units were more active during task performance than during passive cursor playback of the animal's recent performance. In particular, units were more responsive to the cursor when it was at the target, and target-adjacent, positions during the BMI task compared to passive playback, similar to spatial attention boosting evident in previous work (Moran and Desimone, 1985; Engel et al., 2016), and in accordance with evidence that attention can reshape stimulus representations in a manner that more effectively guides decisions (Ruff and Cohen, 2019). Indeed, we found that a classifier trained on neural responses to the different cursor positions could effectively classify responses to these visual cursors, and could do so with significantly better accuracy when the animal was actively engaged in the task versus passively viewing a playback of the same cursors. In the task condition, the classifier ramped up in accuracy even before the cursor was presented, suggesting that neural responses were prospective, but whether this represents an intention (e.g. a control signal) or an expectation (e.g. a passive prediction) remains to be determined in future work. This boosting did not reflect reward expectation, and the animal did not use saccades or overt movements to perform the task. This boosting was also sensitive to the task goal: if the cursor was positioned close to the target and sweeping towards it, responses were boosted relative to when it was sweeping away. If the cursor was far from the target and sweeping farther away from it, responses were also boosted relative to the cursor at the same position sweeping towards the target. This suggests that firing rates reflect intention, and may also reflect a valence of the animal's perceived fluency of cursor control—that is, whether it was successfully or unsuccessfully guiding the cursor towards its intended goal (Lee and Dan, 2012).

We present a novel task and imaging method for exploring the encoding of action-outcome assessments, which allowed us to simultaneously monitor–with both dorsal cortex-wide and cell-level resolution–what activity patterns support the causal control of external objects. We believe imaging-based BMIs can serve as a useful paradigm for

studying the cortical dynamics involved in flexible learning, affording the experimenter control of what is learned, and by what areas or cell types. This makes it an excellent system for studying how the brain deals with sensory and rule manipulations, as well as a promising platform for studying credit assignment. It's also a promising proof-of-principle for exploring the potential–and limitations–of future imaging- and population-activity-based BMIs. While using widefield imaging afforded us a view of the dorsal cortex as animals learned neuroprosthetic control, there are a number of limitations to this method. While widefield fluorescence signals largely reflects neural firing, there is also a contribution from neuropil, as well as hemodynamic effects that were uncorrected in this study, as in our previous work we found it did not significantly impact our findings (Clancy et al., 2019). However, there is clearly some component of the signal contributed by blood vessels (e.g. in Figure 3B), so this is a limitation of the study.

Another limitation is that widefield imaging is limited to recording from dorsal cortex. Cortical neuroprosthetic control requires interactions with basal ganglia (Koralek et al., 2012; Neely et al., 2018), from which we cannot record using this method. Furthermore, we know from work in humans that prefrontal cortex (PFC) is involved in the sense of control over external stimuli, but we cannot record signals from PFC using this preparation due to its obscuration by the frontal sinus.

While we found increased evoked spiking to a visual cursor in the target location using this preparation, we do not know the exact cellular or neuromodulatory mechanisms giving rise to this difference. We found task-related increases in activity emerge over learning specifically in AM, PM and RL, and are not present during passive visual playback or when visual feedback was randomised. However, as we only recorded neuronal responses in AM, we do not know if the changes in neural tuning in AM are specific to this area, nor whether they are specifically required for task performance. Future work may address this by using mesoscale 2-photon imaging to record from or manipulate activity in molecularly defined neural subpopulations in parietal and control areas during task performance.

Acknowledgements

The authors would like to thank Adrianne Zhong for help with the camera control code,
Ivana Orsolic for help with the widefield imaging setup, Lisa Hoermann for performing
surgeries for this project, and Sonja Hofer, Petr Znamenskiy, Spencer Wilson, Alex
Naka, and Adrei Khilkevich for manuscript feedback and discussions of this work. This
work was supported by the European Research Council (NeuroV1sion 616509 to T.D.M-
F.), Swiss National Science Foundation (SNSF 31003A_169802 to T.D.M.- F.), the
Wellcome Trust (090843/E/09/Z core grant to the Sainsbury Wellcome Centre), the
EMBO Long-term Fellowship (ALTF 1481-2014 to K.B.C.), the HFSP Postdoctoral
Fellowship (LT000414/2015-L to K.B.C.), and the Branco Weiss-Society in Science grant
(K.B.C.).

Author contributions

K.B.C. planned and executed the experiments and analysed the data. K.B.C. and
T.D.M.- F. wrote the manuscript.

Competing financial interests statement

The authors declare no competing financial interests.

STAR Methods

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to Kelly Clancy, the lead contact (k.clancy@ucl.ac.uk).

Materials availability

This study did not generate new unique reagents or mouse lines.

Data and code availability

The data and code used in this study are available from the lead contact upon reasonable request.

Experimental model and subject details

Mice

All experimental procedures were carried out in accordance with institutional animal welfare guidelines and licensed by the Swiss cantonal veterinary office. TRE-Gcamp6s mice (Wekselblatt et al., 2016) (Jackson laboratories, https://www.jax.org/strain/024742) were crossed with B6.CBA-Tg(Camk2a-tTA)1Mmay/DboJ mice (Jackson laboratories, JAX 007004), to drive the expression of gCamp6s in CamKII+ pyramidal neurons. Animals were housed in a facility using a reversed light cycle, and recordings were taken during their active period. Eleven female mice were trained on the task, and we took electrophysiological recordings from seven of these, ranging between P55-P75. All animals were healthy, had never undergone previous procedures, and ranged in weight from 15-19g. Sample sizes were not statistically determined, but were consistent with previous papers using related methodology (Clancy et al., 2019; Xiao et al., 2017). Animals were group housed with same-sex littermates in enriched environments (including running wheels, cardboard tubes and chewing toys).

Method details

Surgery

A week before training, mice were prepared for imaging. Animals were anaesthetised with a mixture of fentanyl (0.05 mg per kg), midazolam (5.0 mg per kg), and medetomidine (0.5 mg per kg). The animal's scalp was resected and a head plate was secured to the skull. Four stereotaxically placed marks were made to enable alignment of the imaged brain with the Allen Brain Atlas (http://mouse.brain-map.org/static/atlas) post hoc, using the Allen Brain API (brain-map.org/api/index.html). The exposed skull was cleaned and covered with transparent dental cement to avoid infection, and to cover the cut scalp edges (C&B Metabond). This was polished to enhance the transparency of the preparation. A custom-made 3D printed light shield was cemented to the skull and head plate to avoid light leaks from the visual feedback presented on two computer monitors.

Behavioral setup and recordings

The recording chamber was sound-isolated and shielded from outside light. Mice were head-fixed under the microscope and free to run on a Styrofoam running wheel (diameter = 20 cm, width = 12 cm). The animals' movements were recorded using a rotary encoder in the wheel axis (pulse rate 1000, Kubler). Two monitors were placed side by side in front of the mouse, angled towards one another (21.5" monitors, ~20 cm from mouse, covering ~100x70 degrees of visual space), similarly to the setup described in (Poort et al., 2015). A reward port was place in front of the animal, where reward delivery was triggered via pinch solenoid one second after target hit (NResearch) and animal licks were detected using a custom piezo element coupled to the spout. All behavioral data were recorded using custom MATLAB software and a PCI-6320 acquisition board (National Instruments).

On electrophysiological recording days, pupil recordings were taken by illuminating the animal's right eye with a custom IR-light source and imaging with a CMOS camera (DMK22BUC03, Imaging Source, 30 Hz) using custom MATLAB software. Pupil size was determined as described in (Orsolic et al., 2019): images were first smoothed with a 2-D gaussian filter and thresholded to low luminance areas. These thresholded regions

were then filtered by circularity and size to automatically detect the pupil region. Pupil
edges were detected using the canny method, and ellipses were iteratively fit to the
region, tasked to minimise the geometric distance between the area outline and the fit
ellipse using nonlinear least squares (MATLAB function fitellipse, Richard Brown). The
pupil diameter was taken to be the major axis of the ellipse, then normalised by animal.
Pupil recordings from one animal had to be discarded, as the video was not sufficiently
in focus.

Behavioral training

After recovery, mice were acclimatised to head fixation for a minimum of two days, and
started on food restriction. Awake animals were head-fixed under the microscope and
free to run on a Styrofoam wheel. A baseline of spontaneous activity was taken on every
training day (10-20 minutes) in order to estimate spontaneous hit rates. The decoder
was calibrated such that animals achieved ~25% performance on their first day. Two
small control regions were chosen for real-time read out. In the case of visual feedback
task, these were all located in primary and secondary motor cortex, avoiding ALM. In the
auditory feedback task, control regions were placed in posterior cortex, over visual and
retrosplenial areas. The placement of the two control regions was usually ipsilateral but
sometimes contralateral to each other. The same control regions were used for the first
few days of training, then changed from day to day, or within sessions, so that animals
did not learn a fixed control strategy (see Table 1).

Activity was imaged at 40 Hz and the mean fluorescence from each control region was
transmuted to the cursor's position on screen with a simple transform:

$$p(t) = A_1 F_{R1} - A_2 F_{R2} + B \qquad (1)$$

where p is the cursor position, F_{R1} and F_{R2} are the instantaneous fluorescence ($\Delta F/F$) of
control regions one and two, respectively, and A_1, A_2 and B are coefficients set based on
the daily spontaneous baseline recordings (minimum 10 minutes). P was rounded to the
nearest integer to determine the discrete cursor location. A1 and A2 were determined by
dividing the full dynamic range of each recorded area during the baseline by half the
number of cursor positions:

$$A_1 = [\max(F_{R1})\text{-}\min(F_{R1})]/4; \quad A_2 = [\max(F_{R2})\text{-}\min(F_{R2})]/4; \qquad (2)$$

B represents the activity baseline of both areas. The chance performance was then

assessed by running the baseline data through the decoder to estimate how often the

animal would have achieved the target with spontaneous activity.

ΔF/F was calculated using a moving baseline, set as the tenth percentile of points from

the preceding 20 seconds of data. The raw fluorescence was converted to ΔF/F using a

moving baseline of 5 minutes of activity. The display updated at approximated 10 Hz,

with a latency of 300 ms from camera to screen, measured using a photodiode placed

on one of the monitors (Thorlabs, PDA100A-EC). Activity in R1 would cause the cursor

to move towards the target location in the centre of the animal's visual field, while

increased activity in R2 would cause the cursor to move away from the target zone. The

cursor was presented on two monitors so that the animal could track the cursor with both

eyes; its goal was to move the cursors presented on the two screens on either side to

the middle of its visual field. These changes were binned, such that the cursor could take

one of eight possible locations on the screen. The cursor had to be held at the target

position for 0.3 seconds to count as a hit, at which point the cursor disappeared. When a

target was hit, a MATLAB-controlled Data Acquisition board (National Instruments,

Austin, TX) triggered the administration of a soyamilk reward following a 1 second delay.

The next trial could be initiated within 5 seconds of reward delivery, but only when the

activation of R1 relative to R2 returned to the mean value recording during spontaneous

activity (to ensure enough time had passed for large transients to decay, given slow

calcium dynamics). This was return to baseline condition was rarely triggered (~5% of

trials) and on average lasted under 2 seconds. If the animal did not bring the cursor to

the target within a 30 second trial, the cursor disappeared, and the animal received a

white noise tone and a 10 second 'time out.'

We trained a separate cohort of four mice using an auditory, rather than visual, feedback

cursor, where activity was transmuted to the pitch of a feedback tone (Clancy et al.,

2014). As with the visual feedback task, a spontaneous baseline was recorded every

day (10-20 minutes) to assess chance levels of performance and calibrate the decoder.

Activity from two arbitrarily chosen regions was entered into an online transform

algorithm that related neural activity to the pitch of an auditory cursor:

$$f(t) = A_1 e^{F_{R1}1} - A_2 e^{RF_{R2}} + B \qquad (3)$$

Where f is the cursor frequency, F_{R1} is the instantaneous $\Delta F/F$ of R1, F_{R2} the

instantaneous $\Delta F/F$ of R2, and A1, A2, and B are coefficients set based on the daily

baseline recording. As above, $\Delta F/F$ was calculated using a moving baseline, set as the

tenth percentile of points from the preceding 20 seconds of data. Linear changes in firing

rate resulted in exponential changes in cursor frequency, and frequency changes were

binned in quarter-octave intervals to match rodent psychophysical discrimination

thresholds. As with the visual task, a trial was marked incorrect if the target pitch was not

achieved within 30 seconds of trial initiation. The auditory feedback was played using

speakers mounted on 2 sides of the imaging platform.

Task and control conditions

In the **task condition**, the position of the presented cursor was determined by the

control regions' instantaneous activity, according to equation (1) above. Rewards were

given when the cursor hit the target zone (cursor position 8). Similarly, in the **random**

reward condition, intended to test whether animal's task engagement was goal-

directed or habitual, the position of the presented cursor was determined by the control

regions' instantaneous activity, according to equation (1) above. However, here rewards

were not linked to the cursor position, but given out at random time intervals at a rate

matched to an expertly performing animal (approximately 1.5 rewards/minute). In the

random feedback condition, intended to confirm that the animal was using the visual

feedback to inform its behaviour, the position of the presented cursor was not linked to

the control regions' instantaneous activity, but instead was drawn randomly from a

Gaussian distribution matching the mean and variance of a typical task condition. In the

passive playback condition, the presented cursor position was no longer linked to the

control regions' activity, but was purely a replay of the cursor positions from the training

session the animal had undergone previously. Thus, the cursor positions and timing of

cursor presentations were matched to those of the task condition.

Widefield imaging

Widefield imaging was performed through the intact skull using a custom-built

epifluorescence macroscope with photographic lenses in a face-to-face configuration

(85mm f/1.8D objective, 50mm f/1.4D tube lens; (Ratzlaff and Grinvald, 1991)). Data

were recorded using a CMOS camera (Pco.edge 5.5, PCO, Germany) in global shutter

mode. 16-bit images were acquired at a rate of 40 Hz and binned 2x2 online using

custom-made LABVIEW software. A constant illumination at 470 nm was provided

(M470L3, Thorlands, excitation filter FF02-447/60-25), with average power ~0.05

mW.mm2 (emission filter 525/50-25, Semrock). The imaging site was shielded from light

contamination using a 3D-printed blackout barrier glued to the animal's skull. Signals

from the two control regions were sent via UDP to a computer providing visual or

auditory feedback to the mouse, using custom MATLAB software.

Electrophysiological recordings

The day before recording, mice were anesthetised with isofluorane and a small

craniotomy was opened over AM, which was functionally identified during task

performance, and stereotaxically confirmed. The craniotomy was kept damp with

Ringer's solution and sealed with KwikSil (World Precision Instruments). Recordings

were taken on the following day to avoid residual effects of anesthesia.

On the recording day, animals were head-fixed under a custom-built widefield

microscope, the skull and cortex were cleaned with Ringer's solution, and the KwikSil

plug removed from the craniotomy. A custom-designed silicon probe (64 channels, 2

shanks, Neuronexus, as described in (Clancy et al., 2019)) was inserted at an angle of

~45 degrees from normal to cortex. The probe consisted of two shanks with 64 sites

total, organised into 16 'tetrodes', each consisting of 4 sites located 25 um apart from

each other within-tetrode, and tetrodes spaced 130 um apart from each other. A small

amount of KwikSil or agar was used to cover the exposed cortex after the probe was in

place. After allowing the probe to settle for 20-30 minutes, neural activity was recorded

using the OpenEphys recording system (Siegle et al., 2017). Behavioral and stimulation

data, including pulses representing each camera frame, were recorded using

OpenEphys, enabling the alignment of electrophysiological signals with imaging and

behavioral data. Ephys recordings were filtered between 700 and 7000 Hz, and spikes

detected using the Klustakwik suite (Schmitzer-Torbert et al., 2005). Clusters were

assigned to individual units by manual inspection, excluding any units without a clear

refractory period. Units were separated into fast and broad spiking units by their peak-to-

trough time, using a cutoff of 0.66 ms (Barthó et al., 2004).

Data analysis

Raw imaging data were checked for dropped frames, spatially binned 2x2, and loaded
into MATLAB as a mapped tensor (Muir and Kampa, 2015). The raw fluorescence was
converted to ΔF/F using a moving baseline, calculated as the tenth percentile of points
from the preceding 20 seconds of data. We did not perform hemodynamic correction as
previous work indicates that hemodynamic and flavoprotein signals contribute minimally
compared to the calcium responses (Vanni and Murphy, 2014; Xiao et al., 2017; Clancy
et al., 2019).

Task-activation maps were calculated by taking the normalised average of fluorescence
movies during the task, or visual cursor playback period, subtracted by periods when
animals were not performing the task or viewing any visual stimuli (including periods of
spontaneous activity, and reward collection). To ensure that differences between early
and late in training were not influenced by possible differences in the statistics of the
visual feedback cursor, we randomly excluded success trials on late training days in
order to have comparable numbers of success and failure trials between early and late
training, however, including or excluding these trials did not influence the result. To build
the single-unit affiliation maps (Supp. Figure 6, see also (Clancy et al., 2019)), spike
trains were binned to match imaging frames, and maps were calculated by taking the
correlation of each unit's spike train with each pixel's ΔF/F.

Spectral entropy was calculated in 10 second windows, each overlapping by 5 seconds.
The calcium signal of the control areas was transformed into power spectral density
(PSD) during these windows (the magnitude squared of the signal's Fourier transform).
This was then used to calculate the spectral entropy for that time span:

$$SE = - \sum_{f=\frac{-f_s}{2}}^{f=\frac{f_s}{2}} PSD_n(f) log_2[PSD_n(f)]$$

794 Where SE is the spectral entropy, PSD_n is the normalised PSD, and *f* is frequency.

Classifier analysis

Spike data were binned into 50 ms bins, and split into 40 segments for training: 39 of these splits were used for training the classifier and 1 was used for testing. The data were z-score normalized so that high firing rate units didn't bias the classifier results. This data was then used, along with the cursor position labels, to train the classifier: a mean vector was created for each cursor position class based on the training data, and predictions were made on the test data by choosing the label class with the maximum correlation between the test and training mean vectors. These predicted labels were compared with the true labels to generate an average classifier accuracy over each tested time bin. This process was repeated 20 times using different training/test splits to cross-validate the results. The final reported classification accuracy is the mean of these 20 runs.

Quantification and statistical analysis

All data were analysed using custom code in MATLAB. The statistical tests used in our analyses are indicated in the figure legends, which also includes the value of n, and whether it refers to animals or single units. Differences were tested using Student's t-test, and Bonferroni corrected where apropriate. P-values are reported in the figures as well as legends, and significance is herein defined as $p < 0.05$. The pupil video for one animal, taken on the final day of recording, had to be excluded as it was too out of focus to determine the pupil size. Sample sizes were not statistically determined, but were consistent with previous papers using related methodology (Clancy et al., 2019; Xiao et al., 2017). Data were assumed to be normal but this was not systematically tested. Animals were trained in three separate cohorts (the first, a cohort of four animals trained on the auditory feedback task, the second, a cohort of 5 animals trained on the visual feedback task, and the third, a cohort of 2 animals trained on the visual feedback test) as an internal replication.

References

References

Abdelnour, A.F., and Huppert, T. (2009). Real-time imaging of human brain function by
near-infrared spectroscopy using an adaptive general linear model. NeuroImage *46*, 133–
143.

Aflalo, T., Kellis, S., Klaes, C., Lee, B., Shi, Y., Pejsa, K., Shanfield, K., Hayes-Jackson,
S., Aisen, M., Heck, C., et al. (2015). Neurophysiology. Decoding motor imagery from
the posterior parietal cortex of a tetraplegic human. Science *348*, 906–910.

Andersen, R.A., and Buneo, C.A. (2002). Intentional maps in posterior parietal cortex.
Annu. Rev. Neurosci. *25*, 189–220.

Andersen, R.A., and Cui, H. (2009). Intention, Action Planning, and Decision Making in
Parietal-Frontal Circuits. Neuron *63*, 568–583.

Andersen, R.A., Snyder, L.H., Bradley, D.C., and Xing, J. (1997). Multimodal
representation of space in the posterior parietal cortex and its use in planning movements.
Annu. Rev. Neurosci. *20*, 303–330.

Athalye, V.R., Ganguly, K., Costa, R.M., and Carmena, J.M. (2017). Emergence of
Coordinated Neural Dynamics Underlies Neuroprosthetic Learning and Skillful Control.
Neuron *93*, 955-970.e5.

Bagur, S., Averseng, M., Elgueda, D., David, S., Fritz, J., Yin, P., Shamma, S.,
Boubenec, Y., and Ostojic, S. (2018). Go/No-Go task engagement enhances population
representation of target stimuli in primary auditory cortex. Nat. Commun. *9*, 2529.

Bakay, R.A.E., and Kennedy, P.R. (1998). Cognitive Engineering: Proof of Principle.
Neurosurgery *43*, 706–706.

Barson, D., Hamodi, A.S., Shen, X., Lur, G., Constable, R.T., Cardin, J., Crair, M., and
Higley, M. (2018). Simultaneous mesoscopic and two-photon imaging of neuronal
activity in cortical circuits. BioRxiv 468348.

Barthó, P., Hirase, H., Monconduit, L., Zugaro, M., Harris, K.D., and Buzsáki, G. (2004).
Characterization of Neocortical Principal Cells and Interneurons by Network Interactions
and Extracellular Features. J. Neurophysiol. *92*, 600–608.

Carmena, J.M., Lebedev, M.A., Crist, R.E., O'Doherty, J.E., Santucci, D.M., Dimitrov,
D.F., Patil, P.G., Henriquez, C.S., and Nicolelis, M.A.L. (2003). Learning to Control a
Brain–Machine Interface for Reaching and Grasping by Primates. PLOS Biol. *1*, e42.

Chambon, V., Moore, J.W., and Haggard, P. (2015). TMS stimulation over the inferior
parietal cortex disrupts prospective sense of agency. Brain Struct. Funct. *220*, 3627–3639.

Churchland, A.K., Kiani, R., and Shadlen, M.N. (2008). Decision-making with multiple
alternatives. Nat. Neurosci. *11*, 693–702.

Clancy, K.B., Koralek, A.C., Costa, R.M., Feldman, D.E., and Carmena, J.M. (2014).
Volitional modulation of optically recorded calcium signals during neuroprosthetic
learning. Nat. Neurosci. *17*, 807–809.

Clancy, K.B., Orsolic, I., and Mrsic-Flogel, T.D. (2019). Locomotion-dependent
remapping of distributed cortical networks. Nat. Neurosci.

Collinger, J.L., Wodlinger, B., Downey, J.E., Wang, W., Tyler-Kabara, E.C., Weber,
D.J., McMorland, A.J., Velliste, M., Boninger, M.L., and Schwartz, A.B. (2013). High-
performance neuroprosthetic control by an individual with tetraplegia. The Lancet *381*,
557–564.

Cui, H. (2016). Forward Prediction in the Posterior Parietal Cortex and Dynamic Brain-
Machine Interface. Front. Integr. Neurosci. 10:35.

Desmurget, M., Reilly, K.T., Richard, N., Szathmari, A., Mottolese, C., and Sirigu, A.
(2009). Movement intention after parietal cortex stimulation in humans. Science *324*,
811–813.

Dewey, J. (1896). The reflex arc concept in psychology. Psychol. Rev. *3*, 357–370.

Donoghue, J.P. (2002). Connecting cortex to machines: recent advances in brain
interfaces. Nat. Neurosci. *5 Suppl*, 1085–1088.

Engel, T.A., Steinmetz, N.A., Gieselmann, M.A., Thiele, A., Moore, T., and Boahen, K.
(2016). Selective modulation of cortical state during spatial attention. Science *354*, 1140–
1144.

Fetz, E.E. (1969). Operant conditioning of cortical unit activity. Science *163*, 955–958.

Glickfeld, L.L., and Olsen, S.R. (2017). Higher-Order Areas of the Mouse Visual Cortex.
Annu. Rev. Vis. Sci. *3*, 251–273.

Gnadt, J.W., and Andersen, R.A. (1988). Memory related motor planning activity in
posterior parietal cortex of macaque. Exp. Brain Res. *70*, 216–220.

Haggard, P. (2017). Sense of agency in the human brain. Nat. Rev. Neurosci. *18*, 196–
207.

Harvey, C.D., Coen, P., and Tank, D.W. (2012). Choice-specific sequences in parietal
cortex during a virtual-navigation decision task. Nature *484*, 62–68.

Hochberg, L.R., Bacher, D., Jarosiewicz, B., Masse, N.Y., Simeral, J.D., Vogel, J.,
Haddadin, S., Liu, J., Cash, S.S., Smagt, P. van der, et al. (2012). Reach and grasp by
people with tetraplegia using a neurally controlled robotic arm. Nature *485*, 372–375.

Johnson, P.B., Ferraina, S., Bianchi, L., and Caminiti, R. (1996). Cortical networks for
visual reaching: physiological and anatomical organization of frontal and parietal lobe
arm regions. Cereb. Cortex N. Y. N 1991 *6*, 102–119.

Kahnt, T., Park, S.Q., Haynes, J.-D., and Tobler, P.N. (2014). Disentangling neural
representations of value and salience in the human brain. Proc. Natl. Acad. Sci. *111*,
5000–5005.

Koralek, A.C., Jin, X., Long, J.D., Costa, R.M., and Carmena, J.M. (2012).
Corticostriatal plasticity is necessary for learning intentional neuroprosthetic skills.
Nature *483*, 331–335.

Lacquaniti, F., Guigon, E., Bianchi, L., Ferraina, S., and Caminiti, R. (1995).
Representing spatial information for limb movement: role of area 5 in the monkey.
Cereb. Cortex N. Y. N 1991 *5*, 391–409.

Ledochowitsch, P., Koralek, A.C., Moses, D., Carmena, J.M., and Maharbiz, M.M.
(2013). Sub-mm functional decoupling of electrocortical signals through closed-loop
BMI learning. Conf. Proc. Annu. Int. Conf. IEEE Eng. Med. Biol. Soc. IEEE Eng. Med.
Biol. Soc. Annu. Conf. *2013*, 5622–5625.

Lee, S.-H., and Dan, Y. (2012). Neuromodulation of brain states. Neuron *76*, 209–222.

Licata, A.M., Kaufman, M.T., Raposo, D., Ryan, M.B., Sheppard, J.P., and Churchland,
A.K. (2017). Posterior Parietal Cortex Guides Visual Decisions in Rats. J. Neurosci. Off.
J. Soc. Neurosci. *37*, 4954–4966.

Lyamzin, D., and Benucci, A. (2019). The mouse posterior parietal cortex: Anatomy and
functions. Neurosci. Res. *140*, 14–22.

Makino, H., Ren, C., Liu, H., Kim, A.N., Kondapaneni, N., Liu, X., Kuzum, D., and
Komiyama, T. (2017). Transformation of Cortex-wide Emergent Properties during Motor
Learning. Neuron *94*, 880-890.e8.

Meyers, E. (2013). The neural decoding toolbox. Front. Neuroinformatics *7*.

Mohan, H., Gallero-Salas, Y., Carta, S., Sacramento, J., Laurenczy, B., Sumanovski,
L.T., Kock, C.P.J. de, Helmchen, F., and Sachidhanandam, S. (2018). Sensory
representation of an auditory cued tactile stimulus in the posterior parietal cortex of the
mouse. Sci. Rep. *8*, 7739.

Mohan, H., de Haan, R., Broersen, R., Pieneman, A.W., Helmchen, F., Staiger, J.F.,
Mansvelder, H.D., and de Kock, C.P.J. (2019). Functional Architecture and Encoding of
Tactile Sensorimotor Behavior in Rat Posterior Parietal Cortex. J. Neurosci. Off. J. Soc.
Neurosci. *39*, 7332–7343.

Moran, J., and Desimone, R. (1985). Selective attention gates visual processing in the
extrastriate cortex. Science *229*, 782–784.

Mountcastle, V.B., Lynch, J.C., Georgopoulos, A., Sakata, H., and Acuna, C. (1975).
Posterior parietal association cortex of the monkey: command functions for operations
within extrapersonal space. J. Neurophysiol. *38*, 871–908.

Muir, D.R., and Kampa, B. (2015). FocusStack and StimServer: A new open source
MATLAB toolchain for visual stimulation and analysis of two-photon calcium neuronal
imaging data. Front. Neuroinformatics *8*.

Musall, S., Kaufman, M.T., Juavinett, A.L., Gluf, S., and Churchland, A.K. (2019).
Single-trial neural dynamics are dominated by richly varied movements. Nat. Neurosci.
22, 1677–1686.

Neely, R.M., Koralek, A.C., Athalye, V.R., Costa, R.M., and Carmena, J.M. (2018).
Volitional Modulation of Primary Visual Cortex Activity Requires the Basal Ganglia.
Neuron *97*, 1356-1368.e4.

Nicolelis, M.A. (2001). Actions from thoughts. Nature *409*, 403–407.

Oby, E.R., Golub, M.D., Hennig, J.A., Degenhart, A.D., Tyler-Kabara, E.C., Yu, B.M.,
Chase, S.M., and Batista, A.P. (2019). New neural activity patterns emerge with long-
term learning. Proc. Natl. Acad. Sci. *116*, 15210–15215.

Orsolic, I., Rio, M., Mrsic-Flogel, T.D., and Znamenskiy, P. (2019). Mesoscale cortical
dynamics reflect the interaction of sensory evidence and temporal expectation during
perceptual decision-making. BioRxiv 552026.

Peters, A.J., Steinmetz, N.A., Harris, K.D., and Carandini, M. (2019). Striatal activity
reflects cortical activity patterns. BioRxiv.

Pho, G.N., Goard, M.J., Woodson, J., Crawford, B., and Sur, M. (2018). Task-dependent
representations of stimulus and choice in mouse parietal cortex. Nat. Commun. *9*, 2596.

Pisella, L., Gréa, H., Tilikete, C., Vighetto, A., Desmurget, M., Rode, G., Boisson, D.,
and Rossetti, Y. (2000). An 'automatic pilot' for the hand in human posterior parietal
cortex: toward reinterpreting optic ataxia. Nat. Neurosci. *3*, 729–736.

Poort, J., Khan, A.G., Pachitariu, M., Nemri, A., Orsolic, I., Krupic, J., Bauza, M.,
Sahani, M., Keller, G.B., Mrsic-Flogel, T.D., et al. (2015). Learning Enhances Sensory
and Multiple Non-sensory Representations in Primary Visual Cortex. Neuron *86*, 1478–
1490.

Prsa, M., Galiñanes, G.L., and Huber, D. (2017). Rapid Integration of Artificial Sensory
Feedback during Operant Conditioning of Motor Cortex Neurons. Neuron *93*, 929-
939.e6.

Ratzlaff, E.H., and Grinvald, A. (1991). A tandem-lens epifluorescence macroscope:
hundred-fold brightness advantage for wide-field imaging. J. Neurosci. Methods *36*, 127–
137.

Reimer, J., McGinley, M.J., Liu, Y., Rodenkirch, C., Wang, Q., McCormick, D.A., and
Tolias, A.S. (2016). Pupil fluctuations track rapid changes in adrenergic and cholinergic
activity in cortex. Nat. Commun. *7*, 13289.

Rizzolatti, G., Fogassi, L., and Gallese, V. (1997). Parietal cortex: from sight to action.
Curr. Opin. Neurobiol. *7*, 562–567.

Ruff, D.A., and Cohen, M.R. (2019). Simultaneous multi-area recordings suggest that
attention improves performance by reshaping stimulus representations. Nat. Neurosci. *22*,
1669–1676.

Rushworth, M.F., Ellison, A., and Walsh, V. (2001). Complementary localization and
lateralization of orienting and motor attention. Nat. Neurosci. *4*, 656–661.

Sadtler, P.T., Quick, K.M., Golub, M.D., Chase, S.M., Ryu, S.I., Tyler-Kabara, E.C., Yu,
B.M., and Batista, A.P. (2014). Neural constraints on learning. Nature *512*, 423–426.

Salkoff, D.B., Zagha, E., McCarthy, E., and McCormick, D.A. (2020). Movement and
Performance Explain Widespread Cortical Activity in a Visual Detection Task. Cereb.
Cortex N. Y. N 1991 *30*, 421–437.

Schmitzer-Torbert, N., Jackson, J., Henze, D., Harris, K., and Redish, A.D. (2005).
Quantitative measures of cluster quality for use in extracellular recordings. Neuroscience
131, 1–11.

Serruya, M.D., Hatsopoulos, N.G., Paninski, L., Fellows, M.R., and Donoghue, J.P.
(2002). Instant neural control of a movement signal. Nature *416*, 141–142.

Shevelev, I.A. (1998). Functional imaging of the brain by infrared radiation
(thermoencephaloscopy). Prog. Neurobiol. *56*, 269–305.

Siegle, J.H., Cuevas López, A., Patel, Y., Abramov, K., Ohayon, S., and Voigts, J.
(2017). Open Ephys: An open-source, plugin-based platform for multichannel
electrophysiology. J. Neural Eng.

Sitaram, R., Caria, A., Veit, R., Gaber, T., Rota, G., Kuebler, A., and Birbaumer, N.
(2007). fMRI Brain-Computer Interface: A Tool for Neuroscientific Research and
Treatment. Comput. Intell. Neurosci. *2007*.

Sitaram, R., Ros, T., Stoeckel, L., Haller, S., Scharnowski, F., Lewis-Peacock, J.,
Weiskopf, N., Blefari, M.L., Rana, M., Oblak, E., et al. (2017). Closed-loop brain
training: the science of neurofeedback. Nat. Rev. Neurosci. *18*, 86–100.

Stringer, C., Pachitariu, M., Steinmetz, N., Reddy, C.B., Carandini, M., and Harris, K.D.
(2019). Spontaneous behaviors drive multidimensional, brainwide activity. Science *364*.

Sugrue, L.P., Corrado, G.S., and Newsome, W.T. (2004). Matching behavior and the
representation of value in the parietal cortex. Science *304*, 1782–1787.

Tervo, D.G.R., Proskurin, M., Manakov, M., Kabra, M., Vollmer, A., Branson, K., and
Karpova, A.Y. (2014). Behavioral Variability through Stochastic Choice and Its Gating
by Anterior Cingulate Cortex. Cell *159*, 21–32.

Trautmann, E.M., O'Shea, D.J., Sun, X., Marshel, J.H., Crow, A., Hsueh, B., Vesuna, S.,
Cofer, L., Bohner, G., Allen, W., et al. (2019). Dendritic calcium signals in rhesus
macaque motor cortex drive an optical brain-computer interface. BioRxiv 780486.

Vanni, M.P., and Murphy, T.H. (2014). Mesoscale Transcranial Spontaneous Activity
Mapping in GCaMP3 Transgenic Mice Reveals Extensive Reciprocal Connections
between Areas of Somatomotor Cortex. J. Neurosci. *34*, 15931–15946.

Vinck, M., Batista-Brito, R., Knoblich, U., and Cardin, J.A. (2015). Arousal and
locomotion make distinct contributions to cortical activity patterns and visual encoding.
Neuron *86*, 740–754.

Wang, Q., Gao, E., and Burkhalter, A. (2011). Gateways of Ventral and Dorsal Streams
in Mouse Visual Cortex. J. Neurosci. *31*, 1905–1918.

Weiskopf, N., Veit, R., Erb, M., Mathiak, K., Grodd, W., Goebel, R., and Birbaumer, N.
(2003). Physiological self-regulation of regional brain activity using real-time functional
magnetic resonance imaging (fMRI): methodology and exemplary data. NeuroImage *19*,
577–586.

Wekselblatt, J.B., Flister, E.D., Piscopo, D.M., and Niell, C.M. (2016). Large-scale
imaging of cortical dynamics during sensory perception and behavior. J. Neurophysiol.
115, 2852–2866.

Wisniewski, D., Reverberi, C., Momennejad, I., Kahnt, T., and Haynes, J.-D. (2015). The
Role of the Parietal Cortex in the Representation of Task–Reward Associations. J.
Neurosci. *35*, 12355–12365.

Xiao, D., Vanni, M.P., Mitelut, C.C., Chan, A.W., LeDue, J.M., Xie, Y., Chen, A.C.,
Swindale, N.V., and Murphy, T.H. (2017). Mapping cortical mesoscopic networks of
single spiking cortical or sub-cortical neurons. ELife *6*.

Zacksenhouse, M., Lebedev, M.A., Carmena, J.M., O'Doherty, J.E., Henriquez, C., and
Nicolelis, M.A.L. (2007). Cortical Modulations Increase in Early Sessions with Brain-
Machine Interface. PLOS ONE *2*, e619.

Zapparoli, L., Seghezzi, S., Zirone, E., Guidali, G., Tettamanti, M., Banfi, G., Bolognini,
N., and Paulesu, E. (2020). How the effects of actions become our own. Sci. Adv. *6*,
eaay8301.

1041

1042

1043

1044

1045

1046 **Supplemental Information**

1047 The sensory representation of causally controlled objects

1048 Kelly B. Clancy and Thomas D. Mrsic-Flogel

1049

Supplemental Figure 1. Animals did not use gross motor movements to perform task (related to Figure 1)
A. Example fluorescence trace from R1 and R2 in pre-task spontaneous activity, and during task, indicating a clear change from spontaneous patterns. **B.** Analysis of activity

throughout the task indicated that activity in the two control regions generally became
decoupled during task performance, and not just around hits. Average ΔF/F in the
control regions triggered by bursts in R1 (top row) or R2 (bottom row), for the same three
example animals, for spontaneous activity (left) and during the training session (right).
The burst threshold was taken as the z-scored value of ΔF/F > 3, in order to capture
large events. **C.** Lick averaged around hits (top panel) and misses (bottom panel) on a
day in late training (n = 7 mice, grey traces indicate individual mice, black trace indicates
mean). Pink arrow indicates target hit (top panel) or time out (bottom panel), black arrow
indicates reward delivery (top panel). **D.** Animal's running speed, averaged around hits
(top panel) and misses (bottom panel) on a day in late training (n = 7 mice, grey traces
indicate individual mice, black trace indicates mean). Pink arrow indicates target hit (top
panel) or time out (bottom panel), black arrow indicates reward delivery (top panel). **E.**
Eye saccades averaged around hits (top panel) and misses (bottom panel) on final day
of training (n = 6 mice). Average movement is shown for both x and y directions of pupil
image. Pink arrow indicates target hit (top panel) or time out (bottom panel), black arrow
indicates reward delivery (top panel).
1070

Supplemental Figure 2: Animals exhibit quasi-intentional, specific neural activity control during task performance (related to Figure 2)

A. Hit rate over the course of a training session averaged over 7 mice for three training days (shaded area indicates s.e.m., 7 animals for each day). **B.** Once animals were performing above chance (from day 4 on), there was no significant difference in the time

it took animals to reach 50% performance whether they were using the same ROIs as
the previous day of training, or different ones (t-test, p = 0.4). On day 6, when ROIs were
switched mid-session, there was also no significant difference in the time it took, post-
switch, for the animal to recover their performance to 50% compared to animals using
the same or different control regions to the day before (t-test, p=0.3, 0.7 respectively,
each dot indicates performance on one training day). **C.** On the day control regions
were switched mid-session, there was no significant difference in performance between
animals before their control regions were switched, and their recovery after the switch.
Each line represents one mouse (n = 7 mice). **D.** Example trace of the difference
between control areas' activity during task (R1-R2) for one mouse, indicating an
increase in differential modulation over the course of a training session (left). These
modulations decreased when reward was provided randomly (right). **E.** The slope of the
difference in the two control regions fluorescence over training was positive during the
normal task condition (left, p < 0.01, light-weighted lines each represent one of n = 5
mice that underwent the randomized reward condition on day 8 of training, heavy line
represents mean) and negative over the course of the randomised reward condition
(right, p < 0.01, light-weighted lines each represent one of n = 5 mice that underwent the
randomized reward condition on day 8 of training, heavy line represents mean). **F.** (Left)
Example activity maps locked around reward delivery for three animals performing the
task. One second before the reward delivery marks the target hit, after which the cursor
disappeared during the 1 second wait time. After reward delivery, strong activity is
evident in frontal areas when animals are licking and touching the spout. (Right)
Example activity maps locked around reward delivery for three animals in the random
reward condition. Rewards were given randomly, unlinked to target hits, so the visual
cursor was usually present throughout the reward collection period. After reward
delivery, strong activity is evident in frontal areas.

Supplemental Figure 3. Animals could modestly decorrelate areal brain activity during task performance (related to Figure 3)

A. (Left panel) The distance between control regions did not have an effect on the time it took animals to achieve 50% performance (for 83 training sessions, including all days of above chance performance from all 7 animals, e.g. from day 4 on). (Right panel) The distance between control regions did not have an effect on final performance (for 83 training sessions, including all days of above chance performance from all 7 animals, e.g. from day 4 on). **B.** Correlations between R1 and R2 dropped modestly from spontaneous levels during task performance (104 control region pairs from 7 animals, on all training days: on average 15 days of training per mouse, paired t-test, p = 1e-6). **C.** Left panel: correlations between control regions and activity in V1 dropped between spontaneous activity and task performance, but correlations between control regions and primary somatosensory cortex (S1) were not significantly different between conditions. Areas were identified by registering imaged brain surface with Allen Brain Atlas by stereotaxic marks; 7 animals, for expert level performance on days 9 and 14 of training paired t-test, p = 0.01, p=0.1, respectively for V1, S1). **D.** Correlations between primary visual cortex and primary somatosensory cortex did not change between spontaneous

1120   activity and task performance (areas as identified by registering imaged brain surface
1121   with Allen Brain Atlas by stereotaxic marks; 7 animals, for expert level performance on
1122   days 9 and 14 of training paired t-test, p = 0.02).
1123

1124
Supplemental Figure 4. Expert animals had increased activity in higher visual
areas during task performance (related to Figure 4)
A. Left panel: example task-related activity maps from four animals trained on an
auditory version of the task. Right panel: mean task-related activity map indicating
involvement of RL, a lateral homolog of parietal cortex in mice and a multimodal
associative area (mean from n = 4 mice). Visual areas appear task-activated (despite
there being no visual feedback in this task) likely due to the fact that the control regions
were placed over visual areas. Control regions are shown as slightly larger than they
actually were for better visibility.
1134

1135
Supplemental Figure 5. Pupil and locomotion become uncoupled during task
(related to Figure 5)
A. Responses to the visual cursor at the 8 monitor positions for two single units recorded
during task performance. The red line denotes the time the cursor appeared; the trace
colour denotes cursor position. Here the spiking responses have been smoothed by
convolving with a gaussian kernel with standard deviation of 25 ms. **B.** Responses to the
visual cursor at the 8 monitor positions for two single units recorded during passive

playback. The red line denotes the time the cursor appeared; the trace colour denotes
cursor position. Here the spiking responses have been smoothed by convolving with a
gaussian kernel with standard deviation of 25 ms. **C.** Firing rates for units recorded in 2
animals where the target cursor position was rewarded during passive playback (blue
trace). The enhanced firing for the target-adjacent cursor positions during task
performance (red trace) remained, suggesting the boosting was not simply reward
expectation (95% confidence interval indicated, n = 2 mice, 23 units in the task condition
and 17 units in the passive playback condition). **D.** Average running velocity at each
cursor location during recordings during task (left) and playback (right), (n = 7 mice, final
day of recording). **E**. Normalized pupil diameter at each cursor location during task (left)
and playback (right), (n = 6 mice, final day of recording). **F.** Pupil diameter and running
speed were significantly decorrelated during task performance compared to playback
(reward collection periods and inter-trial intervals excluded, n = 6 mice, final day of
recording, each line represents correlations for one mouse.) **G.** The correlation between
single unit firing rate and pupil diameter was not different from zero during task, but
slightly higher than zero during playback (reward collection periods and inter-trial
intervals excluded, n = 6 mice, N = 105 units task, n= 122 units playback). **H.** The
correlation between single unit firing rate and pupil position was not different from zero
during task or playback conditions (reward collection periods and inter-trial intervals
excluded, n = 6 mice, N = 105 units task, n= 122 units playback). **I.** R1 (left panel) and
R2 (right panel) activity were not significantly correlated with running velocity, either
during task performance or passive playback (reward collection periods and inter-trial
intervals excluded, N = 7 mice, each line represents correlations for one mouse). The
correlation between R1 or R2 and run velocity during task and passive playback were
not significantly different from zero (t-test: for R1, p = 0.08, p=0.6, respectively for task
and playback; for R2: p = 0.08, 0.7 respectively for task and playback), nor significantly
different from each other (paired t-test, p = 0.4). **J.** R1 (left panel) and R2 (right panel)
activity were not significantly correlated with pupil diameter, either during task
performance or passive playback (reward collection periods and inter-trial intervals
excluded, N = 6 mice, t-test; each line represents correlations for one mouse). The
correlation between R1 or R2 and pupil diameter during task and passive playback were
not significantly different than zero (t-test, for R1: p = 0.9, 0.7 for task, playback,
respectively; for R2: p = 0.9, 0.8 for task, and playback, respectively), nor significantly
different from each other (paired t-test, p = 0.4).

1177

Supplemental Figure 6. Cortex-wide affiliations of units tuned to different cursor positions (related to Figures 5 & 6)
A. Spike trains for each unit were correlated with the activity of each pixel to build cortex-wide affiliation maps. These were sorted into bins based on which cursor position units were most responsive to. The average of these maps is shown in the top row for task performance, and the bottom row for passive playback. The reported N under each average map indicates the number of single units in that bin for that condition. **B.** Mean correlation of spiking activity with calcium activity across dorsal cortex, sorted by units' cursor preference for task and playback. **C.** The trend evident in **A** is not due to higher firing rates of target cursor-tuned units, when units were organized into quartiles of mean firing rates rather than cursor preference. The reported N under each average map indicates the number of single units in that bin for that condition.

1190

1191

	[M/L, A/P] (mm)	M1	M2	M3	M4	M5	M6	M7
Day 1	R1	-1.5, 1.5	-2.5, 2	-2, 1.5	-2.5, 2	-2.5, 2.5	-2, 1.5	-1.5, 1
	R2	-2.5, 1.5	-2.6, 1	-2.2, 1	-2.5, 0.5	-2.5, 2	-2.5, 2.5	-2.5, 2
Day 2	R1	-1.5, 1.5	-2.5, 2	-2, 1.5	-2.5, 2	-2.5, 2.5	-2, 1.5	-1.5, 1
	R2	-2.5, 1.5	-2.6, 1	-2.2, 1	-2.5, 0.5	-2.5, 2	-2.5, 2.5	-2.5, 2
Day 3	R1	-1.5, 1.5	-2.5, 2	-2, 1.5	-2.5, 2	-2.5, 2.5	-2, 1.5	-1.5, 1
	R2	-2.5, 1.5	-2.6, 1	-2.2, 1	-2.5, 0.5	-2.5, 2	-2.5, 2.5	-2.5, 2
Day 4	R1	-1.5, 1.5	-2.5, 2	-2, 1.5	-1, 3	-2.5, 2.5	-2, 1.5	-1.5, 1
	R2	-2.5, 1.5	-2.6, 1	-2.2, 1	-2, 1.5	-2.5, 2	-2.5, 2.5	-2.5, 2
Day 5	R1	-1.5, 1.5	-2.5, 2.3	-2, 1.5	-2.5, 2	-2.5, 2.5	-1, 1	-2, 1.5
	R2	-2, 1.5	-2.5, 1.5	-2.2, 1	-2, 1.5	-2.5, 2	-2, 1.3	-1.5, 1.5
Day 6**	R1	-1.5, 1.5	-2.5, 2.3	-2.5, 2.5	-2.5, 2	-2.5, 2.5	-1.5, 1	-2, 1.5
	R2	-2, 1.5	-2.5, 1.5	-2, 2	-2, 1.5	-2.5, 2	-2, 1.5	-1.5, 1.5
Day 7	R1	-1, 1.5	-2.5, 2.5	-2.5, 2.5	-2.5, 2	-2, 2.5	-2.5, 1.5	-2, 1.5
	R2	-1.2, 1.8	-0.5, 3	-2, 2	-1.5, 2.5	-1, 0.5	-2.5, 2	-1.5, 1
Day 8	R1	-0.8, 2.2	-1.7, 1.5	-1, 1.5	-2.5, 2	-2, 2.5	-2.5, 1.5	-2, 2
	R2	-1, 2	-1, 2	-1.5, 2	-1.5, 1.5	-1, 0.5	-2.5, 2	-2.5, 2
Day 9	R1	-1.2, 0.5	-2, 1.5	-1.5, 3	-2.5, 2	-2, 2.5	-2.5, 1.5	-1.5, 1.5
	R2	-2.2, 1	-1.5, 1	-1.5, 1.5	-1.5, 0.5	-1, 0.5	-2.5, 2	-2.5, 2
Day 10	R1	-1.5, 1	-2, 1.5	-2.5, 2.2	-2.5, 2	-2, 1.5	-2.5, 1.5	-1.5, 1.5
	R2	-1.2, 2	-1.5, 1	-1.5, 1.5	-1.5, 0.5	-1.5, 1.5	-2.5, 2	-1.7, 1.7
Day 11	R1	-1.5, 1	-2, 1.5	-2.5, 2	-2.5, 2	-2, 1	-2.5, 1.5	-1.5, 1.5
	R2	-1, +2	-1.5, 1	-1.5, 2	-1.5, 0.5	2, 1	-2.5, 2	-1.7, 1.7
Day 12	R1	-1.5, 1	-2.5, 2	-0.7, 0.5	-2.5, 2	-2, 1	-2, 1.5	-0.5, 0.5
	R2	-1, 1.5	-1.5, 1	-2, 2	-1.5, 0.5	2, 1	-2, 2.5	-1.5, 1.5
Day 13	R1	-1.5, 1	-1.2, 0.5	-1, 0.5	-1.5, 0.5	-2, 2	-1, 0.5	-0.5, 0.5
	R2	-1, 1.5	-1.5, 2	-2, 2	-2.5, 2	-1, 0.5	-2, 1	-1.5, 1.5
Day 14	R1	-1.5, 0.2	-1.2, 0.5	-1, 0.5	-1.5, 0.5	-1.5, 1.5		
	R2	-2, 2	-1.5, 2	-2, 2	-2.5, 2	-2, 2		
Day 15	R1	-1.5, 0.2	-1, 0.2	-1, 0.5	-1.5, 0.5	-1.5, 1.5		
	R2	-2, 2	-1.5, 2	-2, 2	-2.5, 2	-2, 2		
Day 16	R1		-1, 0.2	-1, 0.5	-1.5, 0.5	-1.5, 1.5		
	R2		-1.5, 2	-2, 2	-2.5, 2	-2, 2		
Day 17	R1					-1.5, 1.5		
	R2					-2, 2		

Supplemental Table 1. Control region coordinates (related to Figure 1)
Related to Figure 1B: approximate stereotaxic coordinates, in millimeters, of control
regions on all days of training, reported in [medial/lateral, anterior posterior] pairs, with
bregma = [0, 0], and coordinates to the left of, or posterior to, bregma reported as
negative. Day 6, indicated with **, was the day that control regions were switched
midway through the training session.

Supplemental Video 1. Activity averaged around hits (related to Figure 1)
Video of cortical activity averaged around hits. Time from hit (in seconds) indicated at
the top left, and control regions are indicated (region 1 with a blue box, and region 2 with
a green box). Animal performs task by sweeping activity laterally from region 1 towards
region 2 at the time of the hit. Lick and forepaw activity are then evident as animal
consumes reward.
1218
1219 **Supplemental Video 2. Activity averaged around hits (related to Figure 1)**
1220 Video of cortical activity averaged around hits. Time from hit (in seconds) indicated at
1221 the top, and control regions are indicated (region 1 with a blue box, and region 2 with a
1222 green box). Activity in region 1 increases compared to region 2 at the time of the hit. Lick
1223 and forepaw activity are then evident as animal consumes reward.
1224
1225
Supplemental Video 3. Discovery of successful activity patterns on a day of
training where new control regions were introduced (related to Figures 1 & 2)
(Top) Animals had to sufficiently increase activity in control region 1 (R1) relative to
region 2 (R2) to bring the visual cursor to position 8 for 300 ms to receive reward. Video
of continuous recordings from R1 (blue trace) and R2 (green trace) on a day of training
when new control regions were introduced. If the animal failed to do this within 30
seconds, it was considered a failure trial (only one of every 3 frames is shown). Cyan
vertical bars indicate trial starts; magenta vertical bars indicate a successful target hit;
black vertical bars indicate the end of a failure trial. Top traces in black indicate running
velocity (top trace) and lick bouts (second trace). The R1-R2 trace is indicated in the
bottommost black trace.
(Middle) Reconstruction of the visual feedback cursor position at the time in the
recording indicated by the vertical red bar in the top panel. The visual cursor
disappeared from the screens in between trials.
(Bottom) Activity in dorsal cortex as animal performs the task. The location of control
region 1 is denoted with a blue box, control region 2 with a green box.
1242