Pulmonary acini exhibit complex changes during postnatal rat lung development

David Haberthür[1], Eveline Yao[1], Sébastien F. Barré[1], Tiziana P. Cremona[1], Stefan A. Tschanz[1], and Johannes C. Schittny[1]

[1]Institute of Anatomy, University of Bern, Switzerland

October 29, 2020

Abstract

Pulmonary acini represent the functional gas-exchanging units of the lung. Due to technical limitations, individual acini cannot be identified on microscopic lung sections. To overcome these limitations, we imaged the right lower lobes of instillation-fixed rat lungs from postnatal days P4, P10, P21, and P60 at the TOMCAT beamline of the Swiss Light Source synchrotron facility at a voxel size of 1.48 µm. Individual acini were segmented from the three-dimensional data by closing the airways at the transition from conducting to gas exchanging airways. For a subset of acini (N=268), we followed the acinar development by stereologically assessing their volume and their number of alveoli. We found that the mean volume of the acini increases 23 times during the observed timeframe. The coefficients of variation dropped from 1.26 to 0.49 and the difference between the mean volumes of the fraction of the 20 % smallest to the 20 % largest acini decreased from a factor of 27.26 (day 4) to a factor of 4.07 (day 60), i.e. shows a smaller dispersion at later time points. The acinar volumes show a very large variation early in lung development and homogenize during maturation of the lung by reducing their size distribution by a factor of 7 until adulthood. The homogenization of the acinar sizes hints at an optimization of the gas-exchange region in the lungs of adult animals and that acini of different size are not evenly distributed in the lungs. This likely leads to more homogeneous ventilation at later stages in lung development.

Introduction

Lung development

Lung development starts with the formation of the two lung buds. During the embryonic stage (prenatal days E11-E13 in rats) the major airways and the pleura are formed. During the pseudoglandular stage (E13-E18.5) most of the remaining airway generations are formed by branching morphogenesis. Only few additional branches are added during the later stages (canalicular (E18.5–E20) and (E20 to postnatal day P4)) Schittny and Burri [2008]. Epithelial differentiation during the canalicular stage and angiogenetic activation of the

capillaries lead to the first functional air-blood barriers in the lung Schittny [2018]. The saccular stage represents an intermediate stage where branching morphogenesis ceases and the developmental program switches to the phase of alveolarization. During alveolarization, 90 % of the adult gas-exchange area is formed by lifting off new septa from the existing gas-exchange surface Burri et al. [1974]; Kauffman et al. [1974]; Schittny [2017]. Alveolarization itself is divided into two distinct phases, the so-called classical (P4-P21) and continued alveolarization (P14 to approximately P60) Tschanz et al. [2014]; Schittny et al. [2008]. During classical alveolarization, new septa are formed starting from preexisting immature septa which contain a double-layered capillary network. During continued alveolarization, new septa are formed from preexisting mature septa. In general, during alveolarization the structure of the alveolar septa is changed to increase the efficiency of the lung Tschanz et al. [2014]; Schittny et al. [2008].

Here, the timing of lung development is given for rats while the same stages and developmental mechanisms are observed in every placentalian species studied so far, including humans. However, the timing of lung development and especially the time point of birth relative to lung development is adapted in each species Schittny [2017, 2018]; Schittny et al. [2008].

The functional lung units

Pulmonary acini represent the gas-exchanging units of the bronchial tree and are defined as the airways distal of the terminal bronchioles Haefeli-Bleuer and Weibel [1988]. In humans, the acini contain approximately four generations of respiratory bronchioles before reaching the alveolar ducts. The alveolar ducts start at the so-called bronchioalveolar duct junction (BADJ) where the lining of the inner airway surface abruptly changes from the cuboidal epithelium of the bronchioles into type 1 and type 2 alveolar epithelium which covers the alveoli. The small tree of airways distal of the BADJ is called a ventilatory unit Storey and Staub [1962]. Since murine lungs do not possess respiratory bronchioles, the murine acini consist of one single ventilatory unit Storey and Staub [1962]. In *this* study we used rat lungs; hence we exclusively speak of acini. To translate our results to humans, monkeys, or dogs, one would have to compare a rat acinus to a single ventilatory unit in these species.

Developmentally, the BADJ is of particular interest. The junction is formed during the canalicular stage when the epithelia are differentiating. The generation of the airway in which an individual murine acinus starts is defined during the canalicular stage and remains constant once it is formed. As a consequence, in rats the number of formed acini also remains constant during the phase of alveolarization and thereafter Barré et al. [2016, 2014]. The latter was somehow surprising because in rats the total lung volume increases by roughly a factor of 10 during alveolarization Tschanz et al. [2014]. Therefore, the mean volume of the acini has to increase by approximately the same factor.

However, besides the global numbers little is known about the range of the volume of individual acini throughout lung development. The acinar architecture is important for ventilation, airflow Fishler et al. [2015]; Hasler et al. [2019]; Hofemeier and Sznitman [2014] and particle deposition Semmler-Behnke et al. [2012]; Kreyling et al. [2018]; Tschanz et al. [2014]. However, due to the lack of measured data, simulations are based on relatively simple acinar models. The

question of how well these models represent lung physiology remains open until the necessary data and physiologically correct models are available. To contribute to the validation of the computational fluid dynamics simulations, we extracted individual acini and determined their volume and their number of alveoli throughout rat lung development.

Acinus detection and delineation

On single two-dimensional sections—either histological or virtual from tomographic imaging—it is not possible to unambiguously detect which parts of the airways and alveoli are connected three-dimensionally. To detect and delineate individual acini the full three-dimensional information is needed. Historically, single or very few acini were delineated and extracted from manually traced serial sections Mercer and Crapo [1987]; Woodward and Maina [2005] or sectioned silicone casts Rodriguez et al. [1987] with considerable manual work. Non-destructive three-dimensional imaging is best suited to acquire data sets that can be used to easily detect, delineate, and study large amounts of single acini. This has been shown by Vasilescu et al. Vasilescu et al. [2012]. In their study, they reconstructed 22 mouse acini from tomographic data sets of four 12 weeks old mice (at least postnatal day 84, fully developed lungs) and compared those with scanning electron microscope images of acinar silicone rubber casts. Similarly, Kizhakke Puliyakote et al. Puliyakote et al. [2016] analyzed the volume of 32 mouse acini from six animals, as well as the branching pattern of their internal airways. All prior work known to us has analyzed considerably fewer acini than the 268 acini presented here and focuses only on a time point, at which the lungs of the animals are already fully developed while our data spans the postnatal days 4 to 60.

We developed the hereby presented method to analyze large amounts of acini over the course of postnatal lung development in a time-efficient manner since no data on large amounts of acini is available up to now. Our aim was to understand the size distribution of the acini during lung development including the contribution of the alveoli. The observed range *de*creased during postnatal lung development by a factor of 6-7, which represents a completely unexpected result. The obtained data are essential for further investigations, like the influence of the size of the acini on ventilation or gas-washout, respectively. E.g., we are currently using our computational fluid dynamics model to simulate exactly this influence (own unpublished results and Hasler et al. [2019]).

Materials & Methods

Rat lung samples

In the present study, a superset of the animals described by Haberthür et al. Haberthür et al. [2013] were used. Tschanz et al. Tschanz et al. [2014] described a biphasic formation of new alveoli on the same set of animals. The stereological analysis presented here represents a part of the 3R-initiative (replacement, reduction, and refinement) Flecknell [2002] for the ethical use and the reduction of the number of animals sacrificed for science.

Briefly, we extracted the lungs of Sprague-Dawley rats after having deeply

anesthetized them with a mixture of medetomidine, midazolam, and fentanyl Erhardt et al. [2012]. The rats were euthanized by exsanguination during the removal of the lung. The lungs were fixed with 2.5 % glutaraldehyde by intratracheal instillation and kept under constant pressure (20 cm H_2O) during fixation to prevent recoiling of the tissue. The samples were prepared for tomographic imaging by post-fixation with 1 % osmium tetroxide and staining with 4 % uranyl acetate, as well as by dehydration in a graded series of ethanol and embedding in paraffin. In total we assessed 12 animals, on postnatal days P4, P10, P21, and P60 (N=3 per day. From now on, we will omit the postnatal prefix and only mention the day.).

The animals were housed in the central animal facility of the University of Bern. They received food and water ad libitum at a 12/12 hours day/night cycle. The experiments themselves, as well as the handling of the animals before and during the experiments, were approved and supervised by the Federal Food Safety and Veterinary Office of Switzerland and the Department of Agriculture and Nature of the Kanton of Bern in Switzerland.

Tomographic imaging

Tomographic imaging was performed at the TOMCAT beamline (A beamline for TOmographic Microscopy and Coherent rAdiology experimenTs) Marone et al. [2009, 2011] of the Swiss Light Source at the Paul Scherrer Institute in Villigen, Switzerland. The samples were scanned at an X-ray energy of 20.0 keV. After penetrating the sample, the X-rays were converted into visible light by either a 20 μm-thick LuAG:Ce or 18 μm-thick YAG:Ce scintillator screen (both from Crytur, Turnov, Czech Republic), depending on the date of experiments. The resulting visible light was magnified using a 10-times magnifying, diffraction-limited microscope lens and recorded with a 2048×2048 pixel CCD camera (pco.2000, PCO, Kelheim, Germany) with 14 bits dynamic range operated in 2 by 2 binning mode. As a result, in a single field of view, we were able to image a sample volume of a cube of 1.5 mm side length with a voxel side length of 1.48 μm, with the exposure time of the single projections varying between 160 and 200 ms.

Since our samples were substantially larger than the field of view of the detector, we applied the so-called wide-field scanning method Haberthür et al. [2010] to increase the field of view horizontally. For each sub-scan, we recorded 3578 projections and laterally stitched them so that their combined field of view covered the whole sample width. Additionally, two or three such wide-field scans were stacked vertically to match the sample height.

The resulting data sets for each of the lung samples covered a field of view of a cube with approximately 4.5 mm side length which corresponds to an image stack with approximately 3000×3000×3000 pixels at 1.48 μm side length each. The mean size of the tomographic reconstructions, corresponding to the raw data of each of the 42 analyzed tomographic scans is 8 GB, totaling to 336 GB.

Extraction of acini

For the present manuscript, we extracted single acini from three-dimensional data, acquired without destroying the samples. Details of this semi-automatic extraction of the rat acini are described by Haberthür et al. Haberthür et al.

[2013]. Individual acini were extracted from the tomographic data with a custom image processing network established in MeVisLab (version 2.1, 2010-07-26 release, MeVis Medical Solutions and Fraunhofer MEVIS-Institute for Medical Image Computing, Bremen, Germany). To isolate individual acini, we followed the airway tree in the data and manually placed disk-shaped segmentation stoppers at the acinus entrances. Thus, acini were selected independently of their size and only based on their location in the airway tree, are not selected based on their size. The acinus extraction was performed by segmenting the acinus volume with a gray-level threshold-based region growing algorithm with a seed point perpendicular to the manually placed segmentation stopper. Each individual acinus was then exported as a single DICOM file for portability. For each acinus, its exported DICOM file contained the segmented acinus overlaid with the original background for further analysis. Example images from the DICOM files are shown in the bottom row of figure 1.

This semi-automatic acinus segmentation method was applied to lung samples obtained throughout lung development, at postnatal days 4, 10, 21 and 60. In total we extracted 701 acini from 12 animals. Even if our method is efficient, we could not have stereologically assessed all these 701 acini, we thus performed a subsampling and stereologically analyzed 268 acini, as specified in the section below.

Stereological analysis of alveolar characteristics

The stereological estimation of the alveolar number was performed according to the standards for quantitative assessment of lung structure from the American Thoracic Society Hsia et al. [2010a] to guarantee accurate and unbiased results. As a basis for the estimation the volumes of the lung lobes were first determent by water displacement (Scherle 1017). The lobe-volume was estimated a second time by the Cavalieri method. Both volumes were used to calculate the shrinkage, which was used for the correction of the volume data Tschanz et al. [2014].

To perform the stereological analysis, each DICOM file from the isolated acini was sampled into a systematic random set of single images using a MATLAB script dic. The stereological assessment was performed with the STEPanizer TSCHANZ et al. [2011], an easy-to-use software tool for the stereological assessment of digital images. Details of the stereological assessment have previously been described Haberthur et al. [2013]. Briefly, we counted the appearance or disappearance of alveolar wall interruptions. Under the assumption that these only occur in the region of the alveolar mouth opening and correspond to alveolar entrance rings Hyde et al. [2004] we counted said interruptions on paired images spanning the whole acinus volume. By using the disector method Sterio [1984] we were thus able to estimate the number of alveoli in 268 of the 701 exported acini. The acini we analyzed were subsampled from the full data by systematic uniform random sampling Hsia et al. [2010b]. Briefly, a defined fraction of the acini of one animal were analyzed, e.g. one third. During segmentation, the acini were numbered. Based on these numbers every third acinus was selected, randomly starting with the first, second, or third one.

Previously, we have shown that the stereological assessment of the acinar volume (according to the Cavalieri principle Cruz-Orive [1999]) gives comparable results to an automated volume extraction by counting the segmented pixels in

the tomographic data sets Haberthur et al. [2013]. Due to variations in the gray value, the automatic segmentation underestimates the volume of the single acini. The best-suited approach to obtain unbiased results is to assess the volume manually according to the Cavalieri principle, which is what we did for the 268 acini presented in this manuscript. The sets of JPG slices and the raw results from the stereological analysis with the STEPanizer are available on request.

Data analysis and display

All the stereologically assessed data was processed in a Jupyter Kluyver et al. [2016] notebook, producing all the results, including the statistical data and plots shown below. The notebook with all its history is freely available on GitHub Haberthür. The performed calculations are described in detail in section . In the plots, semitransparent circles mark the single observations. The box shows the 25 % to 75 % quartiles range of the data. The whiskers extend to the rest of the distribution. Outliers are determined as a function of the interquartile range and are shown outside the whiskers. Numerical values in the text are given as (averages±standard deviation). P-values in the text and figure legends are given as precise numbers, as suggested by Amrhein et al. Amrhein et al. [2019].

Usually, we performed a Shapiro-Wilk test for normality Shapiro and Wilk [1965] to test whether we can use an U-test for assessing the differences, namely a two-sided Mann-Whitney rank test Mann and Whitney [1947]. This rank test was used to assess the differences between the possible combinations. An additional Kruskal-Wallis H-test for independent samples Kruskal and Wallis [1952] was used to test for sample independence. The statistical analysis was performed in the aforementioned notebook by using the statistical functions of the Scientific Computing Tools for Python Virtanen et al. [2019] or GraphPad Prism 7.01 (GraphPad Software, San Diego, CA, USA).

Results

The number of alveoli per acinus was assessed for 268 individual acini at days 4, 10, 21 and 60. For day 4, we analyzed 125 acini, for day 10 we analyzed 58 acini, for day 21 we analyzed 42 acini, and for day 60 we analyzed 43 acini.

An overview of our process is shown in figure 1. Panels A-D depict a representative light microscopy image for each assessed time point. Panels E–H of figure 1 each correspond to one slice of the tomographic data sets acquired at TOMCAT. Panels I–L of figure 1 show the middle slice of the datasets used for the stereological analysis of the extracted acini. The acinus is shown in light gray as an overlay over the tomographic data.

[Figure 1 about here.]

Alveoli per acinus

The average entrance ring count, which corresponds to the number of alveoli per acinus for the 125 acini at day 4 is (48 ± 41) alveoli. For the 58 acini at day

10 it is (89 ± 84) alveoli, for the 42 acini at day 21 it is (233 ± 164) alveoli. At day 60 we assessed 43 acini in total, on average they have (702 ± 219) alveoli.

The values for the single animal are given in table 1 and plotted in figure 2.

[Table 1 about here.]

[Figure 2 about here.]

The alveolar number shows highly significant differences Šidák-corrected p-value < 0.00167 Šidák [1967]) between all possible combination of days. All possible combinations of entrance ring counts per day are significantly different (all p-values are better than p=1.9×10^{-5}, which is the p-value of the difference between days 4 and 10). The entrance ring counts for all animals are independent (p=1.6×10^{-34}).

Acinus volume

The stereological assessment resulted in a mean acinar volume of $(0.03 \pm 0.04)\,\mathrm{mm}^3$ for the 125 acini at day 4. For the 58 acini at day 10 we get a volume of $(0.04 \pm 0.05)\,\mathrm{mm}^3$. For the 42 acini at day 21 a volume of $(0.10 \pm 0.09)\,\mathrm{mm}^3$ and for the 43 acini at day 60 a volume of $(0.74 \pm 0.37)\,\mathrm{mm}^3$.

The values for the single animals are shown in table 2 and displayed in figure 3.

[Table 2 about here.]

The mean volume of the $20\,\%$ smallest acini to the $20\,\%$ largest acini at day 4 differs by a factor of 27.47 (from $0.0035\,\mathrm{\mu l}$ to $0.095\,\mathrm{\mu l}$, N=25). At day 10, this increase is 15.28 times (from $0.0071\,\mathrm{\mu l}$ to $0.11\,\mathrm{\mu l}$, N=12). At day 21, this increase is 14.42 times (from $0.018\,\mathrm{\mu l}$ to $0.26\,\mathrm{\mu l}$, N=8). At day 60, we saw an increase of 3.94 times (from $0.33\,\mathrm{\mu l}$ to $1.3\,\mathrm{\mu l}$, N=9).

[Figure 3 about here.]

The acinar volumes show highly significant differences between all possible combinations of days except between days 4 and 10 (p=0.08). All other p-values are better than p=4.5×10^{-6}, which is the p-value for the difference between days 10 and 21. The statistical analysis is equal to what is briefly described for the entrance ring count above. The acinar volumes for animals 21E and all animals of day 60 are non-normally distributed, but all acinar volumes per animal are independent (p=5×10^{-29}).

To assess the volume distribution per day, we normalized the volumes per day to the largest volume per day (figure 4). The whiskers in the plots contain all data points 1.5 times the interquartile range past tow low and high quartiles, points outside the whiskers are considered outliers. The non-outlier data points are closest together on day 4 and slightly more spread out at day 10. At days 21 and 60, we see no outliers for the normalized acinar volumes. At days 4, 10 and 21 the distribution of the normalized volumes is skewed towards below the average (median of normalized volumes at day 4: 0.16, at day 10: 0.18 and at day 21: 0.23), we thus see more smaller than larger acini early in the development. At day 60, the distribution is more homogeneous, with the median of the normalized acinus volumes at exactly 0.50.

[Figure 4 about here.]

Alveoli per volume

The mean number of alveoli for the 125 acini at day 4 is (2166 ± 1132) alveoli/mm^3. For the 58 acini at day 10 we counted (2831 ± 1189) alveoli/mm^3, for the 42 acini at day 21 (2723 ± 771) alveoli/mm^3 and for the 43 acini at day 60 (1080 ± 349) alveoli/mm^3. These numbers were found by dividing the counted entrance rings by the acinus volumes.

The numbers for the single animals are shown in table 3 and displayed in figure 5.

[Table 3 about here.]

[Figure 5 about here.]

The counts per volume, which correspond to the number of alveoli per acinus volume are one of several measure for influencing the complexity of the single acini. The number of alveoli per acinus show highly significant differences between all possible combinations of days except between days 10 and 21 (p=0.7). All other days have a p-value better than p=5×10^{-5}, this being the one between days 4 and 21. The statistical analysis is equal to what is briefly described for the entrance ring count above. The counts per volume for all animals except 04A are non-normally distributed, but all are independent (p=2.9×10^{-21}).

Number of acini

The number of acini for day 4 was $(18\,260 \pm 16\,555)$ acini for day 10 $(14\,359 \pm 5611)$ acini, for day 21 $(11\,203 \pm 2681)$ acini and for day 60 (4277 ± 777) acini. These numbers were obtained by dividing the parenchymal volume of the lungs Tschanz et al. [2014] with the Cavalieri-estimated volume of the acini. To calculate these data we had to use data of two different studies.

Total number of alveoli per lung

The total alveolar number for day 4 was found to be (0.64 ± 0.34) million acini, for day 10 (1.23 ± 0.39) million acini, for day 21 (2.65 ± 0.57) million acini, and for day 60 (3.01 ± 0.34) million acini. The total number of acini was estimated by dividing the mean parenchymal volume of the lungs Tschanz et al. [2014] by the stereologically estimated acinar volume. The number of alveoli for each animal is then calculated by multiplying the average entrance ring count per animal with the estimated number of acini per animal.

Volumes of individual alveoli

The average volume of one alveolus including its part of the alveolar duct was estimated by dividing the mean acinar volume by the mean count of alveoli for each day separately, while correcting for the ductal volume. For day 4 we estimate the volume of an individual alveolus at 4.98×10^5 µm^3, for day 10 at 3.34×10^5 µm^3, for day 21 at 3.07×10^5 µm^3 and for day 60 at 8.12×10^5 µm^3. This corresponds to an average diameter per alveolus of 98 µm, 86 µm, 84 µm and 16 µm for days 4, 10, 21 and 60, respectively. The average diameter was estimated by assuming spherical alveoli and solving the volume equation of a sphere to its diameter.

Discussion

To our best knowledge, the present manuscript is the first to quantitatively describe the number of alveoli in individual acini and changes in the volumes of single rat lung acini. Based on stereological analysis of a large number of individual acini we were able to efficiently extract qualitative and quantitative data of large amounts of single acini from non-destructive three-dimensional tomographic data sets in a highly reproducible manner.

We can easily estimate the total number of acini during lung development from our data. The limitations of this study are mainly twofold: a) due to technical reasons (i.e. available beamtime and limitations of the setup we could not image the entire lung and b) lung location (i.e. lower medial tip of the right lobe). However, our sample is more than suitable for the drawn conclusions because it was previously shown that the right lower lobe is a valid sample for the entire lung Barré et al. [2016] and that information drawn from one lobe is representative of the whole lung Zeltner et al. [1990]. Furthermore, based on stereological data, Zeltner et al. Zeltner et al. [1990] were able to show that the lung parenchyma is quite homogeneous. Looking at individual acini, Kizhakke Puliyakote et al. Puliyakote et al. [2016] observed that the peripheral, pleura facing acini possess an acinar volume which is two thirds larger than the volume of the central ones. However, they did not report additional regional differences, e.g. between acini located cranially and caudally. Own preliminary data point to the same result. We imaged an entire rat lung *in situ* immediately post mortem using high-resolution synchrotron radiation based X-ray tomographic microscopy Borisova et al. [2020]. A first analysis of regional difference shows similar results as mentioned above.

Acinus volumes

The mean acinar volume significantly increases during lung development from postnatal days 10 to 60 with all p-values better than 4.5×10^{-6}. No significant difference was observed between postnatal days 4 and 10 (1.2×increase, p=0.08). A large part of the acinus volume increase happens after day 21 (compare figure 3), which is consistent with previous literature Tschanz et al. [2014]; Burri et al. [1974].

As mentioned in the introduction, Barré et al. Barré et al. [2016] have shown that the number of acini remains constant during lung development, albeit in a different strain of animals (Wistar Bern compared to Sprague-Dawley in this study). The total volume of the alveolar space increases by roughly a factor of 11 during alveolarization Tschanz et al. [2014]. Therefore, we expect the mean volume of the acini to increase by at least the same factor.

We observed a mean increase of the acinus volumes of 22.66 times between days 4 and 60 (p=1.8×10^{-22}). The other combinations of increases were: day 4 to day 21: 3.18×(p=1.3×10^{-9}), day 10 to day 21: 2.60×(p=4.5×10^{-6}), day 10 to day 60: 18.46×(p=1.4×10^{-17}), day 21 to day 60: 7.11×(p=7.1×10^{-15}). The increase from day 4 to day 10 was not significant (p=0.08) at 1.23×).

During the same time, the coefficients of variation of the acinus volumes dropped from 1.26 to 0.49. At day 4, the mean volumes of the 20 % largest acini are 27.47 times larger than the volumes of the 20 % smallest acini (N=25). At day 60, the 20 % largest acini are 3.94 times larger than the 20 % smallest

acini (N=9). The size distribution of the largest to smallest acini thus decreases nearly 7 times over the course of the lung development. This indicates that the dispersion of the value is based on an entirely different pattern of distribution of acinar sizes and that the difference of the mean acinus volumes from the smallest to the largest fraction per day becomes smaller and smaller during postnatal lung development.

When comparing the acinus volumes normalized to the largest volume per day, we found that the volumes are closer together on day 4 and are spread out more on the other days (figure 4). The normalized volumes (without outliers) are most compact on day 4 and slightly more spread out at day 10. At days 21 and 60, we see no outliers for the normalized acinar volumes, which thus range from zero to one. The median of the normalized volumes is smaller than the average at days 4, 10 and 21 while it is exactly at 0.5 for day 60. This means that we have more smaller than larger acini early in the development and a very homogeneous distribution late in lung development. This result suggests that acinar growth is not constant during lung development. It seems that some acini increase their volume faster than others. As a result, we speculate that an increased percentage of acini is somehow dormant at day 4–21, while at day 60 most of the acini reached their final volume.

By dividing the parenchymal volumes of each lung estimated by Tschanz et al. (Table 1, Tschanz et al. [2014]) by the mean number of acini estimated by Barré et al. Barré et al. [2014] we can estimate the mean acinar volume for each day. The acinus volume we estimated by point counting (Cavalieri estimation) are on average 2.07 times lower but show a similar increase over the studied period (see figure. 6). It appears that the comparison between data obtained by stereology based on classical paraffin sections or on any kind of 3D-imaging has its limitation if it comes to the absolute numbers. First, due to the *coast of Britain*-effect the exact value of every surface measurement is depending on the resolution of the imaging method Mandelbrot [1967]. Second, the thickness of the alveolar septa in 3D-imaging data is heavily depending on the segmentation between airspace and tissue—regardless if it is done manually or by a computed algorithm. We observed that the airspace volumes obtained by 3D-imaging has a tendency to be smaller than the ones determined using classical paraffin sections. However, the relative numbers obtained in one study match the ones of other studies quite well.

[Figure 6 about here.]

Number of alveoli per acinus volume

The volume and shape of the alveoli, alveolar ducts, and the acini are critical parameters for ventilation and particle deposition Sznitman et al. [2010]; Henry et al. [2012]; Tsuda et al. [2008]. Particle deposition, for example, is relatively low during the first two weeks of a rats' life, high at postnatal day P21 and medium at days P36–P90 Kreyling et al. [2018]; Semmler-Behnke et al. [2012]. This correlates well with our results, where we found a significantly larger number of alveoli per acinus volume at days 10 and 21 compared to days 4 and 60 (p-values all better than 5.4×10^{-5}, which is the one for day 4 vs. day 21). Kreyling et al. Kreyling et al. [2018] report maximal gold nanoparticle retention at this time point (see their figure 1).

Tschanz et al. Tschanz et al. [2014] stereologically estimated the mean total acinar volume as the sum of ductal and alveolar air spaces. The mean volume of the individual alveoli was found to be the smallest around postnatal day 21 (see their figure 4). If we take the number of alveoli as one possible measure of acinar complexity, we postulate a high acinar complexity at days 10 and 21 while early and late in the lung development, the acinar complexity is relatively low. This also correlates with structural changes. In rats, the alveolar air spaces are large during the first two weeks of life, small at day P21 and medium afterward Tschanz et al. [2014].

Number of acini

Previously, it was shown that the number of acini (e.g. counted acinar entraces) remains constant during postnatal lung development from day 4 to 60 Barré et al. [2016], with a mean of (5840 ± 547) acini per lung (in another strain of animals, full lung data available for days 4 and 60). Even though we did not aim to assess the number of acini in this manuscript, calculating this number helps to validate our assessment and helps us to collate the data that can be derived from our stereological analysis to prior results.

Based on the acinar and parenchymal volume we calculated the number of acini and were able to verify the previously reported results to lie within our margin of error at days 4 and 60 (Barré et al: (6236 ± 497) acini and (5612 ± 547) acini, respectively). As shown in figure 7, our values $\pm$ their standard deviation are overlapping the previously found values, even though we generally see a trend that the number of acini seems to decrease during lung development. This might be due to fusing of several acini to one or simply because we missed some acini in the analysis of the large datasets.

[Figure 7 about here.]

Total alveolar number

By multiplication of the number of acini by the number of alveoli we estimate a total amount of alveoli per lung of 0.64 ± 0.34, 1.23 ± 0.39, 2.65 ± 0.57 and 3.01 ± 0.58 million alveoli for days 4, 10, 21 and 60, respectively. Tschanz et al. estimated these total numbers of alveoli per lung as 0.823 ± 0.010, 3.538 ± 1.260, 14.303 ± 3.080 and 19.297 ± 3.060 million alveoli for days 4, 10, 21 and 60, respectively Tschanz et al. [2014], Table 1). On average, our results are five times smaller but show the same trend when linearly scaled with a factor of 5.04. The values from Tschanz et al. overlap well with our linearly scaled values, as shown in figure 8.

As shown in a study by Osmanagic et al. Osmanagic et al. [2010] the total alveolar number shows a large variability between different labs. E.g. the reported total number of alveoli in adult C57BL6 mice differs by a factor of approximately 8. All the studies mentioned my Osmanagic et al. were performed in labs well-known for their quality of stereology-based investigations of lung morphology. Therefore, any simple bias or methodical error may not account for the reported differences. It seems to be that the total number of alveoli is dependent on unknown small methodical differences and minute differences in how the operator actually does count.

However, relative numbers as reported in this study are more reliable. This is especially the case when the comparison is done in groups of animals processed in parallel and counted by the same operator. We assessed the same lungs as Tschanz et al. Tschanz et al. [2014], but based on a very different imaging method and with two different persons performing the stereological assessment. This may explain the relative factor of 5.04 that we found as a difference.

[Figure 8 about here.]

Volumes of individual alveoli

By dividing the alveolar air space by the number of alveoli (both from Table 1 of Tschanz et al. [2014]) we can estimate a mean alveolar volume of $5.91 \times 10^5 \, \mu m^3$ for day 4, of $2.49 \times 10^5 \, \mu m^3$ for day 10, of $1.34 \times 10^5 \, \mu m^3$ for day 21, and $2.93 \times 10^5 \, \mu m^3$ for day 60. On average, our values (calculated by dividing the mean acinar volume by the mean count of alveoli per acinus) are 1.81 times larger than these estimates of individual alveolar volume (our values day 4: $4.98 \times 10^5 \, \mu m^3$, day 10: $3.34 \times 10^5 \, \mu m^3$, day 21 $3.07 \times 10^5 \, \mu m^3$ and day 60 $8.12 \times 10^5 \, \mu m^3$), but scale equally during postnatal lung development. Tschanz et al. estimated the total alveolar volume without the alveolar ducts. Since we directly estimated the volume fraction of the alveoli per acinus our number is actually expected to be larger.

Additional characteristics of the acinus

We are well aware that there are many more parameters of the acinar characteristics can be extracted from the acinar datasets we segmented for this study. Especially of interest are the microarchitecture in the acinus like the branching pattern of the intra-acinar airways or surface area characteristics of the acinar and alveolar walls. For this study we focused on the easily stereologically available parameters. Ongoing work in our group is aiming to describe additional parameters of the acinus microarchitecture (e.g. fractal dimension, surface curvature, etc.) but the present study did not aim at characterizing these parameters.

Physiological relevance of the acinar size

It has been predicted by computational fluid dynamics simulations and by to-scale experiments that the amount and location of massless particle deposition, as well as ventilation, are depending on the size of the acini Fishler et al. [2015]; Hofemeier and Sznitman [2014]; Sznitman [2009]. The predictions could be confirmed at least in one case, where mice received fluorescent particles during mechanical ventilation. In 3D-visualizations the particles were predominately detected in the proximal half of the acini Yang et al. [2019].

Based on the above-mentioned knowledge and the data of this study, we propose the following. In very small acini as the smallest ones, we observed at postnatal day 4 acinar flows are mainly characterized by radial streamlines Sznitman et al. [2009, 2010]. Massless particle deposition is governed by diffusion Sznitman et al. [2006]. As the acini increase in size, convective transport becomes more and more dominant in the proximal regions of the acini. In larger acini,

the proximal region shows much higher particle retention than the distal one, because in this region the larger air-flow induces a circular current, which facilitates alveolar particle deposition. The proximal regions may even be viewed analogously to a filter capturing the particles whereas the distal regions do not receive significant amounts Fishler et al. [2015]; Hofemeier and Sznitman [2014]; Sznitman [2009]; Tsuda et al. [2008]; Henry et al. [2012]; Semmler-Behnke et al. [2012].

Multi-breath gas-washout is also dependent on the size of the acini. Based on computational simulations of nitrogen multi-breath gas-washout it has been shown that small acini washout faster at the beginning and slower at the end of the washout as compared to larger acini Hasler et al. [2019]. Not surprisingly, the distribution of the acinar sizes in terms of their range has also an influence on the gas-washout. Furthermore, a clustering versus an even distribution of the locations of acinar similar sizes effects gas-washout. Therefore, the question where an acinus of which size is located in the lungs has a significant influence on pulmonary ventilation and particle deposition.

The larger distribution of the acinar sizes at day 4 as compared to day 60 let us speculate what this may mean biologically. Physiologically, a larger distribution implies less homogeneous ventilation and an increased gas-washout time. Therefore, the decrease of the inhomogeneity of the acinar sizes may be understood as optimization of the gas-exchange region in the lungs of adult animals.

The consideration of the location is not only a theoretical one, because it has been shown in adult mice that the acini close to the pleura are larger than the central ones Puliyakote et al. [2016]. Therefore, acini of different size are not evenly distributed in the lungs.

Conclusion

Newly formed acini show a large size distribution which is reduced by a factor of 7 until adulthood. Most likely the latter leads to more homogeneous ventilation of the lung and an increase of gas-washout which is equal to a decreased in the number of breaths needed for gas-washout.

To our best knowledge, no in-depth characterization of the acinar properties as presented here—i.e. spanning the lung development period—was available up to now. The data presented in this study can contribute to improve computational simulations of pulmonary ventilation and particle deposition.

Acknowledgments

We thank Mohammed Ouanella and Bettina de Breuyn for expert technical assistance. Federica Marone and Bernd Pinzer were of paramount importance with their support at the TOMCAT beamline. Fluri Wieland was of great help with the statistical analysis. We are thankful for the support by the Swiss National Science Foundation (grants 310030_153468 and 310030_175953).

Contributions

JS conceived the project. DH, SB and JS performed the tomographic data acquisition. DH segmented the acini. EY and DH stereologically analyzed the acini. DH, ST, TC and JS analyzed the data. DH wrote the manuscript and generated all the figures. ST and JS contributed to writing, all authors proofread the manuscript.

Disclosures

No conflicts of interest, financial or otherwise, are declared by the authors.

References

DicomReaderAndSaverForSTEPanizer.m. https://git.io/JTg4F. URL `https://git.io/JTg4F`. Accessed on Fri, January 05, 2018.

Valentin Amrhein, Sander Greenland, and Blake McShane. Scientists rise up against statistical significance. *Nature*, 567(7748):305–307, mar 2019. doi: 10.1038/d41586-019-00857-9. URL `https://doi.org/10.1038%2Fd41586-019-00857-9`.

Sébastien F. Barré, David Haberthür, Marco Stampanoni, and Johannes C. Schittny. Efficient estimation of the total number of acini in adult rat lung. *Physiological Reports*, 2(7):e12063, jul 2014. doi: 10.14814/phy2.12063. URL `https://doi.org/10.14814%2Fphy2.12063`.

Sébastien F. Barré, David Haberthür, Tiziana P. Cremona, Marco Stampanoni, and Johannes C. Schittny. The total number of acini remains constant throughout postnatal rat lung development. *American Journal of Physiology-Lung Cellular and Molecular Physiology*, 311(6):L1082–L1089, dec 2016. doi: 10.1152/ajplung.00325.2016. URL `https://doi.org/10.1152%2Fajplung.00325.2016`.

E Borisova, G Lovric, A Miettinen, L Fardin, S Bayat, A Larsson, M Stampanoni, JC Schittny, and CM Schlepütz. Micrometer-resolution X-ray tomographic full-volume reconstruction of an intact post-mortem juvenile rat lung. *Histochem Cell Biol*, Mar 2020.

Peter H. Burri, Jaroslav Dbaly, and Ewald R. Weibel. The postnatal growth of the rat lung. I. Morphometry. *The Anatomical Record*, 178(4):711–730, apr 1974. doi: 10.1002/ar.1091780405. URL `https://doi.org/10.1002%2Far.1091780405`.

L. M. Cruz-Orive. Precision of Cavalieri sections and slices with local errors. *Journal of Microscopy*, 193(3):182–198, mar 1999. ISSN 0022-2720. doi: 10.1046/j.1365-2818.1999.00460.x. URL `http://dx.doi.org/10.1046/j.1365-2818.1999.00460.x`.

Wolf Erhardt, Petra Kölle, Julia Henke, Jörg Haberstroh, Christine Baumgartner, and Sabine Tacke. *Anästhesie und Analgesie beim Klein-und Heimtier:*

mit Exoten, Labortieren, Vögeln, Reptilien, Amphibien und Fischen. Schattauer Verlag, 2012.

Rami Fishler, Philipp Hofemeier, Yael Etzion, Yael Dubowski, and Josué Sznitman. Particle dynamics and deposition in true-scale pulmonary acinar models. *Scientific Reports*, 5(1), sep 2015. doi: 10.1038/srep14071. URL https://doi.org/10.1038%2Fsrep14071.

P Flecknell. Replacement, reduction and refinement. *ALTEX*, 19:73–8, 2002.

D. Haberthur, S. F. Barre, S. A. Tschanz, E. Yao, M. Stampanoni, and J. C. Schittny. Visualization and stereological characterization of individual rat lung acini by high-resolution X-ray tomographic microscopy. *Journal of Applied Physiology*, 115(9):1379–1387, aug 2013. doi: 10.1152/japplphysiol. 00642.2013. URL https://doi.org/10.1152%2Fjapplphysiol.00642. 2013.

David Haberthür, Christoph Hintermüller, Federica Marone, Johannes C. Schittny, and Marco Stampanoni. Radiation dose optimized lateral expansion of the field of view in synchrotron radiation X-ray tomographic microscopy. *Journal of Synchrotron Radiation*, 17(5):590–599, sep 2010. ISSN 09090495. doi: 10.1107/S0909049510019618. URL http://doi.org/10. 1107/S0909049510019618.

David Haberthür. habi/acinar-analysis: Zenodo.

David Haberthür, Sébastien F. Barré, Stefan A. Tschanz, Eveline Yao, Marco Stampanoni, and Johannes C. Schittny. Visualization and stereological characterization of individual rat lung acini by high-resolution X-ray tomographic microscopy. *Journal of Applied Physiology*, 115(9):1379–1387, nov 2013. doi: 10.1152/japplphysiol.00642.2013. URL https://doi.org/10. 1152%2Fjapplphysiol.00642.2013.

Beatrice Haefeli-Bleuer and Ewald R. Weibel. Morphometry of the human pulmonary acinus. *The Anatomical Record*, 220(4):401–414, apr 1988. doi: 10. 1002/ar.1092200410. URL https://doi.org/10.1002%2Far.1092200410.

David Hasler, Pinelopi Anagnostopoulou, Sylvia Nyilas, Philipp Latzin, Johannes Schittny, and Dominik Obrist. A multi-scale model of gas transport in the lung to study heterogeneous lung ventilation during the multiple-breath washout test. *PLOS Computational Biology*, 15(6):e1007079, jun 2019. doi: 10.1371/journal.pcbi.1007079. URL https://doi.org/10.1371%2Fjournal. pcbi.1007079.

F. S. Henry, S. Haber, D. Haberthür, N. Filipovic, D. Milasinovic, J. C. Schittny, and A. Tsuda. The Simultaneous Role of an Alveolus as Flow Mixer and Flow Feeder for the Deposition of Inhaled Submicron Particles. *Journal of Biomechanical Engineering*, 134(12), nov 2012. doi: 10.1115/1.4007949. URL https://doi.org/10.1115%2F1.4007949.

Philipp Hofemeier and Josué Sznitman. Role of Alveolar Topology on Acinar Flows and Convective Mixing. *Journal of Biomechanical Engineering*, 136(6), may 2014. doi: 10.1115/1.4027328. URL https://doi.org/10.1115%2F1. 4027328.

CC Hsia, DM Hyde, M Ochs, and ER Weibel. An official research policy statement of the American Thoracic Society/European Respiratory Society: standards for quantitative assessment of lung structure. *Am J Respir Crit Care Med*, 181:394–418, Feb 2010a.

Connie C. W. Hsia, Dallas M. Hyde, Matthias Ochs, and Ewald R. Weibel. An Official Research Policy Statement of the American Thoracic Society/European Respiratory Society: Standards for Quantitative Assessment of Lung Structure. *American Journal of Respiratory and Critical Care Medicine*, 181(4):394–418, feb 2010b. doi: 10.1164/rccm.200809-1522st. URL `https://doi.org/10.1164%2Frccm.200809-1522st`.

DM Hyde, NK Tyler, LF Putney, P Singh, and HJ Gundersen. Total number and mean size of alveoli in mammalian lung estimated using fractionator sampling and unbiased estimates of the Euler characteristic of alveolar openings. *Anat Rec A Discov Mol Cell Evol Biol*, 277:216–26, Mar 2004.

Shirley L. Kauffman, Peter H. Burri, and Ewald R. Weibel. The postnatal growth of the rat lung II. Autoradiography. *The Anatomical Record*, 180(1): 63–76, sep 1974. doi: 10.1002/ar.1091800108. URL `https://doi.org/10.1002%2Far.1091800108`.

Thomas Kluyver, Benjamin Ragan-Kelley, Fernando Pérez, Brian Granger, Matthias Bussonnier, Jonathan Frederic, Kyle Kelley, Jessica Hamrick, Jason Grout, Sylvain Corlay, Paul Ivanov, Damián Avila, Safia Abdalla, and Carol Willing. Jupyter Notebooks − a publishing format for reproducible computational workflows. In F. Loizides and B. Schmidt, editors, *Positioning and Power in Academic Publishing: Players, Agents and Agendas*, pages 87 − 90. IOS Press, 2016.

Wolfgang G. Kreyling, Winfried Möller, Uwe Holzwarth, Stephanie Hirn, Alexander Wenk, Carsten Schleh, Martin Schäffler, Nadine Haberl, Neil Gibson, and Johannes C. Schittny. Age-Dependent Rat Lung Deposition Patterns of Inhaled 20 Nanometer Gold Nanoparticles and their Quantitative Biokinetics in Adult Rats. *ACS Nano*, 12(8):7771–7790, aug 2018. doi: 10.1021/acsnano.8b01826. URL `https://doi.org/10.1021%2Facsnano.8b01826`.

William H. Kruskal and W. Allen Wallis. Use of Ranks in One-Criterion Variance Analysis. *Journal of the American Statistical Association*, 47(260): 583–621, dec 1952. doi: 10.1080/01621459.1952.10483441. URL `https://doi.org/10.1080%2F01621459.1952.10483441`.

B. Mandelbrot. How Long Is the Coast of Britain? Statistical Self-Similarity and Fractional Dimension. *Science*, 156(3775):636–638, may 1967. doi: 10.1126/science.156.3775.636. URL `https://doi.org/10.1126%2Fscience.156.3775.636`.

H. B. Mann and D. R. Whitney. On a Test of Whether one of Two Random Variables is Stochastically Larger than the Other. *The Annals of Mathematical Statistics*, 18(1):50–60, mar 1947. doi: 10.1214/aoms/1177730491. URL `https://doi.org/10.1214%2Faoms%2F1177730491`.

F. Marone, C Hintermüller, S. McDonald, R. Abela, G. Mikuljan, A. Isenegger, and M. Stampanoni. X-ray Tomographic Microscopy at TOMCAT. *Journal of Physics: Conference Series*, 186:012042, sep 2009. ISSN 1742-6596. doi: 10.1088/1742-6596/186/1/012042. URL http://stacks.iop.org/1742-6596/186/i=1/a=012042?key=crossref.7cf7265980f56baa4c39480baaf31367.

F. Marone, R. Mokso, J.L. Fife, S. Irvine, P. Modregger, B.R. Pinzer, K. Mader, A. Isenegger, G. Mikuljan, and M. Stampanoni. Synchrotron-based X-ray Tomographic Microscopy at the Swiss Light Source for Industrial Applications. *Synchrotron Radiation News*, 24(6):24–29, nov 2011. doi: 10.1080/08940886.2011.634315. URL https://doi.org/10.1080%2F08940886.2011.634315.

RR Mercer and JD Crapo. Three-dimensional reconstruction of the rat acinus. *J Appl Physiol (1985)*, 63:785–94, Aug 1987.

E. Osmanagic, A. L. Sukstanskii, J. D. Quirk, J. C. Woods, R. A. Pierce, M. S. Conradi, E. R. Weibel, and D. A. Yablonskiy. Quantitative assessment of lung microstructure in healthy mice using an MR-based 3He lung morphometry technique. *Journal of Applied Physiology*, 109(6):1592–1599, dec 2010. doi: 10.1152/japplphysiol.00736.2010. URL https://doi.org/10.1152%2Fjapplphysiol.00736.2010.

Abhilash S. Kizhakke Puliyakote, Dragoş M. Vasilescu, John D. Newell, Ge Wang, Ewald R. Weibel, and Eric A. Hoffman. Morphometric differences between central vs. surface acini in A/J mice using high-resolution micro-computed tomography. *Journal of Applied Physiology*, 121(1):115–122, may 2016. doi: 10.1152/japplphysiol.00317.2016. URL https://doi.org/10.1152%2Fjapplphysiol.00317.2016.

M. Rodriguez, S. Bur, A. Favre, and E. R. Weibel. Pulmonary acinus: Geometry and morphometry of the peripheral airway system in rat and rabbit. *American Journal of Anatomy*, 180(2):143–155, oct 1987. doi: 10.1002/aja.1001800204. URL https://doi.org/10.1002%2Faja.1001800204.

Johannes Schittny and Peter Burri. Development and growth of the lung. In *Fishman's Pulmonary Diseases and Disorders, 4th Edition*. McGraw-Hill, 2008. ISBN 978-0-07-145739-2.

Johannes C. Schittny. Development of the lung. *Cell and Tissue Research*, 367(3):427–444, jan 2017. doi: 10.1007/s00441-016-2545-0. URL https://doi.org/10.1007%2Fs00441-016-2545-0.

Johannes C. Schittny. How high resolution 3-dimensional imaging changes our understanding of postnatal lung development. *Histochemistry and Cell Biology*, 150(6):677–691, nov 2018. doi: 10.1007/s00418-018-1749-7. URL https://doi.org/10.1007%2Fs00418-018-1749-7.

Johannes C. Schittny, Sonja I. Mund, and Marco Stampanoni. Evidence and structural mechanism for late lung alveolarization. *American Journal of Physiology-Lung Cellular and Molecular Physiology*, 294(2):L246–L254, feb 2008. doi: 10.1152/ajplung.00296.2007. URL https://doi.org/10.1152%2Fajplung.00296.2007.

M. Semmler-Behnke, W. G. Kreyling, H. Schulz, S. Takenaka, J. P. Butler, F. S. Henry, and A. Tsuda. Nanoparticle delivery in infant lungs. *Proceedings of the National Academy of Sciences*, 109(13):5092–5097, mar 2012. doi: 10.1073/pnas.1119339109. URL `https://doi.org/10.1073%2Fpnas.1119339109`.

S. S. Shapiro and M. B. Wilk. An analysis of variance test for normality (complete samples). *Biometrika*, 52(3-4):591–611, dec 1965. doi: 10.1093/biomet/52.3-4.591. URL `https://doi.org/10.1093%2Fbiomet%2F52.3-4.591`.

Zbyněk Šidák. Rectangular Confidence Regions for the Means of Multivariate Normal Distributions. *Journal of the American Statistical Association*, 62 (318):626–633, jun 1967. doi: 10.1080/01621459.1967.10482935. URL `https://doi.org/10.1080%2F01621459.1967.10482935`.

D. C. Sterio. The unbiased estimation of number and sizes of arbitrary particles using the disector. *Journal of microscopy*, 134(Pt 2):127–36, may 1984. ISSN 0022-2720. URL `http://dx.doi.org/10.1111/j.1365-2818.1984.tb02501.x`.

Winnifred F. Storey and Norman C. Staub. Ventilation of terminal air units. *Journal of Applied Physiology*, 17(3):391–397, may 1962. doi: 10.1152/jappl.1962.17.3.391. URL `https://doi.org/10.1152%2Fjappl.1962.17.3.391`.

J. Sznitman, F. Heimsch, D. Altorfer, J.C. Schittny, and T. Rösgen. Alveolar flow simulations during rhythmical breathing motion in reconstructed XTM acinar airspaces. *Journal of Biomechanics*, 39:S441, jan 2006. doi: 10.1016/s0021-9290(06)84801-3. URL `https://doi.org/10.1016%2Fs0021-9290%2806%2984801-3`.

J. Sznitman, R. Sutter, D. Altorfer, M. Stampanoni, T. Rösgen, and J. C. Schittny. Visualization of respiratory flows from 3D reconstructed alveolar airspaces using X-ray tomographic microscopy. *Journal of Visualization*, 13 (4):337–345, jun 2010. doi: 10.1007/s12650-010-0043-0. URL `https://doi.org/10.1007%2Fs12650-010-0043-0`.

Josué Sznitman. Convective gas transport in the pulmonary acinus: Comparing roles of convective and diffusive lengths. *Journal of Biomechanics*, 42(6):789–792, apr 2009. doi: 10.1016/j.jbiomech.2008.12.022. URL `https://doi.org/10.1016%2Fj.jbiomech.2008.12.022`.

Josué Sznitman, Thomas Heimsch, Johannes H. Wildhaber, Akira Tsuda, and Thomas Rösgen. Respiratory Flow Phenomena and Gravitational Deposition in a Three-Dimensional Space-Filling Model of the Pulmonary Acinar Tree. *Journal of Biomechanical Engineering*, 131(3), jan 2009. doi: 10.1115/1.3049481. URL `https://doi.org/10.1115%2F1.3049481`.

S.A. TSCHANZ, P.H. BURRI, and E.R. WEIBEL. A simple tool for stereological assessment of digital images: the STEPanizer. *Journal of Microscopy*, 243(1):47–59, mar 2011. doi: 10.1111/j.1365-2818.2010.03481.x. URL `https://doi.org/10.1111%2Fj.1365-2818.2010.03481.x`.

Stefan A. Tschanz, Lilian A. Salm, Matthias Roth-Kleiner, Sebastien F. Barré, Peter H. Burri, and Johannes C. Schittny. Rat lungs show a biphasic formation of new alveoli during postnatal development. *Journal of Applied Physiology*, 117(1):89–95, jul 2014. doi: 10.1152/japplphysiol.01355.2013. URL `https://doi.org/10.1152%2Fjapplphysiol.01355.2013`.

A. Tsuda, N. Filipovic, D. Haberthür, R. Dickie, Y. Matsui, M. Stampanoni, and J. C. Schittny. Finite element 3D reconstruction of the pulmonary acinus imaged by synchrotron X-ray tomography. *Journal of Applied Physiology*, 105 (3):964–976, sep 2008. doi: 10.1152/japplphysiol.90546.2008. URL `https://doi.org/10.1152%2Fjapplphysiol.90546.2008`.

D. M. Vasilescu, Z. Gao, P. K. Saha, L. Yin, G. Wang, B. Haefeli-Bleuer, M. Ochs, E. R. Weibel, and E. A. Hoffman. Assessment of morphometry of pulmonary acini in mouse lungs by nondestructive imaging using multiscale microcomputed tomography. *Proceedings of the National Academy of Sciences*, 109(42):17105–17110, oct 2012. doi: 10.1073/pnas.1215112109. URL `https://doi.org/10.1073%2Fpnas.1215112109`.

Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, CJ Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and SciPy 1. 0 Contributors. SciPy 1.0–Fundamental Algorithms for Scientific Computing in Python, 2019.

Jeremy D. Woodward and John N. Maina. A 3D digital reconstruction of the components of the gas exchange tissue of the lung of the muscovy duck Cairina moschata. *Journal of Anatomy*, 206(5):477–492, may 2005. doi: 10.1111/j. 1469-7580.2005.00413.x. URL `https://doi.org/10.1111%2Fj.1469-7580.2005.00413.x`.

Lin Yang, Annette Feuchtinger, Winfried Möller, Yaobo Ding, David Kutschke, Gabriele Möller, Johannes C. Schittny, Gerald Burgstaller, Werner Hofmann, Tobias Stoeger, Daniel Razansky, Alex Walch, and Otmar Schmid. Three-Dimensional Quantitative Co-Mapping of Pulmonary Morphology and Nanoparticle Distribution with Cellular Resolution in Nondissected Murine Lungs. *ACS Nano*, jan 2019. doi: 10.1021/acsnano.8b07524. URL `https://doi.org/10.1021%2Facsnano.8b07524`.

TB Zeltner, M Bertacchini, A Messerli, and PH Burri. Morphometric estimation of regional differences in the rat lung. *Exp Lung Res*, 16:145–58, Mar 1990.

List of Figures

Figure 1: Example images for each of the assessed time points. A-D: Representative light microscopy images. E–H: One slice of the tomographic data sets. I-L: Middle slice of one of the extracted acini. The extracted acinus is overlaid on the tomographic dataset. Panels J and K also show the segmentation stopper at the left border of the image. The shown images correspond to the middle slice of the full data sets used for the stereological analysis. The scale bar was burnt in during preparation of the image stacks and is 100 µm long. Panels A, E & I: Day 4. Panels B, F & J: Day 10. Panels C, G & K: Day 21. Panels D, H & L: Day 60.

Figure 2: Distribution of the number of alveoli per acinus (entrance ring count per acinus) for each animal. Left: linear scale, right: logarithmic scale. The entrance ring counts per day are all significantly different (all p-values better than 1.9×10^{-5}, which is the one between days 4 and 10).

Figure 3: Distribution of the acinar volumes for each animal. Left: linear scale, right: logarithmic scale. The acinar volumes are all significantly different (all p-values better than 4.5×10^{-6}, which is the one between days 10 and 21) for each combination of days except between days 4 and 10 (p=0.08).

Figure 4: Normalized acinar volumes. The spread of the volumes is small early in the development and skewed towards the lower end of the volumes. At day 60, the median is exactly in the middle of the normalized volumes.

Figure 5: Number of alveoli per acinus volume. Left: linear scale, right: logarithmic scale. The number of alveoli per acinus volume are all significantly different (all p-values are better than 5×10^{-5}, which is the one between days 4 and 21) for each combination of days except between days 10 and 21 (p=0.7).

Figure 6: Plot of the mean acinar volumes. By dividing the parenchymal volumes of each lung estimated by Tschanz et al. (Table 1, Tschanz et al. [2014]) by the mean number of acini estimated by Barre et al. Barré et al. [2014] we can estimate the mean acinar volume for each measured time point (blue plot). We assessed the volumes of the acini by point counting (Cavalieri estimation, orange plot). Left: original data. Right: Our data, scaled by a linear correction factor. Our volumes of the acini are on average 2.07 times lower but show a similar increase over the studied period. While the estimation based on Tschanz et al. is from the entire right middle lobe and on Barre et al. globally for the whole organ, our estimation is solely based on the stereological analysis of the acini in the lower medial tip of the right lower lung lobe.

Figure 7: Plot of the estimated number of acini. The blue plot shows our estimation, based on the division of the parenchymal volume by the average volume of the acini. The orange plot is based on the data shown by Barre et al. Barré et al. [2014, 2016].

Figure 8: Number of alveoli based on data from Tschanz et al. Tschanz et al. [2014] together with our linearly scaled data (our data multiplied by 5.04). Values are plotted with a band of the width of their standard deviation.

List of Tables

Animal	Assessed acini	Average counts	STD	Minimum	Maximum
04A	51	26.73	23.56	4	111
04B	23	65.04	43.85	15	171
04C	51	60.51	45.42	10	249
10A	27	77.85	56.31	18	245
10B	14	84.57	65.75	23	199
10C	17	108.76	125.45	15	505
21B	14	208.14	197.71	35	781
21D	17	196.0	128.56	50	493
21E	11	323.09	145.94	108	572
60B	24	701.75	230.95	322	1296
60D	10	668.3	121.67	444	810
60E	9	739.11	281.22	272	1204

Table 1: Detailed alveolar numbers for each animal.

Animal	Assessed acini	Average [mm^3]	STD [mm^3]	Minimum [mm^3]	Maximum [mm^3]
04A	51	0.01	0.01	0.0	0.04
04B	23	0.06	0.05	0.01	0.19
04C	51	0.04	0.04	0.01	0.28
10A	27	0.03	0.03	0.0	0.11
10B	14	0.03	0.03	0.0	0.1
10C	17	0.06	0.08	0.0	0.33
21B	14	0.09	0.09	0.01	0.32
21D	17	0.11	0.11	0.01	0.38
21E	11	0.12	0.07	0.03	0.24
60B	24	0.59	0.22	0.23	1.1
60D	10	0.89	0.35	0.36	1.58
60E	9	0.98	0.53	0.19	1.88

Table 2: Detailed volume data for each animal.

Animal	Assessed acini	Average [mm^{-3}]	STD [mm^{-3}]	Minimum [mm^{-3}]	Maximum [mm^{-3}]
04A	51	3124	1152	1618	8208
04B	23	1363	499	679	2704
04C	51	1569	380	799	2775
10A	27	2685	1022	1246	5582
10B	14	3091	1166	1890	5422
10C	17	2848	1462	1182	7009
21B	14	2642	493	1625	3344
21D	17	2534	895	1111	4032
21E	11	3119	775	1604	4379
60B	24	1249	297	530	1912
60D	10	838	271	476	1221
60E	9	898	323	411	1463

Table 3: Details of alveolar density (number per mm^3) per acinus volume for each animal.