Convergent evolution of sex-specific leg ornaments in Drosophilidae – from cells to structures.

K. Tanaka [1*], O. Barmina [1], J. H. Massey [2,3], and A. Kopp [1]

[1] Department of Evolution and Ecology, University of California – Davis, Davis CA USA
[2] Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor MI USA
[3] Janelia Research Campus, Howard Hughes Medical Institute, Ashburn VA USA
* Corresponding author: kohtanaka@ucdavis.edu

Abstract

Sexually dimorphic morphological traits are among the fastest evolving animal features. Similar sex-specific structures have sometimes evolved independently in multiple lineages, presumably as targets of parallel sexual selection. In such cases, comparing the cellular mechanisms that generate these structures in different species can elucidate the interplay between selection and developmental constraint in evolution. In Drosophilidae, male-specific tarsal brushes on the front legs are found in at least four separate lineages. In this study, we combine phylogenetic reconstruction with developmental analyses and behavioral observations to investigate the evolutionary origin of these structures. We show that the sex brush has evolved independently at least three times from sexually monomorphic ancestral morphology. However, all sex brushes have very similar fine structure and develop through indistinguishable cellular processes, providing a striking example of developmental convergence. In all examined species, males use their sex brushes to grasp the female abdomen prior to copulation. We discuss potential reasons why convergent evolution of novel structures is rare even in the face of similar functional demands.

Keywords: *Drosophila* phylogeny; sexual dimorphism; convergent evolution; mating behavior; bristle development

Introduction

Most animals are sexually dimorphic. Perhaps the most fascinating feature of sexual dimorphism is the rapid evolutionary turnover of sex-specific traits. Even among close relatives, the characters that distinguish males from females vary greatly from species to species. This simple observation implies that new sexual characters are gained, and ancestral ones are often lost, during the evolution of many if not most animal lineages. Understanding the genetic and developmental basis of this turnover is necessary to shed light on one of the most important drivers of biological diversity. Examples where similar traits have evolved multiple times are of particular value, as they often provide insights into the general patterns and mechanisms of evolution (Kopp 2009).

Most higher Diptera mate with the male on top of the female, and the male front (T1) legs are often involved in grasping or stimulating the female (Huber et al., 2007; McAlpine, 1981). Perhaps for this reason, male-specific ornaments or grasping structures are found on the T1 legs of many dipteran species (Daugeron et al., 2011; Eberhard, 2001; Ingram et al., 2008; Sivinski, 1997). In Drosophilidae, the most obvious male-specific leg modifications include the sex combs found in the *Drosophila melanogaster* and *obscura* species groups and in the genus *Lordiphosa* (Katoh et al., 2018; Kopp, 2011); branched or spoon-shaped tarsi of some Hawaiian *Drosophila* species (Hardy, 1965; Stark and O'Grady, 2009); and tarsal brushes that are the focus of this study (Figure 1). Sex brushes are found in at least four separate groups within the Drosophilidae: the *Drosophila immigrans* species group, the *loiciana* species complex, *D. repletoides*, and the genus *Zaprionus.* Due to uncertain relationships among these lineages, the evolutionary origin(s) of this structure remain unclear.

In the *immigrans* group, male sex brushes are found in some but not all of the species; the most likely scenario is that the brush was present in the last common ancestor of this clade, but was secondarily lost in the *nasuta* subgroup and greatly reduced in several other species (Rice et al., 2018). In *Zaprionus*, male brushes are present in most African species, with the exception of *Z. neglectus*, *Z. spineus*, and *Z. spinosus* (Tsacas and Chassagnard, 1990; Yassin et al., 2008; Yassin and David, 2010). The *Zaprionus* phylogeny is not fully resolved, but the distant relationship between the first species and the last two

suggests that their lack of brushes is likely to reflect independent secondary losses. The
situation is more complicated among species assigned to the Oriental *Anaprionus* subgenus
of *Zaprionus*. Many of its members, including *Z. lineosus, Z. spinilineosus, Z. orissaensis, Z.*
multistriatus, Z. grandis, and *Z. aungsani*, lack leg brushes (Gupta, 1972; Kikkawa and Peng,
1938; Okada and Carson, 1983; Wynn and Toda, 1988). However, *Anaprionus* is now
thought to be polyphyletic (Yassin, 2007; Yassin et al., 2010), and these species appear to
be more closely related to the genus *Xenophorticella* than to *Zaprionus* sensu stricto (M.
Toda, pers. comm.). Other *Anaprionus* species such as *Z. bogoriensis, Z. obscuricornis,* and *Z.*
pyinoolwinensis have leg brushes (Mainx, 1958; Okada, 1964; Wynn and Toda, 1988) and
likely form a clade with the Afrotropical *Zaprionus* (M. Toda, pers. comm.). Thus, the leg
brush has evolved either at or near the base of *Zaprionus*.

D. pruinosa belongs to the *loiciana* species complex, which also includes *D. loiciana*,
D. allochroa, *D. pachneissa*, *D. semipruinosa*, and *D. xanthochroa*. All of these species have
male leg brushes of different sizes (Tsacas, 2002; Tsacas and Chassagnard, 2000). The
fourth lineage where a male leg brush is found consists of a single species, *D. repletoides,*
which does not have any known close relatives; a species described originally as *D.*
tumiditarsus (Tan et al., 1949) was later synonymized with *D. repletoides* (Hsu, 1943;
Wheeler, 1981). Yassin (Yassin, 2007) suggested that some species currently classified as
Zaprionus (*Z. multistriatus*, *Z. flavofasciatus*, and *Z. cercociliaris*) could in fact be more
closely related to *D. repletoides* than to *Zaprionus*; unfortunately, these species have not
been included in any molecular phylogenies.

The four clades of interest – *Zaprionus*, the *immigrans* species group, *D. repletoides*,
and the *loiciana* complex – have never been included together in the same molecular
phylogeny. Different combinations of these taxa have been examined in several
phylogenetic studies, which were based on a small number of loci and produced different
results. Da Lage et al (Da Lage et al., 2007) and Yassin et al (Yassin et al., 2010) provided
some evidence for a distant relationship among *Zaprionus*, *D. repletoides*, and the
immigrans species group. Russo et al (Russo et al., 2013) placed *D. pruinosa* as sister to *D.*
sternopleuralis (a member of the *histrio* species group that lacks a sex brush), and the
resulting clade as sister to the *immigrans* species group. The phylogenies of Da Lage et al
(Da Lage et al., 2007) and Izumitani et al (Izumitani et al., 2016) did not include *D. pruinosa*,

but did not support a sister-group relationship between *D. sternopleuralis* and the
immigrans species group.

In this study, we used a larger multilocus dataset to test whether the male leg brush
evolved independently in each of these four clades, or whether its distribution could be
better explained by shared origin in some of these lineages. To facilitate this analysis, we
included one or more representatives of each clade, as well as several brush-less species
that have been suggested by previous studies to be closely related to the brush-bearing
clades. In parallel, we compared the cellular mechanisms that produce the male leg
brushes in different species, as well as the role of these ornaments in mating behavior. Our
results strongly suggest that the sex brushes evolved convergently in several distantly
related lineages, but develop through virtually identical mechanisms.

117
118
119 **Materials and Methods**
120
121 *Leg imaging*
122
123 For brightfield imaging, male front legs were dissected, mounted in Hoyers media between
124 two coverslips, and photographed on a Leica DM500B microscope with a Leica DC500
125 camera. For electron microscopy, adult legs were dehydrated in 100% ethanol, critical
126 point dried, and coated with gold. Scanning electron micrographs were taken on Thermo
127 Fisher Quattro S and Philips XL30 TMP.
128
129 *Sequence data collection*
130
The sources of live *Drosophila* strains, fixed specimens, and unpublished genome
assemblies used in this study are listed in Table S1. DNA was extracted from live or
alcohol-fixed flies using an affinity resin based protocol (Hi Yield® Genomic DNA Mini Kit,
Süd-Laborbedarf Gauting, Germany). PCR was carried out using DreamTaq polymerase
(Thermo Fisher) and the following cycling conditions: 95° 5' => (95° 30" => 55° 30" => 72°
80")x35 => 72° 5' =>12°; the loci and primer sequences are listed in Table S2. In some

cases, two rounds of PCR with nested primers were needed to obtain amplicons from fixed

specimens. Amplified fragments were gel-purified and sequenced from both ends using

amplification primers. Sequence chromatograms were trimmed in SnapGene Viewer, and

the two end reads were aligned and edited in Geneious. Heterozygous nucleotide positions,

if present, were represented by IUPAC ambiguity codes. All new sequences were deposited

in Genbank under accession number listed in Table S1. Additional sequences were

obtained from Genbank or extracted from whole-genome assemblies using Blast v2.2.23

(Table S1).

Sequence analysis

The sequences of each locus were aligned using the MUSCLE algorithm (Edgar, 2004)

implemented in Geneious (Kearse et al., 2012). The alignments were trimmed at the ends,

and poorly aligning intronic regions were removed. The alignments of all eight loci were

then concatenated for combined analysis. Combined Bayesian analysis was carried out in

MrBayes v3.2.6 (Ronquist et al., 2012). Two sets of analyses were conducted. In the first,

the dataset was partitioned by gene, and each locus was allowed to follow a different

nucleotide substitution model with empirically estimated parameters. In the second, we

partitioned the dataset by gene and by codon position, and used PartitionFinder with the

PhyML algorithm (Guindon et al., 2010; Lanfear et al., 2012; Lanfear et al., 2017) to identify

the appropriate partitioning scheme; this resulted in a total of 21 character subsets, each of

which was subsequently allowed to follow its own substitution model with empirically

estimated parameters. For each analysis, two parallel runs of 1,500,000 generations, each

starting from a different random tree, were carried out, and convergence was confirmed by

comparing tree likelihoods and model parameters between the two runs. *D. melanogaster*

was used as outgroup. Trees were sampled every 1000 generations and summarized after

a 20% relative burn-in. Each analysis was also repeated after excluding *D. quadrilineata*

(see Results). Samples of probable trees were extracted from the .tprobs file, and a strict

consensus of most probable trees with combined posterior probabilities of 95% or 99%

was constructed from these sets of trees in Geneious. Consensus trees were then formatted

using FigTree v1.3.1 (http://tree.bio.ed.ac.uk/software/figtree/).

Immunocytochemistry and microscopy

Fly cultures were raised on standard *Drosophila* media at room temperature. Since each
species develops at a different rate, the timing of pupal stages was determined empirically
based on the morphology of transverse bristle rows (TBRs). Each species was imaged at an
early stage when TBR bristles of the tibia and the first tarsal segment are separated by one
or more intervening epithelial cells; this stage corresponds to 16 - 21 hours after
pupariation (AP) in *D. melanogaster*. Each species was also imaged at a late stage, after the
packing of TBRs is completed and bristle shaft differentiation is underway (corresponding
to 36+ hrs AP in *D. melanogaster*). Pupal legs were dissected, processed and
immunostained as in Tanaka et al (2009). The primary antibodies used were rat anti-E-
cadherin (DCAD2, from the Developmental Studies Hybridoma Bank, at 1:20) for *D.
pruinosa* and *Z. tuberculatus,* and mouse anti-Armadillo (N2 7A1, DSHB; 1:30) for *D.
immigrans* and *D. repletoides*. AlexaFluor 488 secondary antibodies (Invitrogen) were used
at 1:400. Fluorescent images were taken on an Olympus 1000 confocal microscope and
processed using Image J and Adobe Photoshop.

Analysis of mating behavior

High-speed videos of mating behavior were recorded using a Fascam Photron SA4 mounted
with a 105 mm AF Micro Nikkor lens. In brief, individual virgin males were isolated upon
eclosion in food vials and aged for up to two weeks. Virgin females were isolated upon
eclosion and housed in groups of 20-30. Pairs of males and females were then gently
aspirated into single wells of a 96 well culture plate (Corning 05-539-200) filled halfway
with a hardened 2% agarose solution and sealed using a glass microscope slide and tape.
Video clips were captured at 1000 frames per second (fps) using Photron Fastcam Viewer
software.

Results

Sex brush morphology is highly similar in distantly related species

In all examined species of *Zaprionus*, the *immigrans* species group, *loiciana* species complex, and *D. repletoides*, the male-specific leg brush is found on the anterior-ventral surface of the first (ta1) and sometimes also the second (ta2) tarsal segment of the prothoracic leg. In females and in other species, this area is occupied by transverse bristle rows (TBRs), which the flies use to clean their head and eyes (Tokunaga 1962; Kopp 2011). The male brush replaces the distal TBRs, with a few TBRs remaining at the proximal end of ta1 (Figure 1). In the *immigrans* group (Rice et al. 2018) and in the other species (Figure 1), the brush shows several major differences from the TBRs. The bristles of each TBR are aligned into a straight, tightly packed row that is nearly perpendicular to the proximo-distal (PD) leg axis, while the consecutive TBRs along the PD axis are separated by many cell diameters. In contrast, the modified bristles ("hairs") that make up the brush are packed closely together in all directions and do not show any regular arrangement. The shafts of the TBR bristles are robust and straight, with ridges and grooves running their length and a triangular bract at the base; the brush hairs are thin and wavy with a smooth surface, and lack bracts (Figure 1).

Although the morphology of the male leg brush is similar across species, detailed scanning electron microscopy (SEM) analysis reveals a small but consistent difference. In *D. immigrans* and *D. pruinosa,* the tips of brush hairs are thin and flat, taper to a point, and form hooks that curve toward the base of the leg (Figure 1B, D). In *D. repletoides* and in *Zaprionus*, the hair tips are noticeably thicker and curve away from the leg base (Figure 1 F, H, J, L); in *Zaprionus*, they are also flattened into paddle-like shapes (Figure 1 H, J, L). Thus, while the spatial arrangement of bristles appears to be similar in all species, some differences exist in the morphology of bristle shafts. These differences may reflect the phylogenetic relationships among these taxa, especially the close relationship between *D. pruinosa* and the *immigrans* species group (see next section).

Male leg brushes evolved independently at least three times

We sequenced partial coding sequences of eight nuclear, protein-coding loci: *acon,*

eno, glyp, Amyrel, Ddc, Gpdh, Pepck, and *Pgm*. Separate analyses of each locus produced

very poorly resolved trees. We therefore combined the data from all loci (up to 9060

nucleotides per species) for a partitioned Bayesian analysis where each locus was allowed

to follow its own, empirically estimated substitution model but all loci were constrained to

the same tree topology. The resulting tree (Figure 2A; brush-bearing clades labeled in

blue) suggests a close relationship of *D. pruinosa* to the *immigrans* species group, with the

(*D. sternopleuralis* + *D. trisetosa*) clade, which belongs to the *histrio* species group, as the

next outgroup. In contrast, *D. repletoides* is placed as sister group to the (*D. busckii* + *D.*

brachytarsa) clade, well away from the *immigrans-pruinosa* lineage. Finally, the *Zaprionus*

genus is distantly related to both *D. repletoides* and the *immigrans-pruinosa* lineage, and is

placed near the base of the tree, separately from the *Drosophila* and *Sophophora* subgenera.

We also note that *D. curviceps* and *D. annulipes* appear as sister groups with 100% support,

while there is no support for clustering the *immigrans* species group either with the (*D.*

curviceps + *D. annulipes*) clade or with *D. quadrilineata* (see Supplementary text). A strict

consensus of 11 trees with the cumulative posterior probability of 95% is not resolved near

the base, but does not support a close relationship among the brush-bearing lineages: *D.*

repletoides, the *immigrans-pruinosa* clade, and *Zaprionus* (Figure S1).

We then examined 27 most probable trees, with the cumulative posterior

probability of 99%, to determine the probability of each bipartition of interest to our study.

The *Zaprionus* genus (occasionally together with *D. quadrilineata*; see below) was placed at

the base of the tree, well separated from *D. repletoides* and the *immigrans-pruinosa* lineage

(Figure 2A, Table S3). *D. pruinosa* clustered with the *immigrans* species group, to the

exclusion of the (*D. sternopleuralis* + *D. trisetosa*) clade, with ~87% probability; the

alternative grouping, of *D. pruinosa* with the (*D. sternopleuralis* + *D. trisetosa*) clade to the

exclusion of the *immigrans* species group, was observed with ~13% probability. *D.*

repletoides was grouped with the (*D. busckii* + *D. brachytarsa*) clade with 95% probability,

and this group was separated from the *immigrans-pruinosa* lineage by multiple internal

branches. In contrast, potential groupings of brush-bearing lineages – for example, of

Zaprionus with either *D. repletoides* or the *immigrans-pruinosa* clade, or of the *repletoides-*
busckii-brachytarsa clade with the *immigrans-pruinosa* clade – were never observed among
this set of probable trees.

We noticed that among these 27 probable trees, the position of *D. quadrilineata* was
by far the most unstable. We therefore repeated the analysis after excluding this species.
The resulting tree (Figure 2B, Table S3) shows the same relationships as the full analysis
(Figure 2A), but with stronger support for most basal nodes. This tree, as well as the strict
consensus of the 11 most probable trees with the cumulative posterior probability of 99%
(Figure S2) places *Zaprionus* near the base of the tree, separately from the *Drosophila* and
Sophophora subgenera; *D. pruinosa* together with the *immigrans* species group and the (*D.*
sternopleuralis + *D. trisetosa*) clade; and *D. repletoides* in a clade that is clearly separated
both from *Zaprionus* and from the *immigrans-pruinosa* lineage. The probability of sister-
group relationship between *D. pruinosa* and the *immigrans* species groups is estimated at
93.8% ± 1.32% across these trees.

Finally, we carried out an analysis with a more complicated partitioning scheme,
where each locus and each codon position was allowed to follow its own substitution
model; this analysis was also performed with and without *D. quadrilineata*. In both cases, it
produced a tree with the same topology as in the simpler partitioning scheme, but with
slightly different levels of node support (Figure S3, S4 and Table S3). In summary, we find
substantial though not overwhelming support for a close relationship between *D. pruinosa,*
and by implication the *loiciana* species complex, and the *immigrans* species group.
However, based on our data, the probability of a close relationship among the different
brush-bearing clades – *D. repletoides*, *Zaprionus*, and the *immigrans-pruinosa* lineage – is
very low for each of the possible pairwise relationships.

The same cellular processes underlie sex brush development in all species

In *D. melanogaster,* TBR bristle precursors are specified between 6 and 12 hours
after pupariation (AP) (Joshi and Orenic 2006, Schroff 2007). Initially, these cells are
specified in sparse, loosely organized rows, and separated from one another by several
epithelial cells. By 20-21 hrs AP, the bristle cells that are destined to make each TBR

migrate toward each other to form a straight, contiguous row, while the intervening
epithelial cells are expelled distally and proximally from the TBRs (Atallah et al. 2009;
Tanaka et al. 2009). This mechanism suggests two potential explanations for the tight
packing of brush hairs. One possibility is that the hair progenitor cells are specified with
minimal spacing; in this case, the initial spacing between hair cells is expected to be much
denser than the spacing of TBR progenitors. Alternatively, hair progenitors may first be
specified similarly to TBR bristles, with wide separation by epithelial cells, but expel the
epithelial cells (either laterally or basally) at later stages to form a densely packed brush.
In principle, independently evolved brushes in different species could utilize different
cellular mechanisms to produce adult structures that are essentially indistinguishable.

In order to characterize and compare brush development in different species, we
used antibodies against membrane-localized proteins to visualize cell arrangement in
pupal legs. When labeled with antibodies against the beta-catenin Armadillo (Arm) or the
E-cadherin Shotgun (DE-cad), bristle cells can be distinguished from the surrounding
epithelial cells by their unique membrane shape (Figure 3). We examined brush
development at two timepoints: an early stage roughly corresponding to ~16-21 hr AP in *D.*
melanogaster, when the bristle cells of the future TBRs begin to migrate toward each other
and expel the intervening epithelial cells, and a later stage when cell migration is
completed. We found that at the early stage, most bristle cells in the developing brush are
each surrounded by four to six epithelial cells in all species. In effect, the bristle cells are
separated from one another by one to two epithelial cells (Figure 3). At the late stage, this
spacing remains virtually invariant, although the cells appear more organized compared to
the early stage (Figure 3). We did not see evidence of cell migration in any of the examined
species. These observations indicate that despite the packed appearance similar to TBRs,
the brush cells are not directly adjacent to each other, are specified at high density, and
undergo minimal if any migration during development. Importantly, the cellular
mechanism of brush development is very similar in all species.
319
320
321
322

Leg brushes are used in male mating behavior

High-speed video recordings show that the proximal tarsal segments of the male T1 legs,
including the sex brushes, are used to grab the female abdomen and resist the female's
efforts to dislodge the male during copulation attempts in *Z. tuberculatus*, *D. immigrans*,
and *D. repletoides* (Supplementary movies 1-3); unfortunately, no mating attempts were
observed in *D. pruinosa*. In the *Drosophila melanogaster* species group, males use their sex
combs, which are also located on the T1 tarsus, to grab the female genitalia as in *D.*
melanogaster, or the middle abdominal segments as in *D. kikkawai, D. ananassae*, and *D.*
bipectinata (Massey et al., 2019). In these species, male T2 and T3 legs remain on the
substrate and are not involved in mating. Typically, these males proceed very quickly from
mounting to attempted copulation; females may resist by walking away or using their hind
legs to kick the male off. Interestingly, similar behavior is observed in *D. willistoni*, one of
the closest relatives of the *melanogaster* species group that lacks sex combs (Massey et al.,
2019), suggesting that the origin of the sex comb did not radically change this aspect of
mating. In contrast, *D. immigrans, Z. tuberculatus*, and *D. repletoides* males use their sex
brushes to grab females more anteriorly, near the constriction between the thorax and
abdomen (Supplementary movies 1-3). In the former two species, males also use their T2
legs to grab the female mid-abdomen, while in *D. repletoides* T1, T2 and T3 legs are all used
to grab the female so that the male "rides" on the female and is not in contact with the
substrate. In all these species, females appear to resist mating attempts more vigorously
than in the *melanogaster* group, using side-to-side bucking and wing vibrations in apparent
efforts to dislodge the male, while the males use their legs to resist these efforts. The delay
between mounting and attempted copulation is longer in the brush-bearing species,
especially in *Z. tuberculatus*, than in the comb-bearing species; most mountings result in
the male being eventually dislodged and do not lead to copulation attempts.

A more systematic analysis, including many lineages that lack male-specific leg
modifications, will be needed to test whether morphological evolution correlates with the
evolution of behavior. At this point, we can only speculate that male leg brushes, which
consist of hundreds of thin hairs that are hooked at the tips and have a very large combined

surface area, may have evolved to provide a more secure grip of the female abdomen,
especially if stronger grip is needed to counteract the female attempts to dislodge the male.
355
356
357   **Discussion**
358
359   *Repeated evolution of leg brushes*
360
Our results suggest that the male-specific brush on the tarsal segments of front legs
has evolved independently at least three times in Drosophilidae: in *Zaprionus*, in *D.*
repletoides, and in the common ancestor of the *immigrans* species group and the *loiciana*
complex, which includes *D. pruinosa*. Although our study does not provide unequivocal
support for a sister-group relationship between *D. pruinosa* and the *immigrans* group (a
closer relationship of *D. pruinosa* to the *sternopleuralis-trisetosa* clade cannot be completely
ruled out, and of course there may exist other closely related species that we have not
sampled), independent evolution of brushes in the *immigrans* species group and the
loiciana species complex appears less likely than a single origin in a common ancestor of
these clades. The proximally curving tips of brush hairs in *D. pruinosa* and *D. immigrans*,
distinct from the distally curving tips in *Zaprionus* and *D. repletoides* (Fig. 1), are consistent
with a close relationship between the *immigrans* group and the *loiciana* species complex.
Convergent origin of leg brushes in *Zaprionus*, *D. repletoides*, and the *immigrans-*
loiciana clade is remarkable given the strong structural similarities of these brushes, and
especially the fact that the cellular mechanisms that produce them in different species are
essentially identical. In all species examined, the bristles that make up the sex brush are
specified with only one or two intervening epithelial cells between them.   Bristle
specification in *Drosophila* and other insects is governed by a lateral inhibition mechanism,
which is based on contact signaling between adjacent cells, and prevents two adjacent cells
from both assuming the fate of bristle precursors (Simpson, 1990). Later in development, a
tighter packing of bristles can be achieved through cell migration, as observed for example
in the transverse bristle rows of the front legs (Atallah et al., 2009; Tanaka et al., 2009).
However, no cell migration is observed during sex brush development; instead, the future

brush hairs are always specified at the maximum density allowed by lateral inhibition.
Independent evolution of this highly derived spatial pattern in multiple lineages suggests
that similar selective pressures may elicit not only similar structures, but also similar
changes in the underlying developmental mechanisms.
388
389   *A hierarchy of convergence, from structures to genes*
390
Convergence is a pervasive phenomenon in evolution. Pictures comparing bats and
birds, fish and dolphins, or the vertebrate and cephalopod eyes are textbook clichés. Less
obvious but equally striking examples of convergent forms shaped by a common function
include the asymmetrical ears of owls (Nishikawa, 2002), Mullerian mimicry in butterflies
(Brower, 1994; Naisbit et al., 2003), trophic morphology in cichlid fishes (Ruber and Adams,
2001), and many others (Moore and Willmer, 1997). A fundamental question raised by the
widespread occurrence of convergent traits is to what extent phenotypic convergence
reflects an underlying similarity of the molecular and cellular mechanisms that control
individual development.
The mechanisms of convergence can be viewed in either genetic or developmental
terms. From the genetic standpoint, convergent phenotypes are controlled by the same
mechanism if they are the result of changes in the same loci ("genetic convergence")
(Gompel and Prud'homme, 2009; Martin and Orgogozo, 2013). Alternatively, convergent
changes may be considered to share a similar mechanism if they are produced by the same
changes in development, regardless of the ultimate genetic causes ("developmental
convergence") (Wake, 1991; Wake et al., 2011; Wray, 2002). We can describe convergence
in hierarchical terms by applying three increasingly stringent criteria: phenotypic,
developmental, and genetic. Phenotypic convergence can occur without developmental
convergence: for example, the disproportionally elongated bodies of burrowing
salamanders can be caused either by an increase in the number of vertebrae, or by
increased length of the individual vertebrae – a superficial phenotypic resemblance
produced by entirely different developmental mechanisms (Parra-Olea and Wake, 2001).
Similarly, larval abdominal prolegs of butterflies and sawflies, while superficially similar
and playing similar roles in locomotion, develop from non-homologous leg segments

(Suzuki and Palopoli, 2001). In *D. melanogaster*, latitudinal clines in wing size have evolved

on several continents, but the differences in wing size are due largely to differences in cell

number in some populations, while in other populations differences in the size of

individual cells make a larger contribution (James et al., 1997; Zwaan et al., 2000).

Developmental convergence, in turn, does not necessarily reflect genetic

convergence. In *Drosophila*, similar color patterns can evolve in different species through

changes at different loci (Signor et al., 2016; Wittkopp et al., 2003b; Yassin et al., 2016), but

in the end all of these genetic changes are translated into adult phenotypes through the

same set of enzymatic reactions in the catecholamine metabolism pathway (Kronforst et al.,

2012; Wittkopp and Beldade, 2009; Wittkopp et al., 2003a), presenting an example of

developmental convergence in the absence of genetic convergence. In other cases, the

evolution of similar phenotypes in distantly related species is clearly caused by

independent changes in the same genes – as, for example, in the loss of larval cuticular

hairs through regulatory mutations in the *shavenbaby* gene in two *Drosophila* species

separated by >40 million years (Sucena et al., 2003), or in the evolution of mimetic wing

color patterns in different species of *Heliconius* butterflies through parallel changes in the

optix and *WntA* genes (Gallant et al., 2014; Hines et al., 2011; Martin et al., 2012; Reed et al.,

2011; Supple et al., 2013).

Going even deeper from the genetic to the molecular level, we can ask whether

recurrent involvement of the same gene in multiple instances of convergent evolution

reflects repeated appearance of the same mutations, or whether different mutations can

alter the function of the causative gene in similar ways, leading to the evolution of similar

phenotypes. Again, we can find examples in support of either scenario. A variety of

proteins have evolved similar amino acid residues and active sites repeatedly in different

taxa (Gasparini et al., 1998; Hill et al., 2019; Kriener et al., 2000; Lopreato et al., 2001;

Yokoyama, 2002). Although non-coding DNA sequences are clearly less constrained in

their function than proteins, similar regulatory mutations can nevertheless appear and be

fixed independently in different populations or species in response to similar selective

pressures (Chan et al., 2010; Loehlin et al., 2019; Xie et al., 2019). Conversely, mutations

that affect different amino acids, or different *cis*-regulatory elements, can produce similar

phenotypic outcomes in different populations or species (Gross et al., 2009; Kingsley et al.,

2009; Manceau et al., 2010; Protas et al., 2006; Rosenblum et al., 2009; Yassin et al., 2016).

The sex brush represents a clear case of developmental convergence, but the genetic

control of its development remains to be determined. Specification and positioning of the

sex brush are likely to involve the HOX gene *Scr* and the sex determination gene *doublesex*

(Rice et al., 2018; Tanaka et al., 2011). However, the identity of the downstream genes that

translate the regulatory information provided by *dsx* and *Scr* into the final 3-dimensional

structure is unknown. In principle, these genes need not be the same in different lineages.

Future work may show whether the convergence we observe at the level of cell behavior

extends to the genetic and molecular level.

Convergent innovation and the failure to innovate

The evolution of similar structures in response to similar functional demands is perhaps

not surprising; should we instead be surprised when organisms faced with similar

demands *fail to evolve* convergent phenotypes (Blount et al., 2018)? Males grasp female

abdomen with their front legs in many *Drosophila* species, including those that primitively

lack any male-specific leg ornaments (Massey et al., 2019; Spieth, 1952). Why don't all

Drosophila species evolve sex combs, brushes, or other grasping structures? Development

does not provide any clues. At the level of cell behavior, both the ancestral/female

condition (cell migration that produces tightly packed bristle rows from sparsely spaced

precursors) and the derived/male condition (specification of bristle precursors at the

maximum density permitted by lateral inhibition) are the same in all lineages where the

sex brush is present. There is no a priori reason to think that the transition between these

modes of development is easier in some species than others.

The answer may lie instead in either behavior or population genetics. Although

males of different species use their sex brushes in at least superficially similar ways, we

don't know the female side of the story. If females of different species vary in their

responses to male grasping, the evolution of specialized leg structures in males may not be

universally favored. This may also explain why both the sex brushes and the sex combs

(Kopp, 2011) have been secondarily lost multiple times. Moreover, it is difficult to know

whether the female preferences observed today are the same as they were in the distant
past when the male-specific structures evolved (Watts et al., 2019).

Alternatively, the origin of a new trait such as the leg brush may require such an
unlikely series of genetic changes that it may often fail to occur even in response to strong
selective pressure. For example, it is possible that while a single mutation is sufficient to
modify or eliminate an existing morphological structure, the origin of a *new* structure may
require simultaneous changes in multiple genes. From the population-genetic perspective,
this would mean that functionally novel and positively interacting alleles at multiple loci
must segregate in the same population at the same time in order for selection in favor of a
new structure to be effective. Naturally, this would greatly reduce the probability of
evolutionary innovations compared to other types of phenotypic change. This is of course
pure speculation; we do not know why convergent innovations evolve in some lineages but
fail to evolve in others. We hope that research models where both the functional roles and
the genetic basis of novel traits can be studied in parallel will shed some light on this
intriguing question in the future.

Acknowledgements

We thank the US *Drosophila* species stock center, Ehime University *Drosophila* stock center,
J.-R. David, S. Prigent, and M. Watada for *Drosophila* strains and specimens; Developmental
Studies Hybridoma Bank for antibodies; A. Comeault, A. Gloss, M. Lang, D. Matute, D. Miller,
V. Orgogozo, and N. Whiteman for genome assemblies; M. Toda for advice on *Zaprionus*
phylogeny; G. Rice for *D. immigrans* images; and A. Yassin for comments on the manuscript.
We also thank the MCB Light Microscopy Imaging Facility, Department of Cell Biology and
Human Anatomy EM Lab, and Materials Science and Engineering AMCaT Lab at UC Davis
for equipment and imaging assistance, and Gwyneth Card and W. Ryan Williamson for use
of the Photron camera. The Olympus FV1000 confocal used in this study was purchased
using NIH Shared Instrumentation Grant 1S10RR019266-01. AK and OB are grateful to N.
Gompel for allowing some of this work to be conducted in his lab while on a sabbatical visit,
and to members of the Gompel lab for treating them as their labmates. This work was

supported by NIH grant 5R35GM122592 to AK and by funds from Ludwig-Maximilians-

Universität München to Nicolas Gompel.

 Supplemental text - The *immigrans* species group

The *immigrans* species group and the wider *immigrans-tripunctata* radiation have long

played a pivotal role in *Drosophila* systematics (Throckmorton, 1975; Yassin, 2013).

However, the composition of the *immigrans* group has not been entirely clear. Historically,

this group was proposed to include five subgroups: *immigrans, hypocausta, nasuta,*

quadrilineata, and *curviceps* (Huang et al., 2002; Zhang and Toda, 1988, 1992; Zhang et al.,

1995). Recent studies have shown, however, that both the *hypocausta* and the *immigrans*

subgroups are likely polyphyletic, as some species assigned to these subgroups are actually

closer to the *nasuta* subgroup (Da Lage et al., 2007; Katoh et al., 2007; Rice et al., 2018).

The *quadrilineata* and *curviceps* subgroups have been the source of even greater

complications. Previous phylogenetic studies have provided evidence against a sister-

group relationship between *D. quadrilineata* and the rest of the *immigrans* species group

(Katoh et al., 2007; Morales-Hojas and Vieira, 2012; Yassin, 2013). Moreover, *D. annulipes*,

traditionally assigned to the *quadrilineata* subgroup (Lin and Tseng, 1973), was found to be

distantly related to *D. quadrilineata*, and closer to the *virilis-repleta* radiation (Katoh et al.,

2007). The phylogeny of Huang et al (Huang et al., 2002) showed a close relationship

between *D. annulipes* and the clade composed of *D. curviceps* and *D. oritisa*, another species

assigned to the *curviceps* subgroup. Morales-Hojas and Vieira (Morales-Hojas and Vieira,

2012) confirmed the distant relationship between *D. annulipes* and *D. quadrilineata*, as well

as between *D. annulipes* and the *immigrans* species group *sensu stricto*. Yassin (Yassin,

2013) pointed out important differences between *D. quadrilineata* and the *curviceps*

subgroup and the *immigrans* group *s.s.* Finally, Pradhan et al (Pradhan et al., 2015)

removed the *curviceps* subgroup from the *immigrans* species group, elevating it to the

status of a separate species group.

Our analysis, based on a larger amount of sequence data than previous studies,

confirms these observations, namely that (1) *D. quadrilineata* is not closely related either to

the *immigrans* species group *s.s.* or to *D. annulipes*, (2) *D. curviceps* is not closely related to

the *immigrans* species group *s.s.*, and (3) *D. annulipes* is related to *D. curviceps*. Based on

these results, we support the conclusions on other authors (Katoh et al., 2007; Pradhan et

al., 2015; Yassin, 2013) that the definition of the *immigrans* species group should be

restricted to the *immigrans*, *hypocausta*, and *nasuta* subgroups. Further work, with a more

extensive sampling of the *immigrans* group *s.s.*, is needed to revise its internal taxonomy

and re-establish monophyletic subgroups. Similarly, better taxon sampling will be needed

to determine whether *D. annulipes* could be reassigned to the *curviceps* species group

(Pradhan et al., 2015), or whether it is possible to establish a monophyletic *quadrilineata*

species group.

References

Atallah, J., Liu, N.H., Dennis, P., Hon, A., Godt, D., and Larsen, E.W. (2009). Cell dynamics and developmental bias in the ontogeny of a complex sexually dimorphic trait in Drosophila melanogaster. Evol Dev *11*, 191-204.

Blount, Z.D., Lenski, R.E., and Losos, J.B. (2018). Contingency and determinism in evolution: Replaying life's tape. Science *362*.

Brower, A.V. (1994). Rapid morphological radiation and convergence among races of the butterfly Heliconius erato inferred from patterns of mitochondrial DNA evolution. Proc Natl Acad Sci U S A *91*, 6491-6495.

Chan, Y.F., Marks, M.E., Jones, F.C., Villarreal, G., Jr., Shapiro, M.D., Brady, S.D., Southwick, A.M., Absher, D.M., Grimwood, J., Schmutz, J*., et al.* (2010). Adaptive evolution of pelvic reduction in sticklebacks by recurrent deletion of a Pitx1 enhancer. Science *327*, 302-305.

Da Lage, J.-L., Kergoat, G.J., Maczkowiak, F., Silvain, J.-F., Cariou, M.-L., and Lachaise, D. (2007). A phylogeny of Drosophilidae using the *Amyrel* gene: questioning the *Drosophila melanogaster* species group boundaries. J Zool Syst Evol Res *45*, 47-63.

Daugeron, C., Plant, A., Winkler, I., Stark, A., and Baylac, M. (2011). Extreme male leg polymorphic asymmetry in a new empidine dance fly (Diptera: Empididae). Biology letters *7*, 11-14.

Eberhard, W.G. (2001). The functional morphology of species-specific clasping structures on the front legs of male Archisepsis and Palaeosepsis flies (Diptera, Sepsidae). Zoological Journal of the Linnean Society *133*, 335-368.

Edgar, R.C. (2004). MUSCLE: a multiple sequence alignment method with reduced time and space complexity. BMC Bioinformatics *5*, 113.

Gallant, J.R., Imhoff, V.E., Martin, A., Savage, W.K., Chamberlain, N.L., Pote, B.L., Peterson, C., Smith, G.E., Evans, B., Reed, R.D*., et al.* (2014). Ancient homology underlies adaptive mimetic diversity across butterflies. Nature communications *5*, 4817.

Gasparini, S., Danse, J.M., Lecoq, A., Pinkasfeld, S., Zinn-Justin, S., Young, L.C., de Medeiros, C.C., Rowan, E.G., Harvey, A.L., and Menez, A. (1998). Delineation of the functional site of alpha-dendrotoxin. The functional topographies of dendrotoxins are different but share a conserved core with those of other Kv1 potassium channel-blocking toxins. J Biol Chem *273*, 25393-25403.

Gompel, N., and Prud'homme, B. (2009). The causes of repeated genetic evolution. Dev Biol *332*, 36-47.

Gross, J.B., Borowsky, R., and Tabin, C.J. (2009). A novel role for Mc1r in the parallel evolution of depigmentation in independent populations of the cavefish Astyanax mexicanus. PLoS Genet *5*, e1000326.

Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol *59*, 307-321.

Gupta, J.P. (1972). D. orissaensis, a new species of Drosophila from Orissa, India. Oriental Insects *6*, 561-563.

Hardy, D.E. (1965). Diptera: Cyclorrhapha II, series Schizophora, section Acalypterae I, family Drosophilidae. . In Insects of Hawaii, E.C. Zimmerman, ed. (Honolulu: University of Hawaii Press).

Hill, J., Enbody, E.D., Pettersson, M.E., Sprehn, C.G., Bekkevold, D., Folkvord, A., Laikre, L.,
Kleinau, G., Scheerer, P., and Andersson, L. (2019). Recurrent convergent evolution at
amino acid residue 261 in fish rhodopsin. Proc Natl Acad Sci U S A *116*, 18473-18478.
Hines, H.M., Counterman, B.A., Papa, R., Albuquerque de Moura, P., Cardoso, M.Z., Linares,
M., Mallet, J., Reed, R.D., Jiggins, C.D., Kronforst, M.R.*, et al.* (2011). Wing patterning gene
redefines the mimetic history of Heliconius butterflies. Proc Natl Acad Sci U S A *108*, 19666-
19671.
Hsu, T.C. (1943). Chromosomes of Drosophila repletoides. Kwangsi Agric *4*, 155-160.
Huang, J., Hao, L., Liu, S., Li, L., Zhang, W.-X., and Dai, Z.-H. (2002). Phylogenetic position of
Chinese endemic Drosophila curviceps species subgroup in the Drosophila immigrans
group. Acta Genetica Sinica *29*, 417-423.
Huber, B.A., Sinclair, B.J., and Schmitt, M. (2007). The evolution of asymmetric genitalia in
spiders and insects. Biol Rev *82*, 647-698.
Ingram, K.K., Laamanen, T., Puniamoorthy, N., and Meier, R. (2008). Lack of morphological
coevolution between male forelegs and female wings in Themira (Sepsidae : Diptera :
Insecta). Biological Journal of the Linnean Society *93*, 227-238.
Izumitani, H.F., Kusaka, Y., Koshikawa, S., Toda, M.J., and Katoh, T. (2016). Phylogeography
of the Subgenus Drosophila (Diptera: Drosophilidae): Evolutionary History of Faunal
Divergence between the Old and the New Worlds. PLoS ONE *11*, e0160051.
James, A.C., Azevedo, R.B., and Partridge, L. (1997). Genetic and environmental responses to
temperature of Drosophila melanogaster from a latitudinal cline. Genetics *146*, 881-890.
Katoh, T., Nakaya, D., Tamura, K., and Aotsuka, T. (2007). Phylogeny of the Drosophila
immigrans species group (Diptera: Drosophilidae) based on Adh and Gpdh sequences.
Zoolog Sci *24*, 913-921.
Katoh, T.K., Zhang, G., Toda, M.J., Zhang, W.X., and Gao, J.J. (2018). The Lordiphosa denticeps
species group (Diptera: Drosophilidae) in China, with redescriptions of four known species
and descriptions of nine new species. Zootaxa *4471*, 37-75.
Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper,
A., Markowitz, S., Duran, C.*, et al.* (2012). Geneious Basic: an integrated and extendable
desktop software platform for the organization and analysis of sequence data.
Bioinformatics *28*, 1647-1649.
Kikkawa, H., and Peng, F.T. (1938). Drosophila species of Japan and adjacent localities.
Japanese Journal of Zoology *7*, 507-552.
Kingsley, E.P., Manceau, M., Wiley, C.D., and Hoekstra, H.E. (2009). Melanism in peromyscus
is caused by independent mutations in agouti. PLoS ONE *4*, e6435.
Kopp, A. (2011). Drosophila Sex Combs as a Model of Evolutionary Innovations. Evol Dev *13*,
504-522.
Kriener, K., O'HUigin, C., Tichy, H., and Klein, J. (2000). Convergent evolution of major
histocompatibility complex molecules in humans and New World monkeys.
Immunogenetics *51*, 169-178.
Kronforst, M.R., Barsh, G.S., Kopp, A., Mallet, J., Monteiro, A., Mullen, S.P., Protas, M.,
Rosenblum, E.B., Schneider, C.J., and Hoekstra, H.E. (2012). Unraveling the thread of
nature's tapestry: the genetics of diversity and convergence in animal pigmentation.
Pigment cell & melanoma research *25*, 411-433.

Lanfear, R., Calcott, B., Ho, S.Y., and Guindon, S. (2012). Partitionfinder: combined selection
of partitioning schemes and substitution models for phylogenetic analyses. Mol Biol Evol *29*,
1695-1701.
Lanfear, R., Frandsen, P.B., Wright, A.M., Senfeld, T., and Calcott, B. (2017). PartitionFinder
2: New Methods for Selecting Partitioned Models of Evolution for Molecular and
Morphological Phylogenetic Analyses. Mol Biol Evol *34*, 772-773.
Lin, F.-J., and Tseng, H.-C. (1973). The Drosophila immigrans species group in Taiwan with
descriptions of five new species. Bull Inst Zool, Academia Sinica *12*, 13-26.
Loehlin, D.W., Ames, J.R., Vaccaro, K., and Carroll, S.B. (2019). A major role for noncoding
regulatory mutations in the evolution of enzyme activity. Proc Natl Acad Sci U S A *116*,
12383-12389.
Lopreato, G.F., Lu, Y., Southwell, A., Atkinson, N.S., Hillis, D.M., Wilcox, T.P., and Zakon, H.H.
(2001). Evolution and divergence of sodium channel genes in vertebrates. Proc Natl Acad
Sci U S A *98*, 7588-7592.
Mainx, F. (1958). Zaprionus bogoriensis spec. nov.,eine neue Drosophilide aus Java.
Zoologischer Anzeiger *161*, 126-130.
Manceau, M., Domingues, V.S., Linnen, C.R., Rosenblum, E.B., and Hoekstra, H.E. (2010).
Convergence in pigmentation at multiple levels: mutations, genes and function. Philos T R
Soc B *365*, 2439-2450.
Martin, A., and Orgogozo, V. (2013). The Loci of repeated evolution: a catalog of genetic
hotspots of phenotypic variation. Evolution *67*, 1235-1250.
Martin, A., Papa, R., Nadeau, N.J., Hill, R.I., Counterman, B.A., Halder, G., Jiggins, C.D.,
Kronforst, M.R., Long, A.D., McMillan, W.O.*, et al.* (2012). Diversification of complex
butterfly wing patterns by repeated regulatory evolution of a Wnt ligand. Proc Natl Acad
Sci U S A *109*, 12632-12637.
Massey, J.H., Chung, D., Siwanowicz, I., Stern, D.L., and Wittkopp, P. (2019). The yellow gene
influences Drosophila male mating 1 success through sex comb melanizatio. eLife *in press*.
McAlpine, J.F. (1981). Morphology and terminology - adults. In Manual of Nearctic Diptera,
J.F. McAlpine, B.V. Peterson, G.E. Shewell, H.J. Teskey, J.R. Vockeroth, and D.M. Wood, eds.
(Ottawa: Research Branch, Agriculture Canada), pp. 9--63.
Moore, J., and Willmer, P. (1997). Convergent evolution in invertebrates. Biol Rev Camb
Philos Soc *72*, 1-60.
Morales-Hojas, R., and Vieira, J. (2012). Phylogenetic patterns of geographical and
ecological diversification in the subgenus Drosophila. PLoS ONE *7*, e49552.
Naisbit, R.E., Jiggins, C.D., and Mallet, J. (2003). Mimicry: developmental genes that
contribute to speciation. Evol Dev *5*, 269-280.
Nishikawa, K.C. (2002). Evolutionary convergence in nervous systems: insights from
comparative phylogenetic studies. Brain Behav Evol *59*, 240-249.
Okada, T. (1964). Drosophilidae (Diptera) of Southeast Asia collected by the Thai-Japanese
Biological Expedition 1961-62. In Nature and Life in Southeast Asia, T.a.U. Kira, T., ed.
(Kyoto: Fauna and Flora Research Society), pp. 439-466.
Okada, T., and Carson, H.L. (1983). The genera Phorticella Duda and Zaprionus Coquillett
(Diptera, Drosophilidae) of the Oriental region and New Guinea. Kontyu *51*, 539-553.
Parra-Olea, G., and Wake, D.B. (2001). Extreme morphological and ecological homoplasy in
tropical salamanders. Proc Natl Acad Sci U S A *98*, 7888-7891.

Pradhan, S., Sati, P.C., Fartyal., R.S., Chatterjee, R.N., Sarswat, M., Kandpal, M.C., and Singh, B.K. (2015). Drosophila curviceps Species-Group (Diptera: Drosophilidae) from India, with Description of a New Species and Redescription of a Known Species. Proceedings of the Zoological Society *68*, 178-183.

Protas, M.E., Hersey, C., Kochanek, D., Zhou, Y., Wilkens, H., Jeffery, W.R., Zon, L.I., Borowsky, R., and Tabin, C.J. (2006). Genetic analysis of cavefish reveals molecular convergence in the evolution of albinism. Nat Genet *38*, 107-111.

Reed, R.D., Papa, R., Martin, A., Hines, H.M., Counterman, B.A., Pardo-Diaz, C., Jiggins, C.D., Chamberlain, N.L., Kronforst, M.R., Chen, R., *et al.* (2011). optix drives the repeated convergent evolution of butterfly wing pattern mimicry. Science *333*, 1137-1141.

Rice, G., Barmina, O., Hu, K., and Kopp, A. (2018). Evolving doublesex expression correlates with the origin and diversification of male sexual ornaments in the Drosophila immigrans species group. Evol Dev *20*, 78-88.

Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Hohna, S., Larget, B., Liu, L., Suchard, M.A., and Huelsenbeck, J.P. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Syst Biol *61*, 539-542.

Rosenblum, E.B., Rompler, H., Schoneberg, T., and Hoekstra, H.E. (2009). Molecular and functional basis of phenotypic convergence in white lizards at White Sands. Proc Natl Acad Sci U S A.

Ruber, L., and Adams, D.C. (2001). Evolutionary convergence of body shape and trophic morphology in cichlids from Lake Tanganyika. J Evol Biol *13*, 325.

Russo, C.A., Mello, B., Frazao, A., and Voloch, C.M. (2013). Phylogenetic analysis and a time tree for a large drosophilid data set (Diptera: Drosophilidae). Zoological Journal of the Linnean Society *169*, 765-775.

Signor, S.A., Liu, Y., Rebeiz, M., and Kopp, A. (2016). Genetic Convergence in the Evolution of Male-Specific Color Patterns in Drosophila. Curr Biol *26*, 2423-2433.

Simpson, P. (1990). Lateral inhibition and the development of the sensory bristles of the adult peripheral nervous system of Drosophila. Development *109*, 509-519.

Sivinski, J. (1997). Ornaments in the Diptera. Florida Entomologist *80*, 142-164.

Spieth, H.T. (1952). Mating behavior within the genus Drosophila (Diptera). Bulletin of the American Museum of Natural History *99*, 395-474.

Stark, J.B., and O'Grady, P.M. (2009). Morphological variation in the forelegs of the Hawaiian Drosophilidae. I. The AMC clade. J Morphol.

Sucena, E., Delon, I., Jones, I., Payre, F., and Stern, D.L. (2003). Regulatory evolution of shavenbaby/ovo underlies multiple cases of morphological parallelism. Nature *424*, 935-938.

Supple, M.A., Hines, H.M., Dasmahapatra, K.K., Lewis, J.J., Nielsen, D.M., Lavoie, C., Ray, D.A., Salazar, C., McMillan, W.O., and Counterman, B.A. (2013). Genomic architecture of adaptive color pattern divergence and convergence in Heliconius butterflies. Genome Res *23*, 1248-1257.

Suzuki, Y., and Palopoli, M.F. (2001). Evolution of insect abdominal appendages: are prolegs homologous or convergent traits? Development Genes and Evolution *211*, 486-492.

Tan, C.C., Hsu, T.C., and Sheng, T.C. (1949). Known drosophila species in China with descriptions of twelve new species. In Studies in the genetics of Drosophila VI Articles on genetics, cytology, and taxonomy, J.T. Patterson, ed. (Austin, TX: University of Texas Publications), pp. 196-206.

Tanaka, K., Barmina, O., and Kopp, A. (2009). Distinct developmental mechanisms underlie
the evolutionary diversification of *Drosophila* sex combs. Proceedings of the National
Academy of Sciences *106*, 4764-4769.
Tanaka, K., Barmina, O., Sanders, L.E., Arbeitman, M.N., and Kopp, A. (2011). Evolution of
Sex-Specific Traits through Changes in HOX-Dependent doublesex Expression. PLoS Biol *9*,
e1001131.
Throckmorton, L.H. (1975). The phylogeny, ecology, and geography of Drosophila. In
Handbook of Genetics, R.C. King, ed. (Plenum Press: Plenum Press), pp. 421–469.
Tsacas, L. (2002). Le nouveau complexe africain Drosophila loiciana et l'espèce apparentée
D. matileana n. sp. (Diptera : Drosophilidae). Ann Soc entomol Fr *38*, 57-70.
Tsacas, L., and Chassagnard, M.-T. (2000). Drosophila loiciana, nouvelle espèce africaine et
redescription de son epèce affine D. pruinosa Dudas (Diptera, Drosophilidae). Revue
française d'entomologie *22*, 213-222.
Tsacas, L., and Chassagnard, M.T. (1990). The Species of the Genus Zaprionus with the Fore
Femora Bearing Spines (Diptera, Drosophilidae). Annales De La Societe Entomologique De
France *26*, 461-487.
Wake, D.B. (1991). Homoplasy - the Result of Natural-Selection, or Evidence of Design
Limitations. American Naturalist *138*, 543-567.
Wake, D.B., Wake, M.H., and Specht, C.D. (2011). Homoplasy: from detecting pattern to
determining process and mechanism of evolution. Science *331*, 1032-1035.
Watts, J.C., Flynn, A., Tenhumberg, B., and Hebets, E.A. (2019). Contemporary sexual
selection does not explain variation in male display traits among populations. Evolution.
Wheeler, M.R. (1981). The Drosophilidae. In The Genetics and Biology of Drosophila, M.
Ashburner, H.L. Carson, and J.N. Thompson, eds. (Academic Press), pp. 1-97.
Wittkopp, P.J., and Beldade, P. (2009). Development and evolution of insect pigmentation:
genetic mechanisms and the potential consequences of pleiotropy. Semin Cell Dev Biol *20*,
65-71.
Wittkopp, P.J., Carroll, S.B., and Kopp, A. (2003a). Evolution in black and white: genetic
control of pigment patterns in Drosophila. Trends Genet *19*, 495-504.
Wittkopp, P.J., Williams, B.L., Selegue, J.E., and Carroll, S.B. (2003b). Drosophila
pigmentation evolution: divergent genotypes underlying convergent phenotypes. Proc Natl
Acad Sci U S A *100*, 1808-1813.
Wray, G.A. (2002). Do convergent developmental mechanisms underlie convergent
phenotypes? Brain Behav Evolut *59*, 327-336.
Wynn, S., and Toda, M.J. (1988). Drosophilidae (Diptera) in Burma. IV. The genus Zaprionus.
Kontyu *56*, 843-851.
Xie, K.T., Wang, G., Thompson, A.C., Wucherpfennig, J.I., Reimchen, T.E., MacColl, A.D.C.,
Schluter, D., Bell, M.A., Vasquez, K.M., and Kingsley, D.M. (2019). DNA fragility in the parallel
evolution of pelvic reduction in stickleback fish. Science *363*, 81-84.
Yassin, A. (2007). A revision of the tumiditarsus group of the subgenus Drosophila and its
relation to the genus Zaprionus. Drosophila Information Service *90*, 20-22.
Yassin, A. (2013). Phylogenetic classification of the Drosophilidae Rondani (Diptera): the
role of morphology in the postgenomic era. Systematic Entomology *38*, 349-364.
Yassin, A., Araripe, L.O., Capy, P., Da Lage, J.L., Klaczko, L.B., Maisonhaute, C., Ogereau, D.,
and David, J.R. (2008). Grafting the molecular phylogenetic tree with morphological

branches to reconstruct the evolutionary history of the genus Zaprionus (Diptera:
Drosophilidae). Mol Phylogenet Evol *47*, 903-915.
Yassin, A., Da Lage, J.L., David, J.R., Kondo, M., Madi-Ravazzi, L., Prigent, S.R., and Toda, M.J.
(2010). Polyphyly of the Zaprionus genus group (Diptera: Drosophilidae). Mol Phylogenet
Evol *55*, 335-339.
Yassin, A., and David, J.R. (2010). Revision of the Afrotropical species of Zaprionus (Diptera,
Drosophilidae), with descriptions of two new species and notes on internal reproductive
structures and immature stages. ZooKeys *51*, 33-72.
Yassin, A., Delaney, E.K., Reddiex, A.J., Seher, T.D., Bastide, H., Appleton, N.C., Lack, J.B., David,
J.R., Chenoweth, S.F., Pool, J.E.*, et al.* (2016). The pdm3 Locus Is a Hotspot for Recurrent
Evolution of Female-Limited Color Dimorphism in Drosophila. Curr Biol *26*, 2412-2422.
Yokoyama, S. (2002). Molecular evolution of color vision in vertebrates. Gene *300*, 69-78.
Zhang, W.-X., and Toda, M.J. (1988). The Drosophila immigrans species-group of the
subgenus Drosophila (Diptera: Drosophilidae) in Yunnan, China. Zool Sci *5*, 1095-1103.
Zhang, W.-X., and Toda, M.J. (1992). A new species-subgroup of the Drosophila immigrans
species-group (Diptera, Drosophilidae), with descriptions of two new species from China
and revision of taxonomic terminology. Jpn J Genet *60*, 839-850.
Zhang, W.-X., Toda, M.J., and Watabe, H. (1995). Fourteen new species of the Drosophila
(Drosophila) immigrans species group from the Oriental region. Jpn J Genet *63*, 25-51.
Zwaan, B.J., Azevedo, R.B., James, A.C., Van 't Land, J., and Partridge, L. (2000). Cellular basis
of wing size variation in Drosophila melanogaster: a comparison of latitudinal clines on two
continents. Heredity *84 (Pt 3)*, 338-347.
793

        Figure Legends

Figure 1. Sex brush morphology in distantly related species. (A, C, E, G, I, K)

Brightfield images of the first and second tarsal segments (ta1 and ta2) of the prothoracic

leg in males of six species. (B, D, F, H, J, L) SEM images of the ta1 sex brush. A-B) *D.*

pruinosa. A) The brush occupies the distal ~80% of ta1, replacing most of the transverse

bristle rows (TBRs, shown by brackets in this and other panels), which in females covers

the entire anterior-ventral surface. The ta1 segment is slightly widened at the distal end.

B) The tips of brush hairs are flattened, pointed, and form hooks that curve toward the base

of the leg (see inset). C-D) *D. immigrans.* C) The brush covers the distal ~60% of ta1 and

most of ta2. The shape of the segments is not modified. D) Similar to *D. pruinosa*, the brush

hairs are flattened, pointed, and form proximally curving hooks at the tips (inset). E-F) *D.*

repletoides. E) The brush covers ~70% of ta1 and most of ta2. Both segments are

shortened and have a bulbous shape. F) The tips of brush hairs are flattened but thick and

blunt. They form hooks that curve away from the leg base (inset). G-H) *Z. tuberculatus.* G)

The brush covers the distal ~60% of ta1. H) The tips of brush hairs curve distally and have

a pointed paddle-like shape with a slight depression (inset). I-J) *Z. vittiger.* K-L) *Z.*

bogoriensis. These species have distally curving tips of brush hairs, similar to *Z.*

tuberculatus.

Figure 2. Sex brushes evolved independently at least three times. Species and clades

with male sex brushes are highlighted in blue. A) Bayesian phylogeny of the species

represented in this study, based on the combined dataset. Numbers at each node indicate

the posterior probabilities of the respective taxon bipartitions. B) Bayesian phylogeny

based on the same dataset but excluding *D. quadrilineata*. Posterior probabilities of nodes

A-E in different analyses can be found in Supplement Table 3.

Figure 3. Sex brush development shows strong similarities across species.

Developing brush hair cells were visualized by immunostaining for membrane markers E-

cadherin (DE-cad) or Armadillo (Arm). For each species, the upper panels show the

confocal projections of ta1 segment (A-D, G-H) or both ta1 and ta2 (E-F). The bottom

panels (A'-H') show close-up views. For each species, early developmental stages are
shown on the left and later stages on the right. A-B') *D. pruinosa*, 27 and 42 hrs after
pupariation (AP). C-D') *D. immigrans,* 24 and 43 hrs AP. E-F') *D. repletoides,* 28 and 43 hrs
AP. G-H') *Z. tuberculatus,* 34 and 48 hrs AP. Hair progenitor cells can be distinguished from
epithelial cells by a ring of bright staining with a dense punctum in the middle. In all four
species, each hair cell is surrounded by 4-6 epithelial cells (blue dots), so that neighboring
hair cells are separated from each other by 1-2 cells, at both the early and the late stages.
The proximal TBRs (square brackets) form by expelling the intervening epithelial cells,
with the bristle progenitor cells migrating closer together to make straight rows (Tanaka et
al 2009) (e.g., compare A vs B, and G vs H). In contrast, no cell rearrangement is observed
in the brush.
836
837 **Supplement Figure 1.**   Strict consensus of 11 trees with the cumulative posterior
838 probability of 95% for the complete taxon sample, labeled as in Figure 1. Numbers at each
839 node indicate the posterior probabilities of the respective taxon bipartitions.
840
841 **Supplement Figure 2.**   Strict consensus of 11 trees with the cumulative posterior
842 probability of 99% for the taxon sample without *D. quadrilineata*.
843
844 **Supplement Figure 3.** Bayesian phylogenetic tree for the complete taxon sample, with the
845 dataset partitioned by gene and codon.
846
847 **Supplement Figure 4.** Phylogenetic tree for the taxon sample without *D. quadrilineata*,
848 with the dataset partitioned by gene and codon.

D. pruinosa

A early TBR

B late ta1 DE-cad

A' **B'**

D. immigrans

C early

D late ta1 Arm

C' **D'**

D. repletoides

E early

F late ta1 ta2 Arm

E' **F'**

Z. tuberculatus

G early

H late TBR ta1 DE-cad

G' **H'**