The involvement of neuroimmune cells in adipose innervation

Magdalena Blaszkiewicz[1,2], Elizabeth Wood[1], Sigi Koizar[1], Jake Willows[1], Ryan Anderson[1], Yu-Hua Tseng[3], James Godwin[4,5], Kristy L. Townsend*[1,2,6]

[1]School of Biology and Ecology, University of Maine, Orono, ME
[2]Graduate School of Biomedical Science and Engineering, University of Maine, Orono, ME
[3]Joslin Diabetes Center, Harvard Medical School, Boston MA
[4]Jackson Laboratory, Bar Harbor ME
[5]MDI Biological laboratory, Bar Harbor ME
[6]The Ohio State University, Columbus OH

*Corresponding Author

1014 Biomedical Research Tower
460 W 12[th] Ave
Columbus, OH 43210
kristy.townsend@osumc.edu

Running Title: Myeloid-secreted BDNF maintains adipose innervation
Key Words: Innervation, white adipose tissue (WAT), energy expenditure, browning, CINCs, cold-induced neuroimmune cells, BDNF, monocyte/macrophage

Abstract

Background: Innervation of adipose tissue is essential for the proper function of this critical metabolic organ. Numerous surgical and chemical denervation studies have demonstrated how maintenance of brain-adipose communication through both sympathetic efferent and sensory afferent nerves helps regulate adipocyte size, cell number, lipolysis, and 'browning' of white adipose tissue. Neurotrophic factors are growth factors that promote neuron survival, regeneration and plasticity, including neurite outgrowth and synapse formation. Peripheral blood immune cells have been shown to be a source of neurotrophic factors in humans and mice. Although a number of immune cells reside in the adipose stromal vascular fraction (SVF), it has remained unclear what roles they play in adipose innervation. We previously demonstrated that adipose immune cells secrete brain derived neurotrophic factor (BDNF). Methods: We now show that deletion of this neurotrophic factor from the myeloid lineage led to a 'genetic denervation' of inguinal subcutaneous white adipose tissue (scWAT), thereby causing decreased energy expenditure, increased adipose mass, and a blunted UCP1 response to cold stimulation. Results: We and others have previously shown that noradrenergic stimulation via cold exposure increases adipose innervation in the inguinal depot. Here we have identified a subset of myeloid cells that home to scWAT upon cold exposure and are Ly6C$^+$ CCR2$^+$ Cx3CR1$^+$ monocytes/macrophages that express noradrenergic receptors and BDNF. Conclusions: We propose that these cold induced neuroimmune cells (CINCs) are key players in maintaining adipose innervation as well as promoting adipose nerve remodeling under noradrenergic stimulation, such as cold exposure.

Introduction

Introduction

In order for the central nervous system (CNS) to regulate functions of distal

organs and tissues, peripheral nervous system (PNS) innervation needs to be

maintained and properly coordinated. It has been demonstrated repeatedly that loss of

innervation of the adipose organ (by surgical or chemical means) led to dysfunction of

the tissue and disruption to whole body energy homeostasis. Denervation of brown

adipose tissue (BAT) greatly impairs the energy expending process of adaptive

thermogenesis (1-4), while denervation of white adipose tissue (WAT) results in fat

mass accumulation via hyperplasia and impaired lipolysis (5-7). Furthermore, we have

demonstrated that nerves in subcutaneous white adipose tissue (scWAT) undergo

remodeling in response to environmental stimuli (cold, exercise), and exhibit signs of

neuropathy in obesity/diabetes as well as with aging (8).

The maintenance of proper neural innervation is facilitated by neurotrophic

factors (NFs) both in the CNS (9) and PNS (10). NFs are nerve growth factors that

support nerve health, survival, and plasticity. Brain derived neurotrophic factor (BDNF)

is one member of the neurotrophin family of NFs, which in mammals also includes

nerve growth factor (NGF), Neurotrophin-3 (NT-3), and Neurotrophin-4/5 (NT-4/5).

Neurotrophins signal predominantly through Trk receptors on nerves, through which

they are endocytosed and, in peripheral nerves, transported in a retrograde manner to

the nerve cell body, located in the ganglia. BDNF has been well studied for its role in

hippocampal synaptic plasticity in the adult brain (11-13), as well as learning and

exercise-related neurogenesis (14). BDNF is also an important modulator of energy

balance through its actions in the central nervous system (15-17). Deletion of *Bdnf* in

the ventromedial and dorsomedial regions of the hypothalamus resulted in an obesity

phenotype due to hyperphagic behavior (16). Obesity is associated with lower serum

levels of BDNF in humans (18, 19) while dietary restriction normalizes BDNF deficits in

the brain in a mouse model of Huntington's disease (20). Animal studies have shown

that central and peripheral administration of BDNF reduced food intake and

hyperglycemia and increased energy expenditure, via CNS mediated mechanisms (21-

23). As review by Xu and Xie, genetic mutations in human BDNF and its receptor TrkB

result in morbid early-onset obesity (24); furthermore, genome wide associated studies

(GWAS) have identified single nucleotide polymorphisms (SNPs) in or near *BDNF* to be

associated with increased body mass index (BMI) (24).

Despite these strong correlations between altered function of BDNF and its

receptor TrkB with obesity, this growth factor has been predominately studied only in

the CNS. A few studies have clearly shown that NFs, including BDNF, are present in

adipose tissue (25-27), but the cellular source of BDNF in adipose had not been

determined. In adipose specific (*FABP4-Cre*) knock-outs of BDNF or TrkB, only CNS

effects were apparent (not surprising given off-target brain effects of this cre line) (28).

Using instead an *adipoq*-cre (mature adipocyte-specific line) there was no obesity in a

BDNF or a TrkB knock-out (KO), and the BDNF KO showed no difference in adipose

BDNF levels (28). Together, these findings suggested that mature adipocytes are *not*

the cellular source of BDNF (28), which we have confirmed in this study by

demonstrating predominance of BDNF expression in the SVF. NFs are secreted by glial

cells in the brain (29) and Schwann cells in peripheral tissues (30). However, other cell

types, predominantly immune cells, are also known sources of NFs (31-35) but are less

studied.

As part of the immune response to injury, immune cells are critical players in

wound healing, regeneration, and remodeling of various tissues. They are an important

component of the adipose organ where they modulate the inflammatory response, clear

the tissue of apoptotic cells, and mediate adipose tissue remodeling during obesity

through an influx of monocytes (undifferentiated macrophages), neutrophils, T cells, B

cells and mast cells (36-41). Macrophages and their monocyte precursors are myeloid

lineage immune cells and comprise the highest fraction of immune cells present in

adipose tissue (42). They are highly heterogeneous cells that are polarized by

environmental stimuli to evoke differential responses within a tissue, including secretion

of cytokines.

In a simplistic paradigm, classically activated macrophages (M1) act in a pro-

inflammatory manner, while alternatively activated macrophages (M2) produce an anti-

inflammatory response and initiate tissue remodeling after injury. Both M1 and M2 cells

retain phagocytic behavior. During obesity, M1 adipose tissue macrophages (ATMs)

greatly increase in number without an influx of resolving M2 macrophages, thus

contributing to a chronic state of tissue inflammation (43). Inflamed, insulin-resistant

adipose tissue histology is characterized by macrophage crown-like structures

surrounding hypertrophic, hypoxic and dying adipocytes. On the other hand, it has been

suggested that cold-induced browning of adipose promotes an M2 phenotype in ATMs

(43), possibly promoting tissue remodeling and potentially serving as a source of NFs in

adipose tissue. Importantly, the immunology field has uncovered that M1 and M2

designations are an oversimplification, and in reality many immune cells express

markers of both subtypes and may be inter-converting between the two polarities, but

this has not yet been fully explored in adipose tissues.

Although myeloid cells, including monocytes and macrophages, from peripheral

blood have been shown to store and release NFs (44), it is still unclear what role these

immune cells play in peripheral innervation, and it remains unknown how and if adipose-

resident and infiltrating immune cells are stimulated to release NFs that act locally.

Microglia, the CNS resident myeloid cells that are most similar to macrophages, are an

accepted source of BDNF in the brain (45) and increase secretion of BDNF in response

to neuroinflammation (46). Microglial-derived BDNF in the CNS promotes hippocampal

synaptic plasticity (45) and neurogenesis (47). We hypothesized that myeloid lineage

cells may play a similar role in adipose tissue, and generated a myeloid specific BDNF

knock-out mouse model by crossing *LysMCre$^{+/-}$* and *BDNF$^{fl/fl}$* mice. Here we report a

role for myeloid derived BDNF in the specific maintenance of scWAT innervation, and

identify a subpopulation of monocyte/macrophages (Ly6C$^+$CCR2$^+$Cx3CR1$^+$) that

infiltrate adipose tissue in response to cold stimulation and express BDNF.

Materials & Methods

Mice, Metabolic Phenotyping, and in vivo Analyses

Animals

The following mouse strains were obtained from The Jackson Laboratory: C57BL/6J

(Stock # 000664); LysMCre+/-(B6.129P2-Lyz2/J, Stock # 004781); BDNF$^{fl/fl}$

(Bdnf^{tm3Jae}/J, Stock # 004339); R26R-EYFP (B6.129X1-

Gt(ROSA)26Sortm1(EYFP)Cos/J, Stock # 006148); Cx3CR1-EGFP (B6.129P2(Cg)-

Cx3cr1tm1Litt/J, Stock # 005582). Animals with myeloid specific deletion of BDNF

(*LysMCre$^{+/-}$ ::BDNF$^{-/-}$)* were generated in our facility by crossing mice heterozygous for

the myeloid -specific cre transgene (LysMCre$^{+/-}$) with mice homozygous for floxed

BDNF. Reporter mice were generated by crossing LysMCre$^{+/-}$ x BDNF^{flfl} x R26R-EYFP.

Animals were housed 3-5 to a cage providing for socialization, in a monitored

temperature and humidity-controlled environment with 12/12hr light/dark cycle. Cages

were replaced weekly, Ad libitum access to food and water was maintained. For all

studies animals were sacrificed using CO_2 followed by cervical dislocation.

174   *Dietary Fat Interventions and Food Intake*

Adult male *LysMCre$^{-/-}$::BDNF$^{fl/fl}$* (CON) *and LysMCre$^{+/-}$ ::BDNF$^{-/-}$* (KO) mice were fed a

45% HFD diet from Research Diets (New Brunswick, NJ) for up to 11 weeks. Mice were

housed 2-3 per cage, at room temperature. Body weight was measured weekly. Food

intake was measured daily for 7 days, then weekly until the end of the experiment.

Adiposity was assessed at the end of the study by weighing intact adipose depots after

surgical removal.

181   *CLAMS*

Metabolic cage analyses were conducted in a Comprehensive Laboratory Animal

Monitoring System (CLAMS; Columbus Instruments, Columbus, OH), for measurement

of oxygen consumption (VO$_2$) and carbon dioxide production (VCO$_2$), from which both

respiratory exchange ratio (RER), and energy expenditure (Heat) were calculated:

RER=VCO$_2$/VO$_2$; *Energy expenditure (heat) = CV * VO$_2$ cal/hr, where CV is the "caloric*

*value" as given by CV = (3.815 + 1.232) * RER.* Animals were single housed in a

bedding free cage, at room temperature on a 12hr light/dark cycle.   Mice were

acclimated for 24-48hrs, after which VO$_2$, VCO$_2$, RER, and Heat were measured every

15min for 3 days (72hrs). Waveform analysis of CLAMS data was performed by

matching every 15min measurement across all three 24hr-cycles. Two-way repeated

measures analysis of variance (RM, ANOVA) was performed for average VO$_2$, VCO$_2$,

RER, and Heat per group. An uncorrected Fisher's Least Significance Difference test

was performed for each time point between dietary groups as a post-hoc test.

Interaction P values are reported, which represent differences in 24hr data between

groups, as well as multiple comparison results for differences which were only day/night

phase specific.

CL316,243 Injections

Adult (12-13 week old) male C57BL/6 mice received daily intraperitoneal (i.p.) injections

of ADRβ3 agonist CL316,243 (Tocris Bioscience, Bristol, U.K.; Cat # 1499), at

1.0 mg/kg BW or an equivalent amount of sterile saline, for 10-14 days.

Cold Exposure Experiments

All cold exposure was carried out in a diurnal incubator (Caron, Marietta, OH, USA) at

5°C, and a 12hr light/dark cycle. Animals were housed two to a cage and continuously

cold exposed for 3 - 14 days.

Glucose Tolerance Test (GTT)

Animals were fasted overnight for 16hr, after which they received an i.p. bolus injection

of 1g/kg glucose. Blood glucose was measured using tail vein blood with a hand-held

glucometer (OneTouch UltraMini, LifeScan, Milpitas, CA, Johnson & Johnson, New

Brunswick, NJ), at time 0 and at intervals of 15 minutes, 30 minutes, and 60 minutes

after injection.

Western Blots

Protein lysates were prepared by homogenizing frozen whole adipose depots in RIPA

buffer using a Bullet Blender (Next Advance, Averill Park, NY), followed by Bradford

Assay, and preparation of equal-concentration lysates in Laemmli buffer. 60ug of

protein was loaded per lane of a 10% polyacrylamide gel, and following gel running, proteins were transferred to PVDF membranes for antibody incubation. Primary antibodies used included: anti-PGP9.5 (Abcam, Cambridge, U.K. Cat. #10404 RRID:AB_297145 and #108986 RRID:AB_10891773) used at a 1:1000 and 1:500 dilutions respectively; anti-UCP1 (Abcam, Cambridge, U.K. Cat. #10983; RRID:AB_2241462) used at 1:1000 dilution, anti-TH (Millipore Cat. # AB152; RRID:AB_390204; Merck Millipore, Burlington, MA), anti-β-tubulin (Cell Signaling Technology, Danvers, MA, USA; Cat. # 2146; RRID:AB_2210545), were all used at 1:1000 dilution. Secondary antibody was anti-rabbit HRP (Cell Signaling Ct # 7074; RRID:AB_2099233), used at a 1:3000 dilution. Blots were visualized with enhanced chemiluminescence (ECL; Pierce) on a Syngene G:BOX. Protein expression of PGP9.5, TH, and UCP1 was normalized to either β-tubulin or β-actin and quantified in ImageJ.

Collection of Adipose Secretions and BDNF ELISA

BAT depots were dissected, weighed and minced in a petri dish containing DMEM (high-glucose, serum-free). Minced tissue was transferred to a 15mL conical tube with 5mL DMEM (loosely capped to keep tissue oxygenated) and placed in a shaking water bath at 37°C. Secretions were collected at time 0, 1hr, 2hrs, and 3hrs (1mL collected from conical tube at each time point and replaced with 1mL fresh DMEM). Secretions were stored at -80°C until processing. For ELISA, protein secretions were concentrated using Amicon ® Ultra Centrifugal Filters, Ultracel ® -100K (Millipore, Burlington, MA USA; Cat. # UFC510096), per manufacturer's instructions. Mouse BDNF PicoKine™ ELISA Kit (Boster Biological Technology, Pleasanton, CA, USA; Cat#

EK0309) was used per manufacturer's instruction to determine amount of BDNF
present in adipose active secretions.

Thyroid Hormone ELISA

Mouse sera were used to measure circulating levels of thyroxine (T4) and
triiodothyronine (T3). Circulating concentrations were determined by Enzyme-Linked
Immunosorbant Assays (ELISA) at Maine Medical Center Research Institute's Core
Facilities (Scarborough, ME).

SVF Isolation

Bilateral whole inguinal adipose depots were quickly dissected and weighed. Tissue
was minced in 37°C pre-warmed DMEM (high glucose, serum free) containing 2mg/mL
Roche Collagenase A (Millipore-Sigma, St. Louis, MO; Cat# 10103586001) at a volume
of 10mL/depot. Minced tissue with collagenase containing DMEM was placed in a 50mL
conical tube and transferred to a shaking water bath (350/min rotation) at 37°C. Every
10min cells were dispersed by vortex and pipette mixing. Full dissociation was usually
achieved within 2hrs, when adipocytes were clearly visible and all tissue was
dissociated. Dissociated media was poured through 100um cell strainers, rinsed with
DMEM and centrifuged at 500g for 10min to separate adipocytes from SVF. After
centrifugation adipocytes were collected (found floating on top); remainder of DMEM
was removed sparing the SVF pellet. SVF pellet was incubated with 500uL of RBC lysis
buffer on ice for 2min, after which 2mL of DMEM with serum was added to stop lysis.
Cells were centrifuged at 500g for 5min at 4°C, and either collected for RNA or
resuspended in FACS/MACS buffer for cell sorting.

Magnetic-activated Cell Sorting (MACS)

SVF from bilateral whole inguinal adipose depots was isolated as described above and resuspended in degassed buffer (1XPBS pH7.2, 0.5% BSA and 2mM EDTA). Single-cell suspensions were sorted on the MidiMACS Quadro magnetic-activated cell separator system (Miltenyi Biotec, Bergisch Gladbach, Germany) according to manufactures instructions. Briefly, cells were stained with primary PE-conjugated antibody, CD11b-PE (Cat #130-098-087); a 1:10 antibody dilution per 10^7 cells was used. Cells were incubated for 10min at 4-8°C, washed, and centrifuged at 300g for 10min. Washed cell pellet was resuspended in a 1:10 dilution of anti-PE microbeads included in Anti-PE MultiSort Kit (Cat. #130-090-757). Following 15min incubation at 4-8°C, cells were washed and centrifuged as described above. Cells were resuspended in 500μL of buffer and passed through LS columns of the MACS separator. LS columns were prepped according to manufacturer's suggestion. Cells were washed 3 times, collected cells (CD11b-) were collected. LS columns were removed from the MACS separator and flushed with 5mL of buffer to release the magnetically labeled cell fraction (CD11b+). MicroBeads were removed using MicroSort release reagent included in Anti-PE MultiSort Kit (Cat. #130-090-757). MicroBead free CD11b+ cell fraction was labeled for the secondary marker, Anti-F4/80-APC (Cat. #130-102-942), following the same procedure as described for the primary marker, except that Anti-APC MicroBeads (Cat. #130-090-855) were used.

Fluorescence-activated Cell Sorting (FACS)

SVF from bilateral whole inguinal adipose depots was isolated as described above. For

cell sorting, the following 5 marker panel was used with DAPI exclusion for viability:

Anti-Mouse Ly6C_BV570 (HK1.4), Anti-Mouse CD11b PE (M1/70), Anti-Mouse

CX3CR1 PercP5.5, SA011F11), Anti-Mouse CD45-PE-Cy7 (30-F11), Anti-Mouse CCR2

A700 (475301). Sorting was performed on a BD™ FACSAria II™ cell sorter with SVF

gated on CD45 and CD11b; CD45+CD11b- represented the non-myeloid population;

CD45$^+$CD11b$^+$ myeloid fraction was gated on Ly6C, followed by CCR2 and Cx3CR1.

20 Cell Surface Marker tSNE Analysis of Myeloid Diversity in Male and Female Mice

Adult (13-15 week old) male and female control mice (BDNF$^{fl/fl}$) were cold exposed for

10 days in a diurnal incubator as described above. Following cold exposed SVF from

bilateral whole inguinal adipose depots was isolated as described above. Single cell

suspensions were treated with unlabeled FC receptor blocking antibody cocktail

(CD16/CD32) in 50ul of FACS buffer (HBSS + 5mM EDTA + 2% FCS). Cells were then

incubated in an "staining antibody cocktail" against 20 cell surface markers for 60min at

4°C in 100ul. Cells were then washed in 1mL of FACS buffer and centrifuged at 400g

for 7min, 2 times. Cells were then resuspended and analyzed on a five-laser 30-

parameter FACSymphony A5 cytometer (BD Biocsiences, San Jose, USA) using DAPI

exclusion for cell viability. For compensation of fluorescence spectral overlap,

UltraComp eBeads (eBioscience, Inc.) were used following the manufacturer's

protocols. FCS 3.0 files generated by flow cytometry were initially processed using

FlowJo Software (Tree Star, Ashland, USA) for automated compensation. Standard

manual hierarchical gating was performed to remove debris, cell doublets and (DAPI+)

dead cells from analysis before gating on CD45+ leukocyte populations. In preparation

for performing the 20 marker tSNE leukocyte population analysis, the CD45+ population

from each sample was down-sampled to 3000 events to normalize cellular input

between samples. Using FlowJo, a concatemer of all samples was performed. An

unbiased T-Distributed Stochastic Neighbor Embedding (tSNE) plugin algorithm was

then run using defaults with 22 parameters, on the whole sample pool to obtain a multi-

sample population reference map. Gating of each sample and experimental group was

performed to generate tSNE maps for each condition. Differential cell clusters were

gated and 20 marker histograms plots were used to predict cluster identities.

Antibodies used for adipose flow cytometric panel shown here: Anti-Mouse CD45-

BUV395 (30-F11), Anti-Mouse B220_BUV496 (RA3-6B2), Anti-Mouse Ly-6G BUV563

(Clone 1A8), Anti-Mouse CD19 BUV661 (Clone 1D3), Anti-Mouse CD11b BUV737

(M1/70), Anti-Mouse NKp46_BV421 (29A1.4), Anti-Mouse CD62L BV510 (MEL-14),

Anti-Mouse Ly6C_BV570 (HK1.4), Anti-Mouse CD3 BV605 (17A2), Anti-Mouse

Mrc1_BV650 (C068C2), Anti-Mouse MHCII -I-A/I-E -BV711 (M5/114.15.2), Anti-Mouse

NK1.1 BV785 (PK136), Anti-Mouse CD11c_FITC (HL3), Anti-Mouse CD80 PE (16-

10A1), Anti-Mouse CD115 CF594 (AFS98), Anti-Mouse CX3CR1 PercP5.5

(SA011F11), Anti-Mouse CD64 PE-Cy7 (X54-5/7.1), Anti-Mouse CD14 APC (Sa2-8),

Anti-Mouse CCR2 A700 (475301), Anti-Mouse F480 APC/Cy7 (BM8).

Gene Expression (qPCR)

RNA was isolated from whole tissue depots using Trizol reagent, and total RNA

extracted using a Zymo (Irvine, CA) kit. RNA yield was determined on a Nanodrop;

cDNA was synthesized using a High Capacity Synthesis Kit (Applied Biosystems,

Foster City, CA). Real-time quantitative (q)PCR was performed with SYBR Green (Bio-

Rad, Hercules, CA) on a CFX96 instrument (Bio-Rad, Hercules, CA).

Histology

Adipose

Immunofluorescent staining of 10% buffered formalin fixed, paraffin-embedded, 7um

sections of adipose tissues was performed for detection of UCP1 (RRID:AB_2241462,

1:500, Abcam, Cambridge, UK. Cat. #10983). Alexa 488 (2.5μg.mL, Molecular Probes,

Eugene, OR, USA, Cat. # A11070) was used as secondary antibody. Typogen Black

staining was used to quench tissue autofluorescence (prior to antibody incubation) and

also provided visualization of cell size, browning (multilocularity) and crown-like

structures. Stained sections were mounted using Millipore mounting fluid (Burlington,

MA USA; Cat. # 5013) and 1 1/2 coverslips, and imaged on a Nikon Eclipse E400

epifluorescent microscope equipped with Nikon DS-fi2 camera. Wholemount adipose

staining was performed as previously described (8, 48) and imaged by widefield and

confocal microscopy. Widefield imaging was performed on a Nikon E400

epifluorescence microscope with Hamamatsu ORCA-Flash4.0 V2 Digital CMOS

monochrome camera (Hamamatsu Photonics K.K., C11440-22CU). Objectives used:

Nikon CFI Plan Apochromat Lambda 10X (NA 0.45, WD 4.00mm, dry) and Nikon CFI

Plan Fluor 40X (NA 0.75, WD 0.66mm, dry.) Images were processed using Nikon NIS-

Elements software. Images captured on Hamamatsu ORCA-Flash4.0 V2 were pseudo

colored. For confocal imaging a Leica TCS SP8 DLS microscope with HyD detectors

was used. Fluorophores were excited with a white light laser tuned specifically for the

excitation and emission spectra of EGFP. Objectives used: HC PL APO CS2 20X (NA

0.75, WD 0.62mm, dry) and HC PL APO CS2 63X (NA 1.40, WD 0.14mm, oil). Images

were processed with LASX software and were either pseudo colored or left

monochromatic. Primary antibodies included: PGP9.5 conjugated to Alexa 647 (1:200,

Abcam, Cambridge, U.K. Cat. #196173).

Neuromuscular Junction Immunofluorescence, Imaging, and Analysis

Soleus and medial gastrocnemius muscles were removed and fixed in a 2% PFA at 4°C

for 2 hours. Tissues were rinsed with 1XPBS and incubated in blocking buffer

(1XPBS/2.5%BSA/0.5-1%Triton) at 4°C for at least 24 hours and up to 7 days.

Following blocking muscles were teased, tendons and fat was removed, and tissue was

flattened by being placed between two tightly-bound glass slides for at least 30 minutes

at 4°C. Tissues were transferred to fresh blocking buffer at 4°C for at least 12 hours.

Immunostaining of innervation with primary antibodies was performed overnight at 4°C,

followed by 1XPBS washes on a rotating platform at 4°C replacing PBS every 1hr for a

total of 4-6hrs. Tissues were incubated with secondary fluorescent antibodies in similar

fashion as primary antibodies. Primary antibodies included: neurofilament-M (2H3,

RRID:AB_531793, 1:500) and synaptic vesicles (SV2, RRID:AB_2315387, 1:250) from

Developmental Studies Hybridoma Bank, (University of Iowa, USA). Secondary

antibodies included: Alexa Fluor 488 at 1:500 (A21121) and alpha-bungarotoxin (BTX)-

conjugated to Alexa Fluor 594 at 1:1000 (B13423) from Molecular Probes (Eugene, OR,

USA). Tissues were mounted on microscope slides using Millipore mounting fluid

(Burlington, MA USA; Cat. # 5013) and 1 1/2 coverslips then sealed and allowed to set

overnight. Stained sections were imaged on a Nikon Eclipse E400 epifluorescent

microscope equipped with Nikon DS-fi2 camera. Brightness, contrast, and sharpness

were adjusted in Microsoft PowerPoint. Up to 100 NMJs were counted for each tissue,

statistics were conducted in GraphPad PRISM software (La Jolla, CA, USA) using the

multiple t-tests (one-per row) function.

Statistical Analysis

For all animal experiments, mice were randomized to treatment groups to ensure no

difference in starting body weight. All plots represent mean +/-SEM. Statistical

calculations were carried out in Excel or GraphPad Prism software (La Jolla, CA, USA),

utilizing ANOVA, Linear Regression, or Student's T-test as indications of significance

(specified in Figure legends). Gene and protein expression data were normalized to a

housekeeper and analyzed by either ANOVA or by Student's t-test, two-tailed, using

Welch's correction when variance was unequal. Error bars are SEMs. For all figures, $*p$

< 0.05, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

Ethical Statement

All procedures and handling of animals were performed in accordance with the

University of Maine's Institutional Animal Care and Use Committee (IACUC), to comply

with the guidelines of the PHS Policy on Humane Care and Use of Laboratory Animals,

and Guide for the Care and Use of Laboratory Animals. This study was approved by

the University of Maine's IACUC, under protocol A2017-09-04.

Results

BDNF is expressed primarily in adipose SVF

We previously demonstrated that BDNF secretion increases from scWAT in response to noradrenergic stimulation, after administration of the β-3 adrenergic receptor (ADRβ3) agonist CL316,243 (8). We then endeavored to determine which adipose compartment and cell type was the source of local BDNF secretion in scWAT. Adipose tissue is a heterogeneous organ and although adipocytes are the main cell type, numerous other cell types are contained within the stromal vascular fraction (SVF) of adipose tissue. Adipose SVF consists predominantly of hematopoietic lineage cells including adipose tissue macrophages (ATMs (41, 42)), but also contains numerous other immune cell types, preadipocytes, vascular endothelial cells, and pericytes. To determine the compartmental source of adipose BDNF, adult male C57BL/6 mice were cold exposed and whole SVF was isolated from mature adipocytes of inguinal scWAT by collagenase dispersion. Gene expression analysis revealed that *Bdnf* was almost exclusively expressed in the SVF, while other common growth factors (nerve growth factor; NGF, and vascular endothelial growth factor, VEGFa) showed no difference of expression between SVF and mature adipocyte fractions (Fig. 1A).

Given the known expression of BDNF in immune cells, we sought to determine the contribution of myeloid-lineage BDNF to adipose innervation by creating a knock-out (KO) mouse model using Cre-Lox technology. *LysMCre$^{+/-}$* mice were bred to *BDNF$^{fl/fl}$* mice to generate *LysMCre$^{+/-}$::BDNF$^{-/-}$* (KO) animals, which lacked BDNF in myeloid lineage cells (Fig. 1B). (For the genotyping strategy, see Suppl. Fig. S1A.) Compared to littermate controls, KO animals exhibited a significant, although not complete, decrease of *Bdnf* in adipose SVF as measured by gene expression (Fig. 1C). Likely additional

SVF cell types also contribute to tissue BDNF levels. Since myeloid lineage cells are

also expressed in the brain, and BDNF has been shown to play a role in energy balance

via CNS control of satiety, we also investigated whether our KO model affected BDNF

expression in the hypothalamus. Gene expression of *Bdnf* in the hypothalamus did not

differ between KO animals and their littermate controls (Fig. 1D). Physiological

assessment using metabolic cages performed on KO animals and their littermate

controls in the basal state revealed that KO mice had significantly lower energy

expenditure (Fig. 1E) despite showing no difference in body weight or adiposity (Suppl.

Fig. S1B).

KO mice have a blunted response to cold stimulation

Since a decrease in energy expenditure could be indicative of impaired sympathetic

drive, we stimulated sympathetic nerve activity via cold exposure. When adult (22-23

week old) male mice were cold challenged at 5°C for 4 days, KO and control animals

maintained similar body weight (Fig. 2A, left panel). However, KO animals had a trend

for increased adiposity (Fig. 2A, right panel). As cold exposure stimulates

catecholamine-induced lipolysis mediated through sympathetic nerves, we next

investigated innervation of the inguinal scWAT depot following a 7-day cold challenge.

Protein expression of the pan-neuronal marker PGP9.5 was markedly reduced in

inguinal scWAT of male KO mice compared to littermate controls (Fig. 2B, left panel).

Protein expression of tyrosine hydroxylase (TH), a marker of sympathetic innervation

and activation, was also drastically reduced in inguinal scWAT of KO animals compared

to controls (Fig. 2B, right panel). Gene expression of synaptic/innervation markers

(*Psd95, Sox10, Synapsin I, Synapsin II, Synaptopysisn)* in inguinal scWAT also showed

a coordinated trend to be decreased in KO animals (Suppl. Fig S1C), further indicating

perturbations in scWAT innervation of KO animals. Together, these data indicated a

'genetic denervation' in this model.

Interestingly, gene expression of lipolytic markers *Atgl* and *Hsl* showed an

increased trend in scWAT of KO animals (Suppl. Fig S1C). Loss of scWAT innervation

potentially results in decreased lipolysis due to lack of SNS mediated release of

norepinephrine. However, due to the critical role of lipolysis in facilitating

thermoregulation during cold challenge, we suspect that lipolysis in KO animals is

achieved by alternative, nerve-independent, pathways. For example, glucocorticoid-

mediate lipolysis has been attributed to increased β-adrenergic responsiveness via

direct signaling or indirectly by inducing secretion of angiopoietin-like 4 (Angptl4) (49,

50).

Cold exposure induces expression of uncoupling protein 1 (UCP1) as required

for adaptive thermogenesis, due to its ability to uncouple the mitochondrial respiratory

chain resulting in a proton leak and heat production. UCP1 is therefore a unique marker

of BAT and cold-induced 'browning' in scWAT. Gene expression of *Ucp1* in inguinal

scWAT of 7-day cold exposed mice was reduced in KO animals compared to littermate

controls (Fig. 2C). Thyroid hormone potentiates sympathetic nervous system (SNS)

activation of thermogenesis in BAT, but is produced via a distinct neuronal pathway

from adipose sympathetic drive. We observed no changes in circulating thyroid

hormones between KO and control (CON) animals (Fig. 2D). Histological assessment of

inguinal scWAT revealed what appeared like increased multilocularity suggestive of

browning in KO animals (Fig. 2E, Typogen Black staining). However, there was a

striking lack of UCP1 expression in these areas in KO tissues (Fig. 2E). Typogen Black

(used here to reduce tissue autofluorescence) may also mark immune cells, thus there

is potential for increased immune cell infiltration in the KO tissues as well. However,

perilipin (a lipid droplet associated protein) staining of scWAT from KO animals clearly

showed that the multilocular cells lacking UCP1 expression contained high amounts of

lipid droplets (Fig. 2F, UCP1 immunofluorescence in red, perilipin immunofluorescence

in green on serial sections). Preadipocytes maintain multilocular lipid droplets prior to

differentiation, and considering that denervation of WAT increases hyperplasia, the

observed multilocularity could more likely be areas of increased preadipocytes. In total,

these findings demonstrate reduced thermogenic potential in denervated scWAT.

Genetic denervation of scWAT in LysMCre$^{+/-}$::BDNF$^{-/-}$ (KO) mice is depot specific

We next sought to determine whether genetic denervation of scWAT in KO

animals was restricted to this adipose tissue or extended to other organs. BDNF is a

known myokine, and muscle is an energy expending tissue. We assessed innervation of

fast twitch (gastrocnemius) and slow twitch (soleus) muscle in CON and KO, by

investigating occupancy of neuromuscular junctions (NMJs) at basal conditions in adult

male mice. Immunostaining of the presynaptic nerve and vesicles (neurofilament and

SV2 markers, respectively) with postsynaptic acetylcholine receptors was performed to

allow visualization of NMJ (Suppl. Fig. S2A, left panel). Following counts of occupied,

partially occupied, and unoccupied NMJs it was determined that there was no evidence

of neurodegeneration in the NMJ of KO animals (Suppl. Fig. S2A, right panel). In the

same animals we also assessed axon numbers of spinal (L5 ventral root), motor, and

sensory nerves through cross-section imaging (Suppl. Fig. S2B). A lower axon count

could reflect neuronal death in these larger PNS nerves, but no difference was observed between CON and KO animals.

Myeloid cells are also present in BAT tissue. We therefore wanted to evaluate whether a lack of BDNF in BAT myeloid cells would have an effect on this tissue's function. BAT of 7-day cold exposed 12-25 week old male CON and KO mice was evaluated. Protein expression of neuronal markers TH and PGP9.5 did not differ between KO and control mice (Suppl. Fig. S3A), indicating that BDNF may not have an important neurotrophic role in BAT. Consistent with that, histological assessment of BAT revealed no difference in cellular morphology nor UCP1 expression (Suppl. Fig. S3B). Indeed, when adult male C57BL/6 mice were cold exposed or treated with the pharmacological ADRβ3 agonist, CL316,243, no difference in BDNF secretion from BAT was observed when compared to basal conditions (Suppl. Fig. S3C). Together, these data supported scWAT depot specificity of our genetic denervation model, and BAT neuronal function may simply be due to increased sympathetic nerve activity and not changes in neural plasticity and neurite outgrowth in the tissue. Alternatively, a different neurotrophic factor, like NGF, may be more important for BAT than WAT.

HFD feeding exacerbates fat mass accumulation in LysMCre$^{+/-}$::BDNF$^{-/-}$ (KO) Mice

Loss of sympathetic innervation to inguinal scWAT has been shown to increase depot mass (51), however, in our genetic model of scWAT denervation, no difference in adiposity was observed under basal conditions (Suppl. Fig. S1B) despite the observed decrease in energy expenditure (Fig. 1E). We next metabolically challenged CON and KO mice with a 45% high fat diet (HFD). Adult (25 week old) male CON and KO mice were placed on a HFD for 3-11 weeks, to assess adipose integrity and energy balance.

At 3 weeks of HFD feeding animals were characterized in metabolic cages. HFD

resulted in only a slight decrease in energy expenditure in KO mice compared to

littermate controls (Fig. 3A) as analyzed via waveform analysis over multiple 24hr

periods. However, KO mice showed a higher respiratory exchange ratio (RER) than

CON animals during the light cycle, indicative of preferential metabolism of

carbohydrates over lipids for fuel (Fig. 3B). These data fits with studies demonstrating

that adipose nerves are important for lipolysis (52) and that denervation would shift fuel

preference to carbohydrates. These physiological differences between CON and KO

animals were observed despite no difference in food intake or change in body weight

(Fig. 3C-D) at early time points. After 6 weeks of HFD, KO mice displayed altered

glucose control compared to CON animals (Fig. 3E), perhaps due to a shift in fuel

utilization. By 11 weeks of HFD feeding, KO animals displayed greater adiposity than

littermate controls (Fig. 3F), potentially due to the reduced innervation and loss of neural

control of certain metabolic processes.

Cold-induced neuroimmune cells (CINCs) are recruited to scWAT and express BDNF

After demonstrating the significance of myeloid derived BDNF to scWAT innervation,

we sought to determine which myeloid cells were the source of tissue BDNF. Given

their multifaceted role in adipose tissue, being a source of BDNF in the brain (microglia),

and the phenotype observed in a myeloid-lineage KO, we hypothesized that

monocytes/macrophages were the leading source of BDNF in scWAT. Since previous

studies indicated that BDNF is increased in scWAT with noradrenergic stimulation (8),

$CD11b^+$ $F4/80^+$ macrophages were isolated from SVF of inguinal scWAT of room

temperature and 5-day cold exposed C57BL/6 adult (12 week old) male mice. *Bdnf*

gene expression did not differ between room temperature and cold exposed CD11b$^+$

F4/80$^+$ macrophages (Fig. 4A). F4/80+ is considered a pan-macrophage marker, as

such we considered it too broad to reveal phenotypic changes in the spectrum of

macrophage populations in scWAT. Furthermore, F4/80+ is not the ideal marker of

monocytes (macrophage precursors), which could be infiltrating the tissue in response

to cold exposure. Based on these findings we applied a different approach to

determining which myeloid cells are the source of scWAT BDNF. Adult (12 week old)

female control animals were maintained at room temperature or cold exposed for 10

days. Inguinal scWAT SVF was isolated and flow cytometrically analyzed using a

custom antibody cocktail against immune cells. Surprisingly, cold exposure did not have

an effect on either M1 or M2 ATMs (Fig 4B). Instead, the greatest increase was in

Ly6C$^+$CCR2$^+$ monocytes.   Both Ly6C$^+$ and CCR2$^+$ are markers of inflammatory

monocyte migration (53). Following cold exposure, both Ly6C$^+$CCR2$^+$Cx3CR1$^-$ and

Ly6C$^+$CCR2$^+$Cx3CR1$^+$ populations increased in inguinal scWAT of female mice,

however, only the Ly6C$^+$CCR2$^+$Cx3CR1$^+$ population showed a strong trend for increase

in males (Fig. 4C). In a separate cohort of adult male and female C57BL/6 mice,

unbiased assessment using t-distributed Stochastic Neighbor Embedding (tSNE)

revealed subpopulations of immune cells changing in propensity in scWAT with cold,

and between male and female mice (cold and room temperature). Interestingly, the

Ly6C$^+$CCR2$^+$Cx3CR1$^+$ population, which we call cold-induced neuroimmune cells

(CINCs) again increased in both male and female scWAT with cold (Fig. 4D).

To confirm that Ly6C$^+$CCR2$^+$Cx3CR1$^+$ monocytes/macrophages were the source

of BDNF, we FACS sorted out Ly6C$^+$CCR2$^+$Cx3CR1$^+$ cells from inguinal scWAT SVF of

14 day cold exposed adult (12-13 week old) male C57BL/6 mice. We measured *Bdnf*

gene expression in cold induced Ly6C$^+$CCR2$^+$Cx3CR1$^+$ cells and found that they

showed a trend for increased expression of *Bdnf* compared to cold exposed non-

myeloid cells (Fig. 4E).   We believe the previously observed increase in BDNF in

adipose SVF ((8) and Fig. 1A) was due to infiltration of CINCs homing to scWAT upon

cold exposure, and not an increase in per-cell BDNF levels.

Adrβ3 gene expression in Ly6C$^+$CCR2$^+$Cx3CR1$^+$ cells confirmed the presence of

norepinephrine (NE) receptor on these cells, indicating the potential to be responsive to

SNS stimulation (Suppl. Fig. S4A), a likely mode for promoting BDNF release to the

tissue after cold exposure. Taken together, these data indicated that

Ly6C$^+$CCR2$^+$Cx3CR1$^+$ cells are cold-induced neuroimmune cells (CINCs) that increase

in number in scWAT after cold, have the potential to be stimulated by sympathetic

nerves, and express BDNF.

Adipose lymph node and adipose lymphatics may play a role in recruitment of CINCs

CINCs express the transient markers Ly6C$^+$ and CCR2$^+$, as well as Cx3CR1$^+$, indicating

that they are recruited to adipose tissue following cold stimulation as opposed to being

tissue resident immune cells. Immune cells are generally recruited to tissue through

vasculature upon chemoattract release from target tissue. To further investigate

dynamics of CINC recruitment to adipose tissue we utilized a Cx3CR1-EGFP reporter

mouse.   Whole mount imagining of axillary adipose tissue clearly demonstrated that

Cx3CR1$^+$ macrophages were present at the surface of and within the adipose lymph

node (LN; Fig. 5A-B). 3D reconstruction of adipose LN imaging with depth coding

allowed us to visualize Cx3CR1$^+$ macrophages present on the capsule as well as within

the cortex of the LN (Fig. 5B, middle and right panel). Furthermore, Cx3CR1$^+$

macrophages were enriched in adipose lymphatics (Fig. 5C left panel, yellow arrow)

compared to blood vasculature (Fig. 5C left panel, red arrow). We took advantage of

innate autofluorescence to visualize vasculature in these samples, and although blood

and lymph vasculature are somewhat morphologically similar, the bulbous sacs found at

capillary terminals are a unique feature of lymphatic vasculature only (Fig. C middle and

right panels, bulbous sacs outlined in white). These lymphatics were the vessels that

predominantly contained Cx3CR1^{+} cells, and not the blood vasculature.

Confocal imaging of Cx3CR1^{+} macrophages on lymphatic endothelium clearly

demonstrated that these cells have a distinct morphology (Fig. 5D left panel). The cell

bodies are elongated with cytoplastic extensions creating a spindeloid shape associated

with alternatively activated macrophages (54). 3D reconstruction with depth coding

following confocal imaging of adipose lymph vessels revealed that Cx3CR1^{+}

macrophages did not merely interact with lymph endothelial cells but were present

within the lumen of the vessels (Fig. 5D, middle and right panels). Cx3CR1^{+}

macrophages were present in greater numbers around and within the lymphatics of

adipose (Fig. 5E), and as previously reported were also associated with adipose nerves

(Fig. 5F) (55). Taken together, these data indicated that Cx3CR1+ macrophages are

transported by the lymphatic vasculature to scWAT.

Discussion

Here we present evidence of the necessity for myeloid-derived BDNF in maintaining inguinal scWAT innervation. Loss of BDNF from LysM+ myeloid cells resulted in a decrease in total innervation of the inguinal adipose tissue as measured by protein levels of the pan-neuronal marker PGP9.5, and a near complete denervation of sympathetic nerves (as measured by TH protein expression). We attribute this 'genetic denervation' as a cause for the observed decrease in energy expenditure exhibited under basal conditions, impaired thermogenic potential evidenced by lack of UCP1 induction and increased WAT mass during cold stimulation, and worsened response to a high-fat, high-sugar diet. These phenotypes fit with the known roles of adipose nerves in regulating proper adipogenesis, lipolysis, thermogenesis and overall metabolic health in adipose tissues.

Diet-induced obesity results in chromic low grade adipose inflammation due to infiltration of pro-inflammatory immune cells to the tissue, and ATM content increase from 10-15% (in lean mice) to ~50% with obesity (36, 56). In addition, phenotypic switching of CD4+ T cells and recruitment of T cell and B cells precedes macrophage infiltration, and with obesity macrophage polarity skews to a more inflammatory phenotype.

We initially hypothesized that anti-inflammatory (M2) ATMs were the source of scWAT BDNF. However, we observed no changes in M1 or M2 populations in scWAT after cold exposure. These M1 and M2 designations are an oversimplification of the diversity of macrophage populations and represent diverse subsets of cells with unique molecular signatures. Instead of changes in M1 or M2, we saw an increase in

Ly6C$^+$CCR2$^+$Cx3CR1$^+$ monocytes in scWAT of both male and female cold-exposed

mice. We have named these cells CINCs since they are recruited to scWAT by cold and

express BDNF. Surprisingly, it would appear that CINCs are a pro-inflammatory cell

type as Ly6C$^+$ and CCR2$^+$ are markers of inflammatory monocytes. Although pro-

inflammatory cell infiltration to adipose tissue is usually a harbinger of metabolic

dysfunction, an acute inflammatory response is necessary in wound healing and tissue

remodeling. Only chronic, ongoing inflammation would be detrimental and causative of

insulin resistance.

On the other hand the, the Cx3CR1$^+$ macrophages we identified in adipose

lymphatics exhibited features (such as elongated cell bodies) that are morphologically

attributed to M2 or anti-inflammatory macrophages (54) further underscoring the

incredible diversity of macrophage characterization in adipose. By utilizing an unbiased

20 antibody labeling approach to assess adipose SVF, we have discovered subsets of

myeloid cells that respond to these particular environmental stimuli, providing further

granularity to the myriad of immune cells types active in adipose.

Recently, a CX3CR1$^+$ population of macrophages had been described in

association with sympathetic nerves in adipose tissue (55). These sympathetic nerve

associated macrophages (SAMs) were found to regulate catecholamine levels on

adipose tissue by phagocytosing and degrading norepinephrine (NE) (55). Whether

SAMs are the same cells as our CINCs remains uncertain. Similar to CINCs, SAMs _do_

exhibit pro-inflammatory markers, and a distinct morphology similar to the Cx3CR1$^+$

macrophages we identified in the adipose lymphatics. SAMs home to WAT in the

obese state, while conversely, we see CINCs home to WAT with cold/noradrenergic-

stimulation. Given that these two populations home to WAT in different conditions, this

could indicate that the two immune cell types may have opposing roles, including

potential for phagocytosing nerve debris for SAMs and stimulating nerve regeneration

for CINCs. Fitting with this view, SAMs sequester and degrade NE, while CINCs

express BDNF.

On the other hand, CxCR1$^+$ macrophages have been shown to play diverse and

even opposing roles in the intestines (57), a phenotypic plasticity which may also be

present in adipose tissue. One thing that is clear is the M1/M2 paradigm of ATM

classification is an oversimplification of the functionally distinct populations of

macrophages active in adipose tissue. Further markers beyond Cx3CR1$^+$ are necessary

to fully understand and phenotype the rich variety of macrophages in adipose.

Compounding the difficulty in clearly delineating macrophage populations is that

macrophage activation is a dynamic process. Phenotypic switching appears to be

sequential, responding to microenvironment stimuli (58) and may be dependent on

spatiotemporal differences in tissue resident immune cell subtypes (59).

Since CINCs home into adipose tissue, we used a Cx3CR1-EGFP reporter

mouse to visualize how this happens. Immune cells are recruited to tissues via

vasculature, however, we found it surprising that in adipose it appeared that lymphatic

and not blood vasculature plays a more important recruitment role for CINCs. Strikingly,

the majority of Cx3CR1$^+$ cells were contained within the lymph node and lymphatic

vessels of adipose tissue. Because lymph flows in a unidirectional manner, the

lymphatic system is often overlooked as a source of bidirectional crosstalk. In fact,

lymphatic cells are not all destined for unidirectional trafficking. Leukocytes can also exit

lymphatic vasculature and enter surrounding tissues (60).

We have previously reported increased innervation around LNs in adipose

following cold exposure (8), while others have shown that browning commonly occurs

around LNs (61). Furthermore, all lymph nodes in the body are surrounded by adipose

tissue, which is activated during an immune response (62). Together these data

underscore the potential crosstalk between nerves, adipocytes, and leukocytes in

scWAT. Regardless of whether or not CINCs are recruited to or merely drained by

lymphatic vessels, the importance of the lymphatic system in transporting adipose

immune cells is again underscored by our current findings.

We postulate that CINCs contribute to nerve remodeling under noradrenergic

stimulation, leading to BDNF release (Suppl. Fig. S4B). Cao, *et al.* have recently

suggested that sympathetic nerve plastically is dependent on cold-induced, adipose-

derived NGF (63). Although they provide evidence that NGF is involved to some

degree in promoting cold induced sympathetic nerve density and browning of scWAT,

the cellular source of NGF was ambiguous (our data indicate it is produced more by

mature adipocytes than SVF cells, and thus may be promoting innervation of the

adipocytes and not the SVF). After determining that NGF gene expression increased in

the first couple of days of cold exposure, Cao, *et al.* used an NGF neutralizing antibody

to prevent NGF activity and observed decreased sympathetic nerve density in response

to cold. Their approach in preventing TrkA receptor function in adipose nerves and

showing decreased browning and sympathetic nerve density lends credence to the

notion that NGF plays a role in adipose innervation, similar to the role we observe for

BDNF. Taken together it is clear that locally-derived nerve growth factors are important

for maintaining proper brain-adipose communication and adipose innervation for

metabolic control. Considering the variety of NFs and the multitude of cell types in

adipose tissue it would not be surprising if both NGF and BDNF play a role in adipose

innervation through different mechanisms. Promoting innervation of a specific axonal

subtype to target a specific cell in the tissue may in fact be accomplished by

coordinated secretion of a given growth factor by the target cell type; whereby mature

adipocytes secrete NGF to 'wire up' adipocytes to TrkA+ axons, whereas SVF immune

cells secrete BDNF to 'wire up' immune cells to TrkB+ axons. The role of BDNF in

peripheral nerve health is often overlooked in lieu of the overwhelming focus on central

BDNF action. However, BDNF is known to play an important role in peripheral nerve

regeneration, as reviewed by *McGregor & English* (64). Following peripheral nerve

injury local sources of secreted BDNF include not only dorsal root ganglia neurons but

support cells, including Schwann cells (64). Cx3CR1+ macrophages have also been

shown to play a crucial role in peripheral nerve regeneration by regulating Schwann cell

dynamics in mice and humans (65). These studies align with the data presented here

and offer a new avenue of research in neuroimmune regulation of peripheral nerve

plasticity.

CONCLUSIONS

In conclusion, we have demonstrated the importance of CINC immune cells for

maintaining adipose innervation, particularly after cold exposure, which promotes CINC

movement to adipose (likely via adipose lymphatic vasculature). CINCs express and

likely secrete BDNF, thus promoting axonal survival and metabolic function in the

tissue. CINCs also express adrenergic receptor, making them a likely target for cold-

stimulated NE release in order to promote BDNF secretion to scWAT, once they have

homed to the tissue. Given these findings, it is clear that neuroimmune crosstalk in

scWAT is important for tissue function and metabolic control.

Availability of data and materials

Data available upon request.

List of abbreviations

ADRβ3: β-3 adrenergic receptor

ATMs: Adipose tissue macrophages

BAT: Brown adipose tissue

BDNF: Brain derived neurotrophic factor

BMI: Body mass index

CINCs: Cold induced neuroimmune cells

CNS: Central nervous system

CON: Control

FACS: Fluorescence-activated cell sorting

GWAS: Genome wide associated studies

HFD: High fat diet

KO: Knock-out

LN: Lymph node

MACS: Magnetic-activated cell sorting

NE: Norepinephrine

NFs: neurotrophic factors

NGF: nerve growth factor

NMJs: Neuromuscular junctions

NT-3: Neurotrophin-3

NT-4/5: Neurotrophin-4/5

PGP9.5: Protein gene product 9.5

RER: Respiratory exchange ratio

SAMs: Sympathetic nerve associated macrophages

scWAT: Subcutaneous white adipose tissue

SiLN: Subiliac lymph node

SNPs: Single nucleotide polymorphisms

SNS: Sympathetic nervous system

SVF: Stromal vascular fraction

TH: Tyrosine hydroxylase

tSNE: T-distributed Stochastic Neighbor Embedding

UCP1: Uncoupling protein 1

VEGFa: Vascular endothelial growth factor a

WAT: White adipose tissue

Acknowledgements

The authors wish to thank the following individuals from University of Maine: Brenda

Kennedy-Wade and Prof. James Weber for assistance in the animal facility; Dawna

Beane for paraffin embedding; Callie Greco, Cameron Fudge, and Sarah Berez for

animal care. The authors also wish to thank Drs. Gregory Cox and Robert Burgess

from Jackson Laboratory for sharing their NMJ assessment methods; Dr. Maribel Rios

for expert advice; and Nicholas Cutter and Emma Bragdon for help with spinal, motor,

and sensory nerve assessments.

Funding

788 For this project, the Townsend Lab was funded by start-up funds from the Office of the

789 Vice President for Research at University of Maine, a Rising Tide Seed Grant from

790 University of Maine, a Junior Faculty Award from the American Diabetes Association (1-

791 14-JF-55), and an NIH NIDDK 1R01DK114320-01A1. The Godwin lab is supported by a

792 NIH COBRE award (P20GM104318).

Author Contributions

MB designed experiments, analyzed data, and wrote the manuscript. EW and SK

collected and analyzed histology data. SK and RA collected and analyzed data related

to brown adipose tissue. JG conducted the flow cytometric analysis on adipose immune

cells. YHT oversaw initial pilot experiments. KLT designed experiments, analyzed data,

conceived of and oversaw the project, and wrote the manuscript. KLT is the guarantor

of this work and, as such, had full access to all the data in the study and takes

responsibility of the data and the accuracy of the data analysis.

Declarations

The authors do not have any conflicts of interest to disclose.

- **Ethical Approval and Consent to participate:** All animal studies were IACUC approved at

University of Maine

- **Consent for publication:** All authors have consented for publication.

  Figure Legends

Figure 1. BDNF is expressed in scWAT SVF fraction; *LysMCre$^{+/-}$::BDNF$^{-/-}$* (KO)

mice have lower energy expenditure. Adult male CB57/BL6 mice were cold exposed

at 5°C and SVF was isolated from mature adipocytes of the inguinal subcutaneous

white adipose (scWAT) depot. Differences in gene expression between adipose

compartments of neurotrophic factors, *Bdnf, Ngf,* and *Vegfa,* is shown (A). Data

analyzed by Student's t-test, two-tailed, N=4. Illustration of knock-out model generation

(B). BDNF gene expression in adipose SVF of *LysMCre$^{-/-}$::BDNF$^{fl/fl}$* (CON) versus

LysMCre$^{+/-}$::BDNF$^{-/-}$ (KO) mice (C). Data analyzed by Student's t-test, two-tailed, N=4

per group. Gene expression of *Bdnf* in hypothalamus of *LysMCre$^{-/-}$::BDNF$^{fl/fl}$* (CON)

versus *LysMCre$^{+/-}$ ::BDNF$^{-/-}$* (KO) mice (D). Data analyzed by Student's t-test, two-tailed,

N=5 CON; N=6 KO. Adult (8-12 week old) male CON and KO mice were assessed in

metabolic cages (CLAMS). KO mice displayed lower energy expenditure represented as

heat calculated from measures of VO_2 and VCO_2 over the whole 24hr (E). Waveform

analysis of metabolic cage measurements taken at 15min increments for 48 hrs. Time

of day is indicated on the x-axis, animals were maintained on a 12 hr light/dark cycle

(black bars indicate dark cycle). For all error bars are SEMs.

Figure 2. *LysMCre$^{+/-}$::BDNF$^{-/-}$* **(KO) have increased adiposity and impaired**

response to cold due to genetic denervation of scWAT. Adult (22-23 week old) male

LysMCre$^{-/-}$::BDNF$^{fl/fl}$ (CON) and *LysMCre$^{+/-}$ ::BDNF$^{-/-}$* (KO) mice were cold exposed at

5°C for 4 days; body weight and adiposity were compared between CON and KO

groups (A). Body and tissue weight data were analyzed by two-tailed Student's T-Test,

N=5 CON, N=6 KO. Protein expression of PGP9.5 and tyrosine hydroxylase (TH) in

inguinal scWAT was measured by western blotting from adult (12-25 week old) 7-day

cold (5°C) exposed WT/CON and KO male animals (B). β-tubulin was used as a loading

control for normalization. Data were analyzed by two-tailed Student's T-Test, N=3

WT/CON, N=4 KO, * denotes data that was removed from analysis due to lack of

expression of loading control. Gene expression of *Ucp1* was measured in adult (12-25

week old) 7-day cold (5°C) exposed WT/CON and KO males (C). Data were analyzed

by two-tailed Student's T-Test, N=5 WT/CON, N=5 KO. Circulating thyroid hormones,

thyroxine (T4) and triiodothyronine (T3), were measured by ELISA from serum of adult

(22-23 week old) 4-day cold (5°C) exposed CON and KO male mice (D). Data were

analyzed by One-way ANOVA, with Tukey's multiple comparisons test, N=5 CON, N=6

KO. Immunofluorescent staining for UCP1 was performed on inguinal scWAT sections

of adult (22-23 week old) male CON and KO mice following 4-day cold (5°C) exposure

(E). Immunofluorescent staining for UCP1 and Perilipin was performed on inguinal

scWAT sections of adult (22-23 week old) male KO mice following 4-day cold (5°C)

exposure (F). Typogen Black, used to quench lipid autofluorescence, provided staining

cell morphology which was visualized under brightfield microscopy.   Overlay is

immunofluorescence over brightfield of the same area (F). Images were acquired with a

10X or 40X objective and are representative of N=5 CON, N=8 KO (E) and N=4 KO (F).

For all error bars are SEMs.

Figure 3. *LysMCre$^{+/-}$ ::BDNF$^{-/-}$* (KO) showed accelerated fat accumulation on a 45%

HFD. Adult (25 week old) male *LysMCre$^{-/-}$::BDNF$^{fl/fl}$* (CON) and *LysMCre$^{+/-}$ ::BDNF$^{-/-}$*

(KO) were challenged with a 45% HFD for 3 weeks before undergoing physiological

assessment in metabolic cages (A-B). Energy expenditure as measured by heat was

lower for KO versus CON only for a short period during the dark cycle (A). Respiratory

exchange as a ratio (RER) between the two groups, indicated greater use of

carbohydrates for fuel by KO animals during the light cycle (B). Data presented as

waveform analysis of measurements taken at 15min increments for 3 days. Time of day

is indicated on the x-axis, and animals were maintained on a 12 hr light/dark cycle

(black bars indicate dark cycle)**.** Data analyzed by two-way repeated measures ANOVA

with Fisher's LSD test; N=4 per group. Adult (25 week old) male CON and KO animals

were placed on a 45% HFD, daily food intake (represented as cumulative food intake)

(C) was measured for the first week of HFD feeding. Percent change in body weight (D)

was measured for the first 7 days of HFD feeding. Data were analyzed by two-tailed

Student's T-Test, N=5 CON, N=7 KO. Glucose tolerance testing was performed at 6

weeks of HFD feeding (E). Data were analyzed by Two-way ANOVA, with Tukey's

multiple comparisons test, N=5 CON, N=5 KO. Adiposity was measured for CON and

KO animals after 11 weeks of HFD feeding as a percentage of scWAT over body weight

(F). Data were analyzed by two-tailed Student's T-Test, N=5 CON, N=7 KO. For all error

bars are SEMs. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

Figure 4. Cold induced neuroimmune cells (CINCs) home in to inguinal scWAT and express _Bdnf_. Adult (12 week old) male C57BL/6 were either maintained at room temperature (RT) or cold exposed (5°C) for 5 days, ATMs from inguinal scWAT depots were isolated using magnetic-activated cell sorting (MACS) by positive selection of CD11b$^+$ followed by F4/80$^+$ cells. _Bdnf_ gene expression in doubly labeled CD11b$^+$ F4/80$^+$ macrophages was compared between RT and cold exposed animals (A). Data were analyzed by two-tailed Student's T-Test, N=4 per group. Adult (12 week old) female control animals were either maintained at room temperature (RT) or cold exposed (5°C) for 10 days, SVF from bilateral inguinal scWAT was isolated and FACS sorted using a 20 cell surface marker panel for myeloid lineage immune cells (B-C). Changes in M1/M2 polarity (B) and Ly6C$^+$CCR2$^+$Cx3CR1$^-$ and Ly6C$^+$CCR2$^+$Cx3CR1$^+$ macrophage precursors/monocytes (C, left panel) were measured between RT and cold exposed animals. Adult (12-13 week old) male C57BL/6 were either maintained at room temperature (RT) or cold exposed (5°C) for 14 days; SVF from bilateral inguinal scWAT was isolated and FACS sorted; Ly6C$^+$CCR2$^+$Cx3CR1$^-$ and Ly6C$^+$CCR2$^+$Cx3CR1$^+$ cells were compared between RT and cold exposed animals (C, right panel). Data were analyzed by two-tailed Student's T-Test, N=3-4 per group for both sexes. Adult male (M) and female (F) control animals were either maintained at room temperature (RT) or cold exposed (5°C) for 10 days, SVF from bilateral inguinal scWAT was isolated and FACS sorted using a 20 cell surface marker panel for myeloid lineage immune cells (N=5 per group). t-Distributed Stochastic Neighbor Embedding (tSNE) analysis was performed to identify myeloid lineage cell population changes in response to cold exposure; Ly6C$^+$CCR2$^+$Cx3CR1$^+$ were identified for both sexes as adipose cold induced neuroimmune cells (CINCs) (D). _Bdnf_ gene expression measured in

Ly6C$^+$CCR2$^+$Cx3CR1$^+$ cells indicated that these infiltrating cells express *Bdnf* (E). Data

analyzed by two-tailed Student's T-Test, N=4 per group. For all error bars are SEMs. *p

< 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

Figure 5. Cx3CR1 Cells in Adipose Transit via Lymphatics. Wholemount imaging of

female (A-E) and male (F) axillary scWAT from Cx3CR1-EGFP reporters mice.

Cx3CR1+ cells (green) in lymph node captured by widefield microscopy at 10X (A).

Cx3XR1+ cells were shown to occupy the entire depth of the lymph node, captured by

confocal microscopy at 20X (B), represented as a z-maximum projection (Glow LUT) (B,

left) and 3D reconstruction (depth coded) at two angles (B, middle and right.) Cx3CR1+

cells (green) line lymphatic vasculature (morphologically distinguished from blood

vasculature by the bulbous sacs on initial lymphatics, yellow arrow and outlined in

white) but are absent in blood vasculature (red arrow) (C). Captured by widefield

microscopy at 10X and 40X. Cx3CR1+ cells (Glow LUT) imaged by confocal

microscopy at 63X were shown to reside on lymph vessel endothelium and were

present within the lumen (D). Z-maximum projection of lymph vessel (D, left) and 3D

reconstruction (depth coded) at two angles: looking down z-axis (D, middle) and looking

down the x-axis (D, right) with vessel lumen identified by arrow. Tiled z-maz projection

captured on confocal at 20X demonstrates that Cx3CR1+ cells (white) are found

throughout the lymphatic network in scWAT (E). Cx3CR1+ cells (green) reside around

nerve bundles marked by the pan-neuronal marker PGP9.5 (magenta) as captured by

widefield microscopy at 40X (F).

Supplemental Information
928
929 *Supplemental Figure S1*
930 *Supplemental Figure S2*
931 *Supplemental Figure S3*
932 *Supplemental Figure S4*
933 *Supplemental Figure Legends*
934

References

1.	Dulloo AG, Miller DS. Energy balance following sympathetic denervation of brown adipose tissue. CanJ PhysiolPharmacol. 1984;62(2):235-40.

2.	Minokoshi Y, Saito M, Shimazu T. Sympathetic denervation impairs responses of brown adipose tissue to VMH stimulation. Am J Physiol. 1986;251(5 Pt 2):R1005-R8.

3.	Desautels M, Dulos RA, Mozaffari B. Selective loss of uncoupling protein from mitochondria of surgically denervated brown adipose tissue of cold-acclimated mice. BiochemCell Biol. 1986;64(11):1125-34.

4.	Dulloo A, Miller D. Energy balance following sympathetic denervation of brown adipose tissue. Canadian journal of physiology and pharmacology. 1984;62(2):235-40.

5.	Bowers RR, Festuccia WT, Song CK, Shi H, Migliorini RH, Bartness TJ. Sympathetic innervation of white adipose tissue and its regulation of fat cell number. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology. 2004;286(6):R1167-R75.

6.	Foster MT, Bartness TJ. Sympathetic but not sensory denervation stimulates white adipocyte proliferation. American journal of physiology Regulatory, integrative and comparative physiology. 2006;291(6):R1630-7.

7.	Harris RB. Sympathetic denervation of one white fat depot changes norepinephrine content and turnover in intact white and brown fat depots. Obesity(SilverSpring). 2012:10.

8.	Blaszkiewicz M, Willows JW, Dubois AL, Waible S, DiBello K, Lyons LL, et al. Neuropathy and neural plasticity in the subcutaneous white adipose depot. PLoS One. 2019;14(9):e0221766.

9.	Gray JD, Milner TA, McEwen BS. Dynamic plasticity: the role of glucocorticoids, brain-derived neurotrophic factor and other trophic factors. Neuroscience. 2013;239:214-27.

10.	Terenghi G. Peripheral nerve regeneration and neurotrophic factors. J Anat. 1999;194 (Pt 1):1-14.

11.	An JJ, Gharami K, Liao GY, Woo NH, Lau AG, Vanevski F, et al. Distinct role of long 3' UTR BDNF mRNA in spine morphology and synaptic plasticity in hippocampal neurons. Cell. 2008;134(1):175-87.

12.	Kleim JA, Chan S, Pringle E, Schallert K, Procaccio V, Jimenez R, et al. BDNF val66met polymorphism is associated with modified experience-dependent plasticity in human motor cortex. Nat Neurosci. 2006;9(6):735-7.

13.	Fritsch B, Reis J, Martinowich K, Schambra HM, Ji Y, Cohen LG, et al. Direct current stimulation promotes BDNF-dependent synaptic plasticity: potential implications for motor learning. Neuron. 2010;66(2):198-204.

14.	Tapia-Arancibia L, Rage F, Givalois L, Arancibia S. Physiology of BDNF: focus on hypothalamic function. Front Neuroendocrinol. 2004;25(2):77-107.

15.	Rios M, Fan G, Fekete C, Kelly J, Bates B, Kuehn R, et al. Conditional deletion of brain-derived neurotrophic factor in the postnatal brain leads to obesity and hyperactivity. Mol Endocrinol. 2001;15(10):1748-57.

16.	Unger TJ, Calderon GA, Bradley LC, Sena-Esteves M, Rios M. Selective deletion of Bdnf in the ventromedial and dorsomedial hypothalamus of adult mice results in hyperphagic behavior and obesity. J Neurosci. 2007;27(52):14265-74.

17.	An JJ, Liao GY, Kinney CE, Sahibzada N, Xu B. Discrete BDNF Neurons in the Paraventricular Hypothalamus Control Feeding and Energy Expenditure. Cell Metab. 2015;22(1):175-88.

18.	Roth CL, Elfers C, Gebhardt U, Muller HL, Reinehr T. Brain-derived neurotrophic factor and its relation to leptin in obese children before and after weight loss. Metabolism. 2013;62(2):226-34.

19.   Araki S, Yamamoto Y, Dobashi K, Asayama K, Kusuhara K. Decreased plasma levels of
brain-derived neurotrophic factor and its relationship with obesity and birth weight in obese
Japanese children. Obes Res Clin Pract. 2014;8(1):e63-9.
20.   Duan W, Guo Z, Jiang H, Ware M, Li XJ, Mattson MP. Dietary restriction normalizes
glucose metabolism and BDNF levels, slows disease progression, and increases survival in
huntingtin mutant mice. Proc Natl Acad Sci U S A. 2003;100(5):2911-6.
21.   Nakagawa T, Tsuchida A, Itakura Y, Nonomura T, Ono M, Hirota F, et al. Brain-derived
neurotrophic factor regulates glucose metabolism by modulating energy balance in diabetic mice.
Diabetes. 2000;49(3):436-44.
22.   Nonomura T, Tsuchida A, Ono-Kishino M, Nakagawa T, Taiji M, Noguchi H. Brain-
derived neurotrophic factor regulates energy expenditure through the central nervous system in
obese diabetic mice. Int J Exp Diabetes Res. 2001;2(3):201-9.
23.   Xu B, Goulding EH, Zang K, Cepoi D, Cone RD, Jones KR, et al. Brain-derived
neurotrophic factor regulates energy balance downstream of melanocortin-4 receptor. Nat
Neurosci. 2003;6(7):736-42.
24.   Xu B, Xie X. Neurotrophic factor control of satiety and body weight. Nat Rev Neurosci.
2016;17(5):282-92.
25.   Sornelli F, Fiore M, Chaldakov GN, Aloe L. Adipose tissue-derived nerve growth factor
and brain-derived neurotrophic factor: results from experimental stress and diabetes. Gen Physiol
Biophys. 2009;28 Spec No:179-83.:179-83.
26.   Hausman GJ, Poulos SP, Richardson RL, Barb CR, Andacht T, Kirk HC, et al. Secreted
proteins and genes in fetal and neonatal pig adipose tissue and stromal-vascular cells. J Anim
Sci. 2006;84(7):1666-81.
27.   Bernhard F, Landgraf K, Kloting N, Berthold A, Buttner P, Friebe D, et al. Functional
relevance of genes implicated by obesity genome-wide association study signals for human
adipocyte biology. Diabetologia. 2013;56(2):311-22.
28.   Nakagomi A, Okada S, Yokoyama M, Yoshida Y, Shimizu I, Miki T, et al. Role of the
central nervous system and adipose tissue BDNF/TrkB axes in metabolic regulation. NPJ Aging
Mech Dis. 2015;1:15009.
29.   Goss JR, O'Malley ME, Zou L, Styren SD, Kochanek PM, DeKosky ST. Astrocytes are
the major source of nerve growth factor upregulation following traumatic brain injury in the rat.
Exp Neurol. 1998;149(2):301-9.
30.   Madduri S, Gander B. Schwann cell delivery of neurotrophic factors for peripheral nerve
regeneration. J Peripher Nerv Syst. 2010;15(2):93-103.
31.   Leon A, Buriani A, Dal Toso R, Fabris M, Romanello S, Aloe L, et al. Mast cells
synthesize, store, and release nerve growth factor. Proc Natl Acad Sci U S A. 1994;91(9):3739-
43.
32.   Kobayashi H, Gleich GJ, Butterfield JH, Kita H. Human eosinophils produce
neurotrophins and secrete nerve growth factor on immunologic stimuli. Blood. 2002;99(6):2214-
20.
33.   Noga O, Englmann C, Hanf G, Grutzkau A, Seybold J, Kunkel G. The production,
storage and release of the neurotrophins nerve growth factor, brain-derived neurotrophic factor
and neurotrophin-3 by human peripheral eosinophils in allergics and non-allergics. Clinical and
experimental allergy : journal of the British Society for Allergy and Clinical Immunology.
2003;33(5):649-54.
34.   Lambiase A, Bracci-Laudiero L, Bonini S, Bonini S, Starace G, D'elios MM, et al.
Human CD4+ T cell clones produce and release nerve growth factor and express high-affinity
nerve growth factor receptors. Journal of Allergy and Clinical Immunology. 1997;100(3):408-14.

35.   Kerschensteiner M, Gallmeier E, Behrens L, Leal VV, Misgeld T, Klinkert WE, et al.
Activated human T cells, B cells, and monocytes produce brain-derived neurotrophic factor in
vitro and in inflammatory brain lesions: a neuroprotective role of inflammation? Journal of
Experimental Medicine. 1999;189(5):865-70.
36.   Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW, Jr. Obesity
is associated with macrophage accumulation in adipose tissue. J Clin Invest. 2003;112(12):1796-
808.
37.   Winer DA, Winer S, Shen L, Wadia PP, Yantha J, Paltser G, et al. B cells promote
insulin resistance through modulation of T cells and production of pathogenic IgG antibodies.
Nat Med. 2011;17(5):610-7.
38.   Rausch ME, Weisberg S, Vardhana P, Tortoriello DV. Obesity in C57BL/6J mice is
characterized by adipose tissue hypoxia and cytotoxic T-cell infiltration. Int J Obes (Lond).
2008;32(3):451-63.
39.   Elgazar-Carmon V, Rudich A, Hadad N, Levy R. Neutrophils transiently infiltrate intra-
abdominal fat early in the course of high-fat feeding. J Lipid Res. 2008;49(9):1894-903.
40.   Liu J, Divoux A, Sun J, Zhang J, Clement K, Glickman JN, et al. Genetic deficiency and
pharmacological stabilization of mast cells reduce diet-induced obesity and diabetes in mice.
NatMed. 2009;15(8):940-5.
41.   Strissel KJ, DeFuria J, Shaul ME, Bennett G, Greenberg AS, Obin MS. T-cell
recruitment and Th1 polarization in adipose tissue during diet-induced obesity in C57BL/6 mice.
Obesity (Silver Spring). 2010;18(10):1918-25.
42.   Grant R, Youm YH, Ravussin A, Dixit VD. Quantification of adipose tissue leukocytosis
in obesity. Methods Mol Biol. 2013;1040:195-209.
43.   Hui X, Gu P, Zhang J, Nie T, Pan Y, Wu D, et al. Adiponectin Enhances Cold-Induced
Browning of Subcutaneous Adipose Tissue via Promoting M2 Macrophage Proliferation. Cell
Metab. 2015;22(2):279-90.
44.   Rost B, Hanf G, Ohnemus U, Otto-Knapp R, Groneberg DA, Kunkel G, et al. Monocytes
of allergics and non-allergics produce, store and release the neurotrophins NGF, BDNF and NT-
3. Regul Pept. 2005;124(1-3):19-25.
45.   Parkhurst CN, Yang G, Ninan I, Savas JN, Yates JR, 3rd, Lafaille JJ, et al. Microglia
promote learning-dependent synapse formation through brain-derived neurotrophic factor. Cell.
2013;155(7):1596-609.
46.   Lai SW, Chen JH, Lin HY, Liu YS, Tsai CF, Chang PC, et al. Regulatory Effects of
Neuroinflammatory Responses Through Brain-Derived Neurotrophic Factor Signaling in
Microglial Cells. Mol Neurobiol. 2018;55(9):7487-99.
47.   Gemma C, Bachstetter AD. The role of microglia in adult hippocampal neurogenesis.
Front Cell Neurosci. 2013;7:229.
48.   Willows JW, Blaszkiewicz M, Lamore A, Borer S, Dubois AL, Garner E, et al.
Visualization and Analysis of Whole Depot Adipose Tissue Innervation. bioRxiv. 2019:788885.
49.   Gray NE, Lam LN, Yang K, Zhou AY, Koliwad S, Wang JC. Angiopoietin-like 4
(Angptl4) protein is a physiological mediator of intracellular lipolysis in murine adipocytes. J
Biol Chem. 2017;292(39):16135.
50.   Nielsen TS, Jessen N, Jorgensen JO, Moller N, Lund S. Dissecting adipose tissue
lipolysis: molecular regulation and implications for metabolic disease. J Mol Endocrinol.
2014;52(3):R199-222.
51.   Bartness TJ, Song CK. Brain-adipose tissue neural crosstalk. Physiol Behav.
2007;91(4):343-51.

52. Bartness TJ, Liu Y, Shrestha YB, Ryu V. Neural innervation of white adipose tissue and
the control of lipolysis. Front Neuroendocrinol. 2014;35(4):473-93.
53. Getts DR, Terry RL, Getts MT, Muller M, Rana S, Shrestha B, et al. Ly6c+
"inflammatory monocytes" are microglial precursors recruited in a pathogenic manner in West
Nile virus encephalitis. J Exp Med. 2008;205(10):2319-37.
54. McWhorter FY, Wang T, Nguyen P, Chung T, Liu WF. Modulation of macrophage
phenotype by cell shape. Proc Natl Acad Sci U S A. 2013;110(43):17253-8.
55. Pirzgalska RM, Seixas E, Seidman JS, Link VM, Sánchez NM, Mahú I, et al.
Sympathetic neuron–associated macrophages contribute to obesity by importing and
metabolizing norepinephrine. Nature medicine. 2017;23(11):1309.
56. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, et al. Chronic inflammation in fat
plays a crucial role in the development of obesity-related insulin resistance. J Clin Invest.
2003;112(12):1821-30.
57. Regoli M, Bertelli E, Gulisano M, Nicoletti C. The Multifaceted Personality of Intestinal
CX3CR1(+) Macrophages. Trends Immunol. 2017;38(12):879-87.
58. Stout RD, Jiang C, Matta B, Tietzel I, Watkins SK, Suttles J. Macrophages sequentially
change their functional phenotype in response to changes in microenvironmental influences. J
Immunol. 2005;175(1):342-9.
59. Lumeng CN, Delproposto JB, Westcott DJ, Saltiel AR. Phenotypic switching of adipose
tissue macrophages with obesity is generated by spatiotemporal differences in macrophage
subtypes. Diabetes. 2008;57(12):3239-46.
60. Jackson DG. Leucocyte Trafficking via the Lymphatic Vasculature- Mechanisms and
Consequences. Front Immunol. 2019;10:471.
61. Dichamp J, Barreau C, Guissard C, Carriere A, Martinez Y, Descombes X, et al. 3D
analysis of the whole subcutaneous adipose tissue reveals a complex spatial network of
interconnected lobules with heterogeneous browning ability. Sci Rep. 2019;9(1):6684.
62. Pond CM, Mattacks CA. The activation of the adipose tissue associated with lymph
nodes during the early stages of an immune response. Cytokine. 2002;17(3):131-9.
63. Cao Y, Wang H, Zeng W. Whole-tissue 3D imaging reveals intra-adipose sympathetic
plasticity regulated by NGF-TrkA signal in cold-induced beiging. Protein Cell. 2018;9(6):527-
39.
64. McGregor CE, English AW. The Role of BDNF in Peripheral Nerve Regeneration:
Activity-Dependent Treatments and Val66Met. Front Cell Neurosci. 2018;12:522.
65. Stratton JA, Holmes A, Rosin NL, Sinha S, Vohra M, Burma NE, et al. Macrophages
Regulate Schwann Cell Maturation after Nerve Injury. Cell Rep. 2018;24(10):2561-72 e6.
1113

A.

Gene Expression of Neurotrophic/Growth Factors in Adipose Compartments

Bdnf — *Ngf* — *Vegfa*

B. Generation of Knock-Out (KO) Model

KO mice lack BDNF in myeloid lineage cells, including monocytes/macrophages

C. *Bdnf* Gene Expression in Inguinal scWAT SVF of *LysMCre⁺/⁻:BDNF⁻/⁻* (KO) mice

D. *Bdnf* Gene Expression in Hypothalamus of *LysMCre⁺/⁻:BDNF⁻/⁻* (KO) mice

E.

Basal Heat (CLAMS) Waveform Analysis

Figure 2

A.

Body Weight after Cold Exposure

p=0.0750

CON   KO

scWAT/ Body Weight after Cold Exposure

p=0.1107

CON   KO

B. Inguinal scWAT Innervation after Cold Exposure

p=0.0025

CON   KO

p=0.0200

CON   KO

PGP9.5
Tyrosine Hydroxylase
β-Tubulin

CON   KO

C. *Ucp1* Gene Expression in Inguinal scWAT after Cold Exposure

p=0.0573

CON   KO

D. Circulating Thyroid Hormones after Cold Exposure

p=0.9526   p=0.3471

CON T3   KO T3   CON T4   KO T4

E. Histology and UCP1 Protein in Inguinal scWAT Following Cold Exposure

Typogen Black   UCP1   Typogen Black   UCP1

CON

KO

10X   10X

F. UCP1 Negative Multilocular Cells in KO scWAT Express Perilipin

UCP1 / Perilipin   Typogen Black   Overlay   UCP1 / Perilipin   Typogen Black   Overlay

10 X   40 X

Figure 4

A. *Bdnf* Gene Expression in F4/80⁺ Macrophages

B. Macrophage Polarity

C. CINC Recruitment to Inguinal scWAT Following Cold Exposure

D. tSNE Analysis of Male/Female Cold Induced Immune Changes in Inguinal scWAT

Male RT — MRT1, MRT3, MRT2

Male Cold — MC1, MC3, MC2

Female RT — FRT5, FRT4, FRT1, FRT7, FRT6, FRT8, FRT2, FRT9, FRT3

Female Cold — FC4, FC1, FC3, FC2

Population	%	Predicted Identity	Markers	
MRT1	9.2	Myeloid	CD11b⁺ CD62L⁺	Present in RT, Lost on Cold
MRT2	3.8	APC	CD11b^mid MHC⁺	Present in RT, Lost on Cold
MRT3	2.4	Myeloid	CD11b⁺ CD62L^mid	Present in RT, Lost on Cold
MC1	5.6	T cells	CD11b^mid CD3⁻	Appear on cold
MC2	11.4	**CINCs**	CD11b⁺ CCR2⁻ Cx3CR1⁺	Appear on cold
MC3	5.8	T cells	CD11b^mid CD3+	Appear on cold
FRT1	2.4	B cells	CD11b^mid CD19⁻	Present in RT, Lost on Cold
FRT2	2.9	Monocyte	CD11b⁺ Ly6C⁺	Present in RT, Lost on Cold
FRT3	3.2	APC	CD11b^mid NK1.1⁺ MHC⁺	Present in RT, Lost on Cold
FRT4	0.9	APC	CD11b^mid CD11c⁻	Present in RT, Lost on Cold
FRT5	1.8	APC	CD11b^mid CD62L⁺ Ly6C⁻ MHC^mid	Present in RT, Lost on Cold
FRT6	7.7	T cells	CD11b^mid CD3+	Present in RT, Lost on Cold
FRT7	2.1	T cells	CD11b^mid CD3+	Present in RT, Lost on Cold
FRT8	0.6	T cells	CD11b^mid CD3+	Present in RT, Lost on Cold
FRT9	0.6	APC	CD11b⁺ CD11c⁻	Present in RT, Lost on Cold
FC1	19.2	B cells	CD11b^mid CD19⁻	Appear on cold
FC2	1.1	B cells	CD11b^mid CD19⁻	Appear on cold
FC3	2.3	**CINCs**	CD11b⁺ CCR2⁻ Cx3CR1⁺	Appear on cold
FC4	2.5	T cells	CD11b^mid CD3⁻	Appear on cold

E. *Bdnf* Gene Expression in Ly6C⁺CCR2⁺Cx3CR1⁺ Macrophages (CINCs)

Figure 5

A. Adipose Lymph Node

Cx3CR1 EGFP

10X

B. Imaging with Depth Coding: Cx3CR1 Cells Present on Capsule and Within Cortex of Adipose Lymph Node

Z-max Projection Lymph Node

Cx3CR1 EGFP

20X

Depth Coding: 3D Reconstruction of Lymph Node

Depth Coding: 3D Reconstruction of Lymph Node

Capsule

Cortex

C. Adipose Lymphatic Vessels

Cx3CR1 EGFP

Lymph vessel

Blood vessel

10X

Cx3CR1 EGFP

10X

40X

D. Imaging with Depth Coding: Cx3CR1 Cells on Lymphatic Endothelium and in Lumen

Z-max Projection Lymphatic Vessel

Cx3CR1 EGFP

63X

Depth Coding: 3D Reconstruction Lymphatic Vessel (plane of view: xyz)

Depth Coding: 3D Reconstruction of Lymphatic Vessel (plane of view: zxy)

Endothelium

Endothelium

Lumen

E. Cx3CR1 Cells in Adipose Lymphatics

Cx3CR1 EGFP

20X tiled

F. Cx3CR1 Cells Associated with Nerve Bundles in Adipose

Cx3CR1 EGFP

PGP9.5

40X