Full Title: Facial shape and allometry quantitative trait locus intervals in the Diversity Outbred

mouse are enriched for known skeletal and facial development genes

Short title: The genetics of mouse facial shape variation

Authors

David C. Katz[1*], J. David Aponte[1], Wei Liu[1], Rebecca M. Green[1], Jessica M. Mayeux[2], K. Michael

Pollard[2], Daniel C. Pomp[3], Steven C. Munger[4], Steven A. Murray[4], Charles C. Roseman[5],

Christopher J. Percival[6], James Cheverud[7], Ralph S. Marcucio[8], Benedikt Hallgrímsson[1*]

Affiliations

[1] Department of Cell Biology & Anatomy, Alberta Children's Hospital Research Institute and

McCaig Bone and Joint Institute, Cumming School of Medicine, University of Calgary, AB,

Canada

[2] Department of Molecular Medicine, The Scripps Research Institute, La Jolla, CA, USA

[3] Department of Genetics, University of North Carolina Medical School, Chapel Hill, NC, USA

[4] The Jackson Laboratory, Bar Harbor, ME, USA

[5] Department of Animal Biology, University of Illinois Urbana Champaign, Urbana, IL, USA

[6] Department of Anthropology, Stony Brook University, Stony Brook, NY, USA

[7] Department of Biology, Loyola University Chicago, Chicago, IL, USA

[8] Department of Orthopaedic Surgery, School of Medicine, University of California San

Francisco, San Francisco, CA, USA

* Corresponding authors

bhallgri@ucalgary.ca, david.katz@ucalgary.ca

Author contributions

Conceptualization (Katz, Aponte, Hallgrimsson, Cheverud, Roseman); Data Curation (Pomp,

Murray, Pollard, Munger, Mayeux, Liu, Katz); Formal analysis (Katz); Funding acquisition

(Hallgrimsson); Investigation (Mayeux, Aponte, Katz, Liu); Methodology (Percival, Aponte,

Green, Katz, Cheverud, Roseman, Hallgrimsson); Resources (Murray, Pomp, Pollard, Munger);

Supervision (Cheverud, Roseman, Hallgrimsson); Writing—Original Draft: Katz; Writing—Review

and Editing: Everyone.

Abstract

The biology of how faces are built and come to differ from one another is complex. Discovering the genes that contribute to differences in facial morphology is one key to untangling this complexity, with important implications for medicine and evolutionary biology. This study maps quantitative trait loci (QTL) for skeletal facial shape using Diversity Outbred (DO) mice. The DO is a randomly outcrossed population with high heterozygosity that captures the allelic diversity of eight inbred mouse lines from three subspecies. The study uses a sample of 1147 DO animals (the largest sample yet employed for a shape QTL study in mouse), each characterized by 22 three-dimensional landmarks, 56,885 autosomal and X-chromosome markers, and sex and age classifiers. We identified 37 facial shape QTL across 20 shape principal components (PCs) using a mixed effects regression that accounts for kinship among observations. The QTL include some previously identified intervals as well as new regions that expand the list of potential targets for future experimental study. Three QTL characterized shape associations with size (allometry). Median support interval size was 3.5 Mb. Narrowing additional analysis to QTL for the five largest magnitude shape PCs, we found significant overrepresentation of genes with known roles in growth, skeletal development, and sensory organ development. For most intervals, one or more of these genes lies within 0.25 Mb of the QTL's peak. QTL effect sizes were small, with none explaining more than 0.5% of facial shape variation. Thus, our results are consistent with a model of facial diversity that is influenced by key genes in skeletal and facial development and, simultaneously, is highly polygenic.

Author Summary

The mammalian face is a complex structure serving many functions. We studied the

genetic basis for facial skeletal diversity in a large sample of mice from an experimental

population designed for the study of complex traits. We quantified the contribution of genetic

variation to variation in three-dimensional facial shape across more than 55,000 genetic

markers spread throughout the mouse genome. We found 37 genetic regions which are very

likely to contribute to differences in facial shape. We then conducted a more detailed analysis

of the genetic regions associated with the most variable aspects of facial shape. For these

regions, a disproportionately large number of genes are known to be important to growth and

to skeletal and facial development. The magnitude of these genetic contributions to differences

in facial shape are consistently small. Our results therefore support the notion that facial

skeletal diversity is influenced by many genes of small effect, but that some of these small

effects may be related to genes that are fundamental to skeletal and facial development.

Introduction

The facial skeleton is a complex morphology that serves a diverse set of critical roles. It

houses and protects the forebrain and most sensory organs and provides essential anatomy for

oral communication, feeding, and facial expression. Structured differences (covariance) in facial

shape and form depend upon variation in the biological processes that coordinate facial

development [1-7]. Understanding how these processes act and combine to build a facial

skeleton requires interrogation at multiple levels of the genotype-phenotype (GP) map [8, 9].

Among the most fundamental of these inquiries is the search for genetic variants that

contribute to facial variation. These genetic variants are the basic building blocks of diversity

and, within species, circumscribe the potential for short-term evolutionary change [10, 11].

Over the past decade, dozens of studies in humans, mouse models, and other species have

dissected GP associations for the craniofacial complex using moderate-to-high density genetic

maps (tens of thousands to millions of markers). The focus is typically on additive effects [12],

though some recent work considers interactions among loci [4, 13, 14].

In humans, genome-wide association studies (GWAS) targeting variation in soft tissue

facial form have been completed for European [15-20], African [21], South American [22], and

Asian [23] cohorts. Collectively, the studies report around 50 loci for facial morphology [24].

While there is limited overlap in the candidate gene sets among studies [25], and though many

of the individual genes have no previously known role in facial development [3], the genes that

do replicate across GWAS (*PAX 1*, *PAX3, HOXD* cluster genes, *DCHS2*, *SOX9*) are known to be

important to facial and/or skeletal morphogenesis.

Compared to human studies, mapping the genetics of facial variation in mice offers

several benefits, including greater control over genetic and environmental variation and the

ability to record observations, directly or *in silico*, on the skeleton itself [14, 26-31]. In recent

years, the focus of mouse skull QTL analyses has expanded to include tests of enrichment for

disease phenotypes and roles in skull development, as well as the search for and validation of

candidate genes [32-34]. Dozens of QTL have been identified. As with human facial shape

GWAS candidates, this is encouraging but also a small beginning—on the order of the total

number of loci identified for human height just a decade ago [35]. Quantitative genetics

analyses [32, 36, 37] and expression and epigenomic atlases of facial development both

strongly point to the potential for hundreds or thousands of genetic loci with as-yet-undetected

QTL variance contributions [38-40]. Increased sample sizes, greater mapping resolution, and the

development of new experimental lines all have the potential to expand the facial shape QTL

library and increase the precision of variant localization [41-43].

The present study maps mouse facial shape QTL in what is, to date, the largest sample

yet used for this purpose (n = 1147). We use Diversity Outbred mice (DO; Jackson Laboratory,

Bar Harbor, ME), a random outcross model derived from the Collaborative Cross (CC) founding

lines [43-46]. The DO is specifically designed for genetic mapping of complex traits. Each

animal's genome is a unique mosaic of the genetic diversity present in the CC founders—more

than 45 million segregating SNPs [47]. Random outcrossing (of non-sibs) over many DO

generations maintains this diversity and increases recombination frequency, which is critical for

reducing confidence intervals of localized QTL [41]. Compared to classic backcross and

intercross designs, the combinatorial possibilities for allelic variation within and among DO

animals are vast. The genetic diversity in a DO sample may thus mimic diversity in the natural

populations that animal models are meant to illuminate.

We quantify facial shape with 22 three-dimensional landmarks **(Fig 1)** and estimate its

association with founder haplotype variation at 56,885 genetic markers. The dependent

observations in the QTL scans are scores on shape PCs. For each PC explaining more than 1% of

shape variation, we identify statistically significant marker effects using a mixed effects

regression model that simultaneously accounts for kinship among animals [48, 49]. We then

undertake a more detailed analysis of QTL intervals for shape PCs 1-5. We estimate founder

coefficients, test whether the overall gene set is enriched for biological processes relevant to

facial development and perform association mapping to highlight a list of potential causal

genes. In order to visualize and quantify the magnitude of marker effects on multivariate facial

shape (as opposed to univariate PC scores), for each QTL peak, we fit an additional mixed model

that accommodates high-dimensional dependent observations [50].

127
128   **Fig 1. Landmark map.**

Results and Discussion

QTL and Shape Variable Heritability

We used a mixed effects regression model to quantify GP associations between 56,885

genetic markers and each of the 20 shape PCs explaining >1% of the symmetric component of

facial shape variation **(S1 Fig)**. Evidence for a QTL was evaluated by comparing logs odds (LOD)

scores (likelihood ratios comparing the magnitude of marker model residuals to the residuals of

a null model that excludes the marker effect) [51] at each marker to a distribution of maximum

LOD scores obtained from quantifying marker effects over 1000 random permutations of GP

assignments.

While comparisons to prior studies are complicated by design features unique to each,

the design employed here appears to have conferred benefits in terms of QTL identification and

localization, the primary objectives of QTL analysis [41]. From the 20 genome scans, 37 QTL

surpassed our threshold for statistical significance (95% quantile of the maximum LOD score

distribution). Median support interval width was 3.5 Mb (mean 7.1 Mb) using a 1.5 LOD-drop

threshold to define support interval boundaries. Comparable prior studies identified 17-30 QTL

for the skull, with a lowest median QTL interval of 7.8 Mb (using a less-demanding 1.0 LOD-drop

threshold) [32, 33].

Fig 2 plots the support intervals for these QTL (**see also S1 Table**). The QTL span 15 of 19

autosomes as well as the X-chromosome. On chromosomes 7, 14, 15, and 17, QTL from

multiple PCs have overlapping associations with the same genomic region.

Fig 2. Mouse genome physical map with QTL support intervals. Intervals on the same
chromosome are color coded to match the color of the PC for which the QTL was
identified (listed below chromosome). The dashed line in the middle of chromosome 2
represents a region for which genetic effects were not evaluated due to the potential
effects of transmission distortion on QTL identification (see Materials & Methods for
further explanation).

For each PC, the random effect heritability estimate quantifies the proportion of

phenotypic variance attributable to overall genetic similarity at the founder level [49]. We

excluded from the genetic similarity (relatedness) calculation all markers on the same

chromosome as the QTL marker of interest. This leave-one-chromosome-out (LOCO) strategy

was chosen in order to avoid the loss of power that can arise when relatedness is computed

from the full complement of genotyped markers [52-54]. The LOCO approach produces a

unique heritability estimate for each chromosome on each PC. **Table 1** reports the mean

heritability for each PC, as well as the variance in these heritabilities (averaged over

chromosomes). For most PCs, heritability was moderately high (0.42-0.57), with a few low

heritability outliers. These values are very similar to those Pallares and colleagues estimated for

skull shape PCs in an outbred mouse sample [32], and lie mostly in the middle of the heritability

range for human facial shape (~30-70%) [36, 55].

Table 1. Heritabilities by PC.

	Mean h^2	Variance
PC1	0.5407	0.00012
PC2	0.4389	0.00013
PC3	0.5418	0.00022
PC4	0.3532	0.00022
PC5	0.4627	0.00012
PC6	0.5705	0.00015
PC7	0.4785	0.00014
PC8	0.4604	0.00015
PC9	0.5411	0.00032
PC10	0.5420	0.00017
PC11	0.2439	0.00009
PC12	0.4226	0.00014
PC13	0.5309	0.00013
PC14	0.4364	0.00020
PC15	0.4849	0.00005
PC16	0.4456	0.00010
PC17	0.4658	0.00020
PC18	0.4726	0.00019
PC19	0.4533	0.00006
PC20	0.5040	0.00022

171
172   Mean and variance of heritability for PC 1-20 shape traits. The full set of heritability
173   estimates is provided in **S2 Table**.
174
Based on signal-to-noise evaluation of shape PC eigenvalue decay [56], we limited

further analysis to the first five PCs (effectively four PCs, as no marker exceeded the

permutation significance threshold for the PC 5 shape axis). Each plot in **Fig 3** depicts the LOD

score profile (lower panel) and CC founder haplotype coefficients (upper panel) for a

chromosome harboring one or more QTL for a PC 1-4 shape variable. Shaded rectangular

regions delimit the 1.5-LOD drop support interval boundaries for the QTL. Founder coefficients

were estimated using best linear unbiased predictors [57, 58]. This helps to highlight QTL

regions by imposing a degree of shrinkage on the coefficients chromosome-wide [48].

Fig 3. BLUP effects and LOD panels for all QTL, PCs 1-4. QTL effect colors correspond to CC founders.

We also mapped GP associations for vectors capturing covariation between shape and

two measures of size: centroid size (CS) and body mass. Shape change associated CS and body

mass variation were closely correlated with the primary axis of shape variation in our sample

190 (r_{PC1-CS}: 0.985, $r_{PC1-Mass}$: 0.875, $r_{CS-Mass}$: 0.927). These correlations carry forward to the QTL

191 profiles, as well. We identify two QTL each for CS and body mass allometry, one of which, on

192 chromosome 6, is common to the two allometry vectors. Together, the three QTL—on

193 chromosomes 1, 6, and 7—completely overlap with the PC 1 shape QTL (**Fig 4**).

194
195 **Fig 4.** LOD plots for chromosomes containing allometry (CS and body mass) and PC 1
196 QTL. The permutation significance thresholds for all three phenotypic variables are
197 extremely similar. Only one is plotted.
198
The allometry results suggest a developmental model in which the determinants of

different size measures have partially correlated effects on cranial shape [59]. The near

coincidence of shape PC 1 with our explicitly allometric variables has been observed in many

geometric and classical morphometric investigations, and there is a long history of treating PC 1

as an allometry variable in morphometric analysis [59-64]. With this in mind, though primarily

for economy, we regard PC 1 QTL as allometry QTL in the remainder of this manuscript.

The mouse subspecies that comprise the CC are estimated to have diverged several

hundred thousand years ago [65]. In many DO studies, subspecific divergence in one or more

wild-derived CC lines figures prominently in the founder genetic effects contrasts that underlie

QTL (e.g., [66-69]). Here, the same pattern is evident for most of the PC 1-4 QTL. In six of the

eight QTL (or seven of nine, if the two peaks of the PC 2, chromosome 14 QTL are considered

distinct), the coefficient for a non-*domesticus* wild-derived strain (CAST or PWK) sits at one

extreme of the founder coefficient distribution. Only at the upstream LOD peak of the PC 2,

chromosome 14 QTL is the founder coefficient contrast clearly between *M.m. domesticus*

inbred lines. The prevalence of QTL substantially driven by wild-derived alleles highlights the

value of subspecific genetic diversity in the DO population.

215   The broad region of high LOD scores for PC 1 on chromosome 7 merits individual

attention. This QTL support interval essentially delimits the mouse homolog boundaries of the

human Prader-Willi/Angelman syndromes domain [70]. The founder coefficients point to a

strong subspecies signal, with PWK (*M.m. musculus*) highly differentiated from wild-derived

WSB and the inbred *M.m. domesticus* lines, and CAST (*M.m. castaneus*) intermediate. Though

the QTL interval contains fewer than 40 genes, nine are known to be imprinted [70-74]. Several

more have mixed evidence for imprinting [73], and Prader-Willi and Angelman are parent-of-

origin dependent neurological disorders [75]. In addition, more than a dozen genes in the

interval are expressed in the developing face [40, 76-81]. Our mapping strategy is not

specifically designed to quantify imprinted gene QTL [82-84]. However, closer dissection of this

interval's contribution to facial shape variation, accounting for allelic parent of origin effects,

may be warranted in the future. The characteristic facial phenotype for Prader-Willi includes a

relatively long skull with a narrow upper face [85]. In this regard, the shape effects associated

with the peak marker in this interval, just downstream of the primary Prader-Willi genes, are

suggestive, as they contrast a shorter, wider face with a longer, narrower one (see **Figs 5 and 6**,

below).

Marker Effects on Shape

At the LOD peak marker of each PC 1-4 QTL, we collapsed the 8-state founder

probabilities to 2-state SNP probabilities in order to estimate the shape effect of allelic

substitution at that marker. To estimate effects on total shape (instead of univariate shape

PCs), we employed a Bayesian mixed effects model for highly multivariate dependent

observations [50, 86]. The results are plotted in **Fig 5**. In each figure, QTL effects are magnified

by the scaling factor required to cause the QTL to account for 10% of phenotypic variance.

These effects are shown as displacement vectors emanating from the mean configuration

wireframe. The magnification ensures that even when the effect at a landmark is small, its

direction and relative size (in comparison to effects at other landmarks) is apparent. We further

exploited the Bayesian posterior to plot 80% confidence ellipsoids for the magnified

displacements. The actual proportion of variance explained by the SNP, without magnification,

is noted in the figure (and included in **Table 3**).

Chromosome 1
PC 1 (0.21%)

Chromosome 6
PC 1 (0.21%)

Chromosome 7
PC 1 (0.26%)

Chromosome 6
PC 2 (0.2%)

Chromosome 14
PC 2 (0.25%)

246 **Fig 5.** Superior (left), anterior (middle), and lateral (right) views of the (magnified) effect
247 of allele substitution at the genome scan peak marker. Displacement vectors from
248 wireframe to center of ellipsoids depict the mean effect. Ellipsoids enclose the
249 approximate 80% posterior confidence boundaries.
250

Biological Process Enrichment

We used the human annotations in the online Molecular Signatures Database [87] to

query biological processes relevant to craniofacial variation are overrepresented in the PC 1-4

QTL intervals. Collectively, the intervals contain 243 named, protein coding genes (**S1 File**), 232

of which are annotated in the Entrez Gene library utilized by the Molecular Signatures

Database. **Table 2** provides GO biological process enrichment highlights for these intervals,

including FDR *q*-values adjusted for multiple testing (see **S2 File** for the top 100 *q*-value

processes). Growth, morphogenesis, and skeletal system development are prominent biological

process annotations for the facial shape QTL. The enrichment results suggest that no enriched

biological process is uniquely associated with shape variation on a single PC. Rather, with the

exception of bone development, each process in **Table 2** includes annotations to each PC.

Table 2. Gene ontology enrichment for selected biological processes.

Biological process	FDR *q*-value	GO annotated genes
Animal organ morphogenesis	1.42 e-3	**PC1:** *Aldh1a3, Chsy1, Nab1, Stat1, Trpm1.* **PC2:** *Col1a2, Dlg5, Gngt1, Hesx1, Tnnc1, Wnt5a, Zmiz1 .* **PC3:** *Gdf7, Matn3, Osr1, Sdc1.* **PC4:** *Ccdc66, Mdfi, Ptk7, Runx2, Srf, Vegfa .*
Growth	2.25 e-3	**PC1:** *Apba2, Mstn, Ndn, Nsmce3, Ube3a .* **PC2:** *Bap1, Colq, Il17rb, Mustn1, Sema3g, Tkt, Wnt5a, Zmiz1.* **PC3:** *Gdf7, Matn3, Osr1 .* **PC4:** *Dnph1, Mymx, Ptk7, Srf, Taf8, Vegfa, Hsp90ab1.*
Skeletal system development	1.29 e-2	**PC1:** *Chsy1, Nab1, Foxp1, Lrrk1.* **PC2:** *Col1a2, Mustn1, Wnt5a.* **PC3:** *Matn3, Osr1, Runx2.* **PC4:** *Mdfi, Srf.*
Sensory organ development	1.62 e-2	**PC1:** *Aldh1a3, Gabra5, Gabrb3, Trpm1.* **PC2:** *Ccdc66, Gngt1, Hesx1, Wnt5a.* **PC3:** *Osr1.* **PC4:** *Prph2, Ptk7, Srf, Vegfa.*
Cartilage development	2.4 e-2	**PC1:** *Chsy1.* **PC2:** *Mustn1, Wnt5a .* **PC3:** *Matn3, Osr1.* **PC4:** *Runx2, Srf.*
Bone development	2.54 e-2	**PC1:** *Chsy1, Nab1, Lrrk1, Foxp1.* **PC3:** *Matn3.* **PC4:** *Runx2, Srf .*

263
To consider annotations specific to allometric variation, we queried biological process

enrichment for PC 1 QTL alone (**S3 File**). Collectively, the three allometry QTL intervals were

significantly enriched for genes implicated in bone development (FDR *q* = 0.025). Each of the

intervals included at least one such gene: *Nab1* (chromosome 1) [88], *Foxp1* (chromosome 6)

[89, 90], and *Lrrk1* and *Chsy1* (chromosome 7) [91-93]. The chromosome 1 interval also includes

Myostatin (*Mstn, Gdf8*), a negative regulator of muscle growth. Inactivation of *Mstn* has been

shown to greatly increase facial muscle mass, with corresponding size decreases and shape

changes throughout the bony face and vault [94, 95].

Association Mapping

Finally, to develop a finer-grained impression of potential causal loci, we performed SNP

association mapping within each QTL interval. We imputed a dense map of SNP probabilities to

the sample by inferring their bi-allelic (major/minor) probabilities based on (a) founder

probabilities at flanking, genotyped markers and (b) major/minor strain distribution patterns in

the CC founder variants **(Fig 6)**.

Fig 6. Association mapping results for PC 1-4 QTL intervals. Top panels: LOD scores for all
CC SNP variants in the interval. Magenta points are locations where inferred bi-allelic
SNP probabilities (based on strain distribution patterns in the CC) exceed the 95% LOD
threshold. To control file size, low LOD scores points have been omitted from most
plots. Middle panel: location of genotyped markers in the DO (small black diamonds),
highlighting LOD peak locations for the founder genome scan (large black diamond) and
SNP association scan (large magenta diamond). Bottom panel: named, protein coding
genes for intervals. For legibility, we show the portion of the PC 2, Chromosome 14 map
that contains both peak LOD scores and all SNPs exceeding the LOD threshold. For all
plots, the complete gene list is provided in **S1 File**.

We then queried whether any genes within 0.25 Mb of the SNP or genome scan LOD

peaks are known to be associated with craniofacial development or phenotypes **(Table 3)**. As

with human facial shape GWAS [42], LOD peaks tended to be located in non-coding regions.

While the focal genes we identified all have known roles in skeletal or craniofacial

development, they also tend to be developmentally ubiquitous. *Stat1* (PC1, chr. 1) is implicated

in bone development and remodeling throughout life [96-98]. Overexpression of *Foxp1* (PC1,

chr. 6) can attenuate osteoblast differentiation, chondrocyte hypertrophy, osteoclastogenesis,

and bone resorption, while *Foxp1* deficiency can accelerate ossification [89, 90]. In humans,

some *Foxp1* mutations produce a facial phenotype that includes broad forehead and short nose

[99]. Chondrogenesis is generally disrupted in *Chsy1* null mouse mutants (PC1, chr. 7), and in

humans, *Chsy1* mutations can manifest in micrognathia and short stature [92, 100]. *Kat6b* (PC2,

Chr. 14) activates the transcription factor *Runx2*, which is often referred to as a master

regulator of bone formation [101-103]. For the PC3 QTL at 8.23 – 9.84 Mb on chromosome 12,

the founder and SNP LOD peaks reside at opposite ends of the QTL interval, but each is very

close to an important craniofacial development gene: *Gdf7* (*Bmp12*) contributes to patterning

of the jaw joint and is expressed in the developing incisor pulp and periodontium [104-106];

Osr1 is broadly expressed in the embryonic and developing skull, with overexpression resulting

in intramembranous ossification deficiencies, increased chondrocyte differentiation, and

incomplete suture closure, while under-expression may contribute to cleft palate phenotype

[107-109]. *Osr1* has also previously been identified as a candidate in human facial shape GWAS

[23]. Finally, the SNP LOD peak for PC4, chromosome 17 implicates *Vegfa*, which is critical to

endochondral and intramembranous ossification through its role in angiogenesis [110]. A

distinct SNP LOD peak of nearly equal magnitude is located approximately 1 Mb upstream (at

rs47194347), very close to *Supt3* (human ortholog *Supt3h*) and *Runx2*. In addition to the central

role *Runx2* has in bone formation and development [102], both genes are human facial shape

GWAS candidates [15, 22].

Table 3. Candidate genes.

PC	Chr	Founder Peak (Mb)	Founder ID	Founder type	SNP Peak (Mb)	SNP ID	SNP_type	SNP % Var	Candidates
PC1	1	51.926616	rs48415864	intergenic	51.926623	rs254750887	intergenic	0.21	*STAT1*
PC1	6	99.637221	rs226005706	intron	99.900271	rs31234872	intergenic	0.21	*FOXP1*
PC1	7	65.970847	rs33905059	intron	65.876674	rs32405988	intron	0.26	*CHSY1*
PC2	6	5.704731	rs49620919	intergenic	5.502445	rs32485785	intergenic	0.2	NA
PC2	14	22.058402	rs30739483	intron	22.182166	rs30740474	intron	0.25	*KAT6B*
PC3	12	8.44203	rs48833407	intergenic	9.587123	rs29212735	intergenic	0.18	*OSR1, GDF7*
PC3	12	16.286802	rs49949506	intergenic	17.125642	SV 17121898: 17129386	structural (deletion)	0.38	NA
PC4	17	45.639772	rs107675960	intron	46.106749	rs108721614	upstream gene variant	0.3	*VEGFA* *[RUNX2, SUPT3]*

319 Candidate genes located within 0.25 Mb of founder or SNP QTL peaks. SNP % Var gives
320 the percentage of variation explained by substitution of an additional reference allele at
321 the founder (genome scan) peak.
322

While GP maps for traits influenced by many genes may be quite intricate and layered,

additive effect QTL analyses remain an indispensable means to identify loci that contribute to

phenotypic variation. Our investigation advances understanding of the genetics of facial

skeletal diversity in several ways. The 37 QTL identified herein substantially expand the small

but growing facial shape QTL library. Compared to similar prior studies, the number of QTL is

relatively large and their confidence intervals relatively narrow. These results affirm the value

of larger samples, sophisticated animal models like the outcrossed population utilized here, and

high-resolution genotyping for identification and localization of QTL. Synthesizing our results

with prior insights from developmental biology and medical genetics provides a richer picture

of the sources of facial skeletal diversity. We found a relative abundance of skeletal and

craniofacial development genes in QTL intervals for the largest shape PCs. Moreover, these

genes tend to be located very close to QTL peaks. Thus, our results suggest that genes essential

to building a face are often also the genes that cause faces to vary. Finally, QTL effect sizes were

small. In summary, our results are consistent with a model of facial diversity that is influenced

by key genes in skeletal and facial development and, simultaneously, highly polygenic.

Materials and methods

Diversity Outbred mouse population

The DO (Jackson Laboratory, www.jax.org) are derived from the eight inbred founder lines that contribute to the CC [43, 44], which include three mouse subspecies, and both wild-derived and laboratory strains [46]. Continued random outcrossing of the DO has produced a population with relatively high genetic diversity (~45 million SNPs segregating in the CC founding populations [43]) and increasingly fine mapping resolution in advancing generations.

In combination, these features increase the potential to identify multiple QTL for complex traits [31]. On average, the genomes of the eight CC founders should be equally represented in the DNA of any DO animal. Sequenced genomes for the CC make it possible to characterize and interpret genetic variants in the DO according to 8-state founder identity and as bi-allelic (major/minor allele) SNP variation. In addition, statistical packages designed for analysis of the DO offer features to impute variants at each typed SNP in the CC founders, using the coarser DO marker map as a scaffold to infer a high-density SNP map.

Computing

All analysis was conducted in R [111]. Customized data processing and statistical analysis functions for mapping experiments in DO samples were originally published in the DOQTL package [49]. These routines are now incorporated into the qlt2 package [48], with additional processing and genotype diagnostic tools provided at https://kbroman.org/qtl2/.

Images, landmark data

Our DO sample consists of *n*=1147 male and female adult mice (after exclusion of some records; see *Genotypes*, below) from seven DO generations (**Table 4**). No animals were

sacrificed for this study. Specimens were raised in several labs for unrelated experiments under

approval and conduced in accordance with the guidelines set forth by the Institutional Animal

Care and Use Committee in accordance with protocols #11-299 (University of North Carolina

Chapel Hill), #99066 and #15026 (Jackson Labs), and #08-0150-3 (Scripps Research Institute).

Receipt of animal carcasses and micro-CT (µCT) imaging of mouse skulls were undertaken in

accordance with approved IACUC protocol #AC18-0026 at the University of Calgary.

Table 4. DO sample composition.

	DO generation						
	9	**10**	**15**	**19**	**21**	**23**	**27**
Female	91	221	80	55	80	97	75
Male	0	144	0	58	77	99	70
MUGA Array	Mega	Mega	Mega	Giga	Giga	Giga	Giga

367
368   Overview of the DO sample by generation, sex, and genotyping array.
369
We obtained µCT images of the mouse skulls in the 3D Morphometrics Centre at the

University of Calgary on a Scanco vivaCT40 scanner (Scanco Medical, Brüttisellen, Switzerland)

at 0.035 mm voxel dimensions at 55kV and 72-145 µA. On each skull, one of us (WL) collected

54 3D landmark coordinates from minimum threshold-defined bone surface with ANALYZE 3D

(www.mayo.edu/bir/). Here, we analyzed the 22-landmark subset of the full configuration—a

66 coordinate dimension vector—that captures the shape of the face (**Fig 1**).

Shape data

Landmark configurations were symmetrically superimposed by Generalized Procrustes

Analysis (GPA) [112] using the Morpho package [113]. GPA is a multi-step procedure that

removes location and centroid size differences between configurations, then (iteratively)

rotates each configuration to minimize its squared distance from the sample mean. We chose a

superimposition that decomposes shape variation into its symmetric and asymmetric

components [114, 115], and stored the symmetric configurations for analysis. Working with

symmetric, superimposed data reduced morphological degrees of freedom from 66 landmark

coordinate dimensions to 30 shape dimensions [115].

Genotypes

For all animals, DNA was extracted from ear or tail clippings. Genotyping was performed

by the Geneseek operations of Neogen Corp. (Lincoln, NE). Mice from generations 9, 10, and 15

were genotyped using the MegaMUGA genotyping array (77,808 markers); mice from

generations 19, 21, 23, and 27 were genotyped using the larger GigaMUGA array (143,259

markers) [116]. The two arrays share 58,907 markers in common. We restricted the genotype

data to these shared SNPs [116].

In addition, we excluded 2,022 markers from the center of chromosome 2 (40 - 140

Mb). WSB alleles are highly overrepresented in this genomic region in earlier DO generations.

This overrepresentation has been attributed to female meiotic drive favoring WSB at the *R2d2*

locus [117]. Between DO generations 12 and 21, Jackson Labs systematically purged the meiotic

drive locus and linked effects from the DO [45]. Our samples closely track this history, with WSB

allele frequencies in the middle of chromosome 2 exceeding 60% in generation 9 and falling to

0% by generation 23 (**S2 Fig**). If unaddressed, this directional shift in allele frequencies can

confound statistical analysis of genotype-phenotype associations [118] with potentially false

positive associations between WSB chromosome 2 haplotypes and directional changes in

phenotype over DO generations. After removing this genomic region from consideration, the

final SNP dataset consists of 56,885 markers common to the MegaMUGA and GigaMUGA

arrays, with a mean spacing of 48 kb.

The qtl2 package estimates genotype probabilities, including imputation of sporadically

missing genotypes, with a multipoint Hidden Markov Model (HMM) on the observed marker

genotypes. From these initial genotype probabilities, we derived three probabilistic

representations of genetic state for each observation at each marker: (1) for genome scans,

founder probabilities, a vector encoding the probability of genetic contribution from each of

the eight CC founders to the diplotype state; (2) for association mapping, SNP probabilities,

collapsing the eight founder probabilities to a two-state, major/minor allele strain distribution;

and (3) for mapping multivariate shape effects, additive SNP dosage, twice the estimated major

SNP probability (maximum additive SNP dosage = 2).

We identified problematic genetic samples using the DO genotype diagnostics tools

available at https://kbroman.org/qtl2/ [119]. A total of 11 observations were excluded due to

excessive missing marker data, which we defined as no marker information at >7.5% of

markers. Three probable duplicate sets of genetic data (>99% of markers identical) were

identified. Because the correct genotype-phenotype matches could not be established by other

evidence, all six animals were removed from the sample. Finally, we excluded seven female

samples with low X chromosome array intensity values. These samples are potentially XO [45],

which can affect facial form and body size [120, 121]. The exclusions described above resulted

in a final, retained sample of n = 1147 animals.

Genome scan and association mapping

We mapped QTL for facial shape and facial shape allometry. To identify QTL, we

implemented the linear mixed effects regression model in qtl2. The qtl2 genome scan is

designed to estimate marker effects on one or more univariate observations, but not

covariance among those dimensions. We therefore decomposed the superimposed coordinates

into PCs, treating the specimen scores on each PC as measured traits. For allometry, we treated

common allometric component scores [61] as measured traits. Prior to deriving PCs and

allometry vectors, we centered each observation on its generation mean in order to focus on

shared, within-generation covariance patterns [122, p. 295].

At each marker, we fit a mixed model with fixed effect coefficients for sex, age-at-

sacrifice, and the additive effect of each CC founder haplotype, a random effect of kinship, and

an error term [49]. Evidence for a QTL was evaluated based on marker LOD scores, likelihood

ratios comparing the residuals of the marker model to a null model excluding the marker's

effect [51]. We consider the evidence for a marker-phenotype association to be statistically

significant where the LOD score exceeds the 95% quantile of genome-wide maximum LOD

scores computed from 1000 random permutations of genotype-phenotype associations [123].

We use a 1.5 LOD-drop rule to define the boundaries of QTL support intervals [124], and allow

for multiple QTL peaks on a single chromosome.

The random effect of kinship depends upon pairwise proportions of shared alleles

among observations, encoded in a kinship matrix. Loss of power to detect QTL can arise when

the marker of interest and linked genes are included in the kinship matrix of the null model. To

avoid this problem, we compute kinship matrices using a leave-one-chromosome-out (LOCO)

approach [52-54].

Genome scans, QTL support intervals, and proportions of shape variation attributable to

founder-level kinship (heritability) were estimated for all PCs explaining more than 1% of

symmetric shape variation. However, we limit more in-depth analysis to the five PCs with

reasonably large signal-to-noise ratios (**S2 Fig**). This was assessed using the modified "noise

floor" detection strategy developed by Marroig and colleagues in [56].

In addition, within the PC 1-5 QTL intervals, we used a form of merge analysis to

perform genome-wide association mapping [49, 125, 126]. For each CC founder variant located

between a flanking pair of genotyped markers, the vector of founder allele probabilities is

assumed to be the average of the founder probabilities at those flanking markers. The bi-allelic

SNP probabilities were computed from these founder probabilities and the major/minor allele

strain distribution pattern in the CC. Analysis was again via mixed effects regression model, with

the imputed SNP data as genotype predictors.

Enrichment analysis and candidate genes

Modern GWAS and QTL studies often attempt to contextualize QTL and candidate gene

results post hoc via GO enrichment analysis. Given a list of genes near statistically significant

markers, a GO analysis calculates the probability that a random set of genes of the same size

would contain as many genes under a common annotation term, typically after a family-wise

error rate or FDR adjustment for multiple testing. The outcome of enrichment analysis is a list

of GO terms (or in the absence of any enrichment, no list), which provides a means to consider

whether certain biological processes, cellular components, or molecular functions are over-

represented in the list of genes [127].

For protein coding genes in the PC 1-5 QTL support intervals and for allometry QTL, we

used the Molecular Signatures Database [87] online tool and Entrez Gene library [128] to

identify biological process GO term enrichment [129]. The database application implements a

hypergeometric test of statistical significance, quantifying the probability of sampling, without

replacement, as many (*k*) or more genes annotated to a GO term, given the total number of

genes annotated to that term (*K*) and the number of genes in the QTL and Entrez gene sets (*n*

and *N,* respectively). For each ontology term, after adjustment for multiple testing, we required

a FDR *q*-value < 0.05 to conclude that there was substantial evidence for GO biological process

enrichment. We also inspected the association mapping results in the +/- 0.25 Mb region

centered at each LOD peak for genes with convincing evidence of a role in skeletal, craniofacial,

or dental development or disease. For most QTL, LOD peaks for the genome scan and

association mapping results are virtually coincident. Where they differ substantially, we

examine the region around both peaks.

Marker effects on shape

We considered a significant association between a marker and facial shape along a PC

axis to be evidence of some effect on the multivariate (undecomposed) shape coordinate data.

In order to obtain an estimate of the full shape effect, we fit an additional mixed model for the

maximum LOD score marker. Fitting a mixed model for highly multivariate data presents a

significant computational challenge because the number of parameters that must be estimated

for the random effects covariance matrix scales quadratically with the number of traits (here,

465 parameters for 30 free variables). To address this challenge, we estimated mixed model

covariance matrices using a Bayesian sparse factorization approach [50, 86, 130]. After a

1,000,000 iteration burn-in, we generated 1,000,000 realizations from a single Markov chain,

thinning at a rate of 1,000 to store 1,000 posterior samples for inference. Though the time

needed to fit such a model makes it impractical for a genome scan on a dense array, it is a

useful strategy for disentangling kinship from the effect on multivariate outcomes at a small

number of peak markers. For these regressions, the marker effect predictor was twice the

major SNP probability. Thus, the multivariate mixed model we implemented estimates an

average effect of allele substitution, a standard quantity in quantitative genetics [131]. In

addition, for each regression, we estimated the proportion of shape variation explained using a

version of the coefficient of determination (R^2) that has been adapted for mixed model

parameters [132].

When plotting shape effects, we magnify the allele substitution coefficients by the

scaling factor needed to cause the QTL to explain 5% of phenotypic variance. This magnification

is applied both to the mean effect, in order to depict the mean displacement vector, and to the

posterior distribution of effects, in order to depict confidence ellipsoids. Ellipsoids were

estimated with the 3D ellipse function in the R package rgl [133]. While it is more typical to use

a 95% support boundary to evaluate effect uncertainty, with shape data, morphological

contrasts are best understood over the configuration as a whole [134]. The 80% ellipsoids are

meant to encourage this perspective.

Acknowledgements

Thanks to Xiang Zhao for help with preparation of samples and David Eby for helpful

intervention with the online Molecular Signatures Database. We thank Kunjie Hua, Liyang Zhao

and Kuo-Chen Jung for assistance with UNC mouse experiments. Charles Farber is gratefully

acknowledged for assistance with processing of UNC mouse carcasses. All UNC mouse

procedures were approved and guided by the appropriate institutional animal ethics

committees, including the University of North Carolina at Chapel Hill.

REFERENCES

1.   Hu D, Marcucio RS. A SHH-responsive signaling center in the forebrain regulates
craniofacial morphogenesis via the facial ectoderm. Development. 2009;136(1):107-16. Epub
2008/11/26. doi: 10.1242/dev.026583. PubMed PMID: 19036802; PubMed Central PMCID:
PMCPMC2685963.
2.   Hallgrímsson B, Jamniczky H, Young NM, Rolian C, Parsons TE, Boughner JC, et al.
Deciphering the palimpsest: studying the relationship between morphological integration and
phenotypic covariation. Evolutionary Biology. 2009;36(4):355-76.
3.   Hallgrimsson B, Mio W, Marcucio RS, Spritz R. Let's face it--complex traits are just not
that simple. PLoS Genet. 2014;10(11):e1004724. Epub 2014/11/06. doi:
10.1371/journal.pgen.1004724. PubMed PMID: 25375250; PubMed Central PMCID:
PMCPMC4222688.
4.   Varón-González C, Pallares LF, Debat V, Navarro N. Mouse Skull Mean Shape and Shape
Robustness Rely on Different Genetic Architectures and Different Loci. Front Genet. 2019;10:64.
Epub 2019/02/12. doi: 10.3389/fgene.2019.00064. PubMed PMID: 30809244; PubMed Central
PMCID: PMCPMC6379267.
5.   Young NM, Chong HJ, Hu D, Hallgrímsson B, Marcucio RS. Quantitative analyses link
modulation of sonic hedgehog signaling to continuous variation in facial growth and shape.
Development. 2010;137(20):3405-9. Epub 2010/09/08. doi: 10.1242/dev.052340. PubMed
PMID: 20826528; PubMed Central PMCID: PMCPMC2947484.
6.   Green RM, Fish JL, Young NM, Smith FJ, Roberts B, Dolan K, et al. Developmental
nonlinearity drives phenotypic robustness. Nat Commun. 2017;8(1):1970. Epub 2017/12/06.
doi: 10.1038/s41467-017-02037-7. PubMed PMID: 29213092; PubMed Central PMCID:
PMCPMC5719035.
7.   Mitteroecker P, Bookstein F. The Conceptual and Statistical Relationship between
Modularity and Morphological Integration. Systematic Biology. 2007;56(5):818-36. doi:
10.1080/10635150701648029.
8.   Szabo-Rogers HL, Smithers LE, Yakob W, Liu KJ. New directions in craniofacial
morphogenesis. Dev Biol. 2010;341(1):84-94. Epub 2009/11/24. doi:
10.1016/j.ydbio.2009.11.021. PubMed PMID: 19941846.
9.   Francis-West P, Crespo-Enriquez I. Vertebrate Embryo: Craniofacial Development. eLS
[Internet]. 2016:[1-15 pp.].
10.   Falconer DS, Mackay TF. Introduction to quantitative genetics. 4th ed. Essex: Pearson
Education Limited; 2009.
11.   Huang W, Mackay TF. The Genetic Architecture of Quantitative Traits Cannot Be Inferred
from Variance Component Analysis. PLoS Genet. 2016;12(11):e1006421. Epub 2016/11/03. doi:
10.1371/journal.pgen.1006421. PubMed PMID: 27812106; PubMed Central PMCID:
PMCPMC5094750.
12.   Barton NH, Keightley PD. Understanding quantitative genetic variation. Nat Rev Genet.
2002;3(1):11-21. doi: 10.1038/nrg700. PubMed PMID: 11823787.

13.   Varón-González C, Navarro N. Epistasis regulates the developmental stability of the
mouse craniofacial shape. Heredity (Edinb). 2019;122(5):501-12. Epub 2018/09/12. doi:
10.1038/s41437-018-0140-8. PubMed PMID: 30209292; PubMed Central PMCID:
PMCPMC6461946.
14.   Wolf JB, Leamy LJ, Routman EJ, Cheverud JM. Epistatic pleiotropy and the genetic
architecture of covariation within early and late-developing skull trait complexes in mice.
Genetics. 2005;171(2):683-94. Epub 2005/07/14. doi: 10.1534/genetics.104.038885. PubMed
PMID: 16020793; PubMed Central PMCID: PMCPMC1456798.
15.   Claes P, Roosenboom J, White JD, Swigut T, Sero D, Li J, et al. Genome-wide mapping of
global-to-local genetic effects on human facial shape. Nat Genet. 2018;50(3):414-23. Epub
2018/02/19. doi: 10.1038/s41588-018-0057-4. PubMed PMID: 29459680; PubMed Central
PMCID: PMCPMC5937280.
16.   Lee MK, Shaffer JR, Leslie EJ, Orlova E, Carlson JC, Feingold E, et al. Genome-wide
association study of facial morphology reveals novel associations with FREM1 and PARK2. PLoS
One. 2017;12(4):e0176566. Epub 2017/04/25. doi: 10.1371/journal.pone.0176566. PubMed
PMID: 28441456; PubMed Central PMCID: PMCPMC5404842.
17.   Liu F, van der Lijn F, Schurmann C, Zhu G, Chakravarty MM, Hysi PG, et al. A genome-
wide association study identifies five loci influencing facial morphology in Europeans. PLoS
Genet. 2012;8(9):e1002932. Epub 2012/09/13. doi: 10.1371/journal.pgen.1002932. PubMed
PMID: 23028347; PubMed Central PMCID: PMCPMC3441666.
18.   Paternoster L, Zhurov AI, Toma AM, Kemp JP, St Pourcain B, Timpson NJ, et al. Genome-
wide association study of three-dimensional facial morphology identifies a variant in PAX3
associated with nasion position. Am J Hum Genet. 2012;90(3):478-85. Epub 2012/02/16. doi:
10.1016/j.ajhg.2011.12.021. PubMed PMID: 22341974; PubMed Central PMCID:
PMCPMC3309180.
19.   Shaffer JR, Orlova E, Lee MK, Leslie EJ, Raffensperger ZD, Heike CL, et al. Genome-Wide
Association Study Reveals Multiple Loci Influencing Normal Human Facial Morphology. PLoS
Genet. 2016;12(8):e1006149. Epub 2016/08/25. doi: 10.1371/journal.pgen.1006149. PubMed
PMID: 27560520; PubMed Central PMCID: PMCPMC4999139.
20.   Crouch DJM, Winney B, Koppen WP, Christmas WJ, Hutnik K, Day T, et al. Genetics of the
human face: Identification of large-effect single gene variants. Proc Natl Acad Sci U S A.
2018;115(4):E676-E85. Epub 2018/01/04. doi: 10.1073/pnas.1708207114. PubMed PMID:
29301965; PubMed Central PMCID: PMCPMC5789906.
21.   Cole JB, Manyama M, Kimwaga E, Mathayo J, Larson JR, Liberton DK, et al. Genomewide
Association Study of African Children Identifies Association of SCHIP1 and PDE8A with Facial
Size and Shape. PLoS Genet. 2016;12(8):e1006174. Epub 2016/08/25. doi:
10.1371/journal.pgen.1006174. PubMed PMID: 27560698; PubMed Central PMCID:
PMCPMC4999243.
22.   Adhikari K, Fuentes-Guajardo M, Quinto-Sánchez M, Mendoza-Revilla J, Camilo Chacón-
Duque J, Acuña-Alonzo V, et al. A genome-wide association scan implicates DCHS2, RUNX2,
GLI3, PAX1 and EDAR in human facial variation. Nat Commun. 2016;7:11616. Epub 2016/05/19.
doi: 10.1038/ncomms11616. PubMed PMID: 27193062; PubMed Central PMCID:
PMCPMC4874031.

23.   Cha S, Lim JE, Park AY, Do JH, Lee SW, Shin C, et al. Identification of five novel genetic
loci related to facial morphology by genome-wide association studies. BMC Genomics.
2018;19(1):481. Epub 2018/06/19. doi: 10.1186/s12864-018-4865-9. PubMed PMID: 29921221;
PubMed Central PMCID: PMCPMC6008943.
24.   Richmond S, Howe LJ, Lewis S, Stergiakouli E, Zhurov A. Facial Genetics: A Brief
Overview. Front Genet. 2018;9:462. Epub 2018/10/16. doi: 10.3389/fgene.2018.00462.
PubMed PMID: 30386375; PubMed Central PMCID: PMCPMC6198798.
25.   Cole J, Spritz RA. The genetics of facial morphology. eLS [Internet]. 2017:[1-9 pp.].
26.   Cheverud JM, Routman EJ, Irschick DJ. PLEIOTROPIC EFFECTS OF INDIVIDUAL GENE LOCI
ON MANDIBULAR MORPHOLOGY. Evolution. 1997;51(6):2006-16. doi: 10.1111/j.1558-
5646.1997.tb05122.x. PubMed PMID: 28565113.
27.   Kenney-Hunt JP, Wang B, Norgard EA, Fawcett G, Falk D, Pletscher LS, et al. Pleiotropic
patterns of quantitative trait loci for 70 murine skeletal traits. Genetics. 2008;178(4):2275-88.
doi: 10.1534/genetics.107.084434. PubMed PMID: 18430949; PubMed Central PMCID:
PMCPMC2323815.
28.   Pallares LF, Harr B, Turner LM, Tautz D. Use of a natural hybrid zone for genomewide
association mapping of craniofacial traits in the house mouse. Mol Ecol. 2014;23(23):5756-70.
Epub 2014/11/15. doi: 10.1111/mec.12968. PubMed PMID: 25319559.
29.   Klingenberg CP, Leamy LJ. Quantitative genetics of geometric shape in the mouse
mandible. Evolution. 2001;55(11):2342-52. doi: 10.1111/j.0014-3820.2001.tb00747.x.
30.   Klingenberg CP, Leamy LJ, Cheverud JM. Integration and Modularity of Quantitative Trait
Locus Effects on Geometric Shape in the Mouse Mandible. Genetics. 2004;166(4):1909-21. doi:
10.1534/genetics.166.4.1909.
31.   Flint J, Eskin E. Genome-wide association studies in mice. Nat Rev Genet.
2012;13(11):807-17. Epub 2012/10/09. doi: 10.1038/nrg3335. PubMed PMID: 23044826;
PubMed Central PMCID: PMCPMC3625632.
32.   Pallares LF, Carbonetto P, Gopalakrishnan S, Parker CC, Ackert-Bicknell CL, Palmer AA, et
al. Mapping of Craniofacial Traits in Outbred Mice Identifies Major Developmental Genes
Involved in Shape Determination. PLoS Genet. 2015;11(11):e1005607. Epub 2015/11/02. doi:
10.1371/journal.pgen.1005607. PubMed PMID: 26523602; PubMed Central PMCID:
PMCPMC4629907.
33.   Maga AM, Navarro N, Cunningham ML, Cox TC. Quantitative trait loci affecting the 3D
skull shape and size in mouse and prioritization of candidate genes in-silico. Front Physiol.
2015;6:92. Epub 2015/03/26. doi: 10.3389/fphys.2015.00092. PubMed PMID: 25859222;
PubMed Central PMCID: PMCPMC4374467.
34.   Percival CJ, Green R, Roseman CC, Gatti DM, Morgan JL, Murray SA, et al.
Developmental constraint through negative pleiotropy in the zygomatic arch. Evodevo.
2018;9:3. Epub 2018/01/27. doi: 10.1186/s13227-018-0092-3. PubMed PMID: 29423138;
PubMed Central PMCID: PMCPMC5787316.
35.   Gudbjartsson DF, Walters GB, Thorleifsson G, Stefansson H, Halldorsson BV,
Zusmanovich P, et al. Many sequence variants affecting diversity of adult human height. Nat
Genet. 2008;40(5):609-15. Epub 2008/04/06. doi: 10.1038/ng.122. PubMed PMID: 18391951.
36.   Cole J, Manyama M, Larson JR, Liberton DK, Ferrara TM, Riccardi SL, et al. Human Facial
Shape and Size Heritability and Genetic Correlations. Genetics. 2017;205(2):967-78. Epub

2016/12/14. doi: 10.1534/genetics.116.193185. PubMed PMID: 27974501; PubMed Central
PMCID: PMCPMC5289863.
37.   de Los Campos G, Vazquez AI, Hsu S, Lello L. Complex-Trait Prediction in the Era of Big
Data. Trends Genet. 2018;34(10):746-54. Epub 2018/08/20. doi: 10.1016/j.tig.2018.07.004.
PubMed PMID: 30139641; PubMed Central PMCID: PMCPMC6150788.
38.   Wilderman A, VanOudenhove J, Kron J, Noonan JP, Cotney J. High-Resolution
Epigenomic Atlas of Human Embryonic Craniofacial Development. Cell Rep. 2018;23(5):1581-
97. doi: 10.1016/j.celrep.2018.03.129. PubMed PMID: 29719267; PubMed Central PMCID:
PMCPMC5965702.
39.   Hooper JE, Feng W, Li H, Leach SM, Phang T, Siska C, et al. Systems biology of facial
development: contributions of ectoderm and mesenchyme. Dev Biol. 2017;426(1):97-114. Epub
2017/03/29. doi: 10.1016/j.ydbio.2017.03.025. PubMed PMID: 28363736; PubMed Central
PMCID: PMCPMC5530582.
40.   Diez-Roux G, Banfi S, Sultan M, Geffers L, Anand S, Rozado D, et al. A high-resolution
anatomical atlas of the transcriptome in the mouse embryo. PLoS Biol. 2011;9(1):e1000582.
Epub 2011/01/18. doi: 10.1371/journal.pbio.1000582. PubMed PMID: 21267068; PubMed
Central PMCID: PMCPMC3022534.
41.   Mackay TFC, Stone EA, Ayroles JF. The genetics of quantitative traits: challenges and
prospects. Nat Rev Genet. 2009;10(8):565-77. doi:
http://www.nature.com/nrg/journal/v10/n8/suppinfo/nrg2612_S1.html.
42.   Weinberg SM, Roosenboom J, Shaffer JR, Shriver MD, Wysocka J, Claes P. Hunting for
genes that shape human faces: Initial successes and challenges for the future. Orthod Craniofac
Res. 2019;22 Suppl 1:207-12. doi: 10.1111/ocr.12268. PubMed PMID: 31074157; PubMed
Central PMCID: PMCPMC6550302.
43.   Svenson KL, Gatti DM, Valdar W, Welsh CE, Cheng R, Chesler EJ, et al. High-resolution
genetic mapping using the Mouse Diversity outbred population. Genetics. 2012;190(2):437-47.
doi: 10.1534/genetics.111.132597. PubMed PMID: 22345611; PubMed Central PMCID:
PMCPMC3276626.
44.   Churchill GA, Gatti DM, Munger SC, Svenson KL. The Diversity Outbred mouse
population. Mamm Genome. 2012;23(9-10):713-8. Epub 2012/08/15. doi: 10.1007/s00335-012-
9414-2. PubMed PMID: 22892839; PubMed Central PMCID: PMCPMC3524832.
45.   Chesler EJ, Gatti DM, Morgan AP, Strobel M, Trepanier L, Oberbeck D, et al. Diversity
Outbred Mice at 21: Maintaining Allelic Variation in the Face of Selection. G3 (Bethesda).
2016;6(12):3893-902. Epub 2016/12/07. doi: 10.1534/g3.116.035527. PubMed PMID:
27694113; PubMed Central PMCID: PMCPMC5144960.
46.   Yang H, Wang JR, Didion JP, Buus RJ, Bell TA, Welsh CE, et al. Subspecific origin and
haplotype diversity in the laboratory mouse. Nat Genet. 2011;43(7):648-55. Epub 2011/05/29.
doi: 10.1038/ng.847. PubMed PMID: 21623374; PubMed Central PMCID: PMCPMC3125408.
47.   Consortium CC. The genome architecture of the Collaborative Cross mouse genetic
reference population. Genetics. 2012;190(2):389-401. doi: 10.1534/genetics.111.132639.
PubMed PMID: 22345608; PubMed Central PMCID: PMCPMC3276630.
48.   Broman KW, Gatti DM, Simecek P, Furlotte NA, Prins P, Sen Ś, et al. R/qtl2: Software for
Mapping Quantitative Trait Loci with High-Dimensional Data and Multiparent Populations.

Genetics. 2019;211(2):495-502. Epub 2018/12/27. doi: 10.1534/genetics.118.301595. PubMed
PMID: 30591514; PubMed Central PMCID: PMCPMC6366910.
49.   Gatti DM, Svenson KL, Shabalin A, Wu LY, Valdar W, Simecek P, et al. Quantitative trait
locus mapping methods for diversity outbred mice. G3 (Bethesda). 2014;4(9):1623-33. Epub
2014/09/18. doi: 10.1534/g3.114.013748. PubMed PMID: 25237114; PubMed Central PMCID:
PMCPMC4169154.
50.   Runcie DE, Mukherjee S. Dissecting High-Dimensional Phenotypes with Bayesian Sparse
Factor Analysis of Genetic Covariance Matrices. Genetics. 2013;194(3):753-67. doi:
10.1534/genetics.113.151217.
51.   Lander ES, Botstein D. Mapping mendelian factors underlying quantitative traits using
RFLP linkage maps. Genetics. 1989;121(1):185-99. PubMed PMID: 2563713; PubMed Central
PMCID: PMCPMC1203601.
52.   Listgarten J, Lippert C, Kadie CM, Davidson RI, Eskin E, Heckerman D. Improved linear
mixed models for genome-wide association studies. Nat Methods. 2012;9(6):525-6. Epub
2012/05/30. doi: 10.1038/nmeth.2037. PubMed PMID: 22669648; PubMed Central PMCID:
PMCPMC3597090.
53.   Cheng R, Parker CC, Abney M, Palmer AA. Practical considerations regarding the use of
genotype and pedigree data to model relatedness in the context of genome-wide association
studies. G3 (Bethesda). 2013;3(10):1861-7. Epub 2013/10/03. doi: 10.1534/g3.113.007948.
PubMed PMID: 23979941; PubMed Central PMCID: PMCPMC3789811.
54.   Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL. Advantages and pitfalls in the
application of mixed-model association methods. Nat Genet. 2014;46(2):100-6. doi:
10.1038/ng.2876. PubMed PMID: 24473328; PubMed Central PMCID: PMCPMC3989144.
55.   Tsagkrasoulis D, Hysi P, Spector T, Montana G. Heritability maps of human face
morphology through large-scale automated three-dimensional phenotyping. Sci Rep.
2017;7:45885. Epub 2017/04/19. doi: 10.1038/srep45885. PubMed PMID: 28422179; PubMed
Central PMCID: PMCPMC5395823.
56.   Marroig G, Melo DA, Garcia G. Modularity, noise, and natural selection. Evolution.
2012;66(5):1506-24.
57.   Mrode RA. Linear models for the prediction of animal breeding values. 3rd ed.
Oxfordshire, UK: CABI; 2014.
58.   Robinson GK. That BLUP is a good thing: the estimation of random effects. Statistical
Science. 1991;6(1):15-51. doi: 10.1214/ss/1177011933.
59.   Hallgrimsson B, Katz DC, Aponte JD, Larson JE, Devine J, Gonzalez P, et al. Integration
and the developmental genetics of allometry. Integrative and Comparative Biology. 2019, in
press.
60.   Jolicoeur P, Mosimann JE. Size and shape variation in the painted turtle. A principal
component analysis. Growth. 1960;24:339-54. PubMed PMID: 13790416.
61.   Mitteroecker P, Gunz P, Bernhard M, Schaefer K, Bookstein F. Comparison of cranial
ontogenetic trajectories among great apes and humans. Journal of Human Evolution.
2004;46(6):679-98. doi: 10.1016/j.jhevol.2004.03.006.
62.   Mosimann JE. Size allometry: Size and shape variables with characterizations of the
lognormal and generalized gamma distributions. Journal of the American Statistical Association.
1970;65:930-45.

63.   Cheverud JM. Relationships among ontogenetic, static, and evolutionary allometry. Am J
Phys Anthropol. 1982;59(2):139-49. doi: 10.1002/ajpa.1330590204. PubMed PMID: 7149015.
64.   Larson JR, Manyama MF, Cole JB, Gonzalez PN, Percival CJ, Liberton DK, et al. Body size
and allometric variation in facial shape in children. Am J Phys Anthropol. 2018;165(2):327-42.
Epub 2017/11/27. doi: 10.1002/ajpa.23356. PubMed PMID: 29178597; PubMed Central PMCID:
PMCPMC5777898.
65.   Goios A, Pereira L, Bogue M, Macaulay V, Amorim A. mtDNA phylogeny and evolution of
laboratory mouse strains. Genome Res. 2007;17(3):293-8. Epub 2007/02/06. doi:
10.1101/gr.5941007. PubMed PMID: 17284675; PubMed Central PMCID: PMCPMC1800920.
66.   Logan RW, Robledo RF, Recla JM, Philip VM, Bubier JA, Jay JJ, et al. High-precision
genetic mapping of behavioral traits in the diversity outbred mouse population. Genes Brain
Behav. 2013;12(4):424-37. Epub 2013/03/20. doi: 10.1111/gbb.12029. PubMed PMID:
23433259; PubMed Central PMCID: PMCPMC3709837.
67.   French JE, Gatti DM, Morgan DL, Kissling GE, Shockley KR, Knudsen GA, et al. Diversity
Outbred Mice Identify Population-Based Exposure Thresholds and Genetic Factors that
Influence Benzene-Induced Genotoxicity. Environ Health Perspect. 2015;123(3):237-45. Epub
2014/11/06. doi: 10.1289/ehp.1408202. PubMed PMID: 25376053; PubMed Central PMCID:
PMCPMC4348743.
68.   Smallwood TL, Gatti DM, Quizon P, Weinstock GM, Jung KC, Zhao L, et al. High-
resolution genetic mapping in the diversity outbred mouse population identifies Apobec1 as a
candidate gene for atherosclerosis. G3 (Bethesda). 2014;4(12):2353-63. Epub 2014/10/23. doi:
10.1534/g3.114.014704. PubMed PMID: 25344410; PubMed Central PMCID: PMCPMC4267931.
69.   Shorter JR, Huang W, Beak JY, Hua K, Gatti DM, de Villena FP, et al. Quantitative trait
mapping in Diversity Outbred mice identifies two genomic regions associated with heart size.
Mamm Genome. 2018;29(1-2):80-9. Epub 2017/12/26. doi: 10.1007/s00335-017-9730-7.
PubMed PMID: 29279960; PubMed Central PMCID: PMCPMC6340297.
70.   Bressler J, Tsai TF, Wu MY, Tsai SF, Ramirez MA, Armstrong D, et al. The SNRPN
promoter is not required for genomic imprinting of the Prader-Willi/Angelman domain in mice.
Nat Genet. 2001;28(3):232-40. doi: 10.1038/90067. PubMed PMID: 11431693.
71.   Albrecht U, Sutcliffe JS, Cattanach BM, Beechey CV, Armstrong D, Eichele G, et al.
Imprinted expression of the murine Angelman syndrome gene, Ube3a, in hippocampal and
Purkinje neurons. Nat Genet. 1997;17(1):75-8. doi: 10.1038/ng0997-75. PubMed PMID:
9288101.
72.   Kobayashi S, Kohda T, Ichikawa H, Ogura A, Ohki M, Kaneko-Ishino T, et al. Paternal
expression of a novel imprinted gene, Peg12/Frat3, in the mouse 7C region homologous to the
Prader-Willi syndrome region. Biochem Biophys Res Commun. 2002;290(1):403-8. doi:
10.1006/bbrc.2001.6160. PubMed PMID: 11779183.
73.   Morison IM, Ramsay JP, Spencer HG. A census of mammalian imprinting. Trends Genet.
2005;21(8):457-65. doi: 10.1016/j.tig.2005.06.008. PubMed PMID: 15990197.
74.   Gray TA, Saitoh S, Nicholls RD. An imprinted, mammalian bicistronic transcript encodes
two independent proteins. Proc Natl Acad Sci U S A. 1999;96(10):5616-21. doi:
10.1073/pnas.96.10.5616. PubMed PMID: 10318933; PubMed Central PMCID: PMCPMC21909.
75.   Knoll JH, Nicholls RD, Magenis RE, Graham JM, Lalande M, Latt SA. Angelman and
Prader-Willi syndromes share a common chromosome 15 deletion but differ in parental origin

of the deletion. Am J Med Genet. 1989;32(2):285-90. doi: 10.1002/ajmg.1320320235. PubMed
PMID: 2564739.
76.   Mic FA, Molotkov A, Fan X, Cuenca AE, Duester G. RALDH3, a retinaldehyde
dehydrogenase that generates retinoic acid, is expressed in the ventral retina, otic vesicle and
olfactory pit during mouse development. Mech Dev. 2000;97(1-2):227-30. doi: 10.1016/s0925-
4773(00)00434-2. PubMed PMID: 11025231.
77.   Visel A, Thaller C, Eichele G. GenePaint.org: an atlas of gene expression patterns in the
mouse embryo. Nucleic Acids Res. 2004;32(Database issue):D552-6. doi: 10.1093/nar/gkh029.
PubMed PMID: 14681479; PubMed Central PMCID: PMCPMC308763.
78.   Martin KM, Metcalfe JC, Kemp PR. Expression of Klf9 and Klf13 in mouse development.
Mech Dev. 2001;103(1-2):149-51. doi: 10.1016/s0925-4773(01)00343-4. PubMed PMID:
11335124.
79.   Yokoyama S, Ito Y, Ueno-Kudoh H, Shimizu H, Uchibe K, Albini S, et al. A systems
approach reveals that the myogenesis genome network is regulated by the transcriptional
repressor RP58. Dev Cell. 2009;17(6):836-48. doi: 10.1016/j.devcel.2009.10.011. PubMed PMID:
20059953; PubMed Central PMCID: PMCPMC3110151.
80.   Unda FJ, Pérez-Nanclares G, Le Morvan V, Hernández C, Vilaxa A, De-la-Fuente M, et al.
Dynamic assembly of tight junction-associated proteins ZO-1, ZO-2, ZO-3 and occludin during
mouse tooth development. Histol Histopathol. 2003;18(1):27-38. doi: 10.14670/HH-18.27.
PubMed PMID: 12507281.
81.   Yoshida T, Miyoshi J, Takai Y, Thesleff I. Cooperation of nectin-1 and nectin-3 is required
for normal ameloblast function and crown shape development in mouse teeth. Dev Dyn.
2010;239(10):2558-69. doi: 10.1002/dvdy.22395. PubMed PMID: 21038445.
82.   de Koning DJ, Bovenhuis H, van Arendonk JA. On the detection of imprinted quantitative
trait loci in experimental crosses of outbred species. Genetics. 2002;161(2):931-8. PubMed
PMID: 12072486; PubMed Central PMCID: PMCPMC1462155.
83.   Cheverud JM, Hager R, Roseman C, Fawcett G, Wang B, Wolf JB. Genomic imprinting
effects on adult body composition in mice. Proc Natl Acad Sci U S A. 2008;105(11):4253-8. Epub
2008/03/12. doi: 10.1073/pnas.0706562105. PubMed PMID: 18337500; PubMed Central
PMCID: PMCPMC2393747.
84.   Leamy LJ, Klingenberg CP, Sherratt E, Wolf JB, Cheverud JM. A search for quantitative
trait loci exhibiting imprinting effects on mouse mandible size and shape. Heredity (Edinb).
2008;101(6):518-26. Epub 2008/08/06. doi: 10.1038/hdy.2008.79. PubMed PMID: 18685568.
85.   Holm VA, Cassidy SB, Butler MG, Hanchett JM, Greenswag LR, Whitman BY, et al.
Prader-Willi syndrome: consensus diagnostic criteria. Pediatrics. 1993;91(2):398-402. PubMed
PMID: 8424017; PubMed Central PMCID: PMCPMC6714046.
86.   Katz DC, Grote MN, Weaver TD. Changes in human skull morphology across the
agricultural transition are consistent with softer diets in preindustrial farming groups. Proc Natl
Acad Sci U S A. 2017;114(34):9050-5. Epub 2017/07/24. doi: 10.1073/pnas.1702586114.
PubMed PMID: 28739900; PubMed Central PMCID: PMCPMC5576786.
87.   Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov JP.
Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011;27(12):1739-40. Epub
2011/05/05. doi: 10.1093/bioinformatics/btr260. PubMed PMID: 21546393; PubMed Central
PMCID: PMCPMC3106198.

88.   Le N, Nagarajan R, Wang JY, Svaren J, LaPash C, Araki T, et al. Nab proteins are essential
for peripheral nervous system myelination. Nat Neurosci. 2005;8(7):932-40. PubMed PMID:
16136673.
89.   Shi C, Sakuma M, Mooroka T, Liscoe A, Gao H, Croce KJ, et al. Down-regulation of the
forkhead transcription factor Foxp1 is required for monocyte differentiation and macrophage
function. Blood. 2008;112(12):4699-711. Epub 2008/09/17. doi: 10.1182/blood-2008-01-
137018. PubMed PMID: 18799727; PubMed Central PMCID: PMCPMC2597137.
90.   Zhao H, Zhou W, Yao Z, Wan Y, Cao J, Zhang L, et al. Foxp1/2/4 regulate endochondral
ossification as a suppresser complex. Dev Biol. 2015;398(2):242-54. Epub 2014/12/17. doi:
10.1016/j.ydbio.2014.12.007. PubMed PMID: 25527076; PubMed Central PMCID:
PMCPMC4342236.
91.   Pacheco-Costa R, Kadakia JR, Atkinson EG, Wallace JM, Plotkin LI, Reginato RD.
Connexin37 deficiency alters organic bone matrix, cortical bone geometry, and increases
Wnt/β-catenin signaling. Bone. 2017;97:105-13. Epub 2017/01/16. doi:
10.1016/j.bone.2017.01.010. PubMed PMID: 28096061.
92.   Wilson DG, Phamluong K, Lin WY, Barck K, Carano RA, Diehl L, et al. Chondroitin sulfate
synthase 1 (Chsy1) is required for bone development and digit patterning. Dev Biol.
2012;363(2):413-25. Epub 2012/01/17. doi: 10.1016/j.ydbio.2012.01.005. PubMed PMID:
22280990.
93.   Xing WR, Goodluck H, Zeng C, Mohan S. Role and mechanism of action of leucine-rich
repeat kinase 1 in bone. Bone Res. 2017;5:17003. Epub 2017/03/14. doi:
10.1038/boneres.2017.3. PubMed PMID: 28326224; PubMed Central PMCID:
PMCPMC5348726.
94.   Vecchione L, Byron C, Cooper GM, Barbano T, Hamrick MW, Sciote JJ, et al. Craniofacial
morphology in myostatin-deficient mice. J Dent Res. 2007;86(11):1068-72. doi:
10.1177/154405910708601109. PubMed PMID: 17959898.
95.   Vecchione L, Miller J, Byron C, Cooper GM, Barbano T, Cray J, et al. Age-related changes
in craniofacial morphology in GDF-8 (myostatin)-deficient mice. Anat Rec (Hoboken).
2010;293(1):32-41. doi: 10.1002/ar.21024. PubMed PMID: 19899116; PubMed Central PMCID:
PMCPMC3113544.
96.   Sahni M, Ambrosetti DC, Mansukhani A, Gertner R, Levy D, Basilico C. FGF signaling
inhibits chondrocyte proliferation and regulates bone development through the STAT-1
pathway. Genes & Development. 1999;13:1361-6.
97.   Xiao L, Naganawa T, Obugunde E, Gronowicz G, Ornitz DM, Coffin JD, et al. Stat1
controls postnatal bone formation by regulating fibroblast growth factor signaling in
osteoblasts. J Biol Chem. 2004;279(26):27743-52. Epub 2004/04/08. doi:
10.1074/jbc.M314323200. PubMed PMID: 15073186.
98.   Kim S, Koga T, Isobe M, Kern BE, Yokochi T, Chin YE, et al. Stat1 functions as a
cytoplasmic attenuator of Runx2 in the transcriptional program of osteoblast differentiation.
Genes Dev. 2003;17(16):1979-91. doi: 10.1101/gad.1119303. PubMed PMID: 12923053;
PubMed Central PMCID: PMCPMC196253.
99.   Le Fevre AK, Taylor S, Malek NH, Horn D, Carr CW, Abdul-Rahman OA, et al. FOXP1
mutations cause intellectual disability and a recognizable phenotype. Am J Med Genet A.

2013;161A(12):3166-75. Epub 2013/09/24. doi: 10.1002/ajmg.a.36174. PubMed PMID:
24214399.
100.   Tian J, Ling L, Shboul M, Lee H, O'Connor B, Merriman B, et al. Loss of CHSY1, a secreted
FRINGE enzyme, causes syndromic brachydactyly in humans via increased NOTCH signaling. Am
J Hum Genet. 2010;87(6):768-78. doi: 10.1016/j.ajhg.2010.11.005. PubMed PMID: 21129727;
PubMed Central PMCID: PMCPMC2997365.
101.   Huh A, Horton MJ, Cuenco KT, Raoul G, Rowlerson AM, Ferri J, et al. Epigenetic influence
of KAT6B and HDAC4 in the development of skeletal malocclusion. Am J Orthod Dentofacial
Orthop. 2013;144(4):568-76. doi: 10.1016/j.ajodo.2013.06.016. PubMed PMID: 24075665;
PubMed Central PMCID: PMCPMC3849704.
102.   Vimalraj S, Arumugam B, Miranda PJ, Selvamurugan N. Runx2: Structure, function, and
phosphorylation in osteoblast differentiation. Int J Biol Macromol. 2015;78:202-8. Epub
2015/04/13. doi: 10.1016/j.ijbiomac.2015.04.008. PubMed PMID: 25881954.
103.   Clayton-Smith J, O'Sullivan J, Daly S, Bhaskar S, Day R, Anderson B, et al. Whole-exome-
sequencing identifies mutations in histone acetyltransferase gene KAT6B in individuals with the
Say-Barber-Biesecker variant of Ohdo syndrome. Am J Hum Genet. 2011;89(5):675-81. doi:
10.1016/j.ajhg.2011.10.008. PubMed PMID: 22077973; PubMed Central PMCID:
PMCPMC3213399.
104.   Sena K, Morotome Y, Baba O, Terashima T, Takano Y, Ishikawa I. Gene expression of
growth differentiation factors in the developing periodontium of rat molars. J Dent Res.
2003;82(3):166-71. doi: 10.1177/154405910308200304. PubMed PMID: 12598543.
105.   Nakashima M, Toyono T, Murakami T, Akamine A. Transforming growth factor-beta
superfamily members expressed in rat incisor pulp. Arch Oral Biol. 1998;43(9):745-51. doi:
10.1016/s0003-9969(98)00046-6. PubMed PMID: 9783830.
106.   Cerny R, Cattell M, Sauka-Spengler T, Bronner-Fraser M, Yu F, Medeiros DM. Evidence
for the prepattern/cooption model of vertebrate jaw evolution. Proc Natl Acad Sci U S A.
2010;107(40):17262-7. Epub 2010/09/20. doi: 10.1073/pnas.1009304107. PubMed PMID:
20855630; PubMed Central PMCID: PMCPMC2951391.
107.   So PL, Danielian PS. Cloning and expression analysis of a mouse gene related to
Drosophila odd-skipped. Mech Dev. 1999;84(1-2):157-60. PubMed PMID: 10473132.
108.   Kawai S, Yamauchi M, Amano A. Zinc-finger transcription factor Odd-skipped related 1
regulates cranial bone formation. J Bone Miner Metab. 2018;36(6):640-7. Epub 2017/12/12.
doi: 10.1007/s00774-017-0885-9. PubMed PMID: 29234951.
109.   Jeong J, Cesario J, Zhao Y, Burns L, Westphal H, Rubenstein JL. Cleft palate defect of
Dlx1/2-/- mutant mice is caused by lack of vertical outgrowth in the posterior palate. Dev Dyn.
2012;241(11):1757-69. Epub 2012/09/28. doi: 10.1002/dvdy.23867. PubMed PMID: 22972697;
PubMed Central PMCID: PMCPMC3988582.
110.   Zelzer E, McLean W, Ng YS, Fukai N, Reginato AM, Lovejoy S, et al. Skeletal defects in
VEGF(120/120) mice reveal multiple roles for VEGF in skeletogenesis. Development.
2002;129(8):1893-904. PubMed PMID: 11934855.
111.   R Development Core Team. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/2015.
112.   Mitteroecker P, Gunz P. Advances in Geometric Morphometrics. Evolutionary Biology.
2009;36(2):235-47. doi: 10.1007/s11692-009-9055-x.

113.   Schlager S. Morpho and Rvcg – Shape Analysis in R. In: Zheng G, Li S, Szekely G, editors.
Statistical Shape and Deformation Analysis: Academic Press; 2017. p. 217-56.
114.   Mardia KV, Bookstein FL, Moreton IJ. Statistical assessment of bilateral symmetry of
shapes. Biometrika. 2000;87(2):285-300.
115.   Klingenberg CP, Barluenga M, Meyer A. Shape analysis of symmetric structures:
quantifying variation among individuals and asymmetry. Evolution. 2002;56(10):1909-20.
PubMed PMID: 12449478.
116.   Morgan AP, Fu CP, Kao CY, Welsh CE, Didion JP, Yadgary L, et al. The Mouse Universal
Genotyping Array: From Substrains to Subspecies. G3 (Bethesda). 2015;6(2):263-79. Epub
2015/12/18. doi: 10.1534/g3.115.022087. PubMed PMID: 26684931; PubMed Central PMCID:
PMCPMC4751547.
117.   Didion JP, Morgan AP, Clayshulte AM, Mcmullan RC, Yadgary L, Petkov PM, et al. A
multi-megabase copy number gain causes maternal transmission ratio distortion on mouse
chromosome 2. PLoS Genet. 2015;11(2):e1004850. Epub 2015/02/13. doi:
10.1371/journal.pgen.1004850. PubMed PMID: 25679959; PubMed Central PMCID:
PMCPMC4334553.
118.   Cardon LR, Palmer LJ. Population stratification and spurious allelic association. Lancet.
2003;361(9357):598-604. doi: 10.1016/S0140-6736(03)12520-2. PubMed PMID: 12598158.
119.   Broman KW, Gatti DM, Svenson KL, Sen Ś, Churchill GA. Cleaning Genotype Data from
Diversity Outbred Mice. G3 (Bethesda). 2019;9(5):1571-9. Epub 2019/05/07. doi:
10.1534/g3.119.400165. PubMed PMID: 30877082; PubMed Central PMCID: PMCPMC6505173.
120.   Sybert VP, McCauley E. Turner's syndrome. N Engl J Med. 2004;351(12):1227-38. doi:
10.1056/NEJMra030360. PubMed PMID: 15371580.
121.   Hallgrimsson BA, J David, Katz DC, Bannister JJ, Riccardi SL, Mahasuwan N, McInnes BL,
et al. Automated syndrome diagnosis by three-dimensional facial photogrammetric imaging.
American Journal of Human Genetics. 2019, in review.
122.   Krzanowski WJ. Principles of Multivariate Analysis: A User's Perspective. New York:
Oxford University Press; 2005.
123.   Churchill GA, Doerge RW. Empirical threshold values for quantitative trait mapping.
Genetics. 1994;138(3):963-71. PubMed PMID: 7851788; PubMed Central PMCID:
PMCPMC1206241.
124.   Manichaikul A, Dupuis J, Sen S, Broman KW. Poor performance of bootstrap confidence
intervals for the location of a quantitative trait locus. Genetics. 2006;174(1):481-9. Epub
2006/06/18. doi: 10.1534/genetics.106.061549. PubMed PMID: 16783000; PubMed Central
PMCID: PMCPMC1569776.
125.   Yalcin B, Flint J, Mott R. Using progenitor strain information to identify quantitative trait
nucleotides in outbred mice. Genetics. 2005;171(2):673-81. Epub 2005/08/05. doi:
10.1534/genetics.104.028902. PubMed PMID: 16085706; PubMed Central PMCID:
PMCPMC1456780.
126.   Baud A, Hermsen R, Guryev V, Stridh P, Graham D, McBride MW, et al. Combined
sequence-based and genetic mapping analysis of complex traits in outbred rats. Nat Genet.
2013;45(7):767-75. Epub 2013/05/26. doi: 10.1038/ng.2644. PubMed PMID: 23708188;
PubMed Central PMCID: PMCPMC3821058.

127.   Rhee SY, Wood V, Dolinski K, Draghici S. Use and misuse of the gene ontology
annotations. Nat Rev Genet. 2008;9(7):509-15. Epub 2008/05/13. doi: 10.1038/nrg2363.
PubMed PMID: 18475267.
128.   Maglott D, Ostell J, Pruitt KD, Tatusova T. Entrez Gene: gene-centered information at
NCBI. Nucleic Acids Res. 2011;39(Database issue):D52-7. Epub 2010/11/28. doi:
10.1093/nar/gkq1237. PubMed PMID: 21115458; PubMed Central PMCID: PMCPMC3013746.
129.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene
set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression
profiles. Proc Natl Acad Sci U S A. 2005;102(43):15545-50. Epub 2005/09/30. doi:
10.1073/pnas.0506580102. PubMed PMID: 16199517; PubMed Central PMCID:
PMCPMC1239896.
130.   Katz DC, Grote MN, Weaver TD. A mixed model for the relationship between climate
and cranial form. American Journal of Physical Anthropology. 2016;160:593-603.
131.   Mitteroecker P, Cheverud JM, Pavlicev M. Multivariate Analysis of Genotype-Phenotype
Association. Genetics. 2016. doi: 10.1534/genetics.115.181339.
132.   Nakagawa S, Schielzeth H. A general and simple method for obtaining R^2 from
generalized linear mixed-effects models. Methods in Ecology and Evolution. 2013;4:133-42. doi:
10.1111/j.2041-210x.2012.00261.x.
133.   Adler D, Murdoch D, others. rgl: 3D Visualization Using OpenGL. 0.95.1441 ed2016.
134.   Rohlf FJ. Shape statistics: Procrustes superimpositions and tangent spaces. Journal of
Classification. 1999;16(2):197-223.
968

Supporting Information

S1 Fig. Eigenvalue decay. Percentage of total variation explained for each shape principal

component. We performed genome scans for all PCs explaining more than 1% of shape

variation (red point identifies cutoff at PC 20) and subjected the first five PCs (green point

identifies PC 5 floor) to in-depth analysis.

S2 Fig. Change in chromosome 2 WSB allele frequencies over generations. Sample-wide WSB

allele frequency on chromosome 2 by DO generation. The high WSB allele frequency in earlier

generations reflects meiotic drive for WSB at the *R2d2* locus. Declining frequencies in later

generations reflect the systematic purge of WSB alleles from the DO population.

S1 Table. LOD peaks and support intervals for all QTL on PCs 1-20.

S2 Table. Full array of LOCO mean heritabilities over PCs 1-20.

S1 File. Genes, PCs 1-5. All named, protein-coding genes in the PC 1-5 QTL intervals.

S2 File. Top GO annotations, PCs 1-5. Top 100 FDR q-value GO terms for the PC 1-5 QTL interval

gene set (Molecular Signatures Database output file).

S3 File. Top GO annotations, PC 1. For the PC1 QTL interval gene set, GO terms with FDR q-

value < 0.05 (Molecular Signatures Database output file).