The seagrass methylome memorizes heat stress and is associated with variation in stress performance among clonal shoots

Jueterbock A[1*], Boström C[2], James A Coyer[1,3], Olsen JL[4,] Kopp M[1], Dhanasiri AKS[1,] Smolina I[1] , Arnaud-Haond S[5], Van de Peer Y[6,7], Hoarau G[1]

[1]Marine Molecular Ecology Group, Faculty of Biosciences and Aquaculture, Nord University, Universitetsalleen 11, 8026 Bodø, Norway

[2]Environmental and Marine Biology, Åbo Akademi University, Artillerigatan 6 FI-20520 Åbo, Finland

[3]Shoals Marine Laboratory, University of New Hampshire, Durham, NH 03824, USA

[4]Ecological Genetics-Genomics Group, Groningen Institute of Evolutionary Life Sciences, University of Groningen, 9747 AG Groningen, The Netherlands

[5]UMR MARBEC, Université de Montpellier, Ifremer, IRD, CNRS, Avenue Jean Monnet, CS 30171, 34203 Sète Cedex, France

[6]Department of Plant Biotechnology and Bioinformatics, Ghent University and VIB Center for Plant Systems Biology, Technologiepark 71, 9052, Ghent, Belgium

[7]Center for Microbial Ecology and Genomics, Department of Biochemistry, Genetics and Microbiology, University of Pretoria, Pretoria, South Africa

* Correspondence:

Corresponding Author

Alexander-Jueterbock@web.de

Keywords: DNA methylation, ecological epigenetics, clonality, heat stress, seagrass, *Zostera marina* (eelgrass), priming

Word count: Abstract 253, Main Article 7,239

Number of figures: 6

Number of tables: 0

Number of supplementary figures: 5

Number of supplementary tables: 11

Number of supplementary files: 5

Abstract

Evolutionary theory predicts that clonal organisms are more susceptible to extinction than sexually
reproducing organisms, due to low genetic variation and slow rates of evolution. In agreement,
conservation management considers genetic variation as the ultimate measure of a population's
ability to survive over time. However, clonal plants are among the oldest living organisms on our
planet. Here, we test the hypothesis that clonal seagrass meadows display epigenetic variation that
complements genetic variation as a source of phenotypic variation. In a clonal meadow of the
seagrass *Zostera marina* we characterized DNA methylation among 42 shoots. We also sequenced
the whole genome of 10 shoots to correlate methylation patterns with photosynthetic performance
under exposure to, and recovery from 27°C, while controlling for somatic mutations. Here, we show
for the first time that clonal seagrass shoots display DNA methylation variation that is associated
with variation in fitness-related traits: photosynthetic performance and heat stress resilience. The co-
variation in DNA methylation and phenotype may be linked via gene expression because methylation
patterns varied in functionally relevant genes involved in photosynthesis, and in the repair and
prevention of heat-induced protein damage. A >five week epigenetic heat stress memory may heat-
harden previously heat-exposed shoots. While genotypic diversity has been shown to enhance stress
resilience in seagrass meadows, we suggest that epigenetic variation plays a similar role in meadows
dominated by a single genotype. Consequently, conservation management of clonal plants should
consider epigenetic variation as indicator of resilience and stability, and restoration efforts may
benefit from stress-priming transplanted seeds or shoots.

1   Introduction

Genetic variation is considered key to long-term survival of populations (Bijlsma and Loeschcke, 2012), and is recognized as a main target for the conservation and restoration of biological diversity (Lande, 1988; Spielman et al., 2004; Laikre et al., 2009). In contrast, lack of genetic variation is regarded as an evolutionary dead-end (Lynch and Lande, 1998) but clonal growth challenges the expected relationship between genetic diversity and long-term survival. For example, conservation management follows the rough guideline that within a population >1,000 genetically different individuals must mate randomly to avoid inbreeding depression, and retain evolutionary potential (Frankham et al., 2014). Yet, roughly 40% of all plants can reproduce asexually (Tiffney and Niklas, 1985), mostly by clonal growth where parental genotypes (genets) grow vegetative modules (ramets) that often remain connected via underground stolons or rhizomes.

Although clones benefit from resource sharing, niche specialization, and rapid vegetative growth (Liu et al., 2016), they are predicted to survive only for short periods, and in stable environments (Silvertown, 2008). Asexual reproduction is assumed to lead to slow rates of genetic evolution, and the lack of DNA repair mechanisms afforded by meiosis (Muller's ratchet, mutational meltdown) (Muller, 1964; Gabriel et al., 1993; Lynch et al., 1993). Despite asexual reproduction, many of our most important crops are clones (McKey et al., 2010), including banana, garlic, hops, potatoes, and turmeric, as well as many of the earth's most invasive and oldest plants. For example, genets of Palmer's oak (*Quercus palmeri*) and seagrass (*Posidonia oceanica*) are estimated older than 10,000 years (May et al., 2009; Arnaud-Haond et al., 2012) Thus, long-term survival appears not to rely solely on sexual production.

Somatic mutations can create a certain level of genetic diversity, and may explain some evolutionary potential of clonal organisms (Whitham and Slobodchikoff, 1981; Loxdale et al., 2003; Lushai et al., 2003; Reusch and Boström, 2011). For example, ~7,000 single nucleotide polymorphisms (SNPs), 597 in coding regions and 432 non-synonymous, distinguish ramets of a large Finnish clone of the seagrass *Zostera marina* (Yu et al., 2020). To set this in perspective ,139,321 biallelic SNPs were reported in coding regions among four populations of the same species (Jueterbock et al., 2016). The degree to which epigenetic variation can contribute to phenotypic heterogeneity in ecologically relevant traits, independently from the underlying genetic variation, is a key question in assessing its contribution to stress tolerance, and long-term survival of clonal organisms.

The definition of epigenetics is currently heavily debated (Ptashne, 2007; Greally, 2018). Here,

'epigenetic' implies molecular variations that do not alter the DNA sequence but have the potential to

change gene expression, and include non-coding RNAs (ncRNAs), histone modifications, and DNA

methylation (Bossdorf et al., 2008). DNA methylation is, from an evolutionary perspective, the most

relevant epigenetic mechanism because it can be independent from genetic variation (Bossdorf et al.,

2008; Schmitz et al., 2013; Kilvitis et al., 2014), and transgenerationally stable (Boyko et al., 2010;

Verhoeven et al., 2010; Ou et al., 2012; Bilichak et al., 2015; Williams and Gehring, 2017). DNA

methylation involves the addition of a methyl-group to the C5 position of a cytosine in DNA

sequence motifs (CG, CHG, and CHH in plants, where H stands for A, C, or T) (Kilvitis et al., 2014).

Depending on sequence context, methylation can be associated with gene activation or silencing

(Bossdorf et al., 2008; Niederhuth and Schmitz, 2017). While CG methylation in gene bodies often

correlates with increased gene expression, methylation in promoters and repeat regions, such as

transposable elements (TEs), silences expression (Feng et al., 2010; Seymour et al., 2014; Dubin et

al., 2015; Bewick and Schmitz, 2017; Zhang et al., 2018).

The methylome, or set of DNA methylation modifications in an organism's genome, can change

spontaneously at a rate of 2.5×10^{-4} to 6.3×10^{-4} methylation polymorphisms per CG site per generation,

which is about 7×10^{4} higher than the genetic mutation rate of base substitutions per site per

generation (Schmitz et al., 2011; van der Graaf et al., 2015). That methylome variation can enhance

productivity, and pathogen resistance, has been shown in *Arabidopsis thaliana* plant populations

(Latzel et al., 2013). Moreover, methylation variation explained the rapid invasive success of the

Japanese knotweed (*Fallopia japonica*) by facilitating differentiation in response to new habitats

despite decreased genetic variation (Richards et al., 2012). This suggests that methylation variation

complements genetic variation as a source of phenotypic variation in plant populations deprived of

genotypic diversity (Gao et al., 2010; Richards et al., 2012; Verhoeven and van Gurp, 2012; Latzel et

al., 2013; Zhang et al., 2013; Vanden Broeck et al., 2018).

Unlike genetic variants, methylation variants can also switch state directly in response to the

environment (Dowen et al., 2012) and, if stable enough, establish a molecular memory that can be

involved in stress priming. Priming, also referred to as hardening or conditioning, is well studied in

terrestrial plants (Hilker et al., 2016; Lämke and Bäurle, 2017; Pawar and Laware, 2018), and refers

to the plants' ability to acquire a stress memory that enhances performance under second stress

exposure by responding faster, stronger, or in response to a lower threshold as compared with naïve

plants (Balmer et al., 2015; Lämke and Bäurle, 2017). Stress responsive methylation marks that are transgerationally stable can create adaptive phenotypic changes within a single generation (Bossdorf et al., 2008; Jablonka and Raz, 2009; Nicotra et al., 2010; Verhoeven et al., 2010; Hirsch et al., 2012; Richards et al., 2012; Douhovnikoff and Dodd, 2014; Verhoeven and Preite, 2014).This may explain why in some cases a primed state can improve plant performance even across generations (Rasmann et al., 2012; Slaughter et al., 2012; Kuźnicki et al., 2019).

Under clonal growth, DNA methylation patterns are expected to be more stably inherited than under sexual reproduction (Verhoeven and Preite, 2014), because clonal growth circumvents epigenetic reprogramming during gameto- and embryogenesis. Although stable transmission across asexual generations has been shown for environment-specific phenotypes and DNA methylation patterns in clonal plants (Verhoeven et al., 2010, 2018; Verhoeven and van Gurp, 2012; Vanden Broeck et al., 2018), no link has been demonstrated between methylation variation and fitness-related traits. Thus, while epigenetic mechanisms have been suggested to contribute to clonal plant success (Douhovnikoff and Dodd, 2014; Verhoeven and Preite, 2014; Dodd and Douhovnikoff, 2016; Latzel et al., 2016), empirical evidence is virtually lacking.

Clonal propagation is especially well-developed in aquatic plants (Barrett, 2015). Seagrasses, the only plants to inhabit the marine world, form the foundational basis of some of the most productive and highly diverse coastal marine ecosystems on the planet, and are essential for the health and abundance of economically exploited marine species (Costanza et al., 1997; Larkum et al., 2006; Orth et al., 2006). Ecosystem services are worth more than € 16,000 ha^{-1} year^{-1} (Costanza et al., 1997), including nursery grounds, habitat and food for fish and invertebrates, protection of the coastline from erosion, carbon sequestration of up to 186 g C m^2 yr^{-1} (Fourqurean et al., 2012; Duarte et al., 2013), and nutrient fixation (Orth et al., 2006; Procaccini et al., 2007).

Over the last decades, losses of seagrass ecosystems have been documented worldwide due to increasing anthropogenic stressors such as invasive species, sediment and nutrient runoff, dredging, aquaculture, rising sea levels, and global warming (Orth et al., 2006; Chefaoui et al., 2018). Losses are expected to accelerate under projected global temperature increase, as ocean warming is considered the most severe threat among climate change factors (Repolho et al., 2017; Duarte et al., 2018) and seagrass meadows have tracked temperature changes in the past (Olsen et al., 2004). How

accurate these shifts and losses can be predicted depends on our knowledge of drivers of adaptive

potential, including epigenetic diversity (Duarte et al., 2018).

In this study, we characterize the functional relevance of epigenetic variation for heat stress resilience

in the seagrass *Zostera marina*. *Z. marina* is the most widely distributed seagrass in the northern

hemisphere, inhabiting highly contrasting habitats from sub-arctic to sub-tropical waters (Green and

Short, 2004; Olsen et al., 2004; Boström et al., 2014). Few plants display such dramatic range in

clonal diversity as *Z. marina* (Reusch et al., 2000; Olsen et al., 2004). Its clonal architecture varies

from genetically diverse meadows with high levels of sexual reproduction, to meadows composed of

a single large clone due to exclusive vegetative reproduction (Reusch et al., 1999; Olsen et al., 2004;

Becheler et al., 2010; Reusch and Boström, 2011). Clonality of *Z. marina* peaks in the Baltic Sea,

where large clones were estimated > 1,000 years old (Reusch et al., 1999). These meadows display

remarkable phenotypic variability and persistence across space and time, under perturbations (e.g.

high temperatures) that represent environmental stresspredicted elsewhere for the future (Reusch et

al., 2018). Thus, clonal meadows (such predominated by a single genet) confound experimental

results showing a positive effect of genotypic (i. e. clonal) richness on the productivity and stress

resilience of *Z. marina* (Hughes and Stachowicz, 2004; Reusch et al., 2005; Reusch and Hughes,

2006; Ehlers et al., 2008; Hughes et al., 2008). In other words, if stress resilience and tolerance would

rely strongly on genotypic diversity, clones would not be able to survive for such long time periods.

The resilience, longevity, and adaptive potential of clonal seagrass meadows remain unknown

without a fundamental analysis of their epigenetic variation and its ecological relevance. Therefore,

this study tests the hypothesis that variation in DNA methylation can promote functional phenotypic

diversity and, thus, may explain how clonal seagrass meadows can persist over millennia.

Specific objectives were to: 1) Characterize DNA methylation variation in an ancient clonal meadow

of *Z. marina* (>1,000-years-old); 2) Identify the functional role of this variation in heat-stress

performance–specifically how photosynthetic performance is linked to DNA methylation variation

independently from underlying somatic mutations, and 3) Assess to what extent heat-responsive

methylation patterns are reset versus memorized after the stress is removed.

2   Materials and Methods

2.1 Sampling and cultivation

We sampled seagrass shoots from a clonal meadow (Reusch et al., 1999) in the Baltic Sea (Åland Islands, 60°09'50.4"N, 19°31'48.1"E) in 2015 (Figure 1) by collecting 2-3 shoots attached to the same rhizome (ramets) every 3 meters along a 250 m transect (Supplementary table S1). Shoots were transported in seawater-filled cooling boxes to the field station at Nord University, Norway. Leaf tissue of 42 shoots, randomly chosen from the 2-3 connected ramets, was flash frozen in liquid nitrogen for subsequent DNA extraction (Supplementary table S1).

Ten ramets of the same genet (Supplementary table S1) were replanted in plastic containers (10x15 cm) filled with sand from the sampling site, placed in a 1,280 L aquarium at 15°C, and illuminated with a 16:8 h light:dark cycle, and 200-220 µmol m^{-2} s^{-1} (OSRAM Fluora, 150 W). Seawater at 5.5 PSU, corresponding to the salinity at the collecting site, was obtained by mixing freshwater and natural filtered seawater at 32 PSU. The water, filled to a 40 cm level, was kept in constant motion with airstones. Once a week, 50% of the water was renewed after removing epiphytic algae.

2.2 Heat stress experiments

After two weeks of acclimation, the 10 clonal ramets were exposed to heat stress (Supplementary table S1) in climate chambers (Fitotron, weisstechnik), where they were distributed in aquaria (60 L) filled with aerated brackish water (5.5 PSU) to 30 cm, of which 50% was weekly renewed. Light was kept at 240 µmol m^{-2} s^{-1} and a 16:8 h light:dark cycle. The temperature in the climate chambers was increased with a daily increment of 3°C from 15°C to 27°C, which can be lethal for *Z. marina* (Greve et al., 2003), but is reached in the Baltic Sea during summer heat waves. After three weeks at 27°C, the temperature was decreased by a daily increment of 3°C from 27°C to 15°C, after which all shoots were returned to the 1,280 L aquarium at 15°C for a recovery period of 5.5 weeks.

At three time points (control, stress, recovery), one leaf of each shoot was excised and flash-frozen in liquid nitrogen. Control samples were collected at 15°C on the day before the temperature was increased. Stressed samples were collected at 27°C on the day before the temperature was decreased. Recovery samples were collected after the 5.5 week recovery period at 15°C.

2.3 Photosynthetic performance

At control, stress, and recovery time points, we measured for each shoot (two measurements for each

of the two inner leaves) the increase in chlorophyll a fluorescence upon illumination after a 10 min

dark period (OJIP curve) (Bussotti et al., 2010), with a PAM-Fluorometer (FluorPen FP100, Photon

Systems Instruments) using a saturating pulse of 75% at 455 nm. From the measurements, we

extracted the performance index PiABS (Strasser et al., 2000), reflecting the functionality of

photosystem II (PSII) and photosynthetic performance in general (Živčák et al., 2008; Bussotti et al.,

2010; Stefanov et al., 2011). We tested for correlation of PiABS values (mean values per shoot)

between time points using a two-sided Pearson's product moment correlation test in R v3.4.4 (R Core

Team, 2019).

2.4   DNA extraction

All flash-frozen tissue was freeze-dried for one day, then stored at -80°C until DNA extraction. DNA

was extracted with the HP Plant DNA mini kit (Omega Bio-Tek, protocol version May 2013) from $\geq$

5 mg of lyophilized leaf tissue after grinding with a mixer-mill (Retsch MM 400) in 2 mL Eppendorf

tubes with tungsten beads supplied (60 s at 30 Hz). We added 10 µL beta-mercaptoethanol at step 2,

and equilibrated the columns at step 8 of the standard protocol. The extracted DNA was eluted in 2 x

100 µL EB buffer (Qiagen), cleaned and concentrated with the Clean and Concentrator-5 kit (Zymo

Research, protocol v1.2.1) using 15,000 g for all centrifugation steps, and finally eluted in 2 x 30 µL

EB Buffer (Qiagen) at 60°C.

2.5   Clonality

In order to identify ramets belonging to the same genet, we genotyped each shoot for seven

microsatellite loci (ZosmarGA2, -GA3, -GA6, -CT3, -CT12, -CT19, -CT20) (Reusch and Boström,

2011). PCR was performed using a Veriti 96-Well Thermal Cycler (Applied Biosystems, Life

Technologies) in two 10 µL multiplex reactions containing 2 µL cleaned genomic DNA, and 1X

AccuStart II PCR ToughMix (Quanta bio). One multiplex reaction contained forward and reverse

primers GA2, GA3, and GA6 at 0.5 µM each, and ran at 94°C for 4 min, followed by 30 cycles of

94°C for 60 s, 55°C for 90 s, and 72°C for 90 s, and a final extension at 72°C for 10 min. The other

multiplex reaction contained forward and reverse primers CT3 (0.5 µM), CT12 (0.3 µM), CT19 (0.3

µM), and CT20 (0.3 µM), and ran at 94°C for 3 min, followed by 35 cycles of 94°C for 60 s, 57°C

for 60 s, and 72°C for 60 s, and a final extension at 72°C for 10 min.

DNA fragment lengths were determined on an ABI 3500xl Genetic Analyzer from 1 µL of 1:99

diluted PCR products mixed with 8.9 µL of HiDi Formamide (Life Technologies), and 0.1 µL of

Gene Scan 500 LIZ Size Standard (Life Technologies) after 5 min denaturation at 95°C. Alleles were

called with the GeneMapper v4.1 Software (Applied Biosystems, Thermo Fisher Scientific). The

shoots were assigned to multi-locus genotypes using the R package 'RClone' (Bailleul et al., 2016).

2.6   Whole genome sequencing, SNP detection, and genetic distance

In order to detect SNPs resulting from somatic mutations in the 10 heat-stressed ramets, genomic DNA

libraries were prepared according to the TruSeq DNA PCR-Free (Illumina) protocol, and sequenced on one

Illumina HiSeq 3/4000 lane (2x150bp) at the Norwegian Sequencing Centre (University of Oslo, Norway).

Raw reads (25.7 Million to 44.3 Million per library, Supplementary Table S2, NCBI BioProject

PRJNA575339[1]) were quality-checked with FastQC v0.11.8[2] to control for aberrant read base content, length

distribution, duplication, and over-representation. We used TrimGalore! v0.6.0[3] to remove adapter sequences

with a stringency of 3 bp overlap, and low-quality bases with a Phred score $Q < 20$ (99% base call accuracy).

The high-quality reads (25.5 to 44.0 Million per library, Supplementary Table S2) were mapped to the *Z.*

marina genome v2.1 (Olsen et al., 2016) with BWA v0.7.17 (Li and Durbin, 2009). Read duplicates were

removed with MarkDuplicatesSpark within GATK v4.1.4.1 (Auwera et al., 2013). SNPs were called with

HaplotypeCaller, followed by CombineGVCFs and GenotypeGVCFs within GATK v4.1.4.1 (Auwera et al.,

2013).

Before filtering SNPs, we excluded indels, non-variant sites, and alternate alleles not present in any

genotypes from the vcf file with SelectVariants within GATK v4.1.4.1 (Auwera et al., 2013). This set

of 759,407 raw SNPs was reduced to 105,443 SNPs after hard-filtering with vcffilter from vcflib

(Garrison, 2020) with thresholds that were based on density plots drawn with ggplot2 (Wickham,

2016): QualByDepth (QD < 15.0), FisherStrand (FS >12.0), RMSMappingQuality (MQ < 38),

MappingQualityRankSumTest (MQRankSum < -1.5), ReadPositionRankSumTest

(ReadPosRankSumand < -4.0), and Depth (DP > 4000.0, in order to remove SNPs potentially caused

by genome duplication). Subsequently, we used VCFtools v0.1.15 to remove genotpes with genotype

---

11   [1]Available upon publication

12   [2]http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

13   [3]https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/

quality < 30 (--minGQ 30) or depth < 20 (--minDP 20), and to remove SNPs with more than 2 alleles

(--min-alleles 2 --max-alleles 2), with a minor allele frequency of 0.01 (--maf 0.01), and with any

missing genotype (--max-missing-count 0). From the remaining 15,508 high-quality SNPs

(Supplementary File S1) we excluded all that shared the same genotypes among all 10 heat-stressed

shoots, as these reflected genetic differences only to the reference genome. The remaining 1,079

SNPs (Supplementary File S2) were used to estimate euclidean genetic distances among the 10

shoots using the R package vcfR v1.9.0 (Knaus and Grünwald, 2017) and the *dist* function of the R

package 'stats' v3.6.9 (R Core Team, 2019). We tested for correlation between genetic and physical

distance among the 10 heat-stressed shoots with Mantel tests in the R package 'vegan' v1.4-2

(Oksanen et al. 2019), using 1,000 permutations, and the Pearson's product moment correlation

method.

2.7   Methylome characterization

Sequencing libraries were prepared according to the MethylRAD protocol (Wang et al., 2015) with

few adjustments. MethylRAD is a genome-reduction method based on the methylation-dependent

restriction enzyme FspEI that targets fully methylated CCGG and CCWGG motifs, thus capturing

methylation in CG and CHG sequence contexts. MethylRAD has the potential to reveal genome-wide

DNA methylation patterns that are consistent with those generated from Whole Genome Bisulfite

Sequencing (Wang et al., 2015). First, sense and anti-sense oligos of adapters A1 and A2

(Supplementary File S3) were annealed in 10 µL containing 10 µM of each oligo (Eurofins), 10 mM

Tris HCl (Thermo Fisher), 50 mM NaCl (Thermo Fisher), and 1 mM EDTA (Thermo Fisher).

Library preparation began with digestion of 100 ng cleaned genomic DNA at 37°C for 4 h in 15 µL

containing 4 U FspEI (NEB), 1X CutSmart Buffer (NEB), and 30X Enzyme Activator Solution

(NEB). Digestion was verified on a TapeStation 2200 with a D1000 ScreenTape. Secondly, adapters

were ligated to the digested fragments over night at 4°C in 26µL containing 13 µL digestion solution,

0.1 µM each of two annealed adapters, 1X T4 ligase buffer (NEB), 1.5 µM ATP (NEB) and 1040 U

of T4 DNA ligase (NEB). Ligation products were amplified in 20 µL reactions containing 7 µL

ligated DNA, 0.05 µM of each primer (P1 and P2, Supplementary File S3), 0.3 mM dNTP, 1X

Phusion HF buffer (NEB) and 0.4 U Phusion high-fidelity DNA polymerase (NEB). PCR was

conducted using a Veriti 96-Well Thermal Cycler (Applied Biosystems, Life Technologies) with 16

cycles of 98°C for 5s, 60°C for 20 s, 72°C for 10 s, and a final extension of 5 min at 72°C. The target

band (approx. 100 bp) was extracted from a 2% E-Gel (Thermo Fisher). For multiplex sequencing,

shoot barcodes were introduced by means of PCR. Each 20 µL PCR reaction contained 12 µL of gel-

extracted PCR product, 0.2 µM of each primer (P3 and index primer, Supplementary File S3), 0.3

mM dNTP, 1X Phusion HF buffer (NEB) and 0.4 U Phusion high-fidelity DNA polymerase (NEB).

PCR was conducted with the same PCR cycling program outlined above. PCR products were purified

using AMPURE XP beads (Beckman Coulter) using a 1.8:1 volume ratio of beads to product, and a

final elution in 22 µL EB buffer (Qiagen). The purified fragments were sequenced on an Illumina

NextSeq 500 (1x75bp) using a high-output flow-cell.

The sequenced reads were quality-trimmed with TrimGalore! v0.4.1[4] by removing the adapter

sequences with a stringency of 3 bp overlap, low-quality bases with a Phred score Q < 20, and the

terminal 2 bp from both ends in order to eliminate artifacts that might have arisen at the ligation

position. Quality was checked with FastQC v0.11.8[5] to control for aberrant read base content, length

distribution, duplication and over-representation.

The high-quality reads were mapped with SOAP v1.11 (Li et al., 2008) to 628,255 *in silico* predicted

MethylRAD tags that were extracted from the *Z. marina* genome v2.1 from ORCAE (Sterck et al.,

2012) with the custom python script *InSilicoTypeIIbDigestion.py[6]*. For mapping, we allowed for two

mismatches, filtered reads with >1 N, and used a seed size of 8 bp. Based on the uniquely mapped

reads, we counted the coverage of each methylated site for each shoot using htseq-count (v0.7.2).

Methylation calls were retained only for sites with ≥ 2x coverage, which reduced the false-positive

rate from 1.10% to 0.23% for CG sites and from 2.50% to 0.89% for CHG sites. False-positive rates

were estimated as the percentage of methylation sites supported by at least two reads in the generally

unmethylated chloroplast genome. For each shoot, raw counts were normalized to reads-per-million

by dividing reads per site through the total number of reads per shoot library, times one million.

The methylated sites were annotated with the v2.1 gff3 file from ORCAE (Sterck et al., 2012), and

separated into genes, intergenic regions, and TEs. TEs were located in both genes and intergenic

regions, and contained repeats of the rnd-1/2/3/4 families and some selected repeats (rnd-5-family-

---

15 [4]https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/

16 [5]http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

17 [6]http://marinetics.org/2017/04/11/REdigestions.html

18   11

305 7328, rnd-5-family-287, rnd-5-family-4856). The methylated sites were further separated into such

306 containing CG and CHG recognition sites.

2.8 Methylation variation between the transect shoots

In order to describe the level of intra- and inter-clonal methylation differences among the 42 transect

shoots, we converted the normalized counts to bivariate states (methylated/non-methylated). We then

calculated the number of sites with different methylation states pairwise between all 41 ramets of the

same genet, and between these ramets and the single other seven-microsatellite-genotype (transect

shoot 27, Supplementary table S1).

We tested for correlations between epigenetic and physical distance among the 41 ramets with

Mantel tests using the R package 'vegan' v1.4-2 (Oksanen et al., 2019) with 1,000 permutations, and

the Pearson's product moment correlation method. *P*-values were adjusted according the Benjamini-

Hochberg method (Benjamini and Hochberg, 1995). Epigenetic distance was calculated as the

Euclidean distance between shoots based on their coordinates in the 2-dimensional PCA (Principal

Component Analysis) plot using the *dist* function of the R package 'stats' v3.6.0 (R Core Team,

2019). PCA was done on reads-per-million for all shoots with the PCA function of the R package

'FactoMineR' (Lê et al., 2008).

2.9 Correlation between methylome variation and photosynthetic performance

For the 10 heat-stressed samples, we tested for correlations between epigenetic distance and

performance difference (PiABS values) among shoots while controlling for genetic distance with

Partial Mantel tests using the function *partial.mantel.test* of the R package 'ncf' v1.2.9 (Bjornstad,

2020). Only significant correlations (p<0.05, adjusted for multiple comparisons according to

Benjamini and Hochberg (1995) in R (R Core Team, 2019)) with coefficients $R > 0.65$, when

controlled for genetic distance, were considered strong enough to be reckoned as biologically linked.

The same analysis ran Mantel tests for correlation between genetic and epigenetic distance, and

between genetic distance and performance difference.

2.10 Methylome heat stress response

The methylome of the 10 heat-stressed shoots was characterized under control, stress, and recovery

conditions (3 shoots died from the stress), and compared with the methylome of field transect

samples with PCA on reads-per-million data using the PCA function of the R package 'FactoMineR'

(Lê et al., 2008). Epigenetic distance between the samples was estimated for each sequence context

(gene, intergene, TE, each in CG and CHG regions, respectively) as the Euclidean distance in the 2-

dimensional PCA plot using the *dist* function of the R package 'stats' v3.6.9 (R Core Team, 2019).

2.11 Differential methylation analyses

To estimate the number of sites that changed in methylation state from control to stress and recovery

conditions, we applied differential methylation analyses using the R package 'edgeR' 3.20.9

(Robinson et al., 2009) within the 'SARTools' pipeline v1.6.6 (Varet et al., 2016). Read counts were

normalized using a trimmed mean of M-values (TMM) between each pair of samples (Robinson and

Oshlack, 2010). *P*-values, indicating the significance of increased or decreased methylation, were

adjusted according the Benjamini-Hochberg method (Benjamini and Hochberg, 1995).

To identify how the methylome differed between samples of high and low performance, we used

differential methylation analyses using the R package 'edgeR' 3.20.9 (Robinson et al., 2009) within

the 'SARTools' pipeline v1.6.6 (Varet et al., 2016). Read counts were normalized using a trimmed

mean of M-values (TMM) between each pair of samples (Robinson and Oshlack, 2010). Differential

methylation analysis was done only for conditions and sequence contexts where a positive correlation

was found between performance differences and epigenetic distance. For control conditions, we

compared methylation levels in genes (CG regions) between the two samples of highest performance

(79.1, and 13.2) and the two samples of lowest performance (63.1 and 59.2, Supplementary Figure

S1). For recovery conditions, we compared methylation levels in intergenic TEs (CHG regions)

between the two samples of highest performance (79.1 and 17.1) and the two samples of lowest

performance (57.1 and 59.2).

For differentially methylated sites within gene bodies, we tested with Fisher's exact tests for

enrichment of gene ontology terms of biological processes with the R package 'topgo' (Alexa and

Rahnenführer, 2010) using Fisher's exacts tests. GO terms were obtained from the v2.1 *Zostera*

genome annotation from the ORCAE database (Sterck et al., 2012). To reduce redundancy in the

significantly enriched GO terms (*p*-values < 0.05), we calculated 'sim rel' scores (Schlicker et al.,

2006) (Allowed similarity=0.5), based on the *A. thaliana* GO-term database, using the REVIGO web

server (Supek et al., 2011).

3   Results

3.1   Methylome characterization and variation among the clonal transect shoots

On average, 74 million high-quality reads were obtained per sequencing library, ranging from 0.7 to

151 million (Supplementary Table S3). DNA raw reads are accessible from NCBI under BioProject

number PRJNA575339[7] On average, 35% of the high-quality reads mapped to the *in silico* digested

Z. marina genome, and 11% mapped uniquely (annotated reads-per-million in Supplementary File

S4). In total, 144,420 sites were methylated (covered by at least two reads) across all transect shoots.

Across all transect shoots, 84,640 methylated CG sites and 59,780 methylated CHG sites were

detected, which represents 59% and 41% of all methylated sites, respectively (Supplementary Table

S4). About 23% of all methylated sites fell in gene bodies (of which 41% in TEs), and 77% in

intergenic regions (of which 42% in TEs) (Supplementary Figure S2). In gene bodies, 67% of the

methylated sites fell in CG regions, in intergenic regions only 56%.

Based on their multi-loicrosatellite genotype, 41 of 42 shoots sampled along the transect (except

sample 27 in Supplementary table S1) belonged to the same genet (Supplementary Table S5).

Methylome variation between these 41 clonal ramets exceeded the methylome shift induced by heat

stress (Figure 2), and was not higher between shoots of different genets (Figure 3A,B, Supplementary

Table S4). Methylation differences were generally lower in CHG than in CG sequence contexts

(Figure 3A,B). Epigenetic distance was not significantly correlated with physical distance in any

sequence context (Supplementary Table S6).

3.2   Genetic variation among heat-stressed shoots

Based on whole genome sequences of the 10 heat-stressed shoots, we identified 15,508 high-quality

SNPs (Supplementary File S1). That all 10 shoots shared the same heterozygous state in 14,431

(93%) of all SNPs, and shared the same multi-locus microsatellite genotype (Supplementary Table

____

20   [7]Available upon publication

S5), suggests that they were clones having originated from one ancestral zygote (Supplementary

Figure S3) (Yu et al., 2020).

Despite clonal, the 10 shoots differed in 1.079 SNPs resulting from somatic mutations. Based on

these SNPs, euclidean genetic distances ranged from 9 to 35 (frequency distribution in

Supplementary Figure S4), and were not significantly correlated with physical distances between

shoots (R=0.18, p=0.12).

3.3   Link between epigenetic distance and performance changes under heat stress

Photosynthetic performance (PiABS) declined in all 10 shoots under heat stress (Figure 1,

Supplementary Table S7). Seven of the 10 shoots recovered from the heat stress, yet performances

did not reach pre-stress levels. Performances under control and recovery conditions were positively

correlated (adjusted p<0.05, R=0.93, Figure 4).

Genetic distance correlated moderately (4<R<6, adjusted p<0.05) with performance differences in

recovered samples, and with epigenetic distance in some sequence contexts of stressed and recovered

samples (Figure 4). Nevertheless, epigenetic distance correlated strongly (R>0.65, adjusted p<0.05)

with performance differences even after controlling for genetic distance based on 1,079 SNPs: 1)

epigenetic distance among control samples in CG gene body regions correlated with performance

differences prior to stress, and after recovery (stress resilience, Figure 4); epigenetic distance among

recovered samples in CHG regions of intergenic TEs correlated with performance differences after

recovery (Figure 4).

3.4   Methylome heat stress response

Methylation patterns in all sequence contexts changed under heat stress and did not return to but

instead diverged further from control (pre-stress) patterns during the recovery period (Figure 5 for all

sequence contexts combined, annotated reads-per-million in Supplementary File S5, Supplementary

figure S5 showing the methylation shift for the different sequence contexts). More sites became

hyper- than hypo-methylated in response to stress (Figure 6A,D). Methylation levels differed

significantly at 437 sites between control and stress conditions (257 hyper- and 180 hypo-

methylated), at 1788 sites between control and recovery conditions (1141 hyper-, and 647 hypo-

methylated), and only at 39 sites between stress and recovery conditions (18 hyper-, and 21 hypo-
methylated, Supplementary Table S8). After recovery, CG methylation had been more heat-
responsive in gene body regions, and CHG methylation in intergenic regions (Figure 6B,E).

After recovery, methylation increased in gene bodies with functions including DNA transcription and
replication; catabolism of misfolded proteins, gamma-aminobutyric acid (GABA), and
neurotransmitter; as well as amino acid synthesis (Supplementary Table S9). DNA methylation had
decreased in gene bodies with functions including transmembrane transport of cations, ions, and
protons; transport of ammonium and phospholipids; localization of lipids and organelles; exocytosis;
and secretion (Supplementary Table S9).

3.5   Differential methylation between shoots of high and low photosynthetic performance

Recovered shoots of high and low performance presented highly similar methylation patterns. Only
seven CHG sites in intergenic TEs were hyper-methylated in recovery samples of high
photosynthetic performance (Supplementary Table S10). In contrast, 89 CG gene body sites were
hyper-methylated (1 hypo-methylated) in control samples of high photosynthetic performance
(Supplementary Table S10). Enriched biological processes in the 89 hyper-methylated CG sites
included 'light harvesting in photosystem I' and 'protein folding' (Supplementary Table S11).

4   Discussion

Plant genets persisting >1,000 years challenge the positive correlation between genetic variation
retained through recombination, and long-term survival. Our study shows for the first time that
ramets of the same seagrass genet display DNA methylation variation that is associated with
phenotypic variation in the fitness-related traits of photosynthetic performance and heat stress
resilience. Heat responsive methylation changes were memorized for at least five weeks in the
recovered shoots. This supports the hypothesis (Dodd and Douhovnikoff, 2016) that methylation
variation, via variation in gene regulation, compensates potential costs of clonal reproduction (Lynch
et al., 1993; Lynch and Lande, 1998), and contributes to the long-term survival of clonal seagrass
meadows by increasing temporally stable variation in ecologically relevant traits that can not be
simply explained by the underlying somatic genetic variation.

4.1 Methylome variation of functional relevance

The seagrass methylome is flexible and responds directly to environmental change, given its strong change from field transect samples to acclimated control samples within two weeks (Figure 2). Methylome variation among ramets of the same genet resulted either from random epimutations or from microscale variations in the environment because its correlation with geographic distance between shoots was weak. The change in depth of 3 meters and, thus, gradual changes in environmental factors along the sampled transect may not have been extreme enough to impose a result. The disagreement between methylome variation and transect position may also result from recent uprooting and re-settling of some shoots. This was suggested to explain disagreement between genetic similarity of clonal shoots and their transect position in another clonal meadow of *Z. marina* (Yu et al., 2020), and would further explain the absence of correlation between genetic and physical distance in our heat-stressed ramets. In such case, the re-settled shoots would display a methylome shaped by distant environmental conditions. Whether environmentally induced or not, methylome variation in CG gene regions predicted pre- and post-stress performance (Figure 4) and, thus, is likely functionally relevant for thermal acclimation and stress resilience.

Although difficult to prove (Crisp et al., 2016), the correlations between methylome and photosynthetic performance differences appear to be causal, given that samples of high performance showed hyper-methylated CG sites in gene bodies with relevant functions: 'light harvesting in PSI', 'protein folding', 'protein refolding', and 'misfolded or incompletely synthesized protein catabolic process' (Supplementary Table S11). These functions can prime against heat stress when increased methylation is associated with expression of the underlying genes and, thus, with accumulation of protective molecules like heat-shock proteins that are involved in repair and prevention of heat-induced protein damage (Feder and Hofmann, 2002), and that can facilitate a fast stress response. Indeed, while genes are generally repressed by methylated promoter regions (Lisch, 2009), genes are often activated by CG gene body methylation (Zhang et al., 2006; Schmitz et al., 2013; Yang et al., 2014; Dubin et al., 2015; Bewick and Schmitz, 2017; Niederhuth and Schmitz, 2017) that prevents aberrant expression from intragenic promoters (Takuno and Gaut, 2012; Neri et al., 2017). Thus, via association with gene expression, the observed methylome variation can putatively create ecologically relevant phenotype variation.

Although our study was based on clonal ramets that presented identical seven-locus microsatellite

genotypes, the 10 heat-stressed ramets varied genetically at 1,079 SNPs generated by somatic

mutations. Epigenetic differentiation between populations can be tightly linked to genetic

differentiation (Herrera and Bazaga, 2010; Gáspár et al., 2019). Accordingly, trans-acting SNPs

appeared to explain the correlation between gene body methylation and latitude in *Arabidopsis*

(Dubin et al., 2015). In contrast, part of the correlation between DNA methylation with ecological

factors in landscape epigenomic studies on non-model plant species could not be predicted from the

observed underlying patterns of genetic relatedness (Schulz et al., 2014; Foust et al., 2016; Gugger et

al., 2016). Genetic variation could also not explain the association between epigenetic variation and

phenotypic variation (leaf, petiole and functional traits) in natural populations of the perennial herb

Helleborus foetidus (Medrano et al., 2014), and in clonal populations of the introduced clonal herb

Hydrocotyle vulgaris (Wang et al., 2020). In agreement, the correlation between methylome variation

and stress performance/recovery in our study could not be predicted from the underlying genetic

variation. Independent from genetic variation was also the methylome stress-response that provides

potential for temporally stable phenotypic change at time scales unattainable by somatic mutations.

Thus, our results provide a first indication that DNA methylation variation provides a layer of

ecologically relevant, and potentially selectable, phenotypic variation that is independent from

genetic variation in clonal seagrass meadows.

4.2   Potential of the epigenetic heat stress response for heat-hardening

Stress recovery is a critical period during which the degree of resetting versus memory of epigenetic

stress response is determined (Crisp et al., 2016). *Z. marina* memorized much of the methylome

stress response after a 5-week recovery period (Figure 5). The methylation changes from control to

recovery conditions were initiated in the stress phase, and then continued to diverge from control

conditions. This is illustrated by the strong similarities in methylation profiles between the stress and

recovery phase, with an apparently stronger divergence of recovery profiles from the control ones

(Figure 6B,E, Supplementary Table S8). This agrees with increasing divergence of gene transcription

profiles from heat-stress to recovery conditions in a Danish *Z. marina* population (Franssen et al.,

2011). We speculate that gene expression changes, and other molecular mechanisms involved in the

heat-stress response, could have triggered additional methylation changes after the stress was

removed (e.g. Secco et al. 2015) The memory of the epigenetic heat stress response is long enough to

potentially heat-harden the same generation of previously exposed shoots.

In agriculture, hardening/priming of seeds is a long-standing practice to enhance crop resistance to

environmental challenges, including hot, cold, dry, or saline conditions, or pathogen infections

(Ibrahim, 2016; Wojtyla et al., 2016; Pawar and Laware, 2018). Heat-priming has only very recently

been described in seagrass (Nguyen et al., 2020). *Zostera muelleri* and *Posidonia australis* both

performed better under a second heat-wave, in terms of photosynthetic capacity, leaf growth, and

chlorophyll a content, when they had been previously exposed to a first heatwave as compared with

naïve controls (Nguyen et al., 2020). This could explain why no mortality was reported for the

seagrass *Posidoinia oceanica* after intense and log-lasting heat-waves in 2012, 2015, and 2017

(Darmaraki et al., 2019), although it had suffered high mortality rates after the 2006 heatwave

((Marbà and Duarte, 2010), as discussed in Nguyen et al. (2020).

Molecular mechanisms involved in forming a stress memory include stalled RNA polymerase II,

storage of chemical signaling factors, accumulation and phosphorylation of transcription factors, and

epigenetic mechanisms such as microRNAs, histone modifications, and DNA methylation (Iwasaki

and Paszkowski, 2014; Crisp et al., 2016; Hilker et al., 2016; Gallusci et al., 2017; Lämke and

Bäurle, 2017). In heat-primed seagrass, the significant regulation of methylation related genes, in

particular histone methyltransferases, suggests that epigenetic modifications contribute to form a

thermal stress memory (Nguyen et al., 2020). Our study adds evidence that also DNA methylation is

likely involved in this process. While the memory of histone modifications lasts generally not longer

than hours or days (Cedar and Bergman, 2009; Huang et al., 2013; Lämke and Bäurle, 2017; Kumar,

2018), DNA methylation modifications have been shown to be partly transgenerationally stable

under both sexual (Ou et al., 2012; Herman and Sultan, 2016), and asexual reproduction (Raj et al.,

2011; Verhoeven and van Gurp, 2012; González et al., 2017). For example, that DNA methylation

memories can mediate adaptive transgenerational plasticity was shown in the plant *Polygonium*

persicaria, in which offspring demethylation with zebularine removed the adaptive effect of parental

drought in form of longer root systems and greater biomass (Herman and Sultan 2016).

Across mitotically grown generations, epigenetic patterns can be expected to be more faithfully

inherited than across sexual generations, because clonal growth circumvents epigenetic

reprogramming during meiosis and embryogenesis (Hirsch et al., 2012; Douhovnikoff and Dodd,

2014; Verhoeven and Preite, 2014; Crisp et al., 2016; Latzel et al., 2016). CG methylation,

maintained by a homologue of mammalian DNA METHYLTRANSFERASE (MET1) (Law and

Jacobsen, 2010), is likely more stable than CHG methylation, which is maintained by a dynamic

process involving plant-specific CHROMOMETHYLASE and a feed-forward loop involving histone
3 lysine 9 methylation (Law and Jacobsen, 2010). Thus, we propose that memorized changes in CG
gene body methylation (Figure 6B), are the most likely methylation marks that may contribute to
stress priming with potential for transgenerational plasticity across mitotically grown seagrass
generations. Under global warming, however, seagrasses may increasingly investigate in sexual
reproduction (Ruiz et al., 2018; Marín-Guirao et al., 2019) and, thus, may be able to transmit only
methylome-based stress memories that escape epigenetic reprogramming.

Heat-responsive DNA methylation changes in plants appear not to show a consistent trend across
different species, and little is yet known about their functional role (Liu et al., 2015). In *Brassica*
rapa, heat-responsive DNA methylation was shown to be associated with differential expression of
genes involved in heat stress signal transduction and in RNA metabolic processing (Liu et al., 2018).
In *Z. marina*, biological functions affected by stress-responsive methylation changes suggest that the
plants were investing after stress in the breakdown and repair of misfolded proteins, synthesis of new
proteins, growth (DNA replication), and the regulation of transmembrane transport of ions and
protons, and neurotransmitter levels (Supplementary Table S9). Of particular interest is the CG
hyper-methylation and, thus, potentially constitutive upregulation (Zhang et al., 2006; Schmitz et al.,
2013; Yang et al., 2014; Dubin et al., 2015; Niederhuth and Schmitz, 2017) of genes involved in the
catabolism of misfolded proteins (Supplementary Table S9). Increased investment in the breakdown
of heat-denatured proteins (Feder and Hofmann, 2002), may prime/adapt the affected shoots to future
heat stress. This function was also hyper-methylated in control samples of highest performance and
stress resilience (Supplementary Table S11). Experimental removal of DNA methylation, e.g. using
zebularine or 5-Azacytidine (Griffin et al., 2016), or the targeted change of methylation patterns, e.g.
via CRISPR (Xu et al., 2016), will ultimately allow to identify the relationship between heat-
responsive methylation patterns and adaptive phenotypic changes.

4.3 Conclusions and Perspectives

Our study suggests that DNA methylation is functionally relevant for photosynthetic performance
and heat stress resilience, independent from underlying somatic mutations. A portion of the stress-
responsive changes in methylation appears sufficiently stable (Figure 5) to allow for medium to long-
term acclimation/heat hardening in clonal lineages. In seagrass meadows composed of several
genotypes, stress resilience, growth, and associated invertebrate species diversity is enhanced by

genotypic variation (Hughes and Stachowicz, 2004; Reusch et al., 2005; Ehlers et al., 2008). In clonal

meadows, epigenetic variation may play a similar role in the potential to secure function and

resilience not only of *Z. marina* plants, but also of the entire associated ecosystem.

Due to anthropogenic stressors, nearly one-third of global seagrass area has disappeared over the last

100 years, and the rate of loss accelerated from ca. 1% yr^{-1} before 1940 to 7% yr^{-1} since 1990

(Waycott et al., 2009). At the same time, rising temperatures open up new thermally suitable habitat

in the Arctic (Krause-Jensen and Duarte, 2014). How fast and far warm-temperate and subarctic

range edges will move polewards depends on the ability of seagrass to rapidly acclimate and adapt to

rising temperatures and other environmental changes (Duarte et al., 2018). Thus, future studies are

needed to assess the adaptive value and transgenerational stability of the epigenetic stress response,

and to compare the ability to build up epigenetic variation between seagrass meadows of composed

of a single or multiple clones, as well as between range center *versus* edge populations.

The functional role of methylation variation in plant genets is not only of fundamental interest but

also of applied interest for management programs of clonal organisms designed to assess

evolutionary potential and population stability, and to minimize the loss of biodiversity. Our results

can be relevant for restoration of seagrass ecosystems that largely depends on the success of

replanted shoots to overcome natural variability/stress (Suykerbuyk et al., 2016; van Katwijk et al.,

2016). Together with Nguyen et al. (2020) our study suggests that heat priming may improve

restoration success of meadows damaged after heat waves (e.g. Arias-Ortiz et al. 2018). Given that

40% of all plant species can reproduce clonally (Tiffney and Niklas, 1985), our findings are further

important to other fields, such as invasion biology and crop breeding strategies (Bilichak and

Kovalchuk, 2016) of clonal plants.

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest.

6   Acknowledgements

We thank Marvin Choquet from Nord University for help with re-planting the shoots in the wet lab.

We acknowledge Steinar Johnsen from Nord University for setting up the aquaria and water flow in

the wet lab. This work was supported by the Norwegian Research Council (Havkyst project 243916),

the Åbo Akademi University Foundation sr to CB, and a personal research talents grant from Nord

University to AJ. Open access publication fees were covered by Nord University's Opean Access

fund. The content of this manuscript has previously appeared online as a preprint in bioRxiv

(Jueterbock et al., 2019).

593 7   Author Contributions

594 GH (project leader) and AJ were planning the project and designing the experiments. AJ, CB, IS, and

595 GH collected the shoots. AJ analysed the data and wrote the manuscript. AJ, MK, AKSD, JAC, and

596 IS performed the DNA extraction, library preparation, and sequencing. SAH, JLO, and YVdP were

597 involved in data interpretation. All co-authors read and commented on the manuscript.

598 8   References

Alexa, A., and Rahnenführer, J. (2010). Enrichment analysis for Gene Ontology. *R Packag.*

"topGO" v2.16.0. Available at:

http://www.bioconductor.org/packages/2.11/bioc/html/topGO.html.

Arias-Ortiz, A., Serrano, O., Masqué, P., Lavery, P. S., Mueller, U., Kendrick, G. A., et al. (2018). A

marine heatwave drives massive losses from the world's largest seagrass carbon stocks. *Nat.*

Clim. Chang. 8, 1–7. doi:10.1038/s41558-018-0096-y.

Arnaud-Haond, S., Duarte, C. M., Diaz-Almela, E., Marbà, N., Sintes, T., and Serrão, E. A. (2012).

Implications of extreme life span in clonal organisms: Millenary clones in meadows of the

threatened seagrass *Posidonia oceanica*. *PLoS One* 7. doi:10.1371/journal.pone.0030454.

Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., del Angel, G., Levy-Moonshine, A., et al.

(2013). From FastQ Data to High-Confidence Variant Calls: The Genome Analysis Toolkit Best

Practices Pipeline. *Curr. Protoc. Bioinforma.* 43, 11.10.1-11.10.33.

doi:10.1002/0471250953.bi1110s43.

Bailleul, D., Stoeckel, S., and Arnaud-Haond, S. (2016). RClone: a package to identify MultiLocus
Clonal Lineages and handle clonal data sets in r. *Methods Ecol. Evol.* 7, 966–970.
doi:10.1111/2041-210X.12550.

Balmer, A., Pastor, V., Gamir, J., Flors, V., and Mauch-Mani, B. (2015). The "prime-ome": Towards
a holistic approach to priming. *Trends Plant Sci.* 20, 443–452.
doi:10.1016/j.tplants.2015.04.002.

Barrett, S. C. H. (2015). Influences of clonality on plant sexual reproduction. *Proc. Natl. Acad. Sci.*
112, 8859–8866. doi:10.1073/pnas.1501712112.

Becheler, R., Diekmann, O., Hily, C., Moalic, Y., and Arnaud-Haond, S. (2010). The concept of
population in clonal organisms: Mosaics of temporally colonized patches are forming highly
diverse meadows of *Zostera marina* in Brittany. *Mol. Ecol.* 19, 2394–2407. doi:10.1111/j.1365-
294X.2010.04649.x.

Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate - a Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B-
Methodological 1995.pdf. *J. R. Stat. Soc. Ser. B* 57, 289–300. doi:10.2307/2346101.

Bewick, A. J., and Schmitz, R. J. (2017). Gene body DNA methylation in plants. *Curr. Opin. Plant*
Biol. 36, 103–110. doi:10.1016/j.pbi.2016.12.007.

Bijlsma, R., and Loeschcke, V. (2012). Genetic erosion impedes adaptive responses to stressful
environments. *Evol. Appl.* 5, 117–129. doi:10.1111/j.1752-4571.2011.00214.x.

Bilichak, A., Ilnytskyy, Y., Wóycicki, R., Kepeshchuk, N., Fogen, D., and Kovalchuk, I. (2015). The
elucidation of stress memory inheritance in *Brassica rapa* plants. *Front. Plant Sci.* 6, 1–17.
doi:10.3389/fpls.2015.00005.

Bilichak, A., and Kovalchuk, I. (2016). Transgenerational response to stress in plants and its
application for breeding. *J. Exp. Bot.* 67, 2081–2092. doi:10.1093/jxb/erw066.

Bjornstad, O. N. (2020). ncf: Spatial Covariance Functions. Available at:
https://cran.r-project.org/package=ncf.

Bossdorf, O., Richards, C. L., and Pigliucci, M. (2008). Epigenetics for ecologists. *Ecol. Lett.* 11,
106–115. doi:10.1111/j.1461-0248.2007.01130.x.

Boström, C., Baden, S., Bockelmann, A.-C. C., Dromph, K., Fredriksen, S., Gustafsson, C., et al.

(2014). Distribution, structure and function of Nordic eelgrass (*Zostera marina*) ecosystems:

Implications for coastal management and conservation. *Aquat. Conserv. Mar. Freshw. Ecosyst.*

24, 410–434. doi:10.1002/aqc.2424.

Boyko, A., Blevins, T., Yao, Y., Golubov, A., Bilichak, A., Ilnytskyy, Y., et al. (2010).

Transgenerational adaptation of *Arabidopsis* to stress requires DNA methylation and the

function of dicer-like proteins. *PLoS One* 5. doi:10.1371/journal.pone.0009514.

Bussotti, F., Desotgiu, R., Pollastrini, M., and Cascio, C. (2010). The JIP test: a tool to screen the

capacity of plant adaptation to climate change. *Scand. J. For. Res.* 25, 43–50.

doi:10.1080/02827581.2010.485777.

Cedar, H., and Bergman, Y. (2009). Linking DNA methylation and histone modification: Patterns

and paradigms. *Nat. Rev. Genet.* 10, 295–304. doi:10.1038/nrg2540.

Chefaoui, R. M., Duarte, C. M., and Serrão, E. A. (2018). Dramatic loss of seagrass habitat under

projected climate change in the Mediterranean Sea. *Glob. Chang. Biol.* doi:10.1111/gcb.14401.

Costanza, R., Arge, R., Groot, R. De, Farberk, S., Grasso, M., Hannon, B., et al. (1997). The value of

the world's ecosystem services and natural capital. *Nature* 387, 253–260.

doi:10.1038/387253a0.

Crisp, P. A., Ganguly, D., Eichten, S. R., Borevitz, J. O., and Pogson, B. J. (2016). Reconsidering

plant memory: Intersections between stress recovery, RNA turnover, and epigenetics. *Sci. Adv.*

2, e1501340–e1501340. doi:10.1126/sciadv.1501340.

Darmaraki, S., Somot, S., Sevault, F., and Nabat, P. (2019). Past Variability of Mediterranean Sea

Marine Heatwaves. *Geophys. Res. Lett.* 46, 9813–9823. doi:10.1029/2019GL082933.

Dodd, R. S., and Douhovnikoff, V. (2016). Adjusting to Global Change through Clonal Growth and

Epigenetic Variation. *Front. Ecol. Evol.* 4, 1–6. doi:10.3389/fevo.2016.00086.

Douhovnikoff, V., and Dodd, R. S. (2014). Epigenetics: a potential mechanism for clonal plant

success. *Plant Ecol.* 216, 227–233. doi:10.1007/s11258-014-0430-z.

Dowen, R. H., Pelizzola, M., Schmitz, R. J., Lister, R., Dowen, J. M., Nery, J. R., et al. (2012).

Widespread dynamic DNA methylation in response to biotic stress. *Proc. Natl. Acad. Sci. U. S.*

A. 109, E2183-91. doi:10.1073/pnas.1209329109.

Duarte, B., Martins, I., Rosa, R., Matos, A. R., Roleda, M. Y., Reusch, T. B. H., et al. (2018).
Climate change impacts on seagrass meadows and macroalgal forests: an integrative perspective
on acclimation and adaptation potential. *Front. Mar. Sci.* 5, 190.
doi:10.3389/FMARS.2018.00190.

Duarte, C. M., Kennedy, H., Marbà, N., and Hendriks, I. (2013). Assessing the capacity of seagrass
meadows for carbon burial: Current limitations and future strategies. *Ocean Coast. Manag.* 83,
32–38. doi:10.1016/j.ocecoaman.2011.09.001.

Dubin, M. J., Zhang, P., Meng, D., Remigereau, M. S., Osborne, E. J., Casale, F. P., et al. (2015).
DNA methylation in *Arabidopsis* has a genetic basis and shows evidence of local adaptation.
Elife 4, 1–23. doi:10.7554/eLife.05255.

Ehlers, A., Worm, B., and Reusch, T. B. H. (2008). Importance of genetic diversity in eelgrass
Zostera marina for its resilience to global warming. *Mar. Ecol. Prog. Ser.* 355, 1–7.
doi:10.3354/meps07369.

Feder, M. E., and Hofmann, G. E. (2002). Heat-shock proteins, molecular chaperones, and the stress
response: Evolutionary and ecological physiology. *Annu. Rev. Physiol.* 61, 243–282.
doi:10.1146/annurev.physiol.61.1.243.

Feng, S., Cokus, S. J., Zhang, X., Chen, P.-Y. P.-Y., Bostick, M., Goll, M. G., et al. (2010).
Conservation and divergence of methylation patterning in plants and animals. *Proc. Natl. Acad.*
Sci. 107, 8689–8694. doi:10.1073/pnas.1002720107.

Fourqurean, J. W., Duarte, C. M., Kennedy, H., Marbà, N., Holmer, M., Mateo, M. A., et al. (2012).
Seagrass ecosystems as a globally significant carbon stock. *Nat. Geosci.* 5, 505–509.
doi:10.1038/ngeo1477.

Foust, C. M., Preite, V., Schrey, A. W., Alvarez, M., Robertson, M. H., Verhoeven, K. J. F. F., et al.
(2016). Genetic and epigenetic differences associated with environmental gradients in replicate
populations of two salt marsh perennials. *Mol. Ecol.* 25, 1639–1652. doi:10.1111/mec.13522.

Frankham, R., Bradshaw, C. J. A., and Brook, B. W. (2014). Genetics in conservation management:
Revised recommendations for the 50/500 rules, Red List criteria and population viability
analyses. *Biol. Conserv.* 170, 56–63. doi:10.1016/j.biocon.2013.12.036.

Franssen, S. U., Gu, J., Bergmann, N., Winters, G., Klostermeier, U. C., Rosenstiel, P., et al. (2011).
Transcriptomic resilience to global warming in the seagrass *Zostera marina*, a marine
foundation species. *Proc. …* 108, 19276–19281. doi:10.1073/pnas.1107680108.

Gabriel, W., Lynch, M., and Bürger, R. (1993). Muller's Ratchet and Mutational Meltdowns.
Evolution (N. Y). 47, 1744–1757. doi:10.1111/j.1558-5646.1993.tb01266.x.

Gallusci, P., Dai, Z., Génard, M., Gauffretau, A., Leblanc-Fournier, N., Richard-Molard, C., et al.
(2017). Epigenetics for Plant Improvement: Current Knowledge and Modeling Avenues. *Trends*
Plant Sci. 22, 610–623. doi:10.1016/j.tplants.2017.04.009.

Gao, L., Geng, Y., Li, B., Chen, J., and Yang, J. (2010). Genome-wide DNA methylation alterations
of *Alternanthera philoxeroides* in natural and manipulated habitats: implications for epigenetic
regulation of rapid responses to environmental fluctuation and phenotypic variation. *Plant. Cell*
Environ. 33, 1820–1827. doi:10.1111/j.1365-3040.2010.02186.x.

Garrison, E. (2020). C++ library and cmdline tools for parsing and manipulating VCF files. Available
at: https://github.com/vcflib/vcflib.

Gáspár, B., Bossdorf, O., and Durka, W. (2019). Structure, stability and ecological significance of
natural epigenetic variation: a large-scale survey in *Plantago lanceolata*. *New Phytol.* 221,
1585–1596. doi:10.1111/nph.15487.

González, A. P. R., Dumalasová, V., Rosenthal, J., Skuhrovec, J., and Latzel, V. (2017). The role of
transgenerational effects in adaptation of clonal offspring of white clover (*Trifolium repens*) to
drought and herbivory. *Evol. Ecol.* 31, 345–361. doi:10.1007/s10682-016-9844-5.

Greally, J. M. (2018). A user's guide to the ambiguous word "epigenetics." *Nat. Rev. Mol. Cell Biol.*
19, 207–208. doi:10.1038/nrm.2017.135.

Green, E. P., and Short, F. (2004). *World atlas of seagrasses*. Berkeley, USA: University of
California Press doi:10.1515/BOT.2004.029.

Greve, T. M., Borum, J., and Pedersen, O. (2003). Meristematic oxygen variability in eelgrass
(*Zostera marina*). *Limnol. Oceanogr.* 48, 210–216. doi:10.4319/lo.2003.48.1.0210.

Griffin, P. T., Niederhuth, C. E., and Schmitz, R. J. (2016). A comparative analysis of 5-azacytidine-
and zebularine-induced DNA demethylation. *G3 Genes, Genomes, Genet.* 6, 2773–2780.
doi:10.1534/g3.116.030262.

Gugger, P. F., Fitz-Gibbon, S., PellEgrini, M., and Sork, V. L. (2016). Species-wide patterns of DNA
methylation variation in *Quercus lobata* and their association with climate gradients. *Mol. Ecol.*
25, 1665–1680. doi:10.1111/mec.13563.

Herman, J. J., and Sultan, S. E. (2016). DNA methylation mediates genetic variation for adaptive
transgenerational plasticity. *Proc. R. Soc. B Biol. Sci.* 283, 20160988.
doi:10.1098/rspb.2016.0988.

Herrera, C. M., and Bazaga, P. (2010). Epigenetic differentiation and relationship to adaptive genetic
divergence in discrete populations of the violet *Viola cazorlensis*. *New Phytol.* 187, 867–876.
doi:10.1111/j.1469-8137.2010.03298.x.

Hilker, M., Schwachtje, J., Baier, M., Balazadeh, S., Bäurle, I., Geiselhardt, S., et al. (2016). Priming
and memory of stress responses in organisms lacking a nervous system. *Biol. Rev.* 91, 1118–
1133. doi:10.1111/brv.12215.

Hirsch, S., Baumberger, R., and Grossniklaus, U. (2012). Epigenetic variation, inheritance, and
selection in plant populations. *Cold Spring Harb. Symp. Quant. Biol.* 77, 97–104.
doi:10.1101/sqb.2013.77.014605.

Huang, C., Xu, M., and Zhu, B. (2013). Epigenetic inheritance mediated by histone lysine
methylation: Maintaining transcriptional states without the precise restoration of marks? *Philos.*
Trans. R. Soc. B Biol. Sci. 368. doi:10.1098/rstb.2011.0332.

Hughes, A. R., Inouye, B. D., Johnson, M. T. J., Underwood, N., and Vellend, M. (2008). Ecological
consequences of genetic diversity. *Ecol. Lett.* 11, 609–623. doi:10.1111/j.1461-
0248.2008.01179.x.

Hughes, A. R., and Stachowicz, J. J. (2004). Genetic diversity enhances the resistance of a seagrass
ecosystem to disturbance. *Proc. Natl. Acad. Sci.* 101, 8998–9002.
doi:10.1073/pnas.0402642101.

Ibrahim, E. A. (2016). Seed priming to alleviate salinity stress in germinating seeds. *J. Plant Physiol.*
192, 38–46. doi:10.1016/j.jplph.2015.12.011.

Iwasaki, M., and Paszkowski, J. (2014). Epigenetic memory in plants. *EMBO J.* 33, 1987–1998.
doi:10.15252/embj.201488883.

Jablonka, E., and Raz, G. (2009). Transgenerational Epigenetic Inheritance: Prevalence,
Mechanisms, and Implications for the Study of Heredity and Evolution. *Q. Rev. Biol.* 84, 131–
176. doi:10.1086/598822.

Jueterbock, A., Boström, C., James, A. C., Olsen, J., Kopp, M., Dhanasiri, A., et al. (2019).
Methylation variation promotes phenotypic diversity and evolutionary potential in a millenium-
old clonal seagrass meadow. *bioRxiv,* 787754. doi:10.1101/787754.

Jueterbock, A., Franssen, S. U., Bergmann, N., Gu, J., Coyer, J. A., Reusch, T. B. H. H., et al. (2016).
Phylogeographic differentiation versus transcriptomic adaptation to warm temperatures in
Zostera marina, a globally important seagrass. *Mol. Ecol.* 25, 5396–5411.
doi:10.1111/mec.13829.

Kilvitis, H. J., Alvarez, M., Foust, C. M., Schrey, A. W., Robertson, M., and Richards, C. L. (2014).
"Ecological Epigenetics," in *Ecological Genomics: Ecology and the Evolution of Genes and*
Genomes Advances in Experimental Medicine and Biology., eds. C. R. Landry and N. Aubin-
Horth (Springer Netherlands), 191–210. doi:10.1007/978-94-007-7347-9.

Knaus, B. J., and Grünwald, N. J. (2017). <scp>vcfr</scp> : a package to manipulate and visualize
variant call format data in R. *Mol. Ecol. Resour.* 17, 44–53. doi:10.1111/1755-0998.12549.

Krause-Jensen, D., and Duarte, C. M. (2014). Expansion of vegetated coastal ecosystems in the
future Arctic. *Front. Mar. Sci.* 1, 1–10. doi:10.3389/fmars.2014.00077.

Kumar, S. (2018). Epigenomics of Plant Responses to Environmental Stress. *Epigenomes* 2, 6.
doi:10.3390/epigenomes2010006.

Kuźnicki, D., Meller, B., Arasimowicz-Jelonek, M., Braszewska-Zalewska, A., Drozda, A., and
Floryszak-Wieczorek, J. (2019). BABA-induced DNA methylome adjustment to
intergenerational defense priming in potato to *Phytophthora* infestans. *Front. Plant Sci.* 10, 1–
16. doi:10.3389/fpls.2019.00650.

Laikre, L., Allendorf, F. W., Aroner, L. C., Baker, C. S., Gregovich, D. P., Hansen, M. M., et al.
(2009). Neglect of genetic diversity in implementation of the Convention on Biological
Diversity. *Conserv. Biol.* 24, 86–88. Available at: https://www.treesearch.fs.fed.us/pubs/37018.

Lämke, J., and Bäurle, I. (2017). Epigenetic and chromatin-based mechanisms in environmental
stress adaptation and stress memory in plants. *Genome Biol.* 18, 1–11. doi:10.1186/s13059-017-
1263-6.

Lande, R. (1988). Genetics and demography in biological conservation. *Science (80-.).* 241, 1455–
1460. doi:10.1126/science.3420403.

Larkum, A. W. D. D., Orth, R. J., and Duarte, C. M. (2006). *Seagrasses : Biology, Ecology, and*
Conservation. , eds. A. W. D. D. Larkum, R. J. Orth, and C. M. Duarte Springer Netherlands
doi:10.1111/j.1439-0485.2006.00138.x.

Latzel, V., Allan, E., Bortolini Silveira, A., Colot, V., Fischer, M., and Bossdorf, O. (2013).
Epigenetic diversity increases the productivity and stability of plant populations. *Nat. Commun.*
4, 2875. doi:10.1038/ncomms3875.

Latzel, V., Rendina González, A. P., and Rosenthal, J. (2016). Epigenetic Memory as a Basis for
Intelligent Behavior in Clonal Plants. *Front. Plant Sci.* 7, 1354. doi:10.3389/fpls.2016.01354.

Law, J. A., and Jacobsen, S. E. (2010). Establishing, maintaining and modifying DNA methylation
patterns in plants and animals. *Nat. Rev. Genet.* 11, 204–220. doi:10.1038/nrg2719.

Lê, S., Josse, J., and Husson, F. (2008). FactoMineR: An R Package for Multivariate Analysis. *J.*
Stat. Softw. 25, 1–18. doi:10.1016/j.envint.2008.06.007.

Li, H., and Durbin, R. (2009). Fast and accurate short-read alignment with Burrows-Wheeler
transform. *Bioinformatics* 25, 1754–1760. doi:10.1093/bioinformatics/btp324.

Li, R., Li, Y., Kristiansen, K., and Wang, J. (2008). SOAP: Short oligonucleotide alignment program.
Bioinformatics 24, 713–714. doi:10.1093/bioinformatics/btn025.

Lisch, D. (2009). Epigenetic Regulation of Transposable Elements in Plants. *Annu. Rev. Plant Biol.*
60, 43–66. doi:10.1146/annurev.arplant.59.032607.092744.

Liu, F., Liu, J., and Dong, M. (2016). Ecological Consequences of Clonal Integration in Plants.
Front. Plant Sci. 7, 1–11. doi:10.3389/fpls.2016.00770.

Liu, G., Xia, Y., Liu, T., Dai, S., and Hou, X. (2018). The DNA methylome and association of
differentially methylated regions with differential gene expression during heat stress in *Brassica*
rapa. Int. J. Mol. Sci. 19. doi:10.3390/ijms19051414.

Liu, J., Feng, L., Li, J., and He, Z. (2015). Genetic and epigenetic control of plant heat responses.
Front. Plant Sci. 06, 1–21. doi:10.3389/fpls.2015.00267.

Loxdale, H. D., Lushai, G., Aafc, A. C., Cereal, E., and Ecorc, C. (2003). Intraclonal genetic
variation : ecological and evolutionary aspects. Rapid changes in clonal lines : the death of a
'sacred cow. *Philos. Perspect.*, 3–16.

Lushai, G., Loxdale, H. D., and Allen, J. A. (2003). Intraclonal genetic variation : ecological and
evolutionary aspects . The dynamic clonal genome and its adaptive potential. *Society*, 193–208.

Lynch, M., Butcher, D., Bürger, R., and Gabriel, W. (1993). The mutational meltdown in asexual
populations. *J. Hered.* 84, 339–344. doi:10.1093/oxfordjournals.jhered.a111354.

Lynch, M., and Lande, R. (1998). The critical effective size for a genetically secure population.
Anim. Conserv. 1, 70–72. doi:doi:10.1017/S136794309822110X.

Marbà, N., and Duarte, C. M. (2010). Mediterranean warming triggers seagrass (*Posidonia oceanica*)
shoot mortality. *Glob. Chang. Biol.* 16, 2366–2375. doi:10.1111/j.1365-2486.2009.02130.x.

Marín-Guirao, L., Entrambasaguas, L., Ruiz, J. M., and Procaccini, G. (2019). Heat-stress induced
flowering can be a potential adaptive response to ocean warming for the iconic seagrass
Posidonia oceanica. *Mol. Ecol.* 28, 2486–2501. doi:10.1111/mec.15089.

May, M. R., Provance, M. C., Sanders, A. C., Ellstrand, N. C., and Ross-Ibarra, J. (2009). A
pleistocene clone of palmer's oak persisting in Southern California. *PLoS One* 4, 1–5.
doi:10.1371/journal.pone.0008346.

McKey, D., Elias, M., Pujol, M. E., and Duputié, A. (2010). The evolutionary ecology of clonally
propagated domesticated plants. *New Phytol.* 186, 318–332. doi:10.1111/j.1469-
8137.2010.03210.x.

Medrano, M. M., Herrera, C. M., and Bazaga, P. (2014). Epigenetic variation predicts regional and
local intraspecific functional diversity in a perennial herb. *Mol. Ecol.* 23, 4926–4938.
doi:10.1111/mec.12911.

Muller, H. J. (1964). The relation of recombination to mutational advance. *Mutat. Res. - Fundam.*
Mol. Mech. Mutagen. 1, 2–9. doi:10.1016/0027-5107(64)90047-8.

Neri, F., Rapelli, S., Krepelova, A., Incarnato, D., Parlato, C., Basile, G., et al. (2017). Intragenic
DNA methylation prevents spurious transcription initiation. *Nature* 543, 72–77.
doi:10.1038/nature21373.

Nguyen, H. M., Kim, M., Ralph, P. J., Marín-Guirao, L., Pernice, M., and Procaccini, G. (2020).
Stress memory in seagrasses: first insight into the effects of thermal priming and the role of
epigenetic modifications. *Front. Plant Sci.* 11, 494. doi:10.3389/FPLS.2020.00494.

Nicotra, A. B., Atkin, O. K., Bonser, S. P., Davidson, A. M., Finnegan, E. J., Mathesius, U., et al.
(2010). Plant phenotypic plasticity in a changing climate. *Trends Plant Sci.* 15, 684–692.
doi:10.1016/j.tplants.2010.09.008.

Niederhuth, C. E., and Schmitz, R. J. (2017). Putting DNA methylation in context: from genomes to
gene expression in plants. *Biochim. Biophys. Acta - Gene Regul. Mech.* 1860, 149–156.
doi:10.1016/j.bbagrm.2016.08.009.

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., et al. (2019). vegan:
Community Ecology Package. Available at: https://cran.r-project.org/package=vegan.

Olsen, J. L., Rouzé, P., Verhelst, B., Lin, Y., Bayer, T., Collen, J., et al. (2016). The genome of the
seagrass *Zostera marina* reveals angiosperm adaptation to the sea. *Nature* 530, 331–335.
doi:10.1038/nature16548.

Olsen, J. L., Stam, W. T., Coyer, J. A., Reusch, T. B. H. H., Billingham, M., Boström, C., et al.
(2004). North Atlantic phylogeography and large-scale population differentiation of the seagrass
Zostera marina L. *Mol. Ecol.* 13, 1923–1941. doi:10.1111/j.1365-294X.2004.02205.x.

Orth, R. J., Carruthers, T. J. B., Dennison, W. C., Duarte, C. M., Fourqurean, J. W., Heck, K. L., et
al. (2006). A Global Crisis for Seagrass Ecosystems. *Bioscience* 56, 987–996.
doi:10.1641/0006-3568(2006)56[987:agcfse]2.0.co;2.

Ou, X., Zhang, Y., Xu, C., Lin, X., Zang, Q., Zhuang, T., et al. (2012). Transgenerational Inheritance
of Modified DNA Methylation Patterns and Enhanced Tolerance Induced by Heavy Metal
Stress in Rice (*Oryza sativa* L.). *PLoS One* 7. doi:10.1371/journal.pone.0041143.

Pawar, V. A., and Laware, S. L. (2018). Seed Priming A Critical Review. *Int. J. Sci. Res. Biol. Sci.* 5,
94–101. doi:10.26438/ijsrbs/v5i5.94101.

Procaccini, G., Olsen, J. L., and Reusch, T. B. H. (2007). Contribution of genetics and genomics to
seagrass biology and conservation. *J. Exp. Mar. Bio. Ecol.* 350, 234–259.
doi:10.1016/j.jembe.2007.05.035.

Ptashne, M. (2007). On the use of the word "epigenetic." *Curr. Biol.* 17, 233–236.
doi:10.1016/j.cub.2007.02.030.

R Core Team (2019). R: A Language and Environment for Statistical Computing. Available at:
https://www.r-project.org/.

Raj, S., Bräutigam, K., Hamanishi, E. T., Wilkins, O., Thomas, B. R., Schroeder, W., et al. (2011).
Clone history shapes *Populus* drought responses. *Proc. Natl. Acad. Sci. U. S. A.* 108, 12521–
12526. doi:10.1073/pnas.1103341108.

Rasmann, S., De Vos, M., Casteel, C. L., Tian, D., Halitschke, R., Sun, J. Y., et al. (2012). Herbivory
in the Previous Generation Primes Plants for Enhanced Insect Resistance. *Plant Physiol.* 158,
854–863. doi:10.1104/pp.111.187831.

Repolho, T., Duarte, B., Dionísio, G., Paula, J. R., Lopes, A. R., Rosa, I. C., et al. (2017). Seagrass
ecophysiological performance under ocean warming and acidification. *Sci. Rep.* 7, 41443.
doi:10.1038/srep41443.

Reusch, T. B. H., and Boström, C. (2011). Widespread genetic mosaicism in the marine angiosperm
Zostera marina is correlated with clonal reproduction. *Evol. Ecol.* 25, 899–913.
doi:10.1007/s10682-010-9436-8.

Reusch, T. B. H., Boström, C., Stam, W. T., and Olsen, J. L. (1999). An ancient eelgrass clone in the
Baltic. *Mar. Ecol. Prog. Ser.* 183, 301–304. doi:10.3354/meps183301.

Reusch, T. B. H., Dierking, J., Andersson, H. C., Bonsdorff, E., Carstensen, J., Casini, M., et al.
(2018). The Baltic Sea as a time machine for the future coastal ocean. *Sci. Adv.* 4, eaar8195.
doi:10.1126/sciadv.aar8195.

Reusch, T. B. H., Ehlers, A., Hammerli, A., and Worm, B. (2005). Ecosystem recovery after climatic
extremes enhanced by genotypic diversity. *Proc. Natl. Acad. Sci.* 102, 2826–2831. doi:10.1073/
pnas.0500008102.

Reusch, T. B. H., and Hughes, A. R. (2006). The Emerging Role of Genetic Diversity for Ecosystem
Functioning : Estuarine Macrophytes as Models. *Estuaries and Coasts* 29, 159–164.
doi:10.1007/BF02784707.

Reusch, T. B. H., Stam, W. T., and Olsen J.L. (2000). A microsatellite-based estimation of clonal
diversity and population subdivision in *Zostera marina*, a marine flowing plant. *Mol. Ecol.* 9,
127–140.

Richards, C. L., Schrey, A. W., and Pigliucci, M. (2012). Invasion of diverse habitats by few
Japanese knotweed genotypes is correlated with epigenetic differentiation. *Ecol. Lett.* 15, 1016–
1025. doi:10.1111/j.1461-0248.2012.01824.x.

Robinson, M. D., McCarthy, D. J., and Smyth, G. K. (2009). edgeR: A Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* 26, 139–140.
doi:10.1093/bioinformatics/btp616.

Robinson, and Oshlack (2010). A scaling normalization method for differential expression analysis
of RNA-seq data. *Genome Biol.* 11, R25. Available at:
http://genomebiology.com/2010/11/3/R25.

Ruiz, J. M., Marín-Guirao, L., García-Muñoz, R., Ramos-Segura, A., Bernardeau-Esteller, J., Pérez,
M., et al. (2018). Experimental evidence of warming-induced flowering in the Mediterranean
seagrass *Posidonia oceanica*. *Mar. Pollut. Bull.* 134, 49–54.
doi:10.1016/j.marpolbul.2017.10.037.

Schlicker, A., Domingues, F., Rahnenfuhrer, J., and Lengauer, T. (2006). A new measure for
functional similarity of gene products based on Gene Ontology. *BMC Bioinformatics* 7, 302.

Schmitz, R. J., Schultz, M. D., Lewsey, M. G., O'Malley, R. C., Urich, M. a, Libiger, O., et al.
(2011). Transgenerational epigenetic instability is a source of novel methylation variants.
Science 334, 369–73. doi:10.1126/science.1212959.

Schmitz, R. J., Schultz, M. D., Urich, M. A., Nery, J. R., Pelizzola, M., Libiger, O., et al. (2013).
Patterns of population epigenomic diversity. *Nature* 495, 193–8. doi:10.1038/nature11968.

Schulz, B., Eckstein, R. L., and Durka, W. (2014). Epigenetic variation reflects dynamic habitat
conditions in a rare floodplain herb. *Mol. Ecol.* 23, 3523–3537. doi:10.1111/mec.12835.

Secco, D., Wang, C., Shou, H., Schultz, M. D., Chiarenza, S., Nussaume, L., et al. (2015). Stress
induced gene expression drives transient DNA methylation changes at adjacent repetitive
elements. *Elife* 4. doi:10.7554/eLife.09343.001.

Seymour, D. K., Koenig, D., Hagmann, J., Becker, C., and Weigel, D. (2014). Evolution of DNA
Methylation Patterns in the Brassicaceae is Driven by Differences in Genome Organization.
PLoS Genet. 10, e1004785. doi:10.1371/journal.pgen.1004785.

Silvertown, J. (2008). The Evolutionary Maintenance of Sexual Reproduction: Evidence from the
Ecological Distribution of Asexual Reproduction in Clonal Plants. *Int. J. Plant Sci.* 169, 157–
168. doi:10.1086/523357.

Slaughter, A., Daniel, X., Flors, V., Luna, E., Hohn, B., and Mauch-Mani, B. (2012). Descendants of
primed *Arabidopsis plants* exhibit resistance to biotic stress. *Plant Physiol.* 158, 835–843.
doi:10.1104/pp.111.191593.

Spielman, D., Brook, B. W., and Frankham, R. (2004). Most species are not driven to extinction
before genetic factors impact them. *Proc. Natl. Acad. Sci. U. S. A.* 101, 15261–15264.
doi:10.1073/pnas.0403809101.

Stefanov, D., Petkova, V., and Denev, I. D. (2011). Screening for heat tolerance in common bean
(*Phaseolus vulgaris* L.) lines and cultivars using JIP-test. *Sci. Hortic. (Amsterdam).* 128, 1–6.
doi:DOI: 10. 1016/j. scienta. 2010. 12. 003.

Sterck, L., Billiau, K., Abeel, T., Rouzé, P., and Van de Peer, Y. (2012). ORCAE: online resource for
community annotation of eukaryotes. *Nat. Methods* 9, 1041.
doi:http://www.nature.com/nmeth/journal/v9/n11/abs/nmeth.2242.html#supplementary-
information.

Strasser, R. J., Srivastava, A., and Tsimilli-Michael, M. (2000). "The fluorescence transient as a tool
to characterize and screen photosynthetic samples," in *Probing Photosynthesis: Mechanism,*
Regulation & Adaptation, eds. M. Yunu, U. Pathre, and P. Mohanty (Taylor and Francis), 445–
483. Available at: http://ww.hansatech-instruments.com/docs/the fluorescence transient.pdf.

Supek, F., Bošnjak, M., Škunca, N., and Šmuc, T. (2011). Revigo summarizes and visualizes long
lists of gene ontology terms. *PLoS One* 6, e21800. doi:10.1371/journal.pone.0021800.

Suykerbuyk, W., Govers, L. L., Bouma, T. J., Giesen, W. B. J. T., de Jong, D. J., van de Voort, R., et
al. (2016). Unpredictability in seagrass restoration: analysing the role of positive feedback and
environmental stress on *Zostera noltii* transplants. *J. Appl. Ecol.* 53, 774–784.
doi:10.1111/1365-2664.12614.

Takuno, S., and Gaut, B. S. (2012). Body-methylated genes in *Arabidopsis thaliana* are functionally
important and evolve slowly. *Mol. Biol. Evol.* 29, 219–227. doi:10.1093/molbev/msr188.

Tiffney, B., and Niklas, K. (1985). "Clonal growth in land plants - a paleobotanical perspective," in
Population biology and evolution of clonal organisms, ed. C. R. Jackson JBC, Buss LW (Yale
University Press, New Haven), 35–66.

van der Graaf, A., Wardenaar, R., Neumann, D. A., Taudt, A., Shaw, R. G., Jansen, R. C., et al.
(2015). Rate, spectrum, and evolutionary dynamics of spontaneous epimutations. *Proc. Natl.*
Acad. Sci. U. S. A. 112, 6676–81. doi:10.1073/pnas.1424254112.

van Katwijk, M. M., Thorhaug, A., Marbà, N., Orth, R. J., Duarte, C. M., Kendrick, G. A., et al.
(2016). Global analysis of seagrass restoration: The importance of large-scale planting. *J. Appl.*
Ecol. 53, 567–578. doi:10.1111/1365-2664.12562.

Vanden Broeck, A., Cox, K., Brys, R., Castiglione, S., Cicatelli, A., Guarino, F., et al. (2018).
Variability in DNA Methylation and Generational Plasticity in the Lombardy Poplar, a Single
Genotype Worldwide Distributed Since the Eighteenth Century. *Front. Plant Sci.* 9, 1–11.
doi:10.3389/fpls.2018.01635.

Varet, H., Brillet-Guéguen, L., Coppée, J. Y., and Dillies, M. A. (2016). SARTools: A DESeq2- and
edgeR-based R pipeline for comprehensive differential analysis of RNA-Seq data. *PLoS One* 11,
1–8. doi:10.1371/journal.pone.0157022.

Verhoeven, K. J. F., Jansen, J. J., van Dijk, P. J., and Biere, A. (2010). Stress-induced DNA
methylation changes and their heritability in asexual dandelions. *New Phytol.* 185, 1108–1118.
doi:10.1111/j.1469-8137.2009.03121.x.

Verhoeven, K. J. F., and Preite, V. (2014). Epigenetic variation in asexually reproducing organisms.
Evolution (N. Y). 68, 644–655. doi:10.1111/evo.12320.

Verhoeven, K. J. F., and van Gurp, T. P. (2012). Transgenerational effects of stress exposure on
offspring phenotypes in apomictic dandelion. *PLoS One* 7, e38605.
doi:10.1371/journal.pone.0038605.

Verhoeven, K. J. F., Verbon, E. H., van Gurp, T. P., Oplaat, C., Ferreira de Carvalho, J., Morse, A.
M., et al. (2018). Intergenerational environmental effects: functional signals in offspring
transcriptomes and metabolomes after parental jasmonic acid treatment in apomictic dandelion.
New Phytol. 217, 871–882. doi:10.1111/nph.14835.

Wang, M. Z., Li, H. L., Li, J. M., and Yu, F. H. (2020). Correlations between genetic, epigenetic and
phenotypic variation of an introduced clonal herb. *Heredity (Edinb).* 124, 146–155. doi:10.1038/
s41437-019-0261-8.

Wang, S., Lv, J., Zhang, L., Dou, J., Sun, Y., Li, X., et al. (2015). MethylRAD: a simple and scalable
method for genome-wide DNA methylation profiling using methylation-dependent restriction
enzymes. *Open Biol.* 5, 150130. doi:10.1098/rsob.150130.

Waycott, M., Duarte, C. M., Carruthers, T. J. B., Orth, R. J., Dennison, W. C., Olyarnik, S., et al.
(2009). Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proc.*
Natl. Acad. Sci. U. S. A. 106, 12377–12381. doi:10.1073/pnas.0905620106.

Whitham, T. G., and Slobodchikoff, C. N. (1981). Evolution by individuals, plant-herbivore
interactions, and mosaics of genetic variability: The adaptive significance of somatic mutations
in plants. *Oecologia* 49, 287–292. doi:10.1007/BF00347587.

Wickham, H. (2016). *ggplot2 Elegant Graphics for Data Analysis*. Springer New York
doi:10.1007/978-0-387-98141-3.

Williams, B. P., and Gehring, M. (2017). Stable transgenerational epigenetic inheritance requires a
DNA methylation-sensing circuit. *Nat. Commun.* 8. doi:10.1038/s41467-017-02219-3.

Wojtyla, Ł., Lechowska, K., Kubala, S., and Garnczarska, M. (2016). Molecular processes induced in
primed seeds—increasing the potential to stabilize crop yields under drought conditions. *J.*
Plant Physiol. 203, 116–126. doi:10.1016/j.jplph.2016.04.008.

Xu, X., Tao, Y., Gao, X., Zhang, L., Li, X., Zou, W., et al. (2016). A CRISPR-based approach for
targeted DNA demethylation. *Cell Discov.* 2. doi:10.1038/celldisc.2016.9.

Yang, X., Han, H., DeCarvalho, D. D., Lay, F. D., Jones, P. A., and Liang, G. (2014). Gene body
methylation can alter gene expression and is a therapeutic target in cancer. *Cancer Cell* 26, 577–
590. doi:10.1016/j.ccr.2014.07.028.

Yu, L., Boström, C., Franzenburg, S., Bayer, T., Dagan, T., and Reusch, T. B. H. (2020). Somatic
genetic drift and multilevel selection in a clonal seagrass. *Nat. Ecol. Evol.* 3, 833335.
doi:10.1038/s41559-020-1196-4.

Zhang, H., Lang, Z., and Zhu, J. K. (2018). Dynamics and function of DNA methylation in plants.
Nat. Rev. Mol. Cell Biol. 19, 489–506. doi:10.1038/s41580-018-0016-z.

Zhang, X., Yazaki, J., Sundaresan, A., Cokus, S., Chan, S. W. L., Chen, H., et al. (2006). Genome-
wide High-Resolution Mapping and Functional Analysis of DNA Methylation in *Arabidopsis*.
Cell 126, 1189–1201. doi:10.1016/j.cell.2006.08.003.

Zhang, Y.-Y. Y., Fischer, M., Colot, V., and Bossdorf, O. (2013). Epigenetic variation creates
potential for evolution of plant phenotypic plasticity. *New Phytol.* 197, 314–322.
doi:10.1111/nph.12010.

Živčák, M., Brestič, M., Olšovská, K., and Slamka, P. (2008). Performance index as a sensitive
indicator of water stress in *Triticum aestivum* L. *Plant, Soil Environ.* 54, 133–139.

9   Figure captions

Figure 1: Baltic Sea sampling site of a > 1,000 years old clonal meadow where 42 *Zostera marina*
shoots were collected along a 250 m transect.

Figure 2: Methylation variation of field transect samples of *Zostera marina* exceeded the shift in
methylation patterns along the first two principal components in response to heat stress. The samples
are plotted along the first two principle components (Dim) based on methylation profiles across all
sequence contexts. Circles represent 95% confidence intervals around group means. Bracketed
numbers represent the percentage of explained variation.

Figure 3: Genetic relatedness did not restrict epigenetic differentiation. Bivariate methylation
differences in (A) CG and (B) CHG regions were comparable within ramets of the same genet
(intraclone) and between ramets of different genets (interclone) of *Zostera marina* across all
sequence contexts.

Figure 4: Correlation matrix of photosynthetic performance (PiABS) difference (first three columns)
or of genetic distance (last column) among clonal *Zostera marina* shoots with epigenetic distance
(first 24 rows) or with performance difference at control, stress, or recovery conditions (last three
rows). All correlations between performance difference and epigenetic distance were controlled for
genetic distance. Black squares represent untested correlations that were considered biologically not
meaningful. Pearson product-moment correlation coefficients R are encoded by the color gradient
explained in the bar on the right end, and by the shape of the ellipses. Narrow ellipses represent

stronger correlations as compared with wide ones. Asterisks highlight strong (R>0.65) and significant

(adjusted p<0.05) correlations. CG and CHG: sequence contexts of the methylated cytosine; TE:

Transposable element.

Figure 5: Methylation patterns of all sequence contexts in shoots of a *Zostera marina* clone changed

in response to heat stress and did not return to pre-stress patterns after a >5-week recovery period.

See Supplementary Figure S5 for methylation shifts separated by sequence context. The samples are

plotted along the first two principle components (Dim) based on methylation profiles across all

sequence contexts. Circles represent 95% confidence intervals around group means. Bracketed

numbers represent the percentage of explained variation.

Figure 6: Hyper-methylation dominates the stress response of *Zostera marina*. Hyper- and

hypomethylated sites in CG (A-C) and CHG (D-F) regions in stressed vs. control samples (A,D),

recovered vs. control samples (B,E), and in recovered vs. stressed samples (C,F). Stress-responsive

methylation changes were memorized over the >5-week recovery period. TE: transposable element.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6