# Distinct molecular processes in placentae involved in two major subtypes of preeclampsia

Zhonglu Ren[1,#], Yunfei Gao[1,#], Yue Gao[1,2,3], Qian Chen[1], Sijia Jiang[1], Guanmei Liang[1,2,3], Xiaoxue Yang[1], Cuixia Fan[1], Haizhen Wang[1], Jing Wang[1], Yi-Wu Shi[4], Chaoqun Xiao[1], Mei Zhong[1], Yanhong Yu[1*] and Xinping Yang[1,2,3,5*]

1 Department of Obstetrics and Gynecology, Nanfang Hospital, Southern Medical University, Guangzhou, 510515, China

2 Key Laboratory of Mental Health of the Ministry of Education, Southern Medical University, Guangzhou, 510515, China

3 Department of Bioinformatics, School of Basic Medical Sciences, Southern Medical University, Guangzhou, 510515, China

4 Institute of Neuroscience and Department of Neurology, The Second Affiliated Hospital of Guangzhou Medical University, Guangzhou, 510260, China

5 Lead Contact

# These authors contribute equally to this work

* To whom correspondence should be addressed: xpyang1@smu.edu.cn (X.Y.), yuyh1010@hotmail.com (Y.Y.)

**Running title: Molecular processes in preeclampsia**

Total character: 12098

# Abstract

Patients with preeclampsia display a spectrum of onset time and severity of clinical presentation, yet the underlying molecular bases for the early-onset and late-onset clinical subtypes are not known. Since the root cause of PE is thought to be located in the placentae, we carried out RNA-seq on 65 high-quality placenta samples, including 33 from 30 patients and 32 from 30 control subjects, to search for molecular features. We identified two functionally distinct sets of dysregulated genes in two major subtypes: metabolism-related genes, notably transporter genes, in early-onset severe preeclampsia and immune-related genes in late-onset severe preeclampsia, while the late-onset mild preeclampsia could not be distinguished from normal controls. A small number of dysregulated transcription factors may drive the widespread gene dysregulation in both early-onset and late-onset patients. These results suggest that early-onset and late-onset severe preeclampsia have different molecular mechanisms, whereas the late-onset mild preeclampsia may have no placenta-specific causal factors.

# Introduction

Preeclampsia (PE) is hypertension presenting after 20 weeks of gestation combined with signs of damage to other maternal organs, mostly liver and kidney. The diagnostic criteria include systolic blood pressure higher than 140 mm Hg, and/or diastolic blood pressure above 90 mm Hg, in addition to proteinuria of over 300 mg/day (Mol et al, 2016). It can be classified as early- or late-onset with a gestational age cutoff of 34 weeks (Herzog et al, 2017; Raymond & Peterson, 2011). The global incidence rate is 4.6% of pregnancies (Abalos et al, 2013), leading to 60,000 maternal deaths per year (Logue et al, 2016). The grave consequences of PE on maternal health are mainly attributed to cerebral edema, intracranial hemorrhage and eclampsia (2011; Zeeman, 2009), which some studies have suggested to be caused by abnormal functions of the autonomic nervous system mediated by placenta-derived factors (Eneroth-Grimfors et al, 1994; Melchiorre et al, 2014; Voss et al, 2006). PE has significant consequences on fetal development and growth, often leading to perinatal and infant morbidity or mortality (Mol et al, 2016) and contributes preterm births (Davies et al, 2016) and ~15% cases of fetal growth restriction (FGR) (Mitani et al, 2009; Weiler et al, 2011). In the long term, PE affects brain development and functions of the offspring(Liu et al, 2016), leading to intellectual disability (Griffith et al, 2011), epilepsy (Wu et al, 2008), autism (Dachew et al, 2018b; Trosse et al, 2007; Walker et al, 2015) and schizophrenia (Dalman et al, 1999; Ursini et al, 2018).

Since the symptoms of a pregnant woman with PE usually resolve after delivery and histopathological placental changes can be detected at that time, the placenta has been perceived as the root cause of PE. A widely accepted disease model for the placental origin includes two stages of PE development (Roberts & Gammill, 2005; Roberts & Hubel, 2009). During the first stage, abnormal placentation in early gestation plays a role, such as poor trophoblast invasion and incomplete vascular formation of spiral arteries, which lead to placenta dysfunctions; and in the second stage, the dysfunctional placenta releases factors into the maternal blood, which in turn lead to hypertension and organ damage (Roberts & Gammill, 2005; Roberts & Hubel, 2009). This model

promoted intensive studies on placentae as root cause of PE, and increasing knowledge leads to finer classification of stages (Redman, 2014). However, the PE-associated molecular processes originating in the placenta remain largely unknown, and there are insufficient molecular criteria for distinction of the clinical subtypes.

Some PE cases occur in families, suggesting genetic factors play a role in its pathogenesis. First-degree relatives of women with PE have a 5-fold higher risk of developing the disease (Morgan & Ward, 1999). Men born from a PE pregnancy are also at risk of fathering PE pregnancies (1998; Shiho Nagayama et al, 2014), demonstrating the placental genotype contributes to PE. The heritability of PE is about 50% (Cnattingius et al, 2004; Salonen Ros et al, 2000). Genome-wide linkage studies on pedigrees have identified risk loci on chromosomes 2p13, 2q23, 11q23, 10q22, 22q12, 2p25, 9p13, 4q32 and 9p11 (Arngrimsson et al, 1999; Lachmeijer et al, 2001; Laivuori et al, 2003; Moses et al, 2000; van Dijk et al, 2005). Genome-wide association studies (GWAS) have found some variants associated with PE (Johnson et al, 2012; Zhao et al, 2013).

Patients with early-onset PE have more severe maternal and perinatal complications than those with late-onset PE (Raymond & Peterson, 2011). We do not know, however, if early-onset PE and late-onset PE have a common or distinct underlying pathogenesis at the molecular level. Numerous placental microarray studies failed to identify distinct molecular processes, due to small sample sizes or insufficient grouping of clinical subtypes (Enquobahrie et al, 2008; Junus et al, 2012; Kang et al, 2011; Nishizawa et al, 2011; Nishizawa et al, 2007; Pang & Xing, 2003; Sitras et al, 2009; Tsai et al, 2011). Some analyses use no classification of samples (Enquobahrie et al, 2008; Pang & Xing, 2003) and some focus on severe PE (Nishizawa et al, 2011), or early-onset severe PE (Varkonyi et al, 2011). Although several microarray studies contain both early- and late-onset PE placental samples, these studies investigated differential expression between all PE samples and controls, or between early and late PE (Junus et al, 2012; Nishizawa et al, 2007; Sitras et al, 2009), but did not investigate differential expression in subtypes by comparing them to normal controls. A recent RNA-seq study, done on placentae of 9 PE samples (in three pools) and 9 controls (in three

pools), found 53 PE-associated genes (Kaartokallio et al, 2015). Although these studies point to placental mechanisms underlying PE involved in angiogenesis (Junus et al, 2012; Sitras et al, 2009), immune functions (Tsai et al, 2011) and hypoxia (Nishizawa et al, 2011), it is still not clear whether the molecular pathways involved in the functional defects of placentae are different in clinical subtypes of this complex disorder. Moreover, if transcriptomes are compared between more homogeneous samples of each particular subtype and normal subjects, more complete sets of dysregulated genes might be revealed for each subtype.

In order to systematically search for placental molecular signatures associated with clinical subtypes, we carried out transcriptome sequencing on 65 high quality placenta samples, including 33 from 30 PE patients and 32 from 30 normal subjects, and found dysregulated genes associated with two distinct sets of pathways involved in two the major clinical subtypes.

# Results

**Subtypes based on clinical characteristics and sample clustering based on RNA-seq data.**

We collected 32 placentae from 30 normal individuals and 33 placentae from 30 PE patients, which were carefully classified into subtypes based on clinical characteristics (Fig 1A and Table EV1). The disease group and normal group showed significant differences in systolic and diastolic pressures, delivery time and baby weights. The early-onset severe PE (EOSPE) group showed significant differences in most of these characteristics compared to the late-onset severe PE (LOSPE) cases. Both EOSPE and LOSPE showed significant differences in these characteristics compared to late-onset mild PE (LOMPE), and the LOMPE group had only moderately elevated blood pressures compared to controls (see Table EV1 for statistics and $P$ values, Student's t test). Notably, the LOSPE patients had significantly elevated C-reactive protein (CRP) levels ($P = 0.0203$ between LOSPE and normal control, $P = 0.0268$ between LOSPE and EOPSE, Student's t test), whereas the other two groups were within the normal range (Fig EV1A).

Four placental specimens were dissected from the midsection between the chorionic plate and maternal basal plate of each placenta soon after delivery and rapidly preserved in RNAlater® to prevent RNA degradation. RNA-seq was carried out with Illumina Highseq2500. We also performed RNA-seq on two cord-blood samples for computational removal of contamination of blood cells in the placental tissues (Table EV2). Principal component analysis (PCA) and sample clustering analysis were performed on our dataset, to explore sample features in gene expression levels. The EOSPE samples were clustered together with four LOSPE samples in PCA (Fig 1B) and in the heatmap (Fig 1C). The rest LOSPE and all LOMPE samples were mixed with normal samples (Fig 1B and C). After the normal samples were removed from the clustering analysis on the PE samples, we got similar results (Fig EV1B and Table EV1). Looking into the clinical records, we found that the four LOSPE patients whose samples were clustered together with EOSPE samples, showed characteristics similar to those of the EOSPE patients except for gestational age at PE-onset, indicating these four "LOSPE" patients could in fact be EOSPE patients who were diagnosed later than the actual onset time. We extracted the top 50 genes that contributed to the first principal component (PC1), and found 36 known PE-associated genes such as *BHLHE40*, *ENG*, *FLT1*, *HK2*, *HTRA4*, *INHBA*, *LEP*, *NDRG1*, *PAPPA2*, *SASH1*, *SIGLEC6*, *SLC6A8*. The signatures of gene expression that separate EOSPE from the two other clinical subtypes suggest that they may have different underlying molecular mechanisms.

**Distinct gene expression patterns in PE subtypes**

To identify groups of genes that may be involved in the heterogeneous pathology of the clinical subtypes, we compared the RNA-seq data of all PE samples or each clinical subtype with the normal controls, using two different methods DESeq2 and edgeR (Fig 2B), and found a total number of 3,100 differentially expressed genes (DEGs), including 31 microRNAs and 399 lncRNAs (Table EV3).

We identified 2,977 DEGs (2,298 protein-coding genes) by comparing EOSPE samples with controls, while we only got 1,291 DEGs by comparing all PE samples with controls, suggesting a homogeneity within the EOSPE cases. For the other two subtypes, we found 375 DEGs (322 protein-coding genes) between LOSPE samples and controls, and 42 DEGs (39 protein-coding genes) between LOMPE samples and controls (Fig 2A and EV2A, and Table EV3). The large differences between the numbers of DEGs in the three clinical subtypes (EOSPE, LOSPE and LOMPE) indicate that there are different molecular mechanisms underlying the clinical subtypes (Fig 2B). In general, we found more up-regulated genes than down-regulated genes. The numbers of up-regulated genes are 1.3~2.3 times higher than those of down-regulated genes in the three comparisons. We carried qPCR to validate some of the differentially expressed genes (Fig 2C, EV2B and C).

To assess the functional relevance of our dataset, we compared our DEGs with 1,177 PE-associated genes collected from the literature (Table EV4). Most of the reliable PE-associated genes which appear twice or more in literature are recovered in our dataset (Fig 2D). Of the 1,177 known PE-associated genes, the DEGs in our dataset (protein-coding genes) recover 31.7% (296/934), 59.9% (88/147) and 84.4% (81/96) of those found once, twice, three or more times in literature. These PE-associated genes show significant enrichment in DEGs (protein-coding genes) of EOSPE and LOSPE: 19.3% ($P$ = 2.2e-16, One-sided Fisher's exact test) in all DEGs, 19.5% ($P$ = 2.2e-16, One-sided Fisher's exact test) in DEGs of EOSPE, 26.7% ($P$ = 2.2e-16, One-sided Fisher's exact test) in DEGs of LOSPE, and 7.7% ($P$ = 0.4261, One-sided Fisher's exact test) in DEGs of LOMPE (Fig 2E). Because PE has the most significant consequence on developmental brain impairments in offspring, which lead to intellectual disability (Griffith et al, 2011), epilepsy (Wu et al, 2008), autism (Dachew et al, 2018b; Trosse et al, 2007; Walker et al, 2015) and schizophrenia (Dalman et al, 1999; Ursini et al, 2018), we performed the enrichment analyses of schizophrenia-associated genes in DEGs, and found that the disease-associated genes were significantly enriched (Fig EV2D).

These results suggest that the differentially expressed genes are not only associated with PE, but also with consequence of PE on the offspring in their later life.

**Molecular Pathways involved in PE**

The large differences between DEGs observed in EOSPE and LOSPE suggest that these two major subtypes are due to different molecular mechanisms. To find out the associated pathways and molecular functions of the DEGs of these two major subtypes, we performed enrichment analyses on the Kyoto Encyclopedia of Genes and Genomes (KEGG) and Gene Ontology (GO) terms for the protein-coding DEGs of EOSPE (2,298) and LOSPE (322) (Fig 3). We found that 39 enriched pathways for EOSPE and 35 enriched pathways for LOSPE (Table EV5). Importantly, the enriched pathways for EOSPE and LOSPE were largely distinct, with only 7 pathways in common. The common pathways included the HIF-1 signaling pathway, focal adhesion, carbon metabolism, glycolysis/gluconeogenesis, biosynthesis of amino acids, adipocytokine and the AGE-RAGE signaling pathway (Fig 3A and Table EV5). Most of the enriched pathways of EOSPE were involved with metabolism, such as lysosomal function, AMPK signaling, tight junction pathways, insulin signaling, fatty acid elongation, galactose metabolism, alanine, aspartate and glutamate metabolism pathways (Fig 3A), many of which are reported to be associated with PE in previous studies (Table EV5). By contrast, most of the enriched pathways of LOSPE were involved in immune or autoimmune pathways, such as allograft rejection, inflammatory bowel disease, rheumatoid arthritis, HTLV-I infection and tuberculosis pathways (Fig 3A). The immune genes *HLA-DRB1*, *HLA-DQA1*, *HLA-DRA*, *HLA-DPA1*, *HLA-DPB1* and *HLA-DMB* were up-regulated in LOSPE, but not in EOSPE. These results suggest that the two main subtypes EOSPE and LOSPE have different underlying mechanisms. The subtype EOSPE may be mainly caused by disturbance in metabolism, while the subtype LOSPE by dysfunction of the immune system in the placenta. The metabolic disturbances in EOSPE and the abnormal immune functions in LOSPE may lead to changes in other pathways in the placenta. We obtained similar results in the pathway enrichment

analysis for DEGs of EOSPE-only, LOSPE-only and shared DEGs between the two subtypes (Fig EV3A). We also performed pathway enrichment analysis for up-regulated and down-regulated DEGs respectively, and found that the up-regulated genes and down-regulated genes were involve in different pathways in EOSPE, but no enriched pathway for down-regulated genes in LOSPE (Fig EV3A and B, and Table EV5).

We also performed GO-term enrichment analysis, and found that the DEGs in EOSPE were involved in distinct molecular activities (GO-MF) from those in LOSPE (Fig 3B). We detected significant enrichment of DEGs of EOSPE in 16 molecular function terms of GO, including key molecular activities in constituents of ribosomal structure, differential sugar binding and transmembrane transporter activities, phosphatidylinositol binding, adhesion-molecule binding, virus receptor activities and molecular functions hijacked by virus (Fig 3B). We detected significant enrichment of DEGs of LOSPE in only 5 molecular function terms of GO, three of which are involved in the immune response, and the other two terms are growth factor binding and 3',5'-cyclic-GMP phosphodiesterase activity (Fig 3B). These results are consistent with the results of pathway enrichment analysis we showed above (Fig 3A): the principal molecular disturbance in EOSPE in the placenta involved metabolism, while immune-function abnormalities predominated in LOSPE. The results of enrichment analyses on biological process (BP) terms and cellular component (CC) terms of GO support the same conclusion (Fig EV3C and D, and Table EV5).

**Transporter genes involved in PE**

Because the DEGs of EOSPE were enriched in many transmembrane transporter activities of GO MF terms (Fig 3B), we collected 1,554 transporter genes by searching The Gene Ontology (GO) database with key words "transporter", "carrier" or "substrate exchanging" in order to further look into the dysregulated molecular transportation in the placenta of EOSPE patients (Table EV6). We found 13.2% (205 of 1,554) transporter genes in DEGs of EOSPE, and only 1.5% (23/1,554) transporter genes in DEGs of LOSPE (Fig 4B and Table EV6). Notably, these transporter genes

were significantly enriched in the DEGs of the EOSPE (*P* = 0.027, One-sided Fisher's exact test), but not in LOPSE (*P* = 0.723, One-sided Fisher's exact test) (Fig 4B).

The 205 transporter genes in DEGs of EOSPE were enriched in 397 GO biological process (BP) terms (Adjusted *P*-value ≤ 0.05, hypergeometric test from ClusterProfiler Package). For these genes, we constructed a GO-term-to-gene bipartite network for all enriched BP terms, which contained 5 groups of functionally related genes (Fig 4C and Table EV7). Group I contains genes involved in different ion channels, some of which were previously reported to be associated with PE, e.g. *ATP1A1*, *CACNA1D*, *CALM1*, *EHD3*, *FGF12*, *KCNE1*, *KCNH5*, *NOS1AP*, *PDE4D*, *SCN1B*, *SCN4B* and *SRI* (Table EV7); group II contains genes involved in macro-autophagy, which was also previously proposed to be involved in PE(Saito & Nakashima, 2013; Saito & Nakashima, 2014), such as some up-regulated genes *ATP6V1B1*, *ATP6V1C2*, *ATPV1D1*, *ATP6V0A1*, *ATP6V0E1*, and *ATP6V0D1;* group III, genes in transmembrane transportation of hexose, fatty acids, amino acids, vitamins, organic anions and mitochondrial components; group IV, genes in transmembrane transportation, storage or homeostasis of lipids, sterols, cholesterol and phospholipids, and group V, genes in transportation of hormone, response of cAMP and drug, secretion of insulin and protein. (Fig 4C). The detailed information of all the five groups is included in the Supplemental material (Table EV7). We also performed pathway enrichment analysis for these 205 differentially expressed transporter genes in EOSPE and found that the genes were enriched in 34 pathways (Fig EV4 and Table EV7), which are consistent with the five functional groups in GO-term analysis (Fig 4C). Some of these pathways are known to be involved in PE, such as oxidative phosphorylation (Kang et al, 2011), ABC transporter pathway (Joshi et al, 2016) and circadian entrainment pathway (Valenzuela et al, 2015).

The 23 transporter genes in the DEGs of LOSPE were enriched in 93 GO biological process terms (Adjusted *P*-value ≤ 0.05, hypergeometric test from ClusterProfiler Package). We also constructed a GO-term-to-gene-bipartite network for these 93 terms, which contained 2 groups of functionally related genes (Fig 4D). The group A contains 15 genes involved in ion channels, and

group B only contains 8 genes involved in transportation of chloride, carboxylic acid, cAMP and anion.

The placenta serves as a molecular barrier between maternal and fetal circulation, and transmembrane transporters play a key role in the placental exchange function. On one hand, the fetus receives nutrients from the mother, such as carbohydrates, lipids, amino acids, vitamins and minerals for adequate development and growth, and on the other hand, the fetus can drain waste products via efflux processes into the maternal circulation (Huang et al, 2018). The dysregulation of the quantity, density and activity of membrane transporters could impact the nutrient supply and consequently affect fetal development. We consistently observed low birth weight (LBW) and fetal growth restriction (FGR) in 90% (9 of 10) of the babies born by EOSPE patients and 53% (8 of 15) of the babies born by LOSPE patients (Table EV1). The weights of babies in EOSPE group were significantly lower than those of babies in LOSPE group (Table EV1).

**Transcription factors involved in PE**

We want to know if the distinct differential gene expression patterns in EOSPE and LOSPE might be due to the dysregulation of a set of transcription factors. We used 1,639 reliable human transcription factors (TFs) (Lambert et al, 2018) from literature to search 6 databases (Consortium, 2012; Han et al, 2018; Han et al, 2015; Jiang et al, 2007; Marbach et al, 2016; Neph et al, 2012; Zheng et al, 2008) for their targets, and built a regulatory network containing 996 TFs and 16,538 target genes (Fig 5A). These 996 TFs cover 65.6% (105/160) of the TFs in the DEGs of EOSPE, such as *LTF*, *BCL6*, *ARNTL*, *NFKB1*, *PPARG* and *ATF3* (Table EV8), which target to 81.9% (1,882/2,298) of DEGs in EOSPE. They also cover 81.5% (22/27) of TFs in DEGs of LOSPE, such as *FOSL1*, *FOXO1*, *STAT3*, *YBX1*, *DBP* and *SOX15* (Table EV8), which target 67.1% (216/322) of DEGs in LOSPE (Fig 5B and C, and Table EV8).

These TF-to-target regulation relationships were exemplified as regulatory networks (Fig 5D and E), suggesting that altered expression in a small number of TFs may lead to a large number of

DEGs in PE. We performed pathway enrichment analysis on the two regulatory networks and found that 18 pathways were involved in EOSPE and 47 pathways in LOSPE (Table EV8), which are consistent with the enriched pathways for the differentially expressed genes in EOSPE and LOSPE (Fig 4). Some key PE-related pathways were affected by dysregulated TFs as shown in the two example modules involved in AMPK and FoXO signaling pathways (Fig EV5A and B) in EOSPE and two other example modules in PI3K-Akt signaling pathway and immune function pathways in LOSPE (Fig EV5C and D).

# Discussion

Patients with PE display a wide range of symptoms. Early-onset PE patients usually have severe symptoms; among the late-onset PE patients, however, some show severe symptoms, while others are only mildly affected. Although the clinical characteristics of PE can be measured, the underlying molecular bases for the clinical subtypes are not known. Anecdotal studies on placentae of PE patients point to some abnormal molecular processes, such as defective angiogenesis (Liu et al, 2015; Maynard & Karumanchi, 2011), hypoxia (Rajakumar et al, 2001; Shamshirsaz et al, 2012; Tal, 2012) and inflammation (Centlow et al, 2008; Hoegh et al, 2010; Sitras et al, 2009), but these abnormalities do not unambiguously define the different molecular pathways involved in the clinical subtypes. Additional approaches are needed to identify the genes underlying the differential pathogenesis of the PE subtypes. Numerous microarrays on transcriptomes of placentae (Enquobahrie et al, 2008; Junus et al, 2012; Kang et al, 2011; Nishizawa et al, 2011; Nishizawa et al, 2007; Pang & Xing, 2003; Sitras et al, 2009; Tsai et al, 2011) and a recent RNA-seq study (Kaartokallio et al, 2015) did not identify distinct molecular processes associated with the clinical subtypes, probably due to small sample sizes and inadequate comparisons between the grouped samples. The other transcriptomic study on the PE focuses on the maternal side, the uterine decidua (Tong et al, 2018). In order to systematically search for genes involved in clinical PE subtypes, we

dissected samples from the mid-section placental tissues between the chorionic and maternal basal surfaces of placentae from patients with classical presentation of preeclampsia, which were grouped into three clinical subtypes: early-onset severe PE (EOSPE), late-onset severe PE (LOSPE) and late-onset mild PE (LOMPE) (Table EV1). Our identification of the distinct molecular processes involved in the clinical subtypes was based on initial classification and comparison of RNA-seq data of the well classified samples. Because we noticed that the blood could not be completely washed away from the placental tissues, which might be one factor affecting previous transcriptomic analyses, we performed RNA-seq on whole cord-blood cells and removed cord-blood contamination by computational method (Table EV2).

In the principal component analysis (PCA) and sample clustering analysis, the EOSPE samples grouped together, separate from the clusters of LOSPE and LOMPE samples (Fig 1B). We carried out comparisons between the placental transcriptomes of the three clinical subtypes and normal samples (EOSPE vs. normal, LOSPE vs. normal and LOMPE vs. normal) to identify the differentially expressed genes in the three clinical subtypes. Surprisingly, EOSPE had 2,977 differentially expressed genes (2,298 protein coding genes), while LOSPE had only 375 (322 protein coding genes), indicating different molecular mechanisms for these two subtypes (Fig 2A); both data sets, however, showed significant enrichment of previously reported PE-associated genes (Fig 2C). Moreover, LOMPE had only 42 differentially expressed genes that showed no enrichment for the previously reported PE-associated genes (Fig 2C), suggesting a lesser degree of involvement of genetic factors in the pathogenesis of mild PE. Most interestingly, we found distinct molecular processes involved in the two major subtypes of PE: metabolisms and related activities in the EOSPE and immune activities in LOSPE (Fig 3). These PE-associated pathways were regulated by small number of transcription factors (Fig 5), which might be the driver genes during the abnormal development of the placenta of PE patients.

As the placenta is the supporting system for a developing embryo, the widespread gene expression change in placentae of PE patients might have profound effects on the development of

the fetuses and the health of the offspring in later life. Accumulating evidences point to association of preeclampsia with brain developmental disorders, including intellectual disability (Griffith et al, 2011), epilepsy (Wu et al, 2008), autism (Dachew et al, 2018b; Trosse et al, 2007; Walker et al, 2015) and schizophrenia (Dalman et al, 1999; Ursini et al, 2018). It might be partially due to the placenta-derived factors that affects the development of the brain and thus leads to complex brain disorders (Maher et al, 2018; Ursini et al, 2018). Ursini et al found that, for some loci of schizophrenia, the association between genetic risk and schizophrenia is affected by complicated pregnancies, and the gene set within these loci are differentially expressed in placentae from complicated in comparison with normal pregnancies (Ursini et al, 2018). In order to investigate to what extent the differentially expressed genes in PE may be also involved in schizophrenia, we collected 3,437 schizophrenia-associated genes from classical schizophrenia databases and literature, and found that the schizophrenia-associated genes are significantly enriched in DEGs of EOSPE ($P$ = 1.4e-4, One-sided Fisher's exact test) and LOSPE ($P$ = 0.01, One-sided Fisher's exact test) (Fig EV2D).

In conclusion, this study provides molecular-level evidences that EOSPE and LOSPE are two different subtypes with distinct underlying molecular mechanisms while LOMPE may not be due to placental factors. Expression of metabolism-related genes is significantly affected in EOSPE, whereas in LOSPE it preferentially involves the expression of immune-related genes. Most notably, the transporter genes dysregulated in EOSPE may cause abnormal molecular exchange between maternal and fetal circulation, leading to fetal growth restriction. The TF-target networks demonstrate that only a small number of dysregulated transcription factors may serve as driving genes, causing differential expression of a larger number of genes in EOSPE and LOSPE. Such distinct underlying mechanisms suggest that potentially more effective therapies could be developed to target the different subtypes at their molecular levels.

# Materials and Methods

**Sample collection and criteria for grouping PE cases into different clinical subtypes.**

We collected the tissue samples at the Department of Obstetrics & Gynecology of Nanfang Hospital in China from January 2015 to July 2016. The clinical characteristics of each patient were extracted from the medical records, which strictly followed the American Board of Obstetrics and Gynecology, Williams Obstetrics 24th edition. Samples of placental tissues were collected from the mid-section placental tissues of the placenta, between the chorionic and maternal basal surfaces, at four different positions within 5 minutes after delivery. These placental tissues were placed into RNAlater® solution and stored at -80 °C. This research has been approved by the Ethnics Board of Nanfang Hospital of Southern Medical University, and all patients have signed the informed consent.

Preeclampsia (PE) is characterized by new-onset hypertension (≥140/90 mmHg) and proteinuria at ≥20 weeks of gestation, or in the absence of proteinuria, hypertension with evidence of systemic disease (Chaiworapongsa et al, 2014). If patients with PE have systolic blood pressure ≥160 mmHg or diastolic blood pressure ≥110 mmHg, or other organ failures, we call them "preeclampsia with severe features" or briefly "severe PE". According to gestational age at diagnosis or delivery, PE can be divided as early-onset (< 34 weeks) and late-onset (≥ 34 weeks) (Redman & Sargent, 2005; von Dadelszen et al, 2003). Although other gestational age cut-offs have been suggested, 34 weeks remains the most commonly used (Parra-Cordero et al, 2013; Soto et al, 2012), presumably as the rate of neonatal morbidities declines considerably after reaching this time point.

According to above criteria and clinical record of each case, we grouped our PE patients into three clinical subtypes: late-onset mild PE (LOMPE), early-onset severe PE (EOSPE) and late-onset severe PE (LOPSE). The clinical information of the three clinical subtypes and comparisons between the subtypes were listed in Table EV1.

**RNA-seq.**

Total RNA was isolated using the RNeasy® Plus Universal Mini Kit (Qiagen) according to the manual protocol. RNA-seq was carried out at Berry Genomics Corporation (Beijing, China). Briefly, RNAs with polyA tails were isolated, and double-stranded cDNA libraries were prepared using TruSeq RNA Kit (Illumina) followed by paired-end sequencing using Illumina Hiseq 2500. In order to computationally remove the cord blood contamination, RNA-seq was also done on two samples of cord blood. The reads from the clean data of RNA-seq were aligned to the human reference genome (hg38) with STAR (Dobin et al, 2013) and counted using HTSeq (Anders et al, 2015) with union mode. Raw counts for annotated genes (protein coding genes, rRNA, microRNA, LncRNA, pseudogenes and so on) in the General Transfer Format (GTF) annotation file were obtained. The reference genome sequence and GTF annotation files were downloaded from the Ensembl website (http://asia.ensembl.org/index.html) in the alignment and counting reads steps.

Annotation information, such as gene location, gene types, gene symbols, for differential expression genes were extracted from EnsDb.Hsapiens.v86 package. Four brief categories of gene biotypes were divided following the table listed on the GENCODE website page (https://www.gencodegenes.org/gencode_biotypes.html). Gene Ontology terms overrepresentation, Disease Ontology, Pathway and functional annotation analyses were used clusterProfiler (Yu et al, 2012) and DOSE (Yu et al, 2015) packages. All analyses were done in R-platform, and all packages and public datasets were listed in Table EV9.

**Computational removal of blood contamination**

The contaminated cord blood cells could not be totally washed from the placenta tissue samples, so the extracted RNAs contained some from cord blood cells. To better study the difference in gene expression of placenta tissues between PE patients and controls, we removed the contamination using the expression level of a cord blood marker gene for normalization. Cord blood samples from two normal pregnancy women were sequenced using the same RNA-seq method. Differentially expressed gene analysis was applied to two cord blood samples against two normal placenta

samples to identify marker genes for cord blood cells. We used the Hemoglobin Subunit Mu (HBM), a subunit of hemoglobin that is specifically expressed in red blood cells, and recalibrate the raw count of genes accordingly in the following equation:

$$\text{geneX}_{\text{actual}} = \text{geneX}_{\text{placenta}} - \text{geneX}_{\text{cord blood}} \times \frac{\text{HBM}_{\text{placenta}}}{\text{HBM}_{\text{cord blood}}}$$

The raw counts of genes in each sample before and after removing blood contamination were showed in Table EV2.

**Clustering analysis of PE samples only**

The EOSPE samples were separated from other samples in clustering, whereas LOMPE and LOSPE samples were clustered together with normal samples (Fig 1B). To explore clusters of PE samples, we performed clustering analysis on PE samples only, including EOSPE, LOSPE and LOMPE samples, excluding normal samples. The top 500 most variances (genes) across all PE samples were used for clustering (Fig EV1B). All of the LOMPE samples and most of the LOSPE samples clustered together on the left side of the graph as cluster-1, whereas all EOSPE samples clustered on the right side as cluster-2, together with the four LOSPE samples. In addition, we carried out a comparison of the clinical characteristics of the patients between the two clusters and detected significant differences in blood pressure, gestational age of delivery, baby weight, FGR or LBW and proteinuria (Table EV1). Of the 14 babies (13 placenta samples) in subclass-2, 12 babies (85.7%) had FGR or LBW, whereas 10 of 20 babies (50%) had FGR or LBW in subclass-1. The mean length of gestation in two clusters were 264 (37.7 weeks) and 228.8 (32.5 weeks), and patients in the two clusters showed different weeks at onset of PE. The quantification of proteinuria in the two clusters also had significant differences. In other words, PE samples in cluster-1 showed late-onset, relatively moderate symptom, and less harmful outcomes; PE samples in cluster-2 showed early-onset, more severe symptom and worse outcomes.

**Methods for identifying differentially expressed genes (DEGs)**

Two widely used tools in R platform, DESeq2 (Love et al, 2014) and edgeR (Robinson et al, 2010), were applied to determine differential expression genes in different comparisons. The genes with at least 1 count in every samples were used in DESeq2 method and the genes with CPM (Counts Per Million) value larger than 1 in at least one sample were used in edgeR method.

Samples similarity in each group was calculated using all information of expressed genes (raw count of gene in all samples is more than 1) for heat-map, and top 500 genes with the biggest variations were used for PCA analysis. Four comparisons between the clinical subtypes of PE patients were carried out using both DESeq2 and edgeR: **(i)** all PE placental samples (n = 33) versus all normal placental samples (n = 32); **(ii)** late-onset mild PE placental samples (n = 9) versus all normal placental samples; **(iii)** early-onset severe PE placental samples (n = 9) versus all normal placental samples; and **(iv)** late-onset severe PE placental samples (n = 15) versus all normal placental samples (Fig 2). Adjusted *P*-value in DESeq2 and FDR in edgeR were used together to determine whether a gene is differentially expressed. If adjusted *P*-value ≤ 0.05 of one gene under DESeq2 and FDR ≤ 0.05 under edgeR, the gene was considered as differentially expressed in between two groups of samples. The intersections of differential expression genes from these two methods were taken as the final results of each comparison. Because we only identified a very small number of the differential expression genes in LOMPE, we merge the results of DESeq2 and edgeR as the final result in the comparison of LOMPE samples with normal samples. The numbers of differentially expressed genes identified by the two methods and gene biotypes were showed in Fig EV2A and Table EV3.

**The DEGs overlapped with PE-associated genes from the literature**

There were a lot of studies to investigate differentially expressed genes compared normal placenta tissues with PE placenta tissues. Due to small sample sizes and undistinguished subtypes of PE samples in previous transcriptomic studies, the differentially expressed genes in PE have not been well defined. To increase the sample sizes, several meta-analyses on reported microarray datasets

were published since 2012 (Brew et al, 2016; Kleinrouweler et al, 2013; Moslehi et al, 2013; Tejera et al, 2017; Vaiman et al, 2013; Vaiman & Miralles, 2016). In 2015, Kaartokallio et al reported the first RNA-seq study on PE (Kaartokallio et al, 2015). To obtain a consensus PE-associated genes, we collected 1,177 genes from literature, including 6 meta-analysis papers (Brew et al, 2016; Kleinrouweler et al, 2013; Moslehi et al, 2013; Tejera et al, 2017; Vaiman et al, 2013; Vaiman & Miralles, 2016), 1 literature-curation paper (Kobayashi, 2016) and 1 RNA-seq research paper (Kaartokallio et al, 2015). These 1,177 genes are PE-associated genes and information for genes were listed in Table EV4.

**Pathway enrichment analysis for DEGs of EOSPE-only, LOSPE-only and shared genes**

The numbers of differentially expressed protein-coding genes are 2,061 in EOSPE-only, 85 in LOPSE-only and 237 in common (Venn diagram in Fig EV3A). KEGG pathway enrichment analysis was done by clusterProfiler package (Yu et al, 2012). The cutoff of $q$-value $\leq 0.2$ was chosen to select most significantly enriched pathways (Fig EV3A). There 39 enriched pathways for EOSPE-only DEGs, 35 for LOSPE-only DEGs, and 7 for shared DEGs.

The pathway enrichment for up-regulated and down-regulated DEGs in both EOSPE and LOPSE were done using the same method (Fig EV3B and C).

**GO terms enrichment analysis for DEGs of EOSPE and LOSPE**

Gene Ontology terms, (BP, CC and MF) enrichment analyses were also using clusterProfiler package(Yu et al, 2012), and the $q$-value $\leq 0.05$ and adjusted $p$-value $\leq 0.05$ were set as cutoff to select most significantly enriched terms. The DEGs of EOSPE were statistically significant enriched in 248 terms of biological process (BP), 51 cellular component (CC) and 16 molecular function (MF). The DEGs of LOSPE were statistically significant enriched 169 terms of BP, 22 of CC and 5 of MF. The overlap of three categories of enriched GO terms are 63 (BP), 4 (CC) and 0

(MF) between DEGs of EOSPE and LOSPE, and most of enriched terms are different in EOSPE and LOSPE (Fig EV3D and E, and Table EV5).

We used semantic similarity-based method (REVIGO) (Supek et al, 2011) to summarize the enriched GO terms in the tree components. For EOSPE, 199 of 248 BP terms were summarized into 14 representative entries. The first representative entry is the nuclear-transcribed mRNA catabolism, containing 55 terms. The second is SRP-dependent co-translational protein targeting to membrane, containing 42 terms. Other entries include stress-activated protein kinase signaling cascade, regulation of macro-autophagy, response to metal ion, estrogen and insulin, and superoxide metabolism. For LOSPE, however, 130 of 169 BP terms were summarized into 11 representative entries. The first entry is positive regulation of cell adhesion, containing 57 terms, and the second entry was response to interferon-gamma, containing 27 terms. Other entries include response to oxygen levels, temperature homeostasis, epithelial cell proliferation, and synapse organization.

The Enriched CC and MF terms in EOSPE and LOSPE also showed considerable difference (Table EV5). This suggested that besides "canonical pathways" such as HIF-1 signaling pathway, the enriched terms are associated with metabolisms, substrates transport in EOSPE, whereas enriched terms are associated with immune functions in LOSPE.

We found that there are 5 terms of MF enriched in EOSPE DEGs are related to glucose/carbohydrate transport, and 20 terms of BP enriched in EOSPE DEGs are also associated with materials transmembrane transporters, while 17 terms of BP enriched in LOSPE DEGs are associated with transporters.

**Clustering analysis for GO-term-to-gene-bipartite networks for differentially expressed transporter genes of EOSPE and LOSPE**

We found that transporter genes were significantly enriched in the DEGs of the EOSPE ($P = 0.027$) compared with all annotated human genes (19,738) in GO, but not significantly enriched DEGs of the LOSPE ($P = 0.723$). There were 205 and 23 transporter genes involved in 397 and 93 GO

biological process (GO BP) terms in EOSPE and LOSPE respectively. To summarize the enriched

BP terms, we firstly constructed GO-term-to-gene-bipartite networks for EOSPE and LOSPE in

Cytoscape using yfiles organic layout. Then we calculated Jaccard score using number of shared

genes in the terms, and clustered terms into several clusters based on shared genes. Combining the

layout of networks and the results of clustering analysis, 5 and 2 groups of GO BP terms were

manually determined for EOSPE and LOSPE, respectively. All enriched BP terms in differentially

expressed transporter genes of the EOSPE and the LOSPE were listed in Table EV7.

**Transcription factors and target genes**

To find out the transcription factors that may mostly determine the distinct gene expression patterns

in EOSPE and LOSPE, we matched the target genes collected from in 6 databases (Consortium,

2012; Han et al, 2018; Han et al, 2015; Jiang et al, 2007; Marbach et al, 2016; Neph et al, 2012;

Zheng et al, 2008) to the 1,639 reliable human transcription factors (TFs) downloaded from a recent

literature (Lambert et al, 2018). WE obtained 996 TFs and 16,538 target genes (Table EV8) and

performed TF-target network analysis (Fig 5 and Fig EV5).

**Schizophrenia associated genes enriched in the DEGs of the EOSPE and LOSPE**

Abnormal intra-uterine environment is an important status of preeclampsia, and our data showed

that 90% and 58% babies born by EOSPE and LOSPE patients had FGR/LBW. Recently, a meta-

analysis reported that hypertensive disorders of pregnancy may be associated with an increase in the

risk of ASD (autism spectrum disorder) and ADHD (attention-deficit/hyperactivity disorder)

(Maher et al, 2018). Weinberger's study showed that the set of genes within genomic risk loci for

schizophrenia that have interaction with intra-uterine and perinatal complications is highly

expressed in placenta (Ursini et al, 2018). Other studies also demonstrated that preeclampsia was

associated with increased risk of offspring schizophrenia (Dachew et al, 2018a).

To investigate the relationship between preeclampsia and schizophrenia, we collected 3,437 schizophrenia-associated genes from classical schizophrenia databases and literature (Allen et al, 2008; Cariaso & Lennon, 2012; Cooper et al, 2006; Jia et al, 2017; Ramos et al, 2014; Schizophrenia Working Group of the Psychiatric Genomics, 2014; Wu et al, 2017). Since increasing evidences demonstrate the association of preeclampsia with schizophrenia, we want to know if the differentially expressed genes are enriched for schizophrenia associated genes. In our dataset, the protein coding DEGs (2,405, including DEGs from EOSPE, LOSPE and LOMPE), DEGs of EOSPE (2,298) or LOSPE (322) were significant enriched for schizophrenia-associated genes: 20.7% (497/2,405, $P$ = 5.0e-5), 20.54% (472/2,298, $P$ = 1.4e-4), 22.98% (74/322, $P$ = 0.01), compared with the expected 17.8% (the ratio 3,437 schizophrenia associated genes to 19,351 annotated human coding genes). In protein coding DEGs of LOMPE, there is no significant enrichment of schizophrenia associated genes (15.4%, $P$ > 0.5) (Fig EV2D).

# Acknowledgments

We thank the patients for their generous support for this study. This work was supported by GDSTC | Science and Technology Planning Project of Guangdong Province (S&T Project of Guangdong Province) (No. 2017B020227010 and No. 2015B050501006) and National Natural Science Foundation of China (No.31771434, No. 81671466, No 81471464 and No. 81771609).

# Author contributions

Conception and design: Xinping Y and YY; Recruitment of the patients and normal subjects: Yunfei G and MZ; Sequencing data analysis: ZR, Yue G, GL and Xiaoxue Y; Sample collection: HW, JW; RNA extraction and validation experiments: QC, SJ, CF and YS; Manuscript writing: ZR, Yunfei G, Yue G, Xinping Y, YY and GL.

# Conflict of interests

All authors declare no competing interests.

# The Paper Explained

# PROBLEM

Placenta is one of the most important organs in human. It not only influences both the mothers and fetuses during pregnancy, but also lifelong health (especially brain health) of the offspring. Because it is a temporary organ, its importance has not been well recognized. Preeclampsia, a disease with placenta origin, has a global incidence rate of 4.6% in pregnancies, leading to 60,000 maternal deaths per year. It has significant consequences on fetal development and growth, often leading to perinatal and infant morbidity or mortality and contributes to preterm births and fetal growth restriction (FGR). In the long term, preeclampsia affects brain development and functions of the offspring, leading to intellectual disability, epilepsy, autism and schizophrenia. Preeclampsia patients display a spectrum of onset time and severity of clinical presentation, yet the underlying molecular bases for the early-onset and late-onset clinical subtypes are not known.

# RESULTS

Considering that preeclampsia has placenta origin, we systematically search for placental molecular signatures associated with clinical subtypes and the underlying molecular mechanisms. We carried out RNA-seq on 65 high-quality placenta samples, including 33 from 30 patients with PE and 32 from 30 control subjects. We identified two functionally distinct sets of dysregulated genes in two major subtypes: metabolism-related genes, notably transporter genes, in early-onset severe preeclampsia (EOSPE) and immune-related genes in late-onset severe preeclampsia (LOSPE), while the late-onset mild preeclampsia (LOMPE) could not be distinguished from normal controls.

By looking into differentially expressed transcription factors (TFs) and their targets, we found that a small number of dysregulated transcription factors could contribute to the widespread gene dysregulation in both EOSPE and LOSPE.

# IMPACT

These results demonstrate that early-onset severe PE (EOSPE) and late-onset severe PE (LOSPE) may have different molecular mechanisms, whereas the late-onset mild PE (LOMPE) may have no placenta-specific causal factors. Such distinct underlying mechanisms suggest that potentially more effective therapies could be developed to target the different subtypes at their molecular level.

# For More Information

Detailed analyzed data are included in supplementary excel tables. RNA-seq data are going to be deposited at GEO and will be available. All renewable reagents and detailed protocols will be available upon request.

# References

(1998) Father's genes influence risk of pre-eclampsia. Bmj 316**:** B

(2011)  In WHO Recommendations for Prevention and Treatment of Pre-Eclampsia and Eclampsia. Geneva

Abalos E, Cuesta C, Grosso AL, Chou D, Say L (2013) Global and regional estimates of preeclampsia and eclampsia: a systematic review. European journal of obstetrics, gynecology, and reproductive biology 170**:** 1-7

Allen NC, Bagade S, McQueen MB, Ioannidis JP, Kavvoura FK, Khoury MJ, Tanzi RE, Bertram L (2008) Systematic meta-analyses and field synopsis of genetic association studies in schizophrenia: the SzGene database. Nature genetics 40**:** 827-834

Anders S, Pyl PT, Huber W (2015) HTSeq--a Python framework to work with high-throughput sequencing data. Bioinformatics 31**:** 166-169

Arngrimsson R, Siguroardottir S, Frigge ML, Bjarnadottir RI, Jonsson T, Stefansson H, Baldursdottir A, Einarsdottir AS, Palsson B, Snorradottir S et al (1999) A genome-wide scan reveals a maternal susceptibility locus for pre-eclampsia on chromosome 2p13. Human molecular genetics 8: 1799-1805

Brew O, Sullivan MH, Woodman A (2016) Comparison of Normal and Pre-Eclamptic Placental Gene Expression: A Systematic Review with Meta-Analysis. PloS one 11: e0161504

Cariaso M, Lennon G (2012) SNPedia: a wiki supporting personal genome annotation, interpretation and analysis. Nucleic acids research 40: D1308-1312

Centlow M, Carninci P, Nemeth K, Mezey E, Brownstein M, Hansson SR (2008) Placental expression profiling in preeclampsia: local overproduction of hemoglobin may drive pathological changes. Fertility and sterility 90: 1834-1843

Chaiworapongsa T, Chaemsaithong P, Yeo L, Romero R (2014) Pre-eclampsia part 1: current understanding of its pathophysiology. Nature reviews Nephrology 10: 466-480

Cnattingius S, Reilly M, Pawitan Y, Lichtenstein P (2004) Maternal and fetal genetic factors account for most of familial aggregation of preeclampsia: a population-based Swedish cohort study. American journal of medical genetics Part A 130A: 365-371

Consortium EP (2012) An integrated encyclopedia of DNA elements in the human genome. Nature 489: 57-74

Cooper DN, Stenson PD, Chuzhanova NA (2006) The Human Gene Mutation Database (HGMD) and its exploitation in the study of mutational mechanisms. Current protocols in bioinformatics Chapter 1: Unit 1 13

Dachew BA, Mamun A, Maravilla JC, Alati R (2018a) Association between hypertensive disorders of pregnancy and the development of offspring mental and behavioural problems: A systematic review and meta-analysis. Psychiatry research 260: 458-467

Dachew BA, Mamun A, Maravilla JC, Alati R (2018b) Pre-eclampsia and the risk of autism-spectrum disorder in offspring: meta-analysis. The British journal of psychiatry : the journal of mental science 212: 142-147

Dalman C, Allebeck P, Cullberg J, Grunewald C, Koster M (1999) Obstetric complications and the risk of schizophrenia: a longitudinal study of a national birth cohort. Archives of general psychiatry 56: 234-240

Davies EL, Bell JS, Bhattacharya S (2016) Preeclampsia and preterm delivery: A population-based case-control study. Hypertension in pregnancy 35: 510-519

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29: 15-21

Eneroth-Grimfors E, Westgren M, Ericson M, Ihrman-Sandahl C, Lindblad LE (1994) Autonomic cardiovascular control in normal and pre-eclamptic pregnancy. Acta obstetricia et gynecologica Scandinavica 73: 680-684

Enquobahrie DA, Meller M, Rice K, Psaty BM, Siscovick DS, Williams MA (2008) Differential placental gene expression in preeclampsia. American journal of obstetrics and gynecology 199**:** 566 e561-511

Griffith MI, Mann JR, McDermott S (2011) The risk of intellectual disability in children born to mothers with preeclampsia or eclampsia with partial mediation by low birth weight. Hypertension in pregnancy 30**:** 108-115

Han H, Cho JW, Lee S, Yun A, Kim H, Bae D, Yang S, Kim CY, Lee M, Kim E et al (2018) TRRUST v2: an expanded reference database of human and mouse transcriptional regulatory interactions. Nucleic acids research 46**:** D380-D386

Han H, Shim H, Shin D, Shim JE, Ko Y, Shin J, Kim H, Cho A, Kim E, Lee T et al (2015) TRRUST: a reference database of human transcriptional regulatory interactions. Scientific reports 5**:** 11432

Herzog EM, Eggink AJ, Reijnierse A, Kerkhof MA, de Krijger RR, Roks AJ, Reiss IK, Nigg AL, Eilers PH, Steegers EA et al (2017) Impact of early- and late-onset preeclampsia on features of placental and newborn vascular health. Placenta 49**:** 72-79

Hoegh AM, Borup R, Nielsen FC, Sorensen S, Hviid TV (2010) Gene expression profiling of placentas affected by pre-eclampsia. Journal of biomedicine & biotechnology 2010**:** 787545

Huang X, Anderle P, Hostettler L, Baumann MU, Surbek DV, Ontsouka EC, Albrecht C (2018) Identification of placental nutrient transporters associated with intrauterine growth restriction and pre-eclampsia. BMC genomics 19**:** 173

Jia P, Han G, Zhao J, Lu P, Zhao Z (2017) SZGR 2.0: a one-stop shop of schizophrenia candidate genes. Nucleic acids research 45**:** D915-D924

Jiang C, Xuan Z, Zhao F, Zhang MQ (2007) TRED: a transcriptional regulatory element database, new entries and other development. Nucleic acids research 35**:** D137-140

Johnson MP, Brennecke SP, East CE, Goring HH, Kent JW, Jr., Dyer TD, Said JM, Roten LT, Iversen AC, Abraham LJ et al (2012) Genome-wide association scan identifies a risk locus for preeclampsia on 2q14, near the inhibin, beta B gene. PloS one 7**:** e33666

Joshi AA, Vaidya SS, St-Pierre MV, Mikheev AM, Desino KE, Nyandege AN, Audus KL, Unadkat JD, Gerk PM (2016) Placental ABC Transporters: Biological Impact and Pharmaceutical Significance. Pharmaceutical research 33**:** 2847-2878

Junus K, Centlow M, Wikstrom AK, Larsson I, Hansson SR, Olovsson M (2012) Gene expression profiling of placentae from women with early- and late-onset pre-eclampsia: down-regulation of the angiogenesis-related genes ACVRL1 and EGFL7 in early-onset disease. Molecular human reproduction 18**:** 146-155

Kaartokallio T, Cervera A, Kyllonen A, Laivuori K, Kere J, Laivuori H, Group FCI (2015) Gene expression profiling of pre-eclamptic placentae by RNA sequencing. Scientific reports 5**:** 14107

Kang JH, Song H, Yoon JA, Park DY, Kim SH, Lee KJ, Farina A, Cho YK, Kim YN, Park SW et al (2011) Preeclampsia leads to dysregulation of various signaling pathways in placenta. Journal of hypertension 29**:** 928-936

Kleinrouweler CE, van Uitert M, Moerland PD, Ris-Stalpers C, van der Post JA, Afink GB (2013) Differentially expressed genes in the pre-eclamptic placenta: a systematic review and meta-analysis. PloS one 8**:** e68991

Kobayashi H (2016) Characterization of the down-regulated genes identified in preeclampsia placenta. Hypertension in pregnancy 35**:** 15-21

Lachmeijer AM, Arngrimsson R, Bastiaans EJ, Frigge ML, Pals G, Sigurdardottir S, Stefansson H, Palsson B, Nicolae D, Kong A et al (2001) A genome-wide scan for preeclampsia in the Netherlands. European journal of human genetics : EJHG 9**:** 758-764

Laivuori H, Lahermo P, Ollikainen V, Widen E, Haiva-Mallinen L, Sundstrom H, Laitinen T, Kaaja R, Ylikorkala O, Kere J (2003) Susceptibility loci for preeclampsia on chromosomes 2p25 and 9p13 in Finnish families. American journal of human genetics 72**:** 168-177

Lambert SA, Jolma A, Campitelli LF, Das PK, Yin Y, Albu M, Chen X, Taipale J, Hughes TR, Weirauch MT (2018) The Human Transcription Factors. Cell 172**:** 650-665

Liu H, Li Y, Zhang J, Rao M, Liang H, Liu G (2015) The defect of both angiogenesis and lymphangiogenesis is involved in preeclampsia. Placenta 36**:** 279-286

Liu X, Zhao W, Liu H, Kang Y, Ye C, Gu W, Hu R, Li X (2016) Developmental and Functional Brain Impairment in Offspring from Preeclampsia-Like Rats. Molecular neurobiology 53**:** 1009-1019

Logue OC, George EM, Bidwell GL, 3rd (2016) Preeclampsia and the brain: neural control of cardiovascular changes during pregnancy and neurological outcomes of preeclampsia. Clinical science 130**:** 1417-1434

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology 15**:** 550

Maher GM, O'Keeffe GW, Kearney PM, Kenny LC, Dinan TG, Mattsson M, Khashan AS (2018) Association of Hypertensive Disorders of Pregnancy With Risk of Neurodevelopmental Disorders in Offspring: A Systematic Review and Meta-analysis. JAMA psychiatry 75**:** 809-819

Marbach D, Lamparter D, Quon G, Kellis M, Kutalik Z, Bergmann S (2016) Tissue-specific regulatory circuits reveal variable modular perturbations across complex diseases. Nature methods 13**:** 366-370

Maynard SE, Karumanchi SA (2011) Angiogenic factors and preeclampsia. Seminars in nephrology 31**:** 33-46

Melchiorre K, Sharma R, Thilaganathan B (2014) Cardiovascular implications in preeclampsia: an overview. Circulation 130**:** 703-714

Mitani M, Matsuda Y, Makino Y, Akizawa Y, Ohta H (2009) Clinical features of fetal growth restriction complicated later by preeclampsia. The journal of obstetrics and gynaecology research 35**:** 882-887

Mol BWJ, Roberts CT, Thangaratinam S, Magee LA, de Groot CJM, Hofmeyr GJ (2016) Pre-eclampsia. Lancet 387**:** 999-1011

Morgan T, Ward K (1999) New insights into the genetics of preeclampsia. Seminars in perinatology 23**:** 14-23

Moses EK, Lade JA, Guo G, Wilton AN, Grehan M, Freed K, Borg A, Terwilliger JD, North R, Cooper DW et al (2000) A genome scan in families from Australia and New Zealand confirms the presence of a maternal susceptibility locus for pre-eclampsia, on chromosome 2. American journal of human genetics 67**:** 1581-1585

Moslehi R, Mills JL, Signore C, Kumar A, Ambroggio X, Dzutsev A (2013) Integrative transcriptome analysis reveals dysregulation of canonical cancer molecular pathways in placenta leading to preeclampsia. Scientific reports 3**:** 2407

Neph S, Stergachis AB, Reynolds A, Sandstrom R, Borenstein E, Stamatoyannopoulos JA (2012) Circuitry and dynamics of human transcription factor regulatory networks. Cell 150**:** 1274-1286

Nishizawa H, Ota S, Suzuki M, Kato T, Sekiya T, Kurahashi H, Udagawa Y (2011) Comparative gene expression profiling of placentas from patients with severe pre-eclampsia and unexplained fetal growth restriction. Reproductive biology and endocrinology : RB&E 9**:** 107

Nishizawa H, Pryor-Koishi K, Kato T, Kowa H, Kurahashi H, Udagawa Y (2007) Microarray analysis of differentially expressed fetal genes in placental tissue derived from early and late onset severe pre-eclampsia. Placenta 28**:** 487-497

Pang ZJ, Xing FQ (2003) Expression profile of trophoblast invasion-associated genes in the pre-eclamptic placenta. British journal of biomedical science 60**:** 97-101

Parra-Cordero M, Rodrigo R, Barja P, Bosco C, Rencoret G, Sepulveda-Martinez A, Quezada S (2013) Prediction of early and late pre-eclampsia from maternal characteristics, uterine artery Doppler and markers of vasculogenesis during first trimester of pregnancy. Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology 41**:** 538-544

Rajakumar A, Whitelock KA, Weissfeld LA, Daftary AR, Markovic N, Conrad KP (2001) Selective overexpression of the hypoxia-inducible transcription factor, HIF-2alpha, in placentas from women with preeclampsia. Biology of reproduction 64**:** 499-506

Ramos EM, Hoffman D, Junkins HA, Maglott D, Phan L, Sherry ST, Feolo M, Hindorff LA (2014) Phenotype-Genotype Integrator (PheGenI): synthesizing genome-wide association study (GWAS) data with existing genomic resources. European journal of human genetics : EJHG 22**:** 144-147

Raymond D, Peterson E (2011) A critical review of early-onset and late-onset preeclampsia. Obstetrical & gynecological survey 66**:** 497-506

Redman C (2014) The six stages of pre-eclampsia. Pregnancy hypertension 4**:** 246

Redman CW, Sargent IL (2005) Latest advances in understanding preeclampsia. Science 308**:** 1592-1594

Roberts JM, Gammill HS (2005) Preeclampsia: recent insights. Hypertension 46**:** 1243-1249

Roberts JM, Hubel CA (2009) The two stage model of preeclampsia: variations on the theme. Placenta 30 Suppl A**:** S32-37

Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics 26**:** 139-140

Saito S, Nakashima A (2013) Review: The role of autophagy in extravillous trophoblast function under hypoxia. Placenta 34 Suppl**:** S79-84

Saito S, Nakashima A (2014) A review of the mechanism for poor placentation in early-onset preeclampsia: the role of autophagy in trophoblast invasion and vascular remodeling. Journal of reproductive immunology 101-102**:** 80-88

Salonen Ros H, Lichtenstein P, Lipworth L, Cnattingius S (2000) Genetic effects on the liability of developing pre-eclampsia and gestational hypertension. American journal of medical genetics 91**:** 256-260

Schizophrenia Working Group of the Psychiatric Genomics C (2014) Biological insights from 108 schizophrenia-associated genetic loci. Nature 511**:** 421-427

Shamshirsaz AA, Paidas M, Krikun G (2012) Preeclampsia, hypoxia, thrombosis, and inflammation. Journal of pregnancy 2012**:** 374047

Shiho Nagayama, Akihide Ohkuchi, RieUsui, Shigeki Matsubara, Suzuki M (2014) The Role of the Father in the Occurrence of Preeclampsia. Medical Journal of Obstetrics and Gynecology 2

Sitras V, Paulssen RH, Gronaas H, Leirvik J, Hanssen TA, Vartun A, Acharya G (2009) Differential placental gene expression in severe preeclampsia. Placenta 30**:** 424-433

Soto E, Romero R, Kusanovic JP, Ogge G, Hussein Y, Yeo L, Hassan SS, Kim CJ, Chaiworapongsa T (2012) Late-onset preeclampsia is associated with an imbalance of angiogenic and anti-angiogenic factors in patients with and without placental lesions consistent with maternal underperfusion. The journal of maternal-fetal & neonatal medicine : the official journal of the European Association of Perinatal Medicine, the Federation of Asia and Oceania Perinatal Societies, the International Society of Perinatal Obstet 25**:** 498-507

Supek F, Bosnjak M, Skunca N, Smuc T (2011) REVIGO summarizes and visualizes long lists of gene ontology terms. PloS one 6**:** e21800

Tal R (2012) The role of hypoxia and hypoxia-inducible factor-1alpha in preeclampsia pathogenesis. Biology of reproduction 87**:** 134

Tejera E, Cruz-Monteagudo M, Burgos G, Sanchez ME, Sanchez-Rodriguez A, Perez-Castillo Y, Borges F, Cordeiro M, Paz YMC, Rebelo I (2017) Consensus strategy in genes prioritization and combined bioinformatics analysis for preeclampsia pathogenesis. BMC medical genomics 10**:** 50

Tong J, Zhao W, Lv H, Li WP, Chen ZJ, Zhang C (2018) Transcriptomic Profiling in Human Decidua of Severe Preeclampsia Detected by RNA Sequencing. Journal of cellular biochemistry 119**:** 607-615

Trosse C, Ravneberg H, Stern B, Pryme IF (2007) Vectors encoding seven oikosin signal peptides transfected into CHO cells differ greatly in mediating Gaussia luciferase and human endostatin production although mRNA levels are largely unaffected. Gene regulation and systems biology 1: 303-312

Tsai S, Hardison NE, James AH, Motsinger-Reif AA, Bischoff SR, Thames BH, Piedrahita JA (2011) Transcriptional profiling of human placentas from pregnancies complicated by preeclampsia reveals disregulation of sialic acid acetylesterase and immune signalling pathways. Placenta 32: 175-182

Ursini G, Punzi G, Chen Q, Marenco S, Robinson JF, Porcelli A, Hamilton EG, Mitjans M, Maddalena G, Begemann M et al (2018) Convergence of placenta biology and genetic risk for schizophrenia. Nature medicine 24: 792-801

Vaiman D, Calicchio R, Miralles F (2013) Landscape of transcriptional deregulations in the preeclamptic placenta. PloS one 8: e65498

Vaiman D, Miralles F (2016) An Integrative Analysis of Preeclampsia Based on the Construction of an Extended Composite Network Featuring Protein-Protein Physical Interactions and Transcriptional Relationships. PloS one 11: e0165849

Valenzuela FJ, Vera J, Venegas C, Pino F, Lagunas C (2015) Circadian System and Melatonin Hormone: Risk Factors for Complications during Pregnancy. Obstetrics and gynecology international 2015: 825802

van Dijk M, Mulders J, Poutsma A, Konst AA, Lachmeijer AM, Dekker GA, Blankenstein MA, Oudejans CB (2005) Maternal segregation of the Dutch preeclampsia locus at 10q22 with a new member of the winged helix gene family. Nature genetics 37: 514-519

Varkonyi T, Nagy B, Fule T, Tarca AL, Karaszi K, Schonleber J, Hupuczi P, Mihalik N, Kovalszky I, Rigo J, Jr. et al (2011) Microarray profiling reveals that placental transcriptomes of early-onset HELLP syndrome and preeclampsia are similar. Placenta 32 Suppl: S21-29

von Dadelszen P, Magee LA, Roberts JM (2003) Subclassification of preeclampsia. Hypertension in pregnancy 22: 143-148

Voss A, Baumert M, Baier V, Stepan H, Walther T, Faber R (2006) Autonomic cardiovascular control in pregnancies with abnormal uterine perfusion. American journal of hypertension 19: 306-312

Walker CK, Krakowiak P, Baker A, Hansen RL, Ozonoff S, Hertz-Picciotto I (2015) Preeclampsia, placental insufficiency, and autism spectrum disorder or developmental delay. JAMA pediatrics 169: 154-162

Weiler J, Tong S, Palmer KR (2011) Is fetal growth restriction associated with a more severe maternal phenotype in the setting of early onset pre-eclampsia? A retrospective study. PloS one 6: e26937

Wu CS, Sun Y, Vestergaard M, Christensen J, Ness RB, Haggerty CL, Olsen J (2008) Preeclampsia and risk for epilepsy in offspring. Pediatrics 122: 1072-1078

Wu Y, Yao YG, Luo XJ (2017) SZDB: A Database for Schizophrenia Genetic Research. Schizophrenia bulletin 43**:** 459-471

Yu G, Wang LG, Han Y, He QY (2012) clusterProfiler: an R package for comparing biological themes among gene clusters. Omics : a journal of integrative biology 16**:** 284-287

Yu G, Wang LG, Yan GR, He QY (2015) DOSE: an R/Bioconductor package for disease ontology semantic and enrichment analysis. Bioinformatics 31**:** 608-609

Zeeman GG (2009) Neurologic complications of pre-eclampsia. Seminars in perinatology 33**:** 166-172

Zhao L, Bracken MB, DeWan AT (2013) Genome-wide association study of pre-eclampsia detects novel maternal single nucleotide polymorphisms and copy-number variants in subsets of the Hyperglycemia and Adverse Pregnancy Outcome (HAPO) study cohort. Annals of human genetics 77**:** 277-287

Zheng G, Tu K, Yang Q, Xiong Y, Wei C, Xie L, Zhu Y, Li Y (2008) ITFP: an integrated platform of mammalian transcription factors. Bioinformatics 24**:** 2416-2417

# Figure legends

**Figure 1. Clinical subtypes and sample clustering using RNA-seq data.**

A   Clinical subtypes of PE patients: early-onset severe PE (EOSPE), late-onset severe PE (LOSPE), late-onset mild PE (LOMPE). The numbers under each subtype are the numbers of placentae (not in brackets) and the numbers of patients (in brackets).

B   Principal component analysis (PCA). The top two principal components in this dataset represent up to 30% of all variations. Samples of EOSPE clustered on the right side (within the dashed circle), while samples of LOSPE and LOMPE are clustered together with normal samples on the left side.

C   Heatmap of sample-sample distance clustering using the Ward's method. The EOSPE samples are grouped together as the branch on the left, roughly corresponding to the samples in the right area of the PCA plot. The LOSPE and LOMPE samples are clustered together with normal samples on the right side. Only four LOSPE samples were clustered with EOSPE.

**Figure 2. Differentially expressed genes in preeclamptic placentae.**

A    The differentially expressed genes (DEGs) for each group were identified by comparing data from all PE samples combined or EOSPE, LOSPE and LOMPE samples with normal samples using two methods, DESeq2 and edgeR.

B   The red bars represent up-regulated genes, the blue bars represent down-regulated genes and the grey bars non-coding genes. Numbers represent DEGs in each group, and numbers in brackets represent protein-coding DEGs.

C   Examples of DEGs verified by qPCR. *FOXO3*: RNA-seq quantification boxplot at top-left panel and qPCR validation result at bottom-left panel. *SPX*: RNA-seq quantification boxplot at top-right panel and qPCR validation result at bottom-right panel.

D    The enrichment of PE associated genes in DEGs. PE-associated genes were curated from literature (Table EV4). Of the genes that appear once, twice and three or more times in different literature, 31.7%, 59.9% and 84.4% were recovered by our DEGs. The X-axis represents the number of times of a gene reported in the literature, the Y-axis represents the fraction of reported PE-associated genes under each category, with red bars representing ratio of genes that appear once, twice, and three or more times in different literature, and gray bars the expected ratios.

E   The enrichment of known PE-associated genes in protein-coding DEGs in EOSPE (2,298 genes), LOSPE (322 genes), LOMPE (39 genes) and all DEGs (2,405 genes). We collected 1,177 PE-associated genes from the literature (Table EV4), and calculated the their enrichment in DEGs of different groups: red bar represents the fraction of PE-associated genes in all DEGs (19.3%, $P$ = 2.2e-16), purple bar EOSPE (19.5%, $P$ = 2.2e-16), pink bar LOSPE (26.7%, $P$ = 2.2e-16) and light-green bar LOMPE (7.7%, $P$ = 0.4261), compared to gray bar which represents the expected ratio (6.1%) of 1,177 PE-associated genes to 19,351 human coding genes (GRCH38.p12 ENSEMBL Gene V93). One-sided Fisher's exact test was used to compare the nodes kept with the nodes

removed. Error bars represent the standard error of the fraction, estimated using bootstrapping method with 100 resamplings.

**Figure 3. Enrichment of KEGG pathways and GO molecular function (MF) terms with differentially expressed protein-coding genes of EOSPE and LOSPE.**

A   Enriched KEGG pathways with protein-coding DEGs of EOSPE and LOSPE.

B   Enriched GO MF terms with protein-coding DEGs of EOSPE and LOSPE. Dots represent the enriched KEGG pathways or GO MF terms with description of each pathway and term; colors represent scale of *P*-values, the sizes of dots represent ratio of DEGs in corresponding pathways and GO terms, and red stars indicate transporter terms in GO MF. Detailed information is listed in Table EV5.

**Figure 4. Functional classification of differentially expressed transporter genes.**

A   Venn diagram for the overlapping genes between the transporter genes and DEGs. Of the identified transporter genes, 205 were differentially expressed in EOSPE and 23 genes in LOSPE.

B   Fraction of transport genes in DEGs. The transporter genes were collected from Gene Ontology. Gray bar represents the expected fraction (7.87%) of transporter genes in all GO annotated genes (19,738), the fraction of transporter genes in the DEGs of EOSPE samples (8.92%), the DEGs of LOSPE (7.14%). The transporter genes were significantly enriched in the DEGs of EOSPE, but not in LOSPE. One-sided Fisher's exact test was used to compare the nodes kept with the nodes removed. Error bars represent the standard error of the fraction, estimated using bootstrapping method with 100 resamplings.

C   Gene-GO term bipartite network for enriched GO BP terms and DEGs in EOSPE. Blue dashed circles show 5 biological process clusters involved in different transportation processes of substrates and materials. Group I: ion channels; group II: macro-autophagy; group III: transmembrane

transportation of nonlipid nutrients; group IV: transmembrane transportation of lipids; group V in transportation of hormone.

D Gene-GO term bipartite network for enriched GO BP terms and DEGs in LOSPE. The group A: ion channels; group B: transportation of chloride, carboxylic acid, cAMP and anion. Blue dashed circles show 2 biological process clusters. Red nodes represent up-regulated transporter genes and blue nodes down-regulated transporter genes, and yellow diamond nodes represent GO BP terms. The edges indicate the GO BP terms the genes belong to.

**Figure 5. TF-target networks for EOSPE and LOSPE.**

A Workflow for extracting TF-target network for EOSPE and LOSPE. We collected TF-target data from 6 databases (Consortium, 2012; Han et al, 2018; Han et al, 2015; Jiang et al, 2007; Marbach et al, 2016; Neph et al, 2012; Zheng et al, 2008), and then matched the 1,639 reliable human transcription factors (TFs) (Lambert et al, 2018) to their targets and constructed a TF-target network containing 996 TFs and 16,538 target genes.

B The comparison between curated TFs and DEGs in EOSPE and LOSPE: 91 TFs differentially expressed in EOSPE only; 8 TFs in LOSPE only; 14 TFs shared by both.

C The comparison between TF targets of the DEGs in EOSPE and LOSPE: 1,882 (81.9%) DEGs in EOSPE regulated by the 105 TFs, and 216 (67.1%) DEGs in LOSPE by the 22 TFs.

D TF-target network for EOSPE. The bar chart shows the numbers of differentially expressed TFs (105) and their differentially expressed targets (1,882) in EOSPE.

E TF-target network for LOSPE. The bar chart shows the numbers of differentially expressed TFs (22) and their differentially expressed targets (216) in LOSPE. Red circles represent up-regulated genes, blue circles down-regulated genes, red triangles represent up-regulated TFs and blue triangles down-regulated TFs. Edges with arrows represent regulation of TFs to their target genes.

# Tables and their legends

Tables are provided in excel files.

**Table EV1. Clinical information.**

Sheet1: Clinical information comparisons among groups of patients with EOSPE, LOSPE or LOMPE and normal subjects;

Sheet2: Clinical information comparison between subclass-1 and subclass-2 PE sample groups.

**Table EV2. Raw counts of RNA-seq data.**

Sheet1: Raw counts of each gene for two cord blood samples;

Sheet2: Raw counts of each gene for 65 placental samples before and after removing blood contamination.

**Table EV3. Differentially expressed genes (DEGs).**

Sheet1: Gene biotypes and direction of difference of DEGs;

Sheet2: DEGs of All PE;

Sheet3: DEGs of EOSPE;

Sheet4: DEGs of LOSPE;

Sheet5: DEGs of LOMPE;

Sheet6: Q-PCR validation.

**Table EV4. PE-associated genes curated from literature.**

Sheet1: Literature information, number of PE-associated genes and datasets used in these studies;

Sheet2: PE-associated genes with times found in the literature and the overlap with DEGs of EOSPE, LOPSE and LOMPE.

**Table EV5. Enriched Pathways and GO terms.**

Sheet1: Enriched KEGG pathways in the DEGs of the EOSPE and LOSPE;

Sheet2: Enriched KEGG pathways in the DEGs of the EOSPE-only, LOSPE-only and shared genes in Sheet1;

Sheet3: Enriched KEGG pathways in the up- and down-regulated DEGs of the EOSPE and up-regulated DEGs of the LOSPE;

Sheet4: Enriched GO BP terms;

Sheet5: Enriched GO CC terms;

Sheet6: Enriched GO MF terms;

Sheet7: Representative term groups in BP, CC and MF.

**Table EV6. Transporter genes.**

Sheet1: Information for 1,554 transporter genes downloaded from GO and overlapped genes in the DEGs of the EOSPE and the LOSPE;

Sheet2: Information for differentially expressed transporter genes in the EOSPE and the LOSPE.

**Table EV7. Enriched GO terms and pathways for transporter genes in DEGs of EOSPE and the LOSPE.**

Sheet1: Information in the GO-term-to-gene-bipartite networks of EOPSE and LOPSE;

Sheet2: Enriched 397 BP terms table in EOSPE;

Sheet3: Enriched 93 BP terms table in LOPSE;

Sheet4: Enriched KEGG pathways in differentially expressed transporter genes of the EOSPE.

**Table EV8. Transcription factors and regulatory network.**

Sheet1: Information for 1,639 transcription factors (TFs) and the overlap with the DEGs of the EOSPE and the LOSPE. The first six columns of this table were from Table EV1 of Lambert's study (Lambert et al, 2018);

Sheet2: TF-target regulatory networks for EOSPE. The first column of table is TF and the second column is target regulated by TF;

Sheet3: TF-target regulatory networks for LOSPE. The first column of table is TF and the second column is target regulated by TF;

Sheet4: Enriched biological pathways in TF-target regulatory networks of EOSPE and LOSPE obtained using ReactomeFI plugin of Cytoscape;

Sheet5: TFs involved in enriched pathways.

**Table EV9. Tools, databases and primer sequences.**

Sheet1: Tools for analyses;

Sheet 2: Databases;

Sheet 3: Primer sequences for qPCR.

# Expanded View Figure legends

**Figure EV1. C-reactive protein levels in clinical subtypes and sample clustering analysis based on RNA-seq data of early-onset severe, late-onset severe and mild samples**

A    Box-plot of c-reactive protein levels in LOMPE, EOSPE and LOSPE. The normal range of c-reactive protein levels is 0-5mg/L.

B    Heatmap of sample-sample distance clustering for disease samples (late-onset severe, mild and early-onset severe samples) using Ward.D. Two clear subclasses were observed, where most of late-onset severe together with mild samples clustered as subclass-1 and early-onset severe together with four late-onset severe samples clustered as subclass-2.

**Figure EV2. Identification and validation of differentially expressed genes, and long-term consequence for the offspring.**

A    Venn diagrams for identifying differentially expressed genes using DESeq2 and edgeR methods in four comparison groups. Left Venn diagrams are number of up-regulated genes detected by DESeq2 and edgeR and right Venn diagrams are number of down-regulated genes detected by DESeq2 and edgeR. The overlapped genes were taken as the final differentially expressed genes in each comparison group. Light brown represents method of DESeq2; dark brown method of edgeR.

B    Box-plots for four up-regulated genes identified from RNA-seq and q-PCR validation.

C    Box-plots for six down-regulated genes identified from RNA-seq and q-PCR validation.

D    Schizophrenia-associated genes were enriched in DEGs of preeclampsia. The schizophrenia associated genes collected from classical schizophrenia databases and literature showed significantly enrichment in DEGs in all PE samples combined, EOPSE or LOPSE, whereas no enrichment in DEGs in LOMPE. Gray bar represents the expected fraction (17.8%) of schizophrenia associated genes in all protein-coding genes (19,351), the fraction of schizophrenia-associated genes in DEGs in all PE samples (20.7%), the DEGs of EOSPE (20.54%), LOSPE (22.98%) and LOMPE (15.4%) were presented as red, purple, pink and light green bars. The *P*-

value was calculated using the Fisher's exact test, and error bars represent the standard error of the fraction, estimated using bootstrapping with 100 resamplings.

**Figure EV3. Functional enrichment analysis of differentially expressed genes.**

A   Enrichment of DEGs in KEGG pathways for EOSPE-only, common or LOSPE-only DEGs. Venn-diagram shows the overlaps between DEGs of EOSPE and LOSPE.

B-C   Enrichment of DEGs in KEGG pathways for up- and down-regulated DEGs in the EOSPE (B) and LOPSE (C).

D   Enrichment of DEGs in GO BP terms for DEGs of EOSPE and LOSPE.

E   Enrichment of DEGs in GO CC terms for DEGs of EOSPE and LOSPE. Dot colors indicate enrichment *P*-values and dot sizes represent gene ratios in the enriched pathways.

**Figure EV4. Gene-pathway bipartite network of DEGs and the enriched KEGG pathways in EOSPE.**

Red nodes represent up-regulated genes and green nodes are down-regulated genes in EOSPE; and yellow diamond nodes represent names of KEGG pathways. The edges connect the DEGs to the KEGG pathways the genes belong to.

**Figure EV5. TF-target network modules for DGEs in EOSPE and LOSPE.**

A  TF-target network module for AMPK signaling pathway in EOSPE.

B  TF-target network module for FoXO signaling pathway in EOSPE.

C  TF-target network module for PI3K-Akt signaling pathway in LOSPE.

D   TF-target network module for immune and autoimmune signaling pathway in LOSPE. All genes in TF-target network modules are differentially expressed genes in EOSPE and LOSPE, respectively. The TF targets are members of the pathways.

**Figure 1. Clinical subtypes and sample clustering using RNA-seq data.**

A   Clinical subtypes of PE patients: early-onset severe PE (EOSPE), late-onset severe PE (LOSPE), late-onset mild PE (LOMPE). The numbers under each subtype are the numbers of placentae (not in brackets) and the numbers of patients (in brackets).

B   Principal component analysis (PCA). The top two principal components in this dataset represent up to 30% of all variations. Samples of EOSPE clustered on the right side (within the dashed circle), while samples of LOSPE and LOMPE are clustered together with normal samples on the left side.

C   Heatmap of sample-sample distance clustering using the Ward's method. The EOSPE samples are grouped together as the branch on the left, roughly corresponding to the samples in the right area of the PCA plot. The LOSPE and LOMPE samples are clustered together with normal samples on the right side. Only four LOSPE samples were clustered with EOSPE.

**Figure 2. Differentially expressed genes in preeclamptic placentae.**

A  The differentially expressed genes (DEGs) for each group were identified by comparing data from all PE samples combined or EOSPE, LOSPE and LOMPE samples with normal samples using two methods, DESeq2 and edgeR.

B  The red bars represent up-regulated genes, the blue bars represent down-regulated genes and the grey bars non-coding genes. Numbers represent DEGs in each group, and numbers in brackets represent protein-coding DEGs.

C  Examples of DEGs verified by qPCR. FOXO3: RNA-seq quantification boxplot at top-left panel and qPCR validation result at bottom-left panel. SPX: RNA-seq quantification boxplot at top-right panel and qPCR validation result at bottom-right panel.

D  The enrichment of PE associated genes in DEGs. PE-associated genes were curated from literature (Table EV4). Of the genes that appear once, twice and three or more times in different literature, 31.7%, 59.9% and 84.4% were recovered by our DEGs. The X-axis represents the number of times of a gene reported in the literature, the Y-axis represents the fraction of reported PE-associated genes under each category, with red bars representing ratio of genes that appear once, twice, and three or more times in different literature, and gray bars the expected ratios.

E  The enrichment of known PE-associated genes in protein-coding DEGs in EOSPE (2,298 genes), LOSPE (322 genes), LOMPE (39 genes) and all DEGs (2,405 genes). We collected 1,177 PE-associated genes from the literature (Table EV4 ), and calculated the their enrichment in DEGs of different groups: red bar represents the fraction of PE-associated genes in all DEGs (19.3%, *P* = 2.2e-16), purple bar EOSPE (19.5%, *P* = 2.2e-16), pink bar LOSPE (26.7%, *P* = 2.2e-16%) and light-green bar LOMPE (7.7%, *P* = 0.4261), compared to gray bar which represents the expected ratio (6.1%) of 1,177 PE-associated genes to 19,351 human coding genes (GRCH38.p12 ENSEMBL Gene V93). One-sided Fisher's exact test was used to compare the nodes kept with the nodes removed. Error bars represent the standard error of the fraction, estimated using bootstrapping method with 100 resamplings.

**Figure 3. Enrichment of KEGG pathways and GO molecular function (MF) terms with differentially expressed protein-coding genes of EOSPE and LOSPE.**

A   Enriched KEGG pathways with protein-coding DEGs of EOSPE and LOSPE.

B   Enriched GO MF terms with protein-coding DEGs of EOSPE and LOSPE. Dots represent the enriched KEGG pathways or GO MF terms with description of each pathway and term; colors represent scale of *P*-values, the sizes of dots represent ratio of DEGs in corresponding pathways and GO terms, and red stars indicate transporter terms in GO MF. Detailed information is listed in Table EV5.

**Figure 4. Functional classification of differentially expressed transporter genes.**

A   Venn diagram for the overlapping genes between the transporter genes and DEGs. Of the identified transporter genes, 205 were differentially expressed in EOSPE and 23 genes in LOSPE.

B   Fraction of transport genes in DEGs. The transporter genes were collected from Gene Ontology. Gray bar represents the expected fraction (7.87%) of transporter genes in all GO annotated genes (19,738), the fraction of transporter genes in the DEGs of EOSPE samples (8.92%), the DEGs of LOSPE (7.14%). The transporter genes were significantly enriched in the DEGs of EOSPE, but not in LOSPE. One-sided Fisher's exact test was used to compare the nodes kept with the nodes removed. Error bars represent the standard error of the fraction, estimated using bootstrapping method with 100 resamplings.

C   Gene-GO term bipartite network for enriched GO BP terms and DEGs in EOSPE. Blue dashed circles show 5 biological process clusters involved in different transportation processes of substrates and materials. Group I: ion channels; group II: macro-autophagy; group III: transmembrane transportation of nonlipid nutrients; group IV: transmembrane transportation of lipids; group V in transportation of hormone.

D   Gene-GO term bipartite network for enriched GO BP terms and DEGs in LOSPE. The group A: ion channels; group B: transportation of chloride, carboxylic acid, cAMP and anion. Blue dashed circles show 2 biological process clusters. Red nodes represent up-regulated transporter genes and blue nodes down-regulated transporter genes, and yellow diamond nodes represent GO BP terms. The edges indicate the GO BP terms the genes belong to.

**Figure 5. TF-target networks for EOSPE and LOSPE.**

A   Workflow for extracting TF-target network for EOSPE and LOSPE. We collected TF-target data from 6 databases (Consortium, 2012; Han *et al.*, 2018; Han *et al.*, 2015; Jiang *et al.*, 2007; Marbach *et al.*, 2016; Neph *et al.*, 2012; Zheng *et al.*, 2008), and then matched the 1,639 reliable human transcription factors (TFs) (Lambert *et al.*, 2018) to their targets and constructed a TF-target network containing 996 TFs and 16,538 target genes.

B   The comparison between curated TFs and DEGs in EOSPE and LOSPE: 91 TFs differentially expressed in EOSPE only; 8 TFs in LOSPE only; 14 TFs shared by both.

C   The comparison between TF targets of the DEGs in EOSPE and LOSPE: 1,882 (81.9%) DEGs in EOSPE regulated by the 105 TFs, and 216 (67.1%) DEGs in LOSPE by the 22 TFs.

D   TF-target network for EOSPE. The bar chart shows the numbers of differentially expressed TFs (105) and their differentially expressed targets (1,882) in EOSPE.

E   TF-target network for LOSPE. The bar chart shows the numbers of differentially expressed TFs (22) and their differentially expressed targets (216) in LOSPE. Red circles represent up-regulated genes, blue circles down-regulated genes, red triangles represent up-regulated TFs and blue triangles down-regulated TFs. Edges with arrows represent regulation of TFs to their target genes.