Headwaters fed by subterranean ice: potential climate refugia for mountain stream

communities?

Lusha M. Tronstad[1,*], Scott Hotaling[2,*], J. Joseph Giersch[3], Oliver J. Wilmot[4], and Debra S.

Finn[5]

Affiliations:

[1] Wyoming Natural Diversity Database, University of Wyoming, Laramie, WY, USA

[2] School of Biological Sciences, Washington State University, Pullman, WA, USA;

scott.hotaling@wsu.edu

[3] U.S. Geological Survey, Northern Rocky Mountain Science Center, West Glacier, MT, USA;

jgiersch@usgs.gov

[4] Wyoming Natural Diversity Database, University of Wyoming, Laramie, WY, USA;

owilmot@uwyo.edu

[5] Department of Biology, Missouri State University, Springfield, MO, USA;

DFinn@MissouriState.edu

* Contributed equally

Correspondence: Lusha M. Tronstad, Wyoming Natural Diversity Database, University of

Wyoming, 1000 E. University Ave., Laramie, WY 82071 USA; Phone: (307) 766-3115; Email:

tronstad@uwyo.edu

Running head: Mountain stream climate refugia

ABSTRACT

Near-term extirpations of macroinvertebrates are predicted for mountain streams worldwide as a

warming climate drives the recession of high-elevation ice and snow. However, hydrological

sources likely vary in their resistance to climate change and thus streams fed by more resistant

sources could persist as climate refugia for imperiled biota. In 2015-2016, we measured habitat

characteristics and quantified macroinvertebrate community structure along six alpine streams in

the Teton Range, Wyoming, USA. Strong differences in habitat characteristics (e.g., temperature,

bed stability, conductivity) confirmed three major stream sources: surface glaciers, perennial

snowfields, and subterranean ice. Subterranean ice-fed streams – termed "icy seeps" – appear

common in the Teton Range and elsewhere yet are globally understudied. Midges in the family

Chironomidae dominated our study sites, representing 78.6% of all specimens sampled, with

nematodes, caddisflies (*Neothremma*), and mayflies (*Epeorus*) also common. At the community-

scale, glacier- and snowmelt-fed streams differed significantly in multivariate space, with icy-

seep communities intermediate between them, incorporating components of both assemblages.

Because the thermal environment of subterranean ice, including rock glaciers, is decoupled

from large-scale climatic conditions, we predict that icy seeps will remain intact longer than

streams fed by surface ice and snow. Furthermore, our results suggest that icy seeps are suitable

habitat for many macroinvertebrates occupying streams fed by vulnerable hydrological sources.

Thus, icy seeps may act as key climate refugia for mountain stream biodiversity, an idea in need

of further investigation.

                                    INTRODUCTION

The highest rates of climate change are occurring above the permanent treeline in alpine

and arctic ecosystems (Bradley et al. 2006). In the Rocky Mountains, warming is proceeding two

to three times more quickly than the global average (Hansen et al. 2005, Pederson et al. 2010),

resulting in extensive loss of glaciers and long-term snowpack (Hall & Fagre 2003). Streams fed

by permanent ice may exhibit short-term increases in flow as air temperatures rise and source ice

melting accelerates, but eventually they will shift to reduced flow and warm with the potential

for intermittency or drying permanently (Hotaling et al. 2017). As climate change proceeds,

invertebrate diversity at the mountain range scale is predicted to decrease due to both overall loss

of habitat and summit traps, where the highest-altitude species and communities have nowhere

left to disperse as warmer conditions and lower elevation communities shift upward. Biodiversity

loss will be compounded by the loss of specific aquatic habitat types, particularly the unique

conditions associated with meltwater from once-permanent hydrological sources like glaciers,

snowfields, or subterranean ice (Brown et al. 2007, Milner et al. 2009, Jacobsen et al. 2012, Finn

et al. 2013, 2014, Hotaling et al. 2017). Given that many alpine stream communities appear

uniquely adapted to cold thermal regimes (but see Hotaling et al. 2019c), they are likely to be

highly vulnerable to climate change as meltwater sources are lost (Giersch et al. 2017, Lencioni

2018). However, because alpine streams are heterogeneous with respect to hydrological source,

potential also exists to identify stream types that may be locally buffered from broad-scale

climate patterns and therefore could represent climate refugia for alpine stream biota (Morelli et

al. 2016).

A major, long-term focus in alpine stream biology has been understanding the links

between hydrological sources, the in-stream conditions they promote, and resident biotic

communities (Ward 1994, Hotaling et al. 2017). According to primary hydrological source

following Ward (1994), three types of alpine streams have historically been recognized: surface

glacier-fed, snowmelt-fed, and groundwater-fed streams. A fourth, understudied stream type also

exists – icy seeps – which are fed by subterranean ice (Hotaling et al. 2017, Hotaling et al.

2019a). In mountain ecosystems, the most common form of subterranean ice are rock glaciers,

masses of debris-covered ice that act as conveyor belts moving fallen rock and other debris

slowly downhill (Anderson et al. 2018; Jones et al. 2019). There may be more than 10,000 rock

glaciers in the western United States (Johnson 2018) and they are similarly common worldwide

(e.g., Lilleøren et al. 2011, Scotti et al. 2013, Charbonneau & Smith. 2018). In contrast, there are

~1,250 surface glaciers and ~3,750 perennial snowfields in the western United States (Fountain

et al. 2017). Due to insulating debris cover, rock glaciers are largely decoupled from external

conditions (e.g., warm summer air temperatures; Clark et al. 1994, Anderson et al. 2018, Knight

et al. 2019), and should persist on the landscape longer than surface glaciers and snowfields. Icy

seeps, the outflow of rock glaciers and similar ice features, may therefore act as climate refugia

for cold-adapted stream biodiversity (Brighenti et al. 2019a,b; Hotaling et al. 2019a).

Like many mountain ranges worldwide, virtually nothing is known of alpine stream

ecology and biodiversity in the Teton Range, a granite-dominated subrange of the Rocky

Mountains. Previous studies of montane (2,000-3,000 m), but not alpine, streams revealed

considerable macroinvertebrate diversity in the region's higher elevation streams (Tronstad et al.

2016). Generally speaking, groundwater aquifers appear rare on the Teton Range massif, and

thus groundwater-fed streams are also rare (L.M.T., personal observation). Groundwater-fed

streams are considered the most resistant alpine stream type to warming because they are not

directly influenced by surface ice (e.g., Milner et al. 2009, Jacobsen et al. 2012). With a paucity

of groundwater-fed streams in the region, stream biodiversity in the Teton Range may be

especially vulnerable to climate change.

In this study, we addressed two major objectives. First, we made the first assessment of

alpine stream macroinvertebrate diversity in the Teton Range. Second, we explored associations

between primary hydrological sources (surface glaciers, snowfields, and subterranean ice) and

community structure. Specifically, we asked if benthic communities associated with icy seeps

have substantial taxonomic overlap with communities linked to sources more vulnerable to

climate change (e.g., glacier and snowfields). Our study provides an important first perspective

on an urgent need in freshwater ecology (see Brighenti et al., 2019a); testing whether an

underappreciated but globally common alpine stream type – icy seeps fed by subterranean ice –

may act as key refugia for mountain aquatic biodiversity threatened by global change. Our

results also provide new insight into the biodiversity of one of North America's flagship

protected areas, Grand Teton National Park, and neighboring wilderness areas.

108   **MATERIALS AND METHODS**

109   Study area

During the summers of 2015 and 2016 (26 July-10 August), we sampled six streams in

the Teton Range of Grand Teton National Park and the adjacent Jedediah Smith Wilderness in

northwestern Wyoming, USA (Figure 1; Table 1). Study streams were selected to span the

breadth of alpine hydrological sources we have observed in the Teton Range and included two

streams fed by surface glaciers ('glacier-fed' hereafter), two fed by subterranean ice ('icy seep'

hereafter) and two fed by permanent snowpack ('snowmelt-fed'). In 2015, we sampled both

upstream (near the source) and downstream sites on each stream (Figure 1). On average, upper

sites were 111 m higher in elevation and 690 m in stream distance from lower sites (Table 1). In

2016, we re-sampled the upper sites with the same methods to assess inter-annual variability. We

focused on upper sites because they were as 'true' to primary hydrological source as possible

while lower sites inherently reflected various degrees of mixing among sources. In both years,

snow depth in the range was lower than average (152 cm in May, 1981-2010) with 2015 and

2016 at 63.3% and 80% of normal, respectively (Teton Pass, USDA SNOTEL).

124   Environmental data

At each site, we measured several environmental variables to characterize local habitat

and evaluate whether in-stream environmental conditions varied among stream types. We

measured water temperature for a full year (2015-2016) with *in situ* loggers (HOBO Pro v2,

Onset Computer Corporation) that recorded temperature hourly. We measured specific

conductivity (SPC), oxidation-reduction potential (ORP), pH, and dissolved oxygen (DO) with a

Yellow Springs Instrument (YSI) Professional Plus multiparameter sonde calibrated at the

trailhead (SPC, ORP, and pH) or at each site (DO). We estimated streambed stability with a

modified version of the Pfankuch Index (PI) following Peckarsky et al. (2014). Total suspended

sediments (TSS) were calculated by filtering known volumes of streamwater through pre-

weighed filters (PALL Type A/E glass fiber filters) and measuring dry mass to the 10^{-5} grams.

We used analysis of variance (ANOVA) statistical tests and the R package 'plyr' (R Core

Development Team 2017; Wickham 2011) to characterize differences in environmental variables

(Pfankuch Index, specific conductivity and total suspended solids) among stream types and

between study sites. When stream type was significant ($\alpha = 0.05$), we used Tukey's HSD to

distinguish which stream types differed from one another ($P \leq 0.05$). Our use of Tukey's HSD

tests was highly conservative, given the small sample size of streams in this study.

Using the annual temperature data from each stream, we calculated mean temperature for

the entire year (T_{YEAR}), mean temperature between the summer solstice (21 June) and autumn

equinox (22 September; T_{SUMMER}), and maximum annual temperature range (T_{RANGE}). We also

estimated the date when seasonal snow covered (S_{ON}) and uncovered (S_{OFF}) each site by visually

inspecting thermographs from our *in situ* temperature loggers to approximate the date when

intraday thermal variation ceased and in-stream temperatures became constantly ~0°C (S_{ON}) or

the opposite occurred (S_{OFF}). An example thermograph indicating S_{ON} and S_{OFF} points is

provided in Figure S1. Using S_{ON} and S_{OFF}, we calculated $S_{DURATION}$, the total days a site was

snow covered between study years ($S_{DURATION} = S_{ON} - S_{OFF}$). Finally, we used Principle

Components Analysis (PCA; PC-ORD, McCune and Mefford 2006) to characterize the upper

sites on each stream according to four variables (SPC, TSS, PI, T_{RANGE}) which comprise a

modified 'glaciality index' (Ilg & Castella 2006). Because T_{SUMMER} was correlated with T_{RANGE},

we only included T_{RANGE} in the PCA. The glaciality index has been useful in characterizing

alpine stream hydrological sources globally (e.g., Finn et al. 2013, Cauvy-Fraunié et al. 2015,

Hotaling et al. 2019b).

157   Macroinvertebrate sampling

158   We quantitatively sampled benthic macroinvertebrates at each site using a Surber sampler

159   (Area: 0.09 m², Mesh size: 243 μm). At each location, a composite sample of 5-10 replicates was

160   collected depending on stream size, apparent biomass, and microhabitat diversity. Samples were

161   elutriated in the field to reduce the amount of inorganic substrate and stored in Whirl-Pak bags

162 with ~80% ethanol. In the laboratory, invertebrate samples were divided into large (>2 mm

163 mesh) and small (between 250 µm and 2 mm) fractions. For the large fraction, all invertebrates

164 were identified. The small fraction was subsampled using the record player method when

165 invertebrates were numerous (Waters 1969). Specimens were sorted, identified to the lowest

166 taxonomic level possible using keys in Merritt et al. (2008) and Thorp & Covich (2010), and

167 counted under a dissecting microscope. Insects were typically identified to genus when mature

168 specimens were present except Chironomidae which were classified as either Tanypodinae or

169 non-Tanypodinae. We estimated invertebrate density by summing the total number of individuals

170 for a given site and dividing by the area of streambed sampled. We calculated biomass by

171 measuring the length of the first 20 individuals of each taxon and then using length-mass

172 regressions to estimate individual biomass (Benke et al. 1999). We multiplied the mean

173 individual biomass for each taxon by the total number collected to estimate total biomass.

175             Biological data analysis

176    We used ANOVAs and Tukey's HSD tests performed on data summarized with 'plyr' (R

177 Core Development Team 2017; Wickham 2011) to characterize differences in invertebrate

178 density, biomass, and richness among stream types and study sites. To assess the relationship

179 between taxonomic richness or biomass with stream characteristics of interest, namely snow

180 cover ($S_{DURATION}$), temperature (T_{SUMMER}) and stability (Pfankuch Index), we performed both

181 Pearson and Spearman's rank-order correlations using the R package 'Hmisc' (Harrell Jr. 2013).

182 For correlation analyses, we focused exclusively on upper sites and averaged taxonomic richness

183 and biomass between 2015 and 2016.

We evaluated differences in community structure across streams, sites, and study years

using non-metric multidimensional scaling (NMS) with PC-ORD (McCune & Mefford 2006).

We $\log_{10} (n + 1)$ transformed density data for all taxa, removed rare taxa (either those private to a

single site in the matrix and/or representing < 1% of the total abundance), and used Sørensen's

dissimilarities to create distance matrices. We ran NMS analyses independently on two data

matrices: one including each of the upper and lower sites collected in 2015 only (N = 12 sites)

and the other including only the upper sites (sampled in both 2015 and 2016; N = 12 sites).

Dimensionality of the final solutions was chosen as the number of axes that produced the lowest

stress following 200 iterations. Following NMS we applied multi-response permutation

procedures (MRPP) in PC-ORD to assess whether there were differences in either community

structure and/or mean community distance within the following groups: upstream versus

downstream sites (2015 only) and among stream types for upper sites only (2015 and 2016). We

then used Indicator Species Analysis (ISA) in PC-ORD to assess whether any specific taxa in the

input matrices were indicative of (a) upper or lower sites or (b) a specific stream type. In ISA,

the indicator value (IV) for a taxon is the test statistic. The maximum IV is 100, which indicates

that a taxon is found in one group alone and is absent from other groups. We tested for

significance of all IVs using Monte Carlo randomizations with 15,000 permutations of the data

for each of the two tests (a and b above).

203   **RESULTS**

204   Environmental variation

Our upper sites clearly separated into three groups according to the glaciality index:

glacier-fed streams, snowmelt-fed streams, and icy seeps (Figure 2; Table 2). Glacier-fed streams

had less stable streambeds than both icy seeps and snowmelt-fed streams (F = 19.9, df = 2, P =

0.0001; Tukey's HSD, $P < 0.001$; Table 2). Annual temperature range (T_{RANGE}) was highest in

snowmelt-fed streams and lowest in icy seeps. Specific conductivity was highest in icy seeps

(SPC >100 μS cm^{-1} at upper sites; F = 30.6, df = 2, $P < 0.001$; Tukey's HSD, $P < 0.001$) and

lowest in glacier-fed streams (Table 2). More suspended solids (TSS) were present in glacier-fed

streams (mean = 0.157 g/L) compared to snowmelt-fed streams (F = 3.6, df = 2, $P = 0.44$;

Tukey's HSD, $P = 0.039$; Table 2). Summer temperatures (T_{SUMMER}) were lowest in glacier-fed

streams (mean = 1.6 °C) and icy seeps (mean = 2.2 °C), and higher in snowmelt-fed streams

(mean = 6.9 °C; Table 2). Upper sites were on average 1.2 °C colder in the summer and had less

stable stream beds than lower sites (mean PI, upper = 27; mean PI, lower = 20.33; Table 1).

Other environmental variables (i.e., DO, pH, ORP, days under snow) did not vary among stream

types or between upper and lower sites (Table 2).

220   Biological variation

We collected 35 total invertebrate taxa of which 28 were insects (Supplementary

Appendix 1). Insects composed 95% of the total mean densities and 92% of the total biomass.

While invertebrate densities varied across stream types, biomass and richness were highest in

snowmelt-fed streams (Figure 3A). At upper sites, total macroinvertebrate density did not differ

among stream types (F = 1.3, df = 2, $P = 0.31$; Figure 3A), but biomass was ~7x higher in

snowmelt-fed streams compared to glacier-fed streams and icy seeps (F = 7.1, df = 2, $P < 0.009$;

Tukey's HSD, $P \leq 0.02$; Figure 3B). Additionally, invertebrate richness was ~2x higher in

snowmelt-fed streams than glacier-fed streams and icy seeps (F = 10.4, df = 2, $P < 0.002$,

Tukey's HSD, $P < 0.008$; Figure 3C). We observed generally higher and more variable

invertebrate densities (F = 2.7, df = 1, P = 0.13; Figure 3D) and higher biomass (F = 2.6, df = 1,

P = 0.13; Figure 3E) at lower sites. We also observed ~50% more taxa at lower sites (F = 8.5, df

= 1, P = 0.012; Figure 3F). Only T_{SUMMER} was significantly correlated with richness or biomass

(Richness: Pearson r = 0.81, P = 0.002; Biomass: Pearson r = 0.63, P = 0.038; Figure 4).

Spearman's rank-order correlations of the same relationships exhibited similar patterns, again

with only T_{SUMMER} exhibiting significant correlations with richness (Spearman's r = 0.89, P =

0.003) and biomass (Spearman's r = 0.67, P = 0.023).

The most common invertebrates in high-elevation Teton streams were non-Tanypodinae

midges (78.6% of all specimens sampled; >100 individuals/m^2 at all sites; Appendix 1) followed

by *Neothremma* (caddisflies), nematodes, and *Epeorus* (mayflies). Additionally, 45.7% of taxa

had densities >10 individual/m^2 among sites. Three taxa (Perlodidae, *Zapada* and Empididae)

were present at 67% of sites and twelve taxa (34.3%), including *L. tetonica*, were only found at a

single site (Appendix 1). Overall, we observed more stoneflies (6 taxa) than caddisflies (5 taxa)

and mayflies (4 taxa; Appendix 1). Only two taxa (midges and Collembola) were collected at

upper sites of both glacier-fed streams (Appendix 1). Two taxa were also present at both upper

icy seep sites (midges and the stonefly genus, *Zapada*) and four taxa were present in upper

snowmelt-fed sites [midges, stoneflies (Perlodidae), caddisflies (*Allomyia*), and flatworms

(Turbellaria)]. Of those, only Turbellaria were present at consistently high numbers (e.g., >20

individuals/m^2 per site; Appendix 1). No taxon exclusively occurred in one stream type and three

occurred in upper sites of all stream types (midges, *Allomyia* caddisflies, and *Helodon* black

flies; Appendix 1).

The most stable NMS solution comparing community structure between upper and lower

sites was three-dimensional (stress = 9.6) with the first two axes explaining 88% of the total

variation. Communities from the six lower sites overlapped substantially in ordination space with

those from the six upper sites (MRPP $A = 0.013$; $P = 0.27$). However, the mean pairwise

community distance was greater among upper versus lower sites (0.58 vs. 0.47), a trend that is

apparent in the NMS bi-plot (Figure 5A). Indicator species analyses revealed no taxon with IV

scores approaching the maximum of 100 (Tables S2-S3), an expected result when two

assemblages substantially overlap in ordination space. Mean IV across all taxa was 44.3, and

Empididae had the maximum IV (78.9; $P = 0.025$) to lower sites.

The NMS analysis that included upper sites only (2015 and 2016) converged on a two-

dimensional solution as the most stable result (stress = 8.9). The two axes explained 86% of the

variation, with axis-1 explaining 70% alone. In general, communities occupying glacier-fed

streams had the lowest axis-1 values, communities in snowmelt-fed streams had the highest axis-

1 values and icy seep communities had intermediate axis-1 values (Figure 5B). MRPP results

suggested that communities occupying these three stream types were significantly different from

one another ($A = 0.19$; $P = 0.006$). Pairwise differences were strong between glacier-fed and

snowmelt-fed communities ($A = 0.21$; $P = 0.005$) and were weaker but significant for the two

pairs that included icy seep communities (icy seep vs. glacier-fed: $A = 0.11$; $P = 0.05$; icy seep

vs. snowmelt: $A = 0.12$; $P = 0.03$). Indicator species analysis (Table S3) revealed IV scores >80

for three taxa, of which one was indicative of glacier-fed streams (Collembola; IV = 85.0; $P =$

0.02) and two were indicative of snowmelt streams (Tipulidae, IV = 82.1, $P = 0.004$; *Allomyia*,

IV = 81.5, $P = 0.004$). Turbellaria were also weakly indicative of snowmelt streams (IV = 72.5;

$P = 0.011$).

275   **DISCUSSION**

As climate change proceeds and mountain glaciers recede, there is a need to develop a

clearer understanding of how patterns of extant biodiversity and habitat heterogeneity are linked

in high-elevation ecosystems. In this study, we provide the first description of macroinvertebrate

diversity in the high Teton Range, Wyoming, where three major alpine stream types exist:

glacier-fed streams, snowmelt-fed streams, and icy seeps. To the best of our knowledge, no

streams fed by groundwater aquifers have been documented in the alpine zone of the granitic

Teton Range. From a global perspective, icy seeps, which are fed by subterranean ice (primarily

rock glaciers), rather than aquifers of liquid water, are of particular interest as they are likely to

persist on the landscape longer than surface ice features (Hotaling et al. 2019a). For the Teton

Range, a scarcity of high-elevation, groundwater-fed streams suggests that biodiversity in the

region may be even more reliant on meltwater than similar mountain regions (e.g., European

ranges, Brown et al. 2007; tropical Andes, Finn et al. 2016; Glacier National Park, Giersch et al.

2017). Thus, icy seep-associated climate refugia may be particularly important to Teton alpine

stream biodiversity as climate change proceeds. In the streams sampled for this study, glacier-

and snowmelt-fed streams exhibited significantly different invertebrate communities; however,

icy seeps were intermediate between the two in terms of community structure and invertebrate

density, biomass, and richness. These results suggest that icy seep communities share some

characteristics with both glacier- and snowmelt-fed streams and have the potential to act as

climate refugia for at least a subset of the unique communities present in each of the more

vulnerable stream types. Thus, the potential for icy seeps and ice-influenced terrestrial refugia

(Millar et al. 2015) to buffer climate-induced biodiversity loss has profound, global implications.

The recession of meltwater sources is predicted to strongly affect downstream

invertebrate communities (Jacobsen et al. 2012). In the near term, rising in-stream temperatures

are expected as ice melt comprises ever smaller proportions of stream flow. In alpine streams

worldwide, warmer conditions have been correlated with increased species richness for microbial

diversity (e.g., Wilhelm et al. 2013, Hotaling et al. 2019a), diatoms (Fell et al. 2018), and

macroinvertebrates (Finn & Poff 2005, Jacobsen et al. 2012). Our study adds another line of

evidence to this global pattern as we detected a positive correlation between species richness and

mean summer temperature (T_{SUMMER}). We also observed greater richness at lower (mean = 15

taxa) versus upper (mean = 10 taxa) sites which aligns with, and extends, the conclusions of

Tronstad et al. (2016), the only other study to investigate longitudinal patterns of

macroinvertebrate richness in montane streams of the Teton Range. Indeed, we observed far

fewer taxa at our highest elevation sites (10 taxa at ~3,150 m) versus the highest elevation sites

included in Tronstad et al. (2016): 26 taxa at ~2,700 m.

Although local (alpha) diversity will likely increase with warming water temperatures,

among-stream (beta) diversity may decrease as more diverse, generalist communities shift

upstream and specialized cold-adapted taxa are lost, effectively homogenizing biological

diversity at the regional scale (Jacobsen et al. 2012, Wilhelm et al. 2013, Hotaling et al. 2017). In

the Teton Range, we observed greater beta diversity at upper versus lower sites, and ISA

revealed indicator taxa for snowmelt and glacier-fed streams, but not for icy seeps. Collectively,

these patterns suggest that in the Teton Range, like elsewhere in the world, heterogeneous

hydrological sources bolster regional-scale alpine stream biodiversity. Icy seep communities also

appeared to bridge the taxonomic gap between glacier- and snowmelt-fed streams, and this

pattern remained stable between years (Figure 5B). While clear patterns of community

dissimilarity existed among our sites, ascribing a specific driver to these differences is difficult.

Many factors likely shape community structure in headwater streams (and our estimation of it),

including the primary source and its associated environmental regime, geographic location, and

both the method and timing of sampling (e.g., a site recently uncovered from seasonal snowpack

is likely to have a distinct community when compared to a site that melted out much earlier).

Our study represents the first perspective of macroinvertebrate biodiversity in the highest

elevation streams of the Teton Range. As such, there are many areas where future study will

expand, and refine, our conclusions. First, we did not have the resources to identify midges

(family Chironomidae) to lower taxonomic resolution. Midges are diverse in alpine streams

around the world (e.g., Montagna et al. 2016), and are typically the most common taxon (a

pattern we also observed). Thus, the limited taxonomic resolution of midges in our study may

have influenced our conclusions. For instance, Finn & Poff (2011) found 22 midge species in

snowmelt-fed alpine streams of the Colorado Rockies. Given our focus on three stream types, we

might expect even greater midge species diversity in the Teton Range. Second, our focus on a

limited number of sites, lack of within-year temporal sampling, and use of a single sampling

method/microhabitat (Surber samples from riffle habitats) rather than many approaches/habitats

(e.g., Ghani et al. 2016, Tronstad & Hotaling 2017), limits our ability to confidently describe the

full invertebrate community at a given site. We hope to fill these gaps in the future.

Ultimately, the degree to which alpine streams will be affected by climate change in

terms of flow magnitude and persistence remains largely unknown. In general, studies of alpine

stream ecology operate under the assumption that perennial flow will continue in the decades to

come (e.g., Jacobsen et al. 2012), but this may not be the case (e.g., Haldorsen & Heim 1999,

Herbst et al. 2019, Siebers et al. 2019). Thus, the biological ramifications of declining meltwater

sources in places like the Teton Range, where groundwater aquifer-fed streams are scarce, may

be even more profound than in ranges with a more equitable distribution of alpine stream sources

(e.g., Glacier National Park, Giersch et al. 2017). Indeed, if alpine streams supported by surface

glaciers and permanent snowfields transition to intermittency or dry completely, the future of

biodiversity in these ecosystems may depend almost exclusively on icy seeps. Our study paired

with the broader glaciological literature suggests there is room for optimism. Icy seeps have the

potential to span a wide beta diversity profile, perhaps bridging the taxonomic gap between

glacier- and snowmelt-fed communities (e.g., as observed for microbial diversity, Hotaling et al.

2019a). Moreover, rock glaciers and other subterranean ice forms (e.g., periglacial taluses, Millar

et al. 2013) are common in alpine regions worldwide and likely the most resistant ice form to

future warming. Clearly, research focused on subterranean ice sources and associated icy seeps,

and specifically the biological communities and hydrological flows they support, represent a

pressing need in alpine stream biology. We suggest that future studies incorporate temporal

monitoring of multiple alpine stream types, including icy seeps, to separate year-to-year variation

from true temporal signals. Long-term data will also allow for explicit tests of how biodiversity

and habitat characteristics may be altered among alpine stream types as climate change proceeds.

360   **ACKNOWLEDGEMENTS**

Financial support was provided by The University of Wyoming-National Park Service Research

Station and the Teton Conservation District. We thank Lydia Zeglin for assistance in the field

and Hanna Foster, Logan Fox, Alexis Lester, Jackson Marr, Jake Ruthven, Joe Wannemuehler,

and Kara Wise for their help sorting invertebrate samples. Matthew Green provided helpful

comments on the manuscript. Any use of trade, firm, or product names is for descriptive

purposes only and does not imply endorsement by the U.S. Government.

LITERATURE CITED

Anderson, R. S., L. S. Anderson, W. H. Armstrong, M. W. Rossi & S. E. Crump. 2018.

Glaciation of alpine valleys: the glacier – debris-covered glacier – rock glacier

continuum. Geomorphology 311:127-142.

Benke, A. C., A. D. Huryn, L. A. Smock & J. B. Wallace. 1999. Length-mass relationships for

freshwater macroinvertebrates in North America with particular reference to the

Southeastern United States. Journal of the North American Benthological Society

18:308-343.

Bradley, R. S., M. Vuille, H. F. Diaz & W. Vergara. 2006. Threats to water supplies in the

tropical Andes. Science 312:1755-1756.

Brighenti, S., M. Tolotti, M. C. Bruno, G. Wharton, M. T. Pusch, & W. Bertoldi. 2019.

Ecosystem shifts in Alpine streams under glacier retreat and rock glacier thaw: A

review. *Science of the Total Environment* 675:542-559.

Brighenti, S., M. Tolotti, M. C. Bruno, M. Engel, G. Wharton, L. Cerasino, V. Mair, & W.

Bertoldi. 2019. After the peak water: the increasing influence of rock glaciers on alpine

river systems. *Hydrological Processes* 33:2804-2823.

Brown, L. E., D. M. Hannah & A. M. Milner. 2007. Vulnerability of alpine stream biodiversity

to shrinking glaciers and snowpacks. Global Change Biology 13:958-966.

Cauvy-Fraunié, S., , R. Espinosa, P. Andino, D. Jacobsen & O. Dangles. 2015. Invertebrate

metacommunity structure and dynamics in an Andean glacial stream network facing

climate change. PLoS ONE 10:e0136793.

Charbonneau, A. A. & D. J. Smith. 2018. An inventory of rock glaciers in the central British

390   Columbia Coast Mountains, Canada, from high resolution Google Earth imagery. Arctic,

391   Antarctic, and Alpine Research 50:e1489026.

392   Clark, D. H., M. M. Clark & A. R. Gillespie, 1994. Debris-covered glaciers in the Sierra Nevada,

393   California, and their implications for snowline reconstructions. Quaternary Research

394   41:139-153.

395   Fell, S. C., J. L. Carrivick, M. G. Kelly, L. Füreder & L. E. Brown. 2018. Declining glacier cover

396   threatens the biodiversity of alpine river diatom assemblages. Global Change Biology

397   24(12):5828-5840.

398   Finn, D. S., A. C. Encalada & H. Hampel. 2016. Genetic isolation among mountains but not

399   between stream types in a tropical high-altitude mayfly. Freshwater Biology 61(5):702-

400   714.

401   Finn, D. S., K. Khamis & A. M. Milner. 2013a. Loss of small glaciers will diminish beta

402   diversity in Pyrenean streams at two levels of biological organization. Global Ecology

403   and Biogeography 22:40-51.

404   Finn, D. S. & N. L. Poff. 2005. Variability and convergence in benthic communities along the

405   longitudinal gradients of four physically similar Rocky Mountain streams. Freshwater

406   Biology 50:243-261.

407   Finn, D. S. & N. L. Poff. 2011. Examining spatial concordance of genetic and species diversity

408   patterns to evaluate the role of dispersal limitation in structuring headwater

409   metacommunities. Journal of the North American Benthological Society 30:273-283.

410   Finn, D. S., C. Zamora-Muñoz, C. Múrria, M. Sáinz-Bariáin & J. Alba-Tercedor. 2014. Evidence

411   from recently deglaciated mountain ranges that *Baetis alpinus* (Ephemeroptera) could

412 lose significant genetic diversity as alpine glaciers disappear. Freshwater Science 33:207-

413 216.

414 Fountain, A. G., B. Glenn & H. J. Basagic. 2017. The geography of glaciers and perennial

415 snowfields in the American West. Arctic, Antarctic, and Alpine Research 49:391-410.

416 Ghani, W. M. H. W. A., C. S. M. Rawi, S. A. Hamid & S.A. Al-Shami. 2016. Efficiency of

417 different sampling tools for aquatic macroinvertebrate collections in Malaysian

418 streams. Tropical Life Sciences Research 27:115.

419 Giersch, J. J., S. Hotaling, R. P. Kovach, L. A. Jones & C. C. Muhlfeld. 2017. Climate-induced

420 glacier and snow loss imperils alpine stream insects. Global Change Biology 23:2577–

421 2589.

422 Giersch, J. J., S. Jordan, G. Luikart, L. A. Jones, F. R. Hauer & C. C. Muhlfeld. 2015. Climate-

423 induced range contraction of a rare alpine aquatic invertebrate. Freshwater Science 34:53-

424 65.

425 Haldorsen, S. & M. Heim, 1999. An Arctic groundwater system and its dependence upon

426 climatic change: An example from Svalbard. Permafrost and Periglacial Processes

427 10:137-149.

428 Hall, M. H. P. & D. B. Fagre. 2003. Modeled climate-induced glacier change in Glacier National

429 Park, 1850-2100. Bioscience 53:131-140.

430 Hansen, J., L. Nazarenko, R. Ruedy, M. Sato, J. Willis, A. Del Genio, D. Koch, A. Lacis, K. Lo,

431 S. Menon, T. Novakov, J. Perlwitz, G. Russell, G. A. Schmidt & N. Tausnev. 2005.

432 Earth's energy imbalance: confirmation and implications. Science 308:1431-1435.

433 Harrell, F. E. J. 2013. Hmisc: Harrell miscellaneous. R package version 3.12-2. [Computer

434 software]. Available from http://cran.R-project.Org/web/packages/Hmisc.

Herbst, D. B., S. D. Cooper, R. B. Medhurst, S. W. Wiseman, & C. T. Hunsaker (2019). Drought

ecohydrology alters the structure and function of benthic invertebrate communities in

mountain streams. *Freshwater Biology* 64:886-902.

Hotaling, S., D. S. Finn, J. J. Giersch, D. W. Weisrock & D. Jacobsen. 2017. Climate change

and alpine stream biology: progress, challenges, and opportunities for the future.

Biological Reviews 92:2024-2045.

Hotaling, S., M. Foley, L. Zeglin, D. Finn, L. Tronstad, J. Giersch, C. Muhlfeld & D. W.

Weisrock. 2019a. Microbial assemblages reflect environmental heterogeneity in alpine

streams. Global Change Biology.

Hotaling, S., J. J. Giersch, D. S. Finn, L. M. Tronstad, S. Jordan, L. E. Serpa, R. G. Call, C. C.

Muhlfeld & D. W. Weisrock. 2019b. Congruent population genetic structure but differing

depths of divergence for three alpine stoneflies with similar ecology and geographic

distributions. Freshwater Biology 64:335-347.

Hotaling, S., A. A. Shah, K. L. McGowan, L. M. Tronstad, J.J. Giersch, D. S. Finn, H. A.

Woods, M. E. Dillon & J. L. Kelley (2019c). Mountain stoneflies may tolerate warming

streams: evidence from organismal physiology and gene expression. *bioRxiv*.

Ilg, C. & E. Castella. 2006. Patterns of macroinvertebrate traits along three glacial stream

continuums. Freshwater Biology 51:840-853.

Jacobsen, D., A. M. Milner, L. E. Brown & O. Dangles. 2012. Biodiversity under threat in

glacier-fed river systems. Nature Climate Change 2:361–364.

Johnson, G. F. 2018. Rock Glaciers of the Contiguous United States: Spatial Distribution,

Cryospheric Context, and Riparian Vegetation. Dissertations and Theses 4507:1-147.

Jones, D. B., S. Harrison, K. Anderson, & W. B. Whalley. 2019. Rock glaciers and mountain

hydrology: A review. *Earth-Science Reviews*.

Knight, J., S. Harrison & D. B. Jones. 2019. Rock glaciers and the geomorphological evolution

of deglacierizing mountains. Geomorphology 324:14-24.

Lencioni, V. 2018. Glacial influence and stream macroinvertebrate biodiversity under climate

change: Lessons from the Southern Alps. Science of The Total Environment 622:563-

575.

Lilleøren, K. S. & B. Etzelmüller. 2011. A regional inventory of rock glaciers and ice☐cored

moraines in Norway. Geografiska Annaler: Series A, Physical Geography 93:175-191.

McCune, B. & M. Mefford. 2006. PC-ORD. Multivariate analysis of ecological data. Version 5.

Merritt, R. W., K. W. Cummins & M. B. Berg. 2008. An Introduction To The Aquatic Insects of

North America, 4th Edition. Kendall/Hunt Publishing Company, Dubuque, Iowa.

Millar, C. I., R. D. Westfall & D. L. Delany. 2013. Thermal and hydrologic attributes of rock

glaciers and periglacial talus landforms: Sierra Nevada, California, USA. Quaternary

International 310:169-180.

Millar, C. I., R. D. Westfall, A. Evenden, J. G. Holmquist, J. Schmidt-Gengenbach, R. S.

Franklin, J. Nachlinger & D. L. Delany. 2015. Potential climatic refugia in semi-arid,

temperate mountains: Plant and arthropod assemblages associated with rock glaciers,

talus slopes, and their forefield wetlands, Sierra Nevada, California, USA. Quaternary

International 387:106-121.

Milner, A. M., L. E. Brown & D. M. Hannah. 2009. Hydroecological response of river systems

to shrinking glaciers. Hydrological Processes 23:62-77.

Montagna, M., V. Mereghetti, V. Lencioni & B. Rossaro. 2016. Integrated taxonomy and DNA

barcoding of alpine midges (Diptera: Chironomidae). PLoS One, 11(3), e0149673.

Morelli, T. L., C. Daly, S. Z. Dobrowski, D. M. Dulen, J. L. Ebersole, S. T. Jackson, J. D.

Lundquist, C. I. Millar, S. P. Maher, W. B. Monahan, K. R. Nydick, K. T. Redmond, S.

C. Sawyer, S. Stock & S. R. Beissinger. 2016. Managing climate change refugia for

climate adaptation. PLoS ONE 11:e0159909.

Peckarsky, B. L., A. R. McIntosh, S. C. Horn, K. McHugh, D. J. Booker, A. C. Wilcox, W.

Brown & M. Alvarez. 2014. Characterizing disturbance regimes of mountain streams.

Freshwater Science 33:716-730.

Pederson, G. T., L. J. Graumlich, D. B. Fagre, T. Kipfer & C. C. Muhlfeld. 2010. A century of

climate and ecosystem change in Western Montana: what do temperature trends portend?

Climatic Change 98:133-154.

R Development Core Team. 2017. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org.

Scotti, R., F. Brardinoni, S. Alberti, P. Frattini & G. B. Crosta. 2013. A regional inventory of

rock glaciers and protalus ramparts in the central Italian Alps. Geomorphology 186:136-

149.

Siebers, A. R., A. Paillex, & C. T. Robinson. 2019. Flow intermittency influences the trophic

base, but not the overall diversity of alpine stream food webs. *Ecography*.

Thorp, J. H. & A. P. Covich (eds). 2010. Ecology and Classification of North American

Freshwater Invertebrates. 3rd edition. Academic Press, London.

Tronstad, L. M. & S. Hotaling. 2017. Long-term trends in aquatic ecosystem bioassessment

metrics are not influenced by sampling method: empirical evidence from the Niobrara

River. Knowledge & Management of Aquatic Ecosystems 418:28.

Tronstad, L. M., S. Hotaling & J. C. Bish. 2016. Longitudinal changes in stream invertebrate

assemblages of Grand Teton National Park, Wyoming. Insect Conservation and Diversity

9:320-331.

Ward, J. V., 1994. Ecology of alpine streams. Freshwater Biology 32:277-294.

Waters, T. F., 1969. Subsampler for dividing large samples of stream invertebrate drift.

Limnology and Oceanography 14:813-815.

Wickham, H. 2011. The Split-Apply-Combine Strategy for Data Analysis. 2011 40:29.

Wilhelm, L., G. A. Singer, C. Fasching, T. J. Battin & K. Besemer. 2013. Microbial biodiversity

in glacier-fed streams. ISME Journal 7:1651-1660.

TABLES

Table 1. Key characteristics of our study streams and sites in the Teton Range, Wyoming, USA.

Distances to source (D) are cumulative. Both elevation and D are in meters.

Stream	Code	Site	Type	Lat., Long.	Elev.	D
Petersen Glacier	PG	Upper	Glacier-fed	43.782, -110.846	2,922	51
		Lower	Glacier-fed	43.785, -110.841	2,900	673
Middle Teton	MT	Upper	Glacier-fed	43.728, -110.795	2,955	178
		Lower	Glacier-fed	43.725, -110.790	2,802	780
South Cascade Creek	SCC	Upper	Icy seep	43.722, -110.838	3,152	165
		Lower	Icy seep	43.729, -110.837	2,943	1,051
Wind Cave	WC	Upper	Icy seep	43.666, -110.961	2,692	29
		Lower	Icy seep	43.667, -110.955	2,564	144
S. Fork Teton Creek	SFT	Upper	Snowmelt	43.691, -110.843	2,987	1,227
		Lower	Snowmelt	43.693, -110.859	2,881	2,757
N. Fork Teton Creek	NFT	Upper	Snowmelt	43.777, -110.860	2,955	9
		Lower	Snowmelt	43.775, -110.861	2,910	393

Table 2. Environmental characteristics of the streams and sites included in this study. T_{YEAR}: mean temperature for a calendar year. T_{SUM}: mean temperature between the summer solstice (21 June) and autumn equinox (22 September). T_{RANGE}: the difference between the maximum and minimum temperatures recorded in the stream during the calendar year. S_{ON} and S_{OFF} indicate the day of the year when a site was covered (S_{ON}) or uncovered (S_{OFF}) by snow according to thermographs. $S_{DURATION}$ (S_{DUR} below) is the approximate total number of days each site spent under snow in 2015-2016. Abbreviations and units: temperature (T; °C), dissolved oxygen (DO, percent saturation), specific conductivity (SPC; μS cm^{-1}), oxidation-reduction potential (ORP, mV), total suspended solids (TSS; g/L; upper sites only) and the Pfankuch Index (PI; higher values correspond to a less stable streambed). All data (except for S_{ON}, S_{OFF}, and S_{DUR}) are for 2015 only. Stream codes are provided in Table 1 and parentheses after the stream name indicate the stream type (GF: glacier-fed; SM: snowmelt-fed; IS: icy seep).

Stream	Site	T_{YEAR}	T_{SUM}	T_{RANGE}	S_{ON}	S_{OFF}	S_{DUR}	DO	SPC	pH	ORP	TSS	PI
PG (GF)	Upper	0.6	1.4	13.7	306	195	254	118	3.7	7.9	128.4	0.21	42
	Lower	1.9	6.0	16.2	303	128	190	103	9.9	8.6	24.8	--	23
MT (GF)	Upper	0.7	1.7	10.3	300	128	193	101	4.8	8.4	41.2	0.105	42
	Lower	1.5	4.1	13.6	296	150	219	100	5.7	8.9	62.9	--	30
SCC (IS)	Upper	0.5	1.5	10.0	298	204	271	104	113.2	8.5	-4.6	0.033	21
	Lower	1.5	4.6	13.0	297	155	223	104	110.3	8.4	58.4	--	16
WC (IS)	Upper	1.7	2.9	4.4	No cover			102	176.8	8.6	47	0.023	18
	Lower	1.9	4.2	7.3	309	98	154	99	175.2	8.4	77.2	--	18
SFT (SM)	Upper	1.9	7.1	17.7	300	174	239	101	74.3	8.6	-0.2	0.013	24
	Lower	Logger failed						101	114.7	8.5	23	--	17
NFT (SM)	Upper	2.5	6.6	15.2	300	125	190	101	11.1	8.5	64.9	0.024	15
	Lower	2.5	6.0	10.7	328	133	170	96	11.8	7.8	89.6	--	18

FIGURES

Figure 1. All streams and sites in the Teton Range, Wyoming, USA, included in this study.

Upper sites were selected to be as close to the primary hydrological source as possible. Numbers

in the top panel correspond to satellite imagery (DigitalGlobe, 15 October 2015, bottom panel)

with stream channel (light blue) and sample sites (red dots) marked.

Figure 2. Principal components analysis (PCA) results showing environmental variation among

the upper sites according to the four variables of the glaciality index [specific conductivity,

streambed stability (Pfankuch Index), suspended solids, and temperature (T_{RANGE})].

Figure 3. Macroinvertebrate density, biomass and richness among sites categorized by

hydrologic source (a-c; upper sites only in 2015 and 2016) or site (d-f; 2015 and 2106 data for

upper sites, 2015 data only for lower sites). Bold lines are median values, lower and upper box

limits indicate the 25th and 75th percentiles, respectively, and whiskers represent the lower and

upper limits of the data (excluding outliers which are shown as circles).

Figure 4. Bivariate plots of biomass (a-c) and richness (d-f) versus the Pfankuch Index, a

measure of stream bed stability (higher = less stable), $S_{DURATION}$ (the approximate number of

days each stream was snow-covered in 2015-2016), and T_{SUMMER} (mean stream temperature

between the summer and autumn solstices). Significant Pearson correlations ($P < 0.05$) are in

bold with trendlines shown. The number of data points varies across plots due to a lack of snow

cover at upper Wind Cave during the study period and a failed temperature logger at lower South

Fork Teton Creek (see Table 2).

Figure 5. Non-metric multidimensional scaling (NMS) plots of macroinvertebrate communities

comparing (a) upper (circles) versus lower (triangles) sites in 2015 and (b) upper sites only

between 2015 (circles) and 2016 (squares). Colored polygons in (a) reflect the breadth of NMS

space occupied by upper (dashed lines, green fill) and lower (solid lines, blue fill) sites.