Relationship between spiking activity and simultaneously recorded fluorescence signals in transgenic mice expressing GCaMP6

Lawrence Huang[1]*, Ulf Knoblich[1]*, Peter Ledochowitsch[1], Jérôme Lecoq[1], R. Clay Reid[1], Saskia E. J. de Vries[1], Michael A. Buice[1], Gabe J. Murphy[1], Jack Waters[1], Christof Koch[1], Hongkui Zeng[1]#, Lu Li[1,2,3,4]#

1 Allen Institute for Brain Science, Seattle, WA 98109, USA
2 Guangdong Provincial Key Laboratory of Malignant Tumor Epigenetics and Gene Regulation, Sun Yat-sen Memorial Hospital, Sun Yat-sen University, Guangzhou, Guangdong Prov., China 510120
3 Medical Research Center, Sun Yat-sen Memorial Hospital, Sun Yat-sen University, Guangzhou, Guangdong Prov., China 510120
4 Centre for Brain Science and Brain-Inspired Intelligence, Guangdong-Hongkong-Macao Greater Bay Area, China

*These authors contributed equally
#Correspondence should be addressed to hongkuiz@alleninstitute.org (H.Z.) or lilu67@mail.sysu.edu.cn (L.L.)

Keywords

Abstract

Two-photon calcium imaging is often used with genetically encoded calcium indicators (GECIs)

to investigate neural dynamics, but the relationship between fluorescence and action potentials

(spikes) remains unclear. Pioneering work linked electrophysiology and calcium imaging *in vivo*

with viral GECI expression, albeit in a small number of cells. Here we characterized the spike-

fluorescence transfer function *in vivo* of 91 layer 2/3 pyramidal neurons in primary visual cortex

in four transgenic mouse lines expressing GCaMP6s or GCaMP6f. We found that GCaMP6s

cells have spike-triggered fluorescence responses of larger amplitude, lower variability and

greater single-spike detectability than GCaMP6f cells. Single spike detection rates differed

substantially across neurons in each line. They declined from ~40-90% at 5% false positive rate

under high-resolution imaging to ~10-15% when imaging hundreds of neurons across a larger

field of view. Our dataset thus provides quantitative insights to support more refined inference

of neuronal activity from calcium imaging data.

Introduction

Genetically encoded calcium indicators (GECIs) are widely used with two-photon (2-p) laser scanning microscopy to report neuronal activity within local populations *in vivo* (Luo et al., 2018). By reporting changes in calcium-dependent fluorescence, this optical approach is minimally invasive and enables simultaneous measurement of activity from hundreds or even thousands of neurons at single-cell resolution. As such, calcium imaging has become an attractive alternative to extracellular electrophysiology for surveying large neuronal populations of genetically identified cells, in particular over multiple sessions. Indeed, using a newly developed GECI such as GCaMP6s, fluorescence changes associated with isolated single spikes (action potentials) *in vivo* can be reliably detected when imaged at sufficiently high spatiotemporal resolution (Chen et al., 2013).

However, despite recent advances in imaging approaches and GECI development, calcium imaging remains an indirect measure of a neuron's spiking activity. Inferring the underlying spike train or firing rate from calcium imaging remains challenging (Theis et al., 2016; Berens et al., 2018), in part because the relationship between spiking activity and calcium dynamics is complex and has not been fully characterized. The spike-to-calcium fluorescence transfer function depends on several factors, including (1) the relationship between changes in GECI fluorescence and changes in intracellular calcium concentration, and buffering effects (Rose et al., 2014); (2) the neuronal cell types expressing the GECI, due to differences in intracellular calcium dynamics; (3) GECI expression level (*e.g.* differential viral uptake and expression when using viral GECI delivery); and (4) the behavioral state of the animal, due to changes in calcium dynamics, *e.g.* by neuromodulators (Nadim and Bucher, 2014). Together, these variables underlie the difficulty of linking calcium imaging studies to the large body of existing knowledge concerning spiking activity.

Compared to viral expression, transgenic mouse lines offer convenience (e.g. bypassing virus injection and associated procedures) and achieve more uniform GECI expression in genetically defined neuronal populations (Madisen et al., 2015; Daigle et al., 2018). This should permit a more straightforward comparison of activity across cells and animals. Using our intersectional transgenic mouse lines that enable Cre recombinase-dependent expression of GCaMP6s or GCaMP6f, we simultaneously characterized the spiking activity and fluorescence of individual GECI-expressing pyramidal neurons in layer (L) 2/3 of mouse primary visual cortex (V1). Consisting of 91 neurons from 4 mainstream transgenic lines, this ground truth dataset provides

quantitative insight into the relationship between *in vivo* spiking activity and observed

fluorescence signals, and will aid the interpretation of existing and future calcium imaging

datasets.

Results

Dataset overview

To characterize the single-cell transfer function between observed fluorescence signals and

underlying spikes *in vivo*, we performed simultaneous calcium imaging and cell-attached

recordings in V1 L2/3 excitatory pyramidal neurons in anesthetized mice (**Figure 1A**, see

Methods). The mice were from 4 transgenic lines, including 2 GCaMP6s-expressing lines,

Emx1-IRES-Cre;Camk2a-tTA;Ai94 (referred to as Emx1-s in this study) and *Camk2a-tTA;tetO-*

GCaMP6s (tetO-s), and 2 GCaMP6f-expressing lines, *Emx1-IRES-Cre;Camk2a-tTA;Ai93*

(Emx1-f) and *Cux2-CreERT2;Camk2a-tTA;Ai93* (Cux2-f) (**Table 1**). A total of 237 neurons, all of

which had apparent baseline fluorescence which was clearly excluded from the nuclei, were

randomly selected to record and image for spontaneous activity or visually-evoked responses.

To directly compare our results to virally-expressed GCaMP6f and GCaMP6s (Chen et al.,

2013), calcium imaging was performed at high optical zoom focused on individual cells (field of

view of ~20 x 20 µm at ~158 frames per second; fps). We also patched and imaged a subset of

these neurons at a lower zoom factor, *i.e.* one at which the responses of many neurons can be

characterized in parallel (field of view of ~400 x 400 µm at ~30 fps), allowing direct comparison

of calcium fluorescence responses acquired at higher spatiotemporal resolution with that more

commonly employed by other studies including our Brain Observatory dataset (de Vries et al.,

2019).

A large number (>550) of individual recording/imaging sessions, 2-4 min each, were obtained

from these 237 cells (multiple sessions were collected for some cells in cases where the patch

remained stable). Upon careful inspection of these sessions, we selected 91 cells with high-

quality recording and imaging conditions from the 4 mouse lines (**Table 1**) for analysis, each

with only one recording/imaging session for unbiased sampling.

Mouse line	Acronym	GECI	# Mice	# Cells			
				Spon	Visstim	Total	Low-zoom
Emx1-IRES-Cre;Camk2a-tTA;Ai94	Emx1-s	GCaMP6s	7	21	11	32	10
Camk2a-tTA;tetO-GCaMP6s	tetO-s	GCaMP6s	2	6	0	6	0

Emx1-IRES-Cre;Camk2a-tTA;Ai93	Emx1-f	GCaMP6f	6	9	17	26	10*
Cux2-CreERT2;Camk2a-tTA;Ai93	Cux2-f	GCaMP6f	7	4	23	27	17
Total			22	40	51	91	37

Table 1. Dataset. Sample size for each mouse line. Spon and Visstim: spontaneous and

visually-evoked recordings, respectively. Total: total number of cells in high-zoom dataset. Low-

zoom: cells imaged at both high- and low- optical zooms. *One Emx1-f cell was imaged at the

same low optical zoom but with smaller field of view and at ~60 fps; see **Results**. The duration

of the exclusion and summation windows used for each mouse line is described in **Methods**

and **Figure 1B**.

Constructing single-cell spike-to-calcium fluorescence response curves

Throughout the manuscript, "spike" and "spiking activity" always refer to fast, presumably Na+

mediated, all-or-none electrical events. To delineate the relationship between spiking activity

and GCaMP6 fluorescence, we focused on isolated spiking events whose calcium responses

were well-separated from, and thus minimally contaminated by, those of adjacent spikes. A

spiking event is defined as a group of spikes within a spike summation window (150 ms and 50

ms for GCaMP6s and GCaMP6f, respectively) with no spikes in the pre-event and post-event

exclusion windows (150 ms and 50 ms pre and post respectively for GCaMP6s, and 50 ms both

pre and post (after the last spike) for GCaMP6f) (**Figure 1B**). We summed the number of spikes

within each spiking event, aligned fluorescence responses to the first spike within the event, and

computed the peak fluorescence change (dF/F peak) during the calcium response window (200

ms for GCaMP6s, 50 ms or 75 ms for GCaMP6f in single-spike or multi-spike events) (**Figure**

1B-D; see **Methods** for spike exclusion, spike summation, and calcium response windows).

Spiking events were binned based on the number of spikes within each event. Following this

approach, we constructed single-cell spike-to-calcium fluorescence response curves (dF/F peak

as a function of the number of spikes, with a minimum requirement of 3 events per bin; **Figure**

2). Events with >5 spikes were excluded from analysis due to the low frequency of such events

(*e.g.*, ≥3 6-spike events were observed in 2 out of 32 Emx1-s cells, 0 out of 6 tetO-s cells, 5 out

of 26 Emx1-f cells, and 1 out of 27 Cux2-f cells; also see **Figure 6C, i** below).

For calcium imaging data, we first examined the effect of fluorescence background subtraction

on the peak dF/F signal. Following methods for virally expressed GCaMP6 (Chen et al., 2013),

the background neuropil signal was estimated as the mean fluorescence of all pixels within 20-

μm from the cell center, excluding the selected cell, and neuropil subtraction was performed as:

$F_{corrected}(t) = F_{measured}(t) - r \times F_{neuropil}(t)$, where r was the neuropil contamination ratio (**Figure 2–**

figure supplement 1). Neuropil subtraction using a constant r-value for all cells (ranging from

0.1 to 0.7, where 0.7 was used for virally expressed GCaMP6) did not consistently decrease

within-cell variability of dF/F peak (*i.e.* variability across spiking events) (**Figure 2–figure**

supplement 1D). Given the potential for artificially boosting dF/F due to lower baseline

fluorescence (see example cell in **Figure 2–figure supplement 1B-C**), we chose not to perform

neuropil correction in this study (however, see **Figure 3–figure supplement 3** and **Figure 5–**

figure supplement 3 below).

Population spike-to-calcium fluorescence response curves

To compare the relationship between spikes and fluorescence for the 4 mouse lines, population

response curves were constructed by averaging together responses of individual neurons

(**Figure 3A**). Spontaneous and visually-evoked responses were similar (**Figure 3–figure**

supplement 1) even though they were not recorded from the same cell and/or animal, and were

therefore pooled in the population responses. Data from mice anesthetized with isoflurane (n =

77 cells from 18 mice) and urethane (n = 14 cells from 4 mice; see **Methods**) were also pooled

as their response curves and firing rates were consistent with each other (**Figure 3–figure**

supplement 2).

Of the 4 mouse lines, tetO-s exhibited the largest single-spike fluorescence response (dF/F

peak: 13.6±3.7%, mean ± sd), followed by Emx1-s (7.8±2.6%), Cux2-f (6.0±1.6%), and Emx1-f

(4.5±1.6%) (**Figure 3A-D**). The response curve slopes (representing the average dF/F increase

per spike, computed from linear regression of dF/F peak against the number of spikes) of the 4

lines were comparable to each other, with tetO-s being somewhat higher (**Figure 3E**).

For a more direct comparison with that of viral GCaMP6 (Chen et al., 2013), we additionally

analyzed our data using a neuropil contamination ratio of 0.7 (**Figure 3–figure supplement 3**).

Aggregating spiking events across cells, mean dF/F peak in response to 5-spike events within

150 ms and 50 ms for GCaMP6s and GCaMp6f, respectively, was 135% for Emx1-s, 117% for

tetO-s (130% in response to 4-spike events), 130% for Emx1-f, and 93% for Cux2-f. In

comparison, Chen et al. reported ~200% and ~100% mean dF/F peak in response to 5-spike

events within 250 ms for viral GCaMP6s and GCaMP6f, respectively. Without neuropil

subtraction, mean dF/F peak in response to 5-spike events was 62% for Emx1-s, 73% for tetO-
s, 62% for Emx1-f, and 52% for Cux2-f (**Figure 3B**, bottom).

Fluorescence response variability
Within-cell variability of dF/F peak, as measured by the mean coefficient of variation across all
spiking events, was significantly larger in GCaMP6f-expressing cells compared to that in
GCaMP6s (**Figure 4A**). Likewise, within-cell variability for single-spike events was significantly
larger in GCaMP6f-expressing cells compared to that in GCaMP6s. In GCaMP6f-expressing
cells (but not GCaMP6s-expressing cells), single-spike events exhibited significantly larger
within-cell variability compared to multi-spike events (**Figure 4B**). Between-cell (cell-to-cell)
variability of dF/F peak was generally similar across the 4 lines (**Figure 4C**).

To determine the signal-to-noise ratio (SNR) and whether single-spike events could be
distinguished from zero-spike events (imaging noise), we first identified no-spike intervals of ≥1
s, separated by ≥4 s and ≥1 s from previous and subsequent spikes, respectively, for analysis
(**Figure 4–figure supplement 1A**). We then randomly sampled the corresponding calcium
fluorescence traces to obtain snippets of fluorescence with the same length as single-spike
events, and computed dF/F peak and the standard deviation of dF/F (an estimate of the noise
floor) using the same calcium response windows as single-spike events (50 ms and 200 ms for
GCaMP6f and GCaMp6s, respectively; **Figure 4–figure supplement 1B**). The mean single-
spike response was different from the mean imaging noise in 90% or 27 out of 30 Emx1-s cells
with at least 1 no-spike interval (out of 32 total cells), 100% or 4 out of 4 tetO-s cells with at least
1 no-spike interval (out of 6 total cells), 96% or 22 out of 23 Emx1-f cells with at least 1 no-spike
interval (out of 26 total cells), and 83% or 19 out of 23 Cux-f cells with least 1 no-spike interval
(out of 27 total cells) (**Figure 4–figure supplement 1C-E**).

Emx1-f and tetO-s cells exhibited the lowest and highest noise floor, respectively. Normalizing
the single-spike dF/F peak by the noise floor, the single-spike SNR for GCaMP6s was
significantly larger than that of GCaMP6f (**Figure 4–figure supplement 2A**), although a
difference in single-spike SNR was not observed between Emx1-s and tetO-s as in the single-
spike dF/F peak (**Figure 3D**). The population response curve expressed in SNR (**Figure 4–
figure supplement 2B**) largely resembled that expressed in dF/F peak (**Figure 3B**), with the
exception of tetO-s cells, whose SNR was lower due to a higher noise floor. However, tetO-s,

with a small sample size (n = 4 cells with at least 1 no-spike interval), also had few multi-spike

events (e.g. only 1 cell with 4-spike events).

Spike detection

We next examined spike detection at the single-trial level. Mean (± sd) detection rates were

63±25% and 94±11% for Emx1-s, 87±15% and 100% for tetO-s, 48±21% and 82±26% for

Emx1-f, and 40±27% and 74±22% for Cux2-f for single-spike and 2-spike events, respectively,

at 5% false positive rate (**Figure 5**). Spike detection rates differed substantially between cells,

from 0 to 100% for isolated spikes detected with GCaMP6s at 1% false positive rate (**Figure 5**).

Single-spike detection rate was ≥90% in 13% (4 of 30) Emx1-s cells, 75% (3 of 4) tetO-s cells,

4% (1 of 23) Emx1-f cells and 0% (0 of 22) Cux2-f cells, at 5% false positive rate with no

neuropil subtraction.

Previous studies have documented substantial contamination of somatic traces with

fluorescence from the surrounding neuropil, due to the extended nature of the microscope point

spread function. Neuropil contamination is often removed by subtracting a scaled version of the

neuropil fluorescence from the somatic fluorescence. The scale factor is often referred to as the

r-value and many studies employ the same r-value, often 0.7, for all neurons (Akerboom et al.,

2012). For each neuron, we found the optimum r-value, defined as the r-value which resulted in

the maximum number of detected spikes at the minimum false positive rate and calculated as

the r-value that maximized the area under the ROC curve. Optimal r-values differed greatly

between cells. Mean±sd optimal r-values were 0.39±0.30 for Emx1-s (range 0 to 0.8, with r<0.5

in 14 of 30 cells or 47%), 0.28±0.31 for tetO-s (range 0 to 0.7, with r<0.5 in 3 of 4 cells or 75%),

0.43±0.33 for Emx1-f (range 0 to 1, with r<0.5 in 13 of 22 cells or 59%), and 0.33±0.36 for Cux2-

f (range 0 to 1, with r<0.5 in 15 of 22 cells or 68%) (**Figure 5–figure supplement 1**). In the

Allen Brain Observatory, r-values were calculated individually for each neuron and follow similar

distributions to these optimal r-values. At optimal r-values, the single-spike detection rate at 5%

false positive rate was 76±21% (mean±sd) for Emx1-s, 92±11% for tetO-s, 52±23% for Emx1-f,

and 40±27% for Cux2-f.

Spiking characteristics

Spontaneous or visually evoked firing rates of individual cells, computed from the total number

of spikes detected during a recording, were similar across mouse lines (Emx1-s: 1.17±0.71 Hz,

mean±sd; tetO-s: 0.96±0.40; Emx1-f: 1.52±1.34; Cux2-f: 1.47±1.43; referred to as spikes

detected in **Figure 6A**). After preprocessing (selecting isolated spiking events with ≤5 spikes

within spike summation windows of 150 ms and 50 ms for GCaMP6s and GCaMP6f,

respectively, and excluding atypical spiking events - see **Methods**), 70-78% of detected spikes

were included in the response curve analysis (**Figure 6B**; referred to as spikes analyzed in

Figure 6A). For all mouse lines, >50% of analyzed spikes occurred in multi-spike events and

not as isolated single spikes (**Figure 6C**; see **Figure 6–figure supplement 1** for a comparison

of spiking characteristics between spontaneous and visually-evoked cells). The fraction of

single-spike events within spike summation windows was largest in Cux2-f (48%), followed by

tetO-s (37%), Emx1-s (32%), and Emx1-f (23%).

In multi-spike events, we assessed whether variability in dF/F peak could be attributed to the

mean inter-spike interval (ISI) within the spiking event (here, the maximum ISI corresponded to

the spike summation window: 150 ms and 50 ms for GCaMP6s and GCaMP6f, respectively).

Pooling multi-spike events from all cells, the mean ISI was negatively correlated with dF/F but

explained little of its variance (R^2 ranging from 0.03 to 0.21; **Figure 6–figure supplement 2**).

Breaking this down into individual cells, we found a small fraction of cells having spiking events

with significant correlation coefficients of dF/F peak and mean ISI (**Figure 6–figure**

supplement 3A-B). Within these cells, a negative correlation between mean ISI and dF/F peak

was observed, and for 4- and 5-spike events, R^2 ranged from ~0.4 to 0.9 (**Figure 6–figure**

supplement 3C), likely contributing to the overall negative correlation at the population level.

Comparing spike-to-calcium fluorescence response curves imaged at high and low

spatiotemporal resolutions

For a subset of cells, we performed calcium imaging at both high zoom (focusing on individual

cells; ~20 × 20 μm field of view at 0.2 μm per pixel and ~158 fps) and low zoom (population

imaging; ~400 x 400 μm field of view at 0.8 μm per pixel and ~30 fps, with 1 Emx1-f cell at 1.6

μm per pixel and ~60 fps) (**Table 1, Figure 7A**), enabling a direct comparison of their spike-to-

calcium fluorescence response curves (**Figure 7B**). For Emx1-f and Cux2-f, the slope of the

spike-to-calcium fluorescence response curve was significantly smaller when imaged at lower

spatiotemporal resolution, despite no detected difference in mean firing rates between groups

(**Figure 7C-D; Figure 7–figure supplement 1A**). A similar trend was observed in Emx1-s cells

(p = 0.05, paired t-test) between response curve slopes at high- and low-zoom. Across all 3

lines, imaging at low-zoom consistently resulted in higher variability of dF/F peak, both within

individual cells (**Figure 7B, Figure 7–figure supplement 1B-C**) and between cells (**Figure 7–**

figure supplement 1D). Furthermore, the mean single-spike response was different from the

mean imaging noise in fewer cells at low-zoom compared to that at high-zoom for all mouse

lines (**Figure 7–figure supplement 1E**), with higher noise floors and lower SNRs (**Figure 7–**

figure supplement 1F).

At the single-trial level, detection of single-spikes was 15±12% for Emx1-s (compared to

52±28% for the same cells imaged at high-zoom), 15±8% for Emx1-f (47±18% at high-zoom),

and 9±7% for Cux2-f (42±26% at high-zoom) at 5% false positive rate (**Figure 7–figure**

supplement 2). For 2-spike events, detection rate at low-zoom was 36±22% for Emx1-s

(compared to 92±16% for the same cells imaged at high-zoom), 16±11% for Emx1-f (86±23% at

high-zoom), and 19±12% for Cux-f (73±24% at high-zoom) at 5% false positive rate.

Discussion

Calcium imaging is widely used to report neuronal spiking activity *in vivo*. However, accurate

spike inference from calcium imaging remains a challenge, and there are relatively few ground

truth datasets with simultaneous calcium imaging and electrophysiology to aid the development

of more accurate spike inference algorithms. In a recent challenge (Spike Finder;

http://spikefinder.codeneuro.org/) (Berens et al., 2018), ~40 algorithms were trained and tested

on datasets consisting of 37 GCaMP6-expressing cells, underscoring the need for additional

GCaMP6 calibration data. In addition to supporting efforts toward spike inference, an improved

understanding of the relationship between spiking and observed fluorescence signals is

necessary to further broaden the utility and impact of calcium imaging. To these ends, we

contribute a ground truth dataset consisting of 91 V1 L2/3 excitatory neurons recorded at single-

cell resolution (available at https://brain-map.org/explore/circuits/oephys), and characterized

their spike-to-calcium fluorescence transfer function. Complementing existing datasets with viral

GECI expression (Chen et al., 2013; Theis et al., 2016; Dana et al., 2016), our work facilitates

interpretation of existing and future calcium imaging studies using mainstream transgenic

mouse lines, such as the Allen Institute's Brain Observatory Visual Coding dataset

(http://observatory.brain-map.org/visualcoding) (de Vries et al., 2019).

We used the same ROI extraction and neuropil correction methods as Chen et al. to compare

our results in transgenic mice to that of virally expressed GCaMP6 (Chen et al., 2013). To avoid

contamination of fluorescence responses by subsequent spiking events, we used spike

summation windows of 150 ms and 50 ms for GCaMP6s and GCaMP6f, respectively, instead of

250 ms as reported in Chen et al., a change that was necessitated by differences in firing rates

between our dataset and theirs. Choosing a constant neuropil contamination ratio of 0.7, dF/F

magnitudes of Emx1-f and Cux2-f were comparable to that of viral GCaMP6f (~100% for 5-spike

events in viral GCaMP6f), whereas Emx1-s and tetO-s were lower than that of viral GCaMP6s

(~200% for 5-spike events in viral GCaMP6s). To avoid potential confounding factors

associated with choosing a constant neuropil contamination ratio for all cells (e.g. increased

within-cell variability and artificially boosted dF/F magnitude), we chose not to perform neuropil

subtraction in subsequent analyses, except to quantify the relationship between neuropil

contamination ratio and the efficiency of detecting single-trial spiking events (discussed below).

Consistent with viral GCaMP6 (Chen et al., 2013), GCaMP6s cells in transgenic mice produced

a larger single-spike dF/F peak compared to that of GCaMP6f. Additionally, GCaMP6s cells

exhibited lower within-cell variability compared to GCaMP6f, both for single-spikes and multi-

spike events. Interestingly, tetO-s cells exhibited a larger single-spike dF/F peak and a larger

response curve slope compared to that of Emx1-s (albeit with a small sample size of n=6 cells).

However, tetO-s cells exhibited a higher noise floor (standard deviation of dF/F during no-spike

intervals) compared to that of Exm1-s, such that no difference in single-spike SNR was detected

between the two GCaMP6s lines. For multi-spike events, SNR for tetO-s was lower than that of

Emx1-s and Emx1-f, which could be attributed to a smaller sample size and fewer mutli-spike

events in the tetO-s cells with no-spike intervals (n=4 cells with no-spike intervals out of 6 total

cells). Nevertheless, with the large dF/F response shown here and previous work showing

comparable expression levels to Emx1-s and no observed epileptiform events (Steinmetz et al.,

2017), the tetO-s line represents an alternative to Emx1-s for expressing GCaMP6s.

Spike detection at the single-trial level was the most accurate in tetO-s, with mean single-spike

detection rate of 87±15% (mean±sd) at 5% false positive rate, or 85±16% at 1% false positive

rate. Thus, single spikes could be reliably detected in transgenic mouse lines expressing

GCaMP6s, and in tetO-s the detection rate was not far short of that reported for viral GCaMP6s

(99% at 1% false positive rate) (Chen et al., 2013). The difference in reported values could be

partially attributed to the different spike exclusion windows used: 150/200 ms (pre and post) in

this study and 1 s in Chen et al. Although spike detection rates were variable across cells, we

observed single-spike detection rates of ≥90% in a number of neurons (including 1 Emx1-f cell),

thus validating our experimental setup. Additionally, we found that spike detection rate, similar

to dF/F magnitude, was dependent on the neuropil contamination ratio r. We found optimal r-
values for maximizing single-spike detection rates in individual cells, by maximizing the area
under the ROC curve for classifying spiking events. The effect of optimizing r-values on spike
detection rate was most pronounced in Emx1-s, with an increase of 36±62% per cell compared
to r = 0 (at 5% false positive rate; the corresponding mean single-spike detection rate increased
from 63±25% to 76±20%). These data highlight the potential benefit of tuning r-values for
individual neurons, as was done for our Brain Observatory dataset (de Vries et al., 2019), over
the traditional method of selecting a constant r-value for the population. However, it is worth
noting that increased r-values also typically resulted in increased within-cell variability of dF/F
peak and increased dF/F magnitude, as discussed above.
345
2-p calcium imaging of large neuronal populations in parallel is typically performed on the order
of 30 fps, providing a small fraction (~20% in this study) of the number of temporal samples of
that imaged at high-zoom. Additionally, assuming an 8 × 8 μm square cell, the number of pixels
at low-zoom was 6.25% ($10^2/40^2$) of that at high-zoom. Coupled with differences in the number
of samples per pixel (13 or 18 at high-zoom and 3 at low-zoom), the number of spatial samples
at low-zoom was ~1.5% of that at high-zoom. Not surprisingly, we observed reduced response
curve slope, higher variability, and lower SNR when imaging at the lower spatiotemporal
resolution. Furthermore, single-spike detection rates (at 5% false positive rate) dropped from
~40-50% when imaged at high-zoom to ~10-15% at low-zoom in the same cells. At low-zoom,
detection of 2-spike events remained unreliable: <40% for Emx1-s and <20% for Emx1-f and
Cux2-f. Due to the nature of our experiments (e.g. acute surgical preparation, with brain motion
controlled through agarose, and the presence of a recording micropipette), our low-zoom data
may be underestimating the data quality for a typical population imaging session (a chronic,
more stable preparation with a coverslip cranial window, no micropipette and associated tissue
damage). Nevertheless, these data from the same cells offer a point of comparison between
data acquired under high-zoom and population imaging conditions.
362
In summary, in this study we present a ground truth dataset with simultaneous electrophysiology
and calcium imaging. This unique dataset allowed us to systematically compare multiple
aspects of spike-response properties between different GECIs (GCaMP6s and GCaMP6f) and
among several transgenic lines. For example, we found that GCaMP6s and GCaMP6f cells
have similar spike-fluorescence response curves, but GCaMP6f cells have greater variability
and lower single-spike detection probability than GCaMP6s cells. While both GCaMP6f and

GCaMP6s cells have reasonably good SNR and single-spike detectability at high-zoom conditions, both types of properties deteriorate substantially under the commonly used low-zoom population imaging conditions, with single-spike detection probabilities dropping to 4-20% (at 1-10% false positive rates). This finding has important implications for the interpretation of many large-scale in vivo calcium imaging datasets. By making our data freely available, we hope that it will serve the community as a further resource to better understand quantitatively the link between calcium-evoked fluorescent imaging signals and spiking activity.

Materials and Methods

Experimental procedures were in accordance with NIH guidelines and approved by the Institutional Animal Care and Use Committee (IACUC) of the Allen Institute for Brain Science. All experiments were conducted at the Allen Institute for Brain Science.

Mice. 2-p-targeted electrophysiology and 2-p calcium imaging was conducted in adult transgenic mice (2-6 months old, both sexes, n = 22), including *Emx1-IRES-Cre;Camk2a-tTA;Ai94* (simplified as Emx1-s, n = 7), *Camk2a-tTA;tetO-GCaMP6s* (Wekselblatt et al., 2016) (simplified as tetO-s, n = 2), *Emx1-IRES-Cre;Camk2a-tTA;Ai93* (simplified as Emx1-f, n = 6), and *CreERT2;Camk2a-tTA;Ai93* (simplified as Cux2-f, n = 7). Ai93 and Ai94 containing mice (Madisen et al., 2015) included in this dataset did not show behavioral signs for epileptic brain activity (Steinmetz et al., 2017).

Surgery. Mice were anesthetized with either isoflurane (0.75-1.5% in O_2) or urethane (1.5 g/kg, 30% aqueous solution, intraperitoneal injection), then implanted with a metal head-post. A circular craniotomy was performed with skull thinning over the left V1 centering on 1.3 mm anterior and 2.6 mm lateral to the Lambda. During surgery, the craniotomy was filled with Artificial Cerebrospinal Fluid (ACSF) containing (in mM): NaCl 126, KCl 2.5, NaH_2PO_4 1.25, $MgCl_2$ 1, $NaHCO_3$ 26, glucose 10, $CaCl_2$ 2, in ddH_2O; 290 mOsm; pH was adjusted to 7.3 with NaOH to keep the exposed V1 region from overheating or drying. Durotomy was performed to expose V1 regions of interest that were free of major blood vessels to facilitate the penetration of recording micropipettes. A thin layer of low melting-point agarose (1-1.3% in ACSF, Sigma-Aldrich) was then applied to the craniotomy to control brain motion. The mouse body temperature was maintained at 37°C with a feedback-controlled animal heating pad (Harvard Apparatus).

Calcium imaging. Individual GCaMP6+ neurons within ~100-300 μm underneath the pial surface were visualized under adequate depth of anesthesia (Stage III-3) using a Bruker (Prairie) 2-p microscope with 8 kHz resonant-galvo scanners, coupled with a Chameleon Ultra II Ti:sapphire laser system (Coherent). Fluorescence excited at 920 nm wavelength, with <70 mW laser power measured after the objective, was collected in two spectral channels using green (510/42 nm) and red (641/75 nm) emission filters (Semrock) to visualize GCaMP6 and the Alexa Fluor 594-containing micropipette, respectively. Fluorescence images were acquired at various framerates (117.6-158.3 fps) through a 16x water-immersion objective lens (Nikon, NA 0.8), with or without visual stimulations.

Electrophysiology. 2-p targeted cell-attached recording was performed following established protocols (Margrie et al., 2003; Kitamura et al., 2008; Knoblich et al., 2019). Long-shank borosilicate (KG-33, King Precision Glass) micropipettes (5-10 MΩ) were pulled with a P-97 puller (Sutter) and filled with ACSF and Alexa Fluor 594 to perform cell-attached recordings on GCaMP6+ neurons. Micropipettes were installed on a MultiClamp 700B headstage (Molecular Devices), which was mounted onto a Patchstar micromanipulator (Scientifica) with an approaching angle of 31 degrees from horizontal plane. Minimal seal resistance was 20 MΩ. Data were acquired under "I = 0" mode (zero current injection) with a Multiclamp 700B, recorded at 40 kHz using Multifunction I/O Devices (National Instruments) and custom software written in LabVIEW (National Instruments) and MATLAB (MathWorks). Isoflurane level was intentionally adjusted during recording sessions to keep the anesthesia depth as light as possible, resulting in fluctuation of the firing rates of recorded neurons.

Visual stimulation. Whole-screen sinusoidal static and drifting gratings were presented on a calibrated LCD monitor spanning 60° in elevation and 130° in azimuth to the contralateral eye. The mouse's eye was positioned ~22 cm away from the center of the monitor. For static gratings, the stimulus consisted of 4 orientations (45° increment), 4 spatial frequencies (0.02, 0.04, 0.08, and 0.16 cycles per degree), and 4 phases (0, 0.25, 0.5, 0.75) at 80% contrast in a random sequence with 10 repetitions. Each static grating was presented for 0.25 seconds, with no inter-stimulus interval. A gray screen at mean illuminance was presented randomly a total of 60 times. For drifting gratings, the stimulus consisted of 8 orientations (45° increment), 4 spatial frequency (0.02, 0.04, 0.08, and 0.16 cycle per degree) and 1 temporal frequency (2 Hz), at 80% contrast in a random sequence with up to 5 repetitions. Each drifting grating lasted for 2

seconds with an inter-stimulus interval of 2 seconds. A gray screen at mean illuminance was

presented randomly for up to 15 times.

Data analysis. Electrophysiology and calcium imaging data were analyzed using custom

MATLAB and Python scripts. For electrophysiology, Vm was filtered between 250 Hz and 5 kHz,

and automated spike detection was performed using a threshold criterion (5×std of Vm).

Unusually prolonged transient increases in calcium fluorescence were excluded from analysis

with an adaptive Vm threshold: $0.25 \times V_{baseline}+(V_{peak}-V_{baseline})$, where $V_{baseline}$ was the mean Vm

over 2 ms before the first spike of the spiking event and V_{peak} was the amplitude of the first spike

of the spiking event. The cumulative time above this Vm threshold was compared against a time

threshold (6 ms $\times$ the number of spikes within the spiking event), and the fluorescence

associated with spiking events with cumulative time above the time threshold were not

analyzed.

For calcium imaging, in-plane motion artifacts were corrected (Dombeck et al., 2007), and

cell/region of interest (ROI) selection was performed using a semi-automatic algorithm (Chen et

al., 2013) (kindly provided by Karel Svoboda, Janelia Research Campus). Ring-shaped ROIs

were used to select GCaMP6+ excitatory neurons, with GCaMP6 expression typically excluded

from the nucleus and restricted to the cytoplasm. For 2 of 91 cells, a satisfactory ring-shaped

ROI could not be found automatically, and a circular ROI covering the entire soma was used

instead. Calcium imaging data was smoothed using a local regression method using weighted

linear least squares and a 1st degree polynomial model (built-in "smooth" function in MATLAB

with "rlowess" method), and an averaging window of 5 frames.

To construct spike-calcium fluorescence response curves, we first identified all isolated spiking

events regardless of the number of spikes they contained. For GCaMP6s, spiking events were

separated from previous and subsequent spiking events by >150 ms and >50 ms, respectively,

due to the higher firing rates in our study. The 50 ms post-event exclusion window ensured the

absence of additional spikes in the calcium response window of 200 ms. For GCaMP6f, spiking

events were separated from previous and subsequent spiking events by >50 ms. Within each

spiking event, spike summation windows were 150 ms and 50 ms for GCaMP6s and GCaMP6f,

respectively, chosen based on the rise time of the GECIs (Chen et al., 2013). To determine

spike-triggered calcium fluorescence responses, fluorescence traces were aligned to the first

spike in each spiking event. The change in fluorescence, dF/F, for each spiking event was

calculated as $(F-F_0)/F_0$, where F_0 was computed locally as the mean fluorescence over 50 ms

and 20 ms before the first spike for GCaMP6s and GCaMP6f, respectively. For GCaMP6s, peak

dF/F was found within 200 ms after the first spike. For GCaMP6f, a variable calcium response

window was used: peak dF/F was found within 50 ms and 75 ms after the first spike for single-

spike and multi-spike events, respectively. Bursts of >5 spikes were excluded from analysis due

to the low frequency of such events. Alignment jitter intrinsic to the imaging frame rate was ≤6.3

ms in 86 of 91 cells (imaged at 158.7 fps), with mean expected error of 3.2 ms, and between

≤5.6 ms to ≤8.5 ms in others (imaged at 117.6-178.6 fps), with mean expected error of 2.8 to

4.3 ms.

To compare mean single-spike dF/F peak responses to imaging noise, imaging noise was

estimated as follows: we found no-spike intervals of ≥1 s, separated by ≥4 s and ≥1 s from

previous and subsequent spikes, respectively, and randomly sampled the m corresponding

calcium fluorescence traces n times, where $m×n$ approximately matched the number of single-

spike events. Noise (zero-spike) dF/F peak was computed using the same calcium response

windows as single-spike events (50 ms and 200 ms for GCaMP6f and GCaMp6s, respectively).

To quantify the efficiency of detecting spiking events at a single-trial level, we compared dF/F

traces of the response (single-spike events or 2-spike events) to that of imaging noise. To

estimate imaging noise, calcium fluorescence traces during no-spike intervals were randomly

sampled as described above, for a minimum of 25 times. For each cell, the mean response

trace was used as the template vector. The template vector was normalized after subtracting

the mean to create the unit vector, and the scalar results of projecting the response and noise

traces on the unit vector were computed: r_i and n_i for response and noise scalars, respectively.

The detection threshold was defined as the x^{th} percentile of n_i values, where 1-x represented the

false positive rate (e.g., x = 95 for 95^{th} percentile or 5% false positive rate), and the detection

rate (true positive rate) was the fraction of r_i values above the detection threshold.

Acknowledgements

We are grateful for the Animal Care, Transgenic Colony Management, and Lab Animal Services

teams for mouse husbandry, and Carol Thompson and John Phillips for providing project

management support. We thank Karel Svoboda, Hod Dana and Tsai-Wen Chen for sharing

analysis software. This work was also supported by grants from National Natural Science

Foundation of China (NSFC31871055) and Guangdong Science and Technology Department

(2017B030314026) to L.L. We thank the Allen Institute founders, Paul G. Allen and Jody Allen,

for their vision, encouragement, and support.

Author contributions

509 CK, HZ, RCR conceived the project. LL, UK, LH performed experiments. LL, LH, UK, PL, JW,

510 JL, SEJdV, MAB, GJM analyzed data. LL, HZ, CK, RCR, MAB, GJM supervised the project. LH

511 wrote the paper with input from all co-authors.

Competing interests

514 The authors declare no competing financial interests.

Data availability

517 The dataset is available at https://brain-map.org/explore/circuits/oephys.

References

Akerboom, J., Chen, T.-W., Wardill, T.J., Tian, L., Marvin, J.S., Mutlu, S., Calderón, N.C.,
Esposti, F., Borghuis, B.G., Sun, X.R., et al. (2012). Optimization of a GCaMP calcium indicator
for neural activity imaging. J. Neurosci. Off. J. Soc. Neurosci. *32*, 13819–13840.

Berens, P., Freeman, J., Deneux, T., Chenkov, N., McColgan, T., Speiser, A., Macke, J.H.,
Turaga, S.C., Mineault, P., Rupprecht, P., et al. (2018). Community-based benchmarking
improves spike rate inference from two-photon calcium imaging data. PLoS Comput. Biol. *14*,
e1006157.

Chen, T.-W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R.,
Kerr, R.A., Orger, M.B., Jayaraman, V., et al. (2013). Ultrasensitive fluorescent proteins for
imaging neuronal activity. Nature *499*, 295–300.

Daigle, T.L., Madisen, L., Hage, T.A., Valley, M.T., Knoblich, U., Larsen, R.S., Takeno, M.M.,
Huang, L., Gu, H., Larsen, R., et al. (2018). A Suite of Transgenic Driver and Reporter Mouse
Lines with Enhanced Brain-Cell-Type Targeting and Functionality. Cell *174*, 465-480.e22.

Dana, H., Mohar, B., Sun, Y., Narayan, S., Gordus, A., Hasseman, J.P., Tsegaye, G., Holt,
G.T., Hu, A., Walpita, D., et al. (2016). Sensitive red protein calcium indicators for imaging
neural activity. ELife *5*.

Dombeck, D.A., Khabbaz, A.N., Collman, F., Adelman, T.L., and Tank, D.W. (2007). Imaging
large-scale neural activity with cellular resolution in awake, mobile mice. Neuron *56*, 43–57.

Kitamura, K., Judkewitz, B., Kano, M., Denk, W., and Häusser, M. (2008). Targeted patch-clamp
recordings and single-cell electroporation of unlabeled neurons in vivo. Nat. Methods *5*, 61–67.

Knoblich, U., Huang, L., Zeng, H., and Li, L. (2019). Neuronal cell-subtype specificity of neural
synchronization in mouse primary visual cortex. Nat. Commun. *10*, 2533.

Luo, L., Callaway, E.M., and Svoboda, K. (2018). Genetic Dissection of Neural Circuits: A
Decade of Progress. Neuron *98*, 256–281.

Madisen, L., Garner, A.R., Shimaoka, D., Chuong, A.S., Klapoetke, N.C., Li, L., van der Bourg,
A., Niino, Y., Egolf, L., Monetti, C., et al. (2015). Transgenic mice for intersectional targeting of
neural sensors and effectors with high specificity and performance. Neuron *85*, 942–958.

Margrie, T.W., Meyer, A.H., Caputi, A., Monyer, H., Hasan, M.T., Schaefer, A.T., Denk, W., and
Brecht, M. (2003). Targeted whole-cell recordings in the mammalian brain in vivo. Neuron *39*,
911–918.

Nadim, F., and Bucher, D. (2014). Neuromodulation of neurons and synapses. Curr. Opin.
Neurobiol. *29*, 48–56.

Rose, T., Goltstein, P.M., Portugues, R., and Griesbeck, O. (2014). Putting a finishing touch on
GECIs. Front. Mol. Neurosci. *7*, 88.

Steinmetz, N.A., Buetfering, C., Lecoq, J., Lee, C.R., Peters, A.J., Jacobs, E.A.K., Coen, P.,
Ollerenshaw, D.R., Valley, M.T., de Vries, S.E.J., et al. (2017). Aberrant Cortical Activity in
Multiple GCaMP6-Expressing Transgenic Mouse Lines. ENeuro *4*.

Theis, L., Berens, P., Froudarakis, E., Reimer, J., Román Rosón, M., Baden, T., Euler, T.,
Tolias, A.S., and Bethge, M. (2016). Benchmarking Spike Rate Inference in Population Calcium
Imaging. Neuron *90*, 471–482.

de Vries, S.E.J., Lecoq, J., Buice, M.A., Groblewski, P.A., Ocker, G.K., Oliver, M., Feng, D.,
Cain, N., Ledochowitsch, P., Millman, D., et al. (2019). A large-scale, standardized physiological
survey reveals higher order coding throughout the mouse visual cortex. Nat. Neurosci. *In press*.

Wekselblatt, J.B., Flister, E.D., Piscopo, D.M., and Niell, C.M. (2016). Large-scale imaging of
cortical dynamics during sensory perception and behavior. J. Neurophysiol. *115*, 2852–2866.

567
568
Figure 1. Simultaneous calcium imaging and electrophysiology *in vivo*. (**A**) Experimental design and
example fluorescence and Vm traces from an example Emx1-s neuron. (**B**) Identification of spiking
events. Spike-to-calcium fluorescence analysis was performed for spiking events that were separated
from previous and next spikes by durations of the exclusion windows (red shading) or longer. The
summation window (gray shading) determined the number of spikes within each spiking event. (**C**) Spike-
triggered fluorescence responses. Red lines indicate spikes and numbers indicate the number of spikes
in each spiking event. (**D**) Fluorescence changes (dF/F) in response to representative single-spike events
from the recorded cell shown in (A). For GCaMP6s, the spike summation window (left panel) and calcium
response window (right panel) were 150 ms and 200 ms, respectively. Colors indicate correspondence
between spikes and dF/F. Black line indicates mean dF/F. Circles indicate peak fluorescence changes
(dF/F peak) within the calcium response window.

Figure 2. Example single-cell spike-to-calcium fluorescence response curves. For each type of
mice: (i) Cell ROI and segmentation mask. (ii) Corresponding fluorescence (top) and Vm traces (bottom).
Red lines indicate detected spikes. (iii) dF/F traces (left) and spike-to-calcium fluorescence response
curve (dF/F peak as a function of the number of spikes, with a minimum requirement of 3 events per bin;
right). Horizontal black lines indicate the standard deviation of dF/F during no-spike intervals (i.e.
estimated noise floor; see **Figure 4–figure supplement 2**). Estimated zero-spike dF/F peak were plotted
for comparison (see **Figure 4-figure supplement 1**). Spike summation windows were 150 ms and 50 ms
for GCaMP6s and GCaMP6f, respectively.

590
591
Figure 2–figure supplement 1. Effect of neuropil subtraction on within-cell variability of dF/F peak.
(**A**) An example Emx1-s neuron, with labeled regions where cell and neuropil fluorescence were
measured, and the corresponding mean fluorescence traces. (**B**) Comparison of cell fluorescence without
and with neuropil correction, where the contamination ratio r = 0.5 was chosen. Fluorescence changes
related to spikes (red lines) were also subtracted. (**C**) Comparison of single-cell response curve without
and with neuropil correction (r = 0.5). Red lines indicate mean ± sem. (**D**) Mean coefficient of variation of
dF/F peaks as a function of r, normalized to no neuropil subtraction (r = 0), for all mouse lines. Circles
indicate individual cells. Black crosses represent the Emx1-s cell shown in (A).

Figure 3. Population response. (**A**) Population spike-to-calcium fluorescence response curves. Each line represents one cell. Bold lines indicate mean responses for spontaneous, visually-evoked, and all recordings. Horizontal black lines indicate the mean standard deviation of dF/F during no-spike intervals (i.e. mean noise floor; see **Figure 4–figure supplement 2**). (**B**) Overlay of population response curves. By cell (top): population means from (A), computed from the mean response of individual cells. Emx1-s: n = 32 cells; tetO-s: n = 6; Emx1-f: n = 26; Cux2-f: n = 27. By spiking events (bottom): population response computed from spiking events pooled from all cells as in Chen et al., 2013. Emx1-s: n = 1160, 599, 255, 103, and 52 spiking events for 1-5 spikes; tetO-s: n = 250, 103, 48, 17, and 6; Emx1-f: n = 1204, 921, 416, 157, and 60; Cux2-f: n = 2667, 809, 209, 118, and 46. (**C**) Mean and median dF/F in response to single-spike events. For GCaMP6f, the single-spike calcium response window was 50 ms. Shading corresponds to 2×sem. (**D**) Mean single-spike dF/F peak of each cell. (**E**) Response curve slope of each cell, computed from linear regression of dF/F peaks against the numbers of spikes.

615
616
617 **Figure 3–figure supplement 1. Population response for all genotype and visual stimulation**
618 **groups**. Population spike-to-calcium fluorescence response curves for spontaneous or visually-evoked
619 responses. Bold lines indicate mean responses. No visually-evoked responses were recorded from tetO-s
620 mice. Spike summation windows were 150 ms and 50 ms for GCaMP6s and GCaMP6f, respectively.

621
622
Figure 3–figure supplement 2. Spike-to-calcium fluorescence response curve and mean firing rate
under isoflurane or urethane anesthesia. (**A**) Single-cell spike-to-calcium fluorescence response
curves acquired under isoflurane or urethane anesthesia (left; bold lines indicate mean population
response) and overlay of population response (right). (**B**) Mean firing rate from analyzed spikes acquired
under isoflurane or urethane anesthesia (See **Results** for description of spiking characteristics). Emx1-s:
n = 9 cells from 2 mice using urethane (0 spontaneous and 9 visually-evoked recordings); n = 23 cells
from 5 mice using isoflurane (21 spontaneous and 2 visually-evoked recordings). Cux2-f: n = 5 cells from
2 mice using urethane (3 spontaneous and 2 visually-evoked recordings); n = 22 cells from 5 mice using
isoflurane (1 spontaneous and 21 visually-evoked recordings). Spike summation windows were 150 ms
and 50 ms for GCaMP6s and GCaMP6f, respectively.

633
634
Figure 3–figure supplement 3. Population spike-to-calcium response curves with neuropil
correction. Population spike-to-calcium fluorescence response curves computed from all spiking events
(no exclusion of atypical spiking events) using a neuropil contamination ratio of 0.7. By cell: population
response was computed from the mean response of individual cells (left). Emx1-s: n = 32 cells; tetO-s: n
= 6; Emx1-f: n = 25; Cux2-f: n = 26. By spiking events: population response was computed from spiking
events pooled from all cells as in Chen et al., 2013 (right). Emx1-s: n = 1264, 668, 298, 118, and 57
spiking events for 1-5 spikes; tetO-s: n = 254, 103, 48, 17, and 6; Emx1-f: n = 1300, 1010, 476, 183, and
70; Cux2-f: n = 2673, 788, 207, 118, and 46. For both analyses, only cells with ring-shaped ROI were
included (see **Methods**). Spike summation windows were 150 ms and 50 ms for GCaMP6s and
GCaMP6f, respectively.

645
646
Figure 4. Within-cell and between-cell variability of dF/F peak. (**A**) Within-cell variability. Mean
coefficient of variation of dF/F peaks of each cell computed from all spiking events (left panel; p = 2e-8,
ANOVA) or single-spike events (right panel; p = 3e-7, ANOVA). (**B**) Within-cell variability (coefficient of
variation of dF/F peak) for each spike bin (Emx1-s: p = 0.04; tetO-s: p = 0.03; Emx1-f: p = 3e-11; Cux2-f:
p = 6e-6, ANOVA). ns, p > 0.05; *, p < 0.05; **, p < 0.01; ***, p < 0.001, multiple comparison test using
Tukey's honest significant difference criterion. (**C**) Cell-to-cell variability. Coefficient of variation of dF/F
peak from the population response in Fig. 3B (top). Spike summation windows were 150 ms and 50 ms
for GCaMP6s and GCaMP6f, respectively.

655
656
Figure 4–figure supplement 1. Comparing mean single-spike response and imaging noise. (**A**) No-
spike intervals of ≥1 s, separated by ≥4 s and ≥1 s from previous and subsequent spikes, respectively,
and zoomed-in view of their corresponding calcium fluorescence traces in an example Emx1-s cell. Red
lines indicate detected spikes. Boxed regions indicate no-spike intervals. Zoomed-in views are shown for
numbered boxed regions. (**B**) Fluorescence traces randomly sampled from all no-spike intervals in (A).
(**C**) Kernel density estimates of dF/F peak sampled from no-spike intervals (zero-spike estimate) and in
response to single-spikes. (**D**) Mean zero-spike and single-spike dF/F peak. Circles indicate individual
cells. (**E**) Fraction of cells in which mean single-spike response was different from mean imaging noise (p
< 0.05, t-test).

666
667
Figure 4–figure supplement 2. Noise floor and signal-to-noise ratio. (**A**) Noise floor (standard
deviation of dF/F during no-spike intervals; left; $p = 2e-6$, ANOVA) and single-spike signal-to-noise ratio
(SNR; single-spike dF/F peak normalized by standard deviation of dF/F during no-spike intervals; right; p
$= 2e-5$, ANOVA). *, $p < 0.05$; **, $p < 0.01$, ***, $p < 0.001$, multiple comparison test using Tukey's honest
significant difference criterion. (**B**) SNR as a function of the number of spikes (left panels), and overlay of
mean SNR for the 4 mouse lines (right panel). Spike summation windows were 150 ms and 50 ms for
GCaMP6s and GCaMP6f, respectively.

675
676
677 **Figure 5. Spike detection at the single-trial level.** (i) Receiver operating characteristic (ROC) curves for
678 classifying 1-spike (top) and 2-spike (bottom) events in individual cells. (ii) Mean population response. (iii)
679 Detection rate (true positive rate) at 1%, 5%, and 10% false positive rate. Emx1-s: n = 30 cells with at
680 least 1 no-spike interval out of 32 total cells; tetO-s: n = 4 out of 6; Emx1-f: n = 23 out of 26; Cux2-f: n =
681 23 out of 27.

682
683
Figure 5–figure supplement 1. Effect of neuropil subtraction on single-spike detection rate. (A)
Identifying the optimal neuropil contamination ratio r for maximizing spike detection efficiency. (Top) ROC
curves for classifying single-spike events in an example Emx1-s cell, where the neuropil contamination
ratio r was varied from 0 to 1. (Bottom) Area under ROC curves as a function of r. (B) (Left) Distribution of
optimal r-values in individual cells for maximizing area under ROC curve. (Right) Single-spike detection at
5% false positive rate before (r = 0) and after optimization. (Emx1-s: n = 30 cells with at least 1 no-spike
interval; tetO-s: n = 4; Emx1-f: n = 22; Cux2-f: n = 22. One Emx1-f and 1 Cux2-f cell were not analyzed
because a satisfactory ring-shaped ROI could not be found automatically as described in **Methods**.)

Figure 6. Spiking characteristics. (**A**) Mean firing rate computed from every detected spike or analyzed
spikes within the same time window (i.e. isolated spiking events with atypical spiking events excluded).
Each circle (pair) indicates one cell. (**B**) Fraction of spikes analyzed. Each circle indicates one cell. (**C**) (i)
Fraction of spiking events in each spike bin; (ii) Fraction of number of spikes in each spike bin; and (iii)
Single-spike proportion by cell. Spike summation windows were 150 ms and 50 ms for GCaMP6s and
GCaMP6f, respectively.

Figure 6–figure supplement 1. Spiking characteristics for spontaneous and visually-evoked cells.
(**A**) Mean firing rate from every detected spike or analyzed spikes (i.e. isolated spiking events with
atypical spiking events excluded). Each circle (pair) indicates 1 cell. (**B**) Fraction of spikes analyzed. Each
circle indicates 1 cell. (**C**) (i) Fraction of spiking events in each spike bin; (ii) Fraction of number of spikes
in each spike bin; and (iii) Single-spike proportion by cell. Spon and visstim indicate spontaneous and
visually-evoked responses, respectively. Spike summation windows were 150 ms and 50 ms for
GCaMP6s and GCaMP6f, respectively.

709
710
711 **Figure 6–figure supplement 2. Relationship between mean inter-spike interval and dF/F peak at the**
712 **population level.** Relationship between mean inter-spike interval (ISI) and dF/F peak (normalized to the
713 mean single-spike response) for multi-spike events pooled from all cells. Circles and lines indicate spiking
714 events and linear regression fit. Spike summation windows were 150 ms and 50 ms for GCaMP6s and
715 GCaMP6f, respectively.

716
717
Figure 6–figure supplement 3. Relationship between mean ISI and dF/F peak in individual cells. (**A,**
B) Between-cell variability. (**A**) Fraction of cells with significant correlation coefficients of dF/F peak and
mean ISI ($p < 0.05$). Denominators indicate the number of cells with the corresponding multi-spike event.
Overall, significant correlations were observed in 14 out of 86 (16%) multi-spike events for Emx1-s, 5 out
of 14 (36%) multi-spike events for tetO-s, 16 out of 74 (22%) multi-spike events for Emx1-f, and 12 out of
57 (21%) multi-spike events for Cux2-f. (**B**) R^2 of dF/F peak and mean ISI for cells with significant
correlation coefficients. (**C**) Within-cell relationship between mean ISI and dF/F peak for cells with
significant correlation coefficients. Circles and lines indicate spiking events and linear regression fit;
colors indicate individual cells and do not correspond across figure panels. Spike summation windows
were 150 ms and 50 ms for GCaMP6s and GCaMP6f, respectively.

Figure 7. Spike-to-calcium fluorescence response curves imaged at high and low spatiotemporal
resolutions. (**A**) Fields of view and cell ROIs for an example Cux2-f neuron imaged at both high (white
ROI, inset) and low (green ROI) optical zooms. Scale bars, high-zoom: 10 μm; low-zoom: 100 μm. (**B**)
Corresponding dF/F traces and spike-to-calcium fluorescence response curves for the neuron shown in
(A). Horizontal black lines indicate the standard deviation of dF/F during no-spike intervals (i.e. estimated
noise floor; see **Figure 4–figure supplement 2**). (**C**) Population response of cells imaged at both high-
and low-zooms, expressed as the ratio of high-zoom to low-zoom dF/F peak, without (left) and with
normalization (right) by the mean single-spike response. Colored bold lines indicate the mean response.
One Emx1-f cell with near zero (<0.01) dF/F peak for 2-spikes at low-zoom was excluded from this plot.
(**D**) Response curve slope imaged at high-zoom and low-zoom, computed from linear regression of dF/F
peak against the number of spikes. Blue or red indicates response curve slope was larger when imaged
at high-zoom or low-zoom, respectively. Diagonal line indicates equal response curve slope when imaged
at high-zoom and low-zoom. (***, p < 0.001; **, p < 0.01, paired t-test.) Spike summation windows were
150 ms and 50 ms for GCaMP6s and GCaMP6f, respectively.
743

Figure 7–figure supplement 1. Fluorescence response variability imaged at high and low
spatiotemporal resolutions. (**A**) Mean firing rate from analyzed spikes. (**B**) Mean coefficient of variation
of dF/F peak. Each circle (pair) indicates 1 cell. **, $p < 0.01$; ***, $p < 0.001$, paired t-test. (**C**) Kernel
density estimates of dF/F peak sampled from no-spike intervals (zero-spike estimate) and in response to
single-spikes. (**D**) Mean zero-spike and single-spike dF/F peak. Circles indicate individual cells. (**E**)
Fraction of cells in which mean single-spike response was different from mean imaging noise ($p < 0.05$, t-
test). (**F**) Noise floor (standard deviation of dF/F during no-spike intervals; left) and single-spike SNR
(single-spike dF/F peak normalized by standard deviation of dF/F during no-spike intervals; right).

753
754
755 **Figure 7–figure supplement 2. Spike detection at low spatiotemporal resolution.** (i) ROC curves for
756 classifying 1-spike (top) and 2-spike (bottom) events in individual cells. (ii) Mean population response. (iii)
757 Detection rate (true positive rate) at 1%, 5%, and 10% false positive rate.