Varietal differences in physiological and biochemical responses to salinity stress in six finger millet plants

Asunta Mukami[†1], Alex Ngetich[2], Easter Syombua[3], Richard Oduor[4], Wilton Mbinda[†5*]

[1]Department of Life Sciences, South Eastern Kenya University, Kitui, Kenya

[2]Institute of Biotechnology Research, Jomo Kenyatta University of Agriculture Technology, Nairobi, Kenya

[3]Centre for Biotechnology and Bioinformatics, University of Nairobi, Kenya.

[4]Department of Biochemistry, Microbiology and Biotechnology, Kenyatta University, Nairobi, Kenya

[5]Department of Biochemistry and Biotechnology, Pwani University, Kilifi, Kenya
†contributed equally

[*]Corresponding author: Wilton Mbinda; Email: wilton.mbinda@gmail.com

Abstract

Finger millet is one of the most important cereals that are often grown in semiarid and arid regions of East Africa. However, salinity is known to be a major impediment for the crop growth and production. The goal of this study was to understand the mechanisms of physiological and biochemical responses to salinity stress of six Kenyan finger millet varieties (GBK043137, GBK043128, GBK043124, GBK043122, GBK043094, and GBK043050) grown across different agroecological zones under NaCl-induced salinity stress. Seeds were germinated on the sterile soil and treated using various concentrations of NaCl (100, 200 and 300 mM) for two weeks. Again, the early seedling stage of germinated plants was irrigated with the same salt concentrations for 60 days. Results indicated depression in germination percentage, shoot and root growth rate, leaf relative water content, total chlorophyll content contents, leaf K ion concentration, and leaf K/Na ratios in all varieties studied with increased salt levels. On the contrary, it was observed that proline and malonaldehyde (MDA) contents reduced sugar content and leaf total proteins. At the same time, the leaf Na ion and Cl ion amounts of all plants increased substantially with rising stress levels. Clustering analysis revealed that GBK043094 and GBK043137 were placed together and identified as salt tolerant varieties based on their performance under salt stress. Overall, our findings indicated a significant varietal variability for most of the parameters analysed. These superior varieties identified could be potentially used as promising genetic resources in future breeding programmes development directed towards salt-tolerant finger millet hybrids. However, further analysis at genomic and molecular level need to be undertaken in order to better understand specific mechanisms of the factors that promote salinity tolerance in finger millet.

Introduction

Salinity is the severest environmental stress that adversely affects the growth of plants and their productivity worldwide (Qadir et al. 2014). This abiotic stress mostly characterizes arid and semiarid regions that experience low rainfall and scarcity of good quality water. Salts accumulation in irrigation without proper drainage water system, coupled with underlying rocks rich with high salts contents, leads to gradual salinization of arable land, thereby affecting soil characteristics. The problem is expected to aggravate further owing to effects of rising sea levels and climate change (Tedeschi et al., 2011). It is estimated that if existing salinity stress phenomenon will continue to persist, more than 50% of the current cultivated agricultural land could be lost by the year 2050 (Wang et al., 2003). As of 2013, the global losses in agricultural production due to lands afflicted by salinity had touched US$12 billion and have been steadily rising ever since (Shabala, 2013).

Over time, plants have evolved complex salt tolerance adaptive mechanisms to counteract the harmful effects of salinity through activation of morphological, physiological biochemical, cellular and molecular responses which include changes to metabolic systems, nutritional disproportion, variation and disorder of membranes and reduction in rate of cell division and growth (Munns et al., 2006; Zhu, 2003). Another critical repercussion of salinity stress to plants is the overproduction of reactive oxygen species (ROS) from the pathways such as photosynthesis, mitochondrial respiration and photorespiration. ROS toxicity comes from their reactions with various cell units, which precipitates an avalanche of oxidative reactions and results to enzymes inactivation, protein degradation, lipid peroxidation and DNA damage (reference). Collectively, these effects inhibit growth and development and subsequently reduce crop yields. The most effective way to combat against salinity is the development of the resistant and tolerant crop varieties. It is therefore paramount to identify the genetic resources with high tolerance, and to understand the mechanisms of salinity tolerance in crops.

The resistance and response of plants to a salt stress varies according to its species or varieties or genotypes or variety and environment which could be attributed to the

biological dissimilarities between the species or varieties or genotypes, plant growth,

and composition and concentration of the salt stress conditions (Bertazzini et al.,

2018; Filichkin et al., 2018; Shabala et al., 2013). Many reports have shown that short

term salinity stress significantly affects the germination rate, seedling and root growth

as well as ion composition, levels of relative water content, photosynthetic pigments,

proline content, level of membrane lipid peroxidation as well as the amounts of

reducing sugars and total protein (Dugasa et al., 2019; Sarabi et al., 2017; Kumar and

Khare, 2016). These physiological and biochemical indices multivariate cluster

analysis been used for classification of salt-sensitive and salt-tolerant varieties so that

they can further be used in plant breeding programmes. The prevalent approach to

assess the performance of plants against salinity under laboratory conditions is

through assessing their physiological and biochemical responses on the application of

different concentrations NaCl.

Finger millet, *Eleusine coracana* L. (Geartn), is the 4[th] most important member of

millets after sorghum, pearl millet and foxtail millet, making it one of the most

valuable food cereals cultivated in arid and semi-arid regions of Asia and Africa

(Chivenge et al., 2015). The crop is well adapted to heat, drought and poor soil stress

in marginal and degraded soils. Further, its cereals have comparatively better

antioxidant and nutraceutical properties and superb storage qualities which lack in

other cereals (Kumar et al., 2016). All these attributes make finger millet one of the

important and promising plant genetics resources for agriculture, and food and

nutritional security and alleviation of poverty of poor farmers who live in arid,

infertile and marginal lands. Despite its importance, finger millet potential yields are

adversely affected salinity stress. More specifically, during seed germination and

seedling establishment terminal growth phases are extremely susceptible to salinity

stress (Ibrahim, 2016; Hema et al., 2014; Zhang et al., 2014). Hence, it is imperative

to screen for varieties with intrinsic salinity tolerance for yield improvement breeding

programmes. Salinity tolerance during germination and seedling development is

crucial for the establishment of plants growing in saline soils of arid and semi-arid

regions (Tlig et al., 2008). Accordingly, understanding the physiological and

biochemical salinity responses in finger millet is, therefore, of importance in breeding

salt resistant and tolerant crops. Owing to the wide disparity in agroecological regions across finger millet growing regions, several finger millet landraces exhibit an adaptation to a large range of environmental conditions and subsequently, represent valuable source of useful genetic source that can be exploited to improve salinity tolerance of finger millet varieties belong to distinct geographical zones in Kenya. We therefore investigated the physiological and biochemical responses to salinity stress of six finger millet varieties under NaCl induced salinity stress.

Materials and methods

Plant material, treatments and germination assays

Six Kenyan farmers preferred finger millet varieties (GBK043124, GBK043122, GBK043137, GBK043128, GBK043094 and GBK043050) obtained from Kenya Agricultural and Livestock Research Organization, Gene Bank, Muguga, Kenya were used in this study. Prior to assays, the seeds were sorted by handpicking of the healthy ones, then washed with distilled water to remove dust and other particles. Germination assay was performed using 10 seeds of each variety and at different concentrations of NaCl (100,200 and 300 mM). Seeds were planted in germination trays in round pots containing sterile soil to a depth of approximately 1 cm. The control seeds were irrigated with distilled water. Salinity stress was imposed on treatment groups by irrigating the seeds with various concentrations of NaCl at an interval of 3 days for two weeks. Observations on the rate of germination were scored on the 17[th] day of treatment.

Growth conditions under salinity treatment

Germinated finger millet seedlings were grown for 2 weeks under greenhouse conditions of 25±2 °C and 60-70% humidity, with a 16/8-h photoperiod provided by natural sunlight. To assay salinity stress effects on growth of finger millet, the seedlings were subjected to stress by irrigating with NaCl (100, 200 and 300 mM) for 21 days at an interval of 3 days. Control plants were watered with distilled water. In each experiment, five replications were used for each set of treatment. After treatment, five plants from each treatment were sampled at random and the growth of the plants studied by recording the shoot length and root length.

Relative water content

One leaflet from the first fully expanded leaf of five plants per variety and per treatment was cut from a plant on the 21[st] day. Immediately after cutting, the leaflet was weighed to obtain the fresh weight (FW). Thereafter, the leaflet was immersed in deionized water under normal room temperature for 4 hours. Afterwards, the leaflet was taken out, thoroughly wiped to remove the water on the blade surface and its weight measured to obtain turgid weight (TW). the leaflet was afterwards dried in an oven for 24 hours and its dry weight (DW) measured. The relative water content (RWC %) was calculated using the formula: RWC = [(FW -DW)/ (TW - DW)] x 100.

Determination of chlorophyll content

Chlorophyll a, b and total chlorophylls (a + b) were determined according to Arnon (1949). 0.2g of fresh leaves were taken from 21 days-old NaCl (0-300mM) treated plants, finely ground by vortexing several times to remove chlorophyll efficiently. The extract was centrifuged at 5000 g for 3 minutes. The absorbance of the obtained supernatants was measured at 645 and 663 nm using 1240 UV-Vis Spectrophotometer (Shimadzu, Kyoto, Japan). The total chlorophyll content in each sample, expressed in mg/g fresh mass (FM) was calculated using Arnon's 1949 formula: TC=20.2(A645) $\Box$ 8.02(A663) ×V/1000×W where V corresponds to the volume of total extract per litre and W is the mass of the fresh material.

Proline content measurement

Proline accumulation was determined as described by Bates et al. (1973). Fifty milligrams of fresh leaf tissues from each variety and treatment was homogenized in 10 ml of 3% w/v sulphosalicylic acid and the homogenate was filtrated. The resulting solution was mixed solution of acidic ninhydrin [40% (w/v) acidic ninhydrin (8.8 µM ninhydrin, 10.5 M glacial acetic acid, 2.4 M orthophosphoric acid), 40% (v/v) glacial acetic acid and 20% (v/v) of 3%(v/v) sulphosalicylic acid]. Thereafter, the reaction mixtures were put in a water bath at 100 °C for 60 minutes to develop colors and the reaction was terminated by incubating the mixtures in ice for 5 minutes. Toluene was added to separate chromophores. The optical density was measured at 520 nm using

177 1240 UV-Vis Spectrophotometer. Proline content [µmol/g fresh weight (F. WT)] in

178 leaf tissues was calculated from a standard curve made using 0-100 µg L-proline.

Lipid peroxidation assay

Fresh upper second fully expended leaves (0.3 g) harvested and homogenized in 0.1

% (w/v) trichloroacetic acid and then the homogenates were centrifuged at 10,000 g

for 15 minutes at 4 °C. The supernatant was mixed with 0.5 ml of 1.5 ml 0.5%

thiobarbituric acid diluted in 20% trichloroacetic acid and the mixture was incubated

in water bath at 95 °C for 25 minutes before incubating it on ice for 10 minutes. The

absorbance was measured at 532 and 600 nm using UVmini-1240 UV-Vis

Spectrophotometer with 1% thiobarbituric acid in 20% trichloroacetic acid as control.

The amount of malondialdehyde (µmol/g FW) calculated as a measure of lipid

peroxidation, was determined according to Heath and Packer, (1968).

Estimation of reducing sugar

The amount of reducing sugar in shoots was determined using method describe by

Johnson et al (1964). The sugar was extracted from 1.0 g homogenized tissue using

80% ethanol at 95 °C, then centrifuged for 10 min at 14000 rpm. The resulting

supernatant was dried for 2 hrs at 80 °C, before dissolving the residue in 10 ml of

distilled water and 2.0 ml alkaline copper reagent was added. The mixture was heated

in water bath at 100 °C for 10 min, and then cooled to room temperature. Exactly 1.0

ml of Nelson's reagent was added and the volume was adjusted to 10 ml with double

distilled water. Absorbance of the solution was taken at 520 nm. The amount of

reducing sugar (mg /g FW) was calculated using a standard curve of glucose.

Estimation of leaf total protein

Total sample protein was extracted using the acetone-trichloroacetic acid (TCA)

precipitation method as described by Damerval et al. (1986). In brief, 500 g of leaf

tissue from each treatment was homogenized in 10% TCA in ice and incubated

overnight at 4°C. The homogenate was centrifuged at 14,000 rpm for 15 min at 4°C

and the pellet was washed with100% acetone to remove any contaminating pigments.

To remove phenolic compounds, the pigment-free pellet was first washed with 80%

ethanol, ethanol/trichloromethane (3:1 v/v), then ethanol/ethoxyethane (3:1 v/v) and

finally with ethoxyethane. The washed pellet was then suspended in a volume of 0.1

N sodium hydroxide for protein estimation. The sample proteins were estimated at

750 nm using bovine serum albumin as standard and expressed as gram per dry

weight of tissue.

Measurements of Na and K and Cl content in plant tissue

Mature leaves from randomly selected finger millet plants were powdered and ashed

at 200 °C for 12 hrs. The ashes were dissolved in 5 ml 30% ammonia, and further

diluted with deionized water (Cheng et al. 2004). Concentrations of Na^+ and K^+ ions

were measured using a flame atomic absorption spectrometry. The concentration of

chloride ions was determined after aqueous extraction of 1 g of the plant material in

25 ml of distilled water. Concentrations of Cl^- ions were determined by titration from

the infiltrated solution using silver nitrate in the presence of potassium chromate as

described by Eaton et al. (1995).

Statistical analysis

A completely randomized block design with five replications for each experiment was

used and the results represent mean ± standard error. Analysis of variance (ANOVA)

was performed using the Minitab statistical computer software version 17 (Minitab

Inc., State College, PA, USA) and differences between means were accomplished

using the Fisher's protected LSD test at a confidence level of 95% ($p \leq 0.05$).

Relationships between the assessed features were performed by Pearson's correlation.

Principal component analysis (PCA) and Cluster analysis (CA) were carry out using

the FactoMineR (Factor analysis and data mining with R) package (Husson et al.,

2008).

Results

The present study investigated the changes growth parameters, relative water content,

lipid peroxidation level, proline content, reducing sugar and total protein under NaCl

induced salinity stress in six finger millet varieties. The parameters analyzed exhibited

significant variations among the varieties.

Effects of salt stress on seed germination

The effect of salinity stress on finger millet seeds germination, evaluated by the percentage of germinated seeds after 17 days, is as shown in Table I. Our results indicate that for all varieties, the germination rate decreased with an increase of the NaCl concentration and varied among the varieties. This decrease in germination rate was most profound at 200 mM and 300mM NaCl concentrations where 0 % germination rate were recorded for all six varieties. In contrast, at moderate stress levels (100 mM NaCl), significant differences in germination profile was observed with GBK043122 having the highest germination rate (46.25%) compared to others whose germination rates ranged from 3.75% to 22.50%. The germination percentage under control conditions was also distinct among the six finger millet varieties and ranged ranging from 90.00% for GBK043137 to 56.25% for GBK043122 (Table 1).

Growth characteristics in finger millet varieties under salt stress

After phenotypic observation, chlorosis (yellowish color) was observed in all plants under salinity conditions. Leaf chlorosis (yellowish color), leaf scorch, slowed and delayed growth and enlargement of the leaves were distinctly observed in seedlings of all varieties under salinity stress. Plants growing under control conditions exhibited healthy leaves and normal shoot and root developmental stages (Figure 1). The shoot length progressively retarded with increase in NaCl concentration (Table 2). Particularly, the shoot height of GBK043128 population was significantly reduced at the end of under severe salt stress conditions (300 mM NaCl) by about 72.09% while GBK043124 had the least shoot height reduction rate at 63.33% when compared to the control plants (Table 2). Significance variations on the effect of NaCl on shoot length were only observed at 200 mM NaCl concentration. Higher salt concentrations did not record any varietal difference on shoot length (Table 2). Similarly, increasing salinity stress resulted in gradual reductions in plant root lengths in all studied varieties ranging from 20.9% for GBK043137 to 36.1% for GBK043128 compared to their respective controls (Table 3). We also observed significant differences between varieties in root length values across the salt concentrations, signifying that increased

272 salt stress adversely affected root length growth in the varieties at different degrees

273 (Table 3).

Relative water content

The changes in leaves RWC along with increase in salinity stress are presented in

Table 4. The leaves relative water content of all varieties under control conditions

were similar ranging from 79.44 to 87.86%. Exposition to increasing salinity stress

progressively reduced water potential of leaves in all varieties compared to their

respective control plants leaves and they exhibited variation in their relative water

content. Variety GBK043094 tolerated salinity stress better with the least reduction in

relative water content under severe salinity stress (300 mM NaCl) compared to the

others (Table 4).

Effects of salt stress on chlorophyll content

Analysis of total chlorophyll content demonstrated significant differences in

photochemistry among varieties and the salt treatments (Table 5). More specifically,

for all the varieties, the addition of $NaCl_2$ elicited significant decrease in chlorophyll

content compared to the non-saline treatments and inverse relationship between

salinity stress and total chlorophyll content in all finger millet varieties was observed.

In contrast, plants grown under normal conditions maintained a relatively high levels

total chlorophyll content and interestingly, they did not have similar chlorophyll

content. Under saline conditions, photosynthetic pigment of varieties GBK043137

and GBK043128 were found to be extremely reduced with reduction percentages of

48.22% and 39.54%, respectively. However, GBK043124 retained a relatively higher

chlorophyll content compared to its respective control value, under 300 mM NaCl

stress conditions (Table 5). These findings signified that salinity stress may have

damaged the photochemical apparatus of the plant leaves.

Proline accumulation and lipid peroxidation assay

301 Free proline content was estimated in all six finger millet varieties at early seedling

302 growth stage to evaluated their effect under NaCl induced osmotic stress and the data

303 is shown in Table 6. Increasing salt concentrations from 100 to 200 and 300 mM

NaCl application remarkably induced increased free proline content in the plants by an average of 1.7-, 2.2- and 3.0-fold change, respectively, relative to the levels in the control plants (Table 6). GBK043094 variety had the significantly highest proline content, followed by GBK043137, GBK043124 and GBK043122 while GBK043128 and GBK043050 had the lowest (Table 6). In unstressed plants, proline concentration was similar. As shown in Table 7, we observed continuous increase in malondialdehyde content in leaves of all varieties tested in response to salinity stress relative to their respective controls and the magnitude of response differed among the varieties. A continuous increase in the level of lipid peroxidation was observed with increasing level of salinity in all the varieties. The malondialdehyde levels (μmol/g FW) was elevated to 20.7%, 31.3% and 51.2% at 100, 200 and 300 mM NaCl, respectively, as compared to unstressed plants (Table 7). Malondialdehyde content was significantly elevated in GBK043050, GBK043122 GBK043124 and GBK043128 under severe salinity stress (300 mM NaCl) treatments signifying higher rates of oxidative damage and lipid peroxidation whereas GBK043094 and GBK043137had lower levels of malondialdehyde at corresponding salinity stress (Table 7).

Reducing sugars and protein contents under NaCl stress

The impact of salinity treatment triggered substantial elevation in reducing sugar amounts in the stressed plants when compared to control the experiments (Table 8). Increasing salt concentration caused an increase in reducing sugar amounts in the stressed plant shoots and highest accretion of reducing sugar was found in 100 mM NaCl stress followed by 200 mM and 300 mM NaCl treatments. However, varietal differences difference was seen and the increase was remarkably highest in GBK043094, followed by GBK043050, GBK043137and GBK043122 while GBK043128 had the lowest amount (Table 8). Plants under control conditions had the lowest protein content ranging from 1.20 to 2.23 mg/g FW reducing whereas the highest reducing sugar content protein content of 4.47 to 6.45 mg/g FW was found in plants treated with 300 mM NaCl (Table 8). As showed in Table 9, increasing NaCl concentration had a substantial impact on the protein content of finger millet plants and the response was in a dose dependent relationship. A clear varietal difference was

observed and significantly higher levels of protein were found in GBK043094, GBK043050 and GBK043122 than the rest, under control and also stress conditions (Table 9).

Effect of salinity on shoot Na, K and Cl ion composition

The salinity treatments, varieties and the synergy effects were significant for the concentrations of all leaf ions (Table 10). As expected, the level of Na^+ and Cl^- in all varieties was higher under salt stress but differed in the degree of the increase. The gradual increase of salinity stress triggered a gradual rise of both ion concentration in finger millet leaves. The average levels of Na^+ in leaves ranged from 5.37 to 7.82 mg/g DW for plants grown in control conditions and from 12.3 to 96.2 mg/g DW for salinity stressed plants. Under 300 mM NaCl stress treatments, the different varieties increased their Na ion concentration from 6.8- to 13.1-fold when compared to the controls. GBK043124, GBK043137 and GBK043094 displayed statically the minimum increase of Na under salinity stress (Table 10). On the other hand, the leaf Cl^- levels ranged from 2.5 to 5.1 mg/g DW for finger millet plants under control conditions and from 5.0 to 17.8 mg/g DW for plants under salinity stress. GBK043050 had the lowest concentration of Cl^- under untreated and salinity stress treatments. GBK043124 had the least (3.0.5-fold) increase in Cl ion concentration under salt treatment, while GBK043094 had the largest (4.2-fold) increase (Table 10). In contrast, salinity stress induced significant reduction of K^+ concentration in leaves of finger millet plants irrigated with three NaCl doses (Table 10). In comparison to control experiments, potassium ions concentration decreased by about 18.6, 53.3 and 72.6 % in leaves of plants grown under 100, 200 and 300 mM NaCl respectively. GBK043094 upheld the highest concentration of K^+ and had a 74.0% decline in K concentration while, GBK043050 had the highest decrease in K content (78.9%) under salinity conditions (Table 10). The lowest potassium ion concentration under salinity was found in GBK043128 followed by GBK043124 (Table 10). The leaf K^+/Na^+ ratios differed among the varieties of finger millet studied, ranging from 0.05 in both GBK043094 to 0.02 in GBK043050. Varieties, GBK043094 and GBK043137 presented the greatest K^+/Na^+ ratio under salinity stress owing to low concentration of in the leaves (Table 10).

Discussion

Plants tolerance to salinity stress is a complex trait which is ascribed to a plethora of related morphological, physiological and biochemical adaptive responses and operate synergistically to lessen cell hyperosmolarity and the ensuing ion disequilibrium (Parihar et al., 2015). In this regard, screening and selection finger millet varieties tolerant to salinity stress is essential in order to understand their adaptations under saline soils and for successful production of finger millet in salinity prone areas. In this study, six finger millet varieties from different agroecological zones in Kenya were subjected to different levels of salinity stress, and our findings show tremendous variabilities occur within the tested parameters.

Seed germination and seedling emergence are fundamental biological processes in plant growth and development cycles, and therefore excellent seed germination and emergence are important for attainment of high yields and increasing concentrations of salt adversely affects germination process (Laghmouchi et al., 2017; Anuradha et al., 2001). In the present study, the germination percentage was delayed or constrained under salinity stress compared to control growth conditions. The observed decrease in germination rate under the salinity stress could be attributed to salt toxicity and changes in cellular osmotic potential. We found out that Under higher high hypertonic potential, the reduction in the germination rate was less for the salinity-tolerant variety (GBK043094), compared with most salt sensitive variety (GBK043050). Our finding was in accordance to previous work in lettuce (Ahmed et al., 2019), alfalfa (Sandhu et al., 2017) and wheat (Tounsi et al., 2017) under saline conditions. In addition, a high degree of shoot growth depression in seedlings grown under salinity stress was clearly noticeable, more in the salt-sensitive varieties, which displayed reduced leaf area, leaf chlorosis, leaf burns and plant death, symptoms associated with plant toxicity. Slower growth of both shoots and roots is an adaptive characteristic for plant survival under salinity conditions because this permits the plants to commit numerous resources to mitigate the stress (Soares et al., 2018). Retarded shoot growth under salinity stress could be ascribed to the reduction in osmotic potential due to extra concentration of sodium and chloride ions in the shoot

and root zone resulting to a nutritional imbalance and also due to the deviation of energy destined for growth and development to exclude sodium ions cellular absorption and biosynthesis of solutes for preservation cell turgor during hypertonic saline conditions. The observed reduction of leaf area under salinity treatments compared to control plants also suggests that salinity stress may affect plant growth through reduction in leaf area. Previous works have disclosed that salt tolerant plants displays less growth retardation and have relatively higher growth rate compared to sensitive ones under salinity stress (Carillo et al., 2019; Hussain et al., 2018; Sarabi et al., 2017;). Consequently, our findings suggest that GBK043128 and GBK043137 have a better capacity to sustain growth and development under salt treatments compared to other finger millet varieties studied, to sustain growth and production under salinity conditions (Table 2). Further, roots are often reported to play a key role in the salt tolerance of plants as they represent the first organs that control the uptake and translocation of nutrients and salts throughout the plant. Because of their direct exposure to saline environment, root growth is also vulnerable to salt stress although the extend is less than that of the shoots (Munns and Tester, 2008). The inhibition of root growth in plants adversely affects the survival and productivity of the plants and therefore, root growth under saline conditions may serve as good indicator in the first steps of screening for salinity tolerance programs. The growth of roots varies widely due to soil conditions because the status of all nutrient in plants is maintained from the soil with the help of roots. Root growth rate may be severely affected by saline soils and reduction may even be recorded in salt-tolerant plants. In agreement with previously published studies on the effects of salinity on root elongation (Cirillo et al., 2019; Dugasa et al., 2019), salinity treatments were found to cause stunted root growth. The growth-promoting effect under salinity stress could be due to an increase in the osmotic potential of the cells in the elongation zone coupled with enhanced cell division. The absorbed ions at this point could be quickly compartmentalized into the vacuoles without getting to the maximum capacity, thereby increasing the turgor within the cells and stimulating cell elongation. We also observed that the effects of salinity stress on root grow was much less compared to that of the shoot. This feature could be explained by the fact that roots are less affected by salt salinity due to

transport of ions to other plant organs and hence the stressed roots to maintain

osmotic balance.

Accumulation of ions in plant tissues is regularly used to evaluate the capability of a

plant to resist salt stress and salinity is known to cause fluctuations of macronutrients.

The concentration of sodium, potassium and calcium ions and the K/Na ratio are vital

features that usually used for screening of salt tolerant plants (Sarabi et al., 2017). We

used leaf tissues because they are more sensitive to salt and start displaying toxicity

much earlier compared to other plant organs (Munns and Tester, 2008). Our study

revealed that the NaCl treatments increased the finger millet leaf Na^+ and Cl^-

concentrations. Contrary, salinity treatments caused decease of K^+ in all varieties,

probably due to membranes depolarization and loss of Ca^+ ion due to the

displacement by Na^+ ions. It has also been established that Na^+ and K^+ have similar

cellular effects despite the fact Na^+ inhibits K^+ absorption through binding and

obstructing to its transport system (Flowers and Yeo, 1986). It has been established

that Na^+ and K^+ have similar cellular effects despite the fact Na^+ inhibits K^+

absorption through binding and obstructing to its transport system (Flowers and Yeo

(1986). Many studies have reported that plants growing under high NaCl

concentrations have low ratios of K^+/Na^+ ratio caused by deficiency of intracellular K^+

(Dugasa et al., 2018; Cirillo et al., 2018; Sandhu et al., 2017; Sarabi et al., 2017). The

same phenomenon was also observed in this study, where increment of NaCl

concentration decreased the leaf K^+/Na^+ ratios. In our study, we observed a clear

association between K^+/Na^+ ratio and salinity tolerance and varieties, GBK043137

and GBK043094 showed the highest K^+/Na^+ ratios under both control and NaCl

treatments, however, these varieties were placed at the highest ranking for salinity

tolerance index. Usually, cellular influx of Cl^- ions influx require energy in a reaction

mechanism catalysed by a $Cl^-/2H^+$ coupled antiporters and symporters and it is

typically taken up freely with water uptake, and is therefore accumulated in leaf

organs depending on the transpiration rate (Munns and Tester, 2008). Like Na^+, Cl^-

ions may also be sequestered in cell vacuoles. In our study, the concentration of Cl^- in

leaves was higher than that of Na^+ and this may be justified by the partial control of

Na$^{+,}$ at roots. Comparable results were also exhibited by melon (Sarabi et al., 2017)
and cucumber (Colla et al., 2012).

Several studies suggest chlorophyll content as a biochemical marker of salt tolerance
in plants (Ishikawa Shabala, 2019; Taïbi et al., 2016; Sairam et al., 2005). It is known
that salt tolerant plants show increased or unchanged chlorophyll levels under salinity
conditions whereas chlorophyll contents decreased in salt-sensitive plants (Stepien
and Johonson, 2009; Ashraf and Harris, 2013). In general, decrease of chlorophyll
content under salt stress is considered to be a result of slow synthesis or fast
breakdown of the pigments in cells (Ashraf, 2003). The decrease in total chlorophyll
content may also be observed due to ion accumulation and functional disorders
observed during stoma opening and closing under salinity stress (Nawaz et al., 2010).
Another reason for the decrease of chlorophyll content under salt conditions is stated
to be the rapid maturing of leaves (Yeo et al., 1991). In our study statistically
significant decrease in total chlorophyll content was observed with increasing salt
concentration. Similar results were reported by Ashraf and Yousafali (1998) and Ali
et al., (2004) and showed that the total chlorophyll content of rice leaves was
generally reduced under high salinity. While the other varieties recorded a decrease
over the control plants, variety GBK043094 recorded unchanged total chlorophyll
content with increase in stress (Table4). These results showed that the reduction in
chlorophyll content was variety specific and some varieties showed comparatively
lesser quantum of negative variation in chlorophyll content thus indicating their
potential to grow and perform moderately well even under higher levels of salt stress.
High salt concentration induced reduction of total chlorophyll content indicates that
salt stress induces chlorophyll degradation and destruction of chloroplast structures.

All plants employ complex biochemical defensive mechanisms against oxidative
injury of free radicals and ROS during abiotic stresses. Among these defence systems
is the aggregation of compatible solutes such as proline, an osmoprotectant that
preserves membrane integrity and mitigates oxidative burst in plant challenged by salt
stress Ahmed et al. 2013, Rao et al., 2013). In addition, proline exists in all plant
organs, accumulating in greater proportions compared to other amino acids in salinity

stressed plants (Banu et al., 2009). Although the beneficial outcome of proline

overproduction in plants during salinity stress have been explicated, the definite roles

of proline accretion are still obscure (Banu et al., 2009; Verbruggen and Hermans,

2008). Our study reported an increased concentration of free proline content in all six

finger millet varieties with GBK043094 and GBK043137 displaying higher free

proline amounts at all salinity treatments suggesting that they are comparatively more

tolerant to salinity stress than the rest and which may be related to their competitive

ability under saline stress against oxidative stress. Based on these results, it is worth

noting that increased concentration of free proline content in finger millet plants

subjected to salinity treatments corresponded to improved salinity tolerance.

Degradation of polyunsaturated fatty acids in plants yields malondialdehyde (MDA) a

biomarker for determining the degree of lipid peroxidation and cellular membrane

(Yang et al., 2018). Results from our study reveals that MDA the content in stressed

plants raised with increasing stress levels corroborate with those exhibited in other

plant species like *Lycium ruthenicum* (Li et al., 2019), wheat (Dugasa et al., 2018) and

Cucumis melo L. (Sarabi et al., 2017). Our results indicated that some varieties finger

millet may tolerate saline environments than others depending on the severity of the

stress, by lowering the rate of lipid peroxidation and the cell membrane damage and

therefore have an efficient and effective antioxidant defence mechanism. Moreover,

the strong negative correlation witnessed between MDA and shoot height (r= –0.995),

and root length (r= –0.898) affirms that the NaCl stress triggered lipid peroxidation is

one of the reasons for the observed stunted shoot and root growth in finger millet

plants.

Contrary to other osmolytes such as proline and MDA which are present at very low

amounts except when their biosynthesis is triggered by stress, compatible solutes such

as reducing sugars are elements of metabolism with different cell functional roles,

such as precursors of other metabolites, signalling molecules and major source of

energy. Their levels are highly controlled by various systems to ensure cellular

homeostasis. Reducing sugars therefore play a crucial role in plant cells osmotic

adjustment during salinity stress. The higher reducing sugar levels measured plants

with high salinity tolerant index clearly shows that the sugar contributes to osmotic

adjustment during salt treatments thus cushioning the plants against the toxic effects
of NaCl. These results are substantiated by a remarkable increase in sugar amounts in
salt tolerant genotypes in pigeon pea (Awana et al., 2019), *Juncus* sp (Hassan et al.,
2016) and wheat (Kerepesi and Galiba, 2000). Likewise, accumulation of protein
compounds has essential part in physiological responses of plant to salinity stress.
Increased production of proteins and other nitrogen containing compound may induce
the biosynthesis of osmotically active organic compounds including proteins with
osmoprotective capacities, thereby conferring salinity resistance (Ashraf and Harris,
2004). Generally, plants exposed to NaCl stress have comparatively reduced protein
levels which often results to loss of cellular turgor. Just like in our case, reduction in
the content of soluble protein was observed in maize plants subjected to salinity
treatments (von Alvensleben et al., 2013).

Lastly, it is imperative to note that the results of this study were conducted in a
laboratory set-up (artificial conditions), which may not mirror their complex natural.
However, the findings give suggestive index salinity tolerance to the studied finger
millet varieties.

Conclusions
This study gives a deep analysis of the effect of a NaCl stress treatments on the
physiological and biochemical parameters six finger millet varieties. In conclusion,
our results demonstrated salinity responses on the evaluated features with significant
varietal differences among the plants studied and supported by observations made.
From the responses of GBK043094 and GBK043137 varieties, we hypothesised that
these varieties are promising genetic resources with comparative high tolerance to
salinity and hence they may be utilised for further assessment for breeding programs
of the crop towards enhanced salinity tolerance. Our findings give suggestive salinity
tolerance index to the studied finger millet varieties and should be confirmed for in a
wide range a wide range of environmental conditions and other salt types. Lastly, it is
imperative to note that the results of this study were conducted in a laboratory set-up
(artificial conditions), which may not mirror their complex natural environments
under which the crop is grown.

References

Al-Hassan M., del Pilar López-Gresa, M., Boscaiu, M., & Vicente, O. (2016). Stress tolerance mechanisms in *Juncus*: responses to salinity and drought in three *Juncus* species adapted to different natural environments. Functional Plant Biology. 43(10):949-960. doi.org/10.1071/FP16007

Anuradha S., Rao, S. S. R. (2001). Effect of brassinosteroids on salinity stress induced inhibition of seed germination and seedling growth of rice (*Oryza sativa* L.). Plant Growth Regulation, 33(2):151-153. doi.org/10.1023/A:1017590108484

Anuradha, S., & Rao, S. S. R. (2001). Effect of brassinosteroids on salinity stress induced inhibition of seed germination and seedling growth of rice (Oryza sativa L.). Plant Growth Regulation, 33(2), 151-153.

Arnon, D. (1949). Copper enzyme in isolated chloroplast and chlorophyll expressed in terms of mg per gram. Plant Physiology, 24(1):1-15. https://www.jstor.org/stable/4258165

Ashraf M. P. J. C., & Harris P. J. C. (2004). Potential biochemical indicators of salinity tolerance in plants. Plant science, 166(1), 3-16. doi:10.1016/j.plantsci.2003.10.024

Awana M., Yadav K., Rani K., Gaikwad K., Praveen S., Kumar S., Singh A. (2019). Insights into Salt Stress-Induced Biochemical, Molecular and Epigenetic Regulation of Spatial Responses in Pigeonpea (*Cajanus cajan* L.). Journal of Plant Growth Regulation, 1-17. doi.org/10.1007/s00344-019-09955-4

Bates L. S., Waldren R. P., Teare I. D. (1973). Rapid determination of free proline for water-stress studies. Plant and Soil, 39(1):205-207. doi.org/10.1007/BF00018060

Bertazzini M., Sacchi G. A., Forlani G. (2018). A differential tolerance to mild salt stress conditions among six Italian rice genotypes does not rely on Na^+ exclusion from shoots. Journal of Plant Physiology. 226, 145-153. doi.org/10.1016/j.jplph.2018.04.011

Carillo P., Cirillo C., De Micco V., Arena,C., De Pascale S., & Rouphael, Y. (2019). Morpho-anatomical, physiological and biochemical adaptive responses to saline water of *Bougainvillea spectabilis* Willd. trained to different canopy shapes. Agricultural Water Management. 212:12-22. doi.org/10.1016/j.agwat.2018.08.037

Cheng W., Zhang G., Yao H., Dominy P., Wu W., Wang, R. (2004). Possibility of
predicting heavy-metal contents in rice grains based on DTPA-extracted levels in soil.
Communications in soil science and plant analysis, 35(19-20):2731-2745.
doi.org/10.1081/CSS-200036424

Chivenge P., Mabhaudhi T., Modi A., Mafongoya P. (2015). The potential role of
neglected and underutilised crop species as future crops under water scarce conditions
in Sub-Saharan Africa. International Journal of Environmental Research and Public
Health. 12(6):5685-5711. doi.org/10.3390/ijerph120605685

Cirillo C., Rouphael Y., Caputo R., Raimondi G., Sifola M. I., De Pascale S. (2016).
Effects of high salinity and the exogenous application of an osmolyte on growth,
photosynthesis, and mineral composition in two ornamental shrubs. The Journal of
Horticultural Science and Biotechnology. 91(1), 14-22.
doi.org/10.1080/14620316.2015.1110988

Damerval C, de Vienne D, Zivy M, Thiellement H (1986). Technical improvements in
two-dimensional electrophoresis increase the level of genetic variation detected in
wheat-seedling proteins. Electrophoresis 7(1):52–54.
doi.org/10.1002/elps.1150070108

Eaton A.D., Clesceri, L.S. Greenberg, A.E. (1995). Standard Methods for the
Examination of Water and Wastewater, 19th edition. American Public Health
Association, Washington, USA.

de Freitas P. A. F., de Souza M. R., Marques, E. C., Prisco, J. T., Gomes-Filho, E.
(2018). Salt tolerance induced by exogenous proline in maize is related to low
oxidative damage and favorable ionic homeostasis. Journal of Plant Growth
Regulation, 37(3):911-924. doi.org/10.1007/s00344-018-9787-x

Dugasa M. T., Cao F., Ibrahim W., Wu F. (2019). Differences in physiological and
biochemical characteristics in response to single and combined drought and salinity
stresses between wheat genotypes differing in salt tolerance. Physiologia Plantarum.
165: 134–143. doi:10.1111/ppl.12743

Filichkin S. A., Ansariola M., Fraser V. N., Megraw M. (2018). Identification of
transcription factors from NF-Y, NAC, and SPL families responding to osmotic stress
in multiple tomato varieties. Plant Science, 274, 441-450.
doi.org/10.1016/j.plantsci.2018.06.021

Flowers T. J., & Yeo A. R. (1986). Ion relations of plants under drought and salinity. Functional Plant Biology, 13(1):75-91. doi.org/10.1071/PP9860075

Heath R. L., Packer L. (1968). Photoperoxidation in isolated chloroplasts: I. Kinetics and stoichiometry of fatty acid peroxidation. Archives of biochemistry and biophysics, 125(1):189-198. doi.org/10.1016/0003-9861(68)90654-1

Hema R., Vemanna R. S., Sreeramulu S., Reddy C. P., Senthil K. M., Udayakumar M. (2014). Stable expression of mtlD gene imparts multiple stress tolerance in finger millet. PLoS One, 9(6), e99110. doi.org/10.1371/journal.pone.0099110

Hussain M. I., Al-Dakheel A. J., & Reigosa M. J. (2018). Genotypic differences in agro-physiological, biochemical and isotopic responses to salinity stress in quinoa (*Chenopodium quinoa* Willd.) plants: Prospects for salinity tolerance and yield stability. Plant Physiology and Biochemistry. 129:411-420. /doi.org/10.1016/j.plaphy.2018.06.023

Ibrahim, E. A. (2016). Seed priming to alleviate salinity stress in germinating seeds. Journal of Plant Physiology, 192, 38-46. doi.org/10.1016/j.jplph.2015.12.011

Ishikawa, T., & Shabala, S. (2019). Control of xylem Na^+ loading and transport to the shoot in rice and barley as a determinant of differential salinity stress tolerance. Physiologia Plantarum. 165(3):619-631. doi.org/10.1111/ppl.12758

Johnson, G., Lambert, C., Johnson, D. K. D., & Sunderwirth, S. G. (1964). Colorimetric determination of glucose, fructose, and sucrose in plant materials using a combination of enzymatic and chemical methods. Journal of Agricultural and Food Chemistry, 12, 216-219. doi: 10.1021/jf60133a007

Kerepesi I., & Galiba G. (2000). Osmotic and salt stress-induced alteration in soluble carbohydrate content in wheat seedlings. Crop Science. 40(2):482-487. doi:10.2135/cropsci2000.402482x

Kumar V., Khare T. (2016). Differential growth and yield responses of salt-tolerant and susceptible rice cultivars to individual (Na^+ and Cl^-) and additive stress effects of NaCl. Acta Physiologiae Plantarum. 38(170):1-9. doi: 10.1007/s11738-016-2191-x

Laghmouchi Y., Belmehdi O., Bouyahya A., Senhaji N. S., & Abrini J. (2017). Effect of temperature, salt stress and pH on seed germination of medicinal plant *Origanum compactum*. Biocatalysis and Agricultural Biotechnology. 10:156-160. doi.org/10.1016/j.bcab.2017.03.002

Li Y., Zhang T., Zhang Z., He K. (2019). The physiological and biochemical
photosynthetic properties of *Lycium ruthenicum* Murr in response to salinity and
drought. Scientia Horticulturae. 256(108530):1-9.
doi.org/10.1016/j.scienta.2019.05.057

Munns R., James R. A., Läuchli, A. (2006). Approaches to increasing the salt
tolerance of wheat and other cereals. Journal of Experimental Botany, 57(5):1025-
1043. doi.org/10.1093/jxb/erj100

Munns, R., & Tester, M. (2008). Mechanisms of salinity tolerance. Annu. Rev. Plant
Biol., 59, 651-681.

Parihar, P., Singh, S., Singh, R., Singh, V. P., & Prasad, S. M. (2015). Effect of
salinity stress on plants and its tolerance strategies: a review. Environmental Science
and Pollution Research, 22(6), 4056-4075. doi: 10.1007/s11356-014-3739-1

Qadir M., Quillérou E., Nangia V, Murtaza G., Singh M., Thomas R. J., Drechsel P.,
Noble A. D. Economics of salt☐induced land degradation and restoration. Natural
Resources Forum. 38(4): 282-295. doi: 10.1111/1477-8947.12054

Rouphael, Y., Cardarelli, M., Rea, E., & Colla, G. (2012). Improving melon and
cucumber photosynthetic activity, mineral composition, and growth performance
under salinity stress by grafting onto Cucurbita hybrid rootstocks. Photosynthetica,
50(2):180-188. doi.org/10.1007/s11099-012-0002-1

Sairam R. K., Srivastava G. C., Agarwal S., & Meena R. C. (2005). Differences in
antioxidant activity in response to salinity stress in tolerant and susceptible wheat
genotypes. Biologia Plantarum, 49(1): 85-91. doi.org/10.1007/s10535-005-5091-2

Sandhu D., Cornacchione M. V., Ferreira J. F., Suarez, D. L. (2017). Variable salinity
responses of 12 alfalfa genotypes and comparative expression analyses of salt-
response genes. Scientific Reports, 7(42958):1-18. doi: 10.1038/srep42958

Sarabi B., Bolandnazar S., Ghaderi N., Ghashghaie J. (2017). Genotypic differences
in physiological and biochemical responses to salinity stress in melon (*Cucumis melo*
L.) plants: Prospects for selection of salt tolerant landraces. Plant Physiology and
Biochemistry, 119, 294-311. doi.org/10.1016/j.plaphy.2017.09.006

Shabala S., Hariadi Y., Jacobsen S. E. (2013). Genotypic difference in salinity
tolerance in quinoa is determined by differential control of xylem Na^+ loading and

stomatal density. Journal of Plant Physiology. 170(10):906-914. doi.org/10.1016/j.jplph.2013.01.014

Shabala, S. (2013). Learning from halophytes: physiological basis and strategies to improve abiotic stress tolerance in crops. Annals of botany. 112(7):1209-1221. doi.org/10.1093/aob/mct205

Soares A. C., Geilfus C. M., & Carpentier S. C. (2018). Genotype-specific growth and proteomic responses of maize towards salt stress. Frontiers in plant science, 9(661):1-15. doi.org/10.3389/fpls.2018.00661

Taïbi K., Taïbi F., Abderrahim L. A., Ennajah A., Belkhodja M., & Mulet, J. M. (2016). Effect of salt stress on growth, chlorophyll content, lipid peroxidation and antioxidant defence systems in *Phaseolus vulgaris* L. South African Journal of Botany, 105: 306-312. doi.org/10.1016/j.sajb.2016.03.011

Tedeschi A., Lavini A., Riccardi M., Pulvento, C., d'Andria, R. (2011). Melon crops (*Cucumis melo* L., cv. Tendral) grown in a mediterranean environment under saline-sodic conditions: Part I. Yield and quality. Agricultural water management. 98(9):1329-1338. doi.org/10.1016/j.agwat.2011.04.007

Tlig T., Gorai M., Neffati M. (2008). Germination responses of *Diplotaxis harra* to temperature and salinity. Flora-Morphology, Distribution, Functional Ecology of Plants, 203(5), 421-428. doi.org/10.1016/j.flora.2007.07.002

Tounsi S., Feki K., Hmidi D., Masmoudi K., Brini F. (2017). Salt stress reveals differential physiological, biochemical and molecular responses in *T. monococcum* and *T. durum* wheat genotypes. Physiology and molecular biology of plants, 23(3):517-528. doi.org/10.1007

von Alvensleben N., Stookey K., Magnusson M., & Heimann, K. (2013). Salinity tolerance *of Picochlorum atomus* and the use of salinity for contamination control by the freshwater cyanobacterium *Pseudanabaena limnetica*. PloS ONE, 8(5): e6356. doi.org/10.1371/journal.pone.0063569

Yang C., Shahidi F., & Tsao R. (2018). Biomarkers of oxidative stress and cellular based assays of indirect antioxidant measurement. Measurement of Antioxidant Activity & Capacity. 165-186. doi.org/10.1002/9781119135388.ch9

Zhang H. J., Zhang N. A., Yang R. C., Wang L., Sun Q. Q., Li, D. B., Cao Y.Y, Weeda S, Zhao B, Ren S, Guo Y. D. (2014). Melatonin promotes seed germination

under high salinity by regulating antioxidant systems, ABA and GA 4 interaction in

cucumber (*C sativus* L.). Journal of Pineal Research, 57(3):269-279.

doi.org/10.1111/jpi.12167

Zhu J. K. (2003). Regulation of ion homeostasis under salt stress. Current Opinion in

Plant Biology. 6(5):441-445. doi.org/10.1016/S1369-5266(03)00085-2

Figures

Figure 1. Effect of salinity stress on growth of finger millet. (A) seedling growth

on 300 mM NaCl. (B) seedling growth on 200mM NaCl (C) seedling growth on 100

mM NaCl; **(D)** seedling growth on 0 mM NaCl.

Tables

Table1. Effects of NaCl on germination rate of six finger millet varieties

Variety	Germination rate (%)			
	0 mM	100 mM	200 mM	300 mM
GBK043137	90.0 ± 3.5^a	3.8 ± 3.8^c	0.0 ± 0.0^a	0.0 ± 0.0^a
GBK043128	65.0 ± 3.5^{bc}	18.8 ± 2.4^{bc}	0.0 ± 0.0^a	0.0 ± 0.0^a
GBK043124	80.0 ± 8.4^a	37.5 ± 7.8^{ab}	0.0 ± 0.0^a	0.0 ± 0.0^a
GBK043122	56.3 ± 5.2^c	46.3 ± 9.7^a	0.0 ± 0.0^a	0.0 ± 0.0^a
GBK043094	63.8 ± 1.3^{bc}	22.5 ± 11.6^{bc}	0.0 ± 0.0^a	0.0 ± 0.0^a
GBK043050	76.3 ± 3.7^{ab}	18.85 ± 5.5^{bc}	0.0 ± 0.0^a	0.0 ± 0.0^a

Values within a column marked with different superscript in each column differ significantly at $p < 0.05$ [Fishers LSD]. Each value represented as mean $\pm$ SD are the mean of three replications.

Table2. Effect of NaCl on growth of finger millet

Variety	Seedlings shoot length (cm) under NaCl stress			
	0 mM	100 mM	200 mM	300 mM
GBK043137	3.8 ± 0.4^{ab}	2.5 ± 0.3^a	1.6 ± 0.2^{ab}	1.2 ± 0.1^a
GBK043128	4.3 ± 0.2^a	2.6 ± 0.4^a	1.9 ± 0.2^a	1.2 ± 0.2^a
GBK043124	3.0 ± 0.3^b	2.4 ± 0.2^{ab}	1.6 ± 0.3^{ab}	1.1 ± 0.1^a
GBK043122	3.2 ± 0.4^b	2.5 ± 0.2^a	1.2 ± 0.2^{bc}	1.0 ± 0.0^a
GBK043094	3.1 ± 0.3^b	2.3 ± 0.2^{ab}	1.0 ± 0.0^c	1.1 ± 0.1^a
GBK043050	3.1 ± 0.2^b	1.7 ± 0.1^b	1.3 ± 0.2^{bc}	1.1 ± 0.1^a

Values within a column marked with different superscript in each column differ significantly at $p < 0.05$ [Fishers LSD]. Each value represented as mean $\pm$ SD are the mean of three replications.

Table 3. Effect of NaCl on growth root growth

Variety	Seedlings root length (cm) under NaCl stress			
	0 mM	100 mM	200 mM	300 mM
GBK043137	$6.8+0.4^b$	$6.4+0.4^{ab}$	$5.9+0.2^a$	$5.4+0.2^a$
GBK043128	$8.0+0.4^a$	$7.2+0.4^a$	$5.9+0.3^a$	$5.1+0.3^{ab}$
GBK043124	$6.9+0.6^b$	$6.3+0.6^{ab}$	$5.6+0.4^a$	$5.0+0.46^{ab}$
GBK043122	$6.7+0.2^b$	$6.2+0.5^b$	$5.5+0.3^a$	$4.9+0.2^{ab}$
GBK043094	$6.8+0.5^b$	$6.2+0.7^b$	$5.9+0.4^a$	$5.4+0.4^a$
GBK043050	$7.1+0.4^b$	$6.4+0.5^{ab}$	$5.6+0.2^a$	$4.8+0.3^b$

Values within a column marked with different superscript in each column differ significantly at $p < 0.05$ [Fishers LSD]. Each value represented as mean $\pm$ SD are the mean of three replications.

Table 4. Effect of NaCl on relative water content

Variety	Seedlings relative water content (%) under NaCl stress

	0 mM	100 mM	200 mM	300 mM
GBK043137	85.3±4.1^a	71.5±4.1^a	35.0±3.9^a	37.1±3.3^b
GBK043128	87.9±5.3^a	71.5±4.1^{abc}	35.0±3.9^b	26.8±2.3^c
GBK043124	84.8±4.9^a	67.2±3.4^{bc}	34.2±5.0^b	28.2±2.6^c
GBK043122	83.0±1.8^a	68.0±1.9^{bc}	33.7±3.3^b	46.7±9.2^c
GBK043094	82.1±6.7^a	72.3±3.7^{ab}	51.3±6.1^a	46.7±9.2^a
GBK043050	79.4±4.6^a	65.4±4.8^c	33.2±4.5^b	27.6±3.7^c

750 Values within a column marked with different superscript in each column differ
751 significantly at $p < 0.05$ [Fishers LSD]. Each value represented as mean ± SD are the
752 mean of three replications.
753
754
755 **Table 5. Effect of salinity stress on total chlorophyll content of finger millet**
756 **varieties**

Variety	Seedlings chlorophyll content (mg/g FW) under NaCl stress			
	0 mM	100 mM	200 mM	300 mM
GBK043137	8.4±0.4^a	8.1±1.8^a	5.0±0.4^a	4.4±0.6^a
GBK043128	9.1±1.0^b	7.5±1.5^b	6.4±0.5^b	5.5±0.1^b
GBK043124	5.9±0.1^c	6.8±0.1^c	7.3±0.2^c	5.5±0.1^c
GBK043122	7.3±1.9^d	5.9±0.1^d	6.1±1.0^d	5.8±0.^{7d}
GBK043094	6.2±0.5e	5.0±0.9	5.0±0.8^e	4.7±1.4^e
GBK043050	5.1±1.6^f	4.0±0.9^f	5.0±0.4^f	3.9±0.7^f

757 Values within a column marked with different superscript in each column differ
758 significantly at $p < 0.05$ [Fishers LSD]. Each value represented as mean ± SD are the
759 mean of three replications.
760
761
762 **Table 6. Effect of salinity stress on free proline content of finger millet varieties**

Variety	Proline content (µg/g FW) under NaCl stress			
	0 mM	100 mM	200 mM	300 mM
GBK043137	200.9±2.4^a	411.9±13.4^a	529.3±3.0^{ab}	655.2±28.6^b
GBK043128	224.3±3.6^a	340.3±33.9^b	471.3±63.7^{bc}	571.3±37.1^c
GBK043124	208.4±30.6^a	322.4±34.0^b	417.9±50.6^c	585.1±86.6^{bc}
GBK043122	234.5±16.2^a	344.0±18.2^b	433.2±12.3^c	666.7±2.1^b
GBK043094	208.2±14.4^a	401.6±25.7^a	558.0±12.9^a	801.9±22.8^a
GBK043050	212.8±21.6^a	319.7±7.5^b	404.4±34.9^c	560.5±53.4^c

763 Values within a column marked with different superscript in each column differ
764 significantly at $p < 0.05$ [Fishers LSD]. Each value represented as mean ± SD are the
765 mean of three replications.
766
767
768 **Table 7. Effect of salinity stress on free proline content of finger millet varieties**

Variety	Malondialdehyde content (µg/g FW) under NaCl stress			
	0 mM	100 mM	200 mM	300 mM
GBK043137	1.94±0.1^a	2.2±0.3^b	2.4±0.5^b	2.7±0.4^b
GBK043128	2.21±0.3^a	2.9±0.2^{ab}	2.9±0.4^{ab}	3.6±0.2^a

GBK043124	2.67 ± 0.4^a	3.3 ± 0.3^a	3.4 ± 0.5^a	3.7 ± 0.6^a
GBK043122	2.22 ± 0.4^a	2.8 ± 0.2^{ab}	3.3 ± 0.1^a	3.8 ± 0.4^a
GBK043094	1.96 ± 0.2^a	2.3 ± 0.3^b	2.47 ± 0.2^b	2.8 ± 0.3^b
GBK043050	2.19 ± 0.8^a	2.5 ± 0.9^b	3.0 ± 0.7^{ab}	3.9 ± 0.5^a

Values within a column marked with different superscript in each column differ significantly at $p < 0.05$ [Fishers LSD]. Each value represented as mean ± SD are the mean of three replications.

Table 8. Effect of salt stress on reducing sugars on finger millet

Variety	Reducing sugars content (mg/g FW) under NaCl stress			
	0 mM	100 mM	200 mM	300 mM
GBK043137	1.6 ± 0.4^{bc}	2.1 ± 0.6^{ab}	4.0 ± 0.8^{bc}	4.9 ± 0.9^{bc}
GBK043128	1.2 ± 0.3^c	1.6 ± 0.3^b	3.7 ± 0.7^{bc}	4.6 ± 0.8^c
GBK043124	1.3 ± 0.3^c	1.7 ± 0.3^b	3.3 ± 0.5^c	4.7 ± 0.3^c
GBK043122	1.8 ± 0.4^{abc}	2.1 ± 0.4^{ab}	3.7 ± 0.5^{bc}	5.5 ± 0.2^{bc}
GBK043094	2.2 ± 0.3^a	2.7 ± 0.2^a	5.0 ± 0.0^a	6.5 ± 0.5^a
GBK043050	2.1 ± 0.4^{ab}	2.5 ± 0.4^a	4.4 ± 0.3^{ab}	5.8 ± 0.4^{ab}

Values within a column marked with different superscript in each column differ significantly different at $p < 0.05$ [Fishers LSD]. Each value represented as mean ± SD are the mean of three replications.

Table 9 Effect of salt stress on total protein on finger millet

Variety	Total protein content (mg BSA/g FW) under NaCl stress			
	0 mM	100 mM	200 mM	300 mM
GBK043137	15.2 ± 1.3^b	34.4 ± 1.6^b	73.8 ± 7.3^c	95.7 ± 9.8^b
GBK043128	15.3 ± 2.1^b	33.9 ± 3.0^b	73.1 ± 7.4^c	94.7 ± 8.0^b
GBK043124	13.2 ± 1.9^b	32.1 ± 3.1^b	74.7 ± 7.1^{bc}	85.6 ± 4.1^b
GBK043122	20.0 ± 2.2^a	42.5 ± 5.2^a	95.9 ± 4.1^a	111.9 ± 7.4^a
GBK043094	20.5 ± 3.0^a	45.1 ± 5.7^a	90.5 ± 9.7^a	119.2 ± 6.5^a
GBK043050	20.4 ± 1.2^a	43.3 ± 3.3^a	89.2 ± 11.5^{ab}	117.5 ± 5.4^a

Values within a column marked with different superscript in each column differ significantly at $p < 0.05$ [Fishers LSD]. Each value represented as mean ± SD are the mean of three replications.

Table 10. Effect of salinity stress on shoot Na^+, K^+ and Cl^- contents and Na^+/K^+ ratio of finger millet grown under NaCl treatments

Variety	Treatment	Na^+ (mg/g DW)	K^+ (mg/g DW)	Cl^- (mg/g DW)	Na^+/K^+
GBK043137	Control	$5.8+0.3^b$	$8.8+0.3^{bc}$	$5.1+0.4^a$	
	100	$12.3+1.8^c$	$7.2+0.4^c$	$9.4+0.7^a$	
	200	$37.2+4.3^d$	$4.6+0.8^{ab}$	$13.7+1.7^a$	

	300	$47.0+4.3^c$	$2.2+0.1$	$17.8+2.1^a$	
GBK043128	Control	$5.4+0.4^b$	$7.3+0.4^d$	$4.3+0.2^{ab}$	
	100	$17.1+1.7^b$	$6.0+0.3^c$	$8.2+1.1^{ab}$	
	200	$56.8+4.1^{bc}$	$4.3+1.2^{ab}$	$11.0+0.3^{ab}$	
	300	$70.0+4.1^b$	$2.7+1.0^a$	$13.3+2.0^{bc}$	
GBK043124	Control	$8.5+0.5^a$	$10.4+0.6^a$	$4.9+0.9^a$	
	100	$24.6+3.1^a$	$8.5+0.7^a$	$8.9+1.0^{ab}$	
	200	$51.9+1.9^c$	$4.6+0.4^{ab}$	$8.8+0.5^a$	
	300	$82.7+28.5^{ab}$	$3.2+1.1^a$	$14.6+1.8^b$	
GBK043122	Control	$7.8+1.4^a$	$9.7+0.4^{ab}$	$3.3+0.4^{cd}$	
	100	$22.0+2.1^a$	$7.7+0.6^{ab}$	$6.2+0.8^{cd}$	
	200	$65.6+5.3^a$	$3.2+0.9^b$	$13.1+3.0^{bc}$	
	300	$96.2+19.2^a$	$2.6+0.6^a$	$11.4+1.6^{cd}$	
GBK043094	Control	$7.3+0.5^a$	$9.6+0.7^{ab}$	$3.7+0.5^{bc}$	
	100	$14.3+1.6^{bc}$	$7.5+0.6^b$	$7.5+0.4^{bc}$	
	200	$33.2+2.0^d$	$5.0+1.3^a$	$9.8+1.1^b$	
	300	$49.5+4.8^c$	$2.4+1.0^a$	$15.4+1.6^{ab}$	
GBK043050	Control	$7.6+0.5^a$	$8.6+0.5^c$	$2.5+0.4^d$	
	100	$21.5+1.5^a$	$7.4+0.7^b$	$5.0+0.6^d$	
	200	$63.6+4.6^{ab}$	$3.7+0.9^{ab}$	$6.7+1.3^c$	
	300	$92.1+4.3^a$	$1.8+0.6^a$	$10.2+0.7^d$	

788 For each sorghum NaCl treatment, values within a column sharing same letter
789 comparing NaCl treatments are not significantly different at $p < 0.05$ [Fishers LSD].
790 Each value represented as mean $\pm$ SD are the mean of three replications.
791
792