Anthropogenic disturbance drives dispersal syndromes, demography, and gene flow in

spatially structured amphibian populations

Hugo Cayuela [1,2], Aurélien Besnard[2], Julien Cote[3], Martin Laporte[2], Eric Bonnaire[4], Julian Pichenot[5],

Nicolas Schtickzelle[6], Arnaud Bellec[1], Pierre Joly[1], Jean-Paul Léna[1]

[1]UMR 5023 LEHNA, Laboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés (UMR 5023

LEHNA), 69100 Villeurbanne, France

[2]EPHE, UM, SupAgro, IRD, INRA, UMR 5175 CEFE, CNRS, PSL Research University, Montpellier

F-34293 France

[3]CNRS, Université Toulouse III Paul Sabatier, ENFA ; UMR5174EDB (Laboratoire Évolution &

Diversité Biologique) ; 118 route de Narbonne, F-31062 Toulouse, France

[4]Office National des Forêts, Agence de Verdun, 55100 Verdun, France

[5]CERFE, Centre de Recherche et Formation en Eco-éthologie (CERFE), 08240 Boult-aux-Bois,

France

[6] Earth and Life Institute, Biodiversity Research Centre, Université Catholique de Louvain, 1348

Louvain-la-Neuve, Belgium

Running head: Disturbance and dispersal

Abstract

There is growing evidence that anthropogenic landscapes can strongly influence the evolution of

dispersal, particularly through fragmentation, and may drive organisms into an evolutionary trap by

suppressing dispersal. However, the influence on dispersal evolution of anthropogenic variation in

habitat patch turnover has so far been largely overlooked. In this study, we examined how human-driven

variation in patch persistence affects dispersal rates and distances, determines dispersal-related

phenotypic specialization, and drives neutral genetic structure in spatially structured populations. We

addressed this issue in an amphibian, *Bombina variegata*, using an integrative approach combining

capture–recapture modeling, demographic simulation, common garden experiments, and population

genetics. *B. variegata* reproduces in small ponds that occur either in habitat patches that are persistent

(i.e. several decades or more), located in riverine environments with negligible human activity, or in

patches that are highly temporary (i.e. a few years), created by logging operations in intensively

harvested woodland. Our capture–recapture models revealed that natal and breeding dispersal rates and

distances were drastically higher in spatially structured populations (SSPs) in logging environments than

in riverine SSPs. Population simulations additionally showed that dispersal costs and benefits drive the

fate of logging SSPs, which cannot persist without dispersal. The common garden experiments revealed

that toadlets reared in laboratory conditions have morphological and behavioral specialization that

depends on their habitat of origin. Toadlets from logging SSPs were found to have higher boldness and

exploration propensity than those from riverine SSPs, indicating transgenerationally transmitted

dispersal syndromes. We also found contrasting patterns of neutral genetic diversity and gene flow in

riverine and logging SSPs, with genetic diversity and effective population size considerably higher in

logging than in riverine SSPs. In parallel, intra-patch inbreeding and relatedness levels were lower in

logging SSPs. Controlling for the effect of genetic drift and landscape connectivity, gene flow was found

to be higher in logging than in riverine SSPs. Taken together, these results indicate that anthropogenic

variation in habitat patch turnover may have an effect at least as important as landscape fragmentation

on dispersal evolution and the long-term viability and genetic structure of wild populations.

Keyword: Dispersal, movement behavior, behavioral syndrome, metapopulation, genetic structure,

amphibian

Introduction

In the age of the Anthropocene, a significant proportion of land cover has been replaced by human-dominated landscapes (Foley et al. 2005, Pereira et al. 2010, Gibson et al. 2011, Tilman et al. 2017), with the result that the conditions prevailing in these anthropogenic environments now shape the evolutionary course of almost all species (Otto 2018, Pelletier & Coltman 2018). Land use changes usually have the simultaneous effects of habitat loss, alteration, and/or fragmentation into small habitat patches isolated in a more or less hostile matrix (i.e. unsuitable habitat) (Fahrig 2003, Villard & Metzger 2014). While it is widely accepted that habitat loss is the main factor involved in local extinction and biodiversity loss (Sala et al. 2000, Pereira et al. 2010, Newbold et al. 2015, Tilman et al. 2017), it is also increasingly recognized that habitat alteration and fragmentation are critical ecological and evolutionary drivers in anthropogenic landscapes (Villard & Metzger 2014, Haddad et al 2015). Interestingly, the risk of extirpation from anthropogenic landscapes appears to differ between species (Frishkoff et al. 2014, Edward et al. 2015, Nowakoski et al. 2018) and research is needed to identify the phenotypic traits that allow some species to cope with and succeed in human-dominated contexts.

Dispersal, i.e. the movement from birth to breeding patch (natal dispersal) or between successive breeding patches (breeding dispersal), is a key ecological and evolutionary process. Dispersal provides the demographic supply for population rescuing, habitat (re)colonization (Hanski & Gaggiotti 2004, Bowler & Benton 2005, Gilpin 2012), and range expansion (Travis et al. 2009, Kubisch et al. 2014, Ochocki & Miller 2017). Furthermore, it determines the intensity and direction of gene flow, which has far-reaching consequences for local genetic diversity and adaptive processes (Lenormand 2002, Ronce 2007, Petit & Broquet 2009, Cayuela et al. 2018a). Dispersal is a complex phenotype, partially controlled by genetics and relying on a suite of morphological, behavioral and life history traits that may be subject to joint selection (Saastamoinen et al. 2018). Such associations between dispersal and individual phenotype are called 'dispersal syndromes' and lead to phenotypic specialization within and between populations (Matthysen 2012, Cote et al. 2010, Ronce & Clobert 2012). Dispersal is also influenced by patch and landscape characteristics: individuals are expected to adjust their dispersal decisions according to the fitness prospects of a patch (i.e. 'informed dispersal', Clobert et al. 2009), leading to context-dependent dispersal.

Over the last two decades, an increasing number of studies have suggested that landscape anthropization is an important determinant in dispersal evolution as it affects the balance between fitness benefits and the direct and indirect costs of moving (Bonte et al. 2012) incurred at the different stages of the dispersal process (i.e. emigration, transience, and immigration) (Kokko & Lopez-Sepulcre 2006, Cheptou et al. 2017, Cote et al. 2017). The majority of efforts have been devoted to better understanding the influence of habitat fragmentation on dispersal evolution in anthropogenic landscapes (Cheptou et al. 2017, Cote et al. 2017, Legrand et al. 2017, Atkin et al. 2019). Overall, these studies have reported reduced dispersal propensity or capacity in fragmented landscapes, which is usually attributed to prohibitive costs during the transition phase across the matrix (Cheptou et al. 2017, Cote et al 2017).

This hypothesis is supported by landscape genetic studies, which often report increased spatial genetic

differentiation depending on the harshness of the matrix that separates demes (Baguette et al. 2013,

Cushman et al. 2015). Nevertheless, habitat fragmentation may have a contradictory effect on the

evolution of dispersal: on one hand, it may make transition across the matrix costlier, but on the other,

it may make dispersal profitable due to the increased local extinction risk caused by heightened

demographic stochasticity within severely fragmented landscapes (Ronce 2007, Hanski & Mononen

2011, Cote et al. 2017). The best empirical support of this hypothesis is perhaps the well-documented

selection for dispersal-specialized phenotypes observed in spatially structured populations (SSPs,

Thomas & Kunin 1999) of the Glanville fritillary butterfly subject to highly fragmented landscapes

(Hanski 2011, Hanski et al. 2017). In this unique example, environmental stochasticity makes dispersal

profitable by creating new patches that can be colonized by dispersers. The prime importance of

spatiotemporal patch variability in promoting dispersal is well supported by a number of theoretical

models (Comins et al. 1980, McPeek & Holt 1992, Armsworth & Rougharden 2005), and is also

advanced as the main driver of wing dimorphism observed in insects in a gradient of patch temporality

(Denno et al. 1996).

Habitat alteration in human-dominated landscapes is often associated with shifts in disturbance

regimes (Turner 2010, Newman 2019). For instance, one decade-long worldwide survey revealed a

relatively weak net surface loss of temperate forests, but a high turnover due to forestry practices

(Hansen et al. 2013). Shifts in the disturbance regimes prevailing in habitat remnants could therefore

mitigate or, conversely, magnify the negative effect of habitat fragmentation, depending on their

direction and magnitude. Despite this, apart for aerial dispersal in invertebrates (Denno et al. 1996),

human-induced temporal variation in the spatial distribution of habitat patches has generally been

overlooked when considering dispersal evolution in anthropogenic landscapes. A full appraisal of the

effect of anthropogenic disturbance on dispersal should not only examine whether a dispersal pattern

emerges in a landscape, but whether it gives rise to a genetic footprint throughout successive generations

– and last but not least, to what extent this involves a specialized phenotype. Considering all the facets

of this issue is not a simple task (Kokko & Lopez-Sepulcre 2006, Broquet & Petit 2009, Ronce & Clobert

2012), and to our knowledge has not yet been investigated in vertebrates.

To address this gap, this study examined how human-driven variation in habitat patch turnover

affects dispersal rates and distances, determines dispersal-related phenotypic specialization, and drives

neutral genetic variation in spatially structured populations. We studied this issue in an early

successional amphibian, the yellow-bellied toad (*Bombina variegata*), a species that reproduces in small

waterbodies with a short hydroperiod occurring either in (virtually) undisturbed or anthropogenic

environments (Warren & Büttner 2008; Cayuela et al. 2011, 2015b). In riverine environments with

negligible human activity, the species' habitat patches are groups of rocky pools that result from long-

term geomorphological processes alongside riverbanks (riverine SSPs; Cayuela et al. 2011). This results

in a negligible patch turnover rate and makes patches available and predictable far beyond a toad's

lifespan. In contrast, in harvested woodlands, habitat patches consist of groups of ruts made by logging

vehicles that may appear and disappear yearly as a result of the combined effects of logging operations

and rapid natural silting (we refer to these as logging SSPs hereafter). This leads to a high patch turnover

rate and makes patch location and availability more unpredictable at the scale of a toad's lifespan

(Cayuela et al. 2016a, 2016b). Previous studies have highlighted demographic differences in SSPs from

the two environments and found that individuals in logging SSPs have a faster life history (i.e. a shorter

lifespan and higher fecundity; Cayuela et al. 2016a), experience earlier senescence (Cayuela et al.

2019b), and display higher breeding dispersal probability (Cayuela et al. 2016b) than individuals in

riverine habitats. In this study, our first step was to quantify dispersal probability and distance

throughout an individual's lifetime, as natal dispersal was lacking in previous studies and a review of

recent literature suggested that natal and breeding dispersal patterns can strongly differ in amphibians

(Cayuela et al. 2018c). We expected that (1) both natal and breeding dispersal rates and distances would

be higher in logging than in riverine SSPs. In a second step, we analyzed how patch turnover and related

dispersal costs and benefits affected SSP dynamics and long-term viability using simulations based on

published demographic rates. We hypothesized that (2) dispersal and context-dependent immigration

(i.e. depending on patch age) allows the long-term persistence of logging SSPs. In a third step, we used

common garden experiments to investigate how patch turnover determines dispersal syndromes and

may act as a selective agent on phenotypic specialization in riverine and logging SSPs. We expected (3)

toadlets from logging SSPs to have behavioral traits (i.e. high exploration propensity and boldness) and

morphological traits (i.e. long hind limbs) that generally facilitate dispersal in amphibians (reviewed in

Cayuela et al. 2018c). In a fourth step, we examined how human-driven variation in patch turnover, by

affecting neutral genetic diversity and gene flow, leads to contrasting genetic footprints over the longer

term in riverine and logging SSPs. As genetic differentiation results from the combined effects of genetic

drift and gene flow (Broquet & Petit 2009, Cayuela et al. 2018a), we selected two SSPs per landscape

type in order to determine the relative contribution of each of these drivers. As high dispersal is expected

to increase gene flow and decrease the local effects of genetic drift, we expected (4) a larger effective

population size as well as a lower level of inbreeding and intra-patch relatedness in logging than in

riverine SSPs. We also expected that (5) after controlling for SSP size and landscape connectivity, higher

gene flow would lead to lower genetic structure and weaker genetic Isolation-By-Distance (IBD) pattern

in logging than in riverine SSPs.

Materials and methods

Study area and sampled populations

The study was conducted in 8 SSPs in eastern France, three in riverine environments (R1, R2, and R3)

and five in logging environments (L1, L2, L3, L4 and L5). The SSPs were chosen according to technical

constraints or to minimize bias at each stage of the study – our choices are explained below. The distance

separating SSPs from each other varied from 20–500 km (**Fig. S1**).

Table 1. *Environmental characteristics of the four SSPs (L1, L2, R1 and R2): patch persistence over time, environment type (logging vs riverine), patch isolation (mean distance in meters between two pond networks, and associated variation coefficient) and patch size (mean number of ponds within a patch and associated variation coefficient).*

SSP	Patch persistence	Environment	Patch isolation	Patch size
L1	1 to 10 years	logging activity	2864.50 (57%)	3.70 (70%)
L2	1 to 10 years	logging activity	4424.64 (53%)	5.64 (88%)
R1	>30 years	natural erosion	676.57 (61%)	4.52 (105%)
R2	>30 years	natural erosion	501.62 (61%)	4.71 (93%)

Table 2. *Survey design characteristics for the four SSPs (L1, L2, R1 and R2) considered in the study: study period, survey duration, number of capture sessions performed over the survey period, total number of captures, total number of individuals identified during the survey and total number of sampled patches (pond networks).*

SSP	Study period	Survey duration	Capture sessions	Number of captures	Number of individuals	Number of sampled patches
L1	2000–2008	9 years	29	953	445	28
L2	2012–2016	5 years	15	16477	12192	189
R1	2010–2014	5 years	13	4747	1003	14
R2	2010–2014	5 years	13	3984	769	8

Dispersal patterns throughout toad lifespan in riverine and logging environments

Studied populations

We quantified natal and breeding dispersal rates and distances in four SSPs (L1, L2 and R1, R2; see maps **Fig. S1** and **Fig. S2**) for which breeding rate dispersal had been previously estimated (Cayuela et al. 2016a, 2016b). The environmental characteristics of the four SSPs and the details regarding the survey design are presented in **Table 1** and **Table 2** respectively. The number of individuals captured each year is presented in **Supplementary material**, **Table S1**. A detailed description of the four SSPs can also be found in two previous studies (Cayuela et al. 2016a, 2016b). The number of patches (defined as a group of ruts) occupied by each SSP ranged from 8 to 189. Each SSP was monitored for a period of at least five years in one to five capture sessions per year that were usually between two weeks to one month apart. At each capture session, all the patches were sampled in the daytime and toads were captured by hand or dipnet. Based on previous studies (Cayuela et al. 2016a, 2016c), we considered three life stages: juveniles (i.e. post-wintering metamorphs), subadults (two-year-old immature animals)

and adults (i.e. breeders, three years old or more). This resulted in a total dataset of 12,721 individual

capture–recapture (CR) histories.

<u>Naïve dispersal kernels</u>

We first estimated a dispersal kernel based on distances recorded during sampling using a lognormal

distribution for each population and life stage (juvenile, subadult, and adult). This allowed us to visualize

the form of the kernel from raw data before building complex CR models.

<u>The structure of the multievent model</u>

For the needs of our study, we extended the CR multi-event model proposed by Lagrange et al. (2014),

which allows estimating survival (ϕ) and dispersal (ψ) in numerous sites. By omitting site identity and

distinguishing between individuals that stay and individuals that move, this model circumvents the

computational issues usually encountered in standard multi-site CR models when the number of sites is

large (Lebreton et al. 2009). Lagrange's model includes states that incorporate information about

whether an individual is occupying at t the same site as the one occupied at $t – 1$ ('S' = stayed) or not

('M' = moved). The model also includes information about whether the individual was captured ('+')

or not ('o') at $t – 1$ and t. Recently, Tournier et al. (2017) extended Lagrange's model by breaking down

dispersal (ψ) into distinct parameters of departure (ε) and arrival (α). This new parameterization allows

the estimation of the proportion of individuals arriving in sites of different quality or located at different

distances from the source site.

We adapted this parameterization for our study to consider states incorporating information

about individual capture ('+' and 'o') at $t – 1$ and t as well as movement status. We also included states

with information about the individual's age class: juvenile ('j'), subadult ('s') and adult ('a').

Additionally, we incorporated information about the Euclidian distance covered by dispersers between

the departure and arrival patch using three distance classes: '1' = 100–800 m, '2' = 800–1500 m, '3' >

1500 m. This led to the consideration of 37 states in the model (**Fig.1** and **Fig.2**). For example, an

individual +jS+ was captured at $t – 1$ and t, was a juvenile, and remained in the same patch between $t –$

1 and t. An individual +sM1+ was captured at $t – 1$ and t, was a subadult, did not occupy the same patch

as at $t – 1$, and arrived in a patch located at a distance 100–800 m from the source patch. We distinguished

16 events, which were coded in an individual's capture history and reflect the information available to

the observer at the time of capture (**Fig.2**).

When captured for the first time, the state of an individual could be ojS+, osS+ or oaS+. We

then considered five modeling steps in which the information of the state descriptor was progressively

updated: survival (ϕ), departure (ε), arrival (α), age transition (δ) and recapture (p). Each step was

conditional on all previous steps. In the first step, we updated information about survival. An individual

could survive with a probability of ϕ or die (D) with a probability of $1 - \phi$. This led to a matrix with 37

states of departure and 7 intermediate states of arrival (**Fig.1**). Survival probability could differ between

age classes by allowing differing values for ϕ in lines 1–12, 13–24 and 25–36 of the matrix. In the second

modeling step, departure was updated. Individuals could move (M) from the site they occupied with a

probability of ε or stay (S) with a probability of $1 - \varepsilon$. A matrix of 7 departure states and 13 arrival states

was considered (**Fig.2**). Departure probability could differ between age classes by allowing differing

values for ε in lines 1–2, 3–4 and 5–6 of the matrix. In the third step, we updated the arrival information.

An individual that moved could arrive in a patch located in the first two distance classes (1 or 2) from

the source patch with a probability of α, or arrive in a site located in the third distance class (3) with a

probability of $1 - \alpha$. This resulted in a matrix with 13 departure states and 25 arrival states (**Fig.2**).

Arrival probability could differ between age classes by allowing different values for α in lines 2–4, 7–8

and 11–12 of the matrix. In the fourth step, the information about age was updated. An individual could

reach the next age class (j, s or a) with a probability of δ or remain in the previous age class with a

probability of $1 - \delta$, resulting in a transition matrix with 25 states of departure and 25 states of arrival

(**Fig.2**). The adult individuals (a) were forced to stay in their age class. In the fifth and last step, recapture

was updated (**Fig.2**). An individual could be recaptured with a probability of p or missed with a

probability of $1 - p$, resulting in a transition matrix with 25 states of departure and 37 states of arrival.

The recapture probability could differ between age classes by allowing different values for p in lines 1–

8, 9–16 and 17–24 of the matrix. The last component of the model linked events to states. In this specific

situation, each state corresponded to only one possible event (**Fig.2**).

<u>Biological scenarios in the E-SURGE program</u>

The parameterization was implemented in the E-SURGE program (Choquet et al. 2009). The datasets

for the four SSPs considered in our study were analyzed separately, as the number of study years and

capture sessions for these populations varied. Competing models were ranked through a model-selection

procedure using Akaike information criteria adjusted for a small sample size (AICc) and AICc weights.

Following the recommendation of Burnham & Anderson, we performed model averaging when the

AICc weight of the best-supported model was less than 0.90. The models had a robust design structure

(Pollock 1982). As in previous studies of *B. variegata*, the survival probability was fixed at one between

secondary sessions (Cayuela et al. 2016a, 2016c). The robust design structure allowed both intra-annual

and inter-annual dispersal to be considered. Our hypotheses concerning recapture and state–state

transition probabilities were tested using the general model [ϕ(AGE), ε(AGE), α(AGE), δ(.), p(AGE +

Y)], which included two effects: (1) the three age classes (AGE) coded as states in the model, and (2)

year-specific variation (Y). The notation (.) indicated that the parameter was held constant. We tested

whether survival (ϕ) and departure probabilities (ε) varied between age class (AGE). Moreover, we

hypothesized that the probability of arriving in a patch depends on age (AGE), and on the Euclidean

distance between patches (the distance classes were incorporated as states in the model). Recapture probability was expected to differ between age classes (AGE) and years (Y). We tested our expectations about the model parameters in a stepwise fashion. From this general model, we tested all the possible combinations of effects and ran 16 competing models.

Fig.1. *Model structure: matrices of initial states and state transitions (survival, departure, arrival and age transition). In the transition matrix, the rows correspond to time t − 1, the columns to time t, and whenever a status element is updated to its situation at t, it becomes bold and stays bold throughout the following steps.*

Fig.2. *Model structure: state transitions (recapture) and events (field observations). In the transition matrix, the rows correspond to time t − 1, the columns to time t, and whenever a status element is updated to its situation at t, it becomes bold and stays bold throughout the following steps.*

278
279
280

Simulating the effect of patch turnover and dispersal on SSP dynamics and long-term persistence in
logging contexts

We simulated population trajectories based on different scenarios to investigate the effects of patch

turnover and dispersal costs on the dynamics and long-term persistence of SSPs in habitats subjected to

logging. Adopting the most realistic lifecycle for the yellow-bellied toad (see 'Results', **Fig.4**)

determined in previous studies (e.g. Cayuela et al. 2015, 2018), we used a three age-class (juveniles,

subadults, and adults), female-dominant, prebreeding Leslie matrix (Caswell 2001) (see 'Results',

Fig.4). We used the demographic parameters of a riverine SSP (R1; see Cayuela et al. 2016), which was

considered a reference population whose demographic parameters have not been altered by dispersal

costs (survival in logging SSPs is lower than in riverine SSPs, likely due to dispersal costs, Cayuela et

al. 2018b). Both prebreeding survival probability (juvenile survival, $S1 = 0.70$; subadult survival, $S2 =$

0.77) and adult survival probability ($S3 = 0.92$) were included in the Leslie matrix. Fecundity F was

possible only for adult females and consisted of estimated recruitment: that is, the number of recruited

juvenile females at t per breeding female at $t − 1$ ($F = 0.52$, Cayuela et al. 2016). As Boualit et al. (2019)

found that juvenile recruitment was higher in newly created and disturbed patches than in old,

undisturbed patches in logging SSPs, we specified that F decrease linearly (-5% per year) with the age

of the patch. Furthermore, we also considered the possibility that females may skip breeding

opportunities; in the riverine SSP R1, a previous study showed that the probability of females skipping

breeding (B) was 0.15 (Cayuela et al. 2016). As in Cayuela et al. (2018, 2019c), we considered demographic stochasticity for survival, fecundity and skipping breeding. For each year, demographic parameter values were randomly sampled in a Gaussian distribution centered on mean parameter estimates, and standard deviation was inferred from two previous studies conducted on this species (Cayuela et al. 2016a, 2018b). The standard deviation values were: 0.05 for S1, 0.03 for S2, 0.01 for S3, 0.02 for F, and 0.02 for B.

Three patch turnover scenarios were considered. In scenario 1 (high turnover), a patch disappeared three years after its creation, in scenario 2 (medium turnover) a patch disappeared after six years, and in scenario 3 (low turnover) it disappeared after nine years. These scenarios correspond to the range of patch turnover in logging SSPs reported by forest managers (Eric Bonnaire, *unpublished data*), variation that depends on local management policies and the frequency of forest harvesting operations. We also considered three dispersal scenarios. In scenario 1 (no dispersal), individuals were not able to escape and died when a patch disappeared. In scenario 2 (dispersal with random immigration), individuals could disperse to escape the disappearance of a patch or could disperse by choice (i.e. when the patch remained available). The immigration was random between the patches of the metapopulation and was not influenced by the age of the patch. In scenario 3 (dispersal with informed immigration), individuals could disperse in response to patch disapperance or by choice. Based on an assessment of Boualit et al. (2019), we considered that immigration was not random and that immigration probability linearly decreases with patch age (i.e. a loss of 5% per year). In scenario 2 and 3, in which dispersal was possible, we considered two subsets of scenarios: in subset 1 (non-costly dispersal), individuals did not incur any survival loss when they dispersed. In subset 2 (costly dispersal), we considered that survival loss related to dispersal could be low (-5% of survival), medium (-10%), or high (-15%). As in Cayuela et al. (2018), we made the assumption that survival loss was similar across life stages (i.e. juvenile, subadult, and adult).

Each simulation began with 30 breeding patches over which 1,000 individuals were randomly scattered. The number of individuals in each age class was obtained through the stable stage distribution provided by the three age-class Leslie matrix. Then for each time step (a 1-year interval), we simulated the change in patch availability. We considered that five new patches were created each year. As the patches disappeared in a deterministic way when they reached the age defined in the scenario, the number of available patches remained constant over time (except for the few first years). We simulated the number of individuals in each age class occupying each patch. To do this, we separately considered patches reaching the age of disappearance versus those that did not disappear. In the latter, the number of individuals at $t + 1$ given the number of individuals at t was predicted by the Leslie matrix using the survival probability of individuals occupying an available patch (reported in the reference population R1). To be as realistic as possible, we used demographic stochasticity (fecundity was thus randomly sampled from a Poisson distribution, and survival from a binomial distribution), as in Cayuela et al. (2018). For patches that disappeared, we applied the same procedure, but using the survival probability

(affected or not by dispersal cost, depending on the scenario) for individuals occupying patches that

subsequently disappeared. Surviving individuals from lost patches were then randomly spread over the

available patches at $t + 1$ in the 'dispersal with random immigration' scenario, or they were preferentially

distributed in new patches in the 'dispersal with informed immigration' scenario. In all dispersal

scenarios, we fixed the elective dispersal probability (dispersal when the patch did not disappear) D at

0.15, which was consistent with the annual dispersal rate reported in logging SSPs in the study. We also

considered demographic stochasticity in elective dispersal (the standard deviation value was 0.05). The

modeled population was monitored for 100 years. We did not remove the first few years of the

simulation, when the number of patches progressively increased since none were old enough to

disappear yet, as these had virtually no impact on our results (Cayuela et al. 2018). We performed 1,000

simulations for each scenario. At each time step, we monitored the number of adults in the entire SSP

as well as the proportion of simulations in which the SSP went extinct (the extinction probability).

Phenotypic specialization in riverine and logging environments

<u>Study populations</u>

To compare the morphology and behavior of toadlets in riverine SSPs with those of logging SSPs, we

used a common garden experiment. This involved collecting between 8 and 15 egg clutches (hereafter

referred to as 'family') in three SSPs in each landscape type (riverine = R1, R2, R3, and logging = L3,

L4, L5; **Fig.S1**). These were selected to minimize spatial proximity between SSPs belonging to the same

landscape type and therefore to avoid potential confounding effects resulting from spatial

autocorrelation in environmental conditions. Five siblings per family were randomly chosen just after

hatching and individually reared under controlled laboratory conditions.

<u>Rearing protocol</u>

The egg clutches were carefully transported to the laboratory, where they were individually placed in

aquariums (32×17 cm, height 15 cm) with aged tap water equipped with an oxygen pump. The

aquariums were placed in a climatic room with a light/dark cycle of 18:6, corresponding to the natural

light/dark cycle in the area in the summer, with an ambient temperature varying from 21.5 to 23.5°C.

Embryo development ended 2–5 days after the arrival of the egg clutches at the laboratory. After

hatchling, when tadpoles reached Gosner stage 24–25 (active swimming, external gill atrophy) (Gosner

1960), five siblings per family were randomly chosen and placed individually into plastic containers

(14×8.5 cm, height 13.5 cm) filled with previously aged tap water and aerated in a tank. The plastic

containers were distributed in a predetermined random pattern around the climatically controlled room.

The water was replaced every three days. The tadpoles were fed every day with 150 mg of cooked

lettuce, providing ad libitum feeding. At Gosner stage 44–45 (tail atrophy, mouth posterior to eyes), feeding was terminated, and the water was drained and replaced with a dampened sponge placed in the bottom of the container. The sponge and the walls were sprayed with aged tap water every two days. The individuals were kept until their complete metamorphosis (Gosner stage 46) and were then subjected to behavioral assays.

Experimental arenas

The behavioral assays took place in an arena (70 cm in diameter) made of polyethylene terephthalate, with a central shelter and a 'desiccation' obstacle between this and the 12 possible exits (**Fig. S4**). At the center of the arena, we placed a removable, cylindrical (9 cm diameter, height 90 cm), opaque, covered chamber ('refuge chamber' hereafter). The cylinder had a circular opening (3 cm diameter) covered by a lid. Around the interior edge of the arena, we installed a pit (width 10 cm, depth 0.3 cm) filled with a desiccating mixture of sand and highly active silica gel powder in a weight ratio of 0.8:0.2 ('desiccant zone' hereafter). The arena wall included 12 doors placed at regular intervals around the entire circumference. The arena was confined in an enclosed iron chamber (150×17 cm, height 175 cm) over which a dark opaque sheet was placed to limit potential acoustic and visual interference during behavioral trials.

Behavioral assays

At metamorphosis, each toadlet was subjected to a behavioral assay to quantify their neophobia or exploratory behavior. The behavioral tests were conducted in the circular arena described above. Before each trial, the toadlet and a dampened sponge (already present in the toadlet's rearing container) were gently transferred into an opaque circular release box that was then placed at the center of the arena. The dampened sponge was considered a known object, making the refuge chamber more familiar than the rest of the experimental device. Following previous studies on anuran behavioral syndromes (reviewed in Kelleher et al. 2018), neophobia was quantified as the latency time to enter a novel environment (i.e. the time delay to leave the familiar refuge chamber: BEHAV1). Exploration propensity was assessed using two variables: the latency time to enter a novel but harsh environment (i.e. to reach the desiccant zone after leaving the refuge chamber; BEHAV2), and the latency time to travel the harsh environment and get out of the arena (BEHAV3). The behavior was recorded for 30 min (1800 s) using a digital camera (Sony DCR-SX34).

Extraction of behavioral variables from the videos

The videos were analysed using the BEMOVI R package (Pennekamp et al. 2015) to reconstruct the movement trajectory of the toadlet in the arena, and to extract a series of behavioral variables from this. First, the videos were standardized to a length of 29 min (the minimum duration available for all individuals) removing the initial 60 s and the extra time at the end, if any. Then the videos were converted to a format suitable for analysis in BEMOVI: rectangular pixels (720 x 576) were converted into square pixels (1024 x 576), color information was converted into 256 gray levels, frames per s were decreased from 25 to 5 to limit memory allocation requirements, and the videos were saved as AVI files. These operations were performed using FFMPEG software (ffmpeg.org).

BEMOVI was then run with the following parameters: black and white threshold (40) to discriminate the toadlet from the background; minimum size (20) and maximum size (150), corresponding to the size range of the toadlets; link range (7500 frames) to allow any duration of the 'disappearance' of the toadlet from the video (e.g. when it was in the refuge) while still considering it as a single movement trajectory, and "disp "(100 pixels); in BEMOVI, disp is the maximal distance covered by the toadlet from one frame to the next, corresponding to 0.2s here. This resulted in a database with the movement trajectory of each toadlet, i.e. its X–Y position at each time step. This position was compared to the distance from the center of the arena to determine which zone (refuge chamber, normal zone or desiccant zone) the toadlet was in at each time step. The results were checked for errors in toadlet positioning, which were due to varying light conditions, usually at the beginning of the videos. In a final step, we computed several behavioral variables for each toadlet from this positioning information.

Statistical analyses

We used linear mixed models to test whether morphological traits (body size, body condition and relative hind limb size) of toadlets differed according to the landscape type of origin (logging vs riverine). Each morphological trait was treated as a dependent variable, and the landscape type was introduced as a fixed explanatory term in the model. In the case of body condition and relative leg size, body size and its interactive effect with the landscape type were also introduced as adjustment covariates in the fixed part of the model. Both the SSP of origin and the clutch were introduced as random effects in the model. We also allowed heterogeneity of variance between landscape types by allowing a separate estimation of the residual variance for each landscape type. The estimation method was based on restricted maximum likelihood. Variance heterogeneity was first checked using the likelihood ratio test and removed if non-significant. The significance of each explanatory term was examined with a non-sequential F test based on the Kenward-Roger method to approximate the denominator df (Littel et al. 2006). In the case of covariance analyses (body condition and relative leg size), the interactive effect was discarded if non-significant to obtain the final model. All morphological variables were standardized using Z transformation before the analyses, as recommended by Schielzeth (2010).

We tested whether the behavioral variables BEHAV1 and BEHAV2 varied according to the
landscape type (i.e. logging vs riverine) using a generalized linear mixed model. Each latency time
variable was treated as a dependent variable using a Poisson distribution. The landscape type was
introduced as an explanatory term in the fixed part of the model. We also introduced individual body
size and its interactive effect with the landscape of origin as adjustment covariates in the fixed part of
the model since both locomotion skill and exploratory performance can vary according to individual
size (reviewed in Kelleher et al. 2018). For the analyses of morphological traits, both the SSP of origin
and the clutch were introduced in the model as random effects. Furthermore, a scale parameter was also
introduced to handle data overdispersion and to obtain a corrected statistical test using a quasi-likelihood
approach (McCullagh & Nelder 1989). The estimation method was based on restricted pseudo-
likelihood optimization, and the significance of each explanatory term was examined using the same
methodology as for the morphological analyses. Non-significant terms were successively removed to
obtain the final model, and least square means were used to estimate the difference in latency time
variables according to the landscape of origin.
As the third behavioral variable (BEHAV3) was right censored, it was analyzed using a
proportional hazards mixed-effects model (i.e. a frailty model based on a Cox model, PHREG procedure,
SAS Institute Inc. 2012). We tested whether newborn individuals originating from logging systems were
more prone to exit the assay arena than those from riverine systems. It is not possible to handle multiple
random factors in such a model, so we took into account only the family effect since this was found to
be significant for the other behavioral variables but not for the SSP of origin. The landscape of origin,
the body size and their interactive effect were introduced as explanatory terms. Parameters were
estimated using partial likelihood estimation, and the significance of explanatory terms was assessed
using non-sequential chi-square tests.
470
471   *Neutral genetic variation in riverine and logging environments*
472
473   Study populations
474
We examined neutral genetic variations within two SSPs in riverine environments (R1 and R2) and two
SSPs in logging environments (L3 and L4) (see map in **Fig. S3**) using 15 polymorphic microsatellite
markers (described and tested in Cayuela et al. 2017a). The four SSPs were selected according to the
following criteria: (1) SSPs embedded in a relatively continuous forested matrix to avoid any
confounding effect of matrix composition on gene flow (**Fig. S3**) – woodland is generally considered
highly favorable for the movement of forest amphibians (Cushman 2006a) such as *B. variegata* (Cayuela
et al. 2015b); and (2) two small (R1 and L3) and two large SSPs (R2 and L4) to control for genetic drift.
The number of patches and DNA sampled per SSP are given in **Table S2**. We used the protocol

described in Cayuela et al. (2017) for DNA extraction and amplification, individual genotyping, and
allele scoring.

Estimating basic genetic metrics

We examined basic assumptions (i.e. detection of null alleles, Hardy–Weinberg equilibrium) and
descriptive statistics (mean number of alleles per locus N_a; effective number of alleles per locus N_{ef};
observed heterozygosity H_o; and expected heterozygosity H_e) of the genetic diversity within each SSP.
N_a, N_{ef}, H_o and H_e were calculated using GENALEX (Peakall & Smouse 2006). The inbreeding
coefficient F_{is} and the Hardy–Weinberg deviation were assessed with GENEPOP 4.1 (Rousset 2008).
The detection of null alleles was performed using the program MICRO-CHECKER 2.2.3 (Van
Oosterhout et al. 2004). Two microsatellite markers, Bomvar_Cons470 and bv11.7, were discarded from
the subsequent analyses due to the presence of null alleles in the four SSPs. All descriptive statistics are
provided in **Table S3**.

Estimating relatedness, inbreeding, and effective population size

We estimated relatedness and individual inbreeding using COANCESTRY v1.0.1.8 (Wang 2011). We
performed simulations to identify the best relatedness and inbreeding estimator for our combined
datasets; these consisted of 1,000 dyads spread equally across six categories of relatedness: parent–
offspring (rxy = 0.5), full siblings (rxy = 0.5), half siblings/avuncular/grandparent–grandchild (rxy =
0.25), first cousins (rxy = 0.125), second cousins (rxy = 0.03125), and unrelated (rxy = 0). According to
our simulations, the best rxy estimator for relatedness analysis was DyadML, which showed a strong
correlation of 0.74. We used linear mixed models to evaluate how SSP size (small: R1 and L4; large:
R2 and L3) and the type of environment (logging vs riverine) affected individual inbreeding and intra-
patch relatedness. An SSP's size and environment type were coded as fixed effects (we considered an
interaction between the two factors), whereas patch identity was coded as a random effect. We then
estimated the effective population size of the four SSPs using the linkage disequilibrium method
implemented in Ne Estimator v2.1 (Do et al. 2014).

Clustering approach

We described the genetic differentiation between patches within each SSP using an assignation method
based on the Bayesian clustering algorithm implemented in the software STRUCTURE (Pritchard et al.
2000). Specifically, we estimated the most likely number of genetic clusters (*K*) contained in each SSP
following the hierarchical approach proposed by Balkenhol et al. (2014) to detect additional
substructures within clusters. The STRUCTURE program was run with the admixture model, with a

burn-in period of 100,000 repetitions, and 100,000 subsequent MCMC repetitions. The K values were

tested ranging from 1 to 10 and analyses repeated 10 times for each value. We used STRUCTURE

HARVESTER (Earl 2012) to summarize the results, determining the optimal K value using both log-

likelihood plots and the delta K statistic (Evanno et al. 2005). We followed the hierarchical approach

proposed by Coulon et al. (2008) to test for additional population substructures within clusters.

Accordingly, these analyses were then repeated for each inferred population cluster separately until the

optimal K value was 1 (meaning that no additional structure was found within clusters). To map the

spatial distribution of the different clusters, the individual ancestry values were averaged across the ten

STRUCTURE runs using CLUMMP software (Jakobsson & Rosenberg 2007). The 'greedy' algorithm

in the CLUMMP program was then used to assign the individuals to the cluster in which they showed

the highest Q-values.

<u>Gradient analyses</u>

We used direct gradient analyses (Prunier et al. 2015) to test whether spatial genetic differentiation was

lower within logging than riverine SSPs while controlling for potential differences in the functional

connectivity prevailing within each SSP. To do this, pairwise genetic distances between all individuals

from each SSP were computed using the Bray-Curtis percentage dissimilarity measures (Cushman et al.

2006b, Legendre & Legendre 2012), and were then standardized separately for each SSP. Since by

design all SSPs were embedded in a relatively homogeneous forested matrix, we did not control for the

effect of land cover type in these analyses. Rather, we focused on the effect of topographic roughness

('slope' effect) and the hydrological network topology ('network' effect) since both these landscape

features have often been reported to affect gene flow in amphibians (Lowe 2003, Grant et al. 2010).

For each SSP, resistance layers were produced using ArcGIS Pro. Minimum geographical

bounding of each SSP was determined and extended to a distance of five kilometres to avoid any edge

effects. The elevation raster and hydrographic network shapefile, available through the French National

Institute for Geographical and Forest Information database (BD ALTI®, and BD TOPO®), were

extracted for SSP surface areas. To determine the percentage of steepness raster maps, the slope tool

was used. Due to the spatial resolution imposed by elevation data, all layers were converted to raster

format and homogenized at a spatial resolution of five meters.

Four resistance maps were thus constructed for each SSP. The first map included only the effect

of geographic distance (i.e. isolation by distance, hereafter geographic resistance); it was based on data

extended to include a 5-km buffer and was assigned a uniform resistance of 1 unit for all pixels. The

second map included the topographic roughness (slope), where the resistance of each pixel was a linear

function of the steepness (resistance $= 1 +$ degree of steepness), thus corresponding to the effect of

both geographic distance and topographic roughness. The third map included the hydrological network

topology (network), where the resistance was assigned to 1 unit for all pixels situated in the hydrological
network and to 10 units otherwise. The fourth map included both the slope effect and the network effect.
Pairwise resistance between all patches was then computed from these four resistance maps using the
circuit theory in Circuitscape V4.0 (McRae & Beier 2007).

Linear mixed models for pairwise distance matrices were then used to assess the effect of the
geographic distance between patches (i.e. the geographic resistance) on the genetic distance between
toads within each SSP while controlling for the effect of topographic roughness and/or the hydrological
network topology. The non-independence between pairwise distances was taken into account in the
covariance structure of the models. Specifically, we used the method proposed by Clarke et al. (2002)
using a Toeplitz(1) as a covariance structure to specify the non-independence of pairwise genetic
distance according to the patches of origin (see Selkoe et al. 2010, van Strien et al. 2012 for application
on pairwise Fst distances). We used the extension presented in Prunier et al. (2013) for application on
individual genetic distance in spatially hierarchized sampling schemes as is the case in our study. This
resulted in two covariance parameters for each SSP, one for the patch random effect and the other for
the individual random effect. Since both slope and network resistance were highly correlated to
geographic resistance and as our main aim was to estimate the effect of geographic resistance on genetic
distance while controlling for the effect of functional connectivity, we first regressed each of these
effective resistances on each geographic resistance using simple linear regressions to obtain uncorrelated
effective resistance respectively related to the slope effect, the network effect and their combined effect.
The relative validity of each alternative landscape hypothesis (i.e. including the effect of the slope or of
the stream network or both or neither on the genetic distances) was evaluated using weighted AIC
(Waic), and the model-averaged estimate of the beta weight associated with the effect of geographic
resistance on genetic distance was computed for each SSP.

Results

Dispersal patterns throughout toad lifespan in riverine and logging environments

Model averaging estimates (see **Table S4** for survival and recapture probabilities) indicated that both
natal and breeding dispersal rates were high in logging environments (around 20% per year, **Fig.3**). In
contrast, in riverine environments, natal dispersal rates were null and breeding dispersal rates were very
low (less than 5% per year, **Fig.3**). This higher dispersal propensity in logging than in riverine habitats
is even more remarkable given that the mean inter-patch distance is three times farther in the former
than the latter (Table S1). The few adults that did disperse in riverine environments covered shorter
distances: the median distance was 168 m in R1 and 189 m in R2, while the maximal distance was 455
m and 378 m respectively. This result was further confirmed by multi-event CR models showing that

100% of dispersal occurred over distances ranging from 100 to 800 m in riverine SSPs. In logging
environments, dispersers covered substantially longer distances: the median distance was 431 m in L1
and 568 m in L2, while the maximal distance was 3810 m and 4529 m respectively. This result was
supported by both the observed dispersal kernel and the multi-event model estimates for each life stage
(see **Fig.3**). While these results indicate extremely contrasting dispersal regimes between logging and
riverine landscapes, we also found substantial variations in the dispersal patterns in the two logging
systems. First, while the inter-annual dispersal rate was similar in both logging systems, the intra-annual
dispersal rate was substantially higher in L1 than L2 (i.e. L1 > 10 % and L2 <5%, **Fig.3**; for model
selection procedure see **Table 3** and **Table 4**). Second, the dispersal kernels show a clear leptokurtic
distribution decreasing with age in L2, suggesting a large demographic weight of natal dispersal in this
SSP. This was not the case in L1, in which leptokurtosis was more reduced in juveniles than in adults.
603

604
605
606 **Fig.3.** *Natal and breeding dispersal rates and distances in logging (L1 and L2) and riverine SSPs (R1*
607 *and R2) (A–B). Inter-annual (A) and intra-annual (B) dispersal probabilities are higher in SSPs in*
608 *logging landscapes than in riverine SSPs, regardless of the ontogenetic stage (juvenile, J; subadult, S;*
609 *adult, A). Natal and breeding dispersal distances in logging SSPs (L1 and L2) (C–I). Dispersal event*
610 *frequency decreases with the Euclidean distance between breeding patches at the ontogenetic stages*.
611

r	Model	k	Dev.	wAICc	AICc
SSP L1					
1	ϕ(A), ε(.), α(A), p(A + Y)	26	5161.06	0.81	5214.05
2	ϕ(A), ε(A), α(A), p(A + Y)	28	5160.05	0.17	5217.19
3	ϕ(A), ε(.), α(A), p(Y)	24	5173.01	0.02	5221.86
4	ϕ(A), ε(A), α(A), p(Y)	26	5172.00	0.00	5224.99
5	ϕ(.), ε(.), α(A), p(A + Y)	24	5188.40	0.00	5237.25
6	ϕ(.), ε(A), α(A), p(A + Y)	26	5187.39	0.00	5240.38
7	ϕ(.), ε(.), α(A), p(Y)	22	5199.87	0.00	5254.58
8	ϕ(.), ε(A), α(A), p(Y)	24	5198.86	0.00	5247.70
9	ϕ(A), ε(.), α(A), p(A)	18	5261.02	0.00	5297.50
10	ϕ(A), ε(A), α(A), p(A)	20	5260.01	0.00	5300.60
11	ϕ(A), ε(.), α(A), p(.)	16	5268.73	0.00	5301.11
12	ϕ(A), ε(A), α(A), p(.)	18	5267.72	0.00	5304.20
13	ϕ(.), ε(.), α(A), p(A)	16	5289.52	0.00	5321.90
14	ϕ(.), ε(.), α(A), p(A)	14	5295.71	0.00	5324.01
15	ϕ(.), ε(A), α(A), p(A)	18	5288.51	0.00	5324.99
16	ϕ(.), ε(A), α(A), p(.)	16	5294.70	0.00	5327.08
SSP L2					
1	ϕ(A), ε(A), α(A), p(A + Y)	25	62699.27	0.74	62749.34
2	ϕ(A), ε(.), α(A), p(A + Y)	23	62705.41	0.23	62751.47
3	ϕ(A), ε(A), α(A), p(Y)	23	62738.78	0.02	62784.85
4	ϕ(A), ε(.), α(A), p(Y)	21	62744.92	0.00	62786.98
5	ϕ(.), ε(A), α(A), p(A + Y)	23	62779.96	0.00	62826.03
6	ϕ(.), ε(.), α(A), p(A + Y)	21	62786.10	0.00	62828.16
7	ϕ(.), ε(A), α(A), p(Y)	21	62852.52	0.00	62894.58
8	ϕ(.), ε(.), α(A), p(Y)	19	62858.66	0.00	62896.71
9	ϕ(A), ε(A), α(A), p(A)	21	63103.58	0.00	63145.64
10	ϕ(A), ε(.), α(A), p(A)	19	63109.72	0.00	63147.77
11	ϕ(A), ε(A), α(A), p(.)	19	63144.67	0.00	63182.72
12	ϕ(A), ε(.), α(A), p(.)	17	63150.81	0.00	63184.85
13	ϕ(.), ε(A), α(A), p(A)	19	63185.43	0.00	63223.48
14	ϕ(.), ε(.), α(A), p(A)	17	63191.57	0.00	63225.61
15	ϕ(.), ε(A), α(A), p(.)	17	63272.39	0.00	63306.42
16	ϕ(.), ε(.), α(A), p(.)	15	63278.53	0.00	63308.55

618 ***Table 4**. Multievent models and selection procedure for SSPs R1 and R2. r = model rank, k = number*
619 *of parameters, Dev. = residual deviance, AICc = Akaike information criterion adjusted for small sample*
620 *size, wAICc = AICc weight ϕ = survival probability, ε = departure probability, α = arrival probability,*
621 *p = recapture probability, A = age, . = constant, Y = year.*

r	Model	k	Dev.	wAICc	AICc
SSP R1					
1	$\phi(A), \varepsilon(A), \alpha(A), p(Y)$	19	8044.47	0.47	8082.67
2	$\phi(A), \varepsilon(A), \alpha(A), p(A + Y)$	21	8041.98	0.22	8084.22
3	$\phi(A), \varepsilon(.), \alpha(A), p(Y)$	17	8050.14	0.21	8084.29
4	$\phi(A), \varepsilon(.), \alpha(A), p(A + Y)$	19	8047.65	0.10	8085.84
5	$\phi(.), \varepsilon(A), \alpha(A), p(Y)$	17	8074.54	0.00	8108.70
6	$\phi(.), \varepsilon(.), \alpha(A), p(Y)$	15	8080.20	0.00	8110.32
7	$\phi(.), \varepsilon(A), \alpha(A), p(A + Y)$	19	8073.47	0.00	8111.67
8	$\phi(.), \varepsilon(.), \alpha(A), p(A + Y)$	17	8079.13	0.00	8113.29
9	$\phi(A), \varepsilon(A), \alpha(A), p(.)$	15	8129.11	0.00	8159.23
10	$\phi(A), \varepsilon(A), \alpha(A), p(A)$	17	8125.73	0.00	8159.89
11	$\phi(A), \varepsilon(.), \alpha(A), p(.)$	13	8134.77	0.00	8160.87
12	$\phi(A), \varepsilon(.), \alpha(A), p(A)$	15	8131.39	0.00	8161.52
13	$\phi(.), \varepsilon(A), \alpha(A), p(.)$	13	8157.05	0.00	8183.14
14	$\phi(.), \varepsilon(.), \alpha(A), p(.)$	11	8162.71	0.00	8184.78
15	$\phi(.), \varepsilon(A), \alpha(A), p(A)$	15	8155.63	0.00	8185.75
16	$\phi(.), \varepsilon(.), \alpha(A), p(A)$	13	8161.29	0.00	8187.38
SSP R2					
1	$\phi(A), \varepsilon(A), \alpha(A), p(A + Y)$	21	11041.21	0.64	11083.41
2	$\phi(A), \varepsilon(.), \alpha(A), p(A + Y)$	19	11047.74	0.19	11085.90
3	$\phi(A), \varepsilon(A), \alpha(A), p(A)$	17	11052.43	0.13	11086.56
4	$\phi(A), \varepsilon(.), \alpha(A), p(A)$	15	11058.96	0.04	11089.06
5	$\phi(.), \varepsilon(A), \alpha(A), p(A + Y)$	19	11068.13	0.00	11106.29
6	$\phi(.), \varepsilon(.), \alpha(A), p(A + Y)$	17	11074.66	0.00	11108.78
7	$\phi(.), \varepsilon(A), \alpha(A), p(A)$	15	11078.84	0.00	11108.94
8	$\phi(.), \varepsilon(.), \alpha(A), p(A)$	13	11085.36	0.00	11111.44
9	$\phi(A), \varepsilon(A), \alpha(A), p(Y)$	19	11076.17	0.00	11114.33
10	$\phi(A), \varepsilon(.), \alpha(A), p(Y)$	17	11082.69	0.00	11116.82
11	$\phi(A), \varepsilon(A), \alpha(A), p(.)$	15	11087.00	0.00	11117.10
12	$\phi(A), \varepsilon(.), \alpha(A), p(.)$	13	11093.52	0.00	11119.60
13	$\phi(.), \varepsilon(A), \alpha(A), p(Y)$	17	11111.72	0.00	11145.85
14	$\phi(.), \varepsilon(.), \alpha(A), p(Y)$	15	11118.24	0.00	11148.34
15	$\phi(.), \varepsilon(A), \alpha(A), p(.)$	13	11122.54	0.00	11148.62
16	$\phi(.), \varepsilon(.), \alpha(A), p(.)$	11	11129.07	0.00	11151.12

625
626
Fig.4*. Effect of patch turnover on SSP size (number of adults) based on simulations. We considered*
three turnover rates: high (patch age before disappearance: 3 years, A and D), medium (6 years, B and
E), and low (9 years, C and F). Three dispersal scenarios were also considered: (1) no dispersal
(individuals were not able to escape when a patch disappeared and thus died); (2) dispersal with random
immigration (individuals could disperse to escape the disappearance of a patch or disperse by choice
even if the patch remained available. Immigration was random between the patches in the
metapopulation and was not influenced by the age of the patch); (3) dispersal with informed immigration
(similar to scenario 2, but individuals preferentially immigrated to recently created patches where
fecundity F was highest). In scenarios 2 and 3, we considered two possible options regarding dispersal
cost: (i) individuals did not incur any survival loss during dispersal (i.e. 'non-costly dispersal', A, B,
and C); (ii) survival loss related to dispersal was either low (5% survival cost), medium (10%), or high
(15%) (i.e. 'costly dispersal', D, E, and F). To investigate the demographic consequences of these
scenarios, we used a female-dominant Leslie matrix (G) based on four demographic parameters: F =
the female achieved fecundity, S_1 = juvenile survival, S_2 = subadult survival, and S_3 = adult survival
with both environmental and demographic stochasticity.

Simulating the effect of patch turnover and dispersal on logging SSP dynamics and long-term
persistence

The results showed that patch turnover rate was a critical driver of logging SSP dynamics. The absence
of dispersal within an SSP experiencing patch turnover necessarily led to the extinction of the SSP
(**Fig.4A**, **4B**, and **4C**), and the extinction speed increased with patch turnover rate. In contrast, patch
turnover had a positive effect on SSP size (i.e. number of adults) when dispersal was possible and had

no survival cost. This was caused by a fecundity-related mechanism: high patch turnover led to a decrease in the mean patch age within the SSP, resulting in an increase in average fecundity due to the positive relationship between fecundity and patch age.

Our simulations also showed that an SSP's extinction risk increased with the survival cost related to dispersal and was mitigated by 'informed immigration' (**Fig.4D**, **4E**, and **4F**). In the scenarios with a high patch turnover rate (disappearance of patch after 3 years of availability), SSPs inevitably went extinct when the survival loss was higher than 10%. Random immigration accelerated the SSP's decline compared to informed immigration directed toward recently created patches (where female fecundity was highest). In the scenarios with medium patch turnover (disappearance after 6 years of availability), the SSP decreased when the survival loss was 15% with both random and informed immigration. With a 10% survival loss, informed dispersal mitigated the decline of an SSP, whereas random dispersal drove the SSP to extinction. In the scenarios with low patch turnover (disappearance after 9 years of availability), the SSP experienced a marked decline only when immigration was random and the survival loss was equal to or higher than 10%. In summary, the simulations showed that patch turnover may result in SSP decline when the survival cost of dispersal is relatively high (10–15%) and immigration decisions are not adjusted to patch age and related fitness prospects.

Phenotypic specialization in riverine and logging environments

We focused on three morphological characteristics known to condition movement capacity in amphibians: body size, body condition and relative leg size (Gomes et al. 2009, Hillman et al. 2014). Of the 400 tadpoles, 295 survived until metamorphosis, resulting in a relatively high metamorphosis success rate ($mean \pm sd = 0.75 \pm 0.08$), which did not differ according to the landscape type of origin (n = 400, $F_{1,3.8} = 0, P = 0.95$). Neither body size nor body condition at metamorphosis varied according to the landscape type (n = 295; body size: $F_{1,4} = 0, P = 0.99$; body condition: $F_{1,3.3} = 1.12, P = 0.36$). However, the allometric relationship between leg size and body size varied according to landscape type (n = 295; body size: $F_{1,227} = 776.91$, P < 0.0001; landscape type: $F_{1,3.4} = 3.74$, P = 0.13; body size × landscape type: $F_{1,227} = 6.43$, P = 0.012, Fig. 5). The larger the toadlets, the more those originating from logging SSPs tended to have longer hind limbs than those originating from riverine SSPs. Yet this difference was only just significant for large animals, as indicated by sliced tests (respectively performed at the 1st , 2nd and 3rd quartile of the size distribution: $F_{1,3.8} = 1.64$, P = 0.27; $F_{1,3.4} = 3.28$, P=0.15; $F_{1,4.01} = 6.54$; P = 0.06).

Based on the behavioral trials in the experimental arenas, we measured variables along the boldness–shyness behavioral axis, a personality trait consistently involved in dispersal syndromes across different organisms (Cote et al. 2010). Toadlet behavior was characterized using three measures: one to assess neophobia (BEHAV1 = the time taken to leave the familiar refuge chamber) and the other two to

assess exploration propensity (BEHAV2 = the time taken to reach the desiccant zone after leaving the

refuge chamber; BEHAV3 = the time taken to cross the desiccant zone and get out of the arena). The

findings showed that the neophobia of toadlets significantly varied according to their landscape of origin

(**Fig.5**), but not according to their body size or its interactive effect with landscape (N=295; landscape

type: $F_{1,3.5}$=47.30, P=0.004; body size: $F_{1,13.9}$=0.03, P=0.86; their interactive effect: $F_{1,12.1}$=0.01, P=0.92).

Individuals from riverine landscapes were 2.13+-0.24 times slower to leave the refuge chamber than

those from logging landscapes (respectively 8.58+-0.67 mn and 3.89+-0.33 mn, see **Fig.5**). Once the

toadlet left the refuge chamber, the latency time to reach the desiccating zone was very short (mean 1.83

+- 0.21 mn) and did not vary depending on the landscape of origin, body size, or their interactive effect

(N=295; landscape type: $F_{1,3.9}$ =1.84, P=0.25; body size: $F_{1,30}$=0.42, P=0.52; their interactive effect

$F_{1,33.4}$=0.26, P=0.61, see **Fig.5**). In contrast, the latency time to get out of the arena significantly varied

depending on the landscape of origin, but not according to body size (N=295; landscape type: χ^2_{1df}=9.56,

P =.002; body size: χ^2_{1df}=0.01, P=0.91; their interactive effect: χ^2_{1df}=0.28, P=0.59). The time taken –

and thus the increased hazard – to get out of the arena for toadlets originating from logging SSPs was

1.93 times greater than for toadlets from riverine SSPs.

Fig.5. Behavioral and morphological specialization in six Bombina variegata SSPs occurring in logging

environments (L3, L4, and L5) and riverine environments (R1, R2, and R3). We examined how the type

of environment affected two behavioral traits: the time to exit the refuge chamber (A), a proxy for

neophobia, and the time to reach the desiccation zone (B), a proxy for exploration propensity. We also

 examined how environment type affected hind limb length while considering body length as a control
 covariate in the model. These observations were recorded for 295 toadlets at metamorphosis.

 Fig.6. Inbreeding in the riverine (blue) and logging (green) *Bombina variegata* SSPs.

 Fig.7. Relatedness within patches in the riverine (blue) and logging (green) *Bombina variegata* SSPs.
 Patches where less than 6 relatedness values were available were removed from the plot.

 Neutral genetic variation in riverine and logging environments

 We investigated the neutral genetic footprint associated with each landscape type using microsatellite
 data collected in four SSPs: two from each landscape type (a total of 667 toads genotyped within L3 and
 L4 logging environments and R1 and R2 riverine environments; **Table S2**). Our analyses revealed, first,
 that riverine systems exhibited a higher deviation from the Hardy–Weinberg equilibrium than logging
 systems. Significant deviation from this equilibrium was detected for 75% and 45% of the loci in riverine
 SSPs R1 and R2 respectively, and 0% and 18% in the logging SSPs L3 and L4 (**Table S3**). Second, we

found lower genetic diversity in riverine SSPs than in logging SSPs: both the allelic richness and the

expected heterozygosity (uHE) were substantially lower in riverine SSPs. In contrast, inbreeding

coefficients (F_{IS}) were higher in riverine SSPs: 87% of the loci were found to have a lower uHE in

riverine SSPs compared to logging SSPs (**Table S3**). Similarly, our parentage analyses also showed that

individual inbreeding (LR test, $\chi^2 = 32.55$, P <0.0001; **Fig.6**) and the relatedness level within patches

(LR test, $\chi^2 = 43.14$, P <0.0001, **Fig.7**) were drastically higher in riverine SSPs. In addition, riverine

SSPs had a much smaller effective population size than logging SSPs: the effective population size was

9.7 (95% CI 6.4–13.7) in R1 and 22.6 (95% 19.3–26.3) in R2, while it was 138.9 (95% 80.2–353.6) in

L4 and 168.6 (95% 129.9–229.3) in L3.

Within both types of environment, the size of SSPs (large: R2 and L3; small: R1 and L4) was

also an important predictor of individual relatedness and inbreeding. The relatedness level within

patches was higher (**Fig.7**) in small SSPs than in large SSPs (LR test, $\chi^2 = 214.38$, P <0.0001), and this

difference was larger in riverine SSPs (LR test, $\chi^2 = 6.16$, P = 0.01). Similarly, inbreeding within patches

was higher (**Fig.6**) in small SSPs than in large SSPs (LR test, $\chi^2 = 76.56$, P <0.0001), and the interaction

between these two factors was also confirmed (LR test, $\chi^2 = 8.88$, P = 0.01).

Fig.8. *Patterns of neutral genetic variation and spatial distribution of the hierarchical genetic clusters*

in two riverine (R1 and R2) and logging SSPs (L3 and L4). The analyses were conducted using the

program STRUCTURE. In L3 and L4, no genetic structuring was detected. In contrast, a complex

genetic structure was found in R1 and R2. In R1, the hierarchical analysis revealed the existence of two

initial genetic clusters (A and B), and genetic substructures within cluster A where two nested clusters

were inferred (AA and AB). In R2, the hierarchical analysis highlighted the presence of two initial

genetic clusters (A and B), and a substructure was further detected within cluster B (BA and BB). Genetic substructures were then identified within the cluster BA, with three additional genetic clusters inferred (BAA, BAB, and BAC).

Table 5. *Relative support of the mixed models for each of the four SSPs and their related estimates of geographic resistance (L3 and L4 = logging SSPs, R1 and R2 = riverine SSPs, see Map S1 for details). 'Model' indicates the effect introduced in the model in addition to geographic resistance (null = the model including only the effect of geographic resistance). 'Waic' indicates the weighted AIC of the model.* $\hat{\beta}(sd)$ *indicates the beta weight associated with the effect of geographic resistance and its standard deviation. The variable 'slope' corresponds to the effect of both geographic distance and topographic roughness. The variable 'network' corresponds to the effect of the hydrological network.*

Model	Waic	$\hat{\beta}(sd)$
SSP L3		
null	0.469	1.4763 (0.1544)
slope	0.245	1.4811 (0.1545)
network	0.191	1.4798 (0.1545)
slope+network	0.095	1.4826 (0.1546)
SSP L4		
null	0.169	0.6783 (0.0415)
slope	0.084	0.679 (0.0415)
network	0.507	0.6764 (0.0415)
slope+network	0.240	0.677 (0.0415)
SSP R1		
null	0.391	2.381 (0.0713)
slope	0.276	2.386 (0.0714)
network	0.226	2.3845 (0.0714)
slope+network	0.107	2.3862 (0.0714)
SSP R2		
null	0.070	2.5459 (0.0251)
slope	0.030	2.5461 (0.0251)
network	0.601	2.5479 (0.0251)
slope+network	0.299	2.5482 (0.0251)

We then investigated how patterns of genetic differentiation between patches differed between logging and riverine SSPs. First, we used the Bayesian genetic clustering approach to examine hierarchical genetic structure in the SSPs. In the two SSPs from logging environments, we failed to detect any hierarchical genetic structure (K = 1) (**Fig.8**). In contrast, our analyses revealed a strong hierarchical genetic structure overlaying the spatial distribution of patches in the two SSPs from riverine environments (**Fig.8**). Second, we sought to verify that differences in spatial genetic patterns between landscapes were still significant while controlling for the potential difference of functional connectivity between SSPs using direct gradient analyses. These revealed that genetic differentiation correlated with geographic resistance between individuals whatever the landscape type, indicating substantial genetic isolation by distance even in logging systems (**Fig.9**). There was considerable support for the effect of the hydrological network on genetic differentiation in the large SSPs of both landscape types (SSPs R2 and L3), but not in the small SSPs whatever the landscape type (**Table 5**). Most importantly, even after

correcting for these potential landscape effects, spatial genetic differentiation remained higher in riverine than in logging SSPs as revealed by the model averaged slope estimate associated with the geographic resistance effect ($mean \pm sd$: 2.38 ± 0.07 for R1 and 2.55 ± 0.03 for R2; 1.48 ± 0.15 for L3 and 0.68 ± 0.04 for L4).

Fig.9*. Genetic divergence according to geographic distance in two riverine (R1 and R2) and two logging SSPs (L3 and L4). Each figure represents the contour plot of the kernel density bivariate estimates between the pairwise genetic distance and the pairwise geographic resistance for each SSP. Kernel densities were estimated using a Gaussian distribution. The graduated color contour indicating the (smoothed) observation count is presented on the right side of each plot. The line represents the predicted regression curve between the genetic distance and the geographic resistance from the linear mixed model estimates.*

Discussion

Taken together, the results of this study provide for the first time an extended picture of the effect of anthropogenic disturbance on dispersal in a vertebrate, from dispersal-related phenotypic specialization expressed early in life, through the dispersal pattern emerging in spatially structured populations, to the genetic footprint arising throughout successive generations. Overall, this analysis revealed that anthropogenic disturbance not only strongly promotes dispersal throughout a toad's lifetime, but also prenatally enhances a toadlet's risk-proneness and, to a certain extent, favors longer hind limb length at

metamorphosis. Another finding was that gene flow also substantially increased in anthropogenic
landscapes independently of the SSP's size or functional connectivity.

Natal and breeding dispersal rates and distances in SSPs depend on patch turnover rate

The results revealed contrasting dispersal patterns between riverine and logging SSPs. In the
demographic component of our study, the sampling protocol was weakened by a potential confounding
effect of the population's position along the latitudinal gradient with its status (logging/riverine).
However, we can rule out the possibility of an effect of latitude on the dispersal patterns drawn in our
analyses, as the molecular inferences showed drastically increased gene flow (resulting from dispersal)
in an SSP at a low latitude (L4), which indicates that latitude has a marginal effect on effective dispersal
within SSPs. We are therefore confident in the reliability of our results regarding the effect of turnover
rate on dispersal patterns.

810   In riverine SSPs, we observed a complete suppression of natal dispersal, as well as very low
breeding dispersal, together with a reduced dispersal kernel – the opposite of the dispersal pattern
observed in logging SSPs. In the latter, we nevertheless found some substantial inter-population
variation regarding the contribution of natal dispersal to the overall dispersal process and to seasonal
variation in dispersal rates. Such differences between logging SSPs likely reflect variation in the
anthropogenic disturbance regime resulting from local woodland management practices. Indeed, patch
turnover depends on both the extent and the frequency of the logging activities that create patches, as
well as the post-logging rehabilitation operations that may lead to patch destruction (e.g. filling in of
ruts used as temporary breeding ponds) (Cayuela et al 2018b). This likely results in a continuum of
dispersal strategies along a gradient of patch disturbance, ranging from the near suppression of dispersal
in riverine SSPs to a very high dispersal rate and long dispersal distances in some logging SSPs (e.g. L1
SSP). This pattern is thus very similar to that observed for wing dimorphism in insects alongside patch
temporality gradients (Denno et al. 1996, Roff & Fairban 2007), in which dispersal is suppressed (i.e.
high rate of wingless forms) in persistent habitats, while it is enhanced (i.e. high rate of winged forms)
in more ephemeral habitats according to the (natural) disturbance regime experienced at the landscape
level.

Phenotypic specialization in SSPs depends on patch turnover rate

Previous studies on *Bombina variegata* (Cayuela et al. 2018b) have revealed that dispersal events in
logging SSPs are a mixture of departures conditioned by patch disappearance and unconditional
departures that occur well before patch disappearance. The findings from our common garden
experiments highlight that patch turnover rate prenatally mediates dispersal-related phenotypes and
leads to phenotypic parallelism in logging and riverine SSPs. In particular, toadlets originating from

logging SSPs exhibited higher risk-proneness than those from riverine SSPs, as revealed by their

swiftness in leaving the refuge chamber and in getting out of the arena after crossing a harsh substrate.

Either neophilia or boldness (see Kelleher et al. 2018 for personality traits in amphibians) could explain

the elevated risk-proneness we observed in logging SSPs. Disentangling these two personality traits is

not straightforward (Greggor et al. 2015, Yuen et al. 2017), and further investigations would be useful

to address this. Regardless of the exact composition of personality traits behind our identification of

risk-prone behavior, our results clearly indicate behavioral specialization early in life according to the

disturbance regime prevailing in the landscape.

Concerning morphological traits, we did not find any differences in body size and condition of

toadlets from logging and riverine SSPs. However, the findings showed that toadlets innately have

longer hind limbs in logging than in riverine SSPs. In anurans, hind-limb length is usually positively

associated with locomotor performance (Choi et al. 2003, Philips et al. 2006, Gomes et al 2009, Hudson

et al. 2016), and has also been found to be subject to rapid evolution at the edge of the invasion front in

the introduced species *Rhinella marina* (Philips et al. 2006). Our results thus suggest that long hind

limbs could be a phenotypic trait facilitating dispersal in logging habitats. Yet this difference in leg

length between toadlets from the two environments was only observed in large individuals and was

subject to substantial variation between SSPs. This significant but weak effect of anthropogenic

disturbance on leg length could result from developmental constraints. Limb size mainly depends on the

duration of the larval period in anurans, so species specialized for ephemeral pools (such as *Bombina*

variegata), which are selected for fast larval development, usually exhibit shorter hind limbs compared

to other species (Gomez-Mestre & Buchholz 2006).

Overall, dispersal and related behavioral traits are usually highly plastic phenotypes subject to

partial genetic control (reviewed in Saastamoinen et al. 2018), i.e. determined by $G \times E$ interactions.

Therefore, the phenotypic specialization highlighted in our study may have a genetic basis and/or may

be associated with transgenerational plasticity. In the absence of crossbreed design, we were not able to

disentangle the relative contribution of maternal effect and parental genotypes in the phenotypic

variation observed in logging and riverine SSPs. However, it is possible to rule out the hypothesis of

transgenerational plasticity mediated by toadlet body size at metamorphosis. In amphibians, female

energy investment in breeding influences the size of eggs and the amount of energetic resources

available for the development of embryos and larvae before they become fully heterotrophic (Kaplan

1987, 1992). Studies have reported a positive relationship between egg size and offspring body size at

metamorphosis due to carry-over effects (Laugen et al. 2005, Räsänen et al. 2005, Dziminski et al. 2006),

and body size is an important predictor of behavioral traits (e.g. exploration propensity and risk-taking

behavior) related to dispersal (Kelleher et al. 2017, 2018). In our case study, no difference in body size

was detected in toadlets from logging and riverine environments, which indicates that patch turnover

rate is not likely to alter individual behavior due to the morphological state at metamorphosis in SSPs.

However, epigenetic factors (e.g. DNA methylation, micro-ARN, and histone structure) independent

from female energy investment strategies may lead to transgenerational dispersal plasticity and
contribute to the expression of phenotypic traits that facilitate or hinder dispersal over generations
(Saastamoinen et al. 2018, Cayuela et al. 2019a). Yet it is very likely that a genetic basis partially
determines phenotypic specialization in logging and riverine environments, resulting in incomplete
genetic parallelism between SSPs (Elmer & Meyer 2011, Conte et al. 2012). First, transgenerational
plasticity in dispersal-related traits is usually subject to genetic control (Cayuela et al. 2019a), likely due
to a strong association between epigenetic variation and genetic variants in *cis* and *trans* (Dubin et al.
2015, Zaghlool et al. 2016). Second, as predicted by the dispersal theory and reported in other study
systems, dispersal is partly genetically determined, and behavioral traits related to dispersal often have
a polygenic basis (Saastamoinen et al. 2018). Future studies using Next-Generation Sequencing
approaches (Morozova & Marra 2008, Metzker 2010) could be undertaken to determine the role of
genetic, transcriptional, and epigenetic variation in the disturbance-dependent phenotypic changes and
dispersal evolution in *B. variegata*.

Patch turnover and related dispersal costs and benefits determine SSP dynamics and persistence

Our simulations showed that the absence of dispersal inevitably leads to SSP extinction, and that the
extinction speed increases with patch turnover. When dispersal was possible and had no survival cost,
patch turnover had a positive effect on SSP size. This resulted from an increase in average fecundity due
to the positive relationship between fecundity and patch age reported in logging SSPs (Boualit et al.
2019) and considered in our models. These results are congruent with field observations reporting that
Bombina variegata SSPs may be very large (thousands of adults) in harvested woodlands in western
Europe.
The models also showed that the risk of SSP extinction increased with the survival cost related
to dispersal and was mitigated by informed immigration. This result is congruent with theoretical models
and empirical evidence showing that dispersal only evolves if the benefits of moving outweigh the
related costs (Bonte et al. 2012). In logging SSPs, dispersal seems to be favored as its costs are likely
offset by the benefits of colonizing new patches. Survival (at juvenile, subadult, and adult stage) is lower
in logging than in riverine SSPs (Cayuela et al. 2016a), which likely results from mortality caused by
logging operations and dispersal-related mortality (Cayuela et al. 2018b). This survival cost of dispersal
can be direct, i.e. associated with movement in the landscape matrix (Bonte et al. 2012). It can be also
indirect, resulting from the high energy allocation necessary for recurrent dispersal events over the
toad's lifespan, which might lead to earlier and stronger senescence due to trade-offs (Cayuela et al.
2019b). These direct and indirect costs are obviously offset when animals are forced to disperse
subsequently to patch destruction resulting from logging practices (Cayuela et al 2018b). It should be
noted that even when patches remain available, dispersal may still be strongly favored since their
suitability rapidly declines over time due to the natural silting of ruts if these are not regularly disturbed

by vehicle traffic (Boualit et al. 2019). Furthermore, even if patch suitability is sustained by regular

human disturbance (Boualit et al. 2019), dispersal costs might be mitigated through the colonisation of

recently created patches in which fitness prospects are likely enhanced due to density-dependent

mechanisms. A low density of adults in newly available patches likely reduces the risk of larval

competition, which is an important driver of metamorphosis success (Jasieński 1988) – for this reason

adults preferentially reproduce in tadpole-free waterbodies (Cayuela et al. 2016d, 2017b).

These potential benefits likely favor the evolution of dispersal and dispersal-enhancing

morphological and behavioral traits (i.e. 'dispersal syndromes') in logging SSPs. They also likely

contribute to context-dependent dispersal (Clobert et al. 2009) and matching habitat choice (Edelaar et

al. 2008), implying that individuals adjust their dispersal decisions according to the local fitness

prospects determined by the biotic and abiotic characteristics of breeding patches. This prediction has

been verified by two studies reporting context-dependent dispersal in logging SSPs (Tournier et al. 2017,

Boualit et al. 2019). In particular, these studies showed that adult emigration and immigration decisions

depend on a pond's hydroperiod and the size and annual disturbance of the patch: three factors that

locally affect juvenile recruitment and very likely individual fitness.

In riverine SSPs, the near absence of both natal and breeding dispersal suggests that the benefits

of dispersal do not compensate for its potential costs. First, the absence of patch loss resulting from

anthropogenic or natural processes means individuals are not forced to disperse to survive and

reproduce. In river environments, the local fitness prospects do not deteriorate with patch age, as the

process of natural silting of rocky pools is frequently interrupted by river flooding occurring outside the

breeding period (Cayuela et al. 2011, 2015a). This makes pools available for breeding from one year to

another and limits the risk of larval mortality caused by desiccation. It is also possible that the near

absence of dispersal results from variability in pond characteristics (e.g. hydroperiod and temperature;

Cayuela et al. 2011) within a patch. Indeed, the benefit of dispersing would be low if the variability in

environmental conditions would be similar at intra-patch and inter-patch scales. Overall, the lack of

apparent compensatory benefits should not favor and may even counter-select for dispersal and

dispersal-enhancing traits in riverine SSPs. This is in line with our results, which show that riverine

toadlets innately display low risk-taking behavior and have short hind limbs, two phenotypic traits

hindering dispersal in amphibians (Cayuela et al. 2018c).

Genetic variation patterns in SSPs depend on patch turnover rate

Our study showed for the first time that the spatiotemporal dynamics of habitat patches in a landscape

may have an effect at least as important as landscape fragmentation on gene flow patterns. The findings

highlighted that low dispersal rates and distances are associated with a weaker genetic structure and

lower IBD in riverine than in logging SSPs. Our sampling design allowed us to disentangle the relative

effects of gene flow and genetic drift, which both contribute to genetic differentiation within SSPs

(Slatkin 1977, Broquet & Petit 2009). As well as genetic structure differences caused by patch turnover,

we found higher IBD in the two small SSPs (L4 and R1) than in the two large SSPs in each environment.

Our analysis also took into account the functional connectivity within SSPs by considering two

landscape factors (i.e. topography and hydrological network) that are critically important to the genetic

structure of amphibian populations (reviewed in Cayuela et al. 2018c). As reported in 20 previous

studies (Cayuela et al. 2018c), genetic differentiation increases with increasing topographic slope as this

raises the energy cost of dispersal during the transience phase in amphibians. In addition, our results

showed that genetic differentiation was better explained by the distance of the hydrological network

than the Euclidean distance between two patches. This indicates that the hydrological network improves

the functional connectivity between patches by reducing the cost of displacement in the landscape

matrix, which is congruent with the findings of six previous studies (Cayuela et al. 2018c). Interestingly,

genetic differentiation nevertheless remained higher in riverine than in logging SSPs, despite a denser

hydrological network and therefore weaker landscape resistance in riverine SSPs. This suggests that in

our study system, the rate of patch turnover may be a more important driver of neutral genetic variation

than landscape connectivity.

Our findings suggest that variation in the turnover rate of SSPs has far-reaching consequences

on the evolutionary forces involved in the migration–selection–drift balance. In riverine SSPs, reduced

gene flow between patches leads to lower genetic diversity and smaller effective population size

compared to logging SSPs. Such low standing genetic variation could limit the adaptive response of

riverine SSPs to novel environmental conditions. The probability of allele fixation increases with the

magnitude of the beneficial effect and the effective population size, and this probability is significantly

higher when the allele has a high initial frequency (Barrett & Schluter 2008, Hedrick 2013). Moreover,

standing genetic variation usually allows faster adaptation as beneficial alleles are already present in the

population (Barrett & Schluter 2008, Hedrick 2013). Overall, our results suggest that riverine SSPs

should suffer from a lower capacity of 'evolutionary rescue' (Carlson et al. 2014) than logging SSPs,

which could increase their sensitivity to current global changes.

The results also revealed a high level of inbreeding and relatedness within riverine SSPs,

confirming the results of a previous study (Cayuela et al. 2017a). These findings raise important

questions about the mechanisms associated with the repression of inbreeding depression. As mentioned,

the survival rate at all life stages (juvenile, subadult, and adult) is higher in riverine than in logging SSPs

(Cayuela et al. 2016a), suggesting a marginal effect of inbreeding on postmetamorphic survival and

negligible inbreeding depression. A previous study conducted in the R2 SSP indicated that the absence

of disassortative mating does not seem to mitigate inbreeding risk: females even prefer to reproduce

with related males from their own patch (Cayuela et al. 2017a). Although the effect of inbreeding on

reproductive performance remains as yet unevaluated, those results suggest a low genetic load in riverine

SSPs. This could be due to high efficiency in purging deleterious alleles and the genomic architecture

of genetic load, especially a low linkage of deleterious recessive alleles (Bersabé et al. 2016, Hedrick &

Garcia-Dorado 2016). Future studies could be carried out to identify the mechanisms involved in the

genetic purging of inbreeding depression resulting from limited dispersal in riverine SSPs.

Conclusion

Our study showed that anthropogenic habitat disturbance is likely an important driver of dispersal

evolution. The results found that, like landscape fragmentation, human-driven variation in patch

turnover may promote morphological and behavioral specialization related to dispersal. In particular, it

may lead to phenotypic parallelism affecting dispersal syndromes and patterns (dispersal rate and

distance) in SSPs exposed to contrasting levels of patch turnover. This phenotypic parallelism is likely

underpinned by genetic and/or epigenetic parallelism, for which the molecular basis remains to be

investigated. Our results also revealed that differences in dispersal patterns are associated with variation

in the genetic structure of SSPs, which might affect local eco-evolutionary dynamics (Legrand et al.

2017). In particular, high gene flow and reduced effects of genetic drift allow higher genetic

polymorphism to be maintained in SSPs experiencing high patch turnover than in SSPs with low patch

turnover. In parallel, larger effective population size is expected to increase selection effectiveness in

SSPs exposed to high patch turnover, giving them higher evolutionary potential and increased chances

of evolutionary rescue in the case of environmental change (Carlson et al. 2014). These results

emphasize the central role of anthropogenic disturbance in the spatiotemporal dynamics of landscapes

and the related ecological and evolutionary processes.

1002   In terms of the implications for conservation, the findings suggest that the approach may need

to differ depending on the demographic and genetic characteristics of SSPs driven by the level of patch

turnover. Those SSPs exposed to high patch turnover are likely to be highly sensitive to land use changes

that result in a loss of functional connectivity between patches. An increase in landscape resistance

reduces the success of dispersal following the disappearance of a patch and diminishes the chance of

new patch colonization, which enhances the extinction risk of the whole SSP. In contrast, SSPs

experiencing a low level of patch turnover suffer a loss of genetic diversity and high inbreeding, which

may increase their sensitivity to environmental change (e.g. climate change or the emergence of new

pathogens), potentially speeding up their extinction if they experience a demographic decline (i.e.

inbreeding depression; Reed & Frankham 2003, Frankham 2010). This underlines the tremendous

importance of adjusting conservation decisions to the demographic and genetic characteristics of SSPs

adapted to contrasting regimes of habitat disturbance.

1014

1015   **Acknowledgments**

1016

We would like to thank all the undergraduate students and fieldworkers that contributed to the capture–recapture surveys, DNA samples, and common garden behavioral tests. This research project was funded by the Office National des Forêts, the Lorraine Direction Régionale de l'Environnement, de l'Aménagement et du Logement (DREAL), the Agence de l'Eau Rhin-Meuse, the Conseil Régional de Lorraine, the Conseil Régional de Champagne-Ardenne, the Conseil Régional de Picardie, the Conseil Général de l'Aisne, the Conseil Général d'Ardèche, the Conseil Général d'Isère and the Communauté de Communes de l'Argonne Ardennaise (2C2A). The study was authorized by three government environmental agencies: the Lorraine DREAL, the Rhône-Alpes DREAL and the Agence de l'Eau Rhin-Meuse. The ethical committee of Université Claude Bernard Lyon 1 approved the capture–recapture and common garden protocols.

References

Armsworth, P. R., & Roughgarden, J. E. (2005). Disturbance induces the contrasting evolution of reinforcement and dispersiveness is directed and random movers. Evolution, 59, 2083-2096.

Atkins, J. L., Perry, G. L., & Dennis, T. E. (2019). Effects of mis-alignment between dispersal traits and landscape structure on dispersal success in fragmented landscapes. Royal Society Open Science, 6, 181702.

Baguette, M., Blanchet, S., Legrand, D., Stevens, V. M., & Turlure, C. (2013). Individual dispersal, landscape connectivity and ecological networks. Biological Reviews, 88, 310-326.

Balkenhol, N., Holbrook, J. D., Onorato, D., Zager, P., White, C., & Waits, L. P. (2014). A multi-method approach for analyzing hierarchical genetic structures: a case study with cougars Puma concolor. Ecography, 37, 552-563.

Barrett, R. D., & Schluter, D. (2008). Adaptation from standing genetic variation. Trends in Ecology & Evolution, 23, 38-44.

Bersabé, D., Caballero, A., Pérez-Figueroa, A., & García-Dorado, A. (2016). On the consequences of purging and linkage on fitness and genetic diversity. G3: Genes, Genomes, Genetics, 6, 171-181.

Bonte, D., Van Dyck, H., Bullock, J. M., Coulon, A., Delgado, M., Gibbs, M., Lehouck, V., Matthysen, E., Mustin, K., Saastamoinen, M., Schtickzelle, N., Stevens, V. M., Vandewoestijne, S., Baguette, M., Barton, K., Benton, T.G., Chaput-Bardy, A., Clobert, J., Dytham, C., Hovestadt, T., Meier, C. M., Palmer, S.C.F., Turlure, C., Travis, J. M. J. (2012). Costs of dispersal. Biological Reviews, 87, 290-312.

Boualit, L., Pichenot, J., Besnard, A., Helder, R., Joly, P., & Cayuela, H. (2019). Environmentally mediated reproductive success predicts breeding dispersal decisions in an early successional amphibian. Animal behaviour, 149, 107-120.

Bowler, D. E., & Benton, T. G. (2005). Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. Biological Reviews, 80, 205-225.

Broquet, T., & Petit, E. J. (2009). Molecular estimation of dispersal for ecology and population genetics.
Annual Review of Ecology, Evolution, and Systematics, 40, 193-216.

Carlson, S. M., Cunningham, C. J., & Westley, P. A. (2014). Evolutionary rescue in a changing world.
Trends in Ecology & Evolution, 29, 521-530.

Caswell H. (2001) Matrix population models: constraints, analysis and interpretation, Sinauer
Associates, Sunderland.

Cayuela, H., Cheylan, M., & Joly, P. (2011). The best of a harsh lot in a specialized species: breeding
habitat use by the yellow-bellied toad (*Bombina variegata*) on rocky riverbanks. Amphibia-Reptilia,
32, 533-539.

Cayuela, H., Arsovski, D., Boitaud, S., Bonnaire, E., Boualit, L., Miaud, C., Joly, P., & Besnard, A.
(2015a). Slow life history and rapid extreme flood: demographic mechanisms and their consequences
on population viability in a threatened amphibian. Freshwater Biology, 60, 2349-2361.

Cayuela, H., Lambrey, J., Vacher, J. P., & Miaud, C. (2015b). Highlighting the effects of land-use
change on a threatened amphibian in a human-dominated landscape. Population Ecology, 57, 433-
443.

Cayuela, H., Arsovski, D., Thirion, J. M., Bonnaire, E., Pichenot, J., Boitaud, S., Brison, A.-L., Miaud,
C., Joly, P., & Besnard, A. (2016a). Contrasting patterns of environmental fluctuation contribute to
divergent life histories among amphibian populations. Ecology, 97, 980-991.

Cayuela, H., Boualit, L., Arsovski, D., Bonnaire, E., Pichenot, J., Bellec, A., Miaud, C., Léna, J.-P.,
Joly, P., & Besnard, A. (2016b). Does habitat unpredictability promote the evolution of a colonizer
syndrome in amphibian metapopulations?. Ecology, 97, 2658-2670.

Cayuela, H., Arsovski, D., Thirion, J. M., Bonnaire, E., Pichenot, J., Boitaud, S., Miaud, C., Joly, P., &
Besnard, A. (2016c). Demographic responses to weather fluctuations are context dependent in a long-
lived amphibian. Global Change Biology, 22, 2676-2687.

Cayuela, H., Lengagne, T., Kaufmann, B., Joly, P., & Léna, J. P. (2016d). Larval competition risk shapes
male–male competition and mating behavior in an anuran. Behavioral Ecology, arw100.

Cayuela, H., Léna, J. P., Lengagne, T., Kaufmann, B., Mondy, N., Konecny, L., Dumet, A., Vienney,
A., & Joly, P. (2017a). Relatedness predicts male mating success in a pond-breeding amphibian.
Animal Behaviour, 130, 251-261.

Cayuela, H., Lengagne, T., Joly, P., & Léna, J. P. (2017b). Females trade off the uncertainty of breeding
resource suitability with male quality during mate choice in an anuran. Animal Behaviour, 123, 179-
185.

Cayuela, H., Rougemont, Q., Prunier, J. G., Moore, J. S., Clobert, J., Besnard, A., & Bernatchez, L.
(2018a). Demographic and genetic approaches to study dispersal in wild animal populations: a
methodological review. Molecular Ecology, 27, 3976-4010.

Cayuela, H., Besnard, A., Quay, L., Helder, R., Léna, J. P., Joly, P., & Pichenot, J. (2018b).
Demographic response to patch destruction in a spatially structured amphibian population. Journal
of Applied Ecology, 55, 2204-2215.

Cayuela, H., Valenzuela-Sanchez, A., Teulier, L., Martínez-Solano, Í., Léna, J. P., Merilä, J., Muths, E.,
Shine, R., Quay, L., Denoël, M., Clobert, J., Schmidt, B. R. (2018c). Determinants and consequences
of dispersal in vertebrates with complex life cycles: a review of pond-breeding amphibians. PeerJ
Preprints, e27394v1.

Cayuela, H., Jacob, S., Schtickzelle, N., Verdonck, R., Philippe, H., Laporte, M., Bernatchez, L.,
Legrand, D. (2019a). Transgenerational dispersal plasticity and its fitness consequences are under
genetic control. bioXiv, doi: ...

Cayuela, H., Lemaitre, J. F., Bonnaire, E., Pichenot, J., & Schmidt, B. (2019b). Population position
along the fast-slow life-history continuum predicts intraspecific variation in actuarial senescence.
bioRxiv, 621425.

Cayuela, H., Gillet, L., Laudelout, A., Besnard, A., Bonnaire, E., Levionnois, P., Muths, E., Dufrêne,
M., & Kinet, T. (2019c). Survival cost to relocation does not reduce population self-sustainability in
an amphibian. Ecological Applications, 29, e01909.

Cheptou, P. O., Hargreaves, A. L., Bonte, D., & Jacquemyn, H. (2017). Adaptation to fragmentation:
evolutionary dynamics driven by human influences. Philosophical Transactions of the Royal Society
B: Biological Sciences, 372, 20160037.

Choquet, R., Rouan, L., & Pradel, R. (2009). Program E-SURGE: a software application for fitting
multievent models. In: Thomson, D. L., Cooch, E. G., Conroy, M. J. (eds). Modeling demographic
processes in marked populations. Pp. 845-865. Springer, Boston, US.

Clarke, R., Rothery, P. & Raybould, A. (2002). Confidence limits for regression relationships between
distance matrices: Estimating gene flow with distance. Journal of Agricultural, Biological, and
Environmental Statistics, 7, 361-372.

Clobert, J., Galliard, L., Cote, J., Meylan, S., & Massot, M. (2009). Informed dispersal, heterogeneity
in animal dispersal syndromes and the dynamics of spatially structured populations. Ecology letters,
12, 197-209.

Comins, H. N., Hamilton, W. D., & May, R. M. (1980). Evolutionarily stable dispersal strategies.
Journal of Theoretical Biology, 82, 205-230.

Conte, G. L., Arnegard, M. E., Peichel, C. L., & Schluter, D. (2012). The probability of genetic
parallelism and convergence in natural populations. Proceedings of the Royal Society B: Biological
Sciences, 279, 5039-5047.

Cote, J., Clobert, J., Brodin, T., Fogarty, S., & Sih, A. (2010). Personality-dependent dispersal:
characterization, ontogeny and consequences for spatially structured populations. Philosophical
Transactions of the Royal Society B: Biological Sciences, 365, 4065-4076.

Cote, J., Bestion, E., Jacob, S., Travis, J., Legrand, D., & Baguette, M. (2017). Evolution of dispersal
strategies and dispersal syndromes in fragmented landscapes. Ecography, 40, 56-73.

Coulon, A., Fitzpatrick, J. W., Bowman, R., Stith, B. M., Makarewich, C. A., Stenzler, L. M., & Lovette,
I. J. (2008). Congruent population structure inferred from dispersal behaviour and intensive genetic
surveys of the threatened Florida scrub-jay (*Aphelocoma cœrulescens*). Molecular Ecology, 17,
1685-1701.

Cushman, S. A. (2006a). Effects of habitat loss and fragmentation on amphibians: a review and
prospectus. Biological conservation, 128, 231-240.

Cushman, S. A., McKelvey, K. S., Hayden, J., & Schwartz, M. K. (2006b). Gene flow in complex
landscapes: testing multiple hypotheses with causal modeling. The American Naturalist, 168, 486-
499.

Cushman, S., Storfer, A., & Waits, L. (2015). Landscape genetics: concepts, methods, applications. John
Wiley & Sons.

Denno, R. F., Roderick, G. K., Peterson, M. A., Huberty, A. F., Dobel, H. G., Eubanks, M. D., Losey,
J. E., & Langellotto, G. A. (1996). Habitat persistence underlies intraspecific variation in the
dispersal strategies of planthoppers. Ecological Monographs, 66, 389-408.

Do, C., Waples, R. S., Peel, D., Macbeth, G. M., Tillett, B. J. & Ovenden, J. R. (2014) NeEstimator V2:
re-implementation of software for the estimation of contemporary effective population size (Ne)
from genetic data. Molecular Ecology Resources, 14, 209-214.

Dubin, M. J., Zhang, P., Meng, D., Remigereau, M. S., Osborne, E. J., Casale, F. P., Drewe, P. Kahles,
A., Jean, G., Vilhjalmsson, B., Jagoda, J., Irez, S., Voronin, V., Song, Q., Long, Q., Ratsch, G.,
Stegle, O., Clark, R. M., & Nordborg, M. (2015). DNA methylation in Arabidopsis has a genetic
basis and shows evidence of local adaptation. elife, 4, e05255.

Dziminski, M. A., & Roberts, J. D. (2006). Fitness consequences of variable maternal provisioning in
quacking frogs (Crinia georgiana). Journal of Evolutionary Biology, 19, 144-155.

Earl, D. A. (2012). STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE
output and implementing the Evanno method. Conservation Genetics Resources, 4, 359-361.

Edelaar, P., Siepielski, A. M., & Clobert, J. (2008). Matching habitat choice causes directed gene flow:
a neglected dimension in evolution and ecology. Evolution, 62, 2462-2472.

Edwards, D. P., Gilroy, J. J., Thomas, G. H., Uribe, C. A. M., & Haugaasen, T. (2015). Land-sparing
agriculture best protects avian phylogenetic diversity. Current biology, 25, 2384-2391.

Elmer, K. R., & Meyer, A. (2011). Adaptation in the age of ecological genomics: insights from
parallelism and convergence. Trends in Ecology & Evolution, 26, 298-306.

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the
software STRUCTURE: a simulation study. Molecular Ecology, 14, 2611-2620.

Fahrig, L. (2003). Effects of habitat fragmentation on biodiversity. Annual Review of Ecology,
Evolution, and Systematics, 34, 487-515.

Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S., Coe, M.
T., Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A., Kucharik, C. J.,
Monfreda, C., Patz, J. A., Prentice, I. C., Ramankutty, N., & Snyder, P. K. (2005). Global
consequences of land use. Science, 309, 570-574.

Frankham, R. (2010). Inbreeding in the wild really does matter. Heredity, 104, 124.

Frishkoff, L. O., Hadly, E. A., & Daily, G. C. (2015). Thermal niche predicts tolerance to habitat
conversion in tropical amphibians and reptiles. Global Change Biology, 21, 3901-3916.

Gibson, L., Lee, T. M., Koh, L. P., Brook, B. W., Gardner, T. A., Barlow, J., Peres, A. C., Bradshaw,
C. J. A., Laurance, W. F., Lovejoy, T. E., & Sodhi, N. S. (2011). Primary forests are irreplaceable
for sustaining tropical biodiversity. Nature, 478, 378.

Gilpin, M. (Ed.). (2012). Metapopulation dynamics: empirical and theoretical investigations. Academic
Press.

Gomes, F. R., Rezende, E. L., Grizante, M. B. & Navas, C. A. (2009). The evolution of jumping
performance in anurans morphological correlates and ecological implications Journal of
Evolutionary Biology, 22, 1088-1097

Grant, E. H. C., Nichols, J. D., Lowe, W. H., & Fagan, W. F. (2010). Use of multiple dispersal pathways
facilitates amphibian persistence in stream networks. Proceedings of the National Academy of
Sciences, 201000266.

Greggor, A. L., Thornton, A., & Clayton, N. S. (2015). Neophobia is not only avoidance: improving
neophobia tests by combining cognition and ecology. Current Opinion in Behavioral Sciences, 6, 82-
89.

Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., Lovejoy, T. E.,
Sexton, J. O., Austin, M. P., Collins, C. D., Cook, W. M., Damschen, E. I., Ewers, R. M., Foster, B.
L., Jenkins, C. N., King, A. J., Laurance, W. F., Levey, D. J., Margules, C. R., Melbourne, B. A.,
Nicholls, A. O., Orrock, J. L., Song, D. X., & Townshend, J. R (2015). Habitat fragmentation and its
lasting impact on Earth's ecosystems. Science Advances, 1, e1500052.

Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A. A., Tyukavina, A., Thau,
D., Stehman, S. V., Goetz, S. J., Loveland, T. R., Kommareddy, A., Egorov, A., Chini, L., Justice,
C. O., & Townshend, J. R. G. (2013). High-resolution global maps of 21st-century forest cover
change. Science, 342, 850-853.

Hanski, I., & Gaggiotti, O. E. (Eds.). (2004). Ecology, genetics, and evolution of metapopulations.
Academic Press.

Hanski, I. A. (2011). Eco-evolutionary spatial dynamics in the Glanville fritillary butterfly. Proceedings
of the National Academy of Sciences, 108, 14397-14404.

Hanski, I., & Mononen, T. (2011). Eco-evolutionary dynamics of dispersal in spatially heterogeneous
environments. Ecology Letters, 14, 1025-1034.

Hanski, I., Schulz, T., Wong, S. C., Ahola, V., Ruokolainen, A., & Ojanen, S. P. (2017). Ecological and

genetic basis of metapopulation persistence of the Glanville fritillary butterfly in fragmented

landscapes. Nature Communications, 8, 14504.

Hedrick, P. W. (2013). Adaptive introgression in animals: examples and comparison to new mutation

and standing variation as sources of adaptive variation. Molecular Ecology, 22, 4606-4618.

Hedrick, P. W., & Garcia-Dorado, A. (2016). Understanding inbreeding depression, purging, and

genetic rescue. Trends in Ecology & Evolution, 31, 940-952.

Hillman, S.S., Drewes, R.C., Hedrick, M.S. & Hancock, T.V. (2014). Physiological vagility and its

relationship to dispersal and neutral genetic heterogeneity in vertebrates. The Journal of

Experimental Biology, 217, 3356-3364.

Jakobsson, M., & Rosenberg, N. A. (2007). CLUMPP: a cluster matching and permutation program for

dealing with label switching and multimodality in analysis of population structure. Bioinformatics,

23, 1801-1806.

Kaplan, R. H. (1987). Developmental plasticity and maternal effects of reproductive characteristics in

the frog, *Bombina orientalis*. Oecologia, 71, 273-279.

Kaplan, R. H. (1992). Greater maternal investment can decrease offspring survival in the frog *Bombina*

orientalis. Ecology, 73, 280-288.

Kelleher, S. R., Silla, A. J., Dingemanse, N. J., & Byrne, P. G. (2017). Body size predicts between-

individual differences in exploration behaviour in the southern corroboree frog. Animal behaviour,

129, 161-170.

Kelleher, S. R., Silla, A. J., & Byrne, P. G. (2018). Animal personality and behavioral syndromes in

amphibians: a review of the evidence, experimental approaches, and implications for conservation.

Behavioral Ecology and Sociobiology, 72, 79.

Kokko, H., & López-Sepulcre, A. (2006). From individual dispersal to species ranges: perspectives for

a changing world. Science, 313, 789-791.

Kubisch, A., Holt, R.D., Poethke, H.J. & Fronhofer, E.A. 2014. Where am I and why? Synthesising

range biology and the eco-evolutionary dynamics of dispersal. Oikos 123:5–22.

Gosner, K. L. (1960). A simplified table for staging anuran embryos and larvae with notes on

identification. Herpetologica, 16, 183-190.

Lagrange, P., Pradel, R., Bélisle, M., & Gimenez, O. (2014). Estimating dispersal among numerous sites

using capture–recapture data. Ecology, 95, 2316-2323.

Laugen, A. T., Kruuk, L. E., Laurila, A., Räsänen, K., Stone, J., & Merilä, J. (2005). Quantitative

genetics of larval life-history traits in Rana temporaria in different environmental conditions.

Genetics Research, 86, 161-170.

Lebreton, J. D., Nichols, J. D., Barker, R. J., Pradel, R., & Spendelow, J. A. (2009). Modeling individual

animal histories with multistate capture–recapture models. Advances in Ecological Research, 41, 87-

173.

Legendre P., & Legendre A. (1998) Complex ecological data sets. In: Legendre, P., Legendre, A. (eds)
Numerical Ecology. Pp. 247–302. Elsevier, Amsterdam, Netherlands.

Legrand, D., Cote, J., Fronhofer, E. A., Holt, R. D., Ronce, O., Schtickzelle, N., Travis, J. M. J., &
Clobert, J. (2017). Eco-evolutionary dynamics in fragmented landscapes. Ecography, 40, 9-25.

Lenormand, T. (2002). Gene flow and the limits to natural selection. Trends in Ecology & Evolution,
17, 183-189.

Littell, R. C., Milliken, G. A. , Stroup, W. W. , Wolfinger, R. D. (2006). SAS system for mixed models
2nd ed. SAS Institute, Inc., Cary, North Carolina, USA.

Lowe, W. H. (2003). Linking dispersal to local population dynamics: a case study using a headwater
salamander system. Ecology, 84, 2145-2154.

Matthysen, E. (2012). Multicausality of dispersal: a review. In Clobert, J., Baguette, M., Benton, T. G.,
& Bullock, J. M.: Dispersal Ecology and Evolution. Pp. 3–18, Oxford University Press, Oxford, UK.

McCullagh, P., & Nelder, J. A. (1989). Generalized linear models. CRC press.

McPeek, M. A., & Holt, R. D. (1992). The evolution of dispersal in spatially and temporally varying
environments. The American Naturalist, 140, 1010-1027.

McRae, B. H., & Beier, P. (2007). Circuit theory predicts gene flow in plant and animal populations.
Proceedings of the National Academy of Sciences of the United States of America, 104, 19885–
19890.

Metzker, M. L. (2010). Sequencing technologies—the next generation. Nature Reviews Genetics, 11,
31.

Morozova, O., & Marra, M. A. (2008). Applications of next-generation sequencing technologies in
functional genomics. Genomics, 92, 255-264.

Newbold, T., Hudson, L. N., Hill, S. L., Contu, S., Lysenko, I., Senior, R. A., Börger, L., Bennett, D. J.,
Choimes, A., Collen, B., Day, J., De Palma, A., Díaz, S., Echeverria-Londoño, S., Edgar, M. J.,
Feldman, A., Garon, M., Harrison, M. L. K., Alhusseini, T., Ingram, D. J., Itescu, Y., Kattge, J.,
Kemp, V., Kirkpatrick, L., Kleyer, M., Laginha Pinto Correia, D., Martin, C. D., Meiri, S.,
Novosolov, M., Pan, Y., Phillips, H. R. P., Purves, D. W., Robinson, A., Simpson, J., Tuck, S. L.,
Weiher, E., White, H. J., Ewers, R. M., Mace, G. M., Scharlemann, J. P. W., & Purvis, A. (2015).
Global effects of land use on local terrestrial biodiversity. Nature, 520, 45.

Newman, E. A. (2019). Disturbance ecology in the Anthropocene. Frontiers in Ecology and Evolution,
7, 147.

Nowakowski, A. J., Frishkoff, L. O., Thompson, M. E., Smith, T. M., & Todd, B. D. (2018).
Phylogenetic homogenization of amphibian assemblages in human-altered habitats across the globe.
Proceedings of the National Academy of Sciences, 115, E3454-E3462.

Ochocki, B. M., & Miller, T. E. X. (2017). Rapid evolution of dispersal ability makes biological
invasions faster and more variable. Nature Communications, 8, 14315.

Otto, S. P. (2018). Adaptation, speciation and extinction in the Anthropocene. Proceedings of the Royal
Society B, 285, 20182047.

Peakall, R. O. D., & Smouse, P. E. (2006). GENALEX 6: genetic analysis in Excel. Population genetic
software for teaching and research. Molecular Ecology Notes, 6, 288-295.

Pelletier, F., & Coltman, D. W. (2018). Will human influences on evolutionary dynamics in the wild
pervade the Anthropocene?. BMC Biology, 16, 7.

Pennekamp, F., Schtickzelle, N., & Petchey, O. L. (2015). BEMOVI, software for extracting behavior
and morphology from videos, illustrated with analyses of microbes. Ecology and Evolution, 5, 2584-
2595.

Pereira, H. M., Leadley, P. W., Proença, V., Alkemade, R., Scharlemann, J. P., Fernandez-Manjarrés, J.
F., Araújo, M. B., Balvanera, P., Biggs, R., Cheung, W. M. L., Chini, L., Cooper, H. D., Gilman, E.
L., Guénette, S., Hurtt, G. C., Huntington, H. P., Mace, G. M., Oberdorff, T., Revenga, C., Rodrigues,
P., Scholes, R. J., Sumaila, U. R., Walpole, M. (2010). Scenarios for global biodiversity in the 21st
century. Science, 330, 1496-1501.

Pollock, K. H. (1982). A capture-recapture design robust to unequal probability of capture. The Journal
of Wildlife Management, 46, 752-757.

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus
genotype data. Genetics, 155, 945-959.

Prunier, J. G., Kaufmann, B., Fenet, S., Picard, D., Pompanon, F., Joly, P., & Lena, J. P. (2013).
Optimizing the trade-off between spatial and genetic sampling efforts in patchy populations:
Towards a better assessment of functional connectivity using an individual-based sampling scheme.
Molecular Ecology, 22, 5516-5530.

Prunier, J. G., Colyn, M., Legendre, X., Nimon, K. F., & Flamand, M. C. (2015). Multicollinearity in
spatial genetics: separating the wheat from the chaff using commonality analyses. Molecular
Ecology, 24, 263-283.

Räsänen, K., Laurila, A., & Merilä, J. (2005). Maternal investment in egg size: environment-and
population-specific effects on offspring performance. Oecologia, 142, 546-553.

Reed, D. H., & Frankham, R. (2003). Correlation between fitness and genetic diversity. Conservation
Biology, 17, 230-237.

Roff, D. A., & Fairbairn, D. J. (2007). The evolution and genetics of migration in insects. AIBS Bulletin,
57, 155-164.

Ronce, O. (2007). How does it feel to be like a rolling stone? Ten questions about dispersal evolution.
Annual Review of Ecology, Evolution, and Systematics, 38, 231-253.

Ronce, O., & Clobert, J. (2012). Dispersal syndromes. In Clobert, J., Baguette, M., Benton, T. G., &
Bullock, J. M.: Dispersal Ecology and Evolution. Pp. 119–138, Oxford University Press, Oxford,
UK.

Rousset, F. (2008). genepop'007: a complete re-implementation of the genepop software for Windows
and Linux. Molecular Ecology Resources, 8, 103-106.

Saastamoinen, M., Bocedi, G., Cote, J., Legrand, D., Guillaume, F., Wheat, C.W., Fronhofer, E.A.,
Garcia, C., Henry, R., Husby, A., Baguette, M., Bonte, D., Coulon, A., Kokko, H., Matthysen, E.,
Niitepõld, K., Nonaka, E., Stevens, V.M., Travis, J.M., Donohue, K., Bullock, J.M., del Mar
Delgado, M. (2018) Genetics of dispersal. Biological Reviews, 93, 574-599.

Sala, O. E., Chapin, F. S., Armesto, J. J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-Sanwald, E.,
Huenneke, L. F., Jackson, R. B., Kinzig, A., Leemans, R., Lodge, D. M., Mooney, H. A., Oesterheld,
M., LeRoy Poff, N., Sykes, M. T., Walker, B. H., Walker, M., & Wall, D. H. (2000). Global
biodiversity scenarios for the year 2100. Science, 287, 1770-1774.

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients. Methods
in Ecology and Evolution, 1, 103-113.

Selkoe, K. A., Watson, J. R., White, C., Horin, T. B., Iacchei, M., Mitarai, S., Siegel, D.A., Gaines, S.D.
& Toonen, R. J. (2010). Taking the chaos out of genetic patchiness: Seascape genetics reveals
ecological and oceanographic drivers of genetic patterns in three temperate reef species. Molecular
Ecology, 19, 3708–3726.

Slatkin, M. (1977). Gene flow and genetic drift in a species subject to frequent local extinctions.
Theoretical Population Biology, 12, 253-262.

Thomas, C. D., & Kunin, W. E. (1999). The spatial structure of populations. Journal of Animal Ecology,
68, 647-657.

Tilman, D., Clark, M., Williams, D. R., Kimmel, K., Polasky, S., & Packer, C. (2017). Future threats to
biodiversity and pathways to their prevention. Nature, 546, 73.

Tournier, E., Besnard, A., Tournier, V., & Cayuela, H. (2017). Manipulating waterbody hydroperiod
affects movement behaviour and occupancy dynamics in an amphibian. Freshwater Biology, 62,
1768-1782.

Travis, J.M.J., Mustin, K., Benton, T.G. & Dytham, C. 2009. Accelerating invasion rates result from the
evolution of density-dependent dispersal. Journal Theoretical Biology, 259, 151–158.

Turner, M. G. (2010). Disturbance and landscape dynamics in a changing world. Ecology, 91, 2833-
2849.

Van Oosterhout, C., Hutchinson, W. F., Wills, D. P., & Shipley, P. (2004). MICRO-CHECKER:
software for identifying and correcting genotyping errors in microsatellite data. Molecular Ecology
Notes, 4, 535-538.

Van Strien, M. J., Keller, D., & Holderegger, R. (2012). A new analytical approach to landscape genetic
modelling: Least-cost transect analysis and linear mixed models. Molecular Ecology, 21, 4010–4023.

Villard, M. A., & Metzger, J. P. (2014). Beyond the fragmentation debate: a conceptual model to predict
when habitat configuration really matters. Journal of Applied Ecology, 51(2), 309-318.

Wang J (2011) COANCESTRY: a program for simulating, estimating and analysing relatedness and

inbreeding coefficients. Molecular Ecology Resources, 11, 141–145.

Warren, S. D., & Büttner, R. (2008). Relationship of endangered amphibians to landscape disturbance.

The Journal of Wildlife Management, 72, 738-744.

Yuen, C. H., Schoepf, I., Schradin, C., & Pillay, N. (2017). Boldness: are open field and startle tests

measuring the same personality trait?. Animal Behaviour, 128, 143-151.

Zaghlool, S. B., Al-Shafai, M., Al Muftah, W. A., Kumar, P., Gieger, C., Waldenberger, M., Falchi, M.,

& Suhre, K. (2016). Mendelian inheritance of trimodal CpG methylation sites suggests distal cis-

acting genetic effects. Clinical Epigenetics, 8, 124.

1352