Anthropogenic disturbance drives dispersal syndromes, demography, and gene flow in
spatially structured amphibian populations

Hugo Cayuela [1,2], Aurélien Besnard[2], Julien Cote[3], Martin Laporte[2], Eric Bonnaire[4], Julian Pichenot[5],
Nicolas Schtickzelle[6], Arnaud Bellec[1], Pierre Joly[1], Jean-Paul Léna[1]

[1]UMR 5023 LEHNA, Laboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés (UMR 5023
LEHNA), 69100 Villeurbanne, France
[2]EPHE, UM, SupAgro, IRD, INRA, UMR 5175 CEFE, CNRS, PSL Research University, Montpellier
F-34293 France
[3]CNRS, Université Toulouse III Paul Sabatier, ENFA ; UMR5174EDB (Laboratoire Évolution &
Diversité Biologique) ; 118 route de Narbonne, F-31062 Toulouse, France
[4]Office National des Forêts, Agence de Verdun, 55100 Verdun, France
[5]CERFE, Centre de Recherche et Formation en Eco-éthologie (CERFE), 08240 Boult-aux-Bois,
France
[6] Earth and Life Institute, Biodiversity Research Centre, Université Catholique de Louvain, 1348
Louvain-la-Neuve, Belgium

Running head: Disturbance and dispersal

Abstract

There is growing evidence that anthropogenic landscapes can strongly influence the evolution of
dispersal, particularly through fragmentation, and may drive organisms into an evolutionary trap by
suppressing dispersal. However, the influence on dispersal evolution of anthropogenic variation in
habitat patch turnover has so far been largely overlooked. In this study, we examined how human-
driven variation in patch persistence affects dispersal rates and distances, determines dispersal-related
phenotypic specialization, and drives neutral genetic structure in spatially structured populations. We
addressed this issue in an amphibian, *Bombina variegata*, using an integrative approach combining
capture–recapture modeling, demographic simulation, common garden experiments, and population
genetics. *B. variegata* reproduces in small ponds that occur either in habitat patches that are persistent
(i.e. several decades or more), located in riverine environments with negligible human activity, or in
patches that are highly temporary (i.e. a few years), created by logging operations in intensively
harvested woodland. Our capture–recapture models revealed that natal and breeding dispersal rates
and distances were drastically higher in spatially structured populations (SSPs) in logging
environments than in riverine SSPs. Population simulations additionally showed that dispersal costs

and benefits drive the fate of logging SSPs, which cannot persist without dispersal. The common garden experiments revealed that toadlets reared in laboratory conditions have morphological and behavioral specialization that depends on their habitat of origin. Toadlets from logging SSPs were found to have higher boldness and exploration propensity than those from riverine SSPs, indicating transgenerationally transmitted dispersal syndromes. We also found contrasting patterns of neutral genetic diversity and gene flow in riverine and logging SSPs, with genetic diversity and effective population size considerably higher in logging than in riverine SSPs. In parallel, intra-patch inbreeding and relatedness levels were lower in logging SSPs. Controlling for the effect of genetic drift and landscape connectivity, gene flow was found to be higher in logging than in riverine SSPs. Taken together, these results indicate that anthropogenic variation in habitat patch turnover may have an effect at least as important as landscape fragmentation on dispersal evolution and the long-term viability and genetic structure of wild populations.

Keyword: Dispersal, movement behavior, behavioral syndrome, metapopulation, genetic structure, amphibian

Introduction

In the age of the Anthropocene, a significant proportion of land cover has been replaced by human-dominated landscapes (Foley et al. 2005, Pereira et al. 2010, Gibson et al. 2011, Tilman et al. 2017), with the result that the conditions prevailing in these anthropogenic environments now shape the evolutionary course of almost all species (Otto 2018, Pelletier & Coltman 2018). Land use changes usually have the simultaneous effects of habitat loss, alteration, and/or fragmentation into small habitat patches isolated in a more or less hostile matrix (i.e. unsuitable habitat) (Fahrig 2003, Villard & Metzger 2014). While it is widely accepted that habitat loss is the main factor involved in local extinction and biodiversity loss (Sala et al. 2000, Pereira et al. 2010, Newbold et al. 2015, Tilman et al. 2017), it is also increasingly recognized that habitat alteration and fragmentation are critical ecological and evolutionary drivers in anthropogenic landscapes (Villard & Metzger 2014, Haddad et al 2015). Interestingly, the risk of extirpation from anthropogenic landscapes appears to differ between species (Frishkoff et al. 2014, Edward et al. 2015, Nowakoski et al. 2018) and research is needed to identify the phenotypic traits that allow some species to cope with and succeed in human-dominated contexts.

Dispersal, i.e. the movement from birth to breeding patch (natal dispersal) or between successive breeding patches (breeding dispersal), is a key ecological and evolutionary process. Dispersal provides the demographic supply for population rescuing, habitat (re)colonization (Hanski & Gaggiotti 2004, Bowler & Benton 2005, Gilpin 2012), and range expansion (Travis et al. 2009, Kubisch et al. 2014, Ochocki & Miller 2017). Furthermore, it determines the intensity and direction of gene flow, which has far-reaching consequences for local genetic diversity and adaptive processes (Lenormand 2002, Ronce 2007, Petit & Broquet 2009, Cayuela et al. 2018a). Dispersal is a complex phenotype, partially controlled by genetics and relying on a suite of morphological, behavioral and life history traits that may be subject to joint selection (Saastamoinen et al. 2018). Such associations between dispersal and individual phenotype are called 'dispersal syndromes' and lead to phenotypic specialization within and between populations (Matthysen 2012, Cote et al. 2010, Ronce & Clobert 2012). Dispersal is also influenced by patch and landscape characteristics: individuals are expected to adjust their dispersal decisions according to the fitness prospects of a patch (i.e. 'informed dispersal', Clobert et al. 2009), leading to context-dependent dispersal.

Over the last two decades, an increasing number of studies have suggested that landscape anthropization is an important determinant in dispersal evolution as it affects the balance between fitness benefits and the direct and indirect costs of moving (Bonte et al. 2012) incurred at the different stages of the dispersal process (i.e. emigration, transience, and immigration) (Kokko & Lopez-Sepulcre 2006, Cheptou et al. 2017, Cote et al. 2017). The majority of efforts have been devoted to better understanding the influence of habitat fragmentation on dispersal evolution in anthropogenic landscapes (Cheptou et al. 2017, Cote et al. 2017, Legrand et al. 2017, Atkin et al. 2019). Overall, these studies have reported reduced dispersal propensity or capacity in fragmented landscapes, which

is usually attributed to prohibitive costs during the transition phase across the matrix (Cheptou et al.

2017, Cote et al 2017). This hypothesis is supported by landscape genetic studies, which often report

increased spatial genetic differentiation depending on the harshness of the matrix that separates demes

(Baguette et al. 2013, Cushman et al. 2015). Nevertheless, habitat fragmentation may have a

contradictory effect on the evolution of dispersal: on one hand, it may make transition across the

matrix costlier, but on the other, it may make dispersal profitable due to the increased local extinction

risk caused by heightened demographic stochasticity within severely fragmented landscapes (Ronce

2007, Hanski & Mononen 2011, Cote et al. 2017). The best empirical support of this hypothesis is

perhaps the well-documented selection for dispersal-specialized phenotypes observed in spatially

structured populations (SSPs, Thomas & Kunin 1999) of the Glanville fritillary butterfly subject to

highly fragmented landscapes (Hanski 2011, Hanski et al. 2017). In this unique example,

environmental stochasticity makes dispersal profitable by creating new patches that can be colonized

by dispersers. The prime importance of spatiotemporal patch variability in promoting dispersal is well

supported by a number of theoretical models (Comins et al. 1980, McPeek & Holt 1992, Armsworth &

Rougharden 2005), and is also advanced as the main driver of wing dimorphism observed in insects in

a gradient of patch temporality (Denno et al. 1996).

Habitat alteration in human-dominated landscapes is often associated with shifts in disturbance

regimes (Turner 2010, Newman 2019). For instance, one decade-long worldwide survey revealed a

relatively weak net surface loss of temperate forests, but a high turnover due to forestry practices

(Hansen et al. 2013). Shifts in the disturbance regimes prevailing in habitat remnants could therefore

mitigate or, conversely, magnify the negative effect of habitat fragmentation, depending on their

direction and magnitude. Despite this, apart for aerial dispersal in invertebrates (Denno et al. 1996),

human-induced temporal variation in the spatial distribution of habitat patches has generally been

overlooked when considering dispersal evolution in anthropogenic landscapes. A full appraisal of the

effect of anthropogenic disturbance on dispersal should not only examine whether a dispersal pattern

emerges in a landscape, but whether it gives rise to a genetic footprint throughout successive

generations – and last but not least, to what extent this involves a specialized phenotype. Considering

all the facets of this issue is not a simple task (Kokko & Lopez-Sepulcre 2006, Broquet & Petit 2009,

Ronce & Clobert 2012), and to our knowledge has not yet been investigated in vertebrates.

To address this gap, this study examined how human-driven variation in habitat patch turnover

affects dispersal rates and distances, determines dispersal-related phenotypic specialization, and drives

neutral genetic variation in spatially structured populations. We studied this issue in an early

successional amphibian, the yellow-bellied toad (*Bombina variegata*), a species that reproduces in

small waterbodies with a short hydroperiod occurring either in (virtually) undisturbed or

anthropogenic environments (Warren & Büttner 2008; Cayuela et al. 2011, 2015b). In riverine

environments with negligible human activity, the species' habitat patches are groups of rocky pools

that result from long-term geomorphological processes alongside riverbanks (riverine SSPs; Cayuela

et al. 2011). This results in a negligible patch turnover rate and makes patches available and
predictable far beyond a toad's lifespan. In contrast, in harvested woodlands, habitat patches consist of
groups of ruts made by logging vehicles that may appear and disappear yearly as a result of the
combined effects of logging operations and rapid natural silting (we refer to these as logging SSPs
hereafter). This leads to a high patch turnover rate and makes patch location and availability more
unpredictable at the scale of a toad's lifespan (Cayuela et al. 2016a, 2016b). Previous studies have
highlighted demographic differences in SSPs from the two environments and found that individuals in
logging SSPs have a faster life history (i.e. a shorter lifespan and higher fecundity; Cayuela et al.
2016a), experience earlier senescence (Cayuela et al. 2019b), and display higher breeding dispersal
probability (Cayuela et al. 2016b) than individuals in riverine habitats. In this study, our first step was
to quantify dispersal probability and distance throughout an individual's lifetime, as natal dispersal
was lacking in previous studies and a review of recent literature suggested that natal and breeding
dispersal patterns can strongly differ in amphibians (Cayuela et al. 2018c). We expected that (1) both
natal and breeding dispersal rates and distances would be higher in logging than in riverine SSPs. In a
second step, we analyzed how patch turnover and related dispersal costs and benefits affected SSP
dynamics and long-term viability using simulations based on published demographic rates. We
hypothesized that (2) dispersal and context-dependent immigration (i.e. depending on patch age)
allows the long-term persistence of logging SSPs. In a third step, we used common garden
experiments to investigate how patch turnover determines dispersal syndromes and may act as a
selective agent on phenotypic specialization in riverine and logging SSPs. We expected (3) toadlets
from logging SSPs to have behavioral traits (i.e. high exploration propensity and boldness) and
morphological traits (i.e. long hind limbs) that generally facilitate dispersal in amphibians (reviewed in
Cayuela et al. 2018c). In a fourth step, we examined how human-driven variation in patch turnover, by
affecting neutral genetic diversity and gene flow, leads to contrasting genetic footprints over the longer
term in riverine and logging SSPs. As genetic differentiation results from the combined effects of
genetic drift and gene flow (Broquet & Petit 2009, Cayuela et al. 2018a), we selected two SSPs per
landscape type in order to determine the relative contribution of each of these drivers. As high
dispersal is expected to increase gene flow and decrease the local effects of genetic drift, we expected
(4) a larger effective population size as well as a lower level of inbreeding and intra-patch relatedness
in logging than in riverine SSPs. We also expected that (5) after controlling for SSP size and landscape
connectivity, higher gene flow would lead to lower genetic structure and weaker genetic Isolation-By-
Distance (IBD) pattern in logging than in riverine SSPs.
157
158   **Materials and methods**
159
160   *Study area and sampled populations*
161

The study was conducted in 8 SSPs in eastern France, three in riverine environments (R1, R2, and R3) and five in logging environments (L1, L2, L3, L4 and L5). The SSPs were chosen according to technical constraints or to minimize bias at each stage of the study – our choices are explained below. The distance separating SSPs from each other varied from 20–500 km (**Fig. S1**).

Table 1. *Environmental characteristics of the four SSPs (L1, L2, R1 and R2): patch persistence over time, environment type (logging vs riverine), patch isolation (mean distance in meters between two pond networks, and associated variation coefficient) and patch size (mean number of ponds within a patch and associated variation coefficient).*

SSP	Patch persistence	Environment	Patch isolation	Patch size
L1	1 to 10 years	logging activity	2864.50 (57%)	3.70 (70%)
L2	1 to 10 years	logging activity	4424.64 (53%)	5.64 (88%)
R1	>30 years	natural erosion	676.57 (61%)	4.52 (105%)
R2	>30 years	natural erosion	501.62 (61%)	4.71 (93%)

Table 2. *Survey design characteristics for the four SSPs (L1, L2, R1 and R2) considered in the study: study period, survey duration, number of capture sessions performed over the survey period, total number of captures, total number of individuals identified during the survey and total number of sampled patches (pond networks).*

SSP	Study period	Survey duration	Capture sessions	Number of captures	Number of individuals	Number of sampled patches
L1	2000–2008	9 years	29	953	445	28
L2	2012–2016	5 years	15	16477	12192	189
R1	2010–2014	5 years	13	4747	1003	14
R2	2010–2014	5 years	13	3984	769	8

Dispersal patterns throughout toad lifespan in riverine and logging environments

Studied populations

We quantified natal and breeding dispersal rates and distances in four SSPs (L1, L2 and R1, R2; see maps **Fig. S1** and **Fig. S2**) for which breeding rate dispersal had been previously estimated (Cayuela et al. 2016a, 2016b). The environmental characteristics of the four SSPs and the details regarding the survey design are presented in **Table 1** and **Table 2** respectively. The number of individuals captured each year is presented in **Supplementary material**, **Table S1**. A detailed description of the four SSPs can also be found in two previous studies (Cayuela et al. 2016a, 2016b). The number of patches (defined as a group of ruts) occupied by each SSP ranged from 8 to 189. Each SSP was monitored for a period of at least five years in one to five capture sessions per year that were usually between two weeks to one month apart. At each capture session, all the patches were sampled in the daytime and

toads were captured by hand or dipnet. Based on previous studies (Cayuela et al. 2016a, 2016c), we

considered three life stages: juveniles (i.e. post-wintering metamorphs), subadults (two-year-old

immature animals) and adults (i.e. breeders, three years old or more). This resulted in a total dataset of

12,721 individual capture–recapture (CR) histories.

<u>Naïve dispersal kernels</u>

We first estimated a dispersal kernel based on distances recorded during sampling using a lognormal

distribution for each population and life stage (juvenile, subadult, and adult). This allowed us to

visualize the form of the kernel from raw data before building complex CR models.

<u>The structure of the multievent model</u>

For the needs of our study, we extended the CR multi-event model proposed by Lagrange et al. (2014),

which allows estimating survival ($\square$) and dispersal (ψ) in numerous sites. By omitting site identity and

distinguishing between individuals that stay and individuals that move, this model circumvents the

computational issues usually encountered in standard multi-site CR models when the number of sites

is large (Lebreton et al. 2009). Lagrange's model includes states that incorporate information about

whether an individual is occupying at t the same site as the one occupied at $t - 1$ ('S' = stayed) or not

('M' = moved). The model also includes information about whether the individual was captured ('+')

or not ('o') at $t - 1$ and t. Recently, Tournier et al. (2017) extended Lagrange's model by breaking

down dispersal (ψ) into distinct parameters of departure (ε) and arrival (α). This new parameterization

allows the estimation of the proportion of individuals arriving in sites of different quality or located at

different distances from the source site.

We adapted this parameterization for our study to consider states incorporating information

about individual capture ('+' and 'o') at $t - 1$ and t as well as movement status. We also included

states with information about the individual's age class: juvenile ('j'), subadult ('s') and adult ('a').

Additionally, we incorporated information about the Euclidian distance covered by dispersers between

the departure and arrival patch using three distance classes: '1' = 100–800 m, '2' = 800–1500 m, '3' >

1500 m. This led to the consideration of 37 states in the model (**Fig.1** and **Fig.2**). For example, an

individual +jS+ was captured at $t - 1$ and t, was a juvenile, and remained in the same patch between $t -$

1 and t. An individual +sM1+ was captured at $t - 1$ and t, was a subadult, did not occupy the same

patch as at $t - 1$, and arrived in a patch located at a distance 100–800 m from the source patch. We

distinguished 16 events, which were coded in an individual's capture history and reflect the

information available to the observer at the time of capture (**Fig.2**).

When captured for the first time, the state of an individual could be ojS+, osS+ or oaS+. We

then considered five modeling steps in which the information of the state descriptor was progressively

updated: survival (□), departure (ε), arrival (α), age transition (δ) and recapture (*p*). Each step was

conditional on all previous steps. In the first step, we updated information about survival. An

individual could survive with a probability of □ or die (D) with a probability of 1 − □. This led to a

matrix with 37 states of departure and 7 intermediate states of arrival (**Fig.1**). Survival probability

could differ between age classes by allowing differing values for □ in lines 1–12, 13–24 and 25–36 of

the matrix. In the second modeling step, departure was updated. Individuals could move (M) from the

site they occupied with a probability of ε or stay (S) with a probability of 1 − ε. A matrix of 7

departure states and 13 arrival states was considered (**Fig.2**). Departure probability could differ

between age classes by allowing differing values for ε in lines 1–2, 3–4 and 5–6 of the matrix. In the

third step, we updated the arrival information. An individual that moved could arrive in a patch located

in the first two distance classes (1 or 2) from the source patch with a probability of α, or arrive in a site

located in the third distance class (3) with a probability of 1 − α. This resulted in a matrix with 13

departure states and 25 arrival states (**Fig.2**). Arrival probability could differ between age classes by

allowing different values for α in lines 2–4, 7–8 and 11–12 of the matrix. In the fourth step, the

information about age was updated. An individual could reach the next age class (j, s or a) with a

probability of δ or remain in the previous age class with a probability of 1 − δ, resulting in a transition

matrix with 25 states of departure and 25 states of arrival (**Fig.2**). The adult individuals (a) were forced

to stay in their age class. In the fifth and last step, recapture was updated (**Fig.2**). An individual could

be recaptured with a probability of *p* or missed with a probability of 1 − *p*, resulting in a transition

matrix with 25 states of departure and 37 states of arrival. The recapture probability could differ

between age classes by allowing different values for *p* in lines 1–8, 9–16 and 17–24 of the matrix. The

last component of the model linked events to states. In this specific situation, each state corresponded

to only one possible event (**Fig.2**).

<u>Biological scenarios in the E-SURGE program</u>

The parameterization was implemented in the E-SURGE program (Choquet et al. 2009). The datasets

for the four SSPs considered in our study were analyzed separately, as the number of study years and

capture sessions for these populations varied. Competing models were ranked through a model-

selection procedure using Akaike information criteria adjusted for a small sample size (AICc) and

AICc weights. Following the recommendation of Burnham & Anderson, we performed model

averaging when the AICc weight of the best-supported model was less than 0.90. The models had a

robust design structure (Pollock 1982). As in previous studies of *B. variegata*, the survival probability

was fixed at one between secondary sessions (Cayuela et al. 2016a, 2016c). The robust design

structure allowed both intra-annual and inter-annual dispersal to be considered. Our hypotheses

concerning recapture and state–state transition probabilities were tested using the general model

[□(AGE), ε(AGE), α(AGE), δ(.), *p*(AGE + Y)], which included two effects: (1) the three age classes

(AGE) coded as states in the model, and (2) year-specific variation (Y). The notation (.) indicated that

the parameter was held constant. We tested whether survival ($\square$) and departure probabilities (ε) varied

between age class (AGE). Moreover, we hypothesized that the probability of arriving in a patch

depends on age (AGE), and on the Euclidean distance between patches (the distance classes were

incorporated as states in the model). Recapture probability was expected to differ between age classes

(AGE) and years (Y). We tested our expectations about the model parameters in a stepwise fashion.

From this general model, we tested all the possible combinations of effects and ran 16 competing

models.

Fig.1. *Model structure: matrices of initial states and state transitions (survival, departure, arrival and*

age transition). In the transition matrix, the rows correspond to time t – 1, the columns to time t, and

whenever a status element is updated to its situation at t, it becomes bold and stays bold throughout
the following steps.

Fig.2. *Model structure: state transitions (recapture) and events (field observations). In the transition*
matrix, the rows correspond to time t – 1, the columns to time t, and whenever a status element is
updated to its situation at t, it becomes bold and stays bold throughout the following steps.

Simulating the effect of patch turnover and dispersal on SSP dynamics and long-term persistence in
logging contexts

We simulated population trajectories based on different scenarios to investigate the effects of patch

turnover and dispersal costs on the dynamics and long-term persistence of SSPs in habitats subjected

to logging. Adopting the most realistic lifecycle for the yellow-bellied toad (see 'Results', **Fig.4**)

determined in previous studies (e.g. Cayuela et al. 2015, 2018), we used a three age-class (juveniles,

subadults, and adults), female-dominant, prebreeding Leslie matrix (Caswell 2001) (see 'Results',

Fig.4). We used the demographic parameters of a riverine SSP (R1; see Cayuela et al. 2016), which

was considered a reference population whose demographic parameters have not been altered by

dispersal costs (survival in logging SSPs is lower than in riverine SSPs, likely due to dispersal costs,

Cayuela et al. 2018b). Both prebreeding survival probability (juvenile survival, $S1 = 0.70$; subadult

survival, $S2 = 0.77$) and adult survival probability ($S3 = 0.92$) were included in the Leslie matrix.

Fecundity F was possible only for adult females and consisted of estimated recruitment: that is, the

number of recruited juvenile females at t per breeding female at $t - 1$ ($F = 0.52$, Cayuela et al. 2016).

As Boualit et al. (2019) found that juvenile recruitment was higher in newly created and disturbed
patches than in old, undisturbed patches in logging SSPs, we specified that F decrease linearly (-5%
per year) with the age of the patch. Furthermore, we also considered the possibility that females may
skip breeding opportunities; in the riverine SSP R1, a previous study showed that the probability of
females skipping breeding (B) was 0.15 (Cayuela et al. 2016). As in Cayuela et al. (2018, 2019c), we
considered demographic stochasticity for survival, fecundity and skipping breeding. For each year,
demographic parameter values were randomly sampled in a Gaussian distribution centered on mean
parameter estimates, and standard deviation was inferred from two previous studies conducted on this
species (Cayuela et al. 2016a, 2018b). The standard deviation values were: 0.05 for S1, 0.03 for S2,
0.01 for S3, 0.02 for F, and 0.02 for B.

Three patch turnover scenarios were considered. In scenario 1 (high turnover), a patch
disappeared three years after its creation, in scenario 2 (medium turnover) a patch disappeared after six
years, and in scenario 3 (low turnover) it disappeared after nine years. These scenarios correspond to
the range of patch turnover in logging SSPs reported by forest managers (Eric Bonnaire, *unpublished*
data), variation that depends on local management policies and the frequency of forest harvesting
operations. We also considered three dispersal scenarios. In scenario 1 (no dispersal), individuals were
not able to escape and died when a patch disappeared. In scenario 2 (dispersal with random
immigration), individuals could disperse to escape the disappearance of a patch or could disperse by
choice (i.e. when the patch remained available). The immigration was random between the patches of
the metapopulation and was not influenced by the age of the patch. In scenario 3 (dispersal with
informed immigration), individuals could disperse in response to patch disapperance or by choice.
Based on an assessment of Boualit et al. (2019), we considered that immigration was not random and
that immigration probability linearly decreases with patch age (i.e. a loss of 5% per year). In scenario
2 and 3, in which dispersal was possible, we considered two subsets of scenarios: in subset 1 (non-
costly dispersal), individuals did not incur any survival loss when they dispersed. In subset 2 (costly
dispersal), we considered that survival loss related to dispersal could be low (-5% of survival),
medium (-10%), or high (-15%). As in Cayuela et al. (2018), we made the assumption that survival
loss was similar across life stages (i.e. juvenile, subadult, and adult).

Each simulation began with 30 breeding patches over which 1,000 individuals were randomly
scattered. The number of individuals in each age class was obtained through the stable stage
distribution provided by the three age-class Leslie matrix. Then for each time step (a 1-year interval),
we simulated the change in patch availability. We considered that five new patches were created each
year. As the patches disappeared in a deterministic way when they reached the age defined in the
scenario, the number of available patches remained constant over time (except for the few first years).
We simulated the number of individuals in each age class occupying each patch. To do this, we
separately considered patches reaching the age of disappearance versus those that did not disappear. In
the latter, the number of individuals at $t + 1$ given the number of individuals at t was predicted by the

Leslie matrix using the survival probability of individuals occupying an available patch (reported in

the reference population R1). To be as realistic as possible, we used demographic stochasticity

(fecundity was thus randomly sampled from a Poisson distribution, and survival from a binomial

distribution), as in Cayuela et al. (2018). For patches that disappeared, we applied the same procedure,

but using the survival probability (affected or not by dispersal cost, depending on the scenario) for

individuals occupying patches that subsequently disappeared. Surviving individuals from lost patches

were then randomly spread over the available patches at $t + 1$ in the 'dispersal with random

immigration' scenario, or they were preferentially distributed in new patches in the 'dispersal with

informed immigration' scenario. In all dispersal scenarios, we fixed the elective dispersal probability

(dispersal when the patch did not disappear) D at 0.15, which was consistent with the annual dispersal

rate reported in logging SSPs in the study. We also considered demographic stochasticity in elective

dispersal (the standard deviation value was 0.05). The modeled population was monitored for 100

years. We did not remove the first few years of the simulation, when the number of patches

progressively increased since none were old enough to disappear yet, as these had virtually no impact

on our results (Cayuela et al. 2018). We performed 1,000 simulations for each scenario. At each time

step, we monitored the number of adults in the entire SSP as well as the proportion of simulations in

which the SSP went extinct (the extinction probability).

Phenotypic specialization in riverine and logging environments

<u>Study populations</u>

To compare the morphology and behavior of toadlets in riverine SSPs with those of logging SSPs, we

used a common garden experiment. This involved collecting between 8 and 15 egg clutches (hereafter

referred to as 'family') in three SSPs in each landscape type (riverine = R1, R2, R3, and logging = L3,

L4, L5; **Fig.S1**). These were selected to minimize spatial proximity between SSPs belonging to the

same landscape type and therefore to avoid potential confounding effects resulting from spatial

autocorrelation in environmental conditions. Five siblings per family were randomly chosen just after

hatching and individually reared under controlled laboratory conditions.

<u>Rearing protocol</u>

The egg clutches were carefully transported to the laboratory, where they were individually placed in

aquariums (32×17 cm, height 15 cm) with aged tap water equipped with an oxygen pump. The

aquariums were placed in a climatic room with a light/dark cycle of 18:6, corresponding to the natural

light/dark cycle in the area in the summer, with an ambient temperature varying from 21.5 to 23.5°C.

Embryo development ended 2–5 days after the arrival of the egg clutches at the laboratory. After

hatchling, when tadpoles reached Gosner stage 24–25 (active swimming, external gill atrophy)

(Gosner 1960), five siblings per family were randomly chosen and placed individually into plastic

containers (14×8.5 cm, height 13.5 cm) filled with previously aged tap water and aerated in a tank.

The plastic containers were distributed in a predetermined random pattern around the climatically

controlled room. The water was replaced every three days. The tadpoles were fed every day with 150

mg of cooked lettuce, providing ad libitum feeding. At Gosner stage 44–45 (tail atrophy, mouth

posterior to eyes), feeding was terminated, and the water was drained and replaced with a dampened

sponge placed in the bottom of the container. The sponge and the walls were sprayed with aged tap

water every two days. The individuals were kept until their complete metamorphosis (Gosner stage

46) and were then subjected to behavioral assays.

<u>Experimental arenas</u>

The behavioral assays took place in an arena (70 cm in diameter) made of polyethylene terephthalate,

with a central shelter and a 'desiccation' obstacle between this and the 12 possible exits (**Fig. S4**). At

the center of the arena, we placed a removable, cylindrical (9 cm diameter, height 90 cm), opaque,

covered chamber ('refuge chamber' hereafter). The cylinder had a circular opening (3 cm diameter)

covered by a lid. Around the interior edge of the arena, we installed a pit (width 10 cm, depth 0.3 cm)

filled with a desiccating mixture of sand and highly active silica gel powder in a weight ratio of 0.8:0.2

('desiccant zone' hereafter). The arena wall included 12 doors placed at regular intervals around the

entire circumference. The arena was confined in an enclosed iron chamber (150×17 cm, height 175

cm) over which a dark opaque sheet was placed to limit potential acoustic and visual interference

during behavioral trials.

<u>Behavioral assays</u>

At metamorphosis, each toadlet was subjected to a behavioral assay to quantify their neophobia or

exploratory behavior. The behavioral tests were conducted in the circular arena described above.

Before each trial, the toadlet and a dampened sponge (already present in the toadlet's rearing

container) were gently transferred into an opaque circular release box that was then placed at the

center of the arena. The dampened sponge was considered a known object, making the refuge chamber

more familiar than the rest of the experimental device. Following previous studies on anuran

behavioral syndromes (reviewed in Kelleher et al. 2018), neophobia was quantified as the latency time

to enter a novel environment (i.e. the time delay to leave the familiar refuge chamber: BEHAV1).

Exploration propensity was assessed using two variables: the latency time to enter a novel but harsh

environment (i.e. to reach the desiccant zone after leaving the refuge chamber; BEHAV2), and the

latency time to travel the harsh environment and get out of the arena (BEHAV3). The behavior was
recorded for 30 min (1800 s) using a digital camera (Sony DCR-SX34).

Extraction of behavioral variables from the videos

The videos were analysed using the BEMOVI R package (Pennekamp et al. 2015) to reconstruct the
movement trajectory of the toadlet in the arena, and to extract a series of behavioral variables from
this. First, the videos were standardized to a length of 29 min (the minimum duration available for all
individuals) removing the initial 60 s and the extra time at the end, if any. Then the videos were
converted to a format suitable for analysis in BEMOVI: rectangular pixels (720 x 576) were converted
into square pixels (1024 x 576), color information was converted into 256 gray levels, frames per s
were decreased from 25 to 5 to limit memory allocation requirements, and the videos were saved as
AVI files. These operations were performed using FFMPEG software (ffmpeg.org).
BEMOVI was then run with the following parameters: black and white threshold (40) to
discriminate the toadlet from the background; minimum size (20) and maximum size (150),
corresponding to the size range of the toadlets; link range (7500 frames) to allow any duration of the
'disappearance' of the toadlet from the video (e.g. when it was in the refuge) while still considering it
as a single movement trajectory, and "disp "(100 pixels); in BEMOVI, disp is the maximal distance
covered by the toadlet from one frame to the next, corresponding to 0.2s here. This resulted in a
database with the movement trajectory of each toadlet, i.e. its X–Y position at each time step. This
position was compared to the distance from the center of the arena to determine which zone (refuge
chamber, normal zone or desiccant zone) the toadlet was in at each time step. The results were
checked for errors in toadlet positioning, which were due to varying light conditions, usually at the
beginning of the videos. In a final step, we computed several behavioral variables for each toadlet
from this positioning information.

Statistical analyses

We used linear mixed models to test whether morphological traits (body size, body condition and
relative hind limb size) of toadlets differed according to the landscape type of origin (logging vs
riverine). Each morphological trait was treated as a dependent variable, and the landscape type was
introduced as a fixed explanatory term in the model. In the case of body condition and relative leg
size, body size and its interactive effect with the landscape type were also introduced as adjustment
covariates in the fixed part of the model. Both the SSP of origin and the clutch were introduced as
random effects in the model. We also allowed heterogeneity of variance between landscape types by
allowing a separate estimation of the residual variance for each landscape type. The estimation method
was based on restricted maximum likelihood. Variance heterogeneity was first checked using the

likelihood ratio test and removed if non-significant. The significance of each explanatory term was

examined with a non-sequential F test based on the Kenward-Roger method to approximate the

denominator df (Littel et al. 2006). In the case of covariance analyses (body condition and relative leg

size), the interactive effect was discarded if non-significant to obtain the final model. All

morphological variables were standardized using Z transformation before the analyses, as

recommended by Schielzeth (2010).

We tested whether the behavioral variables BEHAV1 and BEHAV2 varied according to the

landscape type (i.e. logging vs riverine) using a generalized linear mixed model. Each latency time

variable was treated as a dependent variable using a Poisson distribution. The landscape type was

introduced as an explanatory term in the fixed part of the model. We also introduced individual body

size and its interactive effect with the landscape of origin as adjustment covariates in the fixed part of

the model since both locomotion skill and exploratory performance can vary according to individual

size (reviewed in Kelleher et al. 2018). For the analyses of morphological traits, both the SSP of origin

and the clutch were introduced in the model as random effects. Furthermore, a scale parameter was

also introduced to handle data overdispersion and to obtain a corrected statistical test using a quasi-

likelihood approach (McCullagh & Nelder 1989). The estimation method was based on restricted

pseudo-likelihood optimization, and the significance of each explanatory term was examined using the

same methodology as for the morphological analyses. Non-significant terms were successively

removed to obtain the final model, and least square means were used to estimate the difference in

latency time variables according to the landscape of origin.

As the third behavioral variable (BEHAV3) was right censored, it was analyzed using a

proportional hazards mixed-effects model (i.e. a frailty model based on a Cox model, PHREG

procedure, SAS Institute Inc. 2012). We tested whether newborn individuals originating from logging

systems were more prone to exit the assay arena than those from riverine systems. It is not possible to

handle multiple random factors in such a model, so we took into account only the family effect since

this was found to be significant for the other behavioral variables but not for the SSP of origin. The

landscape of origin, the body size and their interactive effect were introduced as explanatory terms.

Parameters were estimated using partial likelihood estimation, and the significance of explanatory

terms was assessed using non-sequential chi-square tests.

Neutral genetic variation in riverine and logging environments

<u>Study populations</u>

We examined neutral genetic variations within two SSPs in riverine environments (R1 and R2) and

two SSPs in logging environments (L3 and L4) (see map in **Fig. S3**) using 15 polymorphic

microsatellite markers (described and tested in Cayuela et al. 2017a). The four SSPs were selected

according to the following criteria: (1) SSPs embedded in a relatively continuous forested matrix to

avoid any confounding effect of matrix composition on gene flow (**Fig. S3**) – woodland is generally

considered highly favorable for the movement of forest amphibians (Cushman 2006a) such as *B.*

variegata (Cayuela et al. 2015b); and (2) two small (R1 and L3) and two large SSPs (R2 and L4) to

control for genetic drift. The number of patches and DNA sampled per SSP are given in **Table S2**. We

used the protocol described in Cayuela et al. (2017) for DNA extraction and amplification, individual

genotyping, and allele scoring.

Estimating basic genetic metrics

We examined basic assumptions (i.e. detection of null alleles, Hardy–Weinberg equilibrium) and

descriptive statistics (mean number of alleles per locus N_a; effective number of alleles per locus N_{ef};

observed heterozygosity H_o; and expected heterozygosity H_e) of the genetic diversity within each

SSP. N_a, N_{ef}, H_o and H_e were calculated using GENALEX (Peakall & Smouse 2006). The inbreeding

coefficient F_{is} and the Hardy–Weinberg deviation were assessed with GENEPOP 4.1 (Rousset 2008).

The detection of null alleles was performed using the program MICRO-CHECKER 2.2.3 (Van

Oosterhout et al. 2004). Two microsatellite markers, Bomvar_Cons470 and bv11.7, were discarded

from the subsequent analyses due to the presence of null alleles in the four SSPs. All descriptive

statistics are provided in **Table S3**.

Estimating relatedness, inbreeding, and effective population size

We estimated relatedness and individual inbreeding using COANCESTRY v1.0.1.8 (Wang 2011). We

performed simulations to identify the best relatedness and inbreeding estimator for our combined

datasets; these consisted of 1,000 dyads spread equally across six categories of relatedness: parent–

offspring (rxy = 0.5), full siblings (rxy = 0.5), half siblings/avuncular/grandparent–grandchild (rxy =

0.25), first cousins (rxy = 0.125), second cousins (rxy = 0.03125), and unrelated (rxy = 0). According

to our simulations, the best rxy estimator for relatedness analysis was DyadML, which showed a

strong correlation of 0.74. We used linear mixed models to evaluate how SSP size (small: R1 and L4;

large: R2 and L3) and the type of environment (logging vs riverine) affected individual inbreeding and

intra-patch relatedness. An SSP's size and environment type were coded as fixed effects (we

considered an interaction between the two factors), whereas patch identity was coded as a random

effect. We then estimated the effective population size of the four SSPs using the linkage

disequilibrium method implemented in Ne Estimator v2.1 (Do et al. 2014).

Clustering approach

We described the genetic differentiation between patches within each SSP using an assignation

method based on the Bayesian clustering algorithm implemented in the software STRUCTURE

(Pritchard et al. 2000). Specifically, we estimated the most likely number of genetic clusters (K)

contained in each SSP following the hierarchical approach proposed by Balkenhol et al. (2014) to

detect additional substructures within clusters. The STRUCTURE program was run with the

admixture model, with a burn-in period of 100,000 repetitions, and 100,000 subsequent MCMC

repetitions. The K values were tested ranging from 1 to 10 and analyses repeated 10 times for each

value. We used STRUCTURE HARVESTER (Earl 2012) to summarize the results, determining the

optimal K value using both log-likelihood plots and the delta K statistic (Evanno et al. 2005). We

followed the hierarchical approach proposed by Coulon et al. (2008) to test for additional population

substructures within clusters. Accordingly, these analyses were then repeated for each inferred

population cluster separately until the optimal K value was 1 (meaning that no additional structure was

found within clusters). To map the spatial distribution of the different clusters, the individual ancestry

values were averaged across the ten STRUCTURE runs using CLUMMP software (Jakobsson &

Rosenberg 2007). The 'greedy' algorithm in the CLUMMP program was then used to assign the

individuals to the cluster in which they showed the highest Q-values.

<u>Gradient analyses</u>

We used direct gradient analyses (Prunier et al. 2015) to test whether spatial genetic differentiation

was lower within logging than riverine SSPs while controlling for potential differences in the

functional connectivity prevailing within each SSP. To do this, pairwise genetic distances between all

individuals from each SSP were computed using the Bray-Curtis percentage dissimilarity measures

(Cushman et al. 2006b, Legendre & Legendre 2012), and were then standardized separately for each

SSP. Since by design all SSPs were embedded in a relatively homogeneous forested matrix, we did not

control for the effect of land cover type in these analyses. Rather, we focused on the effect of

topographic roughness ('slope' effect) and the hydrological network topology ('network' effect) since

both these landscape features have often been reported to affect gene flow in amphibians (Lowe 2003,

Grant et al. 2010).

For each SSP, resistance layers were produced using ArcGIS Pro. Minimum geographical

bounding of each SSP was determined and extended to a distance of five kilometres to avoid any edge

effects. The elevation raster and hydrographic network shapefile, available through the French

National Institute for Geographical and Forest Information database (BD ALTI®, and BD TOPO®),

were extracted for SSP surface areas. To determine the percentage of steepness raster maps, the slope

tool was used. Due to the spatial resolution imposed by elevation data, all layers were converted to

raster format and homogenized at a spatial resolution of five meters.

Four resistance maps were thus constructed for each SSP. The first map included only the
effect of geographic distance (i.e. isolation by distance, hereafter geographic resistance); it was based
on data extended to include a 5-km buffer and was assigned a uniform resistance of 1 unit for all
pixels. The second map included the topographic roughness (slope), where the resistance of each pixel
was a linear function of the steepness ($resistance = 1 + degree\ of\ steepness$), thus corresponding to
the effect of both geographic distance and topographic roughness. The third map included the
hydrological network topology (network), where the resistance was assigned to 1 unit for all pixels
situated in the hydrological network and to 10 units otherwise. The fourth map included both the slope
effect and the network effect. Pairwise resistance between all patches was then computed from these
four resistance maps using the circuit theory in Circuitscape V4.0 (McRae & Beier 2007).

Linear mixed models for pairwise distance matrices were then used to assess the effect of the
geographic distance between patches (i.e. the geographic resistance) on the genetic distance between
toads within each SSP while controlling for the effect of topographic roughness and/or the
hydrological network topology. The non-independence between pairwise distances was taken into
account in the covariance structure of the models. Specifically, we used the method proposed by
Clarke et al. (2002) using a Toeplitz(1) as a covariance structure to specify the non-independence of
pairwise genetic distance according to the patches of origin (see Selkoe et al. 2010, van Strien et al.
2012 for application on pairwise Fst distances). We used the extension presented in Prunier et al.
(2013) for application on individual genetic distance in spatially hierarchized sampling schemes as is
the case in our study. This resulted in two covariance parameters for each SSP, one for the patch
random effect and the other for the individual random effect. Since both slope and network resistance
were highly correlated to geographic resistance and as our main aim was to estimate the effect of
geographic resistance on genetic distance while controlling for the effect of functional connectivity,
we first regressed each of these effective resistances on each geographic resistance using simple linear
regressions to obtain uncorrelated effective resistance respectively related to the slope effect, the
network effect and their combined effect. The relative validity of each alternative landscape
hypothesis (i.e. including the effect of the slope or of the stream network or both or neither on the
genetic distances) was evaluated using weighted AIC (Waic), and the model-averaged estimate of the
beta weight associated with the effect of geographic resistance on genetic distance was computed for
each SSP.
589
590   **Results**
591
592   *Dispersal patterns throughout toad lifespan in riverine and logging environments*
593

Model averaging estimates (see **Table S4** for survival and recapture probabilities) indicated that both

natal and breeding dispersal rates were high in logging environments (around 20% per year, **Fig.3**). In

contrast, in riverine environments, natal dispersal rates were null and breeding dispersal rates were

very low (less than 5% per year, **Fig.3**). This higher dispersal propensity in logging than in riverine

habitats is even more remarkable given that the mean inter-patch distance is three times farther in the

former than the latter (Table S1). The few adults that did disperse in riverine environments covered

shorter distances: the median distance was 168 m in R1 and 189 m in R2, while the maximal distance

was 455 m and 378 m respectively. This result was further confirmed by multi-event CR models

showing that 100% of dispersal occurred over distances ranging from 100 to 800 m in riverine SSPs.

In logging environments, dispersers covered substantially longer distances: the median distance was

431 m in L1 and 568 m in L2, while the maximal distance was 3810 m and 4529 m respectively. This

result was supported by both the observed dispersal kernel and the multi-event model estimates for

each life stage (see **Fig.3**). While these results indicate extremely contrasting dispersal regimes

between logging and riverine landscapes, we also found substantial variations in the dispersal patterns

in the two logging systems. First, while the inter-annual dispersal rate was similar in both logging

systems, the intra-annual dispersal rate was substantially higher in L1 than L2 (i.e. L1 > 10 % and L2

<5%, **Fig.3**; for model selection procedure see **Table 3** and **Table 4**). Second, the dispersal kernels

show a clear leptokurtic distribution decreasing with age in L2, suggesting a large demographic weight

of natal dispersal in this SSP. This was not the case in L1, in which leptokurtosis was more reduced in

juveniles than in adults.

616
617 ***Fig.3.*** *Natal and breeding dispersal rates and distances in logging (L1 and L2) and riverine SSPs (R1*
618 *and R2) (A–B). Inter-annual (A) and intra-annual (B) dispersal probabilities are higher in SSPs in*
619 *logging landscapes than in riverine SSPs, regardless of the ontogenetic stage (juvenile, J; subadult, S;*
620 *adult, A). Natal and breeding dispersal distances in logging SSPs (L1 and L2) (C–I). Dispersal event*
621 *frequency decreases with the Euclidean distance between breeding patches at the ontogenetic stages.*
622

r	Model	k	Dev.	wAICc	AICc
SSP L1					
1	□(A), ε(.), α(A), p(A + Y)	26	5161.06	0.81	5214.05
2	□(A), ε(A), α(A), p(A + Y)	28	5160.05	0.17	5217.19
3	□(A), ε(.), α(A), p(Y)	24	5173.01	0.02	5221.86
4	□(A), ε(A), α(A), p(Y)	26	5172.00	0.00	5224.99
5	□(.), ε(.), α(A), p(A + Y)	24	5188.40	0.00	5237.25
6	□(.), ε(A), α(A), p(A + Y)	26	5187.39	0.00	5240.38
7	□(.), ε(.), α(A), p(Y)	22	5199.87	0.00	5254.58
8	□(.), ε(A), α(A), p(Y)	24	5198.86	0.00	5247.70
9	□(A), ε(.), α(A), p(A)	18	5261.02	0.00	5297.50
10	□(A), ε(A), α(A), p(A)	20	5260.01	0.00	5300.60
11	□(A), ε(.), α(A), p(.)	16	5268.73	0.00	5301.11
12	□(A), ε(A), α(A), p(.)	18	5267.72	0.00	5304.20
13	□(.), ε(.), α(A), p(A)	16	5289.52	0.00	5321.90
14	□(.), ε(.), α(A), p(A)	14	5295.71	0.00	5324.01
15	□(.), ε(A), α(A), p(A)	18	5288.51	0.00	5324.99
16	□(.), ε(A), α(A), p(.)	16	5294.70	0.00	5327.08
SSP L2					
1	□(A), ε(A), α(A), p(A + Y)	25	62699.27	0.74	62749.34
2	□(A), ε(.), α(A), p(A + Y)	23	62705.41	0.23	62751.47
3	□(A), ε(A), α(A), p(Y)	23	62738.78	0.02	62784.85
4	□(A), ε(.), α(A), p(Y)	21	62744.92	0.00	62786.98
5	□(.), ε(A), α(A), p(A + Y)	23	62779.96	0.00	62826.03
6	□(.), ε(.), α(A), p(A + Y)	21	62786.10	0.00	62828.16
7	□(.), ε(A), α(A), p(Y)	21	62852.52	0.00	62894.58
8	□(.), ε(.), α(A), p(Y)	19	62858.66	0.00	62896.71
9	□(A), ε(A), α(A), p(A)	21	63103.58	0.00	63145.64
10	□(A), ε(.), α(A), p(A)	19	63109.72	0.00	63147.77
11	□(A), ε(A), α(A), p(.)	19	63144.67	0.00	63182.72
12	□(A), ε(.), α(A), p(.)	17	63150.81	0.00	63184.85
13	□(.), ε(A), α(A), p(A)	19	63185.43	0.00	63223.48
14	□(.), ε(.), α(A), p(A)	17	63191.57	0.00	63225.61
15	□(.), ε(A), α(A), p(.)	17	63272.39	0.00	63306.42
16	□(.), ε(.), α(A), p(.)	15	63278.53	0.00	63308.55

629
630
631
632

Table 4. *Multievent models and selection procedure for SSPs R1 and R2. r = model rank, k = number of parameters, Dev. = residual deviance, AICc = Akaike information criterion adjusted for small sample size, wAICc = AICc weight □ = survival probability, ε = departure probability, α = arrival probability, p = recapture probability, A = age, . = constant, Y = year.*

r	Model	k	Dev.	wAICc	AICc
SSP R1					
1	□(A), ε(A), α(A), p(Y)	19	8044.47	0.47	8082.67
2	□(A), ε(A), α(A), p(A + Y)	21	8041.98	0.22	8084.22
3	□(A), ε(.), α(A), p(Y)	17	8050.14	0.21	8084.29
4	□(A), ε(.), α(A), p(A + Y)	19	8047.65	0.10	8085.84
5	□(.), ε(A), α(A), p(Y)	17	8074.54	0.00	8108.70
6	□(.), ε(.), α(A), p(Y)	15	8080.20	0.00	8110.32
7	□(.), ε(A), α(A), p(A + Y)	19	8073.47	0.00	8111.67
8	□(.), ε(.), α(A), p(A + Y)	17	8079.13	0.00	8113.29
9	□(A), ε(A), α(A), p(.)	15	8129.11	0.00	8159.23
10	□(A), ε(A), α(A), p(A)	17	8125.73	0.00	8159.89
11	□(A), ε(.), α(A), p(.)	13	8134.77	0.00	8160.87
12	□(A), ε(.), α(A), p(A)	15	8131.39	0.00	8161.52
13	□(.), ε(A), α(A), p(.)	13	8157.05	0.00	8183.14
14	□(.), ε(.), α(A), p(.)	11	8162.71	0.00	8184.78
15	□(.), ε(A), α(A), p(A)	15	8155.63	0.00	8185.75
16	□(.), ε(.), α(A), p(A)	13	8161.29	0.00	8187.38
SSP R2					
1	□(A), ε(A), α(A), p(A + Y)	21	11041.21	0.64	11083.41
2	□(A), ε(.), α(A), p(A + Y)	19	11047.74	0.19	11085.90
3	□(A), ε(A), α(A), p(A)	17	11052.43	0.13	11086.56
4	□(A), ε(.), α(A), p(A)	15	11058.96	0.04	11089.06
5	□(.), ε(A), α(A), p(A + Y)	19	11068.13	0.00	11106.29
6	□(.), ε(.), α(A), p(A + Y)	17	11074.66	0.00	11108.78
7	□(.), ε(A), α(A), p(A)	15	11078.84	0.00	11108.94
8	□(.), ε(.), α(A), p(A)	13	11085.36	0.00	11111.44
9	□(A), ε(A), α(A), p(Y)	19	11076.17	0.00	11114.33
10	□(A), ε(.), α(A), p(Y)	17	11082.69	0.00	11116.82
11	□(A), ε(A), α(A), p(.)	15	11087.00	0.00	11117.10
12	□(A), ε(.), α(A), p(.)	13	11093.52	0.00	11119.60
13	□(.), ε(A), α(A), p(Y)	17	11111.72	0.00	11145.85
14	□(.), ε(.), α(A), p(Y)	15	11118.24	0.00	11148.34
15	□(.), ε(A), α(A), p(.)	13	11122.54	0.00	11148.62
16	□(.), ε(.), α(A), p(.)	11	11129.07	0.00	11151.12

Fig.4. *Effect of patch turnover on SSP size (number of adults) based on simulations. We considered three turnover rates: high (patch age before disappearance: 3 years, A and D), medium (6 years, B and E), and low (9 years, C and F). Three dispersal scenarios were also considered: (1) no dispersal (individuals were not able to escape when a patch disappeared and thus died); (2) dispersal with random immigration (individuals could disperse to escape the disappearance of a patch or disperse by choice even if the patch remained available. Immigration was random between the patches in the metapopulation and was not influenced by the age of the patch); (3) dispersal with informed immigration (similar to scenario 2, but individuals preferentially immigrated to recently created patches where fecundity F was highest). In scenarios 2 and 3, we considered two possible options regarding dispersal cost: (i) individuals did not incur any survival loss during dispersal (i.e. 'non-costly dispersal', A, B, and C); (ii) survival loss related to dispersal was either low (5% survival cost), medium (10%), or high (15%) (i.e. 'costly dispersal', D, E, and F). To investigate the demographic consequences of these scenarios, we used a female-dominant Leslie matrix (G) based on four demographic parameters: F = the female achieved fecundity, = juvenile survival, = subadult survival, and = adult survival with both environmental and demographic stochasticity.*

Simulating the effect of patch turnover and dispersal on logging SSP dynamics and long-term persistence

The results showed that patch turnover rate was a critical driver of logging SSP dynamics. The absence of dispersal within an SSP experiencing patch turnover necessarily led to the extinction of the SSP (**Fig.4A**, **4B**, and **4C**), and the extinction speed increased with patch turnover rate. In contrast, patch turnover had a positive effect on SSP size (i.e. number of adults) when dispersal was possible

and had no survival cost. This was caused by a fecundity-related mechanism: high patch turnover led

to a decrease in the mean patch age within the SSP, resulting in an increase in average fecundity due to

the positive relationship between fecundity and patch age.

Our simulations also showed that an SSP's extinction risk increased with the survival cost

related to dispersal and was mitigated by 'informed immigration' (**Fig.4D**, **4E**, and **4F**). In the

scenarios with a high patch turnover rate (disappearance of patch after 3 years of availability), SSPs

inevitably went extinct when the survival loss was higher than 10%. Random immigration accelerated

the SSP's decline compared to informed immigration directed toward recently created patches (where

female fecundity was highest). In the scenarios with medium patch turnover (disappearance after 6

years of availability), the SSP decreased when the survival loss was 15% with both random and

informed immigration. With a 10% survival loss, informed dispersal mitigated the decline of an SSP,

whereas random dispersal drove the SSP to extinction. In the scenarios with low patch turnover

(disappearance after 9 years of availability), the SSP experienced a marked decline only when

immigration was random and the survival loss was equal to or higher than 10%. In summary, the

simulations showed that patch turnover may result in SSP decline when the survival cost of dispersal

is relatively high (10–15%) and immigration decisions are not adjusted to patch age and related fitness

prospects.

Phenotypic specialization in riverine and logging environments

We focused on three morphological characteristics known to condition movement capacity in

amphibians: body size, body condition and relative leg size (Gomes et al. 2009, Hillman et al. 2014).

Of the 400 tadpoles, 295 survived until metamorphosis, resulting in a relatively high metamorphosis

success rate ($mean \pm sd = 0.75 \pm 0.08$), which did not differ according to the landscape type of

origin (n = 400, $F_{1,3.8} = 0, P = 0.95$). Neither body size nor body condition at metamorphosis varied

according to the landscape type (n = 295; body size: $F_{1,4} = 0, P = 0.99$; body condition: $F_{1,3.3} =$

$1.12, P = 0.36$). However, the allometric relationship between leg size and body size varied according

to landscape type (n = 295; body size: $F_{1,227} = 776.91$, P < 0.0001; landscape type: $F_{1,3.4} = 3.74$, P =

0.13; body size × landscape type: $F_{1,227} = 6.43$, P = 0.012, Fig. 5). The larger the toadlets, the more

those originating from logging SSPs tended to have longer hind limbs than those originating from

riverine SSPs. Yet this difference was only just significant for large animals, as indicated by sliced

tests (respectively performed at the 1st , 2nd and 3rd quartile of the size distribution: $F_{1,3.8} = 1.64$, P =

0.27; $F_{1,3.4} = 3.28$, P=0.15; $F_{1,4.01} = 6.54$; P = 0.06).

Based on the behavioral trials in the experimental arenas, we measured variables along the

boldness–shyness behavioral axis, a personality trait consistently involved in dispersal syndromes

across different organisms (Cote et al. 2010). Toadlet behavior was characterized using three

measures: one to assess neophobia (BEHAV1 = the time taken to leave the familiar refuge chamber) and

the other two to assess exploration propensity (BEHAV2 = the time taken to reach the desiccant zone

after leaving the refuge chamber; BEHAV3 = the time taken to cross the desiccant zone and get out of

the arena). The findings showed that the neophobia of toadlets significantly varied according to their

landscape of origin (**Fig.5**), but not according to their body size or its interactive effect with landscape

(N=295; landscape type: $F_{1,3.5}$ =47.30, P=0.004; body size: $F_{1,13.9}$=0.03, P=0.86; their interactive effect:

$F_{1,12.1}$=0.01, P=0.92). Individuals from riverine landscapes were 2.13+-0.24 times slower to leave the

refuge chamber than those from logging landscapes (respectively 8.58+-0.67 mn and 3.89+-0.33 mn,

see **Fig.5**). Once the toadlet left the refuge chamber, the latency time to reach the desiccating zone was

very short (mean 1.83 +- 0.21 mn) and did not vary depending on the landscape of origin, body size,

or their interactive effect (N=295; landscape type: $F_{1,3.9}$ =1.84, P=0.25; body size: $F_{1,30}$=0.42, P=0.52;

their interactive effect $F_{1,33.4}$=0.26, P=0.61, see **Fig.5**). In contrast, the latency time to get out of the

arena significantly varied depending on the landscape of origin, but not according to body size

(N=295; landscape type: χ^2_{1df}=9.56, P =.002; body size: χ^2_{1df}=0.01, P=0.91; their interactive effect:

χ^2_{1df}=0.28, P=0.59). The time taken – and thus the increased hazard – to get out of the arena for

toadlets originating from logging SSPs was 1.93 times greater than for toadlets from riverine SSPs.

Fig.5. Behavioral and morphological specialization in six Bombina variegata SSPs occurring in

logging environments (L3, L4, and L5) and riverine environments (R1, R2, and R3). We examined how

the type of environment affected two behavioral traits: the time to exit the refuge chamber (A), a proxy
for neophobia, and the time to reach the desiccation zone (B), a proxy for exploration propensity. We
also examined how environment type affected hind limb length while considering body length as a
control covariate in the model. These observations were recorded for 295 toadlets at metamorphosis.

721
722   **Fig.6**. Inbreeding in the riverine (blue) and logging (green) *Bombina variegata* SSPs.
723

724
725   **Fig.7**. Relatedness within patches in the riverine (blue) and logging (green) *Bombina variegata* SSPs.
726   Patches where less than 6 relatedness values were available were removed from the plot.
727

Neutral genetic variation in riverine and logging environments

We investigated the neutral genetic footprint associated with each landscape type using microsatellite
data collected in four SSPs: two from each landscape type (a total of 667 toads genotyped within L3
and L4 logging environments and R1 and R2 riverine environments; **Table S2**). Our analyses
revealed, first, that riverine systems exhibited a higher deviation from the Hardy–Weinberg

equilibrium than logging systems. Significant deviation from this equilibrium was detected for 75%

and 45% of the loci in riverine SSPs R1 and R2 respectively, and 0% and 18% in the logging SSPs L3

and L4 (**Table S3**). Second, we found lower genetic diversity in riverine SSPs than in logging SSPs:

both the allelic richness and the expected heterozygosity (uHE) were substantially lower in riverine

SSPs. In contrast, inbreeding coefficients ( ) were higher in riverine SSPs: 87% of the loci were

found to have a lower uHE in riverine SSPs compared to logging SSPs (**Table S3**). Similarly, our

parentage analyses also showed that individual inbreeding (LR test, = 32.55, P <0.0001; **Fig.6**) and

the relatedness level within patches (LR test, = 43.14, P <0.0001, **Fig.7**) were drastically higher in

riverine SSPs. In addition, riverine SSPs had a much smaller effective population size than logging

SSPs: the effective population size was 9.7 (95% CI 6.4–13.7) in R1 and 22.6 (95% 19.3–26.3) in R2,

while it was 138.9 (95% 80.2–353.6) in L4 and 168.6 (95% 129.9–229.3) in L3.

Within both types of environment, the size of SSPs (large: R2 and L3; small: R1 and L4) was

also an important predictor of individual relatedness and inbreeding. The relatedness level within

patches was higher (**Fig.7**) in small SSPs than in large SSPs (LR test, = 214.38, P <0.0001), and

this difference was larger in riverine SSPs (LR test, = 6.16, P = 0.01). Similarly, inbreeding within

patches was higher (**Fig.6**) in small SSPs than in large SSPs (LR test, = 76.56, P <0.0001), and the

interaction between these two factors was also confirmed (LR test, = 8.88, P = 0.01).

752
753 ***Fig.8****. Patterns of neutral genetic variation and spatial distribution of the hierarchical genetic clusters*
754 *in two riverine (R1 and R2) and logging SSPs (L3 and L4). The analyses were conducted using the*
755 *program STRUCTURE. In L3 and L4, no genetic structuring was detected. In contrast, a complex*

genetic structure was found in R1 and R2. In R1, the hierarchical analysis revealed the existence of two initial genetic clusters (A and B), and genetic substructures within cluster A where two nested clusters were inferred (AA and AB). In R2, the hierarchical analysis highlighted the presence of two initial genetic clusters (A and B), and a substructure was further detected within cluster B (BA and BB). Genetic substructures were then identified within the cluster BA, with three additional genetic clusters inferred (BAA, BAB, and BAC).

Table 5. *Relative support of the mixed models for each of the four SSPs and their related estimates of geographic resistance (L3 and L4 = logging SSPs, R1 and R2 = riverine SSPs, see Map S1 for details). 'Model' indicates the effect introduced in the model in addition to geographic resistance (null = the model including only the effect of geographic resistance). 'Waic' indicates the weighted AIC of the model. $\hat{\beta}(sd)$ indicates the beta weight associated with the effect of geographic resistance and its standard deviation. The variable 'slope' corresponds to the effect of both geographic distance and topographic roughness. The variable 'network' corresponds to the effect of the hydrological network.*

Model	Waic	$\hat{\beta}(sd)$
SSP L3		
null	0.469	1.4763 (0.1544)
slope	0.245	1.4811 (0.1545)
network	0.191	1.4798 (0.1545)
slope+network	0.095	1.4826 (0.1546)
SSP L4		
null	0.169	0.6783 (0.0415)
slope	0.084	0.679 (0.0415)
network	0.507	0.6764 (0.0415)
slope+network	0.240	0.677 (0.0415)
SSP R1		
null	0.391	2.381 (0.0713)
slope	0.276	2.386 (0.0714)
network	0.226	2.3845 (0.0714)
slope+network	0.107	2.3862 (0.0714)
SSP R2		
null	0.070	2.5459 (0.0251)
slope	0.030	2.5461 (0.0251)
network	0.601	2.5479 (0.0251)
slope+network	0.299	2.5482 (0.0251)

We then investigated how patterns of genetic differentiation between patches differed between logging and riverine SSPs. First, we used the Bayesian genetic clustering approach to examine hierarchical genetic structure in the SSPs. In the two SSPs from logging environments, we failed to detect any hierarchical genetic structure (K = 1) (**Fig.8**). In contrast, our analyses revealed a strong hierarchical genetic structure overlaying the spatial distribution of patches in the two SSPs from riverine environments (**Fig.8**). Second, we sought to verify that differences in spatial genetic patterns between landscapes were still significant while controlling for the potential difference of functional connectivity between SSPs using direct gradient analyses. These revealed that genetic differentiation correlated with geographic resistance between individuals whatever the landscape type, indicating substantial genetic isolation by distance even in logging systems (**Fig.9**). There was considerable

784 support for the effect of the hydrological network on genetic differentiation in the large SSPs of both

785 landscape types (SSPs R2 and L3), but not in the small SSPs whatever the landscape type (**Table 5**).

786 Most importantly, even after correcting for these potential landscape effects, spatial genetic

787 differentiation remained higher in riverine than in logging SSPs as revealed by the model averaged

788 slope estimate associated with the geographic resistance effect (:   for R1 and

789   for R2;   for L3 and   for L4).

791
Fig.9. Genetic divergence according to geographic distance in two riverine (R1 and R2) and two
logging SSPs (L3 and L4). Each figure represents the contour plot of the kernel density bivariate
estimates between the pairwise genetic distance and the pairwise geographic resistance for each SSP.
Kernel densities were estimated using a Gaussian distribution. The graduated color contour
indicating the (smoothed) observation count is presented on the right side of each plot. The line
represents the predicted regression curve between the genetic distance and the geographic resistance
from the linear mixed model estimates.

Discussion

Taken together, the results of this study provide for the first time an extended picture of the effect of

anthropogenic disturbance on dispersal in a vertebrate, from dispersal-related phenotypic

specialization expressed early in life, through the dispersal pattern emerging in spatially structured

populations, to the genetic footprint arising throughout successive generations. Overall, this analysis

revealed that anthropogenic disturbance not only strongly promotes dispersal throughout a toad's

lifetime, but also prenatally enhances a toadlet's risk-proneness and, to a certain extent, favors longer

hind limb length at metamorphosis. Another finding was that gene flow also substantially increased in

anthropogenic landscapes independently of the SSP's size or functional connectivity.

Natal and breeding dispersal rates and distances in SSPs depend on patch turnover rate

The results revealed contrasting dispersal patterns between riverine and logging SSPs. In the

demographic component of our study, the sampling protocol was weakened by a potential

confounding effect of the population's position along the latitudinal gradient with its status

(logging/riverine). However, we can rule out the possibility of an effect of latitude on the dispersal

patterns drawn in our analyses, as the molecular inferences showed drastically increased gene flow

(resulting from dispersal) in an SSP at a low latitude (L4), which indicates that latitude has a marginal

effect on effective dispersal within SSPs. We are therefore confident in the reliability of our results

regarding the effect of turnover rate on dispersal patterns.

In riverine SSPs, we observed a complete suppression of natal dispersal, as well as very low

breeding dispersal, together with a reduced dispersal kernel – the opposite of the dispersal pattern

observed in logging SSPs. In the latter, we nevertheless found some substantial inter-population

variation regarding the contribution of natal dispersal to the overall dispersal process and to seasonal

variation in dispersal rates. Such differences between logging SSPs likely reflect variation in the

anthropogenic disturbance regime resulting from local woodland management practices. Indeed, patch

turnover depends on both the extent and the frequency of the logging activities that create patches, as

well as the post-logging rehabilitation operations that may lead to patch destruction (e.g. filling in of

ruts used as temporary breeding ponds) (Cayuela et al 2018b). This likely results in a continuum of

dispersal strategies along a gradient of patch disturbance, ranging from the near suppression of

dispersal in riverine SSPs to a very high dispersal rate and long dispersal distances in some logging

SSPs (e.g. L1 SSP). This pattern is thus very similar to that observed for wing dimorphism in insects

alongside patch temporality gradients (Denno et al. 1996, Roff & Fairban 2007), in which dispersal is

suppressed (i.e. high rate of wingless forms) in persistent habitats, while it is enhanced (i.e. high rate

of winged forms) in more ephemeral habitats according to the (natural) disturbance regime

experienced at the landscape level.

Phenotypic specialization in SSPs depends on patch turnover rate

Previous studies on *Bombina variegata* (Cayuela et al. 2018b) have revealed that dispersal events in

logging SSPs are a mixture of departures conditioned by patch disappearance and unconditional

departures that occur well before patch disappearance. The findings from our common garden

experiments highlight that patch turnover rate prenatally mediates dispersal-related phenotypes and

leads to phenotypic parallelism in logging and riverine SSPs. In particular, toadlets originating from
logging SSPs exhibited higher risk-proneness than those from riverine SSPs, as revealed by their
swiftness in leaving the refuge chamber and in getting out of the arena after crossing a harsh substrate.
Either neophilia or boldness (see Kelleher et al. 2018 for personality traits in amphibians) could
explain the elevated risk-proneness we observed in logging SSPs. Disentangling these two personality
traits is not straightforward (Greggor et al. 2015, Yuen et al. 2017), and further investigations would
be useful to address this. Regardless of the exact composition of personality traits behind our
identification of risk-prone behavior, our results clearly indicate behavioral specialization early in life
according to the disturbance regime prevailing in the landscape.

Concerning morphological traits, we did not find any differences in body size and condition of
toadlets from logging and riverine SSPs. However, the findings showed that toadlets innately have
longer hind limbs in logging than in riverine SSPs. In anurans, hind-limb length is usually positively
associated with locomotor performance (Choi et al. 2003, Philips et al. 2006, Gomes et al 2009,
Hudson et al. 2016), and has also been found to be subject to rapid evolution at the edge of the
invasion front in the introduced species *Rhinella marina* (Philips et al. 2006). Our results thus suggest
that long hind limbs could be a phenotypic trait facilitating dispersal in logging habitats. Yet this
difference in leg length between toadlets from the two environments was only observed in large
individuals and was subject to substantial variation between SSPs. This significant but weak effect of
anthropogenic disturbance on leg length could result from developmental constraints. Limb size
mainly depends on the duration of the larval period in anurans, so species specialized for ephemeral
pools (such as *Bombina variegata*), which are selected for fast larval development, usually exhibit
shorter hind limbs compared to other species (Gomez-Mestre & Buchholz 2006).

Overall, dispersal and related behavioral traits are usually highly plastic phenotypes subject to
partial genetic control (reviewed in Saastamoinen et al. 2018), i.e. determined by $G \times E$ interactions.
Therefore, the phenotypic specialization highlighted in our study may have a genetic basis and/or may
be associated with transgenerational plasticity. In the absence of crossbreed design, we were not able
to disentangle the relative contribution of maternal effect and parental genotypes in the phenotypic
variation observed in logging and riverine SSPs. However, it is possible to rule out the hypothesis of
transgenerational plasticity mediated by toadlet body size at metamorphosis. In amphibians, female
energy investment in breeding influences the size of eggs and the amount of energetic resources
available for the development of embryos and larvae before they become fully heterotrophic (Kaplan
1987, 1992). Studies have reported a positive relationship between egg size and offspring body size at
metamorphosis due to carry-over effects (Laugen et al. 2005, Räsänen et al. 2005, Dziminski et al.
2006), and body size is an important predictor of behavioral traits (e.g. exploration propensity and
risk-taking behavior) related to dispersal (Kelleher et al. 2017, 2018). In our case study, no difference
in body size was detected in toadlets from logging and riverine environments, which indicates that
patch turnover rate is not likely to alter individual behavior due to the morphological state at

metamorphosis in SSPs. However, epigenetic factors (e.g. DNA methylation, micro-ARN, and histone

structure) independent from female energy investment strategies may lead to transgenerational

dispersal plasticity and contribute to the expression of phenotypic traits that facilitate or hinder

dispersal over generations (Saastamoinen et al. 2018, Cayuela et al. 2019a). Yet it is very likely that a

genetic basis partially determines phenotypic specialization in logging and riverine environments,

resulting in incomplete genetic parallelism between SSPs (Elmer & Meyer 2011, Conte et al. 2012).

First, transgenerational plasticity in dispersal-related traits is usually subject to genetic control

(Cayuela et al. 2019a), likely due to a strong association between epigenetic variation and genetic

variants in *cis* and *trans* (Dubin et al. 2015, Zaghlool et al. 2016). Second, as predicted by the

dispersal theory and reported in other study systems, dispersal is partly genetically determined, and

behavioral traits related to dispersal often have a polygenic basis (Saastamoinen et al. 2018). Future

studies using Next-Generation Sequencing approaches (Morozova & Marra 2008, Metzker 2010)

could be undertaken to determine the role of genetic, transcriptional, and epigenetic variation in the

disturbance-dependent phenotypic changes and dispersal evolution in *B. variegata*.

Patch turnover and related dispersal costs and benefits determine SSP dynamics and persistence

Our simulations showed that the absence of dispersal inevitably leads to SSP extinction, and that the

extinction speed increases with patch turnover. When dispersal was possible and had no survival cost,

patch turnover had a positive effect on SSP size. This resulted from an increase in average fecundity

due to the positive relationship between fecundity and patch age reported in logging SSPs (Boualit et

al. 2019) and considered in our models. These results are congruent with field observations reporting

that *Bombina variegata* SSPs may be very large (thousands of adults) in harvested woodlands in

western Europe.

906   The models also showed that the risk of SSP extinction increased with the survival cost related

to dispersal and was mitigated by informed immigration. This result is congruent with theoretical

models and empirical evidence showing that dispersal only evolves if the benefits of moving outweigh

the related costs (Bonte et al. 2012). In logging SSPs, dispersal seems to be favored as its costs are

likely offset by the benefits of colonizing new patches. Survival (at juvenile, subadult, and adult stage)

is lower in logging than in riverine SSPs (Cayuela et al. 2016a), which likely results from mortality

caused by logging operations and dispersal-related mortality (Cayuela et al. 2018b). This survival cost

of dispersal can be direct, i.e. associated with movement in the landscape matrix (Bonte et al. 2012). It

can be also indirect, resulting from the high energy allocation necessary for recurrent dispersal events

over the toad's lifespan, which might lead to earlier and stronger senescence due to trade-offs

(Cayuela et al. 2019b). These direct and indirect costs are obviously offset when animals are forced to

disperse subsequently to patch destruction resulting from logging practices (Cayuela et al 2018b). It

should be noted that even when patches remain available, dispersal may still be strongly favored since

their suitability rapidly declines over time due to the natural silting of ruts if these are not regularly
disturbed by vehicle traffic (Boualit et al. 2019). Furthermore, even if patch suitability is sustained by
regular human disturbance (Boualit et al. 2019), dispersal costs might be mitigated through the
colonisation of recently created patches in which fitness prospects are likely enhanced due to density-
dependent mechanisms. A low density of adults in newly available patches likely reduces the risk of
larval competition, which is an important driver of metamorphosis success (Jasieński 1988) – for this
reason adults preferentially reproduce in tadpole-free waterbodies (Cayuela et al. 2016d, 2017b).

These potential benefits likely favor the evolution of dispersal and dispersal-enhancing
morphological and behavioral traits (i.e. 'dispersal syndromes') in logging SSPs. They also likely
contribute to context-dependent dispersal (Clobert et al. 2009) and matching habitat choice (Edelaar et
al. 2008), implying that individuals adjust their dispersal decisions according to the local fitness
prospects determined by the biotic and abiotic characteristics of breeding patches. This prediction has
been verified by two studies reporting context-dependent dispersal in logging SSPs (Tournier et al.
2017, Boualit et al. 2019). In particular, these studies showed that adult emigration and immigration
decisions depend on a pond's hydroperiod and the size and annual disturbance of the patch: three
factors that locally affect juvenile recruitment and very likely individual fitness.

In riverine SSPs, the near absence of both natal and breeding dispersal suggests that the
benefits of dispersal do not compensate for its potential costs. First, the absence of patch loss resulting
from anthropogenic or natural processes means individuals are not forced to disperse to survive and
reproduce. In river environments, the local fitness prospects do not deteriorate with patch age, as the
process of natural silting of rocky pools is frequently interrupted by river flooding occurring outside
the breeding period (Cayuela et al. 2011, 2015a). This makes pools available for breeding from one
year to another and limits the risk of larval mortality caused by desiccation. It is also possible that the
near absence of dispersal results from variability in pond characteristics (e.g. hydroperiod and
temperature; Cayuela et al. 2011) within a patch. Indeed, the benefit of dispersing would be low if the
variability in environmental conditions would be similar at intra-patch and inter-patch scales. Overall,
the lack of apparent compensatory benefits should not favor and may even counter-select for dispersal
and dispersal-enhancing traits in riverine SSPs. This is in line with our results, which show that
riverine toadlets innately display low risk-taking behavior and have short hind limbs, two phenotypic
traits hindering dispersal in amphibians (Cayuela et al. 2018c).

Genetic variation patterns in SSPs depend on patch turnover rate

Our study showed for the first time that the spatiotemporal dynamics of habitat patches in a landscape
may have an effect at least as important as landscape fragmentation on gene flow patterns. The
findings highlighted that low dispersal rates and distances are associated with a weaker genetic

structure and lower IBD in riverine than in logging SSPs. Our sampling design allowed us to

disentangle the relative effects of gene flow and genetic drift, which both contribute to genetic

differentiation within SSPs (Slatkin 1977, Broquet & Petit 2009). As well as genetic structure

differences caused by patch turnover, we found higher IBD in the two small SSPs (L4 and R1) than in

the two large SSPs in each environment.

Our analysis also took into account the functional connectivity within SSPs by considering

two landscape factors (i.e. topography and hydrological network) that are critically important to the

genetic structure of amphibian populations (reviewed in Cayuela et al. 2018c). As reported in 20

previous studies (Cayuela et al. 2018c), genetic differentiation increases with increasing topographic

slope as this raises the energy cost of dispersal during the transience phase in amphibians. In addition,

our results showed that genetic differentiation was better explained by the distance of the hydrological

network than the Euclidean distance between two patches. This indicates that the hydrological network

improves the functional connectivity between patches by reducing the cost of displacement in the

landscape matrix, which is congruent with the findings of six previous studies (Cayuela et al. 2018c).

Interestingly, genetic differentiation nevertheless remained higher in riverine than in logging SSPs,

despite a denser hydrological network and therefore weaker landscape resistance in riverine SSPs.

This suggests that in our study system, the rate of patch turnover may be a more important driver of

neutral genetic variation than landscape connectivity.

Our findings suggest that variation in the turnover rate of SSPs has far-reaching consequences

on the evolutionary forces involved in the migration–selection–drift balance. In riverine SSPs, reduced

gene flow between patches leads to lower genetic diversity and smaller effective population size

compared to logging SSPs. Such low standing genetic variation could limit the adaptive response of

riverine SSPs to novel environmental conditions. The probability of allele fixation increases with the

magnitude of the beneficial effect and the effective population size, and this probability is significantly

higher when the allele has a high initial frequency (Barrett & Schluter 2008, Hedrick 2013). Moreover,

standing genetic variation usually allows faster adaptation as beneficial alleles are already present in

the population (Barrett & Schluter 2008, Hedrick 2013). Overall, our results suggest that riverine SSPs

should suffer from a lower capacity of 'evolutionary rescue' (Carlson et al. 2014) than logging SSPs,

which could increase their sensitivity to current global changes.

The results also revealed a high level of inbreeding and relatedness within riverine SSPs,

confirming the results of a previous study (Cayuela et al. 2017a). These findings raise important

questions about the mechanisms associated with the repression of inbreeding depression. As

mentioned, the survival rate at all life stages (juvenile, subadult, and adult) is higher in riverine than in

logging SSPs (Cayuela et al. 2016a), suggesting a marginal effect of inbreeding on postmetamorphic

survival and negligible inbreeding depression. A previous study conducted in the R2 SSP indicated

that the absence of disassortative mating does not seem to mitigate inbreeding risk: females even

prefer to reproduce with related males from their own patch (Cayuela et al. 2017a). Although the

effect of inbreeding on reproductive performance remains as yet unevaluated, those results suggest a
low genetic load in riverine SSPs. This could be due to high efficiency in purging deleterious alleles
and the genomic architecture of genetic load, especially a low linkage of deleterious recessive alleles
(Bersabé et al. 2016, Hedrick & Garcia-Dorado 2016). Future studies could be carried out to identify
the mechanisms involved in the genetic purging of inbreeding depression resulting from limited
dispersal in riverine SSPs.

Conclusion

1000

Our study showed that anthropogenic habitat disturbance is likely an important driver of dispersal
evolution. The results found that, like landscape fragmentation, human-driven variation in patch
turnover may promote morphological and behavioral specialization related to dispersal. In particular, it
may lead to phenotypic parallelism affecting dispersal syndromes and patterns (dispersal rate and
distance) in SSPs exposed to contrasting levels of patch turnover. This phenotypic parallelism is likely
underpinned by genetic and/or epigenetic parallelism, for which the molecular basis remains to be
investigated. Our results also revealed that differences in dispersal patterns are associated with
variation in the genetic structure of SSPs, which might affect local eco-evolutionary dynamics
(Legrand et al. 2017). In particular, high gene flow and reduced effects of genetic drift allow higher
genetic polymorphism to be maintained in SSPs experiencing high patch turnover than in SSPs with
low patch turnover. In parallel, larger effective population size is expected to increase selection
effectiveness in SSPs exposed to high patch turnover, giving them higher evolutionary potential and
increased chances of evolutionary rescue in the case of environmental change (Carlson et al. 2014).
These results emphasize the central role of anthropogenic disturbance in the spatiotemporal dynamics
of landscapes and the related ecological and evolutionary processes.

In terms of the implications for conservation, the findings suggest that the approach may need
to differ depending on the demographic and genetic characteristics of SSPs driven by the level of
patch turnover. Those SSPs exposed to high patch turnover are likely to be highly sensitive to land use
changes that result in a loss of functional connectivity between patches. An increase in landscape
resistance reduces the success of dispersal following the disappearance of a patch and diminishes the
chance of new patch colonization, which enhances the extinction risk of the whole SSP. In contrast,
SSPs experiencing a low level of patch turnover suffer a loss of genetic diversity and high inbreeding,
which may increase their sensitivity to environmental change (e.g. climate change or the emergence of
new pathogens), potentially speeding up their extinction if they experience a demographic decline (i.e.
inbreeding depression; Reed & Frankham 2003, Frankham 2010). This underlines the tremendous
importance of adjusting conservation decisions to the demographic and genetic characteristics of SSPs
adapted to contrasting regimes of habitat disturbance.

1028

1029 **Acknowledgments**

1030

We would like to thank all the undergraduate students and fieldworkers that contributed to the

capture–recapture surveys, DNA samples, and common garden behavioral tests. This research project

was funded by the Office National des Forêts, the Lorraine Direction Régionale de l'Environnement,

de l'Aménagement et du Logement (DREAL), the Agence de l'Eau Rhin-Meuse, the Conseil Régional

de Lorraine, the Conseil Régional de Champagne-Ardenne, the Conseil Régional de Picardie, the

Conseil Général de l'Aisne, the Conseil Général d'Ardèche, the Conseil Général d'Isère and the

Communauté de Communes de l'Argonne Ardennaise (2C2A). The study was authorized by three

government environmental agencies: the Lorraine DREAL, the Rhône-Alpes DREAL and the Agence

de l'Eau Rhin-Meuse. The ethical committee of Université Claude Bernard Lyon 1 approved the

capture–recapture and common garden protocols.

1041

1042   **References**
1043

Armsworth, P. R., & Roughgarden, J. E. (2005). Disturbance induces the contrasting evolution of

reinforcement and dispersiveness is directed and random movers. Evolution, 59, 2083-2096.

Atkins, J. L., Perry, G. L., & Dennis, T. E. (2019). Effects of mis-alignment between dispersal traits

and landscape structure on dispersal success in fragmented landscapes. Royal Society Open

Science, 6, 181702.

Baguette, M., Blanchet, S., Legrand, D., Stevens, V. M., & Turlure, C. (2013). Individual dispersal,

landscape connectivity and ecological networks. Biological Reviews, 88, 310-326.

Balkenhol, N., Holbrook, J. D., Onorato, D., Zager, P., White, C., & Waits, L. P. (2014). A

multi☐method approach for analyzing hierarchical genetic structures: a case study with cougars

Puma concolor. Ecography, 37, 552-563.

Barrett, R. D., & Schluter, D. (2008). Adaptation from standing genetic variation. Trends in Ecology

& Evolution, 23, 38-44.

Bersabé, D., Caballero, A., Pérez-Figueroa, A., & García-Dorado, A. (2016). On the consequences of

purging and linkage on fitness and genetic diversity. G3: Genes, Genomes, Genetics, 6, 171-181.

Bonte, D., Van Dyck, H., Bullock, J. M., Coulon, A., Delgado, M., Gibbs, M., Lehouck, V.,

Matthysen, E., Mustin, K., Saastamoinen, M., Schtickzelle, N., Stevens, V. M., Vandewoestijne, S.,

Baguette, M., Barton, K., Benton, T.G., Chaput-Bardy, A., Clobert, J., Dytham, C., Hovestadt, T.,

Meier, C. M., Palmer, S.C.F., Turlure, C., Travis, J. M. J. (2012). Costs of dispersal. Biological

Reviews, 87, 290-312.

Boualit, L., Pichenot, J., Besnard, A., Helder, R., Joly, P., & Cayuela, H. (2019). Environmentally

mediated reproductive success predicts breeding dispersal decisions in an early successional

amphibian. Animal behaviour, 149, 107-120.

Bowler, D. E., & Benton, T. G. (2005). Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. Biological Reviews, 80, 205-225.

Broquet, T., & Petit, E. J. (2009). Molecular estimation of dispersal for ecology and population genetics. Annual Review of Ecology, Evolution, and Systematics, 40, 193-216.

Carlson, S. M., Cunningham, C. J., & Westley, P. A. (2014). Evolutionary rescue in a changing world. Trends in Ecology & Evolution, 29, 521-530.

Caswell H. (2001) Matrix population models: constraints, analysis and interpretation, Sinauer Associates, Sunderland.

Cayuela, H., Cheylan, M., & Joly, P. (2011). The best of a harsh lot in a specialized species: breeding habitat use by the yellow-bellied toad (*Bombina variegata*) on rocky riverbanks. Amphibia-Reptilia, 32, 533-539.

Cayuela, H., Arsovski, D., Boitaud, S., Bonnaire, E., Boualit, L., Miaud, C., Joly, P., & Besnard, A. (2015a). Slow life history and rapid extreme flood: demographic mechanisms and their consequences on population viability in a threatened amphibian. Freshwater Biology, 60, 2349-2361.

Cayuela, H., Lambrey, J., Vacher, J. P., & Miaud, C. (2015b). Highlighting the effects of land-use change on a threatened amphibian in a human-dominated landscape. Population Ecology, 57, 433-443.

Cayuela, H., Arsovski, D., Thirion, J. M., Bonnaire, E., Pichenot, J., Boitaud, S., Brison, A.-L., Miaud, C., Joly, P., & Besnard, A. (2016a). Contrasting patterns of environmental fluctuation contribute to divergent life histories among amphibian populations. Ecology, 97, 980-991.

Cayuela, H., Boualit, L., Arsovski, D., Bonnaire, E., Pichenot, J., Bellec, A., Miaud, C., Léna, J.-P., Joly, P., & Besnard, A. (2016b). Does habitat unpredictability promote the evolution of a colonizer syndrome in amphibian metapopulations?. Ecology, 97, 2658-2670.

Cayuela, H., Arsovski, D., Thirion, J. M., Bonnaire, E., Pichenot, J., Boitaud, S., Miaud, C., Joly, P., & Besnard, A. (2016c). Demographic responses to weather fluctuations are context dependent in a long‐lived amphibian. Global Change Biology, 22, 2676-2687.

Cayuela, H., Lengagne, T., Kaufmann, B., Joly, P., & Léna, J. P. (2016d). Larval competition risk shapes male–male competition and mating behavior in an anuran. Behavioral Ecology, arw100.

Cayuela, H., Léna, J. P., Lengagne, T., Kaufmann, B., Mondy, N., Konecny, L., Dumet, A., Vienney, A., & Joly, P. (2017a). Relatedness predicts male mating success in a pond-breeding amphibian. Animal Behaviour, 130, 251-261.

Cayuela, H., Lengagne, T., Joly, P., & Léna, J. P. (2017b). Females trade off the uncertainty of breeding resource suitability with male quality during mate choice in an anuran. Animal Behaviour, 123, 179-185.

Cayuela, H., Rougemont, Q., Prunier, J. G., Moore, J. S., Clobert, J., Besnard, A., & Bernatchez, L.
(2018a). Demographic and genetic approaches to study dispersal in wild animal populations: a
methodological review. Molecular Ecology, 27, 3976-4010.

Cayuela, H., Besnard, A., Quay, L., Helder, R., Léna, J. P., Joly, P., & Pichenot, J. (2018b).
Demographic response to patch destruction in a spatially structured amphibian population. Journal
of Applied Ecology, 55, 2204-2215.

Cayuela, H., Valenzuela-Sanchez, A., Teulier, L., Martínez-Solano, Í., Léna, J. P., Merilä, J., Muths,
E., Shine, R., Quay, L., Denoël, M., Clobert, J., Schmidt, B. R. (2018c). Determinants and
consequences of dispersal in vertebrates with complex life cycles: a review of pond-breeding
amphibians. PeerJ Preprints, e27394v1.

Cayuela, H., Jacob, S., Schtickzelle, N., Verdonck, R., Philippe, H., Laporte, M., Bernatchez, L.,
Legrand, D. (2019a). Transgenerational dispersal plasticity and its fitness consequences are under
genetic control. bioXiv, doi: ...

Cayuela, H., Lemaitre, J. F., Bonnaire, E., Pichenot, J., & Schmidt, B. (2019b). Population position
along the fast-slow life-history continuum predicts intraspecific variation in actuarial senescence.
bioRxiv, 621425.

Cayuela, H., Gillet, L., Laudelout, A., Besnard, A., Bonnaire, E., Levionnois, P., Muths, E., Dufrêne,
M., & Kinet, T. (2019c). Survival cost to relocation does not reduce population self‐sustainability
in an amphibian. Ecological Applications, 29, e01909.

Cheptou, P. O., Hargreaves, A. L., Bonte, D., & Jacquemyn, H. (2017). Adaptation to fragmentation:
evolutionary dynamics driven by human influences. Philosophical Transactions of the Royal
Society B: Biological Sciences, 372, 20160037.

Choquet, R., Rouan, L., & Pradel, R. (2009). Program E-SURGE: a software application for fitting
multievent models. In: Thomson, D. L., Cooch, E. G., Conroy, M. J. (eds). Modeling demographic
processes in marked populations. Pp. 845-865. Springer, Boston, US.

Clarke, R., Rothery, P. & Raybould, A. (2002). Confidence limits for regression relationships between
distance matrices: Estimating gene flow with distance. Journal of Agricultural, Biological, and
Environmental Statistics, 7, 361-372.

Clobert, J., Galliard, L., Cote, J., Meylan, S., & Massot, M. (2009). Informed dispersal, heterogeneity
in animal dispersal syndromes and the dynamics of spatially structured populations. Ecology
letters, 12, 197-209.

Comins, H. N., Hamilton, W. D., & May, R. M. (1980). Evolutionarily stable dispersal strategies.
Journal of Theoretical Biology, 82, 205-230.

Conte, G. L., Arnegard, M. E., Peichel, C. L., & Schluter, D. (2012). The probability of genetic
parallelism and convergence in natural populations. Proceedings of the Royal Society B: Biological
Sciences, 279, 5039-5047.

Cote, J., Clobert, J., Brodin, T., Fogarty, S., & Sih, A. (2010). Personality-dependent dispersal:
characterization, ontogeny and consequences for spatially structured populations. Philosophical
Transactions of the Royal Society B: Biological Sciences, 365, 4065-4076.

Cote, J., Bestion, E., Jacob, S., Travis, J., Legrand, D., & Baguette, M. (2017). Evolution of dispersal
strategies and dispersal syndromes in fragmented landscapes. Ecography, 40, 56-73.

Coulon, A., Fitzpatrick, J. W., Bowman, R., Stith, B. M., Makarewich, C. A., Stenzler, L. M., &
Lovette, I. J. (2008). Congruent population structure inferred from dispersal behaviour and
intensive genetic surveys of the threatened Florida scrub□jay (*Aphelocoma cœrulescens*).
Molecular Ecology, 17, 1685-1701.

Cushman, S. A. (2006a). Effects of habitat loss and fragmentation on amphibians: a review and
prospectus. Biological conservation, 128, 231-240.

Cushman, S. A., McKelvey, K. S., Hayden, J., & Schwartz, M. K. (2006b). Gene flow in complex
landscapes: testing multiple hypotheses with causal modeling. The American Naturalist, 168, 486-
499.

Cushman, S., Storfer, A., & Waits, L. (2015). Landscape genetics: concepts, methods, applications.
John Wiley & Sons.

Denno, R. F., Roderick, G. K., Peterson, M. A., Huberty, A. F., Dobel, H. G., Eubanks, M. D., Losey,
J. E., & Langellotto, G. A. (1996). Habitat persistence underlies intraspecific variation in the
dispersal strategies of planthoppers. Ecological Monographs, 66, 389-408.

Do, C., Waples, R. S., Peel, D., Macbeth, G. M., Tillett, B. J. & Ovenden, J. R. (2014) NeEstimator
V2: re-implementation of software for the estimation of contemporary effective population size
(Ne) from genetic data. Molecular Ecology Resources, 14, 209-214.

Dubin, M. J., Zhang, P., Meng, D., Remigereau, M. S., Osborne, E. J., Casale, F. P., Drewe, P. Kahles,
A., Jean, G., Vilhjalmsson, B., Jagoda, J., Irez, S., Voronin, V., Song, Q., Long, Q., Ratsch, G.,
Stegle, O., Clark, R. M., & Nordborg, M. (2015). DNA methylation in Arabidopsis has a genetic
basis and shows evidence of local adaptation. elife, 4, e05255.

Dziminski, M. A., & Roberts, J. D. (2006). Fitness consequences of variable maternal provisioning in
quacking frogs (Crinia georgiana). Journal of Evolutionary Biology, 19, 144-155.

Earl, D. A. (2012). STRUCTURE HARVESTER: a website and program for visualizing
STRUCTURE output and implementing the Evanno method. Conservation Genetics Resources, 4,
359-361.

Edelaar, P., Siepielski, A. M., & Clobert, J. (2008). Matching habitat choice causes directed gene
flow: a neglected dimension in evolution and ecology. Evolution, 62, 2462-2472.

Edwards, D. P., Gilroy, J. J., Thomas, G. H., Uribe, C. A. M., & Haugaasen, T. (2015). Land-sparing
agriculture best protects avian phylogenetic diversity. Current biology, 25, 2384-2391.

Elmer, K. R., & Meyer, A. (2011). Adaptation in the age of ecological genomics: insights from
parallelism and convergence. Trends in Ecology & Evolution, 26, 298-306.

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. Molecular Ecology, 14, 2611-2620.

Fahrig, L. (2003). Effects of habitat fragmentation on biodiversity. Annual Review of Ecology, Evolution, and Systematics, 34, 487-515.

Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S., Coe, M. T., Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A., Kucharik, C. J., Monfreda, C., Patz, J. A., Prentice, I. C., Ramankutty, N., & Snyder, P. K. (2005). Global consequences of land use. Science, 309, 570-574.

Frankham, R. (2010). Inbreeding in the wild really does matter. Heredity, 104, 124.

Frishkoff, L. O., Hadly, E. A., & Daily, G. C. (2015). Thermal niche predicts tolerance to habitat conversion in tropical amphibians and reptiles. Global Change Biology, 21, 3901-3916.

Gibson, L., Lee, T. M., Koh, L. P., Brook, B. W., Gardner, T. A., Barlow, J., Peres, A. C., Bradshaw, C. J. A., Laurance, W. F., Lovejoy, T. E., & Sodhi, N. S. (2011). Primary forests are irreplaceable for sustaining tropical biodiversity. Nature, 478, 378.

Gilpin, M. (Ed.). (2012). Metapopulation dynamics: empirical and theoretical investigations. Academic Press.

Gomes, F. R., Rezende, E. L., Grizante, M. B. & Navas, C. A. (2009). The evolution of jumping performance in anurans morphological correlates and ecological implications Journal of Evolutionary Biology, 22, 1088-1097

Grant, E. H. C., Nichols, J. D., Lowe, W. H., & Fagan, W. F. (2010). Use of multiple dispersal pathways facilitates amphibian persistence in stream networks. Proceedings of the National Academy of Sciences, 201000266.

Greggor, A. L., Thornton, A., & Clayton, N. S. (2015). Neophobia is not only avoidance: improving neophobia tests by combining cognition and ecology. Current Opinion in Behavioral Sciences, 6, 82-89.

Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., Lovejoy, T. E., Sexton, J. O., Austin, M. P., Collins, C. D., Cook, W. M., Damschen, E. I., Ewers, R. M., Foster, B. L., Jenkins, C. N., King, A. J., Laurance, W. F., Levey, D. J., Margules, C. R., Melbourne, B. A., Nicholls, A. O., Orrock, J. L., Song, D. X., & Townshend, J. R (2015). Habitat fragmentation and its lasting impact on Earth's ecosystems. Science Advances, 1, e1500052.

Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A. A., Tyukavina, A., Thau, D., Stehman, S. V., Goetz, S. J., Loveland, T. R., Kommareddy, A., Egorov, A., Chini, L., Justice, C. O., & Townshend, J. R. G. (2013). High-resolution global maps of 21st-century forest cover change. Science, 342, 850-853.

Hanski, I., & Gaggiotti, O. E. (Eds.). (2004). Ecology, genetics, and evolution of metapopulations. Academic Press.

Hanski, I. A. (2011). Eco-evolutionary spatial dynamics in the Glanville fritillary butterfly.
Proceedings of the National Academy of Sciences, 108, 14397-14404.

Hanski, I., & Mononen, T. (2011). Eco☐evolutionary dynamics of dispersal in spatially heterogeneous
environments. Ecology Letters, 14, 1025-1034.

Hanski, I., Schulz, T., Wong, S. C., Ahola, V., Ruokolainen, A., & Ojanen, S. P. (2017). Ecological
and genetic basis of metapopulation persistence of the Glanville fritillary butterfly in fragmented
landscapes. Nature Communications, 8, 14504.

Hedrick, P. W. (2013). Adaptive introgression in animals: examples and comparison to new mutation
and standing variation as sources of adaptive variation. Molecular Ecology, 22, 4606-4618.

Hedrick, P. W., & Garcia-Dorado, A. (2016). Understanding inbreeding depression, purging, and
genetic rescue. Trends in Ecology & Evolution, 31, 940-952.

Hillman, S.S., Drewes, R.C., Hedrick, M.S. & Hancock, T.V. (2014). Physiological vagility and its
relationship to dispersal and neutral genetic heterogeneity in vertebrates. The Journal of
Experimental Biology, 217, 3356-3364.

Jakobsson, M., & Rosenberg, N. A. (2007). CLUMPP: a cluster matching and permutation program
for dealing with label switching and multimodality in analysis of population structure.
Bioinformatics, 23, 1801-1806.

Kaplan, R. H. (1987). Developmental plasticity and maternal effects of reproductive characteristics in
the frog, *Bombina orientalis*. Oecologia, 71, 273-279.

Kaplan, R. H. (1992). Greater maternal investment can decrease offspring survival in the frog
Bombina orientalis. Ecology, 73, 280-288.

Kelleher, S. R., Silla, A. J., Dingemanse, N. J., & Byrne, P. G. (2017). Body size predicts between-
individual differences in exploration behaviour in the southern corroboree frog. Animal behaviour,
129, 161-170.

Kelleher, S. R., Silla, A. J., & Byrne, P. G. (2018). Animal personality and behavioral syndromes in
amphibians: a review of the evidence, experimental approaches, and implications for conservation.
Behavioral Ecology and Sociobiology, 72, 79.

Kokko, H., & López-Sepulcre, A. (2006). From individual dispersal to species ranges: perspectives for
a changing world. Science, 313, 789-791.

Kubisch, A., Holt, R.D., Poethke, H.J. & Fronhofer, E.A. 2014. Where am I and why? Synthesising
range biology and the eco-evolutionary dynamics of dispersal. Oikos 123:5–22.

Gosner, K. L. (1960). A simplified table for staging anuran embryos and larvae with notes on
identification. Herpetologica, 16, 183-190.

Lagrange, P., Pradel, R., Bélisle, M., & Gimenez, O. (2014). Estimating dispersal among numerous
sites using capture–recapture data. Ecology, 95, 2316-2323.

Laugen, A. T., Kruuk, L. E., Laurila, A., Räsänen, K., Stone, J., & Merilä, J. (2005). Quantitative
genetics of larval life-history traits in Rana temporaria in different environmental conditions.
Genetics Research, 86, 161-170.

Lebreton, J. D., Nichols, J. D., Barker, R. J., Pradel, R., & Spendelow, J. A. (2009). Modeling
individual animal histories with multistate capture–recapture models. Advances in Ecological
Research, 41, 87-173.

Legendre P., & Legendre A. (1998) Complex ecological data sets. In: Legendre, P., Legendre, A. (eds)
Numerical Ecology. Pp. 247–302. Elsevier, Amsterdam, Netherlands.

Legrand, D., Cote, J., Fronhofer, E. A., Holt, R. D., Ronce, O., Schtickzelle, N., Travis, J. M. J., &
Clobert, J. (2017). Eco□evolutionary dynamics in fragmented landscapes. Ecography, 40, 9-25.

Lenormand, T. (2002). Gene flow and the limits to natural selection. Trends in Ecology & Evolution,
17, 183-189.

Littell, R. C., Milliken, G. A. , Stroup, W. W. , Wolfinger, R. D. (2006). SAS system for mixed
models 2nd ed. SAS Institute, Inc., Cary, North Carolina, USA.

Lowe, W. H. (2003). Linking dispersal to local population dynamics: a case study using a headwater
salamander system. Ecology, 84, 2145-2154.

Matthysen, E. (2012). Multicausality of dispersal: a review. In Clobert, J., Baguette, M., Benton, T.
G., & Bullock, J. M.: Dispersal Ecology and Evolution. Pp. 3–18, Oxford University Press, Oxford,
UK.

McCullagh, P., & Nelder, J. A. (1989). Generalized linear models. CRC press.

McPeek, M. A., & Holt, R. D. (1992). The evolution of dispersal in spatially and temporally varying
environments. The American Naturalist, 140, 1010-1027.

McRae, B. H., & Beier, P. (2007). Circuit theory predicts gene flow in plant and animal populations.
Proceedings of the National Academy of Sciences of the United States of America, 104, 19885–
19890.

Metzker, M. L. (2010). Sequencing technologies—the next generation. Nature Reviews Genetics, 11,
31.

Morozova, O., & Marra, M. A. (2008). Applications of next-generation sequencing technologies in
functional genomics. Genomics, 92, 255-264.

Newbold, T., Hudson, L. N., Hill, S. L., Contu, S., Lysenko, I., Senior, R. A., Börger, L., Bennett, D.
J., Choimes, A., Collen, B., Day, J., De Palma, A., Díaz, S., Echeverria-Londoño, S., Edgar, M. J.,
Feldman, A., Garon, M., Harrison, M. L. K., Alhusseini, T., Ingram, D. J., Itescu, Y., Kattge, J.,
Kemp, V., Kirkpatrick, L., Kleyer, M., Laginha Pinto Correia, D., Martin, C. D., Meiri, S.,
Novosolov, M., Pan, Y., Phillips, H. R. P., Purves, D. W., Robinson, A., Simpson, J., Tuck, S. L.,
Weiher, E., White, H. J., Ewers, R. M., Mace, G. M., Scharlemann, J. P. W., & Purvis, A. (2015).
Global effects of land use on local terrestrial biodiversity. Nature, 520, 45.

Newman, E. A. (2019). Disturbance ecology in the Anthropocene. Frontiers in Ecology and Evolution,
7, 147.

Nowakowski, A. J., Frishkoff, L. O., Thompson, M. E., Smith, T. M., & Todd, B. D. (2018).
Phylogenetic homogenization of amphibian assemblages in human-altered habitats across the
globe. Proceedings of the National Academy of Sciences, 115, E3454-E3462.

Ochocki, B. M., & Miller, T. E. X. (2017). Rapid evolution of dispersal ability makes biological
invasions faster and more variable. Nature Communications, 8, 14315.

Otto, S. P. (2018). Adaptation, speciation and extinction in the Anthropocene. Proceedings of the
Royal Society B, 285, 20182047.

Peakall, R. O. D., & Smouse, P. E. (2006). GENALEX 6: genetic analysis in Excel. Population genetic
software for teaching and research. Molecular Ecology Notes, 6, 288-295.

Pelletier, F., & Coltman, D. W. (2018). Will human influences on evolutionary dynamics in the wild
pervade the Anthropocene?. BMC Biology, 16, 7.

Pennekamp, F., Schtickzelle, N., & Petchey, O. L. (2015). BEMOVI, software for extracting behavior
and morphology from videos, illustrated with analyses of microbes. Ecology and Evolution, 5,
2584-2595.

Pereira, H. M., Leadley, P. W., Proença, V., Alkemade, R., Scharlemann, J. P., Fernandez-Manjarrés,
J. F., Araújo, M. B., Balvanera, P., Biggs, R., Cheung, W. M. L., Chini, L., Cooper, H. D., Gilman,
E. L., Guénette, S., Hurtt, G. C., Huntington, H. P., Mace, G. M., Oberdorff, T., Revenga, C.,
Rodrigues, P., Scholes, R. J., Sumaila, U. R., Walpole, M. (2010). Scenarios for global biodiversity
in the 21st century. Science, 330, 1496-1501.

Pollock, K. H. (1982). A capture-recapture design robust to unequal probability of capture. The
Journal of Wildlife Management, 46, 752-757.

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using
multilocus genotype data. Genetics, 155, 945-959.

Prunier, J. G., Kaufmann, B., Fenet, S., Picard, D., Pompanon, F., Joly, P., & Lena, J. P. (2013).
Optimizing the trade☐off between spatial and genetic sampling efforts in patchy populations:
Towards a better assessment of functional connectivity using an individual☐based sampling
scheme. Molecular Ecology, 22, 5516-5530.

Prunier, J. G., Colyn, M., Legendre, X., Nimon, K. F., & Flamand, M. C. (2015). Multicollinearity in
spatial genetics: separating the wheat from the chaff using commonality analyses. Molecular
Ecology, 24, 263-283.

Räsänen, K., Laurila, A., & Merilä, J. (2005). Maternal investment in egg size: environment-and
population-specific effects on offspring performance. Oecologia, 142, 546-553.

Reed, D. H., & Frankham, R. (2003). Correlation between fitness and genetic diversity. Conservation
Biology, 17, 230-237.

Roff, D. A., & Fairbairn, D. J. (2007). The evolution and genetics of migration in insects. AIBS
Bulletin, 57, 155-164.

Ronce, O. (2007). How does it feel to be like a rolling stone? Ten questions about dispersal evolution.
Annual Review of Ecology, Evolution, and Systematics, 38, 231-253.

Ronce, O., & Clobert, J. (2012). Dispersal syndromes. In Clobert, J., Baguette, M., Benton, T. G., &
Bullock, J. M.: Dispersal Ecology and Evolution. Pp. 119–138, Oxford University Press, Oxford,
UK.

Rousset, F. (2008). genepop'007: a complete re☐implementation of the genepop software for
Windows and Linux. Molecular Ecology Resources, 8, 103-106.

Saastamoinen, M., Bocedi, G., Cote, J., Legrand, D., Guillaume, F., Wheat, C.W., Fronhofer, E.A.,
Garcia, C., Henry, R., Husby, A., Baguette, M., Bonte, D., Coulon, A., Kokko, H., Matthysen, E.,
Niitepõld, K., Nonaka, E., Stevens, V.M., Travis, J.M., Donohue, K., Bullock, J.M., del Mar
Delgado, M. (2018) Genetics of dispersal. Biological Reviews, 93, 574-599.

Sala, O. E., Chapin, F. S., Armesto, J. J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-Sanwald, E.,
Huenneke, L. F., Jackson, R. B., Kinzig, A., Leemans, R., Lodge, D. M., Mooney, H. A.,
Oesterheld, M., LeRoy Poff, N., Sykes, M. T., Walker, B. H., Walker, M., & Wall, D. H. (2000).
Global biodiversity scenarios for the year 2100. Science, 287, 1770-1774.

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients.
Methods in Ecology and Evolution, 1, 103-113.

Selkoe, K. A., Watson, J. R., White, C., Horin, T. B., Iacchei, M., Mitarai, S., Siegel, D.A., Gaines,
S.D. & Toonen, R. J. (2010). Taking the chaos out of genetic patchiness: Seascape genetics reveals
ecological and oceanographic drivers of genetic patterns in three temperate reef species. Molecular
Ecology, 19, 3708–3726.

Slatkin, M. (1977). Gene flow and genetic drift in a species subject to frequent local extinctions.
Theoretical Population Biology, 12, 253-262.

Thomas, C. D., & Kunin, W. E. (1999). The spatial structure of populations. Journal of Animal
Ecology, 68, 647-657.

Tilman, D., Clark, M., Williams, D. R., Kimmel, K., Polasky, S., & Packer, C. (2017). Future threats
to biodiversity and pathways to their prevention. Nature, 546, 73.

Tournier, E., Besnard, A., Tournier, V., & Cayuela, H. (2017). Manipulating waterbody hydroperiod
affects movement behaviour and occupancy dynamics in an amphibian. Freshwater Biology, 62,
1768-1782.

Travis, J.M.J., Mustin, K., Benton, T.G. & Dytham, C. 2009. Accelerating invasion rates result from
the evolution of density-dependent dispersal. Journal Theoretical Biology, 259, 151–158.

Turner, M. G. (2010). Disturbance and landscape dynamics in a changing world. Ecology, 91, 2833-
2849.

Van Oosterhout, C., Hutchinson, W. F., Wills, D. P., & Shipley, P. (2004). MICRO☐CHECKER:

software for identifying and correcting genotyping errors in microsatellite data. Molecular Ecology

Notes, 4, 535-538.

Van Strien, M. J., Keller, D., & Holderegger, R. (2012). A new analytical approach to landscape

genetic modelling: Least☐cost transect analysis and linear mixed models. Molecular Ecology, 21,

4010–4023.

Villard, M. A., & Metzger, J. P. (2014). Beyond the fragmentation debate: a conceptual model to

predict when habitat configuration really matters. Journal of Applied Ecology, 51(2), 309-318.

Wang J (2011) COANCESTRY: a program for simulating, estimating and analysing relatedness and

inbreeding coefficients. Molecular Ecology Resources, 11, 141–145.

Warren, S. D., & Büttner, R. (2008). Relationship of endangered amphibians to landscape disturbance.

The Journal of Wildlife Management, 72, 738-744.

Yuen, C. H., Schoepf, I., Schradin, C., & Pillay, N. (2017). Boldness: are open field and startle tests

measuring the same personality trait?. Animal Behaviour, 128, 143-151.

Zaghlool, S. B., Al-Shafai, M., Al Muftah, W. A., Kumar, P., Gieger, C., Waldenberger, M., Falchi,

M., & Suhre, K. (2016). Mendelian inheritance of trimodal CpG methylation sites suggests distal

cis-acting genetic effects. Clinical Epigenetics, 8, 124.

1370