Access to unexplored regions of sequence space in directed enzyme evolution *via* insertion/deletion mutagenesis

Stephane Emond[1,*], Maya Petek[1], Emily Kay[1], Brennen Heames[1], Sean Devenish[1], Nobuhiko Tokuriki[2] and Florian Hollfelder[1,*]

[1] Department of Biochemistry, University of Cambridge, Cambridge CB2 1GA, UK

[2] Michael Smith Laboratories, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

* To whom correspondence should be addressed. Tel: +44 (0)1223 766048 Email: fh111@cam.ac.uk.
Correspondence may also be addressed to Stephane Emond. Email: emond.stephane@gmail.com.

Present addresses:

Stephane Emond, Evonetix Ltd, Chesterford Research Park, Little Chesterford CB10 1XL, UK.

Emily Kay, Cancer Research UK Beatson Institute, Glasgow G61 1BD, UK.

Brennen Heames, Institute for Evolution and Biodiversity, Westfälische Wilhelms-Universität, Hüfferstrasse 1, 48149 Münster, Germany.

Sean Devenish, Fluidic Analytics, The Paddocks Business Centre, Cherry Hinton Road, Cambridge, CB1 8DH, UK.

ABSTRACT

Insertions and deletions (InDels) are frequently observed in natural protein evolution, yet their potential remains untapped in laboratory evolution. Here we introduce a transposon mutagenesis approach (TRIAD) to generate libraries of random variants with short in-frame InDels, and screen TRIAD libraries to evolve a promiscuous arylesterase activity in a phosphotriesterase. The evolution exhibits features that are distinct from previous point mutagenesis campaigns: while the *average activity* of TRIAD variants is more deleterious, a *larger proportion* has successfully adapted for the new activity, exhibiting different functional profiles: (*i*) both strong and weak trade-off in original vs promiscuous activity are observed; (*ii*) trade-off is more severe (10- to 20-fold increased k_{cat}/K_M in arylesterase with ~100-fold decreases in the original phosphotriesterase activity) and (*iii*) improvements show up in k_{cat} rather than K_M, suggesting novel adaptive solution. These distinct features make TRIAD an alternative to widely used point mutagenesis, providing access to functional innovations and traversing unexplored fitness landscape regions.

INTRODUCTION

Directed evolution aims at identifying proteins with new functional traits by mimicking the natural process of genetic variation through mutations, followed by selection of improved variants. A major challenge for the success of this approach remains that only a very small fraction of the theoretically and practically accessible sequence space can be sampled during any screening or selection process. As such, expanding the diversity and the quality of gene libraries has been a major research focus to increase the chances of identifying new variants with desired functions. So far, most directed evolution (and, more generally, protein engineering) experiments have been performed using point substitutions for gene diversification, while insertions or deletions (InDels) remain an overlooked source of variation despite their frequent and functionally beneficial occurrence in natural protein evolution (1). Combinatorial approaches to incorporate InDels at predefined positions, based on phylogenetic and/or structural analyses, have been developed to alter catalytic specificities of enzymes (2-5) or to improve the binding affinities of engineered antibodies (6). Several methods for incorporating InDels randomly within a gene of interest have been developed but show many limitations in terms of library quality and diversity. Most of these approaches generate frame-shifting InDels at high frequency (>66%) (*e.g.*, using error-prone DNA polymerases (7,8), terminal deoxynucleotidyl transferase (9), exonucleases (10,11), tandem duplication insertion (12) or truncation (13)), resulting in libraries consisting mostly of non-functional variants, which must be removed by high-throughput selection or screening. Methods based on the use of engineered transposons are designed to avoid frameshifts but, so far, have been limited to the generation of deletions (14,15) or insertions of fixed length and defined sequences (16).

In the present work, a set of procedures (dubbed TRIAD: Transposition-based Random Insertion And Deletion mutagenesis) for random introduction of single short in-frame InDels of one, two or three nucleotide triplets (± 3, 6 or 9 bp) into a given DNA sequence was established and validated by generating libraries of InDel variants of *Brevundimonas diminuta* phosphotriesterase (*wt*PTE), a highly efficient paraoxon-hydrolyzing enzyme (17) with promiscuous esterase and lactonase activities (18). The resulting TRIAD libraries were used to investigate the fitness effects of InDels on *wt*PTE and compare it to that of substitutions. Moreover, screening these libraries for improved arylesterase activity revealed several hits that would have been inaccessible using traditional point substitution mutagenesis approaches, demonstrating that the introduction of InDels can harvest functional diversity in previously unexplored areas of protein sequence space.

MATERIALS AND METHODS

Materials

Paraoxon, 4-nitrophenyl butyrate (pNPB), 1-naphthyl butyrate (1-NB), 2-naphthyl hexanoate (2-NH), and Fast Red were purchased from Sigma. FastDigest restriction endonucleases, MuA transposase and T4 DNA ligase were purchased from Thermo Fisher Scientific. DNA Polymerase I, Large (Klenow) Fragment was purchased from New England Biolabs. All DNA modifying enzymes were used according to the manufacturer's conditions.

Gene synthesis and cloning experiments

Detailed procedures and sequences can be found in the online Supplementary Information for the design and construction of transposons (TransDel and TransIns), cloning cassettes (Del2, Del3, Ins1, Ins2 and Ins3) and vectors (pID-T7, pID-Tet). The *wt*PTE gene lacking MlyI and AcuI sites was synthesised by GenScript (NJ, USA). InDel libraries of *wt*PTE prepared in the pID-Tet vector were subcloned with NcoI and HindIII into pET-strep vector (19) to express the strep-tag–PTE fusion protein for screening experiments and purification for the enzyme kinetics and stability assay. The trinucleotide substitution library of *wt*PTE used to compare the functional impact of InDels *vs.* point substitutions was generated following the TriNEx method (20) as described previously (21).

Construction of InDel variant libraries of wtPTE

(1) Generation of transposon insertion libraries. The transposons TransDel and TransIns were extracted from pUC57 by BglII digestion and recovered by gel electrophoresis and purification. Insertion of TransDel or TransIns in the pID-Tet plasmid containing *wt*PTE was performed using *in vitro* transposition using 300 ng of plasmid, 50 ng of transposon and 0.22 µg MuA transposase in a 20 µL reaction volume. After incubation for 2 h at 30°C, the MuA transposase was heat-inactivated for 10 min at 75 °C. DNA products were purified and concentrated in 7 µL deionized water using a DNA clean concentrator kit (Zymo Research). 2 µL of the purified DNA was used to transform *E. coli* E. cloni® 10G cells (> 10^{10} CFU/µg pUC19; Lucigen) by electroporation. The transformants (typically 10,000 to 50,000 CFU) were selected on LB agar containing ampicillin (amp; 100 µg/mL) and chloramphenicol (cam; 34 µg/mL). The resulting colonies were pooled, and their plasmid DNA extracted. The fragments corresponding to *wt*PTE containing the inserted transposon were obtained by double restriction digestion (NcoI/HindIII) followed by gel extraction and ligated in pID-Tet (50-100 ng). The ligation products were then transformed into electrocompetent *E. coli* E. cloni® 10G cells. Upon selection on LB-agar-amp-cam, transformants (generally 1 to $2x10^6$ CFU) were pooled and their plasmid DNA extracted, yielding transposon (either TransDel or TransIns) insertion libraries.

(2) Generation of libraries of deletion variants. The construction of libraries with triplet deletion variants (-3 bp) was performed as previously described (14). TransDel insertion library plasmids were first digested with MlyI to remove TransDel. The fragments corresponding to linear pID-Tet-*wt*PTE plasmids (with a -3 bp deletion in *wt*PTE) were isolated by gel electrophoresis and purified. Self-

circularization was then performed using T4 DNA ligase (Thermo Scientific) and 10-50 ng linearized plasmid (final concentration: ≥ 1 ng/µL). Upon purification and concentration, the ligation products were transformed into electrocompetent *E. coli* Ecloni® 10G cells subsequently selected on LB-agar-amp, yielding a library of gene of interest variants with -3bp random deletions. For the construction of libraries of -6 and -9 bp deletion variants, cassettes Del2 and Del3 were ligated into the MlyI linearized plasmid (50-100 ng) in a 1:3 molar ratio. After purification and concentration, these ligation products were transformed into electrocompetent *E. coli* Ecloni® 10G. The transformants (generally 1-3 x 10^6 colony forming units, CFU) were selected on LB agar containing ampicillin (100 µg/L) and kanamycin (Kan; 50 µg/mL). The plasmids (corresponding to Del2 and Del3 insertion libraries) were extracted from the colonies and subsequently digested using MlyI to remove the Del2 and Del3 cassettes. The resulting linear pID-Tet-*wt*PTE products (containing the gene of interest with -6 or -9 bp deletions) were recovered by gel electrophoresis, purified and subsequently self-circularized. The resulting products were transformed into electrocompetent *E. coli* Ecloni® 10G cells subsequently plated on LB-agar-amp, yielding libraries of *wt*PTE variants with -6 bp or -9 bp random deletions. All libraries were purified and stored in the form of plasmid solutions.

(3) Generation of libraries of insertion variants. TransIns insertion library plasmids were digested with NotI and MlyI to remove TransIns. The linearized pID-Tet-*wt*PTE plasmids were recovered by gel electrophoresis and purification. Cassettes Ins1, Ins2 and Ins3 were then inserted in the pID-Tet-*wt*PTE plasmid (50-100 ng) in a 1:3 molar ratio. After purification and concentration, these ligation products were transformed into electrocompetent *E. coli* Ecloni® 10G and the transformants (generally 1.10^6 to 3.10^6 CFU) were selected on LB-agar-Amp-Kan. After extraction from the resulting colonies, the plasmids corresponding to Ins1, Ins2 and Ins3 insertion libraries were digested with AcuI. The linearized pID-Tet-*wt*PTE plasmids (with an insertion of 3, 6 and 9 bp in *wt*PTE) were recovered by gel electrophoresis, purified and subsequently treated with the Klenow fragment of DNA Polymerase I to remove 3' overhangs created by AcuI digestion. After that blunting step, the plasmids were self-circularized. The resulting products were transformed into electrocompetent *E. coli* E. cloni® 10G cells subsequently plated on LB-agar-amp, yielding libraries of *wt*PTE variants with +3, +6 or +9 bp random insertions. All the libraries were purified and stored in the form of plasmid solutions.

Sequencing analysis of *wt*PTE libraries

The mutagenesis efficiency of TRIAD was analysed both by Sanger sequencing (Supplementary Tables S1-3) and deep sequencing. For the sequencing of individual *wt*PTE InDel variants obtained upon the transformation of libraries into *E. coli* (see above), individual colonies (~20 per library; Supplementary Tables S1 and S2) were randomly picked for plasmid extraction and subsequent Sanger sequencing. For deep sequencing, libraries were digested from pID-Tet with FastDigest restriction enzymes Bpu1102I and Van91I to give a pool of 1.3 kb linear fragments, which were processed using Nextera DNA Library Preparation Kit according to manufacturer's instructions and sequenced on Illumina MiSeq using 2x75 bp paired-end sequencing. The reads were de-multiplexed, adaptors trimmed and assembled using PEAR (22). Assembled and unassembled reads were

mapped to the reference using Bowtie2 (23) and re-aligned to reference using the accurate Needleman-Wunsch algorithm with gap open penalty 15 and gap extend penalty 0.5 (24). Placing InDels in particular sequence contexts may be inherently ambiguous because of potential InDel redundancy: when two or more InDels inserted at *different* positions in the target gene result in *identical* final sequence, no algorithm will be able to distinguish between them and the resulting InDel is always assigned to a *single* arbitrarily chosen original insertion or deletion site (see the discussion of examples in the Supplementary Methods 2.7). No attempt was made to correct for such ambiguity at this point. Resulting alignments were used to count the number of reads in which the mutations occur, their type and position using in-house developed Python scripts (see Supplementary Methods 2.4 and 2.5). To analyse the sequence preference for TransDel transposition, the counts were corrected for codon ambiguity by dividing the observed count equally between all positions where the deletion could have originated.

Screening procedures for *wt*PTE libraries

Prior to screening, InDel variant libraries of *wt*PTE were excised by NcoI/HindIII double digestion and subcloned into pET-Strep vector. The resulting libraries were transformed into *E. coli* BL21 (DE3) containing pGro7 plasmid for overexpression of the GroEL/ES chaperone system. Transformed cells (typically 2000-10000 CFU) were plated on agar-amp-cam. The colonies were replicated using a filter paper (BioTrace NT Pure Nitrocellulose Transfer Membrane 0.2 μm, PALL Life Sciences), which was placed onto a second plate containing IPTG (1 mM), $ZnCl_2$ (200 μM) and arabinose (20% (w/v)) for chaperone overexpression. After overnight expression at room temperature, the filter paper was placed into an empty Petri dish and cells were lysed prior to the activity assay by alternating three times between storage at -20 °C and 37 °C. Subsequently, top agar (0.5% agar in 100 mM Tris–HCl pH 7.5) containing either 1-NB or 2-NH (200 μM) and FAST Red (200 μM) was layered and a red precipitate (resulting from the complex formation between Fast Red and the naphthol product) developed within ~30 minutes. Colonies expressing an active PTE variant were picked, transferred in 96-well plates containing 200 μL LB-Amp-Cam per well and re-grown overnight at 30°C. Subsequently, 25 μL of the resulting cultures were used to inoculate 425 μL LB-Amp-Cam arabinose (20% (w/v)) in deep 96-well plates. After growth for 2 to 3 h at 30°C, expression of PTE variants was induced by adding IPTG (1 mM final concentration) and cultures were incubated for an additional 2 hours at 30 °C. Cells were then pelleted by centrifugation at 4 °C at maximum speed (3320×g) for 5-10 minutes and the supernatant removed. Pellets were frozen overnight at -80°C and, after thawing, lysed in 200 μL 50 mM Tris-HCl pH 7.5 supplemented with 0.1% (w/v) Triton-X100, 200 μM $ZnCl_2$, 100 μg/mL lysozyme and 0.8 U/mL benzonase (Novagen). After 30 minutes of lysis, cell debris were spun down at 4 °C at 3320×g for 20 minutes. Enzyme assays were performed in 96-well plates containing a volume of 200 μL per well (20 μL pre-diluted lysate + 180 μL of 200 μM substrate in 50 mM Tris-HCl, pH 7.5 supplemented with Triton-X100 (0.02% in the case of paraoxon and 0.1% in the case of 2-NH/FR). The hydrolysis of paraoxon and 4-NPB were monitored by absorbance readings at 405 nm. The complex formation between 2-Naphthol and Fast Red was monitored at 500 nm.

Purification of Strep-tagged PTE variants

pET-Strep-PTE plasmids were transformed into *E. coli* BL21 (DE3) grown for 8 h at 30 °C in Overnight Express Instant TB medium (Novagen) containing 100 μg/mL ampicillin and 200 μM ZnCl2 before lowering the temperature to 16 °C and continuing incubation overnight. Cells were harvested by centrifugation, resuspended and lysed using a 1:1 mixture of B-PER Protein Extraction Reagent (Thermo Scientific: 50 mM Tris-HCl buffer, pH 7.5 containing 200 μM $ZnCl_2$, 100 μg/mL lysozyme and approx. 1 μL of benzonase per 100 mL. Cell debris was removed by centrifugation and the clarified lysate passed through a 45 μm filter before loading onto a Strep-Tactin Superflow High capacity column (1 mL). Strep-PTE variants were eluted with Elution buffer (100 mM Tris-HCl, pH 7.5, 200 μM $ZnCl_2$ and 2.5 mM desthiobiotin) according to the manufacturer's instructions (IBA Lifesciences).

Kinetic characterization of PTE variants

Initial velocities (v_0) were determined at 12 different substrate concentrations in the range of 2–2,000 μM in Tris-HCl (100 mM, pH 7.5) and $ZnCl_2$ (200 μM). Rates of reaction were monitored by following the complex formation between the product and FastRed at 500 nm for 2-NH hydrolysis (in the presence of 2 mM FAST Red) and product formation at 405 nm for paraoxon hydrolysis. Purified protein was diluted to a concentration of 10 nM for paraoxon and 500 nM for 2-NH. K_M and k_{cat} were determined by fitting the initial rates at each concentration to the Michaelis-Menten model using KaleidaGraph (Synergy Software).

RESULTS

A strategy for creation of random InDel libraries

TRIAD consists of a single transposition reaction followed by successive cloning steps for the generation of deletions or insertions (Figure 1; see also Supplementary Figure S1 for a more detailed illustration). TRIAD's first step is an *in vitro* Mu transposition reaction (25) that ultimately determines the location of the forthcoming single InDel event in each variant. The reaction is performed using engineered mini-Mu transposons, dubbed TransDel and TransIns (Supplementary Figure S2A), that are inserted randomly within the target DNA sequence during the first step of TRIAD; resulting in the generation of transposon insertion libraries. The ends of TransDel and TransIns were designed to bring about deletion and insertion libraries, respectively. TransDel is functionally equivalent to the previously described MuDel transposon (14) with recognition sites for the type IIS restriction enzyme MlyI at both ends. The positioning of MlyI sites within TransDel enables the deletion of 3 bp at random positions within the target sequence upon MlyI digestion and self-ligation (Figure 2A), as previously described (14). This strategy was extended to the generation of longer contiguous deletions (*i.e.*, -6 and -9 bp) with a second stage, involving the insertion and subsequent MlyI-mediated removal of custom-made cassettes (dubbed Del2 and Del3; Figure 1A and Figure 2A). For the generation of insertions, a new transposon, TransIns, was designed as – in contrast to TransDel – an *asymmetric* transposon (Figure 1B and Figure 2B**)**, bearing different end sequences (NotI on one end and MlyI on the other). The latter site marks subsequent insertion sites for the ligation of custom-made shuttle cassettes: Ins1, Ins2 and Ins3 carrying one, two and three randomized nucleotide triplets, respectively. Further digestion using a type IIS restriction enzyme (AcuI) removes most of the shuttle sequence but leaves triplet insertions behind (Figure 1B and Figure 2B).

Generation of random InDel libraries by TRIAD

To validate TRIAD, we generated InDel libraries from the gene encoding a highly expressed variant of phosphotriesterase (*wt*PTE) that had been previously used as starting point to generate an efficient arylesterase by laboratory evolution (26,27). To enable TRIAD, any recognition sequences for MlyI, NotI and AcuI in the target sequence or the plasmid containing the target sequence must be removed. A synthetic gene encoding *wt*PTE (Supplementary Figure S3) as well as dedicated cloning vectors (namely pID-T7 and pID-Tet; Supplementary Figure S4 and Supplementary Methods) were therefore designed and assembled prior to the construction of libraries. The generation of transposition insertion libraries with TransDel or TransIns was performed as previously described (28). Briefly, *in vitro* transposition was performed to integrate TransDel or TransIns (~1 kbp) within the plasmid (pID-Tet, ~2.7 kbp) carrying the *wt*PTE synthetic gene (~1 kbp). Transformation of the transposition products into electrocompetent *E. coli* yielded >30,000 colonies per transposition reaction. The fraction of transformants with the transposon inserted into *wt*PTE (~27% of the entire plasmid length) corresponds to >8,000 colonies, corresponding to >8-fold coverage of possible insertion sites (~1,000) within *wt*PTE. The transposon is inserted randomly throughout the entire plasmid, so the fragments corresponding to the target sequence carrying the transposon (~2 kbp) were isolated by

restriction digestion and subcloned back into intact pID-Tet, thereby generating TransDel and TransIns insertion libraries. At this stage, transformation of these libraries into *E. coli* typically yielded >10^6 colonies, maintaining oversampling of insertion sites without skewing the distribution due to sampling.

The TransDel and TransIns insertion libraries were then used as starting material to generate six independent libraries of *wt*PTE InDel variants: three deletion (-3, -6 and -9 bp) and three insertion libraries (+3, +6 and +9 bp). The theoretical diversity (*i.e.*, the total number of possible variants) accessible *via* such modifications will depend both on the type of InDel that is introduced and on the target sequence (Supplementary Figure 5). For instance, as deletions can occur once at each position of the target DNA sequence, the maximal possible theoretical diversity of deletion libraries is identical to the number of nucleotides (Supplementary Figure S5A), *e.g.*, ~1,000 possible deletion variants for a ~1 kbp target sequence (*e.g.*, *wt*PTE). By contrast, since TRIAD inserts degenerate nucleotide triplets (*e.g.*, [NNN]$_{1,\ 2\ \text{or}\ 3}$ corresponding to 64 (=4^3), 4,096 (=4^6) and 262,144 (=4^9) possible sequences, respectively), the number of possible insertion variants will depend both on the length of the target sequence and the size of the insertions. The maximal possible theoretical diversity for insertion libraries generated from *wt*PTE is 6.4×10^4 (=64×10^3), ~4.1×10^6 (=64^2×10^3), and ~2.6×10^8 (=64^3×10^3), corresponding to +3, +6 and +9 bp insertions, respectively (Supplementary Figure S5A). Because of potential InDel redundancy depending on the target sequence (*i.e.*, two or more neighbouring InDels can result in the same DNA variant; Supplementary Figure S5B), the theoretical diversities accessible from a given DNA sequence are usually lower (see Supplementary Figure S5C in the specific case of *wt*PTE). Theoretical diversities at the protein level (*i.e.*, the number of protein variants that have the intended InDel length) are further reduced due to codon degeneracy and occurrence of stop codons as a result of certain InDels (Supplementary Figure S5C). Practically, the size of our libraries was limited by transformation efficiency, achieving > 10^6 variants upon transformation into *E. coli.* Therefore, deletions as well as +3 bp insertions were oversampled such that the library diversity was maintained between transformations, while the diversity of sampled transposition sites was maintained in larger +6 bp and +9 bp insertion libraries, with only a fraction of theoretical library diversity generated from the outset.

Quality assessment of TRIAD libraries

First, the accuracy of the TRIAD approach (specifically the number of intended in-frame InDels, unwanted frameshifts and incidental mutations) was assessed using Sanger sequencing to analyse individual *wt*PTE InDel variants. After the final transformation step into *E. coli*, around 20 colonies from each naïve InDel library of individual *wt*PTE variants were randomly picked and sequenced (121 variants in total; Supplementary Tables S1-3). All the sequenced variants displayed only a single modification resulting from the initial transposon insertion and 90 among them (74%; corresponding to 89 unique InDels) showed anticipated in-frame InDel mutations (86% of the deletion variants; 61% of the insertion variants; Supplementary Tables S1-2). Most in-frame InDels were observed only once and were distributed throughout the *wt*PTE sequence (Supplementary Table S3). No frameshift was observed among sequenced variants from the -3 bp library, which is generated without shuttle cloning

steps in contrast to the other libraries. Frameshifts were more frequent among variants from the +3, +6 and +9 bp insertion libraries (~40% of the sequenced variants). Higher frameshift frequency in insertion libraries may be due to exonuclease over-digestion by the Klenow fragment of DNA polymerase I, which removes 3' overhangs left by AcuI digestion (Figure 2B). Note that no incidental additional base pair point mutations located elsewhere in the variants' sequence (*i.e.*, at positions different to that of the initial transposon insertion sites) and resulting from the TRIAD cloning process were detected. On the protein level, TRIAD may generate a secondary point substitution contiguous to the introduced InDel (14), depending on the point of insertion of TransDel or TransIns in the reading frame of the target sequence. This occurs when the InDel is not inserted at previous codon boundaries (statistically in two of three cases, although not all such events lead to amino acid substitution). As a result, 22% of the InDels observed in individual *wt*PTE variants exhibited such an adjacent substitution.

While the accuracy of the TRIAD procedure could be assessed based on a small number of sequences (n = 121, giving 89 unique in-frame InDels), deep next-generation sequencing is required to gain quantitative insight into the distribution of InDels over the target sequence and assess the coverage afforded by the initial transposition step prior to diversification *via* InDel mutagenesis (Table 1). Deep sequencing yielded a much expanded dataset containing ~1×10^6 total 75-bp reads per deletion library and $>3\times10^6$ reads per insertion library, 95% of which aligned to the reference sequence (Supplementary Methods; Supplementary Figure S6 and Supplementary Table S4). The reads were assembled, aligned and parsed to extract and count mutations to determine the composition of libraries. In all libraries, the targeted in-frame InDels were found in high abundance, reaching more than 10^5 variants detected by deep sequencing in the most diverse +6 bp and +9 bp libraries (> 10^3 unique deletions and > 10^5 unique insertions overall; Table 1). In agreement with the sequencing of individual variants (Supplementary Table S1), frameshifts were rare in the -3 bp deletion library (4%) and more frequent (>20%) in the others (Supplementary Table S4B). Likewise, on the protein level, an average of 39% of the InDels observed in the deep sequencing dataset of *wt*PTE variants exhibited an adjacent amino acid substitution (Table 1).

The in-frame InDels detected by deep sequencing were spread throughout *wt*PTE's DNA sequence (Figure 3), although the exact sampling of positions may depend on the initial transposon (either TransDel or TransIns) used to generate the libraries. Analysis of -3 bp and +3 bp libraries showed that TransDel and TransIns insertion accessed 85% and 95% of all possible DNA positions, respectively. An analysis of Mu transposon target site preference (29) suggests a strong preference for pyrimidines in position 2 and purines in position 4 of the 5 bp transposition site, based on 806 observed transpositions. Here we show that the sequence bias of Mu transposons (see insert in Figure 3A; Supplementary Table S5) is less pronounced than previously thought (*i.e.*, 5'N-Py-G/C-Pu-N) (29,30). Comparative frequencies were observed for most deletions (with 52% of all detected deletions having between 10 and 99 reads per variant; Supplementary Figure S7), with 213 individual deletion variants or 11% of all deletions (Supplementary Table S7) occurring more frequently more across all three libraries combined (200 reads or more per variant; see distribution in Supplementary Figure S7). Good coverage of possible positions in the insertion libraries translates into high diversity

at most positions in *wt*PTE (Figure 3C; Supplementary Figure S8A): 10 or more distinct insertions were observed at between 66% (+3 bp) and 80% (+6 and +9 bp) of positions; furthermore, 100 or more insertions were detected in 34% (+6 bp) and 31% (+9 bp) of positions (Supplementary Figure S8B). While insertion libraries were sequenced with a higher loading onto the flow cell, this was still insufficient to fully capture the diversity in the +6 bp and +9 bp libraries (24% and 2% at the protein level, respectively; Table 1), where each variant was observed only once or twice (Supplementary Figure S7), and so the true diversity may be higher. Note that no significant bias was observed in the nucleotide composition of the in-frame insertions (Supplementary Figure S9), indicating that TRIAD generates diverse insertion variants.

Overall, our quality assessment of the TRIAD libraries showed that beyond a weak bias during transposon insertion, TRIAD libraries show excellent coverage of > 85% of positions in the DNA sequence of *wt*PTE. These results confirm that the TRIAD approach enabled the generation of diverse and large libraries of InDel variants through a set of simple cloning procedures that spanned just over 5 days (Supplementary Figure S1).

Comparison of fitness effects between InDels and point substitutions

To compare the distribution of fitness effects of InDels *vs.* point substitutions, the levels of native phosphotriesterase (PTE; substrate: paraoxon; Figure 4) and promiscuous arylesterase (AE; substrate: 4-nitrophenyl butyrate, 4-NPB; Figure 4) activities were determined for several hundred *wt*PTE variants from each TRIAD library and from a trinucleotide substitution library (Figure 5; Supplementary Tables S8-9). Considering *wt*PTE is an evolutionarily "optimized" enzyme as a phosphotriesterase (based on the observation that it is operating near the diffusion limit for its native activity (17)), it is to be expected that very few mutations would be beneficial and that InDels are more deleterious than point substitutions overall. This expectation is underlined by the observation that 83% of deletions and 77% of insertions are strongly deleterious (<0.1 PTE activity), compared to only 24% in the substitution library (Figure 5A). The average fitness change similarly favours substitutions and is an order of magnitude more deleterious for InDels (Figure 5C). However, of 485 screened deletions and 351 insertions, a total of 12 were beneficial (>1.5-fold PTE activity increase) against a background of already high catalytic efficiency. By contrast, no beneficial substitutions were found amongst the 342 substitutions screened. Similar frequencies were observed with respect to deleterious fitness changes induced by InDels *versus* point substitutions in *wt*PTE's promiscuous arylesterase activity, with 76% of deletions and 62% of insertions strongly deleterious in comparison to only 19% of substitutions (Figure 5B; Supplementary Table S8). The frequency of InDels beneficial for arylesterase activity was found to be at least 3-fold higher than that of beneficial substitutions (6% and 7.7% for deletions and insertions, respectively *versus* 1.8% for substitutions; Figure 5B).

Mapping the observed mutations to the 3D structure of *wt*PTE provided insight into the location of InDels in comparison with point substitutions. While substitutions selected for ≥ 50% of *wt*PTE activity are found throughout the protein, the positions of InDels triggering similar functional effect appear more clustered in loops and on the surface (Figure 5D). Analysis of surface-accessible solvent area (SASA) suggests that mutations affecting the buried residues are more detrimental than surface-

exposed ones (Supplementary Figure S10 and Supplementary Table S10). This observation holds for both InDels *and* substitutions. For substitutions, the correlation between SASA and fitness effects on activity is weak, while only ~20% of neutral or beneficial InDels affect buried residues (cf. ~40% of substitutions), readily explained by the larger impact of InDels on presumably optimised packing in the protein core.

Screening and identification of adaptive InDels in *wt*PTE

To demonstrate that TRIAD libraries allow access to functional innovation *via* adaptive InDels, all the libraries generated from the full-length *wt*PTE gene (six libraries in total: -3, -6, -9, +3, +6 and +9 bp) were subjected to two parallel screening campaigns to identify variants with enhanced arylesterase activity against either 4-nitrophenyl butyrate (4-NPB) or 2-naphthyl hexanoate (2-NH) (Figure 4). Both screening campaign consisted of a general two-step assay workflow. Upon transformation of the TRIAD libraries into *E. coli*, the resulting colonies (around 1 to 3×10^4 per library) were first screened for either 1-naphthyl butyrate (prior to subsequent screening against 4-NPB in crude cell lysates) or 2-NH hydrolysis (using the FAST Red indicator that reacts with the released naphthol product). Colonies expressing an active variant (300 to 600 per library) were subsequently grown, lysed and tested for enzymatic activity (for either 4-NPB or 2-NH) in 96-well plates. Note that screening assays on colonies and in cell lysates were both performed after expression of *wt*PTE variants in the presence of overexpressed GroEL/ES chaperonin as described previously (19).

Overall, 81 hits (55 insertions and 26 deletions) were identified based on improved arylesterase activity against 2-NH or 4-NPB in cell lysates, with increases ranging from 2- to 140-fold in lysate activity compared to *wt*PTE (Table 2; Supplementary Table S11). In contrast with the adaptive substitutions previously identified (27), these adaptive InDels appeared to have a more drastic effect on the native phosphotriesterase activity, indicating a more severe trade-off *on average* between maintaining original and enhancing promiscuous activity (average specificity ratio ~ 260; Supplementary Figure S11). However, numerous *individual* mutants that do not show such strong negative trade-off were also identified (*e.g.*, 64 variants out of 81 showed a specificity ratio < 100; Figures 6A-B; Supplementary Figure S12).

Sequence analysis of the nature and the location of the InDels responsible for the improvement in arylesterase activity (Table 2) showed that all the adaptive InDels (apart from one double triplet insertion, *e.g.*, V99G/Q99aI99b) were clustered in two flexible regions of *wt*PTE, namely loop 7 (residues L252 to Q278) and loop 8 (residues S299 to P322)(Figure 6C). Activity against 2-NH was improved by single InDels present in both loops while activity against 4-NPB was essentially enhanced by InDels clustered in loop 7 (Table 2; Supplementary Figure S13)**.** Unexpectedly, the best variant (10.5-fold improvement in AE) found in the -9 bp deletion library exhibited a 12 bp deletion (presumably as a result of a rearrangement during the transposition step in the TRIAD process) resulting in a four-amino acid residue deletion (*i.e.*, ΔA270-G273).

To further demonstrate that the identified InDels genuinely improve the arylesterase activity of *wt*PTE, the four variants exhibiting the strongest improvement against the 2-NH substrate (*i.e.*, ΔA270-G273, P256R/G256aA256b, S256aG256b and G311a) were purified and characterized to give

a 10- to 20-fold increased k_{cat}/K_M for 2-NH, while decreasing paraoxon hydrolysis by around 100-fold (Figure 6B; Table 3). The quantitation of the improvements varies between cell lysates and purified protein (*e.g.*, 140-fold in cell lysates versus 14-fold with the purified protein for P256R/G256aA256b), which may be ascribed to variation in expression levels.

DISCUSSION

A simple and versatile approach to generate libraries of short InDel variants

Access to reliable methods for gene randomization is essential for success in directed evolution experiments. Popular and practically useful methods must meet several key requirements: a good library generation protocol should create a large number of variants, avoid bias in gene composition or type of variant introduced, and be technically straightforward. When it comes to amino acid substitutions, several approaches (e.g. error-prone PCR, site-saturation mutagenesis starting with synthetic oligonucleotides) have been developed that partially or fully meet these criteria. In contrast, the use of InDels in directed evolution experiments has been curtailed by practical limitations in existing methodologies to randomly incorporate insertions and/or deletions (see Supplementary Table S13). Consequently, their application in protein engineering has been sparse, with very few directed evolution campaigns on record that originate from such libraries. For example, the RID protocol (31), the first attempt towards creating InDel libraries, relies on a complex protocol involving random cleavage of single stranded DNA, so that random substitutions are introduced unintentionally alongside the target mutations. Two other early methods, segmental mutagenesis (11) and RAISE (9), do not control for the length of the InDel and consequently produce libraries that primarily contain frameshifted variants. In contrast, a codon-based protocol dubbed COBARDE (32) gives a pool of multiple codon-based deletions with <5% frameshifts but requires custom reprogramming of an oligonucleotide synthesizer to create mutagenic oligonucleotides. Alternatively, the viability of transposon-based protocols has been established for generating deletions of various sizes, up to gene truncation variants (13-15). However, the only reported such protocol to create insertions, namely pentapeptide scanning mutagenesis (16,33), merely gains access to insertions of defined size and sequence.

Improving on existing methodology, the TRIAD protocol meets all major requirements outlined above and gives easy access to large, diverse InDel libraries. The random insertion of a transposon gives excellent sampling of the entire target sequence (Figure 3; Supplementary Figure S8). Extensive sequencing shows that Mu transposon is even less biased than previously thought, so that functional effects upon insertion/deletion in *any* region of the protein can be taken advantage of. Access to library sizes upwards of 10^5 variants was possible, covering most of the theoretical diversity of up to two randomised amino acid insertions (Supplementary Figure S5). Finally, the procedure is technically straightforward, consisting of transposition and cloning steps, and does not require access to specialized DNA synthesis equipment (as in (32)). The TRIAD workflow is a versatile process that can be adapted to create libraries focused on a specific region of a protein (*e.g.,* in cases where screening throughput is limited). This approach would be analogous to other procedures (although only a few (32,34) have directly exemplified this case). In the case of TRIAD, this was typically achieved by adding an in-frame seamless cloning step using a type IIS restriction enzyme such as SapI (see Supplementary Methods and Results; Supplementary Figure S12; Supplementary Table S12). InDel libraries constructed in this way showed good coverage of the target region, albeit with slightly more pronounced bias than whole-gene TRIAD, presumably due to increased sensitivity to

preferential transposon insertions on a short target sequence. Alternatively, TRIAD can be further expanded with a recombination protocol (*e.g.*, DNA shuffling or Staggered Extension Process) to generate variants combining multiple InDels, which can be screened in a high-throughput assay (35).

How do InDels compare to substitutions as sources of functional innovation?

Point substitutions, small insertions and deletions account for most evolutionary changes among natural proteins (1). The ratio of InDels to point substitutions covers a wide variety of ratios across different species, ranging from 1:5 in humans and primates (36) to 1:20 in bacteria (37), which indicates that InDels are typically subject to stronger purifying selection. Additionally, protein sequence alignments have established that the majority of InDels fixed in protein-coding genes are short (*i.e.*, encompassing 1 to 5 residues) and occur almost exclusively in loops linking secondary structure elements at the solvent-exposed surfaces of proteins (38-43). While a large body of experimental evidence reports on the effects of substitutions, the impact of InDels on structural stability and functional divergence in protein evolution is still imperfectly understood, no doubt in part because convenient methods to introduce them in library experiments were missing. Substitutions, being merely side chain alterations, tend to have local effects with typically minor consequences for the overall structure of a protein. By contrast, InDels alter the length of the backbone, opening the way to dramatically larger changes in the packing and orientation of domains that may result in more global effect on the protein structure (44-46). Examples of InDels that cause significant repositioning of the backbone and nearby side chains to accommodate the extra or lost residues are on record (47-49). If such rearrangements occur near the active site of a protein, the resulting structural changes can change specificity and activity (4,50,51). Additionally, short InDels occurring at oligomerisation interfaces have also been shown to have important effects on the stability and/or specificity of protein complexes (52,53). A corollary of the comparatively drastic effect of InDels on protein structure is the perception that they are more deleterious. Indeed, this view is now experimentally corroborated by our work on *wt*PTE (Figure 5) as well as a recent deep mutational scanning study investigating the fitness effects of single amino acid InDels on TEM-1 β-lactamase (54). However, InDels have also been shown to be contribute to functional divergence in several enzyme families, such as lactate and malate dehydrogenases (55), tRNA nucleotidyltransferases (56), nitroreductases (57), o-succinylbenzoate synthases (53), and phosphotriesterase-like lactonases (2,4). Accordingly, our comparative analysis of the fitness effect of InDels and point substitutions indicated that InDels are more likely to yield *wt*PTE variants with improved arylesterase activity than substitutions (Figure 5B), thus emphasising that an InDel strategy is a viable alternative for protein engineering.

The potential of InDel mutagenesis strategies in directed protein evolution

Having developed an accessible route to a wide diversity of InDel variants, we set out to interrogate what effect they may have on the activity of the model enzyme phosphotriesterase (*wt*PTE). The promiscuous esterase activity of *wt*PTE has previously been used as the starting point of an evolutionary trajectory, which generated an arylesterase that hydrolysed 2-NH with high efficiency (27). The mutation H254R, selected after the first round of mutagenesis, appeared to be a mutation

on which the rest of the trajectory was highly contingent. InDel mutagenesis and selection puts us in a position to address the question whether alternative initial mutations would enable access to different evolutionary trajectories leading towards the same functional outcome. Based on the hypothesis that the use of a wider genetic and functional diversification (*i.e.*, by both substitutions and InDels) might lead to a wider diversity of possible evolutionary trajectories, the first objective was to identify new adaptive mutations improving the promiscuous arylesterase activity of *wt*PTE by screening InDel libraries of *wt*PTE generated *via* TRIAD. This resulted in the identification of multiple beneficial deletions and insertions, confirming that introduction of InDels can give rise to functional and improved catalysts.

We further observed that four of these adaptive InDels increase arylesterase activity 10- to 20-fold (in k_{cat}/K_M) against 2-NH, which is more than brought about by the initial H254R mutation from the previous directed evolution experiment (in which only a 2.6-fold difference was observed) (27). For all four InDel variants, the improvement in 2-NH catalytic efficiency appears to be due to increased k_{cat} (from 13- to 67-fold), which outweighs an increased K_M in all four (from 2 to 6-fold). Similarly, all four variants increased in K_M for paraoxon (from 2 to 45-fold). On the other hand, the substitution H254R showed a different profile: it decreased K_M for paraoxon 8-fold, while hardly increasing it for 2-NH (1.4-fold) (27). Therefore, the top InDel hits in the cell lysate screening are more disruptive for both the binding of paraoxon and arylester (2-NH) substrates than substitutions, as may be expected for mutations that alter the backbone structure, while remaining beneficial overall.

Despite the scarcity of facile random InDel mutagenesis methods until recently, several examples of an adaptive role of InDels in protein directed evolution have been observed. Work on TEM-1 β-lactamase using the original Mu transposon-based triplet deletion libraries identified variants with increased resistance towards the antibiotic ceftazidime, up to 64-fold in minimum inhibitory concentration (58). A similar campaign that selected for eGFP variants with increased brightness in a colony screen identified the surprising eGFP-ΔGly4 deletion, which has significantly more cellular fluorescence likely due to increased refolding efficiency (59). Finally, a recent focused library approach in a PTE-like lactonase with insertions into loop 7 (being shorter in lactonases) led to variants enhanced in phosphotriesterase activity, with increased k_{cat} and decreased K_M for paraoxon (k_{cat}/K_M increased up to 600-fold) (4). Native lactonase activity was strongly affected in those variants with up to 10^4-fold decreases in catalytic efficiency. These results in an enzyme closely related to PTE are very similar to our observations of the mixed effect of InDels on *wt*PTE, as explored based on the larger diversity of adaptive variants rendered available by TRIAD.

In conclusion, our results suggest that unprecedented evolutionary trajectories become accessible by screening InDel libraries obtained *via* TRIAD, establishing a new paradigm that complements current strategies following the 'one amino acid at the time' adage (60) which are believed to lead to successful outcomes slowly, yet steadily. The effect of InDels is *on average* more deleterious than substitutions (Figure 5A), while the *fraction* of hits is increased in InDel libraries (Figure 5B), suggesting that InDel library strategies tend to 'polarize' properties of library members towards extremes. For thermodynamically more difficult reactions than those studied here, this trend to more extreme outcomes may practically imply low hit rates, in which case high throughput screening would

become crucial. For example, ultrahigh-throughput screening based on droplet microfluidics (61,62) could be combined with InDel mutagenesis to powerfully explore sequence space for evolutionary trajectories and individual variants that would not arise from epPCR mutagenesis libraries. It remains to be seen whether this new way of 'jumping' (rather than 'tiptoeing') across sequence space yields functionally better catalysts - or just different ones.

AVAILABILITY

Illumina raw sequencing reads were deposited with European Nucleotide Archive (https://www.ebi.ac.uk/ena) at accession number PRJEB28011. The source code along with instructions for all scripts involved in data processing are freely available at https://github.com/fhlab/TRIAD.

SUPPLEMENTARY DATA

Results describing focused InDel libraries generated by TRIAD, supplementary methods, supplementary figures S1-S13 and supplementary tables S1-14 are available as Supplementary Information.

ACKNOWLEDGEMENTS

This work was performed using resources provided by the Cambridge Service for Data Driven Discovery (CSD3) operated by the University of Cambridge Research Computing Service (http://www.csd3.cam.ac.uk/), provided by Dell EMC and Intel using Tier-2 funding from the Engineering and Physical Sciences Research Council (capital grant EP/P020259/1), and DiRAC funding from the Science and Technology Facilities Council (www.dirac.ac.uk). We would like to thank Christopher Kitching for assistance with analysis scripts and Charlotte Miton for helpful comments on the manuscript.

FUNDING

This work was supported by the Biotechnology and Biological Sciences Research Council through a research grant [BB/L002469/1] and a studentship to MP [BB/M011194/1]. SE, NT and SRAD held EU Marie Curie FP7 fellowships, FH is an H2020 ERC Advanced Investigator [695669].

REFERENCES

1. Chothia, C., Gough, J., Vogel, C. and Teichmann, S.A. (2003) Evolution of the protein repertoire. *Science*, **300**, 1701-1703.
2. Afriat-Jurnou, L., Jackson, C.J. and Tawfik, D.S. (2012) Reconstructing a missing link in the evolution of a recently diverged phosphotriesterase by active-site loop remodeling. *Biochemistry*, **51**, 6047-6055.
3. Herman, A. and Tawfik, D.S. (2007) Incorporating Synthetic Oligonucleotides via Gene Reassembly (ISOR): a versatile tool for generating targeted libraries. *Protein Eng Des Sel*, **20**, 219-226.
4. Hoque, M.A., Zhang, Y., Chen, L., Yang, G., Khatun, M.A., Chen, H., Hao, L. and Feng, Y. (2017) Stepwise Loop Insertion Strategy for Active Site Remodeling to Generate Novel Enzyme Functions. *ACS Chem Biol*, **12**, 1188-1193.
5. Park, H.S., Nam, S.H., Lee, J.K., Yoon, C.N., Mannervik, B., Benkovic, S.J. and Kim, H.S. (2006) Design and evolution of new catalytic activity with an existing protein scaffold. *Science*, **311**, 535-538.
6. Lamminmaki, U., Pauperio, S., Westerlund-Karlsson, A., Karvinen, J., Virtanen, P.L., Lovgren, T. and Saviranta, P. (1999) Expanding the conformational diversity by random insertions to CDRH2 results in improved anti-estradiol antibodies. *J Mol Biol*, **291**, 589-602.
7. Emond, S., Mondon, P., Pizzut-Serin, S., Douchy, L., Crozet, F., Bouayadi, K., Kharrat, H., Potocki-Veronese, G., Monsan, P. and Remaud-Simeon, M. (2008) A novel random mutagenesis approach using human mutagenic DNA polymerases to generate enzyme variant libraries. *Protein Eng Des Sel*, **21**, 267-274.
8. Kashiwagi, K., Isogai, Y., Nishiguchi, K. and Shiba, K. (2006) Frame shuffling: a novel method for in vitro protein evolution. *Protein Eng Des Sel*, **19**, 135-140.
9. Fujii, R., Kitaoka, M. and Hayashi, K. (2014) Random insertional-deletional strand exchange mutagenesis (RAISE): a simple method for generating random insertion and deletion mutations. *Methods Mol Biol*, **1179**, 151-158.
10. Hida, K., Won, S.Y., Di Pasquale, G., Hanes, J., Chiorini, J.A. and Ostermeier, M. (2010) Sites in the AAV5 capsid tolerant to deletions and tandem duplications. *Arch Biochem Biophys*, **496**, 1-8.
11. Pikkemaat, M.G. and Janssen, D.B. (2002) Generating segmental mutations in haloalkane dehalogenase: a novel part in the directed evolution toolbox. *Nucleic Acids Res*, **30**, E35-35.
12. Kipnis, Y., Dellus-Gur, E. and Tawfik, D.S. (2012) TRINS: a method for gene modification by randomized tandem repeat insertions. *Protein Eng Des Sel*, **25**, 437-444.
13. Morelli, A., Cabezas, Y., Mills, L.J. and Seelig, B. (2017) Extensive libraries of gene truncation variants generated by in vitro transposition. *Nucleic Acids Res*, **45**, e78.
14. Jones, D.D. (2005) Triplet nucleotide removal at random positions in a target gene: the tolerance of TEM-1 beta-lactamase to an amino acid deletion. *Nucleic Acids Res*, **33**, e80.
15. Liu, S.S., Wei, X., Ji, Q., Xin, X., Jiang, B. and Liu, J. (2016) A facile and efficient transposon mutagenesis method for generation of multi-codon deletions in protein sequences. *J Biotechnol*, **227**, 27-34.
16. Hallet, B., Sherratt, D.J. and Hayes, F. (1997) Pentapeptide scanning mutagenesis: random insertion of a variable five amino acid cassette in a target protein. *Nucleic Acids Res*, **25**, 1866-1867.
17. Caldwell, S.R., Newcomb, J.R., Schlecht, K.A. and Raushel, F.M. (1991) Limits of diffusion in the hydrolysis of substrates by the phosphotriesterase from Pseudomonas diminuta. *Biochemistry*, **30**, 7438-7444.
18. Roodveldt, C. and Tawfik, D.S. (2005) Shared promiscuous activities and evolutionary features in various members of the amidohydrolase superfamily. *Biochemistry*, **44**, 12728-12736.
19. Tokuriki, N. and Tawfik, D.S. (2009) Chaperonin overexpression promotes genetic variation and enzyme evolution. *Nature*, **459**, 668-673.
20. Baldwin, A.J., Busse, K., Simm, A.M. and Jones, D.D. (2008) Expanded molecular diversity generation during directed evolution by trinucleotide exchange (TriNEx). *Nucleic Acids Res*, **36**, e77.
21. Kaltenbach, M., Emond, S., Hollfelder, F. and Tokuriki, N. (2016) Functional Trade-Offs in Promiscuous Enzymes Cannot Be Explained by Intrinsic Mutational Robustness of the Native Activity. *PLoS Genet*, **12**, e1006305.

22. Zhang, J., Kobert, K., Flouri, T. and Stamatakis, A. (2014) PEAR: a fast and accurate Illumina Paired-End reAd mergeR. *Bioinformatics*, **30**, 614-620.
23. Langmead, B. and Salzberg, S.L. (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods*, **9**, 357-359.
24. Rice, P., Longden, I. and Bleasby, A. (2000) EMBOSS: the European Molecular Biology Open Software Suite. *Trends Genet*, **16**, 276-277.
25. Haapa, S., Taira, S., Heikkinen, E. and Savilahti, H. (1999) An efficient and accurate integration of mini-Mu transposons in vitro: a general methodology for functional genetic analysis and molecular biology applications. *Nucleic Acids Res*, **27**, 2777-2784.
26. Kaltenbach, M., Jackson, C.J., Campbell, E.C., Hollfelder, F. and Tokuriki, N. (2015) Reverse evolution leads to genotypic incompatibility despite functional and active site convergence. *Elife*, **4**.
27. Tokuriki, N., Jackson, C.J., Afriat-Jurnou, L., Wyganowski, K.T., Tang, R. and Tawfik, D.S. (2012) Diminishing returns and tradeoffs constrain the laboratory optimization of an enzyme. *Nat Commun*, **3**, 1257.
28. Baldwin, A.J., Arpino, J.A., Edwards, W.R., Tippmann, E.M. and Jones, D.D. (2009) Expanded chemical diversity sampling through whole protein evolution. *Mol Biosyst*, **5**, 764-766.
29. Haapa-Paananen, S., Rita, H. and Savilahti, H. (2002) DNA transposition of bacteriophage Mu. A quantitative analysis of target site selection in vitro. *J Biol Chem*, **277**, 2843-2851.
30. Mizuuchi, M. and Mizuuchi, K. (1993) Target site selection in transposition of phage Mu. *Cold Spring Harb Symp Quant Biol*, **58**, 515-523.
31. Murakami, H., Hohsaka, T. and Sisido, M. (2002) Random insertion and deletion of arbitrary number of bases for codon-based random mutation of DNAs. *Nat Biotechnol*, **20**, 76-81.
32. Osuna, J., Yanez, J., Soberon, X. and Gaytan, P. (2004) Protein evolution by codon-based random deletions. *Nucleic Acids Res*, **32**, e136.
33. Hayes, F. and Hallet, B. (2000) Pentapeptide scanning mutagenesis: encouraging old proteins to execute unusual tricks. *Trends Microbiol*, **8**, 571-577.
34. Tizei, P.A.G., Harris, E., Renders, M. and Pinheiro, V.B. (2017) InDel assembly: A novel framework for engineering protein loops through length and compositional variation. *bioRxiv*, 10.1101/127829, 14-July-2017, pre-print: not peer-reviewed.
35. Skamaki, K., Emond, S., Chodorge, M., Andrews, J., Rees, G., Cannon, D., Popovic, B., Buchanan, A., Minter, R. and Hollfelder, F. (2019) In vitro Evolution of Antibody Affinity via Insertional Mutagenesis Scanning of an Entire Antibody Variable Region. *bioRxiv*, DOI TBA, E-PUB Date TBA, pre-print: not peer-reviewed.
36. Ng, P.C., Levy, S., Huang, J., Stockwell, T.B., Walenz, B.P., Li, K., Axelrod, N., Busam, D.A., Strausberg, R.L. and Venter, J.C. (2008) Genetic variation in an individual human exome. *PLoS Genet*, **4**, e1000160.
37. Chen, J.Q., Wu, Y., Yang, H., Bergelson, J., Kreitman, M. and Tian, D. (2009) Variation in the ratio of nucleotide substitution and indel rates across genomes in mammals and bacteria. *Mol Biol Evol*, **26**, 1523-1531.
38. Ajawatanawong, P. and Baldauf, S.L. (2013) Evolution of protein indels in plants, animals and fungi. *BMC Evol Biol*, **13**, 140.
39. Benner, S.A., Cohen, M.A. and Gonnet, G.H. (1993) Empirical and structural models for insertions and deletions in the divergent evolution of proteins. *J Mol Biol*, **229**, 1065-1082.
40. Lin, M., Whitmire, S., Chen, J., Farrel, A., Shi, X. and Guo, J.T. (2017) Effects of short indels on protein structure and function in human genomes. *Sci Rep*, **7**, 9313.
41. Pascarella, S. and Argos, P. (1992) Analysis of insertions/deletions in protein structures. *J Mol Biol*, **224**, 461-471.
42. Taylor, M.S., Ponting, C.P. and Copley, R.R. (2004) Occurrence and consequences of coding sequence insertions and deletions in Mammalian genomes. *Genome Res*, **14**, 555-566.
43. Toth-Petroczy, A. and Tawfik, D.S. (2013) Protein insertions and deletions enabled by neutral roaming in sequence space. *Mol Biol Evol*, **30**, 761-771.
44. Grishin, N.V. (2001) Fold change in evolution of protein structures. *J Struct Biol*, **134**, 167-185.
45. Studer, R.A., Dessailly, B.H. and Orengo, C.A. (2013) Residue mutations and their impact on protein structure and function: detecting beneficial and pathogenic changes. *Biochem J*, **449**, 581-594.
46. Toth-Petroczy, A. and Tawfik, D.S. (2014) Hopeful (protein InDel) monsters? *Structure*, **22**, 803-804.

47. Heinz, D.W., Baase, W.A., Dahlquist, F.W. and Matthews, B.W. (1993) How amino-acid insertions are allowed in an alpha-helix of T4 lysozyme. *Nature*, **361**, 561-564.
48. O'Neil, K.T., Bach, A.C., 2nd and DeGrado, W.F. (2000) Structural consequences of an amino acid deletion in the B1 domain of protein G. *Proteins*, **41**, 323-333.
49. Stott, K.M., Yusof, A.M., Perham, R.N. and Jones, D.D. (2009) A surface loop directs conformational switching of a lipoyl domain between a folded and a novel misfolded structure. *Structure*, **17**, 1117-1127.
50. Shortle, D. and Sondek, J. (1995) The emerging role of insertions and deletions in protein engineering. *Curr Opin Biotechnol*, **6**, 387-393.
51. Tawfik, D.S. (2006) Biochemistry. Loop grafting and the origins of enzyme species. *Science*, **311**, 475-476.
52. Hashimoto, K. and Panchenko, A.R. (2010) Mechanisms of protein oligomerization, the critical role of insertions and deletions in maintaining different oligomeric states. *Proc Natl Acad Sci U S A*, **107**, 20352-20357.
53. Odokonyero, D., Sakai, A., Patskovsky, Y., Malashkevich, V.N., Fedorov, A.A., Bonanno, J.B., Fedorov, E.V., Toro, R., Agarwal, R., Wang, C. *et al.* (2014) Loss of quaternary structure is associated with rapid sequence divergence in the OSBS family. *Proc Natl Acad Sci U S A*, **111**, 8535-8540.
54. Gonzalez, C.E., Roberts, P. and Ostermeier, M. (2019) Fitness Effects of Single Amino Acid Insertions and Deletions in TEM-1 beta-Lactamase. *J Mol Biol*, **431**, 2320-2330.
55. Boucher, J.I., Jacobowitz, J.R., Beckett, B.C., Classen, S. and Theobald, D.L. (2014) An atomic-resolution view of neofunctionalization in the evolution of apicomplexan lactate dehydrogenases. *Elife*, **3**.
56. Neuenfeldt, A., Just, A., Betat, H. and Morl, M. (2008) Evolution of tRNA nucleotidyltransferases: a small deletion generated CC-adding enzymes. *Proc Natl Acad Sci U S A*, **105**, 7953-7958.
57. Akiva, E., Copp, J.N., Tokuriki, N. and Babbitt, P.C. (2017) Evolutionary and molecular foundations of multiple contemporary functions of the nitroreductase superfamily. *Proc Natl Acad Sci U S A*, **114**, E9549-E9558.
58. Simm, A.M., Baldwin, A.J., Busse, K. and Jones, D.D. (2007) Investigating protein structural plasticity by surveying the consequence of an amino acid deletion from TEM-1 beta-lactamase. *FEBS Lett*, **581**, 3904-3908.
59. Arpino, J.A., Reddington, S.C., Halliwell, L.M., Rizkallah, P.J. and Jones, D.D. (2014) Random single amino acid deletion sampling unveils structural tolerance and the benefits of helical registry shift on GFP folding and structure. *Structure*, **22**, 889-898.
60. Tracewell, C.A. and Arnold, F.H. (2009) Directed enzyme evolution: climbing fitness peaks one amino acid at a time. *Curr Opin Chem Biol*, **13**, 3-9.
61. Colin, P.Y., Zinchenko, A. and Hollfelder, F. (2015) Enzyme engineering in biomimetic compartments. *Curr Opin Struct Biol*, **33**, 42-51.
62. Mair, P., Gielen, F. and Hollfelder, F. (2017) Exploring sequence space in search of functional enzymes using microfluidic droplets. *Curr Opin Chem Biol*, **37**, 137-144.

TABLES AND FIGURES CAPTIONS

Table 1. Mutagenesis efficiency of TRIAD analysed by deep sequencing

Unique in-frame InDels (*i.e.*, InDels of multiple of three nucleotides) were counted both at the DNA and the protein level. Adjacent amino acid substitutions and truncations (resulting from the occurrence of stop codons) may occur depending on the insertion point of the transposon, resulting in a lower value for the number of observed unique protein InDels. The proportion relative to the theoretical diversity accessible from the *wt*PTE sequence (both at the DNA and the protein level) was calculated as the ratio between the number of unique in-frame InDels observed by deep sequencing and the theoretical diversity for a given TRIAD library (see Supplementary Figure S5).

Table 2. Analysis of InDel *wt*PTE variants with at least 2-fold improved arylesterase activity.

Values refer to the activity change of all or AE positive variants relative to wtPTE obtained by comparing the initial rates v_0 for the hydrolysis of paraoxon, 4-NPB or 2-NH to that of *wt*PTE at 200 µM substrate concentration, resulting in a dimensionless ratio (recorded in Table 2). The average effect value was determined as the geometric mean of the relative activities of all the variants listed in Supplementary Table S11. The maximum, median and minimum changes correspond to the maximum, median and minimum relative activities for each substrate among the variants (See also Supplementary Figure S12).

Table 3. Kinetic properties of *wt*PTE variants.

Figure 1. Schematic outline of TRIAD.

(A) Generation of deletion libraries. *Step 1*: The TransDel insertion library is generated by *in vitro* transposition of the engineered transposon TransDel into the target sequence. *Step 2*: *Mly*I digestion removes TransDel together with 3 bp of the original target sequence and generate a single break per variant. *Step 3a*: self-ligation results in the reformation of the target sequence minus 3 bp, yielding a library of single variants with a deletion of one triplet (14). *Step 3b*: DNA cassettes dubbed Del2 and Del3 are then inserted between the break in the target sequence to generate Del2 and Del3 insertion libraries. *Step 4b*: *Mly*I digestion removes Del2 and Del3 together with 3 and 6 additional bp of the original target sequence, respectively. *Step 5b*: self-ligation results in the reformation of the target sequence minus 6 and 9 bp, yielding libraries of single variants with a deletion of 2 and 3 triplets, respectively. Deletions are indicated by red vertical lines.

(B) Generation of insertion libraries. Step 1: The TransIns insertion library is generated by *in vitro* transposition of the engineered transposon TransIns into the target sequence. Step 2: digestion by *Not*I and *Mly*I removes TransIns. Step 3: DNA cassettes dubbed Ins1, Ins 3 and Ins3 (with respectively 1, 2 and 3 randomized bp triplets at one of their extremities; indicated by purple triangles) are then inserted between the break in the target sequence to generate the corresponding Ins1, Ins2 and Ins3 insertion libraries. Step 4: *Acu*I digestion and 3'-end digestion by the Klenow fragment

remove the cassettes, leaving the randomized triplet(s) in the original target sequence. Step 5: Self-ligation results in the reformation of the target sequence plus 3, 6 and 9 random bp, yielding libraries of single variants with an insertion of 1, 2 and 3 triplets, respectively.

Figure 2. Mechanism for the generation of InDels using TRIAD.

(A) Generation of single, double and triple triplet nucleotide deletions. Step 1. Two MlyI recognition sites (5'GAGTC(N)$_5\downarrow$) are positioned at each end of TransDel, 1 bp away from the site of transposon insertion. Transposition with TransDel results in the duplication of 5 bp ($N_4N_5N_6N_7N_8$) of the target DNA at the insertion point. TransDel carries a selection marker (resistance gene against chloramphenicol; CamR) enabling the recovery of *in vitro* transposition products after transformation into *E. coli*. Step 2. MlyI digestion removes TransDel together with 8 bp of the target DNA (4 bp at each end), leaving blunt ends and resulting in the removal of a contiguous 3 bp sequence from the target DNA ($N_5N_6N_7$). Step 3a. Self-ligation reforms the target DNA minus 3 bp, as previously described (14). Step 3b. Alternatively, blunt-ended cassettes Del2 or Del3 are ligated into the gap left upon TransDel removal for the generation of 6 and 9 bp deletions, respectively. Both Del2 and Del3 also contain two MlyI recognition sites advantageously positioned towards the ends of the cassettes. These cassettes also contain a different marker than TransDel (resistance gene against kanamycin; KanR) to avoid cross-contamination. Step 4b. MlyI digestion removes Del2 and Del3 together with respectively 3 and 6 additional bp of the original target DNA. In the case of Del2, MlyI digestion results in the removal of a 3 bp sequence ($N_2N_3N_4$) on one side of the cassette. In the case of Del3, MlyI digestion results in the removal of two 3 bp sequence ($N_2N_3N_4$) on both side of the cassette ($N_2N_3N_4$ and $N_8N_9N_{10}$). Step 5b. Self-ligation reforms the target DNA minus 6 or 9 bp.

(B) Generation of single, double and triple randomized triplet nucleotide insertions. Step 1. TransDel is an asymmetric transposon with MlyI at one end and NotI at the other end. Both recognition sites are positioned 1bp away from TransIns insertion site. Upon transposition, 5 bp ($N_1N_2N_3N_4N_5$) of the target DNA are duplicated at the insertion point of TransIns. Step 2. Double digestion with NotI and MlyI results in the removal of TransIns. Digestion with MlyI removes TransIns with 4 bp ($N_1N_2N_3N_4$) of the duplicated sequence at the transposon insertion site. Digestion with NotI leaves a 5', 4-base cohesive overhang. Step 3. DNA cassettes Ins1, Ins2 and Ins3 (Ins1/2/3) carrying complementary ends are ligated in the NotI/MlyI digested TransIns insertion site. Ins1, Ins2 and Ins3 carry respectively 1, 2 and 3 randomized bp triplets at their blunt-ended extremities ([NNN]$_{1,2 \text{ or } 3}$; indicated in purple). Ins1/2/3 contain two AcuI recognition sites (5'CTGAAG(16/14)) strategically positioned towards their ends. One site is located so that AcuI will cleave at the point where the target DNA joins Ins1/2/3. The other site is positioned so that AcuI will cut inside Ins1/2/3 to leave the randomized triplet(s) with the target DNA. Step 4. Digestion with AcuI removes Ins1/2/3 leaving 3', 2-base overhangs with the target DNA (*i.e.*, 5'N$_5$T on one end and 5'TC on the end carrying the randomized triplet(s)). Digestion with the Large Klenow fragment generates blunt ends by removing the overhangs. This step also enables to discard the extra nucleotide (N_5) from the sequence

duplicated during the transposition. Step 5. Self-ligation reforms the target DNA with one, two or three randomized nucleotide triplets.

Figure 3. Mutagenesis efficiency of TRIAD. The composition of InDel libraries in the *wt*PTE gene was determined by deep sequencing and validated using Sanger sequences from randomly chosen variants. **(A)** Relative frequency of TransDel transposon insertion across *wt*PTE, derived from -3 bp deletions observed in deep sequencing and normalized for sequencing depth and InDel redundancy in DNA sequence (see Supplementary Methods 2.7). Insert: Relative site preference converted into the TransDel consensus insertion site. **(B)** Distribution and number of detected distinct DNA deletions in -3, -6 and -9 bp libraries combined per *wt*PTE position. **(C)** Distribution and number of observed +3 bp mutations per DNA position in +3 bp library, compared to the median 20.85 variants per position (horizontal line). Due to varying InDel redundancy depending on sequence context, the theoretical DNA diversity per position is between 42 and 48 variants (see Supplementary Figure S5). Analogous plots for +6 and +9 bp libraries are shown in Supplementary Figure S8.

Figure 4. Structures of substrates. *wt*PTE catalyzes the hydrolysis of paraoxon (native substrate) and possess promiscuous activity against arylester substrates, *e.g.* 4-nitrophenyl butyrate and 2-naphthyl hexanoate.

Figure 5. Fitness effects of InDels *versus* substitutions on *wt*PTE phosphotriesterase. (A) Distribution of fitness effects on phosphotriesterase activity (paraoxon). **(B)** Distribution of fitness effects on arylesterase activity (4-NPB). Fitness effects are classified as strongly deleterious (>10-fold activity decrease relative to *wt*PTE), mildly deleterious (10-fold to 1.5-fold decrease), neutral (<1.5-fold change), and beneficial (>1.5-fold increase). **(C)** Average fitness change in phosphotriesterase activity by deletions, insertions and substitutions. The average fitness change refers to the change in initial rates as a consequence of mutation and is calculated as the geometric mean of the relative activities of the variants (see Supplementary Table S8). Error bars indicate the corresponding confidence intervals (5% risk of error). **(D)** Structural mapping of protein changes observed in variants retaining ≥ 50% of *wt*PTE activity level (PDB ID 4PCP).

Figure 6. Identification of InDels improving the promiscuous arylesterase activity of *wt*PTE. (A)(B) Changes in phosphotriesterase (native; PTE) and arylesterase (promiscuous; AE) activities among *wt*PTE. InDel variants identified upon screening against butyrate (4-NPB; panel A) and hexanoate (2-NH; panel B) esters, respectively. The enzymatic activities for each variant (shown as grey dots) were measured in cell lysates and are plotted relative to those of *wt*PTE. Data are averages of triplicate values from three independent experiments and error bars represent +/- 1 SEM. The dashed diagonal lines (in blue) demarcate the different trade-off regimes. Variants below the diagonal show a strong negative trade-off, with a large detriment to the original PTE activity (specificity ratio > 100), as AE activity is improved. Above the diagonal variants with weak trade-off

emerge as generalists (specificity ratio < 100). **(C)** Position of the adaptive InDels in the PTE structure highlighting the frequency of mutations in loops 8, 7 and 2.

Figure 1

Figure 2

Figure 3

Paraoxon

4-NPB (4-nitrophenyl butyrate)

2-NH (2-naphthyl hexanoate)

Figure 4

Figure 5

A

Insertions
Deletions

*wt*PTE

Relative PTE activity

Relative AE activity
(4-NPB)

B

*wt*PTE

Relative PTE activity

Relative AE activity
(2-NH)

C

Loop 8

310
303
308

311
313-314
316-318

260-263

Loop 7

256-258

270-273

Loop 2

Figure 6