Loss of AKAP1 triggers Drp1 dephosphorylation-mediated mitochondrial fission and loss

in retinal ganglion cells

Genea Edwards,[1]* Guy A. Perkins,[2]* Keun-Young Kim,[2]* YeEun Kong,[1] Yonghoon Lee,[1] Soo-

Ho Choi,[3] Yujia Liu,[4] Dorota Skowronska-Krawczyk,[1] Robert N. Weinreb,[1] Linda Zangwill,[1]

Stefan Strack[4] and Won-Kyu Ju[1]

[1]Hamilton Glaucoma Center and Viterbi Family Department of Ophthalmology, Shiley Eye

Institute, University of California San Diego, La Jolla, CA, USA

[2]National Center for Microscopy and Imaging Research and Department of Neuroscience,

University of California San Diego, La Jolla, CA, USA

[3]Department of Medicine, University of California San Diego, La Jolla, CA, USA

[4]Department of Pharmacology and Iowa Neuroscience Institute, University of Iowa, Iowa City,

IA, USA

Running title: Loss of AKAP1 in retinal ganglion cells

*Genea Edwards, Guy A. Perkins and Keun-Young Kim contributed equally to this work.

Corresponding author: Won-Kyu Ju, Ph.D.

Hamilton Glaucoma Center

Viterbi Family Department of Ophthalmology

Shiley Eye Institute

University of California, San Diego

9415 Campus Point Drive

La Jolla, CA 92037

Email: wju@ucsd.edu

Abstract

Impairment of mitochondrial structure and function is strongly linked to glaucoma pathogenesis. Despite the widely appreciated disease relevance of mitochondrial dysfunction and loss, the molecular mechanisms underlying mitochondrial fragmentation and metabolic stress in glaucoma are poorly understood. We demonstrate here that glaucomatous retinal ganglion cells (RGCs) show loss of A-kinase anchoring protein 1 (AKAP1), activation of calcineurin (CaN) and reduction of dynamin-related protein 1 (Drp1) phosphorylation at serine 637 (Ser637). These findings suggest that AKAP1-mediated phosphorylation of Drp1 at Ser637 has a critical role in RGC survival in glaucomatous neurodegeneration. Male mice lacking AKAP1 show increases of CaN and total Drp1 level, as well as a decrease of Drp1 phosphorylation at Ser637 in the retina. Ultrastructural analysis of mitochondria shows that loss of AKAP1 triggers mitochondrial fragmentation and loss, as well as mitophagosome formation in RGCs. Loss of AKAP1 deregulates oxidative phosphorylation (OXPHOS) complexes (Cxs) by increasing CxII and decreasing CxIII-V, leading to metabolic and oxidative stress. Also, loss of AKAP1 decreases Akt phosphorylation at Serine 473 (Ser473) and threonine 308 (Thr308) and activates the Bim/Bax signaling pathway in the retina. These results suggest that loss of AKAP1 has a critical role in RGC dysfunction by decreasing Drp1 phosphorylation at Ser637, deregulating OXPHOS, decreasing Akt phosphorylation at Ser473 and Thr308, and activating the Bim/Bax pathway in glaucomatous neurodegeneration. Thus, we propose that overexpression of AKAP1 or modulation of Drp1 phosphorylation at Ser637 are potential therapeutic strategies for neuroprotective intervention in glaucoma and other mitochondria-related optic neuropathies.

Introduction

Primary open-angle glaucoma (POAG) is characterized by a slow and progressive degeneration

of retinal ganglion cells (RGCs) and their axons, leading to loss of visual function[1]. The factors

contributing to degeneration of the RGC and its axon degeneration in POAG are not well

understood. Recent studies have shown that POAG patients have mitochondrial abnormalities[2-7].

Evidence from our group and others indicates that compromised mitochondrial dynamics,

metabolic stress and mitochondrial dysfunction by glaucomatous insults such as elevated

intraocular pressure (IOP) and oxidative stress are critical to RGC loss in experimental

glaucoma[8-14]. Despite the widely appreciated disease relevance of mitochondrial dysfunction

and loss, the molecular mechanisms underlying the impairment of mitochondrial structure and

function in glaucoma are poorly understood.

A-kinase anchoring protein 1 (AKAP1), a scaffold protein, is part of the elaborate

architecture which binds to the regulatory subunit of protein kinase A (PKA) along with other

signaling molecules such as Src kinase, protein phosphatase 1 (PP1), Ca^{2+}-dependent

phosphatase calcineurin (CaN) and phosphodiesterase 4 (PDE4)[15-20]. By recruiting the PKA

holoenzyme to the outer mitochondrial membrane (OMM), AKAP1 is strategically positioned to

integrate multiple signal transduction cascades, including cyclic adenosine 3´5´-monophosphate

(cAMP) signaling, in the regulation of mitochondrial shape and function. AKAP1 facilitates

dynamin-related protein 1 (Drp1) phosphorylation and subsequent mitochondrial elongation. It

also increases adenosine triphosphate (ATP) generation and mitochondrial membrane

potential[21,22], and these collective changes are neuroprotective [23,24]. Emerging evidence

indicates that AKAP1 protects from cerebral ischemic stroke by maintaining respiratory chain

activity, inhibiting Drp1 dependent mitochondrial fission and superoxide production, and

delaying Ca^{2+} deregulation [24].

Excessive mitochondrial fission-mediated dysfunction has been implicated in various

neurodegenerative diseases including glaucoma[8,11,25,26]. Complexes of Drp1 assemble from the

cytosol onto mitochondria at focal sites of mitochondrial fission[27,28]. Recent evidence indicates

that post-translational modifications of Drp1 are linked to mitochondrial dysfunction-mediated

bioenergetic failure, synaptic injury and neuronal cell death[25,29-32]. Phosphorylation of Drp1 at

serine 637 (Ser637) by the cAMP-dependent PKA promotes mitochondrial fusion, whereas

dephosphorylation by the CaN promotes mitochondrial fission by decreasing Drp1 activity[33-35].

Previous studies showed that inhibition of Drp1 activity prevents mitochondrial fission and

protects RGCs and their axons in experimental glaucoma[11,36,37]. However, it remains unknown

whether AKAP1-mediated Drp1 phosphorylation at Ser637 play a critical role in glaucomatous

neurodegeneration. Along this line, the existence of AKAP1 in RGCs and its role in glaucoma

are completely unknown.

To address these questions, we investigated AKAP1 protein expression and Drp1

phosphorylation at Ser637 in retinas from glaucomatous DBA/2J mice, and also evaluated the

effect of AKAP1 loss in the retina using AKAP1 knockout (AKAP1$^{-/-}$) mice.

Results

AKAP1 deficiency in glaucomatous RGCs.

We examined AKAP1 protein expression in the retina of a mouse model of glaucoma DBA/2J mice, which spontaneously develop elevated IOP and glaucomatous damage with age[8,11,38]. Control DBA/2J-*Gpnmb*[+]/*SjJ* (D2-*Gpnmb*[+]) was used in which the gene for Gpnmb that has been altered to wild type sequence by targeted gene mutation. It is otherwise identical to the DBA/2J mouse strain, but maintains normal IOP during aging and does not develop optic nerve axon loss[39,40]. Our results demonstrated for the first time that elevated IOP significantly decreased AKAP1 protein expression in the retinas of 10-month-old glaucomatous DBA/2J mice compared with the retina of age-matched control D2-*Gpnmb*[+] mice (Fig. 1a). We observed that AKAP1 immunoreactivity was highly present in the outer plexiform layer (OPL) and ganglion cell layer (GCL) in D2-*Gpnmb*[+] retina (Fig. 1b). More specifically, AKAP1 immunoreactivity was colocalized with neuronal class III β-tubulin (TUJ1)-positive RGCs in the GCL of D2-*Gpnmb*[+] retina. Of interest, however, AKAP1 immunoreactivity was decreased in the OPL and TUJ1-positive RGCs in the GCL of glaucomatous DBA/2J retina (Fig. 1b and c).

Activation of CaN and dephosphorylation of Drp1 at Ser637 in glaucomatous retina

AKAP1 binds with two Serine/Threonine phosphatases, PP1 and CaN[41,42]. Loss of AKAP1 causes Drp1-mediated mitochondrial fission and decreases Drp1phsophorylation at Ser637 in neuronal cells of the brain[21,23,24,43,44]. More importantly, AKAP1 protects brain neuronal cells against cerebral ischemic stroke by inhibiting Drp1-dependent mitochondrial fission[24]. Since elevated IOP increased CaN and total Drp1 protein expression[11,45], as well as Drp1 inhibition rescued RGCs and their axons by preserving mitochondrial integrity in the retina and/or glial lamina of glaucomatous DBA/2J mice[11], we examined the expression levels of CaN and total Drp1, as well as phosphorylation of Drp1 at Ser637 in the retina of 10-month-old glaucomatous DBA/2J mice. We observed a significant increase of CaN protein expression in glaucomatous DBA/2J retina (Fig. 2a). Consistently, our results showed an increase of CaN immunoreactivity in RNA-binding protein with multiple splicing (RBPMS)-positive RGCs as well as neurons in the inner nuclear layer (INL) of glaucomatous DBA/2J retina (Fig. 2b and c). We also observed a significant increase of total Drp1 protein expression but a significant dephosphorylation of Drp1 Ser637 in glaucomatous DBA/2J retina (Fig. 2d). Consistently, our results showed an

increase of Drp1 immunoreactivity in TUJ1-positive RGCs of glaucomatous DBA/2J retina (Fig.

2e and f). These results suggest that elevated IOP-induced CaN activation is associated with

dephosphorylation of Drp1 at Ser637 in glaucomatous RGCs, leading to mitochondrial fisison[11].

Loss of AKAP1 triggers activation of CaN and dephosphorylation of Drp1 at Ser637 in the

retina

We next used the retinas from mice lacking 10-month-old AKAP1 (AKAP1[-/-])[24] and age-

matched wild-type (WT) mice and examined the expression level of CaN. Following the

confirmation of a significant decrease of *AKAP1* gene expression (Fig. 3a), we observed a

significant increase of CaN protein expression in AKAP1[-/-] retina compared with WT retina (Fig.

3b). In addition, increased CaN immunoreactivity was present in RBPMS-positive RGCs as well

as neurons in the INL in AKAP1[-/-] retina (Fig. 3c and d). We further examined expression levels

of AMP-activated protein kinase (AMPK) and Drp1 protein, as well as phosphorylation of

AMPK at threonine 172 (Thr172) and Drp1 at Ser637 in AKAP1[-/-] retina. There was no

significant difference in the phosphorylation of AMPK at Thr172 between WT and AKAP1[-/-]

retinas, whereas we observed a significant decrease of total AMPK protein expression in

AKAP1[-/-] retina (Fig. 4a). Interestingly, we observed a significant increase of total Drp1 protein

expression in in AKAP1[-/-] retina (Fig. 4b). Also, AKAP1[-/-] retina showed a significant

dephosphorylation of Drp1at Ser637, whereas there was no significant difference in the

phosphorylation of Drp1 at Ser616 between WT and AKAP1[-/-] retinas (Fig. 4b). Consistent with

this finding, increased Drp1 immunoreactivity was present in RBPMS-positive RGCs of

AKAP1[-/-] retina (Fig. 4c and d).

AKAP1 loss causes mitochondrial fission and mitophagosome formation in RGCs

We further examined the expression levels of optic atrophy type 1 (OPA1) and mitofusin (Mfn)

1 and 2 in AKAP1[-/-] retina and the alteration of mitochondrial ultrastructure in AKAP1[-/-] RGC

somas. We observed no significant difference in *OPA1* gene expression, but a significant

decrease of total OPA1 protein expression in AKAP1[-/-] retina compared with WT (Fig. 5a).

More specifically, small isoform (80 kDa) of OPA1 protein expression was significantly

decreased, whereas there was no significant difference in the large isoform (100 kDa) of OPA1

protein expression in AKAP1[-/-] retina (Fig. 5a). We also observed a significant decrease of *Mfn1*

gene expression but a significant increase of Mfn1 protein expression, whereas there was no

significant difference in Mfn2 protein expression in AKAP1[-/-] retina (Fig. 5a). Using EM and

electron microscope (EM) tomography analyses, we further assessed changes of mitochondrial

ultrastructure by measuring mitochondrial volume density, number, cross-sectional area and

cristae abundance in AKAP1[-/-] RGC somas. In comparison with WT mice, representative

volumes generated by EM tomography provided evidence for mitochondrial fission and loss of

mitochondrial mass (Fig. 5, b-d). Quantitative analysis showed significant decreases of

mitochondrial volume density ($13 \pm 0.7\%$), and mitochondrial cross-sectional area (0.14 ± 0.02

μm^2), but a significant increase of mitochondrial number (0.92 ± 0.06 per μm^2) in AKAP1[-/-]

RGC somas compared with WT (22 ± 2 %, mitochondrial volume density; 0.20 ± 0.01 μm^2,

mitochondrial cross-sectional area ; 0.67 ± 0.05 per μm^2, mitochondrial number) (Fig. 5c).

However, there was no significant difference in mitochondrial cristae abundance, which is the

ratio of cristae membrane surface area divided by the mitochondrial outer membrane surface

area, between WT and AKAP1[-/-] RGC somas (Fig. 5c).   Using three-dimensional (3D)

reconstruction of tomographic volume, we observed that AKAP1[-/-] mitochondria are

significantly smaller than the WT mitochondria. Smaller mitochondria should have fewer

cristae, given that cristae shape doesn't change a lot, but the density of cristae was about the same

between WT and AKAP1[-/-] (Fig. 5d). These results suggest that AKAP1 loss induces an

imbalance of mitochondrial dynamics by triggering mitochondrial fission and loss in RGCs.

AKAP1 loss promotes mitochondrial abnormalities and mitophagy in cardiac injury[46]. Based on

our current findings of mitochondrial fission and loss, we determined whether AKAP1 loss

induces mitophagosome formation. Indeed, AKAP1 loss significantly enhanced LC3-II protein

expression but decreased p62 protein expression in AKAP1[-/-] retina (Fig. 6a). Our results

showed strong LC3 immunoreactivities in RGC somas and axons in the GCL of AKAP1[-/-] mice

(Fig. 6b). Notably, segmented volumes from 3D tomographic reconstructions using EM

tomography revealed examples of degraded mitochondria with severe cristae depletion that were

engulfed in mitophagosomes in RGC soma of AKAP1[-/-] mice (Fig. 6c-j).

AKAP1 loss results in OXPHOS dysfunction and induces oxidative stress in the retina

Since AKAP1 deletion increased superoxide production in primary hippocampal and cortical

neurons in response to glutamate excitotoxicity as well as resulted in OXPHOS complex (Cx) II

dysfunction[24], we determined whether loss of AKAP1 alters OXPHOS Cxs and induces

oxidative stress in the retina. We further observed that AKAP1 loss significantly increases

OXPHOS Cx II but decreased OXPHOS Cx III-V protein expression in AKAP1[-/-] retina (Fig.

7a). However, there was no significant difference in OXPHOS Cx I protein expression between

WT and AKAP1[-/-] retinas (Fig. 7a). We observed a significant increase of superoxide dismutase

2 (SOD2) protein expression in AKAP1[-/-] retina (Fig. 7b). We also observed that SOD2 protein

expression was increased in the inner retina, including RGCs in the GCL, of AKAP1[-/-] mice (Fig.

7c and d). These results suggest that AKAP1 loss compromises OXPHOS and may contribute to

oxidative stress.

AKAP1 loss inactivates Akt and activates Bim/Bax pathway in the retina

We examined the expression levels of Akt protein expression, and Akt phosphorylation at serine

473 (Ser473) and threonine 308 (Thr308) in AKAP1[-/-] retina. We observed a significant increase

of total Akt protein expression but a significant dephosphorylation Akt at Ser473 and Thr308

protein expression in AKAP1[-/-] retina (Fig. 8a). Consistently, our results showed that Akt

immunoreactivity was increased in the INL and GCL of AKAP1[-/-] retina (Fig. 8b and c). More

specifically, increased Akt immunoreactivity was prominent in Brn3a-positive RGCs of AKAP1[-]

[/-] mice (Fig. 8b and c). Because Akt inactivation is associated with Bim/Bax activation and

intrinsic apoptotic cell death[47,48], we further determined whether AKAP1 loss activates Bim/Bax

in the retina. We observed significant increases of Bim and Bax protein expression in AKAP1[-/-]

retina (Fig. 8d). In addition, we observed that Bax immunoreactivity was increased in the GCL

of AKAP1[-/-] retina (Fig. 8d). More specifically, increased Bax immunoreactivity was prominent

in RBPMS-positive RGCs of AKAP1[-/-] mice (Fig. 8e and f).

Discussion

We here demonstrated for the first time that RGCs showed AKAP1 immunoreactivity in D2-*Gpnmb+* mice and elevated IOP triggered loss of AKAP1 in glaucomatous RGCs. In addition, elevated IOP increased CaN and total Drp1 levels in glaucomatous RGCs. It also impaired Drp1 phosphorylation at Ser637 in glaucomatous retina. These results suggest that AKAP1 and Drp1 phosphorylation at Ser637 have critical roles in RGC mitochondria and survival against glaucomatous damage. We also observed that loss of AKAP1 increased CaN and total Drp1 levels in RGCs, and compromised Drp1 phosphorylation at Ser637 in the retina, leading to mitochondrial fission and loss, and mitophagosome formation in RGCs. In addition, loss of AKAP1 resulted in OXPHOS dysfunction, as well as Akt inactivation and Bim/Bax activation; these effects contribute to glaucomatous neurodegeneration.

Impairment of mitochondrial dynamics is critically involved in glaucomatous RGCs and their axon degeneration[8,9,11,36], and inhibition of Drp1 rescues RGCs and their axons by preserving mitochondrial integrity in experimental glaucoma[11,36]. Our previous study demonstrated that elevated IOP showed a significant increase of total Drp1 level, while it did not alter phosphorylation of Drp1 at Ser616 in glaucomatous retina[11]; these effects suggest that elevated IOP-mediated mitochondrial fission activity is not dependent on Drp1 phosphorylation at Ser616 in glaucomatous RGC degeneration. Hence, our novel finding that glaucomatous RGCs lacking AKAP1 trigger CaN-mediated dephosphorylation of Drp1 at Ser637 suggest an important rationale to further investigate the impact of AKAP1 loss, mitochondrial fission and its related signaling pathway in RGCs.

AKAP1 is an important regulator of mitochondrial function by PKA anchoring and Drp1 phosphorylation at Ser637, and overexpression of AKAP1 is neuroprotective in neuronal cells against the mitochondrial toxin rotenone[20,21,23,24,43]. Emerging evidence showed that mice lacking AKAP1 exhibit increased infarct following transient cerebral ischemia as well as increased Drp1 localization to mitochondria from the forebrain tissues and decreased Drp1 phosphorylation at Ser637[24]. Our findings of enhanced CaN-mediated Drp1 phosphorylation at Ser637, and extensive mitochondrial fission and loss in the retina of AKAP1[-/-] mice suggest that AKAP1 loss directly contributes to Drp1 phosphorylation-dependent mitochondrial fission and loss in glaucomatous RGCs. This notion is also supported by the evidence of increasing LC3-II and decreasing p62 levels, as well as mitophagosome formation that contains degrading

mitochondria in RGC somas of AKAP1[-/-] mice. In addition, it is supported by the evidence of

our previous report showing autophagosome/mitophagosome formation in RGC axons of

glaucomatous DBA/2J mice[11].

AMPK is a nutrient and energy sensor that maintains energy homeostatsis[49] and mediates

mitochondrial fission in response to energy stress[50]. It has been reported that there was evidence

of AMPK activation by increasing AMPK phosphorylation at Thr172 in the retina and optic

nerve in glaucomatous DBA/2J mice[14,51]. Also, it has been proposed that AMPK uncouples

PKA from D-AKAP1 (AKAP140/149 and other splice variants AKAP121, sAKAP84) to

promote mitochondrial fission and mitophagy[52]. Furthermore, AMPK is associated with the

regulation of Drp1 phosphorylation[53,54]. In the current study, our findings showed no significant

evidence of AMPK activation, suggesting that AKAP1 loss may not directly contribute to

AMPK activation in the retina. Because energy stress activates AMPK[14,55], it is also possible

that AKAP1 loss may not trigger an excessive energy stress and substantial energy deficit in the

retina. In the line of this notion, our results showed that the level of dephosphorylation of Drp1

Ser637 in AKAP1[-/-] retina was relatively modest compared to the level of dephosphorylation of

Drp1 Ser637 in glaucomatous retina. Hence, our results intriguingly raise a possibility that

combined induction of both AMPK activation and AKAP1 loss may exacerbate mitochondrial

fission and loss in glaucomatous RGCs and their axons, leading to the promotion of energy crisis

and cell death. We believe that further studies will be required to investigate the relationship

between AMPK activation and AKAP1 loss in glaucomatous RGCs.

The increase in LC3-II accumulates autophagosomes, but does not guarantee autophagic

degradation. If, however, the amount of LC3-II further accumulates in the presence of lysosomal

protease inhibitors, this would indicate enhancement of autophagic flux[56]. On the other hand, the

protein expression level of p62/SQSTM1, a polyubiquitin binding protein, serves as autophagic

receptor and is a good indicator of autophagic flux[57-59]. Also, p62 protein expression has been

used as autophagic marker with LC3-II[60]. Although p62 level alone is not indicative of

activation of autophagy, p62 is also a marker for the degradation phase of autophagosomes and

has a lot of functions beyond autophagy[59]. In the current study, our findings showed that loss of

AKAP1 increased LC3-II protein expression but decreased p62 protein expression in AKAP1[-/-]

retina. Intriguingly, AKAP1[-/-] heart reduced p62 protein expression and enhanced mitophagy and

apoptosis[46]. Collectively, our results suggest that the increased LC3-II and decreased p62

contribute to the induction of autophagy/mitophagy in AKAP1[-/-] retina. Hence, we raise a

possibility that loss of AKAP1 may not only induce autophagosome/mitophagosome formation,

but also trigger increase of autophagic flux and degradation in RGCs. Also, whether this

autophagic/mitophagic process is a compensatory or pathological pathway in AKAP1[-/-] retina

should be considered in the future studies.

Previous study indicated the notion that AKAP1 regulates mitochondrial dynamics by

posttranslational modifications, but levels of Drp1, OPA1 and Mfn2 were unaltered in total

forebrain homogenates from AKAP1[-/-] mice[24]. Of interest, however, we observed a decrease of

total OPA1, but an increase of total Drp1 and Mfn1 in the retina of AKAP1[-/-] mice. Inner

membrane-anchored long OPA1 (L-OPA1) undergoes proteolytic cleavage resulting in short

OPA1 (S-OPA1)[61]. Cells contain a mixture of L-OPA1, required for mitochondrial fusion and S-

OPA, which facilitate mitochondrial fission under normal conditions. Disruption of this balance

in times of stress, result in mitochondrial fragmentation[61]. Thus, these unexpected, but important

results propose the possibility that AKAP1 may contribute to the regulation of mitochondrial

dynamics not only by posttranslational modifications, but also by alteration of fusion/fission

protein expression in the retina. Future studies will be needed to clarify the molecular

mechanisms underlying the alteration of fusion/fission proteins in the retina lacking AKAP1.

Emerging evidence showed that AKAP1 deletion results in OXPHOS Cx II dysfunction

in neuronal cells[24]. OXPHOS Cx II is a source of the mitochondrial reserve respiratory capacity

that is regulated by metabolic sensor and promotes cell survival against hypoxia. In the present

study, the observation that increasing OXPHOS Cx II, but decreasing OXPHOS Cxs III-V

activities triggered by loss of AKAP1 in the retina is likely to imply that AKAP1 plays a critical

role in OXPHOS function in RGCs. Moreover, increasing OXPHOS Cx II activity may be an

important endogenous compensatory defense mechanism in response to oxidative stress.

Further, this imbalance of OXPHOS Cxs suggests another possibility that AKAP1 loss may

contribute to an increase of ROS production and decrease of ATP production in the retina that

are previously reported in pressure and/or oxidative stress-induced RGC degeneration[11]. In

particular, association of POAG with polymorphism of mitochondrial cytochrome c oxidase

subunit 1 of OXPHOS Cx IV suggests a potential role of abnormal OXPHOS-mediated

mitochondrial dysfunction in glaucoma pathogenesis[3,4]. Therefore, these results collectively

suggest that AKAP1 loss in glaucomatous retina contributes to OXPHOS dysfunction that is

associated with metabolic and oxidative stress by increasing ROS production and decreasing

ATP reduction, leading to RGC death during glaucomatous neurodegeneration.

Loss of AKAP1 inactivates Akt signaling and increases apoptosis in cardiac

dysfunction[62,63]. Akt pathway controls the expression of apoptosis-related gene through

transcription factors such as FoxO (FoxO1 and FoxO3a)[64,65] and Akt activation promotes cell

survival by inhibiting FoxO3a/Bim/Bax pathway[48,64,65]. The pro-apoptotic Bcl-2 homology

domain 3-only protein Bim induces apoptosis, primarily through its increased binding activity

toward multiple pro-survival Bcl-2-like proteins, whose dissociations activate Bak and Bax[66,67].

In addition, Bax-mediated mitochondrial outer membrane permeabilization is an important

pathophysiological mechanism for metabolic dysfunction and cell death[68,69], and Bax activation

plays a critical role in mitochondrial dysfunction-mediated RGC degeneration[11,12]. Because

Drp1 contributes to Bax oligomerization, mitochondrial fission and the cellular apoptotic

pathway[70-72], increased total Drp1 and Bax levels in AKAP1[-/-] retinas suggest that AKAP1 loss

may contribute to Drp1-mediated Bax oligomerization, leading to mitochondrial fission. Recent

evidence from our group and others indicated that activation of cAMP/PKA pathway promotes

RGC survival [48,73-75] and that AKAP1-mediated neuroprotection requires PKA anchoring and

Drp1 phosphorylation at Ser637[21,23,24,43,76]. In this regard, our observation that AKAP1[-/-] retinas

dephosphorylates Akt at Ser473 and SerT308, and activates Bim/Bax pathway raise the

possibility that AKAP1 may be critical to RGC protection by modulating Akt/Bim/Bax pathway

against glaucomatous damage. Future studies will be necessary to examine whether

overexpression of AKAP1 rescues RGCs by promoting Drp1 phosphorylation at S637,

maintaining metabolic activity, activating Akt pathway and inhibiting Bim/Bax pathway.

In summary, our results represent the first direct evidence of a Drp1 phosphorylation-

mediated mitochondrial pathogenic mechanism that leads to mitochondrial fission and metabolic

dysfunction in glaucomatous RGC degeneration. Also, we provide evidence for the first time that

AKAP1 loss-mediated CaN activation and Drp1 dephosphorylation at Ser637 result in

mitochondrial fission, mitophagosome formation, OXPHOS dysfunction and oxidative stress, as

well as Akt inactivation and Bim/Bax activation in RGCs. Therefore, we propose

overexpression of AKAP1 and modulation of Drp1 phosphorylation at Ser637 as therapeutic

strategies for neuroprotective intervention in glaucoma and other mitochondria-related optic

neuropathies.

Materials and methods

Animals

Adult female DBA/2J and D2-*Gpnmb*$^+$ mice (The Jackson Laboratory), and adult male WT and AKPA1$^{-/-}$ mice were housed in covered cages, fed with a standard rodent diet ad libitum, and kept on a12 h light/12 h dark cycle. Animals were assigned randomly to experimental and control groups. The following PCR primers were used to determine AKAP1 genotype: GGAGGCGATCACAGCAACAACCG (R), ATACAGAAGCAGATCACTCAGGAGG (F-WT), CAGTCCCAAGGCTCATTTCAGGCC (F-KO)[24]. To investigate the effect of IOP elevation, 20 10 month-old DBA/2J and 10 age-matched D2-*Gpnmb*$^+$ mice were used. To investigate the effect of AKAP1 loss, 10 10 month-old AKAP1$^{-/-}$ and 10 age-matched WT mice were used. All procedures concerning animals were in accordance with the Association for Research in Vision and Ophthalmology Statement for the Use of Animals in Ophthalmic Vision Research and under protocols approved by Institutional Animal Care and Use Committee at the University of California, San Diego.

Tissue preparation

Mice were anesthetized by an intraperitoneal (IP) injection of a cocktail of ketamine (100 mg/kg, Ketaset; Fort Dodge Animal Health) and xylazine (9 mg/kg, TranquiVed; VEDCO Inc.) prior to cervical dislocation. The investigators were not blinded during tissue and data collection from mice. Blinding was used during data analysis. For immunohistochemistry, the retinas were dissected from the choroids and fixed with 4% paraformaldehyde (Sigma-Aldrich) in phosphate buffered saline (PBS, pH 7.4) for 2 h at 4 °C. Retinas were washed several times with PBS then dehydrated through graded levels of ethanol and embedded in polyester wax. For Western blot and PCR analyses, extracted retinas were immediately used.

Western blot analyses

Harvested retinas ($n = 4$ retinas/group) were homogenized for 1 min on ice with a modified RIPA lysis buffer (150 mM NaCl, 1 mM EDTA, 1% NP-40, 0.1% SDS, 1 mM DTT, 0.5% sodium deoxycholate and 50 mM Tris-Cl, pH 7.6), containing complete protease inhibitors. The lysates were then centrifuged at 15,000 g for 15 min and protein amounts in the supernatants were measured by Bradford assay. Proteins (10-20 μg) were separated by SDS/PAGE and

electrotransferred to polyvinylidene difluoride membranes (GE Healthcare Bio-Science). The

membranes were blocked with 5% non-fat dry milk and PBS/0.1% Tween-20 (PBS-T) for 1 h at

room temperature and incubated with primary antibodies for overnight at 4 °C. Primary

antibodies are mouse monoclonal anti-Actin antibody (1:20,000; Millipore; Cat#, MAB1501),

rabbit monoclonal anti-AKAP1 (1:1000; Cell Signaling; Cat# 5203), rabbit polyclonal anti-Akt

antibody (1:5000; Cell Signaling; Cat# 92725), mouse monoclonal anti-phospho-Akt Ser473

antibody (1:100, Cell Signaling; Cat# 9271), rabbit monoclonal anti-phospho-Akt Thr308

antibody (1:1000, Cell Signaling; Cat# 4056), rabbit polyclonal anti-AMPKα1/2 antibody

(1:1000, Santa Cruz Biotechnology; Cat# sc25792), rabbit monoclonal anti-AMPKα Thr172

antibody (1:1000, Cell Signaling Cat# 2535), mouse monoclonal anti-Bax antibody (6A7; 1:500;

Santa Cruz Biotechnology; Cat# sc-23959), rabbit polyclonal anti-Bim antibody (1:100, Santa

Cruz Biotechnology; Cat# sc-8265), mouse monoclonal anti-CaN antibody (1:200, BD

Bioscience; Cat# 610259), mouse monoclonal anti-Drp1 antibody (1:1000; BD Biosciences;

Cat# 611113), rabbit polyclonal phospho-Drp1 Ser637 antibody (1:750; Cell Signaling; Cat#

4867), rabbit polyclonal phospho-Drp1 Ser616 antibody (1:750; Cell Signaling; Cat# 3455),

rabbit polyclonal anti-LC3 antibody (1:2000; MBL International; Cat# PM036), mouse

monoclonal anti-Mfn1 antibody (1:20000; Abcam; Cat# ab57602), mouse monoclonal anti-Mfn2

antibody (1:2000; Abcam; Cat# ab56889), mouse monoclonal anti-OPA1 antibody (1:2000; BD

Biosciences; Cat# 612607), rat/mouse monoclonal anti-OXPHOS Cx Cocktail Kit (1:1000;

Invitrogen; Cat# 458099), rabbit polyclonal p62 antibody (1:1000; MBL International; Cat#

PM045) and rabbit polyclonal anti-SOD2 antibody (1:5000; Santa Cruz; Cat# sc-30080).

Membrane were washed several times with PBS-T then incubated with peroxidase-conjugated

goat anti-mouse or rabbit IgG (1:1000-7000; Bio-Rad; Cat# 1721011 or 1706515) for 1 h at

room temperature and developed using enhanced chemiluminescence substrate system. The

images were captured and quantified by using ImageQuant™ LAS 4000 system (GE Healthcare

Bio-Science) and band densities were normalized to that of actin, total Drp1, total Akt, or total

AMPK.

Immunohistochemistry

Immunohistochemical staining of 7 μm wax sections of full thickness retina were performed.

Sections from wax blocks from each group ($n = 4$ retinas/group) were used for

immunohistochemical analysis. Primary antibodies included rabbit monoclonal anti-AKAP1

antibody (1:100, Cell Signaling; Cat# 5203), rabbit polyclonal anti-Akt antibody (1:100, Cell

Signaling; Cat# 92725), mouse monoclonal anti-Bax antibody (6A7; 1:50, Santa Cruz

Biotechnology; Cat# sc-23959), goat polyclonal anti-Brn3a antibody (1:500, Santa Cruz

Biotechnology; Cat# sc-23959), mouse monoclonal anti-CaN (1:200, BD Bioscience; Cat#

610259), mouse monoclonal anti-Drp1 antibody (1:25; BD Biosciences; Cat# 611113), rabbit

polyclonal anti-Drp1 (1:50, Santa Cruz Biotechnology, Cat# sc-32898), rabbit polyclonal anti-

LC3 antibody (1:2000; MBL International) ; Cat# PM036, rabbit polyclonal anti-RBPMS (1:300,

Novus; Cat# NBP2-20112), rabbit polyclonal anti-SOD2 antibody (1:300, Santa Cruz

Biotechnology; Cat# sc-30080) and mouse monoclonal anti-TUJ1 (1:300, BioLegend; Ct#

801202). To prevent non-specific background, tissues were incubated in 1% bovine serum

albumin/PBS for 1 h at room temperature before incubation with the primary antibodies for 16 h

at 4°C. After several wash steps, the tissues were incubated with the secondary antibodies,

Alexa Fluor 488 or 568 dye-conjugated donkey anti-mouse IgG (1:100, Invitrogen; Cat# R37114

or A10037), Alexa Fluor 488 or 568 dye-conjugated donkey anti-rabbit IgG (1:100, Invitrogen;

Cat# R37118 or A10042) and Alexa Fluor 568 dye-conjugated donkey anti-goat IgG (1:100,

Invitrogen; Cat# A11057) for 4 h at 4°C and subsequently washed with PBS. The sections were

counterstained with the nucleic acid stain Hoechst 33342 (1 μg/mL; Invitrogen) in PBS. Images

were acquired with confocal microscopy (Olympus FluoView1000; Olympus, Tokyo, Japan).

Each target protein fluorescent integrated intensity in pixel per area was measured using the

ImageJ software[77-79]. All imaging parameters was remained the same and corrected with the

background subtraction.

RT-PCR and quantitative real-time RT-PCR

Total RNA was isolated from retinas of each group (n = 4 retinas/group) using the Absolutely

RNA Miniprep Kit (Stratagene), according to the manufacturer's protocol as previously

described[48]. cDNAs were synthesized from total RNAs with iScript cDNA Synthesis Kit (Bio-

Rad) according to the manufacturer's protocols. Reverse Transcription (RT)-PCR was

performed with cDNAs synthesized from 1 μg of the total RNA of each group as a template and

specific primers for: AKAP1, CTGCCAGTCAGTACTCAGCC (F),

CTTTGGCACCTCGATCTCCC (R), OPA1, CATCTACCTTCCAGCTGCCC (F),

TCTCCTCCTTCACAGCCTCC (R), and Mfn1, GTTTTCCCTGGGCTGGTCTT (F),

TTTTCCAAATCACGCCCCCA (R). For the quantification of the relative mRNA expressions

of each group, real-time PCR was carried out using MX3000P real-time PCR system

(Stratagene) as follows. cDNAs were amplified using SSO Adv. Universal SYBR Green

SuperMix (Bio-Rad) and the specific primers for 40 cycles (95 °C for 5 min, and 40 cycles

(95 °C for 30 sec, 60 °C for 30 sec, and 72 °C for 30 sec)). Output data were obtained

as *Ct* values and the differential mRNA expression of the target genes among groups was

calculated using the comparative *Ct* method. GAPDH mRNA, an internal control, was amplified

along with the target gene, and the *Ct* value of GAPDH was used to normalize the expression of

the target gene.

Transmission electron microscopy

For conventional EM, two eyes from each group ($n = 2$ mice) were fixed via cardiac perfusion

with 2% paraformaldehyde, 2.5% glutaraldehyde (Ted Pella, Redding, CA) in 0.15M sodium

cacodylate (pH 7.4, Sigma) solution at 37°C and placed in pre-cooled fixative of the same

composition on ice for 1 h. The following procedure was used to optimize mitochondrial

structural preservation and membrane contrast. The retinas were dissected in a solution of

0.15M sodium cacodylate plus 3mM calcium chloride (pH 7.4) on ice and then post-fixed with a

1% osmium tetroxide, 0.8% potassium ferrocyanide, 3mM calcium chloride in 0.1M sodium

cacodylate solution (pH 7.4) for 1 h, washed with ice-cold distilled water, poststained with 2%

uranyl acetate solution at 4°C, dehydrated using graded ethanols, and embedded in Durcupan

resin (Fluka). A similar procedure was used for cultured primary RGCs from each group.

Ultrathin (70 nm) sections were post-stained with uranyl acetate and lead salt solutions and

evaluated using a JEOL 1200FX transmission EM operated at 80kV. Images were recorded on

film at 8,000X magnification. The negatives were digitized at 1800 dpi using a Nikon Cool scan

system, giving an image size of 4033 x 6010 pixel array and a pixel resolution of 1.77 nm. For

quantitative analysis, the number of mitochondria was normalized to the total area occupied by

axons in each image, which was measured using ImageJ (http://rsb.info.nih.gov/ij/).

Mitochondrial lengths were measured with ImageJ. The mitochondrial volume density, defined

as the volume occupied by mitochondria divided by the volume occupied by the axoplasm, was

estimated using stereology as follows. A 112 x 112 square grid (112 x 112 chosen for ease of use

with Photoshop) was overlaid on each image loaded in Photoshop (Adobe Systems Inc.), and

mitochondria and axoplasm lying under intercepts were counted. The relative volume of

mitochondria was expressed as the ratio of intercepts coinciding with this organelle relative to

the intercepts coinciding with axoplasm.

Electron microscope tomography

Sections of retina tissues from each group were cut at a thickness of 400 nm. For each

reconstruction, a double-tilt series of images at one degree tilt increment was collected with an

FEI titan hibase electron microscope operated at 300 kV. Images were recorded with a Gatan

4Kx4K CCD camera. The magnification was 11,000x and the pixel resolution was 0.81 nm.

The IMOD package was used for alignment, reconstruction and volume segmentation. Volume

segmentation was performed by manual tracing of membranes in the planes of highest resolution

with the Drawing Tools and Interpolator plug-ins[80]. The mitochondrial reconstructions and

surface-rendered volumes were visualized using 3DIMOD. Measurements of mitochondrial

outer, inner boundary, and cristae membrane surface areas and volumes were made within

segmented volumes using IMODinfo. These were used to determine the cristae abundance,

defined as the ratio: sum of the cristae membrane surface areas divided by the mitochondrial

outer membrane surface area. Movies of the tomographic volume were made using IMODmovie.

Statistical analyses

Data were shown as mean ± S.D. Comparison of experimental conditions was evaluated using

the two-tailed unpaired Student's t-test between groups. $P < 0.05$ was considered to be

statistically significant.

Acknowledgements

This work was supported, in part, by NIH grants EY018658, P30 EY022589, T32 EY026590,

5P41GM103412, and an unrestricted grant from Research to Prevent Blindness (New York, NY)

and UCSD Academic Senate.

Conflict of Interest

The authors declare no conflict of interest.

References

559 1   Weinreb, R. N. & Khaw, P. T. Primary open-angle glaucoma. *Lancet* **363**, 1711-1720
560   (2004).
561

562 2   Abu-Amero, K. K. & Bosley, T. M. Mitochondrial abnormalities in patients with LHON-
563   like optic neuropathies. *Invest Ophthalmol Vis Sci* **47**, 4211-4220 (2006).
564

565 3   Collins, D. W. *et al.* The MT-CO1 V83I Polymorphism is a Risk Factor for Primary
566   Open-Angle Glaucoma in African American Men. *Invest Ophthalmol Vis Sci* **59**, 1751-
567   1759 (2018).
568

569 4   Collins, D. W. *et al.* Association of primary open-angle glaucoma with mitochondrial
570   variants and haplogroups common in African Americans. *Mol Vis* **22**, 454-471 (2016).
571

572 5   Bosley, T. M. *et al.* Down-regulation of OPA1 in patients with primary open angle
573   glaucoma. *Mol Vis* **17**, 1074-1079 (2011).
574

575 6   Lee, S. *et al.* Impaired complex-I-linked respiration and ATP synthesis in primary open-
576   angle glaucoma patient lymphoblasts. *Invest Ophthalmol Vis Sci* **53**, 2431-2437 (2012).
577

578 7   Singh, L. N. *et al.* Mitochondrial DNA Variation and Disease Susceptibility in Primary
579   Open-Angle Glaucoma. *Invest Ophthalmol Vis Sci* **59**, 4598-4602 (2018).
580

581 8   Ju, W. K. *et al.* Intraocular pressure elevation induces mitochondrial fission and triggers
582   OPA1 release in glaucomatous optic nerve. *Invest Ophthalmol Vis Sci* **49**, 4903-4911
583   (2008).
584

585 9   Ju, W. K. *et al.* Increased optic atrophy type 1 expression protects retinal ganglion cells
586   in a mouse model of glaucoma. *Mol Vis* **16**, 1331-1342 (2010).
587

588 10   Lee, D. *et al.* Coenzyme Q10 inhibits glutamate excitotoxicity and oxidative stress-
589   mediated mitochondrial alteration in a mouse model of glaucoma. *Invest Ophthalmol Vis*
590   *Sci* **55**, 993-1005 (2014).
591

592 11   Kim, K. Y. *et al.* DRP1 inhibition rescues retinal ganglion cells and their axons by
593   preserving mitochondrial integrity in a mouse model of glaucoma. *Cell Death Dis* **6**,
594   e1839 (2015).
595

596 12   Shim, M. S. *et al.* Mitochondrial pathogenic mechanism and degradation in optineurin
597   E50K mutation-mediated retinal ganglion cell degeneration. *Sci Rep* **6**, 33830 (2016).
598

599 13   Williams, P. A. *et al.* Vitamin B3 modulates mitochondrial vulnerability and prevents
600   glaucoma in aged mice. *Science* **355**, 756-760 (2017).
601

602   14   Harun-Or-Rashid, M. *et al.* Structural and Functional Rescue of Chronic Metabolically
603   Stressed Optic Nerves through Respiration. *J Neurosci* **38**, 5122-5139 (2018).
604

605   15   Herberg, F. W., Maleszka, A., Eide, T., Vossebein, L. & Tasken, K. Analysis of A-kinase
606   anchoring protein (AKAP) interaction with protein kinase A (PKA) regulatory subunits:
607   PKA isoform specificity in AKAP binding. *J Mol Biol* **298**, 329-339 (2000).
608

609   16   Affaitati, A. *et al.* Essential role of A-kinase anchor protein 121 for cAMP signaling to
610   mitochondria. *J Biol Chem* **278**, 4286-4294, doi:10.1074/jbc.M209941200 (2003).
611   17   Rubin, C. S. A kinase anchor proteins and the intracellular targeting of signals carried by
612   cyclic AMP. *Biochim Biophys Acta* **1224**, 467-479 (1994).
613

614   18   Colledge, M. & Scott, J. D. AKAPs: from structure to function. *Trends Cell Biol* **9**, 216-
615   221 (1999).
616

617   19   Huang, L. J., Durick, K., Weiner, J. A., Chun, J. & Taylor, S. S. D-AKAP2, a novel
618   protein kinase A anchoring protein with a putative RGS domain. *Proc Natl Acad Sci U S*
619   *A* **94**, 11184-11189 (1997).
620

621   20   Merrill, R. A. & Strack, S. Mitochondria: a kinase anchoring protein 1, a signaling
622   platform for mitochondrial form and function. *Int J Biochem Cell Biol* **48**, 92-96 (2014).
623

624   21   Dickey, A. S. & Strack, S. PKA/AKAP1 and PP2A/Bbeta2 regulate neuronal
625   morphogenesis via Drp1 phosphorylation and mitochondrial bioenergetics. *J Neurosci*
626   **31**, 15716-15726 (2011).
627

628   22   Gomes, L. C., Di Benedetto, G. & Scorrano, L. During autophagy mitochondria elongate,
629   are spared from degradation and sustain cell viability. *Nat Cell Biol* **13**, 589-598 (2011).
630

631   23   Merrill, R. A. *et al.* Mechanism of neuroprotective mitochondrial remodeling by
632   PKA/AKAP1. *PLoS Biol* **9**, e1000612 (2011).
633

634   24   Flippo, K. H. *et al.* AKAP1 Protects from Cerebral Ischemic Stroke by Inhibiting Drp1-
635   Dependent Mitochondrial Fission. *J Neurosci* **38**, 8233-8242 (2018).
636

637   25   Knott, A. B., Perkins, G., Schwarzenbacher, R. & Bossy-Wetzel, E. Mitochondrial
638   fragmentation in neurodegeneration. *Nat Rev Neurosci* **9**, 505-518 (2008).
639

640   26   Burte, F., Carelli, V., Chinnery, P. F. & Yu-Wai-Man, P. Disturbed mitochondrial
641   dynamics and neurodegenerative disorders. *Nat Rev Neurol* **11**, 11-24 (2015).
642

643   27   Chen, H. & Chan, D. C. Emerging functions of mammalian mitochondrial fusion and
644   fission. *Hum Mol Genet* **14 Spec No. 2**, R283-289 (2005).
645

646   28   Okamoto, K. & Shaw, J. M. Mitochondrial morphology and dynamics in yeast and
647   multicellular eukaryotes. *Annu Rev Genet* **39**, 503-536 (2005).

648 29   Cho, D. H. *et al.* S-nitrosylation of Drp1 mediates beta-amyloid-related mitochondrial
649   fission and neuronal injury. *Science* **324**, 102-105 (2009).
650
651 30   Kim, J. *et al.* Mitochondrial loss, dysfunction and altered dynamics in Huntington's
652   disease. *Hum Mol Genet* **19**, 3919-3935 (2010).
653
654 31   Song, W. *et al.* Mutant huntingtin binds the mitochondrial fission GTPase dynamin-
655   related protein-1 and increases its enzymatic activity. *Nat Med* **17**, 377-382 (2011).
656
657 32   Dowding, J. M. *et al.* Cerium oxide nanoparticles protect against Abeta-induced
658   mitochondrial fragmentation and neuronal cell death. *Cell death and differentiation* **21**,
659   1622-1632 (2014).
660
661 33   Cereghetti, G. M. *et al.* Dephosphorylation by calcineurin regulates translocation of Drp1
662   to mitochondria. *Proc Natl Acad Sci U S A* **105**, 15803-15808 (2008).
663
664 34   Cribbs, J. T. & Strack, S. Reversible phosphorylation of Drp1 by cyclic AMP-dependent
665   protein kinase and calcineurin regulates mitochondrial fission and cell death. *EMBO Rep*
666   **8**, 939-944 (2007).
667
668 35   Chang, C. R. & Blackstone, C. Cyclic AMP-dependent protein kinase phosphorylation of
669   Drp1 regulates its GTPase activity and mitochondrial morphology. *The Journal of
670   biological chemistry* **282**, 21583-21587 (2007).
671
672 36   Park, S. W. *et al.* A selective inhibitor of drp1, mdivi-1, increases retinal ganglion cell
673   survival in acute ischemic mouse retina. *Invest Ophthalmol Vis Sci* **52**, 2837-2843 (2011).
674
675 37   Ju, W. K. *et al.* Elevated hydrostatic pressure triggers mitochondrial fission and decreases
676   cellular ATP in differentiated RGC-5 cells. *Invest Ophthalmol Vis Sci* **48**, 2145-2151
677   (2007).
678
679 38   John, S. W., Hagaman, J. R., MacTaggart, T. E., Peng, L. & Smithes, O. Intraocular
680   pressure in inbred mouse strains. *Invest Ophthalmol Vis Sci* **38**, 249-253 (1997).
681 39   Libby, R. T. *et al.* Inherited glaucoma in DBA/2J mice: pertinent disease features for
682   studying the neurodegeneration. *Vis Neurosci* **22**, 637-648 (2005).
683
684 40   Libby, R. T. *et al.* Susceptibility to neurodegeneration in a glaucoma is modified by Bax
685   gene dosage. *PLoS Genet* **1**, 17-26 (2005).
686
687 41   Abrenica, B., AlShaaban, M. & Czubryt, M. P. The A-kinase anchor protein AKAP121 is
688   a negative regulator of cardiomyocyte hypertrophy. *J Mol Cell Cardiol* **46**, 674-681
689   (2009).
690
42   Rogne, M., Stokka, A. J., Tasken, K., Collas, P. & Kuntziger, T. Mutually exclusive
binding of PP1 and RNA to AKAP149 affects the mitochondrial network. *Hum Mol
Genet* **18**, 978-987 (2009).

43 Dagda, R. K. *et al.* Mitochondrially localized PKA reverses mitochondrial pathology and
dysfunction in a cellular model of Parkinson's disease. *Cell Death Differ* **18**, 1914-1923
(2011).
697

698 44 Kim, H. *et al.* Fine-tuning of Drp1/Fis1 availability by AKAP121/Siah2 regulates
699   mitochondrial adaptation to hypoxia. *Mol Cell* **44**, 532-544 (2011).
700

701 45 Huang, W. *et al.* Calcineurin cleavage is triggered by elevated intraocular pressure, and
702   calcineurin inhibition blocks retinal ganglion cell death in experimental glaucoma. *Proc*
703   *Natl Acad Sci U S A* **102**, 12242-12247 (2005).
704

705 46 Schiattarella, G. G. *et al.* Akap1 Deficiency Promotes Mitochondrial Aberrations and
706   Exacerbates Cardiac Injury Following Permanent Coronary Ligation via Enhanced
707   Mitophagy and Apoptosis. *PLoS One* **11**, e0154076 (2016).
708

709 47 Kong, D. *et al.* Insulin-like growth factor 1 rescues R28 retinal neurons from apoptotic
710   death through ERK-mediated BimEL phosphorylation independent of Akt. *Exp Eye Res*
711   **151**, 82-95 (2016).
712

713 48 Shim, M. S. *et al.* Elevated intracellular cAMP exacerbates vulnerability to oxidative
714   stress in optic nerve head astrocytes. *Cell Death Dis* **9**, 285 (2018).
715

716 49 Hardie, D. G., Ross, F. A. & Hawley, S. A. AMPK: a nutrient and energy sensor that
717   maintains energy homeostasis. *Nat Rev Mol Cell Biol* **13**, 251-262 (2012).
718

719 50 Toyama, E. Q. *et al.* Metabolism. AMP-activated protein kinase mediates mitochondrial
720   fission in response to energy stress. *Science* **351**, 275-281 (2016).
721

722 51 Harun-Or-Rashid, M. & Inman, D. M. Reduced AMPK activation and increased HCAR
723   activation drive anti-inflammatory response and neuroprotection in glaucoma. *J*
724   *Neuroinflammation* **15**, 313 (2018).
725

726 52 Zhang, J., Wang, Y., Liu, X., Dagda, R. K. & Zhang, Y. How AMPK and PKA Interplay
727   to Regulate Mitochondrial Function and Survival in Models of Ischemia and Diabetes.
728   *Oxid Med Cell Longev* **2017**, 4353510 (2017).
729

730 53 Li, J. *et al.* Pharmacological activation of AMPK prevents Drp1-mediated mitochondrial
731   fission and alleviates endoplasmic reticulum stress-associated endothelial dysfunction. *J*
732   *Mol Cell Cardiol* **86**, 62-74 (2015).
733

734 54 Zhou, H. *et al.* Empagliflozin rescues diabetic myocardial microvascular injury via
735   AMPK-mediated inhibition of mitochondrial fission. *Redox Biol* **15**, 335-346 (2018).
736

737 55 Garcia, D. & Shaw, R. J. AMPK: Mechanisms of Cellular Energy Sensing and
738   Restoration of Metabolic Balance. *Mol Cell* **66**, 789-800 (2017).

739   56   Mizushima, N. & Yoshimori, T. How to interpret LC3 immunoblotting. *Autophagy* **3**,
740   542-545 (2007).
741

742   57   Shin, J. P62 and the sequestosome, a novel mechanism for protein metabolism. *Arch*
743   *Pharm Res* **21**, 629-633 (1998).
744

745   58   Bitto, A. *et al.* P62/SQSTM1 at the interface of aging, autophagy, and disease. *Age*
746   *(Dordr)* **36**, 9626 (2014).
747

748   59   Sanchez-Martin, P. & Komatsu, M. p62/SQSTM1 - steering the cell through health and
749   disease. *J Cell Sci* **131**, jcs.222836 (2018).
750

751   60   Klionsky, D. J. *et al.* Guidelines for the use and interpretation of assays for monitoring
752   autophagy (3rd edition). *Autophagy* **12**, 1-222 (2016).
753

754   61   MacVicar, T. & Langer, T. OPA1 processing in cell death and disease - the long and
755   short of it. *J Cell Sci* **129**, 2297-2306 (2016).
756

757   62   Schiattarella, G. G. *et al.* Loss of Akap1 Exacerbates Pressure Overload-Induced Cardiac
758   Hypertrophy and Heart Failure. *Front Physiol* **9**, 558 (2018).
759

760   63   Schiattarella, G. G. *et al.* Akap1 Regulates Vascular Function and Endothelial Cells
761   Behavior. *Hypertension* **71**, 507-517 (2018).
762

763   64   Zhang, X., Tang, N., Hadden, T. J. & Rishi, A. K. Akt, FoxO and regulation of apoptosis.
764   *Biochim Biophys Acta* **1813**, 1978-1986, doi:10.1016/j.bbamcr.2011.03.010 (2011).
765   65   Brunet, A. *et al.* Akt promotes cell survival by phosphorylating and inhibiting a Forkhead
766   transcription factor. *Cell* **96**, 857-868 (1999).
767

768   66   Bouillet, P. *et al.* Proapoptotic Bcl-2 relative Bim required for certain apoptotic
769   responses, leukocyte homeostasis, and to preclude autoimmunity. *Science* **286**, 1735-
770   1738 (1999).
771

772   67   Willis, S. N. *et al.* Apoptosis initiated when BH3 ligands engage multiple Bcl-2
773   homologs, not Bax or Bak. *Science* **315**, 856-859 (2007).
774

775   68   Kirkland, R. A., Saavedra, G. M., Cummings, B. S. & Franklin, J. L. Bax regulates
776   production of superoxide in both apoptotic and nonapoptotic neurons: role of caspases. *J*
777   *Neurosci* **30**, 16114-16127 (2010).
778

779   69   Jiang, X., Jiang, H., Shen, Z. & Wang, X. Activation of mitochondrial protease OMA1 by
780   Bax and Bak promotes cytochrome c release during apoptosis. *Proc Natl Acad Sci U S A*
781   **111**, 14782-14787 (2014).
782

783   70   Karbowski, M. *et al.* Spatial and temporal association of Bax with mitochondrial fission
784   sites, Drp1, and Mfn2 during apoptosis. *J Cell Biol* **159**, 931-938 (2002).

785 71 Youle, R. J. & Strasser, A. The BCL-2 protein family: opposing activities that mediate
786 cell death. *Nat Rev Mol Cell Biol* **9**, 47-59 (2008).
787

788 72 Montessuit, S. *et al.* Membrane remodeling induced by the dynamin-related protein Drp1
789 stimulates Bax oligomerization. *Cell* **142**, 889-901 (2010).
790

791 73 Corredor, R. G. *et al.* Soluble adenylyl cyclase activity is necessary for retinal ganglion
792 cell survival and axon growth. *J Neurosci* **32**, 7734-7744 (2012).
793

794 74 Hellstrom, M. & Harvey, A. R. Cyclic AMP and the regeneration of retinal ganglion cell
795 axons. *Int J Biochem Cell Biol* **56**, 66-73 (2014).
796

797 75 Cueva Vargas, J. L., Belforte, N. & Di Polo, A. The glial cell modulator ibudilast
798 attenuates neuroinflammation and enhances retinal ganglion cell viability in glaucoma
799 through protein kinase A signaling. *Neurobiol Dis* **93**, 156-171 (2016).
800

801 76 Pryde, K. R., Smith, H. L., Chau, K. Y. & Schapira, A. H. PINK1 disables the anti-fission
802 machinery to segregate damaged mitochondria for mitophagy. *J Cell Biol* **213**, 163-171
803 (2016).
804

805 77 Joachim, S. C. *et al.* Retinal ganglion cell loss is accompanied by antibody depositions
806 and increased levels of microglia after immunization with retinal antigens. *PLoS One* **7**,
807 e40616 (2012).
808

809 78 Ju, W. K. *et al.* Increased mitochondrial fission and volume density by blocking
810 glutamate excitotoxicity protect glaucomatous optic nerve head astrocytes. *Glia* **63**, 736-
811 753 (2015).
812

813 79 Yoshida, M. *et al.* Overexpression of neurotrophin-3 stimulates a second wave of
814 dopaminergic amacrine cell genesis after birth in the mouse retina. *J Neurosci* **31**, 12663-
815 12673 (2011).
816

817 80 Perkins, G. *et al.* Electron tomography of neuronal mitochondria: three-dimensional
818 structure and organization of cristae and membrane contacts. *Journal of structural*
819 *biology* **119**, 260-272 (1997).
820

Figure legend

Fig. 1 AKAP1 deficiency in glaucomatous RGCs. (a) Western blot analysis for AKAP1 in the

retinas of 10-month-old glaucomatous DBA/2J and age-matched D2-*Gpnmb⁺* mice. (b)

Representative images from immunohistochemical analyses for AKAP1 (green) and TUJ1 (red)

in the retina of D2-*Gpnmb⁺* and glaucomatous DBA/2J mice. Arrowheads indicate accumulation

of AKAP1 co-labeled with TUJ1 in RGC somas and arrows indicate TUJ1-labeled axon bundles.

Note that glaucomatous RGCs showed decrease of AKAP1 protein expression. Blue color

indicates nucleus. (c) Quantitative analysis for fluorescent intensity showed a significant

decrease of AKAP1 immunoreactivity in the retina of glaucomatous DBA/2J mice. GCL,

ganglion cell layer; IPL, inner plexiform layer; INL, inner nuclear layer; OPL, outer plexiform

layer; ONL, outer nuclear layer. Mean $\pm$ SD; $n = 3$ (a) and $n = 10$ (c); $*P < 0.05$ and $***P <$

0.001 (two-tailed unpaired Student's t-test). Scale bar: 20 μm.

Fig. 2 CaN-mediated dephosphorylation of Drp1 at S637 in glaucomatous retina. (a)

Western blot analyses for CaN in the retinas of 10-month-old glaucomatous DBA/2J and age-

matched D2-*Gpnmb⁺* mice. (b) Representative images from immunohistochemical analyses for

CaN (green, arrowheads) co-labeled with RBPMS (red, arrowheads) in RGCs. Note that

glaucomatous RGCs showed increases of CaN protein expression. Blue color indicates nucleus.

(c) Quantitative analysis for fluorescent intensity showed a significant increase of CaN

immunoreactivity in the retina of glaucomatous DBA/2J mice. (d) Western blot analyses for

total Drp1 and phospho-Drp1 Ser637 in the retinas of glaucomatous DBA/2J and age-matched

D2-*Gpnmb⁺* mice. (e) Representative images from immunohistochemical analyses for total

Drp1 (green, arrowheads) co-labeled with TUJ1 (red, arrowheads) in RGCs. Note that

glaucomatous RGCs showed increases of total Drp1 protein expression. Blue color indicates

nucleus. (f) Quantitative analysis for fluorescent intensity showed a significant increase of Drp1

immunoreactivity in the retina of glaucomatous DBA/2J mice. GCL, ganglion cell layer; IPL,

inner plexiform layer; INL, inner nuclear layer; OPL, outer plexiform layer; ONL, outer nuclear

layer. Mean $\pm$ SD; $n = 3$ (a and d) and $n = 10$ (c and f); $**P < 0.01$ and $***P < 0.001$ (two-

tailed unpaired Student's t-test). Scale bar: 20 μm.

Fig. 3 CaN expression in AKAP1$^{-/-}$ retina. (a) Quantitative RT-PCR analysis for AKAP1 in
the retinas of AKAP1$^{-/-}$ and WT mice. (b) Western blot analysis for CaN in the retinas of
AKAP1$^{-/-}$ and WT mice. (c) Representative images from immunohistochemical analyses for
CaN (green, arrowheads) co-labeled with RBPMS (red, arrowheads) in RGCs. Note that
glaucomatous RGCs showed increases of CaN protein expression in the GCL and INL. Blue
color indicates nucleus. (d) Quantitative analysis for fluorescent intensity showed a significant
increase of CaN immunoreactivity in the retina of AKAP1$^{-/-}$ mice. GCL, ganglion cell layer;
IPL, inner plexiform layer; INL, inner nuclear layer; OPL, outer plexiform layer; ONL, outer
nuclear layer. Mean $\pm$ SD; $n = 3$ or 4 (a and b) and $n = 10$ (d); $**P < 0.01$ and $***P < 0.001$
(two-tailed unpaired Student's t-test). Scale bar: 20 μm.
868
Fig. 4 Phosphorylation of AMPK at Thr172 and Drp1 at Ser637 in AKAP1$^{-/-}$ retina. (a)
Western blot analysis for AMPK and phospho-AMPK Thr172 in the retinas of AKAP1$^{-/-}$ and
WT mice. (b) Western blot analysis for total Drp1, phospho-Drp1 Ser637 and phospho-Drp1
Ser616 in the retinas of AKAP1$^{-/-}$ and WT mice. (c) Representative images from
immunohistochemical analyses for total Drp1 (green, arrowheads) co-labeled with RBPMS (red,
arrowheads) in RGCs. Note that glaucomatous RGCs showed an increase of total Drp1 protein
expression in RGCs. Blue color indicates nucleus. (d) Quantitative analysis for fluorescent
intensity showed a significant increase of Drp1 immunoreactivity in the retina of AKAP1$^{-/-}$ mice.
GCL, ganglion cell layer; IPL, inner plexiform layer; INL, inner nuclear layer; OPL, outer
plexiform layer; ONL, outer nuclear layer. Mean $\pm$ SD; $n = 3$ or 6 (a) and $n = 10$ (d); $*P < 0.05$
(two-tailed unpaired Student's t-test). Scale bar: 20 μm.
880
Fig. 5 Mitochondrial fission in AKAP1$^{-/-}$ RGC somas. (a) Quantitative RT-PCR analyses for
OPA1 and Mfn1 and Western blot analysis for total OPA1 (L-OPA1 & S-OPA1), and Mfn1 and
2 in the retinas of AKAP1$^{-/-}$ and WT mice. (b-d) EM and EM tomography analyses for
mitochondrial fission and loss in AKAP1$^{-/-}$ RGC somas. (b) 1.6-nm thick slices through the
middle of tomographic volumes of WT RGC somas show an abundance of larger mitochondria.
In contrast, the mitochondria are smaller with a greater number in AKAP1$^{-/-}$ RGC somas, but
show no alteration of cristae architecture. (c) Quantitative analyses of mitochondrial volume
density, number and cross-sectional area and cristae abundance in WT and AKAP1$^{-/-}$ RGC

somas. Note that AKAP1$^{-/-}$ RGC somas show significant loss of mitochondrial mass and cross-

sectional area, and a significant increase in mitochondrial number. However, there was no

significant difference in cristae abundance of AKAP1$^{-/-}$ RGC somas. (d) Top view of the

surface-rendered volume showing 3 adjacent mitochondria from a WT tomographic volume

(left) and an AKAP1$^{-/-}$ volume (right). Comparing the side view with the top view provides

perspective on the distribution of the predominantly tubular cristae (in an assortment of colors)

and reveals substantial heterogeneity of cristae size. The mitochondrial outer membranes were

made translucent to better visualize the cristae. The number of cristae in each mitochondrion is

indicated. Mean $\pm$ SD (a) and Mean $\pm$ SEM (c); $n = 3$ (a), $n = 11$ (mitochondrial volume,

number and cristae abundance for WT and AKAP1$^{-/-}$, c) and $n = 22$ or 39 (mitochondrial size for

WT or AKAP1$^{-/-}$, c); *$P < 0.05$ and **$P < 0.01$ (two-tailed unpaired Student's t-test). Scale

bars: 200 nm.

Fig. 6 Mitophagosome formation in AKAP1$^{-/-}$ RGC somas. (a) Western blot analyses for

LC3 and p62 in the retinas of AKAP1$^{-/-}$ and WT mice. (b) Representative images from

immunohistochemical analyses for LC3 (green) co-labeled with TUJ1 (red) in RGCs and their

axons. Note that LC3 immunoreactivity was localized in RGC somas (arrowheads) and their

axons (arrows) in WT and AKAP1$^{-/-}$ retina. Blue color indicates nucleus. GCL, ganglion cell

layer; IPL, inner plexiform layer; INL, inner nuclear layer; OPL, outer plexiform layer; ONL,

outer nuclear layer. Scale bar, 20 μm. (c-j) EM tomography and 3D reconstruction analysis.

(c) A 1.6 nm-thick slice through the tomographic volume of the RGC showing an

autophagosome with a degenerating mitochondrial fragment inside. A mitochondrion is nearby

with a strand of ER positioned in between the autophagosome and mitochondrion. ER is

typically involved with mitochondrial fission. The double membrane of the autophagosome is

only partially visualized and so is indicated by white arrowheads on opposite sides. For

comparison, the double membrane of the mitochondrial periphery is indicated by black

arrowheads on opposite sides to emphasize that the spacing between individual autophagosomal

membranes is about the same as the spacing between individual mitochondrial peripheral

membranes, as it should be. Labels: A, autophagosome; M, mitochondrion; PM, RGC plasma

membrane; ER, endoplasmic reticulum strand. Scale = 200 nm. (d) Another slice through the

RGC volume showing a small stretch of clearly separated components of the autophagosomal

double membrane (white arrowheads). For comparison, the spacing of the mitochondrial double

membrane is indicated by the black arrowheads. (e) A third slice through the RGC volume that

best shows the double membrane of the mitochondrial fragment (white arrowheads) inside the

autophagosome. Perhaps the reason that less than half the mitochondrial double membrane is

seen is because it was being degraded at the time the tissue was fixed. The spacing between the

individual membranes of the fragment is the same as the spacing between the individual

membranes of the mitochondrion (black arrowheads). (f) Side view with the mitophagosome

membrane made transparent to see the engulfed mitochondrion. (g) Side view with the outer

mitochondrial membrane made transparent to see the 2 fragments of the inner boundary

membrane (IBM) and the one crista that remain. (h) Top view showing the crista and right-hand

portion of the IBM inside the mitochondrion, which is inside the mitophagosome. (i and j) An

oblique view with the mitophagosome and outer mitochondrial membranes made transparent to

see the crista and two IBM fragments. Note that mitochondria in the AKAP1$^{-/-}$ RGC soma

demonstrate autophagosome/mitophagosome formation. Mean $\pm$ SD; $n = 3$ or 4 (a); *$P < 0.05$

and **$P < 0.01$ (two-tailed unpaired Student's t-test). Scale bar: 500 nm

Fig. 7 OXPHOS dysfunction and oxidative stress in AKAP1$^{-/-}$ retinas. (a) Western blot

analyses for OXPHOS Cxs in the retinas of AKAP1$^{-/-}$ and WT mice. (b) Western blot analyses

for SOD2 in the retinas of AKAP1$^{-/-}$ and WT mice. (c) Representative images from

immunohistochemical analyses for SOD2 (green, arrowheads) co-labeled with Brn3a (red,

arrowheads) in the retina. Note that AKAP1$^{-/-}$ retina shows increases of SOD2 protein

expression in the inner retinal layer compared with WT retina. Blue color indicates nucleus. (d)

Quantitative analysis for fluorescent intensity showed a significant increase of SOD2

immunoreactivity in the retina of AKAP1$^{-/-}$ mice. GCL, ganglion cell layer; IPL, inner plexiform

layer; INL, inner nuclear layer; OPL, outer plexiform layer; ONL, outer nuclear layer.

Mean $\pm$ SD; $n = 4$ or 10 (a), $n = 4$ (b) and $n = 10$ (d). *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$

(two-tailed unpaired Student's t-test). Scale bar: 20 μm.

Fig. 8 Akt inactivation and Bim/Bax activation in AKAP1$^{-/-}$ retinas. (a) Western blot

analyses for Akt, phospho-Akt Ser473 and phospho-Akt Thr308 in the retinas of AKAP1$^{-/-}$ and

WT mice. (b) Representative images from immunohistochemical analyses for Akt (green,

arrowheads) co-labeled with Brn3a (red, arrowheads) in the retina. Note that AKAP1$^{-/-}$ retina

show increases of Akt protein expression in the INL and GCL compared with WT retina. Blue

color indicates nucleus. (c) Quantitative analysis for fluorescent intensity showed a significant

increase of Akt immunoreactivity in the retina of AKAP1$^{-/-}$ mice. (d) Western blot analyses for

Bim and Bax in the retinas of AKAP1$^{-/-}$ and WT mice. (e) Representative images from

immunohistochemical analyses for Bax (green, arrowheads) co-labeled with RBPMS (red,

arrowheads) in the retina. Note that AKAP1$^{-/-}$ retina show increases of Bax protein expression in

the GCL compared with WT retina. Blue color indicates nucleus. (f) Quantitative analysis for

fluorescent intensity showed a significant increase of Bax immunoreactivity in the retina of

AKAP1$^{-/-}$ mice. GCL, ganglion cell layer; IPL, inner plexiform layer; INL, inner nuclear layer;

OPL, outer plexiform layer; ONL, outer nuclear layer. Mean $\pm$ SD; $n = 3$ or 4 (a and d) and $n =$

10 (c and f). *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$ (two-tailed unpaired Student's t-test).

Scale bar: 20 μm.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

a

WT   AKAP1⁻/⁻

CxV
CxIII
CxIV

CxII

CxI

Actin

- 100
- 70
- 55
- 40
- 35
- 25
- 15
- 55
- 40

☐ *WT*   ■ *AKAP1⁻/⁻*

Fold

2.0
1.5
1.0
0.5

CNT   CxI   CxII   CxIII   CxIV   CxV

b

WT   AKAP1⁻/⁻

SOD2

Actin

- 25
- 15
- 55
- 40

Fold

2
1

SOD2

☐ *WT*
■ *AKAP1⁻/⁻*

c

SOD2   Brn3a   Nucleus   Merged

WT

ONL
OPL
INL
IPL
GCL

IPL
GCL

AKAP1⁻/⁻

ONL
OPL
INL
IPL
GCL

IPL
GCL

d

Intensity of total immunofluorescence (10⁶ a.u.)

SOD2

☐ *WT*
■ *AKAP1⁻/⁻*

Figure 8.