Resource conservation manifests in the genetic code

Liat Shenhav[1,2,*] and David Zeevi[1,*,^]

[1]Center for Studies in Physics and Biology, Rockefeller University, New York, NY, USA

[2]Department of Computer Science, University of California Los Angeles, Los Angeles, CA, USA

[*]denotes equal contribution

[^]dzeevi@rockefeller.edu

Abstract

Nutrient limitation is a strong selective force, driving competition for resources. However, much is

unknown about how selective pressures resulting from nutrient limitation shape microbial coding

sequences. Here, we study this 'resource-driven' selection using metagenomic and single-cell data of

marine microbes, alongside environmental measurements. We show that a significant portion of the

selection exerted on microbes is explained by the environment and is strongly associated with nitrogen

availability. We further demonstrate that this resource conservation optimization is encoded in the structure

of the standard genetic code, providing robustness against mutations that increase carbon and nitrogen

incorporation into protein sequences. Overall, we demonstrate that nutrient conservation exerts a significant

selective pressure on coding sequences and may have even contributed to the evolution of the genetic code.

Introduction

Nitrogen and carbon are major limiting factors in many ecosystems, with recent studies linking their availability to core genomic properties (*1, 2*). In low-nitrogen environments there is a lower incorporation of nitrogen-rich side-chains into proteins, a strong A+T bias in nucleotide sequences, and smaller genome sizes (*3–5*). An opposite trend was shown for carbon (*2*) and indeed nitrogen and carbon concentrations are typically inversely correlated (*6*). These studies suggest a purifying selective pressure associated with resource conservation, which we term 'resource-driven' selection.

This resource-driven selection postulates that DNA mutations which result in excess incorporation of nutrients such as nitrogen and carbon into proteins, are disfavored. However, not all mutations have the same effect on protein sequences due to constraints imposed by the structure of the genetic code. The standard genetic code, common to virtually all of life on earth, is known to be highly efficient in mitigating deleterious effects of mistranslation errors and point mutations (*7*), specifically those leading to radical changes in amino acids. This error minimization is prominent among theories regarding the origin of the genetic code (*8–11*), which propose that the code had evolved through selection to minimize potential adverse effects of mutations on protein structure and function (*12–14*). To quantify code optimality, some theories provide structurally-informed amino acid metrics based on hydropathy and stereochemistry [e.g., the polar requirement scale (*11*) and hydropathy index (*15*)]. To our knowledge, an optimization of nutrient conservation in the genetic code has not been studied thus far.

Here, we use publicly available metagenomic and single-cell data from global marine habitats, coupled with their corresponding measurements of environmental conditions, to comprehensively characterize and quantify the selective pressure resulting from nutrient limitation. We estimate selection in 746 marine samples (**Fig. S1A**) and quantify its 'resource-driven' component, which we show to be a ubiquitous purifying selective force. We further demonstrate that the structure of the genetic code, known to minimize the effects of DNA mutations on the structure and function of proteins, also minimizes nutrient expenditure given a random mutation. We show that this optimization for resource conservation is orthogonal to known

mechanisms of error minimization in protein structure and function encoded in the genetic code. This

analysis exposes a hitherto unknown facet of the genetic code suggesting resource conservation as an

important contributor to its evolution.

Results

Widespread purifying selection in the marine environment

To comprehensively characterize how coding sequences of marine microbes are affected by resource

availability, we devised a computational pipeline that calculates selection metrics from marine

metagenomic samples (**Fig. S1B**). We downloaded 746 samples from Tara oceans (*16*) (n=136),

bioGEOTRACES (*17*) (n=480), and the Hawaii Ocean- (HOT; n=68); and Bermuda Atlantic- (BATS;

n=62) Time-series (*17*) [**Fig. S1A**; (*18*)]. We aligned these reads to the Ocean Microbiome Reference Gene

Catalog (OM-RGC) (*16*), a database of genes from marine environments that is accompanied by functional

information. We then searched for single nucleotide polymorphisms (SNPs) in genes that had sufficient

high-quality coverage [**Fig. S1B**; (*18*)]. Overall, we found 71,921,864 high-confidence SNPs (*18*), in a total

of 1,590,843 genes.

To quantify purifying selection on different gene functions, we annotated genes from the OM-RGC

database using either KEGG orthology [KO; (*19*)] or eggNOG orthologous groups [OG; (*18*, *20*)]. Using

called SNPs, we calculated for each orthologous group and in each sample, the ratio of nonsynonymous

polymorphisms which lead to a change in the coded amino acid (pN), to synonymous polymorphisms,

which maintain it (pS) [pN/pS; (*18*)]. Across all samples, we found pN/pS ratios to be close to zero with an

average of 0.074 in OGs (CI [0.072, 0.075]) and 0.079 in KOs (CI [0.077, 0.080]; **Fig. 1A**), indicating

widespread purifying selection in the marine environment. To corroborate the validity of calculating

selection metrics from metagenomic samples, we compared nonsynonymous mutations leading to

"conservative" amino acid substitutions to those leading to "radical" substitutions and found conservative

mutations to be significantly more common (**Supplementary text**).

Significant resource-driven selection across marine microbial genes

Based on recent studies (*2–5*), we hypothesized that nutrient availability is a central driver of this purifying

selection. We sought to quantify the environmental contribution to this selective pressure. We considered

environmental measurements taken alongside each sample, including depth, water temperature and salinity,

and the concentrations of nitrate, nitrite, oxygen, phosphate and silicate [**Fig. S2A-H**; (*18*)].

These environmental measurements presented consistent correlation patterns with the pN/pS of many

KEGG and eggNOG orthologs (**Fig. S3**). However, as they are also highly correlated with each other (**Fig.**

S2I), we cannot accurately estimate their individual effects. We therefore sought to estimate the fraction of

variance in pN/pS explained by the environment while accounting for these covariations. We used a linear

mixed model (LMM) with variance components (*18*), which allows to estimate the fraction of variance in

pN/pS ratios (dependent variable) that is explained by the environment (random effect) while controlling

for the complex correlation structure between the environmental parameters. We term the fraction of

variance in pN/pS explained by resource availability 'environmental explained variance' [EEV; (*18*)].

Across both KEGG and eggNOG orthologs we found that a significant fraction of the variance in pN/pS

can be attributed to the environment (**Fig. 1B,C, S4A,B**; Mann-Whitney *U* test $P<10^{-16}$), with nitrate being

more strongly correlated with pN/pS ratios than any other environmental parameter (**Fig. S5**; Kolmogorov-

Smirnov test; $P<10^{-30}$ for all comparisons). Examining typical DNA mutations and amino acid substitutions

in nitrate-rich versus nitrate-poor environments, we find that environmental nitrate is associated with

specific changes to both DNA and protein sequences, favoring lower nitrogen incorporation into protein

sequences when nitrate is scarce (**Supplementary text**).

This association between environmental measurements and the magnitude of purifying selection is

significant even after controlling for potential confounders such as time or effective population size (**Fig.**

S4C,D; Mann-Whitney U test $P<10^{-16}$; **Supplementary text**) as well as in specific environmental niches

(**Fig. S6**; Mann-Whitney U test $P<10^{-20}$; **Supplementary text**). Additionally, these results were replicated

using benchmarking data of assembled genomes from uncultivated single cells from three dominant

lineages of the surface ocean [SAR-11, SAR-86 and Prochlorococcus; **Fig. S4E-G; Supplementary text**;

(*21*)]. These validations demonstrate that the association between selective pressure and environmental

conditions is robust to both data type and selection metric and is not confounded by population properties

and clade-specific metabolism.

Figure 1. Analysis of pN/pS reveals resource-driven selection. (A) Histogram of pN/pS ratios for eggNOG

orthologous groups [OG (20); top] and KEGG orthologs [KO (19); bottom] across all marine samples.

(B) Box plot (line, median; box, IQR; whiskers, 5th and 95th percentiles) of the variance of eggNOG OG

pN/pS that was explained by the environment in a linear mixed model [LMM;(18)] as compared to the same

data with shuffled labels. (C) Same as B, for KEGG KO pN/pS. P, Wilcoxon signed-rank test. (D) Box plot

of variance in pN/pS explained by the environment in the 100 lowest and highest expressed KEGG KOs. P,

Mann-Whitney U test.

Environmental association is stronger in resource-consuming genes

With nitrate being the environmental factor most strongly associated with pN/pS, we sought to provide a population-level perspective to the theory and experiments demonstrating selection against mutations that increase the nitrogen requirements of cells, especially in nitrogen-limited conditions (*3*, *5*, *22*). This implies stronger purifying selection in highly expressed genes (*3*, *4*), where one DNA mutation could translate to thousands of proteins, each consuming more resources (illustrated in **Fig. S7**). We used an additional expression dataset for marine microbial genes (*23*) to rank KEGG orthologs by their mean expression (*18*) and found that the 100 most highly expressed KEGG KOs had a significantly higher EEV as compared to the 100 least expressed ones (**Fig. 1D, S8**; Mann-Whitney U test $P<10^{-9}$). We replicated these results using single-cell data pertaining to specific bacterioplankton lineages (Mann-Whitney U test $P<10^{-7}$). Additionally, we found that extracellular protein-coding genes, where resources excreted from the cell cannot be recycled, had significantly higher EEV as compared to other gene groups [**Fig. S9**; $P<0.05$; (*18*)]. This higher EEV in resource-consuming genes further strengthens our results regarding the breadth of resource-driven selection.

Resource-conservation as an optimization mechanism in the genetic code

We observed selection against DNA mutations which result in excess incorporation of nutrients, such as nitrogen and carbon, into proteins. However, not all mutations have the same effect on protein sequences due to constraints imposed by the structure of the genetic code. Indeed, the genetic code was shown to minimize the impact of point mutations on protein structure and function (*12–14*). We hypothesized that the genetic code also minimizes the impact of point mutations on nutrient incorporation into proteins. Specifically, that the genetic code acts as a buffer between the DNA level, where mutations occur, and the protein level, where resource-driven selection is exerted, such that the expected cost of a random DNA mutation, in terms of added resources to proteins, is minimized.

To rigorously test this hypothesis, we first defined, for each element *e* (e.g., carbon, nitrogen), a function quantifying the cost of a single mutation as the added number of amino acid atoms resulting from it. For example, a missense mutation from codon CCA to CGA results in an amino acid substitution from proline to arginine, with an increase of one carbon and three nitrogen atoms, setting the nitrogen cost of such mutation to 3 and the carbon cost to 1 (**Fig. 2A**).

Figure 2. Resource-conservation is facilitated by the genetic code. *(A) Nitrogen (left), carbon (center) and oxygen (right) content of different amino acids depicted along their positions in the standard genetic code. (B) Histograms of the expected random mutation cost (ERMC), in 1,000,000 random permutations of the genetic code for nitrogen (left, blue), carbon (center, black) and oxygen (right, red). Black-yellow bar marks the ERMC of the standard genetic code, ERMC(Vs), for each of the elements.*

To estimate the cost of random mutations across the entire genetic code, we calculated, per element (e.g., nitrogen), the Expected Random Mutation Cost (ERMC):

$$ERMC_e(V) = \sum_{v,v' \in V_S} P(v)P(mut(v,v'))c_e^+(v,v'),$$

Where V is a genetic code; P(v) the abundance of codon v, here calculated from all marine samples; P(mut(v,v')) the probability of mutation from codon v to codon v', set to be the relative abundance of the single nucleotide mutation driving this codon change (e.g., for mutation from GCA to CCA, we use the

abundance of G-to-C transversions); and the cost function $c_e^+(v,v')$ the number of atoms of element e added

post-mutation (*18*). For the standard genetic code, and codon abundances and mutation rates calculated

from marine microbes, we report an ERMC of 0.440, 0.158 and 0.163 for carbon, nitrogen and oxygen,

respectively, corresponding to an average increase of this number of atoms per random mutation (**Fig. 2B**).

To check if the standard genetic code, along with codon abundances and mutation rates, is indeed robust to

resource-consuming mutations, we compared it to other hypothetical codes. We simulated alternative

genetic codes by randomizing the first and second position in all codons one million times, to create a null

distribution of ERMC (*18*). We found that the standard genetic code, common to most life forms, is

parsimonious in terms of carbon and nitrogen utilization, given a random mutation. This is marked by a

significantly low ERMC for nitrogen (**Fig. 2B**; $ERMC_N$ P=0.0142) and carbon ($ERMC_C$ P=0.0119), but not

oxygen ($ERMC_O$ P=0.8704).

We compared the extent of robustness against an addition of carbon or nitrogen to well-studied structure

and function error minimization mechanisms (*12–14*). We calculated ERMC for changes in hydropathy

(*15*) and polar requirement [PR; (*11*)], both structurally-informed amino acid properties used to determine

code error minimization (*18*). We found that these optimization mechanisms are of a similar magnitude to

those shown here for nitrogen and carbon [**Fig. S10C,D**; $ERMC_{hyd}$ P=0.0152; $ERMC_{PR}$ P=0.0137; (*18*)].

We next sought to confirm that optimization for carbon and nitrogen is not confounded by these known

optimization mechanisms. To this end, we devised a hierarchical model which examined the subset of

genetic codes (out of 1M hypothetical codes tested) that have a lower ERMC than the standard code for PR

or hydropathy, and tested whether this subset is also optimized for nitrogen or carbon (*18*). If nutrient

optimization exists separately from structural optimization, we expect the standard code to be optimized

for carbon and nitrogen even as compared to this subset, featuring significantly lower $ERMC_N$ and $ERMC_C$.

We find that out of 15,223 hypothetical codes that have $ERMC_{hyd}$ lower than the standard code, only 270

have lower $ERMC_N$ (**Fig. 3, S10E**; P=0.019) and only 249 have lower $ERMC_C$ (**Fig. 3, S10E**; P=0.0208).

Similarly, out of the 13,729 hypothetical codes that have $ERMC_{PR}$ lower than the standard code, only 83

have lower $ERMC_N$ (**Fig. S10F**; P=0.0058) and only 442 have lower $ERMC_C$ (**Fig. S10F**; P=0.037). This

is in contrast with the observed overlap between hydropathy and PR: out of the 15,223 hypothetical codes

that have $ERMC_{hyd}$ lower than the standard code, 6,736 have lower $ERMC_{PR}$ (**Fig. S10E**; P=0.4425). These

results indicate that the detected carbon and nitrogen optimization is not confounded by previously reported

optimization properties such as hydropathy and PR.

Figure 3. Optimization for carbon and nitrogen is not confounded by hydropathy. *(A) Hydropathy of*
different amino acids depicted along their positions in the standard genetic code. (B; top) As in Fig. 2B,
for hydropathy; (B; bottom) histograms of ERMC for nitrogen (left) and carbon (right) for the subset of
hypothetical genetic codes with ERMC lower than the standard code for hydropathy. P, permutation test.

Remarkably, only 128 out of 1M randomized genetic codes were better than the standard code in

conservation of nitrogen and carbon together (**Fig. S10G,H**; $ERMC_{CN}$ P=1.28×10^{-4}). We note that this

number is significantly smaller than the number of hypothetical codes expected to have both a lower

$ERMC_C$ and $ERMC_N$ (Chi-square test of independence P=0.0013; **Table S1**). This is possibly driven by a

small overlap between the positions of high-nitrogen and high-carbon amino acids. This property of the

standard code potentially enables concurrent optimization for both carbon and nitrogen. These results

highlight a new optimization principle of the genetic code that is of similar magnitude- and independent of-

previously proposed principles.

The genetic code facilitates resource conservation across kingdoms

To show that the resource-robustness of the genetic code was not limited to our dataset and analytic

approach, we calculated the ERMC of 187 strains of genera *Prochlorococcus* and *Synechococcus*. We

computed codon abundances and mutation rates using published protein-coding sequences (*1*) and the

accepted transition:transversion rate of 2:1 (*18*, *24*). We corroborated significant conservation of carbon,

nitrogen, and both elements combined (**Fig. S11A**; $ERMC_C$ mean P=0.013, P=0.020; $ERMC_N$ mean

P=0.049, P=0.032; $ERMC_{CN}$ P=0.0004, P=0.0007 for *Prochlorococcus* and *Synechococcus*, respectively).

To explore whether this nutrient conservation optimization mechanism of the standard code extends across

organisms, we performed a similar calculation using codon abundances from 39 organisms across all

domains of life, including all human protein-coding sequences, and a range of transition:transversion rates

(*18*). Similarly to marine microbes, we find that the standard code is optimized in terms of resource

utilization for all tested organisms, marked by a significantly low ERMC for nitrogen and carbon combined,

across all transition:transversion rates (P<0.01, **Fig. 4**). Moreover, we find significant optimization even in

the theoretical case where all codon abundances are the same (**Fig. S11B**). The codon abundances of a great

majority of organisms also demonstrate significantly low ERMC for nitrogen (**Fig. S11C**) and carbon (**Fig.

S11D**), for a wide range of transition:transversion rates. These results indicate that resource optimization

in the genetic code transcends taxonomy, codon choices, and mutation rates.

Figure 4. ***The genetic code is optimized for resource conservation across organisms.*** *Heat map of ERMC$_{CN}$ P-values across 39 organisms and 11 transition:transversion rates. Organisms in each group are ordered by the GC content of their coding sequences.*

Resource conservation may bias codon usage

We hypothesized that variation in the third codon position for a single amino acid and may also be biased

due to the differential cost of a random mutation for each codon. We therefore focused on codon usage of

the amino acid threonine. We note that a C-to-G transversion in the second position for codons ACT and

ACC yields serine (AGT and AGC, respectively), but the same mutation for codons ACA and ACG yields

arginine (AGA and AGG; **Fig. 5A**, inset). Arginine has higher carbon and nitrogen content than serine. We

thus hypothesized that for a cell to conserve nutrients in case of a random mutation, codons ACT and ACC

should have a higher abundance than codons ACA and ACG, respectively, given a known GC-bias that is

a genomic quantity. To test this hypothesis, we examine codon usage in 187 *Prochlorococcus* and

Synechococcus strains, and show a significantly higher use of ACT as compared to ACA (**Fig. 5A**;

Wilcoxon signed-rank test $P<10^{-20}$) and ACC as compared to ACG (**Fig. 5A**; Wilcoxon signed-rank test

$P<10^{-20}$). Similarly, isoleucine codon ATT has higher abundance as compared to ATA (**Fig. 5A**; Wilcoxon

signed-rank test $P<10^{-20}$). These results demonstrate that resource conservation may guide codon usage and

thereby affect not only protein sequence but also cellular translation efficiency.

Structural principles drive optimization in the genetic code

We next examined the organizing principles underlying code resource optimization. We observed that

codons of the nitrogen-rich amino acids histidine, glutamine, asparagine, lysine and arginine span only two

nucleotides in their first position and two in their second position. We define this organization to be a

'square' arrangement, and hypothesize that it amplifies nitrogen conservation [**Fig. 5B**; (*18*)]. Specifically,

in the square arrangement, some codons require at least two mutations to increase the number of nitrogen

atoms (e.g., those coding for alanine and valine). This is in contrast to other hypothetical arrangements,

including a 'diagonal' one in which nitrogen-rich amino acid codons span all possible nucleotides in the

first and second positions [**Fig. 5B**; (*18*)]. We suggest that this arrangement would be nutrient-wasteful, as

in such arrangements a single mutation in either the first or second codon position could increase the

nitrogen content of a protein. To test this, we generated 10,000 hypothetical codes, with 220 arrangements

happening to embody a square structure, and 127 a diagonal one. We found that, when compared to all

other possible arrangements, square arrangements present a significantly lower $ERMC_N$ while diagonal

arrangements exhibit a significantly higher $ERMC_N$ (**Fig. 5B**; Mann-Whitney U test $P<10^{-10}$ for both). This

result demonstrates that resource optimization in the standard code is driven by structural principles,

perhaps underlying the optimization observed across kingdoms.

Figure 5. Structural properties and codon usage bias underlying optimality in the genetic code. *(A) Violin*

plot of codon usage among 187 species of Prochlorococcus and Synechococcus showing significant

preference of threonine codons ACT and ACC as compared to ACA and ACG, and of isoleucine codon ATT

as compared to ACA. P, Wilcoxon signed-rank test. (B) Box plot (line, median; box, IQR; whiskers, 5th and

95th percentiles) of $ERMC_N$ of square arrangements (left) and diagonal arrangements [right, (18)], as

compared to all other arrangements (center) out of 10,000 hypothetical arrangements. P, Mann-Whitney

U test.

Discussion

In this work, we characterize and quantify the selective forces exerted by nutrient availability on protein

coding genes in marine environments. We provide a data-driven, population-level perspective and show

that resource-driven selection is a ubiquitous force. We further show that a significant portion of DNA

mutations may not result in increased nutrient incorporation into protein sequences, due to the structure of

the standard genetic code. We provide evidence that the genetic code, shared among most lifeforms, along

with organismal codon frequencies, is robust against the incorporation of additional atoms of nitrogen and

carbon given a random mutation. We quantitatively demonstrate that this nutrient conservation is not

confounded by known error minimization mechanisms of the genetic code.

In light of these results, we hypothesize that resource-driven selection is equally exerted on all parts of the

protein-coding gene. This sets it apart from selection to maintain the structural integrity of a protein or the

function of its active site, which occur predominantly in structurally important regions (*25*). Thus,

accounting for resource-driven selection may improve the identification of alternative translation start sites,

alternatively spliced introns, or readthrough stop codons, as intermittently translated regions of the protein

may be under weaker resource-driven selection as compared to constitutively translated ones (*26*).

We note that in addition to resource-driven selection there are other forces associated with the environment,

driving both selective and neutral mechanisms. Genomic GC-content is associated with exposure to UV

radiation (*27*) and nitrogen availability (*2, 28*) yet nonsynonymous mutations, especially in codons

encoding structurally and functionally important protein positions, are likely to be less affected by GC-bias.

Furthermore, codon usage bias was also shown to be associated with the environment (*29*).

We show that the genetic code optimizes nutrient conservation, calling upon an examination of code

evolution theories. Hypotheses regarding the origin of the genetic code include stereochemical affinity

between a codon or anticodon and their amino acid (*11, 30*); a frozen accident theory (*31*); a co-evolution

of the genetic code with the emergence of amino acid-producing biosynthetic pathways (*9*); and an early

fixation of an optimal genetic code, suggesting selection for error minimization (*10, 32*). Our results are in

line with the latter theory and suggest that the genetic code may have also been optimized for nutrient

conservation. This implies that the genetic code can be viewed as a buffer between the evolutionary forces

of mutation and selection, the former occurring in DNA sequences and the latter predominantly in proteins.

In the case of nutrient conservation, many DNA mutations do not result in the incorporation of additional

nutrients into proteins, and are thus not selected against, which may allow more 'freedom' to explore the

mutation-space. Notwithstanding, while optimized for resource conservation, the genetic code is also near-

immutable. This is evident in heterotrophic eukaryotes, which while being net-nitrogen producers, still

harbor a nitrogen-conservative code.

Since the genetic code is common to virtually all organisms, we can assume that it was already fixed in its

current form, or one very similar, in the last universal common ancestor. Thus, if the genetic code evolved

to optimize resource conservation, it may reflect selective pressures in the beginning of life on earth. An

organism harboring a nitrogen- and carbon-efficient genetic code would have had a selective advantage

over its peers, especially in the absence of fully evolved DNA mutation repair mechanisms.

References

1. P. M. Berube, A. Rasmussen, R. Braakman, R. Stepanauskas, S. W. Chisholm, Emergence of trait
variability through the lens of nitrogen assimilation in Prochlorococcus. *Elife*. **8** (2019),
doi:10.7554/eLife.41043.

2. D. R. Mende *et al.*, Environmental drivers of a microbial genomic transition zone in the ocean's
interior. *Nature Microbiology*. **2** (2017), pp. 1367–1373.

3. J. J. Grzymski, A. M. Dussaq, The significance of nitrogen cost minimization in proteomes of
marine microorganisms. *ISME J.* **6**, 71–80 (2012).

4. H. Akashi, T. Gojobori, Metabolic efficiency and amino acid composition in the proteomes of
Escherichia coli and Bacillus subtilis. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 3695–3700 (2002).

5. F. L. Hellweger, Y. Huang, H. Luo, Carbon limitation drives GC content evolution of a marine
bacterium in an individual-based genome-scale model. *The ISME Journal*. **12** (2018), pp. 1180–
1187.

6. E. R. M. Druffel, S. Griffin, A. I. Coppola, B. D. Walker, Radiocarbon in dissolved organic carbon
of the Atlantic Ocean. *Geophys. Res. Lett.* **43**, 5279–5286 (2016).

7. R. T. Hinegardner, J. Engelberg, RATIONALE FOR A UNIVERSAL GENETIC CODE. *Science*.
142, 1083–1085 (1963).

8. G. Gamow, Possible Relation between Deoxyribonucleic Acid and Protein Structures. *Nature*. **173**,
318–318 (1954).

9. J. T. Wong, A co-evolution theory of the genetic code. *Proc. Natl. Acad. Sci. U. S. A.* **72**, 1909–1912
(1975).

10. S. J. Freeland, L. D. Hurst, The genetic code is one in a million. *J. Mol. Evol.* **47**, 238–248 (1998).

11. C. R. Woese, D. H. Dugre, W. C. Saxinger, S. A. Dugre, The molecular basis for the genetic code.
Proc. Natl. Acad. Sci. U. S. A. **55**, 966–974 (1966).

12. S. E. Massey, A neutral origin for error minimization in the genetic code. *J. Mol. Evol.* **67**, 510–516
(2008).

13. A. S. Novozhilov, E. V. Koonin, Exceptional error minimization in putative primordial genetic
codes. *Biol. Direct*. **4**, 44 (2009).

14. N. Torabi, H. Goodarzi, H. Shateri Najafabadi, The case for an error minimizing set of coding amino
acids. *J. Theor. Biol.* **244**, 737–744 (2007).

15. D. Haig, L. D. Hurst, A quantitative measure of error minimization in the genetic code. *J. Mol. Evol.*
33, 412–417 (1991).

16. S. Sunagawa *et al.*, Structure and function of the global ocean microbiome. *Science*. **348**, 1261359
(2015).

17. S. J. Biller *et al.*, Marine microbial metagenomes sampled across space and time. *Sci Data*. **5**,

180176 (2018).

18.   Materials and methods are available as supplementary materials at the Science website.

19.   M. Kanehisa, S. Goto, KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* **28**,
27–30 (2000).

20.   J. Huerta-Cepas *et al.*, Fast Genome-Wide Functional Annotation through Orthology Assignment by
eggNOG-Mapper. *Mol. Biol. Evol.* **34**, 2115–2122 (2017).

21.   M. G. Pachiadaki *et al.*, Charting the Complexity of the Marine Microbiome through Single-Cell
Genomics. *Cell.* **179** (2019), pp. 1623–1635.e11.

22.   A. Dufresne, L. Garczarek, F. Partensky, Accelerated evolution associated with genome reduction in
a free-living prokaryote. *Genome Biol.* **6**, R14 (2005).

23.   B. C. Kolody *et al.*, Diel transcriptional response of a California Current plankton microbiome to
light, low iron, and enduring viral infection. *ISME J.* (2019), doi:10.1038/s41396-019-0472-2.

24.   E. Urbach, D. J. Scanlan, D. L. Distel, J. B. Waterbury, S. W. Chisholm, Rapid Diversification of
Marine Picophytoplankton with Dissimilar Light-Harvesting Structures Inferred from Sequences of
Prochlorococcus and Synechococcus (Cyanobacteria). *Journal of Molecular Evolution.* **46** (1998),
pp. 188–201.

25.   J. Liu, Y. Zhang, X. Lei, Z. Zhang, Natural selection of protein structural and functional properties: a
SNP perspective. *Genome Biology.* **9** (2008), p. R69.

26.   R. W. Read *et al.*, Nitrogen cost minimization is promoted by structural changes in the transcriptome
of N-deprived Prochlorococcus cells. *ISME J.* **11**, 2267–2278 (2017).

27.   C. E. Singer, B. N. Ames, Sunlight ultraviolet and bacterial DNA base ratios. *Science.* **170**, 822–825
(1970).

28.   C. E. A. Mcewan, D. Gatherer, N. R. Mcewan, Nitrogen-fixing aerobic bacteria have higher genomic
GC content than non-fixing species within the same genus. *Hereditas.* **128**, 173–178 (1998).

29.   M. Botzman, H. Margalit, Variation in global codon usage bias among prokaryotic organisms is
associated with their lifestyles. *Genome Biol.* **12**, R109 (2011).

30.   M. Yarus, J. J. Widmann, R. Knight, RNA–Amino Acid Binding: A Stereochemical Era for the
Genetic Code. *J. Mol. Evol.* **69**, 406 (2009).

31.   F. H. Crick, The origin of the genetic code. *J. Mol. Biol.* **38**, 367–379 (1968).

32.   S. J. Freeland, R. D. Knight, L. F. Landweber, L. D. Hurst, Early fixation of an optimal genetic code.
Mol. Biol. Evol. **17**, 511–518 (2000).

33.   D. Zeevi *et al.*, Structural variation in the gut microbiome associates with host health. *Nature.* **568**,
43–48 (2019).

34.   1000 Genomes Project Consortium, G. R. Abecasis *et al.*, A map of human genome variation from
population-scale sequencing. *Nature.* **467**, 1061–1073 (2010).

35. J. Huerta-Cepas *et al.*, eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated
orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.* **47**, D309–D314
(2019).

36. T. Seemann, Prokka: rapid prokaryotic genome annotation. Bio-Informatics 30: 2068--2069 (2014).

37. A. Bairoch, R. Apweiler, The SWISS-PROT protein sequence database and its supplement TrEMBL
in 2000. *Nucleic Acids Res.* **28**, 45–48 (2000).

38. J. H. Ward, Hierarchical Grouping to Optimize an Objective Function. *J. Am. Stat. Assoc.* **58**, 236–
244 (1963).

39. K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7: improvements
in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

40. A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics*. **30**, 1312–1313 (2014).

41. N. Goldman, Z. Yang, A codon-based model of nucleotide substitution for protein-coding DNA
sequences. *Mol. Biol. Evol.* **11**, 725–736 (1994).

42. R. Schlitzer *et al.*,The GEOTRACES Intermediate Data Product 2017. *Chem. Geol.* **493**, 210–223
(2018).

43. J. Honaker, G. King, M. Blackwell, Others, Amelia II: A program for missing data. *J. Stat. Softw.*
45, 1–47 (2011).

44. A. L. Price, N. A. Zaitlen, D. Reich, N. Patterson, New approaches to population stratification in
genome-wide association studies. *Nat. Rev. Genet.* **11**, 459–463 (2010).

45. S. Schloissnig *et al.*, Genomic variation landscape of the human gut microbiome. *Nature*. **493**, 45–50
(2013).

46. M. R. E. Symonds, S. P. Blomberg, A Primer on Phylogenetic Generalised Least Squares. *Modern
Phylogenetic Comparative Methods and Their Application in Evolutionary Biology* (2014), pp. 105–
130.

47. M. Westoby, M. R. Leishman, J. M. Lord, On misinterpreting thephylogenetic correction'. *J. Ecol.*
83, 531–534 (1995).

48. F. E. Rheindt, T. U. Grafe, Rapidly evolving traits and the comparative method: how important is
testing for phylogenetic signal? *Evol. Ecol.* (2004) (available at http://www.evolutionary-
ecology.com/abstracts/v06/1507.html).

49. J. Wang, L. Raskin, D. C. Samuels, Y. Shyr, Y. Guo, Genome measures used for quality control are
dependent on gene function and ancestry. *Bioinformatics*. **31**, 318–323 (2015).

50. J. Athey *et al.*, A new and updated resource for codon usage tables. *BMC Bioinformatics*. **18**, 391
(2017).

51. J. Zhang, Rates of conservative and radical nonsynonymous nucleotide substitutions in mammalian
nuclear genes. *J. Mol. Evol.* **50**, 56–68 (2000).

52. S. J. Giovannoni *et al.*, Genome streamlining in a cosmopolitan oceanic bacterium. *Science*. **309**,
1242–1245 (2005).

53. F. L. Hellweger, E. van Sebille, N. D. Fredrick, Biogeographic patterns in ocean microbes emerge in
a neutral agent-based model. *Science*. **345**, 1346–1349 (2014).

54. B. K. Swan *et al.*, Prevalent genome streamlining and latitudinal divergence of planktonic bacteria in
the surface ocean. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 11463–11468 (2013).

55. R. Milo, What is the total number of protein molecules per cell volume? A call to rethink some
published values. *Bioessays*. **35**, 1050–1055 (2013).

56. Y. Taniguchi, Quantifying E. coli Proteome and Transcriptome with Single-molecule Sensitivity in
Single Cells. *Seibutsu Butsuri*. **51** (2011), pp. 136–137.

57. T. R. Anderson, P. J. le B. Williams, A one-dimensional model of dissolved organic carbon cycling
in the water column incorporating combined biological-photochemical decomposition. *Global
Biogeochemical Cycles*. **13** (1999), pp. 337–349.

58. B. Charlesworth, Effective population size and patterns of molecular evolution and variation. *Nat.
Rev. Genet.* **10**, 195–205 (2009).

59. M. Lynch, J. S. Conery, The evolutionary fate and consequences of duplicate genes. *Science*. **290**
(2000), pp. 1151–1155.

60. A. L. Hughes, R. Friedman, P. Rivailler, J. O. French, Synonymous and nonsynonymous
polymorphisms versus divergences in bacterial genomes. *Mol. Biol. Evol.* **25**, 2199–2209 (2008).

61. J. H. McDonald, M. Kreitman, Adaptive protein evolution at the Adh locus in Drosophila. *Nature*.
351, 652–654 (1991).

Acknowledgements: We thank Tal Korem and Nimrod D. Rubinstein for discussions, suggestions and
help in analyses. We thank Jean-Pierre Eckmann, Otto X. Cordero, Itzhak Mizrahi and Niv Antonovsky for
their suggestions. **Funding:** DZ was funded by the James S. McDonnell Foundation. **Authors**
contributions: LS and DZ conceived the project, performed analyses and wrote this manuscript.
Competing interests: The authors declare no competing interests. **Data and materials availability:** All
data is available in the manuscript, supplementary materials and referenced datasets. Code is available at:
https://github.com/zeevilab/resource-conservation

Supplementary Materials

Materials and Methods

Supplementary text

Fig S1 – S16

Table S1 – S2

References (33-61)

Materials and Methods

<u>Marine microbiome samples</u>

Marine samples collected with Tara oceans (*16*), bioGEOTRACES (*17*), the Hawaii Ocean Timeseries (HOT) and the Bermuda Atlantic Timeseries (BATS) (*17*) were downloaded from ENA with accessions ENA:PRJEB1787 (TARA oceans prokaryotic fraction), ENA:PRJNA385854 (bioGEOTRACES) and ENA:PRJNA385855 (HOT/BATS), each sample with a minimum of 5 million reads. For all samples for which a size parameter was available, we selected the 0.22μm+ size category (typically between 0.22 and 5μm). This includes TARA ocean samples (prokaryote size) and data pertaining to expression (*22*). In samples from the bioGEOTRACES, HOT and BATS, we used the available size fraction, larger than 0.2μm.

<u>Mapping of Illumina reads to reference gene sequences</u>

Samples were mapped to nucleotide sequences from the Ocean Microbiome Reference Gene Catalog (OM-RGC) (*16*) using bowtie2 with parameters --sensitive -a 20 --quiet -p 8 and saved as a bam file using the 'samtools view' command. As gene sequences are relatively short, reads from both ends of the metagenomic sequencing samples were mapped separately, and reunited prior to variant calling.

<u>Determining metagenomic read assignment probability</u>

We determined the probability of assignment of metagenomic reads to marine microbial genes using the Iterative Coverage-based Read-assignment Algorithm (ICRA) (*32*) with parameters max_mismatch=12, consider_lengths=True, epsilon=1e-6, max_iterations=30, min_bins=4, max_bins=100, min_reads=10, dense_region_coverage=60, length_minimum=300, length_maximum=2e5, use_theta=False. To prevent spurious mapping, alignments were considered for downstream analysis only if the probability of alignment was higher than 0.9.

<u>Variant calling</u>

Alignments from both ends of all sequencing runs pertaining to the same sample were united using the samtools cat command and sorted using the samtools sort command, with default parameters. To facilitate variant calling in tractable timescales, each filtered, untied and sorted bam file was split into chunks, each encompassing 10,000 reference sequences (out of about 40 million reference sequences). For each such batch of reference sequences, we called variants across all samples using the following command: bcftools mpileup --threads 4 -a FORMAT/AD -Q 15 -L 1000 -d 100000 -m 2 -f <OM-RGC

fasta> <bam filenames> | bcftools call --threads 4 -Ov -mv -o <output vcf>, where <OM-RGC fasta> is

the fasta file of OM-RGC nucleotide sequences, <bam filenames> are the filenames of all bam files

pertaining to the reference sequence chunk in question, and <output vcf> is the output vcf file pertaining

to that same chunk.

To ensure that sequencing errors do not skew our results, we considered single nucleotide variants as

SNPs for downstream analysis only if they had a GATK quality score of at least 30, and an allele

frequency of at least 1% (*33*). While the GATK quality requirements filter out sequencing errors

randomly distributed across the genes, the 1% criterion eliminates random sequencing errors that

accumulate in the same position when depth of coverage reaches very high numbers.

For a sample mapped to a reference gene to be considered for downstream analysis, we demanded that at

least 60% of SNPs called along the length of the reference gene for that sample would be supported by at

least 4 reads, thereby enabling accurate calculation of pN/pS ratios. For a gene to be considered for

downstream analysis, we demanded for that gene to have at least one SNP common to 20 or more

samples.

Calculation of pN/pS in single genes

While comparing SNP patterns across samples, it is instrumental to avoid biases due to differences in

coverage. We therefore downsampled the read coverage (depth) to the minimum across all samples, for

each position in a sample that was supported by more than 4 reads. For positions that had minimum

support of fewer than 4 reads, no subsampling was performed. Subsampling was performed by drawing

from a multinomial distribution, with n trials and variant probabilities p, where n was set to the calculated

minimum depth and p set to the relative abundance of each variant in a given sample.

The ratio of non-synonymous and synonymous polymorphisms was calculated by considering all called

SNPs in every gene. First, we calculate a consensus sequence for each gene by taking, for each SNP

position, the variant that was overall more common across all samples (after the subsampling performed

above). We counted, for each gene, the number of non-synonymous and synonymous sites across the

consensus sequence. For each SNP position in each sample, we counted the number of synonymous and

nonsynonymous substitutions. As more than one variant can exist in a single sample, we considered the

relative abundance of synonymous to nonsynonymous substitutions dictated by the different variants. For

example, if the reference codon was CAC, coding for histidine, one variant, a C-to-G transversion in the

third position abundant at 50%, led to a nonsynonymous mutation that resulted in glutamine (CAG) while

another variant in the same sample and the same position was a synonymous C-to-T transition, we

counted 0.5 synonymous substitutions and 0.5 nonsynonymous substitutions. We followed by calculating

the pN/pS ratio:

488
$$pN/pS = \frac{nonsynonymous\ substitutions}{nonsynonymous\ sites} \bigg/ \frac{synonymous\ substitutions}{synonymous\ sites}$$

<u>Aggregation of calculated metrics using KEGG and eggNOG orthologies</u>

Functional assignments to KEGG KOs and eggNOG OGs for all OM-RGC genes were computed using

eggNOG-mapper v2 based on the eggNOG v5.0 database (*19, 34*). For each functional assignment in each

sample, all OM-RGC genes annotated with the same functional assignment were concatenated and treated

as one long genomic sequence per the calculation of pN/pS ratios. To reduce noise in pN/pS calculation,

we considered only KOs and OGs that had at least 5 genes per sample, in at least 50 samples.

<u>Calculating nonsynonymous mutations leading to radical and conservative amino acid replacements</u>

We calculated the ratio of nonsynonymous mutations leading to radical amino acid replacements as

compared to those leading to conservative replacements using two known amino acid properties: (1) the

polar requirement scale (PR) and (2) amino-acid hydropathy index, and multiple thresholds of "radical"

and "conservative" (**Table S2**). For each KEGG ortholog and each radical/conservative threshold, we

compared the proportion of radical nonsynonymous substitutions (out of all possible radical sites) to the

proportion of conservative ones (out of all conservative sites) across all samples using a Mann-Whitney *U*

test. P-values were adjusted to control for multiple hypotheses.

Formally, our statistical analysis postulates:

505   H_0: proportion of radical changes $\geq$ proportion of conservative changes

506   H_1: proportion of radical changes $<$ proportion of conservative changes

For each radical/conservative threshold, we calculated the proportion of KEGG orthologs in which we

reject this null hypothesis and conducted a permutation test. We permuted the sample labels of each

radical/conservative threshold across 10,000 iterations, where in each iteration we calculated the

proportion of KEGG orthologs in which we reject the null hypothesis. The P-value, per

radical/conservative threshold, is defined to be the proportion of iterations in which the ratio of KEGG

orthologs in which we reject the null hypothesis is greater than in the original data.

To examine whether highly expressed genes have fewer radical amino acid substitutions as compared to

lowly expressed genes, we used data from Pachiadaki et al. (*22*) to rank KEGG orthologs by their mean

expression.

Benchmarking data of assembled genomes

Benchmarking data was employed in the form of uncultivated single cells of dominant and representative

lineages of the surface ocean (SAR-11, SAR-86 and Prochlorococcus) from different parts of the world.

We used genomes varying at the strain level (*20*), with potentially minimal batch effects, as those have

been analyzed in a single laboratory.

Calculation of dN/dS in single genes

For every gene ortholog in each one of the lineages dN/dS was calculated as follows:

1. We extracted all gene sequences pertaining to each ortholog from all strains within each

clade. We determined orthologous genes using the annotation that was published

alongside the single-cell data. For each clade, we chose the annotation type that had the

highest coverage of genes: prokka (*35*) for Prochlorococcus, and Swissprot (*36*) for SAR-

11 and SAR-86.

2. To account for the possibility that gene sequences may be misannotated resulting in the

grouping of more than one orthologous group under the same name, we performed

additional curation by clustering genes using Ward hierarchical clustering (*37*) on protein

sequences using normalized edit distance (Levenshtein) as a dissimilarity metric. We cut

off clusters that had dissimilarity lower than 30%, to ensure structural similarity.

3. For each of the resulting clusters that had more than 20 members (i.e., genes

corresponding to strains within the clade) we generated a multiple sequence alignment

(MSA) of protein sequences using MAFFT (*38*) , with the following command:

mafft --quiet --maxiterate 1000 --localpair <input_fasta> > <output_msa>

Following that, we computed a codon alignment using the generated MSA and nucleotide

sequences, with the biopython AlignIO package.

4. Using the MSA, we computed a phylogenetic tree of the sequences using raxml (*39*)

(raxmlHPC command line version) with model GTRGAMMA; rooted it using the -f I

modifier; and computed the ancestral sequence using the -f A modifier.

5. For each leaf node (strain ortholog) in the phylogenetic tree of gene sequences, we

calculated dN/dS with respect to the tree root, using the biopython cal_dn_ds function

with the "ML" model (Goldman and Yang 1994)(*40*) and default parameters.

6. To quantify purifying selection on different gene functions, we annotated genes from the
OM-RGC database using KEGG orthology [KO; (*18*)] and aggregated the dN/dS across
genes from the same KO using a geometric mean.

<u>Environmental variables</u>
For each sample, we compiled measurements pertaining to the following environmental parameters:
Depth [m], Nitrate [µmol/kg], Nitrate+Nitrite [µmol/kg], Oxygen [µmol/kg], Phosphate [µmol/kg],
Silicate [µmol/kg], Temperature [°C] and Salinity [g/kg].
Tara oceans metadata was downloaded from PANGAEA (https://doi.pangaea.de/10.1594/) with accession
numbers PANGAEA.875575 (Nutrients) and PANGAEA.875576 (Watercolumn sensor), and recorded
median values for all the above nutrients were extracted. Tara nutrient concentrations were given as
[µmol/l]. Conversion to [µmol/kg] was done by dividing the measured concentration by the measured
specific gravity for the same sample.
bioGEOTRACES metadata was compiled from CTD sensor data and discrete sample data from the
GEOTRACES intermediate data product v.2 (*41*). HOT metadata was downloaded from the ftp server of
the University of Hawai'i at Manoa (ftp://ftp.soest.hawaii.edu/hot/) and BATS metadata was downloaded
from the Bermuda Institute of Ocean Sciences (http://bats.bios.edu/bats-data/). As
GEOTRACES/HOT/BATS ocean water samples are not linked to specific biological samples as is the
case with Tara oceans samples, we considered only water samples from the exact same geographic
location, within a day from biological sample collection time, and within 5% difference in depth of
collection, and chose the closest sample in terms of time and depth of collection. As all these
measurements of environmental conditions are highly correlated with each other (**Fig. S2I**), we utilized
this correlation structure to impute missing values using the EM algorithm (*42*).

<u>Linear mixed models</u>
We used a linear mixed model (LMM) with variance components, commonly used in population genetics
(*43*), which allows to estimate the fraction of variance in pN/pS ratios (dependent variable) that is
explained by the environment (random effect) while controlling for the complex correlation structure
between the environmental parameters, in a data-driven manner.

<u>*Generative model.*</u> Consider a collection of M_i, $where\ i \in \{1,2\}$ features (i.e., $M_{1,2}$ number of KEGG and
eggNOG orthologs respectively), each measured across N samples. We get as input an $(M_i \times N)$ matrix
O^i, where $O_{mn}{}^i$ is the pN/pS of ortholog m in sample n. Let $y^i{}_m = (O_{m1}{}^i, \ldots, O_{mN}{}^i)$ be a $N \times 1$ vector

representing the pN/pS in ortholog m, according to grouping i, across N samples (e.g., pN/pS in KO

K02274 across N samples). Let W be a $(N \times q)$ normalized matrix of environmental measurements.

This included the depth of the sample, water temperature and salinity, as well as concentration of the key

molecules nitrate, nitrite, oxygen, phosphate and silicate.

With these notations, we assume the following generative linear model:

$$y^i{}_m = W u_m + \epsilon_m, \qquad (1)$$

581 where u_m and ϵ_m are independent random variables distributed as $u_m \sim N\left(0, \sigma^2{}_{u_m} I\right)$ and $\epsilon_m \sim$

582 $N(0, \sigma^2_{\epsilon_m} I)$. The parameters of the model are $\sigma^2{}_{u_m}$ and $\sigma^2_{\epsilon_m}$.

584 An equivalent mathematical representation of model (1) is given by

$$y^i{}_m \sim N(0, \sigma^2_K K + \sigma^2_{\epsilon_m} I), \qquad (2)$$

586 where $\sigma^2_K = M_i \sigma^2{}_{u_m}$, $K = \frac{1}{M_i} W W^T$. We will refer to K as the environmental kinship matrix, which

587 represents the similarity, in terms of environmental covariates, between every pair of samples across

588 grouping i (i.e., represent the correlation structure to the data).

590 *Environmental explained variance (EEV).* The explained variance of a specific feature $y^i{}_m$ by the

591 environmental measurements

$$\chi^i{}_m = \frac{\sigma^2_K}{\sigma^2_K + \sigma^2_{\epsilon_m}}. \qquad (3)$$

594 Estimating effective population size

595 To approximate effective population size (N_e) we use fourfold degenerate synonymous diversity.

596 Calculation was adapted from Schloissnig et al.(*44*):

$$\pi_{ffsyn}(S, G) = \frac{1}{|G_{ffsyn}|} \sum_{i=1}^{|G_{ffsyn}|} \sum_{B_1 \in \{A,C,G,T\}} \sum_{B_2 \in \{A,C,G,T\} \setminus B_1} \frac{x_{i,B_1}}{c_i} \times \frac{x_{i,B_2}}{c_i - 1}, \qquad (4)$$

where S is the sample, G is the gene of interest, $|G_{ffsyn}|$ is the number of fourfold degenerate sites, x_{i,B_j} is

the number of times nucleotide B_j was seen in the fourfold degenerate position i, and c_i is the coverage of

position i.

<u>Controlling for the effects of time and effective population size</u>

To control for the effects of time and effective population size, we added these two potential confounders

to our LMM as fixed effects. We formulate our LMM as follows:

$$y^i_m = \beta_{0_i} + \beta_{1_i} pS_i + \beta_{2_i} SD_i + Wu_m + \epsilon_m \ , \qquad (5)$$

where β_{0_i} is the intercept, pS (as a proxy for time) is the first fixed effect with β_{1_i} as its coefficient, SD_i

(as a proxy for effective population size) is the second fixed effect with β_{2_i} as its coefficient. W is a

normalized matrix of environmental factors, u_m and ϵ_m are independent random variables distributed as

$u_m \sim N(0, \sigma^2_{u^n_m} I)$ and $\epsilon_m \sim N(0, \sigma^2_{\epsilon_m} I)$. The parameters of the model are β_{0_i} , β_{1_i} , β_{2_i} $\sigma^2_{u_m}$ and

$\sigma^2_{\epsilon_m}$.

<u>LMM on samples from the epipelagic zone</u>

To demonstrate that the association of pN/pS with the environment is not confounded by depth, we reran

the LMM (eq. 5) only on samples from the epipelagic zone (up to 76m, median depth across 746

samples).

<u>Linear model for dN/dS rates</u>

We modeled the benchmarking data by constructing a linear model for dN/dS (the dependent variable)

with the environmental factors as independent variables:

$$dN/dS_m = \beta_{0_m} + Wu_m + \epsilon_m, \qquad (6)$$

where β_{0_m} is the intercept, W is a normalized matrix of environmental factors and u_m is a vector of

environmental coefficients. $\epsilon_m \sim N(0, \sigma^2_{\epsilon_m} I)$. $m \in \{1, \ldots, M\}$, where M is the number of genes tested.

To account for the effect of time as potential confounder we constructed the following linear model for

dN/dS (the dependent variable) with the environmental factors and dS (as a proxy for time) as

independent variables.

624
$$dN/dS_m = \beta_{0_m} + \beta_{1_m} dS_m + Wu_m + \epsilon_{2_m}, (7)$$

where β_{0_m} is the intercept, dS (as a proxy for time) is a fixed effect with β_{1_m} as its coefficient, W is a

normalized matrix of environmental factors and u_m is a vector of environmental coefficients that were

available with these samples (phosphate, depth, nitrate + nitrite). $\epsilon_{2_m} \sim N(0, \sigma^2_{\epsilon_{2_m}} I)$. $m \in$

$\{1, \dots, M\}$, where M is the number of orthologs tested. In this model we only used depth, NO_2+NO_3 and

phosphate, as these were the environmental factors available for the benchmarking data.

To account for potential underlying structure in terms of phylogeny, we applied phylogenetic generalized

least squares (45) (PGLS) regression in addition to ordinary least squares (OLS) regression to model

dN/dS. Phylogenetic regression models are used when there is an underlying phylogenetic clustering

structure of the observations. In such case, samples may be lowly or highly phylogenetically diverged

leading to residuals that are not *i.i.d.*, which can affect the fit of the model. On the other hand, the caveat

of PGLS is an underestimation of the R^2 as compared to linear models (46, 47). In our data PGLS and

OLS essentially gave the same results and we therefore only report the latter.

KEGG KO expression data as a ranking metric

Using expression data from 4,092 KEGG KOs collected by Kolody et al. (22), we ranked the KO genes in

our marine samples in the following way. We first represent the expression data in relative abundance

space (normalize each sample by its read counts). Next, for each focal KO i, where $i \in \{1, \dots, I\}$, we

sum across its different instances within each sample. The input is an $m \times n$ matrix, where m is the

number of different instances of KO i, and n is the number of samples. The output is a vector of length

n: $(x_1, \dots, x_n)$. Finally, we average the expression levels $(x_1, \dots, x_n)$ across samples: $\frac{1}{n}\sum_{j=1}^{n} x_j$ and rank

the KOs based on the calculated average expression. We limited the scope of our analysis to samples

collected only in small fraction filters ($0.22 \mu m$).

Determination of extracellular genes

To determine extracellular gene groups, we searched the eggNOG v.5 OG database for the words

'secreted' or both words 'extracellular' and 'protein' in their description. We demanded that the words

'autoinducer', 'expression', 'role' and 'hypothetical' are not in the description to prevent instances where

(a) the OG in question describes a hypothetical protein; and (b) where the OG produces a secreted particle

but is not secreted by itself, as is the case with autoinducer producing genes. To ensure robust pN/pS

calculations, descriptions encompassing 10 OGs or more were assigned a group name, while descriptions

encompassing less than 10 OGs were grouped together.

<u>Calculation of mutation flux in divergent nitrate concentrations</u>

To calculate the differential rate of nutrient incorporation between environments, we compare, for each

pair of codons a and b, the ratio of mutations that result in a becoming b and the ratio of reverse

mutations, that result in b becoming a. We term this quantity mutation flux. To prevent simplex-related

confounding effects, mutation flux is estimated using the log-odds ratio between the two (e.g.,

$log\,(p(AAA \to AAC)/p(AAC \to AAA))$. We created a matrix $H^{(U)}$ as follows: consider a set U of all

genes for which SNP measurements exist and a subset $T \subseteq U$ of this set across K samples. Let $G^{(T)}{}_j =$

(V, E) be a codon graph for subset T and sample j, where $v \in V$ is a codon (e.g., CUU coding for

Alanine) and $(v, v') \in E$ if and only if v and v' are one mutation apart (e.g., CUU for Alanine and CAU

for Histidine). Let $w^{(T)}{}_j : (v, v') \to [0,1]$ be a weight function where

664
$$w^{(T)}{}_j(v, v') = \frac{Number\ of\ mutations\ turning\ v\ to\ v\prime\ in\ subset\ T\ and\ sample\ j}{Number\ of\ occurrences\ of\ codon\ v\ in\ subset\ T\ and\ sample\ j}.$$

Let $H^{(T)}$ be a matrix of dimension $(|E| \times K)$, where $H^{(T)}{}_{(v,v\prime),j} = w^{(T)}{}_j(v, v')$. We next sum-normalized

H per each sample and compared the codon mutation frequencies between the 40 lowest- and 40 highest-

nitrate samples. Despite significant differences in codon mutation frequencies between low-nitrate and

high-nitrate samples, some of the difference could be driven by the simplex properties of the sum-

normalized codon mutation frequencies, and some could be attributed to the different rates of

synonymous mutations between the high- and low-nitrate groups, which, combined with simplex

properties, may affect observed nonsynonymous mutation rates. To address these simplex properties, we

employed a centered log-ratio (CLR) normalization on H. The CLR transformation is a mapping, per

codon composition, from the simplex to a Euclidean vector subspace. This log transforms each value and

then centers them around zero as given below:

675
$$CLR(V) = [log\frac{v_1}{g(v)},\dots,log\frac{v_D}{g(v)}] = log(v) - log(g(v))$$

where g(v) is the geometric mean of all of the codons.

To address differences in rates of the different types of mutations, for each mutation (v, v') in the CLR

normalized matrix $H'^{(U)}$ we calculated the log odds ratio between the mutation and its reverse mutation.

Namely, we computed mutation flux matrix $F^{(U)}$ where $F^{(U)}{}_{(v,v\prime),j} = H^{(U)}{}_{(v,v\prime),j} - H^{(U)}{}_{(v\prime,v),j}$. We

compared differences in codon mutation flux between low- and high nitrate samples using the Mann-

Whitney U test.

<u>Calculation of expected random mutation cost per genetic code</u>

Let V be a genetic code with a set $V_s \subset V$ of stop codons. Let $P(v)$ be the abundance of codon $v \in V$ in a

sample and $P(mut(v, v'))$ the probability of a single mutation from codon v to v'. Let $c_e: V \times V \to$

$[0,1]$ be a cost function for element e, where:

$$c_e(v, v') = \# \ of \ atoms \ of \ e \ in \ v' \ - \ \# \ of \ atoms \ of \ e \ in \ v.$$

With these notations, we define the expected cost of genetic code V for element e as follows:

$$E[C_e(V)] \ = \sum_{v,v' \in V} P(v) P(mut(v, v')) c_e(v, v'),$$

and the ERMC cost as:

$$ERMC_e(V) \ = \ \sum_{v,v' \in V} P(v) P(mut(v, v')) c_e^+(v, v'),$$

where $c^+_e = max(0, c_e)$. When testing the random mutation cost on hydrophobicity or PR we consider

the absolute difference in the amino-acid hydropathy index or PR value pre- and post-mutation.

We estimate $ERMC_e(V)$ as follows:

1.   We define $P(v)$ as the median abundance of all codons v' coding for the same amino acid as v.

2.   We calculate mutation rates in sites that are under minimal selection. To this end, we estimate

696            $P(mut(v, v'))$ by calculating, from fourfold-degenerate synonymous mutation sites the average

697   abundance of each single nucleotide mutation (e.g. A to C) across all genes in which there are

698   called SNPs in all ocean samples, excluding stop codons. We then estimate $P(mut(v, v'))$ using

699   the relative abundances of all pairs of single nucleotide mutations. We estimate $P(mut(v, v'))$ for

700   *Prochlorococcus* and *Synechococcus* using published transition:transversion rates (*1, 48*).

3.   We calculate c using information on the amino acids which each codon codes for.

To compute a P-value, we generate a null distribution by calculating $ERMC_e(V)$ for hypothetical genetic

codes. We randomize the first and second position of all codons, while maintaining that the two sets of

first and second positions in which the stop codons reside are separated by a single transition mutation.

<u>Confounding effects between cost functions for the structure of the genetic code</u>

To confirm that our elemental cost function is not confounded by traditional properties of amino acids

such as the polar requirement (PR) and hydropathy index (*11*, *15*), we calculated the expected random

mutation cost (ERMC), per hypothetical genetic code, using these common cost functions across 1

million codes. To this end, we randomized the first and second position of all codons, while maintaining

that the two sets of first and second positions in which the stop codons reside are separated by a single

transition mutation. We next calculated a contingency table for each pair of cost functions for both

nitrogen and carbon (i.e., $ERMC_N(V){:}ERMC_{PR}(V)$, $ERMC_N(V){:}ERMC_{hyd}(V)$, $(V){:}ERMC_{PR}(V)$,

$ERMC_C(V){:}ERMC_{hyd}(V)$). We assign each code to one of four bins in the following way: (1) surpassing

the standard code in both cost functions (e.g., nitrogen and PR), (2) surpassing the standard code only in

element *e* cost (e.g. only nitrogen), (3) surpassing the standard genetic only in the traditional cost function

(e.g., PR), (4) not surpassing the standard code in either. We applied the Chi-square test of independence

with two degrees of freedom to each contingency table.

Finally, we examined the subset of genetic codes (out of the 1M hypothetical codes) that have a lower

ERMC than the standard code in terms of PR ($ERMC_{PR}$) or hydropathy ($ERMC_{hyd}$), and tested whether

this subset is also optimized for nitrogen and carbon. To this end, we created a null model in which we

compute EMRC only accounting for hydropathy or PR robustness and an expanded, nested model that

adds minimization of nitrogen and carbon utilization. Under the null, we expect that in the subset of codes

optimizing for hydropathy and PR, nitrogen and carbon utilization will also be minimized. Under the

alternative, if such nutrient optimization indeed exists separately from structural optimization, we expect

that nitrogen carbon and utilization will not be optimized. Exact P-values are defined to be the fraction of

shuffled codes with ERMC smaller or equal to the standard code (one for each element - carbon and

nitrogen) in this subset of codes that have a lower ERMC than the standard code in terms of PR

($ERMC_{PR}$) or hydropathy ($ERMC_{hyd}$).

<u>Determination of 'square' and 'diagonal' arrangements of nitrogen in the genetic code</u>

We define a 'square' arrangement of the codons coding for nitrogen-rich amino acids histidine,

glutamine, asparagine, lysine and arginine as one where their codons span only two nucleotides in the first

position and two nucleotides in the second position. In the standard code, these amino acids are coded by

CAN, CGN, AAN, AGR, following a square configuration. In contrast, a 'diagonal' arrangement of the

codons coding for these amino acids is one where they span all possible nucleotides in the first position

and all possible nucleotides in the second position. For example, a genetic code where TTY codes for

histidine, TTR for glutamine, CCN and AAR for arginine, GGY for asparagine and GGR for lysine

constitutes a 'diagonal' arrangement of nitrogen amino acids. In each of the alternative genetic codes we

generated, we tested whether either of these conditions hold and, if so, designated the code as 'square' or

'diagonal' accordingly.

Prochlorococcus and *Synechococcus* genomic data and mutation rates

We downloaded *Prochlorococcus* and *Synechococcus* protein-coding gene sequences (where available)

from the Joint Genome Institute (https://genome.jgi.doe.gov/portal/) following accession numbers

published by Berube et al. (*1*). To estimate codon relative abundance $P(v)$, we counted and sum-

normalized codons in all protein-coding genes for each strain. To estimate codon mutation rate

$P(mut(v, v'))$, we used the published transition:transversion rate of 2:1 for *Prochlorococcus* and

Synechococcus (*23*).

Multiple taxa ERMC calculation

To calculate ERMC for 39 taxa across multiple transition:transversion rates, we downloaded codon usage

and GC-content data collected by Athey et al. (*49*). We used codon usage counts to estimate $P(v)$ and 11

transition:transversion rates (1:5, 1:4, 1:3, 1:2, 2:3, 1:1, 3:2, 2:1, 3:1, 4:1, 5:1) to estimate $P(mut(v, v'))$.

Supplementary Text

To corroborate the validity of calculating selection metrics from metagenomic samples, we sought to
replicate a known property of mutations in amino acids, namely that mutations resulting in "conservative"
amino acid replacements are more common than those resulting in "radical" ones (*50*). To this end, we
calculated the ratio of nonsynonymous mutations leading to radical amino acid replacements and
compared them to those leading to conservative replacements. To best define "radical" and
"conservative" nonsynonymous substitutions, we used two known amino acid properties: (1) the Polar
Requirement (PR) scale and (2) amino-acid hydropathy index (**Table S2**). Since the definition of radical
and conservative may be subjective, we considered multiple thresholds as radical and conservative for
each property. For example, a radical substitution could be one that results in a difference of more than 3
units on the hydropathy index whereas a "conservative" one results in just one or less (**Table S2**;
hydropathy case 4).
We then calculated, for each KEGG ortholog and for each radical/conservative threshold, the proportion
of radical nonsynonymous substitutions (out of all possible radical sites) as compared to the proportion of
conservative ones across samples using a Mann-Whitney U test. P-values were adjusted to control for
multiple hypotheses.
Formally, our statistical analysis postulates:

$$H_0 : proportion\ of\ radical\ changes\ \geq proportion\ of\ conservative\ changes$$
$$H_1 : proportion\ of\ radical\ changes\ < proportion\ of\ conservative\ changes$$

Next we calculated, for each radical/conservative threshold, the proportion of KEGG orthologs in which
we reject the null hypothesis (i.e., the proportion of KOs in which nonsynonymous mutations leading to
conservative amino acid replacements are significantly more common than nonsynonymous mutations
leading to radical amino acid replacements) and conducted a permutation test. We permuted the sample
labels of each radical/conservative threshold across 10,000 iterations, where in each iteration we
calculated the proportion of KEGG orthologs in which we reject the null hypothesis. The P-value, per
radical/conservative threshold, is defined to be the proportion of iterations in which the ratio of KEGG
orthologs in which we reject the null hypothesis is higher as compared to the original data. We find that

these comparisons are significant for all cases, regardless of the amino acid property in question or the

implemented threshold and that this is the case for most KOs (P<0.0001 for all comparisons; **Table S2**).

Finally, we examined whether highly expressed genes have fewer radical amino acid replacements as

compared to lowly expressed genes. To this end, we used an additional expression dataset for marine

microbial gene expression to rank KEGG orthologs by their mean expression (similar to the approach

implemented in **Fig. 1D** (*22*)). We find that KEGG orthologs which exhibit significantly more

conservative mutations than radical ones have significantly higher expression as compared to the

alternative case (Wilcoxon signed-rank test $P<10^{-5}$). These results demonstrate that metagenomic

sequences are adequate for analyzing selective pressures.

Supplementary Note 2. Resource-driven selection drives specific amino acid changes

Our results indicate that mutations leading to an addition of resources would be more common in nutrient-

rich environments. This is supported by previous observations regarding genomic and proteomic changes

associated with environmental concentrations of nitrate (*2, 5, 21, 51–53*). We thus sought to quantify how

resource conservation manifests in molecular changes to DNA and protein sequences. Mutations in

nitrogen-rich and, typically, carbon-poor environments were shown to drive an increase in genomic GC

content (which increases nitrogen requirements), as well as by higher nitrogen and lower carbon in

protein sequences.

Using SNP derived from metagenomic data, we corroborate previous findings (*5, 21*), and show that

mutations leading to higher GC content and those leading to higher nitrogen proteins are more prevalent

in high-nitrate waters as compared to low-nitrate (**Fig. S12A,B**). We note that mutations leading to higher

GC content (higher nitrogen DNA) often result in higher nitrogen proteins (*2, 3*). We further deconvolve

these two types of mutations and find that the ratio of mutations that lead to higher nitrogen DNA but do

not change protein nitrogen-content increases with environmental nitrate (R=0.333; **Fig. S12C**).

Additionally, the ratio of mutations that increase protein nitrogen but do not change DNA nitrogen-

content is even more strongly correlated with nitrate concentrations (R=0.473; **Fig. S12D**). We

hypothesize that this stronger association of the environment with protein sequences can be attributed to

the higher copy number of protein as compared to DNA and RNA, often in the order of 10^2-10^5 protein

copies per RNA molecule (*54, 55*) leading to higher cost per gene, and thus to stronger selective pressure.

Our results suggest a differential effect of mutations on nutrient incorporation in DNA as compared to

protein sequences.

To study amino acid substitutions typical to low- and high-nitrate environments, we examined mutations

leading to changes in protein nutrient content by comparing codon mutation frequencies in low- and high-

nitrate samples, after accounting for simplex-related confounders (Methods). We found significant

differences in codon mutation frequencies between environments (**Fig. S13A-C**). We next sought to

examine the typical change in nutrient consumption in varying nitrate concentrations. We therefore

defined mutation flux as the ratio between a codon mutation and its reverse, thereby measuring the typical

flux of nutrients; and estimated it using the log-odds ratio between the two (e.g., $log\ (p(AAA \rightarrow$

$AAC)/p(AAC \rightarrow AAA))$. Notably, across all the mutations significantly more prevalent in samples from

high nitrate environments (Methods), averaged across amino acids, we find a total increase in nitrogen

(**Fig. S13D**; 18 N atoms summed across all significant amino acid changes, P=0.0082), decrease in carbon

(**Fig S13D**; -37 atoms, P=0.0165), decrease in sulfur (-6 atoms, P=0.009), a decrease in molecular weight

(-508.91 g/mol, P=0.0193) and a non-significant decrease in oxygen (-6 atoms, P=0.1505). These results

provide additional evidence that the higher the nitrate concentrations are, the weaker the selection against

mutations leading to higher nitrogen incorporation into protein sequences. While nitrate concentrations

increase with depth (**Fig. S2I**), labile dissolved organic carbon concentrations typically decrease (*56*). We

therefore also show the specific codon mutations reducing carbon incorporation into proteins in nitrogen

rich and typically carbon poor, environments (**Fig. S13D,E**).

<u>Supplementary Note 3. The association between the environment and pN/pS is robust to time, effective</u>

<u>population size and environmental niche</u>

The magnitude of purifying selection, approximated by pN/pS ratios, may increase as a function of both

time and effective population size (*57, 58*). These two mechanisms might also be associated with

environmental conditions and we thus sought to control for them as potential confounders. Time may

affect the magnitude of pN/pS ratios via random genetic drift. Effective population size may affect

selection via competition, as in large populations, any slightly deleterious mutation is rapidly selected

against. We used the ratio of synonymous substitutions (i.e., pS) calculated from our data as an

approximation of dS and representing neutral polymorphism rate, as a proxy for time (*59*). To

approximate effective population size, we relied on population genetic theory suggesting that effective

population size is scaled to diversity at the silent sites when assuming constant mutation rate between

populations. As our data is composed of closely related sequences and hence similar mutation rate, we

used fourfold-degenerate synonymous diversity (SD) as a proxy for the effective population size (*44*)

(Methods).

We first tested whether SD is associated with environmental conditions, and compared the SD of

populations from different oceanic regions with different environmental parameters. We found that pN/pS

was more strongly associated with most environmental parameters than SD (Kolmogorov-Smirnov test

$P < 10^{-20}$ for depth, nitrate, oxygen, phosphate, silicate and temperature; $P < 10^{-5}$ for $NO_2 + NO_3$

concentrations; non-significant for salinity; **Fig. S3**) suggesting that selection on protein sequences may

be more strongly affected by the environment than effective population sizes are. We further found that in

KOs with the highest EEV these differences are the most pronounced (Wilcoxon signed-rank test $P < 10^{-15}$;

Fig. S14).

As LMMs also allow controlling for potential confounders as fixed effects, we next added our

approximations of time and the effective population size to our LMM in order to account for their effect

on the magnitude of pN/pS ratios. Accounting for these potential confounders as fixed effects would

enable a more accurate quantification of the environmental effect. Similar to the results of our original

LMM, across both KEGG and eggNOG orthologs, we found that a significant fraction of the variance in

pN/pS can be attributed to the environment, even after controlling for these confounders (**Fig. S4C,D**;

Mann-Whitney U test $P < 10^{-16}$). We found a high correlation between EEV estimated by the two models

($R^2 = 0.947$ and 0.929 for KEGG and eggNOG, respectively; **Fig. S15**).

To demonstrate that our results are not confounded by depth, the environmental variable with the second

strongest correlation to pN/pS (after nitrate), we recalculated our LMM only on samples from the

epipelagic zone (Methods). Similar to the original results of the LMM, we show significant variance

explained in pN/pS in this environmental niche (**Fig. S6**; Mann-Whitney U test $P < 10^{-20}$). Taken together,

these results demonstrate that the association between the environment and pN/pS is robust to time,

effective population size and environmental niche.

<u>Supplementary Note 4. The association between environmental measurements and the magnitude of</u>

<u>purifying selection is replicated using single cell data and dN/dS</u>

pN/pS, the ratio of nonsynonymous polymorphisms (pN) to synonymous polymorphisms (pS),

approximates dN/dS which quantifies the magnitude and type of selection (purifying, neutral or positive)

exerted on protein-coding sequences across a phylogenetic group (*44*, *60*). While dN/dS calculation

requires genomic sequences at the strain-level that are available for only a small number of samples and

from limited strains, it allows for a more accurate quantification of purifying selection in the taxa

represented in the phylogeny. We therefore sought to confirm our findings using dN/dS. To this end, we

used benchmarking data in the form of assembled genomes from uncultivated single cells from dominant

and representative lineages of the surface ocean (SAR-11, SAR-86 and Prochlorococcus) (*20*), and their

corresponding environmental measurements (Methods).

We calculated dN/dS for KEGG orthologs in each lineage. Given the small number of samples, we

estimated the fraction of variance explained by the environment using a linear model with environmental

covariates (Methods). Across all bacterioplankton lineages comprising the benchmarking data, we found

that a significant portion of the variance in dN/dS ratios calculated from single-cell data is explained by

the environment (SAR-11, SAR-86, Prochlorococcus; Wilcoxon signed-rank test $P<10^{-16}$; **Fig. S4E-G**).

Similar to our original LMM, we note that this association holds, across all three bacterioplankton

lineages, even after controlling for both time as a confounder (Wilcoxon signed-rank test $P<10^{-16}$; **Fig.**

S16) and for phylogeny, using phylogenetic generalized least-square regression (Methods). These results

validate our findings using pN/pS calculated from metagenomic sequences and demonstrate the

association between selective pressure and environmental conditions, even in a single clade. This

indicates that associations of pN/pS or dN/dS with environmental variables are likely not confounded by

organismal properties which may differ in their metabolism across environments, but rather exhibit a

consistent trend even within a single clade.

Fig. S1.

(A) Geographical overview of the samples used in this study. (B) Illustration of our computational

pipeline.

Fig. S2

(A-H) Distribution of measurements taken alongside marine microbial samples for depth (A), nitrate (B), nitrate and nitrite (C), oxygen (D), phosphate (E), salinity (F), silicate (G) and temperature (H). (I) Spearman correlation coefficients between all pairs of environmental measurements across all available samples.

 Fig. S3

 (A-P) Histograms of Spearman correlations between synonymous diversity (red) or pN/pS (blue) and

 environmental variables, for both KEGG KOs and eggNOG OGs. Panels A, C, E, G, I, K, M and O depict

 correlations between OG calculated parameters and depth, nitrate, nitrite and nitrite, oxygen, phosphate,

 salinity, silicate and temperature, respectively. Panels B, D, F, H, J, L, N and P depict correlations

 between KO calculated parameters and depth, nitrate, nitrite and nitrite, oxygen, phosphate, salinity,

 silicate and temperature, respectively. P, one-sided Kolmogorov-Smirnov test.

Fig. S4

(A) Left, box plot (line, median; box, IQR; whiskers, 5th and 95th percentiles) of the variance of eggNOG

OG pN/pS rates that was explained by the environment in a basic linear mixed model (LMM) setting

(Methods) as compared to the same data with shuffled labels; Right, mean variance explained in

909 unshuffled data (red) as compared to a histogram (blue) of mean variance explained in 500 executions

910 with shuffled data. (B) Same as A, for KEGG KO pN/pS. (C) Same as A, for variance in eggNOG OG

911 pN/pS in a LMM corrected for confounding factors of time and effective population size. (D) Same as C,

912 for KEGG KO pN/pS. (E) Box plot (line, median; box, IQR; whiskers, 5th and 95th percentiles) of the

913 variance of KEGG KO dN/dS in genus *Prochlorococcus* that was explained by the environment in a

914 linear model. (F) Same as E, for the SAR-11 clade. (G) Same as E, for the SAR-86 clade. ****, Wilcoxon

915 signed-rank test $P<10^{-16}$.

Fig. S5

Histograms of Spearman correlations between environmental variables (blue) or their inverses (red) and

pN/pS ratios of KEGG KOs as compared to a Spearman correlation between the concentration of nitrate

and pN/pS ratios (green). P, one-sided Kolmogorov-Smirnov test. We note that unmeasured variables

correlated with nitrate, such as labile dissolved organic carbon, may also be correlated with pN/pS.

Fig. S6

Box plot (line, median; box, IQR; whiskers, 5th and 95th percentiles) of the variance of KEGG KO

pN/pS ratios that was explained by the environment in epipelagic samples (up to 76m) as compared to the

same data with shuffled labels.

Fig. S7

Illustration of the potential environmental effect of nitrate on protein-coding sequences. Depicted are two microbes, wild-type (WT; A) and a mutant with a T-to-G transversion in the 14th position of the yjcG gene, leading to a leucine to arginine substitution, thereby consuming three extra nitrogen atoms per protein molecule (B). If environmental nitrate is scarce and growth-limiting, A is likely to have a selective advantage over B, which over multiple generations would translate into an extinction of the B variant. In highly expressed genes such a mutation is likely to have an even stronger effect, leading to a quicker extinction. On the molecular level, this phenomenon would translate to lower pN/pS in nitrate-limited waters as compared to nitrate-rich ones.

Fig. S8

(top, red) 24 highly expressed KOs, exhibiting high correlation between pN/pS and nitrate concentrations.

(bottom, blue) 24 lowly expressed KOs, exhibiting low correlation between pN/pS and nitrate

concentrations. R, Spearman rho.

943 **Fig. S9**

944 Box plot (line, median; box, IQR; whiskers, 5th and 95th percentiles) of variance in pN/pS explained by

945 the environment in extracellular gene groups versus all eggNOG OGs.

Fig. S10

Upper panel: (Top) Nitrogen content (A; blue color scale), carbon content (B; black), the polar requirement (C; purple) and hydropathy (D; green-red) of different amino acids depicted along their positions in the standard genetic code. (Bottom) Histograms of the expected random mutation cost (ERMC), in 1,000,000 random permutations of the genetic code for nitrogen (blue), carbon (black), hydropathy (green) and polar requirements (purple). Lower panel: Histograms of ERMC for nitrogen (blue), carbon (black), hydropathy (green) and polar requirements (purple) for the subset of shuffled genetic codes with ERMC lower than the standard genetic code for (E) hydropathy, (F) the polar requirement, (G) carbon and (H) nitrogen. Vertical black-yellow line, ERMC of the standard genetic code for each property. P, permutation test.

Fig. S11

(A) Box plots (line, median; box, IQR; whiskers, 5th and 95th percentiles) of P-values for the ERMC of the standard genetic code for nitrogen (left), carbon (center) and both (right) across 187 Prochlorococcus (P, blue) and Synechococcus (S, red) strains. (B) Heat map of ERMC P-values for nitrogen, carbon and both, for a theoretical case in which all codons are of the same abundance. (C,D) Same as Fig. 2C for $ERMC_N$ (D) and $ERMC_C$ (E) P-values.

Fig. S12

Scatter plots of nitrate concentrations (x-axis) versus percent of mutations (y-axis) for (A) all AT to GC

mutations; (B) All mutations in coding sequences changing an amino acid with no nitrogen in its side-

chain to one with nitrogen in the side-chain (non-N to N mutations); (C) AT to GC mutations not

changing protein nitrogen; (D) non-N to N mutations not changing GC content. R and P, Spearman

correlation coefficient and P-value; line, linear fit; shaded area, 95% confidence interval.

Fig. S13

(A-C) Histograms of the distribution of Mann-Whitney *U* P-values of codon-to-codon mutations
compared between (A) 40 low-nitrate samples and 40 other low-nitrate samples selected randomly out of
the 80 lowest-nitrate samples; (B) 40 high-nitrate samples and 40 other high-nitrate samples selected
randomly out of the 80 highest-nitrate samples; (C) 40 low-nitrate samples and 40 high-nitrate samples
selected randomly out of the 80 lowest and highest nitrate samples. (D) Depiction of mutation flux
(Methods) common in high versus low environmental nitrate concentrations, affecting amino acid

979   nitrogen (left) and carbon (right) content. Yellow arrows, increase in resource; blue arrows, decrease in

980   resource; arrow thickness corresponds to number of atoms changed by mutation. (E) Same as D, showing

981   all fluxes, regardless of change in resources. Two-headed arrows mark cases where different codons of

982   the same amino acid have opposite mutation fluxes.

983

984 **Fig. S14**

985 Box plots (line, median; box, IQR; whiskers, 10th and 90th percentiles) of P-values (t-test, multiple

986 hypothesis adjustment) between samples with high-N (more than 19.6 μmol/kg) and low-N (0 μmol/kg),

987 for KEGG KOs ordered by their EEV in the LMM, for both pN/pS (blue) and synonymous diversity (red).

988 ****, Wilcoxon signed-rank test P<10^{-15}.

Fig. S15

(A,B) scatter plot of explained variance in a linear mixed model considering only environmental factors as random effects (x-axis) versus a model considering environmental factors as random effects plus dS (as a proxy for time) and synonymous diversity (as a proxy for effective population size) as fixed effects (y-axis) in all (A) KEGG KOs and (B) eggNOG OGs.

Fig. S16

Box plots (line, median; box, IQR; whiskers, 10th and 90th percentiles) of explained variance in dN/dS using linear models considering the environment only (left, red), dS only (middle, blue), or both (right, green). (A) Prochlorococcus, (B) SAR-11 clade and (C) SAR-86 clade. **, Wilcoxon signed-rank test $P<0.01$; ***, $P<10^{-5}$; ****, $P<10^{-16}$.

Table S1.

Contingency tables for each pair of cost functions for both nitrogen (n+) and carbon (c+), compared to PR

(pr) and Hydropathy index (hyd), across 1 million simulated genetic codes. Each code is assigned to one

of four bins: (1) surpassing the standard genetic code in both cost functions (<; <), (2) surpassing the

standard genetic code only in element e cost (<; >=), (3) surpassing the standard genetic code only in the

traditional cost function (>=; <), (4) not surpassing the standard genetic code in neither (>=; >=). Chi-

square test of independence was applied to each contingency table.

Cost function a	Cost function b	Sign a	Sign b	Alternative code counts	Chi-squared P-value
n+	c+	<	<	128	1.32E-03
			>=	14106	
		>=	<	11802	
			>=	973964	
n+	hyd	<	<	270	2.71E-04
			>=	13964	
		>=	<	14953	
			>=	970813	
n+	pr	<	<	83	4.68E-16
			>=	14151	
		>=	<	13646	
			>=	972120	
c+	hyd	<	<	249	4.86E-07
			>=	11681	
		>=	<	14974	
			>=	973096	
c+	pr	<	<	442	4.29E-107
			>=	11488	
		>=	<	13287	
			>=	974783	

1008

1009 **Table S2.**

1010 Comparisons between "radical" and "conservative" nonsynonymous substitutions using the polar

1011 requirement scale and amino-acid hydropathy index and multiple thresholds for each property. P,

1012 permutation test.

	Radical Threshold	Conservative Threshold	Proportion of significant Kos	Permutation test P-value
PR: Case 1	Absolute difference > 1	Absolute difference < 1	0.704	< 0.0001
PR: Case 2	Absolute difference > 2	Absolute difference < 1	0.737	< 0.0001
PR: Case 3	Absolute difference > 2	Absolute difference < 2	0.737	< 0.0001
PR: Case 4	Absolute difference > 3	Absolute difference < 1	0.928	< 0.0001
PR: Case 5	Absolute difference > 3	Absolute difference < 2	0.925	< 0.0001
PR: Case 6	Absolute difference > 3	Absolute difference < 3	0.913	< 0.0001
Hyd: Case 1	Absolute difference > 1	Absolute difference < 1	0.885	< 0.0001
Hyd: Case 2	Absolute difference > 2	Absolute difference < 1	0.880	< 0.0001
Hyd: Case 3	Absolute difference > 2	Absolute difference < 2	0.832	< 0.0001
Hyd: Case 4	Absolute difference > 3	Absolute difference < 1	0.950	< 0.0001
Hyd: Case 5	Absolute difference > 3	Absolute difference < 2	0.934	< 0.0001
Hyd: Case 6	Absolute difference > 3	Absolute difference < 3	0.932	< 0.0001
Hyd: Case 7	pre-mutation < -1 & post-mutation > 1 || pre-mutation > 1 & post-mutation < -1	Otherwise	0.943	< 0.0001
Hyd: Case 8	pre-mutation < -2 & post-mutation > 2 || pre-mutation > 2 & post-mutation < -2	Otherwise	0.968	< 0.0001

1013