Enhancing autophagy by redox regulation extends lifespan in *Drosophila*

Authors

Claudia Lennicke[1,2], Ivana Bjedov[3], Sebastian Grönke[4], Katja E. Menger[5], Andrew M. James[5], Jorge Iván Castillo-Quan[6], Lucie A. G. van Leeuwen[1,2], Andrea Foley[1,2], Marcela Buricova[1,2], Jennifer Adcott[6], Alex Montoya[1,2], Holger B. Kramer[1,2], Pavel V. Shliaha[1,2], Angela Logan[5], Filipe Cabreiro[2,7], Michael P. Murphy[5], Linda Partridge[4,6]*, Helena M. Cochemé[1,2]*

Affiliations

[1] MRC Laboratory of Medical Sciences, Du Cane Road, London W12 0HS, UK.

[2] Institute of Clinical Sciences, Imperial College London, Hammersmith Hospital Campus, Du Cane Road, London W12 0HS, UK.

[3] UCL Cancer Institute, 72 Huntley Street, London WC1E 6DD, UK.

[4] Max Planck Institute for Biology of Ageing, Joseph Stelzmann Strasse 9b, 50931 Cologne, Germany.

[5] MRC Mitochondrial Biology Unit, University of Cambridge, Cambridge Biomedical Campus, Hills Road, Cambridge CB2 0XY, UK.

[6] Institute of Healthy Ageing and GEE, University College London, Gower Street, London WC1E 6BT, UK.

[7] CECAD Research Cluster, University of Cologne, Joseph Stelzmann Strasse 26, 50931 Cologne, Germany.

* correspondence: linda.partridge@ucl.ac.uk; helena.cocheme@lms.mrc.ac.uk

Abstract

Dysregulation of redox homeostasis has been implicated in the ageing process and the pathology of age-related diseases. To study redox signalling by H_2O_2 *in vivo*, we established a redox-shifted model by manipulating levels of the H_2O_2-degrading enzyme catalase in *Drosophila*. Here we report that ubiquitous over-expression of catalase robustly extends lifespan in flies. As anticipated, these flies were strongly resistant to a range of oxidative stress challenges, but interestingly were sensitive to starvation, which could not be explained by differences in levels of energy reserves. This led us to explore the contribution of autophagy, which is an important mechanism for organismal survival in response to starvation. We show that autophagy is essential for the increased lifespan upon catalase up-regulation, as the survival benefits were completely abolished upon global autophagy knock-down. Furthermore, using a specific redox-inactive knock-in mutant, we highlight the *in vivo* role of a key regulatory cysteine residue in Atg4a, which is required for the lifespan extension in our catalase model. Altogether, these findings confirm the redox regulation of autophagy *in vivo* as an important modulator of longevity.

Introduction

Redox signalling is a fundamental process that impacts a diverse range of biological pathways[1-4]. Redox regulation can operate through the selective post-translational modification (PTM) of redox-sensitive cysteine residues in target proteins[1-4]. Cysteines are typically present in a reduced, thiol state *in vivo* (–SH)[5-8], but can undergo oxidation in response to specific stimuli, including levels of reactive oxygen species (ROS) or changes in the ratios of redox cofactors, such as NADPH/NADP$^+$ and glutathione/glutathione disulfide. The nature of this oxidative PTM will vary according to the physiological context, but typically proceeds via a sulfenic acid (–SOH) intermediate[1-4].

The redox reactivity of a given cysteine will depend on a combination of physical and biochemical properties. For instance, the position of the cysteine within the tertiary protein structure will determine its accessibility to pro-oxidants, and ability to interact with other redox-dependent binding partners. Additionally, the cysteine pK_a will be influenced by adjacent charged amino acid residues and the local cellular environment, which governs its protonation state and therefore susceptibility to oxidative PTM[1-4]. Critically, redox signalling is distinct from irreversible oxidative damage and functions as a reversible 'redox switch' to regulate target proteins. Dysregulation of redox homeostasis has long been implicated in the pathophysiology of many age-related diseases, as well as in the ageing process itself, however the underlying mechanisms remain largely unclear[9,10].

H_2O_2 acts as the major effector of redox signalling, both directly and through intracellular thiol redox relays[11,12]. The oxidative PTMs induced by H_2O_2 lead to specific changes in cellular pathways, including many processes implicated in ageing[13,14]. To study redox signalling by H_2O_2 *in vivo* and explore its involvement in health and longevity, we used the fruit fly *Drosophila* as a model organism, with its tractable lifespan, powerful genetic tools, and strong evolutionary conservation of many central metabolic pathways with mammals[15]. Here we report that inducing an endogenous redox-shift, by manipulating levels of the H_2O_2-degrading enzyme catalase, improves health and robustly extends lifespan in flies, independently of oxidative stress resistance and dietary restriction. We find that the catalase redox-shifted flies are sensitive to starvation, which relies on autophagy as a vital survival mechanism. Importantly, we show that autophagy is essential for the lifespan extension of the catalase up-regulated flies, which is abolished in an autophagy-deficient background. Furthermore, using a redox-inactive knock-in mutant of Atg4a, a major effector of autophagy, we show that the lifespan extension in response to catalase requires a key redox-regulatory cysteine residue, Cys102 in *Drosophila* Atg4a. These findings demonstrate that redox regulation of autophagy can extend lifespan, confirming the importance of redox signalling in ageing and as a potential pro-longevity target.

Results

Catalase over-expression extends lifespan

To explore the role of endogenous redox signalling by H_2O_2 *in vivo*, we used the binary UAS/GAL4 expression system to up-regulate catalase in wild-type (WT) flies. Global up-regulation of catalase under control of the ubiquitous *daughterless* promoter (da-GAL4>UAS-cat) extended the median and maximum lifespan of female flies (typically by ~10-15%; Fig. 1a, Supplementary Table 1). Importantly, for these experiments, we used the *white Dahomey* (w^{Dah}) background, which is a long-lived and outbred WT, hence we are extending healthy lifespan and not rescuing a short-lived defect. Catalase was over-expressed ~5-10-fold at the mRNA level in whole flies (Supplementary Fig. 1a), which led to corresponding increases in catalase protein (Supplementary Figs. 1b,c) and enzymatic activity (Supplementary Fig. 1d). Interestingly, lifespan was not extended in males (Fig. 1a), despite similar catalase over-expression (Supplementary Figs. 1a,b,d). However, interventions such as modulating nutrient-sensing in *Drosophila* often show gender-specific effects on survival in females, that are absent or marginal in males[16]. Furthermore, over-expression of a mitochondria-targeted catalase (da-GAL4>UAS-mito-cat) did not lead to lifespan extension in either females or males (Supplementary Figs. 1e,f), consistent with a previous study[17], suggesting that altered mitochondrial metabolism is not implicated.

The da-GAL4>UAS-cat flies were mildly delayed in eclosing, without affecting the overall proportion of larvae surviving to adulthood (Supplementary Fig. 1g). To exclude developmental effects, we showed that the lifespan extension could be fully recapitulated using the inducible GeneSwitch system, with over-expression from d2 of adulthood onwards (da-GS>UAS-cat ±RU; Fig. 1b and Supplementary Fig. 1a). Varying the dose of the inducer drug RU (50-400 µM) still did not extend lifespan in males and had only marginal further effects on lifespan extension in females (Supplementary Fig. 1h), suggesting that the catalase benefits are already maximal. Over-expressing catalase under the control of an alternative ubiquitous driver (actin5c-GAL4>UAS-cat) also extended lifespan (Supplementary Fig. 1i). Furthermore, catalase-mediated lifespan extension was independent of *Wolbachia* status (Supplementary Figs. 1j,k), which can influence fly longevity and physiology[18,19]. In addition to lifespan extension, the catalase females also exhibited increased healthspan[20], as inferred from their enhanced climbing ability with age (Fig. 1c). To explore effects on age-specific mortality, trajectories derived from the survival curves revealed a shift in the intercept, but not the slope (Fig. 1d), indicating that catalase up-regulation decreased the overall risk of death, rather than slowing its rate of increase with age[21]. Therefore, the catalase over-expressing flies were healthier for longer.

Using the inducible GeneSwitch system showed that induction of catalase from middle-age (d28 and d42) or old-age (d56) was sufficient to extend lifespan (Fig. 1e and Supplementary Fig. 1l),

although not to the full extent from induction at d2. By d56, the –RU control flies had already started dying, yet switching to +RU treatment even at this late stage still enhanced survival. This implies that for full benefits the redox shift needs to occur early in life, yet late-onset still offers protection. The level of catalase up-regulation induced by RU was equivalent at all ages, as were the levels of endogenous catalase in the controls, eliminating any contribution from changes in RU consumption or endogenous catalase expression with age (Supplementary Fig. 1m). We conclude that ubiquitous up-regulation of catalase improves healthspan and extends lifespan in female flies. Interestingly, tissue-specific catalase up-regulation using a range of drivers (e.g. renal tubules, insulin-producing cells, pan-neuronal, intestine, fat body, pan-muscular; Supplementary Figs. 1n-s), did not recapitulate the strong lifespan extension obtained by the ubiquitous drivers, suggesting that catalase is either acting in an untested tissue (or combination of tissues), or alternatively is required at a global, organismal level. The catalase over-expressors were exceptionally resistant to multiple modes of oxidative stress - by exogenous H_2O_2 (Fig. 1f), the redox cycler paraquat both upon feeding (Fig. 1g) and injection (Fig. 1h), as well as hyperoxia (Fig. 1i and Supplementary Fig. 1t). However, this enhanced oxidative stress resistance is unlikely to explain the lifespan extension in females, because catalase over-expression protected males to a similar extent against oxidative stress without increasing longevity.

Catalase-induced lifespan extension is not through dietary restriction (DR)

To explore the mechanism underlying the catalase-mediated lifespan extension, we examined its relationship to DR, which is a robust and evolutionary conserved nutritional intervention known to have health and longevity benefits[22]. We measured the lifespan response of catalase over-expressor females to DR by varying the yeast content (i.e. protein source) in the food, while maintaining the sugar content constant[23]. This generated a typical tent-shaped response (Fig. 2a), with lifespan decreased at very low yeast levels (0.1x), highest under restricted conditions (0.5x), then gradually shortened towards more fully-fed conditions (1.5x). The lifespan of the catalase flies was enhanced relative to controls at all yeast levels (Fig. 2a), while fecundity increased with yeast content throughout the 0.1-1.5x range for both the control and catalase females (Supplementary Fig. 2a). Therefore, the catalase over-expressor females exhibit a normal DR response, and the lifespan extension upon catalase up-regulation is not mediated by the activation of DR pathways.

While the DR experiment revealed improved survival compared to control at a range of yeast concentrations (Fig. 2a), including extremely poor nutritional conditions (0.1x-yeast; Fig. 2b), we unexpectedly observed that the catalase over-expressing females, but not males, were more sensitive than control to complete starvation (Fig. 2c). There was no difference in triacylglyceride (TAG) levels both basally (t=0) and during a starvation time course, either at d7 (Fig. 2d) or d28

(Supplementary Fig. 2b). Similarly, the levels of glycogen storage and mobilisation were the same
in control and over-expressor females (Fig. 2e). Therefore, the starvation sensitivity of the catalase
flies was not due to differences in metabolic energy reserves or their mobilisation.

Autophagy is up-regulated and required for the catalase-mediated lifespan extension
Autophagy is a known longevity assurance process, involved in the response to nutritional
challenges such as starvation[24-28]. Furthermore, there is evidence for redox-regulation of
autophagy[29]. Therefore, we next explored the involvement of autophagy in the differential starvation
response and longevity of the catalase flies. To monitor autophagy status, we stained fly midguts
with LysoTracker Red to label autophagolysosomes (Fig. 2f). In parallel, we also monitored
autophagic flux using the dye CytoID Green, which accumulates when autophagy is blocked[30]
(Supplementary Fig. 2c). The number of LysoTracker punctae was significantly increased in the
catalase over-expressors (Fig. 2f), without detectable CytoID staining (Supplementary Fig. 2d),
indicating that autophagy is induced without blocking flux. Mitophagy levels were unchanged upon
catalase up-regulation as inferred from the genetically-encoded mito-QC fluorescent reporter[31]
(Supplementary Fig. 2e).

To assess autophagy by an orthologous approach, we quantified the levels of Atg8 (LC3 in
mammals), a major autophagosome marker, by immunoblotting. The levels of both the unlipidated
(Atg8-I) and lipidated (Atg8-II) forms were strongly elevated in the catalase over-expressors
(Fig. 2g), confirming that autophagy was induced. Indeed, levels of autophagy are physiologically
fine-tuned, with both autophagy inhibition and excessive activation shown to induce starvation
sensitivity *in vivo*[32-34]. Furthermore, similarly to the starvation stress assay, the catalase flies were
also sensitive to treatment with the autophagy inhibitor chloroquine (Supplementary Fig. 2f).

To test the involvement of autophagy in the longevity of the catalase flies, we down-
regulated autophagy by RNAi of Atg5 (Supplementary Fig. 2g), which impairs autophagy[32,34]. Atg5
knock-down enhanced sensitivity to starvation (Supplementary Fig. 2h), but did not affect the
lifespan of control females under fed conditions (da-GS>UAS-Atg5RNAi ±RU; Fig. 2h), consistent
with previous reports[34]. Importantly, the lifespan extension by catalase over-expression was
abolished in an Atg5-RNAi background (da-GS>UAS-Atg5RNAi+UAS-cat ±RU; Fig. 2i).
Therefore, autophagy is required for the enhanced longevity by catalase up-regulation.

Redox proteomics of the catalase up-regulated flies reveals an oxidising shift in bulk thiols
Redox regulation of autophagy has been described for Atg4 in the context of starvation-induced
ROS production *in vitro*[35]. Atg4 is the only cysteine peptidase amongst the autophagy components,
and is essential for autophagosome biogenesis[29]. Atg4 regulates autophagy by processing Atg8 at

two critical stages: 1) the initial cleavage of Atg8, mediated by the redox-insensitive catalytic cysteine of Atg4, therefore this first step promoting Atg8 lipidation is redox-independent; and 2) the subsequent redox-dependent de-lipidation of Atg8, which is selectively inactivated upon oxidation of an adjacent redox-regulatory cysteine in Atg4. Under oxidising conditions, lipidated Atg8 accumulates due to the redox-mediated suppression of de-conjugation by Atg4, therefore enhancing autophagosome biogenesis and promoting Atg4-mediated autophagy (Fig. 3a).

We previously showed that fasting for 24 h is associated with a strong oxidising shift of bulk cysteine residues *in vivo* in *Drosophila*[6]. We therefore hypothesised that the starvation sensitivity of the da-GAL4>UAS-cat females may be attributed to such thiol redox changes. To explore the effects of catalase up-regulation on global thiol redox state, we applied the same redox proteomic technique (oxidative isotope-coded affinity tags, OxICAT)[6] to the catalase over-expressing females. In OxICAT, samples undergo differential labelling of cysteine residues according to their redox status, followed by trypsin proteolysis and enrichment for cysteine-containing peptides, and finally detection by tandem mass spectrometry (Supplementary Fig. 3a). This allows both the identification of redox-responsive cysteine residues, as well as determination of their redox state. The bulk redox state of cysteines in control flies does not change with age, with the majority remaining at ~10-15% oxidised[6]. In contrast, the catalase over-expressors displayed an oxidising shift in cysteine redox state relative to controls with increasing age (Figs. 3b-d, Supplementary Figs. 3b-e, and Supplementary Tables 2-3). This finding is surprising, since we are over-expressing an antioxidant enzyme. Indeed, we confirmed that global ROS levels were decreased upon catalase up-regulation, as inferred from the redox-sensitive fluorescent dye CellROX (Supplementary Figs. 3f,g). Mitochondrial H_2O_2 levels, measured using the *in vivo* mitochondria-targeted ratiometric mass spectrometry probe MitoB[36], were unchanged in the da-GAL4>UAS-cat females compared to UAS-cat/+ controls (Supplementary Fig. 3h), reflecting the fact that this catalase is not mitochondria-targeted.

To reconcile catalase up-regulation with the oxidising shift in bulk cysteine redox state, we tested the hypothesis that by quenching cytoplasmic H_2O_2, catalase over-expression blocks physiological H_2O_2-mediated redox signalling that up-regulates other antioxidant systems and redox couples. The Keap1/Nrf2 signalling pathway is an appealing candidate for this process, as it is an oxidative stress response pathway that enhances the expression of multiple redox systems and is known to be redox-regulated in *Drosophila*[37]. To assess this pathway, we used a transgenic reporter for Keap1/Nrf2 activity, gstD-GFP. This pathway was up-regulated with age in controls, but not in the long-lived catalase flies (Fig. 3e), suggesting that catalase over-expression prevented the induction of Keap1/Nrf2 signalling with age, and thus a range of downstream redox processes. Consistent with this finding, total levels of glutathione - an important cellular antioxidant and redox

cofactor, whose synthesis is up-regulated by Nrf2 signalling - were decreased in the catalase overexpressors (Fig. 3f). Altogether, we have shown that catalase flies undergo an unexpected global oxidising thiol redox shift with age. This oxidation is consistent with the enhancement of autophagy via redox-regulation of Atg4.

Cys102 in Atg4a is required for the redox-mediated lifespan extension

The protein sequence of Atg4a is evolutionarily conserved, with both the catalytic cysteine (Cys98 in *Drosophila*) and the adjacent redox-regulatory cysteine (Cys102 in *Drosophila*) present in flies and mammals. This preservation is highlighted by structural modelling of *Drosophila* Atg4a and alignment analysis against human ATG4A (Figs. 4a,b and Supplementary Fig. 4a). To dissect the physiological role of Atg4 redox-regulation *in vivo*, we generated a transgenic knock-in fly line by CRISPR, where the regulatory cysteine in endogenous Atg4a was replaced by a redox-inactive serine residue (C102S mutant). This C102S knock-in mutation did not affect Atg4a expression (Supplementary Fig. 4b), or levels of catalase activity both basally and upon genetic up-regulation (Supplementary Fig. 4c). Catalase over-expression in Atg4a-C102S knock-in flies displayed the same strong resistance to H_2O_2-induced oxidative stress as in the Atg4a-WT background (Supplementary Fig. 4d).

Basal levels of autophagy were not affected under control conditions in the redox knock-in line (UAS-cat/+, Atg4a-WT *v.* UAS-cat/+, Atg4a-C102S), whereas autophagy induction by catalase over-expression was abolished (da-GAL4>UAS-cat, Atg4a-C102S), as assessed both by Western blotting against Atg8 (Fig. 4c and Supplementary Fig. 4e) and LysoTracker staining (Fig. 4d and Supplementary Fig. 4f). Therefore, this redox-regulatory cysteine in Atg4a is required for autophagy induction by redox signalling *in vivo*, as previously reported *in vitro*[35].

To interrogate the role of Atg4a-Cys102 in mediating the longevity of the catalase flies, we performed survival assays with the Atg4a-WT CRISPR control line, and reproduced the catalase lifespan extension in this background (Fig. 4e). The Atg4a-C102S point mutation did not affect survival of control flies, confirming that this knock-in alone is not deleterious (Fig. 4f). Critically, in contrast to the Atg4a-WT control, the lifespan extension upon catalase up-regulation was fully abolished in the Atg4a-C102S mutant background (Fig. 4f), as well as the climbing benefits (Supplementary Fig. 4g). While catalase up-regulation led to decreased egg laying (Supplementary Fig. 4h), this was observed in both the Atg4a-WT and C102S backgrounds, therefore any changes in fecundity cannot explain the lifespan extension. Overall, redox-regulation of autophagy via Atg4a-Cys102 mediates the longevity upon catalase over-expression. Enhancing autophagy is an evolutionarily conserved intervention associated with health and survival benefits, and here we demonstrate that selective redox-mediated up-regulation of autophagy can extend lifespan.

Discussion

Many attempts have been made to extend lifespan in model organisms by enhancing their antioxidant capacity, notably through the over-expression of antioxidant enzymes, including catalase[38-41]. These trials have been largely unsuccessful, casting doubt on the causative role of ROS and oxidative damage in ageing[42]. Therefore, our finding that catalase over-expression extends lifespan was at first surprising. The original study over-expressing catalase in *Drosophila* found no effect on lifespan and only modest resistance to oxidative stress by H_2O_2[38]. However, the study used an extra chromosomal copy under its endogenous promoter, resulting in lower over-expression of catalase (~1.75-fold at the mRNA level and ~1.5-fold increased enzyme activity). The degree of catalase over-expression is therefore likely to be important for the lifespan extension. Furthermore, the earlier study used only males, while our findings show a robust effect specifically in females. The results of these studies are therefore reconcilable.

The sex-specificity of lifespan benefits upon catalase up-regulation is striking. Indeed, we find that ubiquitous catalase over-expression does not extend lifespan or induce starvation sensitivity in males. A recent study showed that basal autophagy levels are significantly higher in male flies[43], shedding light on this sex-specificity of the catalase effect. This suggests that females have a broader 'metabolic plasticity', and are more likely to benefit from interventions that enhance autophagy.

Importantly, our study also challenges widespread assumptions regarding the directionality of antioxidant interventions. Indeed, we find that *in vivo* up-regulation of the antioxidant catalase, while directly conferring strong resistance to acute oxidative stress challenges as expected, also unexpectedly leads to an indirect oxidising shift in bulk cysteine redox state by interfering with homeostatic redox signalling. Consistent with our observations, a recent study in *C. elegans* has shown that excess dietary supplementation with *N*-acetyl cysteine (NAC) accelerates ageing by inhibiting *skn-1*-mediated transcription, the worm orthologue of Nrf2[44]. Therefore, caution is required in the interpretation of experiments involving the genetic or dietary manipulation of antioxidants.

Our unbiased redox proteomic analysis has revealed that the catalase flies undergo an oxidising shift in bulk cysteine redox state with age. Interestingly, this pattern is similar to our earlier observations in control flies under starvation[6]. Nutrient deprivation can directly affect redox homeostasis by depleting the provision of important building blocks for reducing equivalents, such as NADPH and glutathione, mediating an intracellular oxidising shift[45]. Therefore, we suggest that the catalase-induced thiol oxidising shift is perceived as an internal state of starvation, which triggers the induction of autophagy as a protective response (Fig. 4g). Autophagy plays a fundamental role in healthy physiology, such as cellular differentiation, tissue remodelling, and mitochondrial homeostasis, as well as in the response to stress and the clearance of cellular damage[27].

Consequently, enhanced autophagy is a common denominator of many evolutionary conserved interventions that extend lifespan[26,27], both genetically, such as down-regulation of insulin signalling[46], and pharmacologically, for instance rapamycin treatment[47]. Furthermore, direct up-regulation of autophagy has been shown to exert health benefits and extend lifespan in a range of model organisms including worms, flies and mice[34,48-50]. Several components of the autophagy pathway are known to be redox-regulated, including Atg3 and Atg7[51], as well as the focus of our study Atg4[35,52].

Overall, we have shown that shifting the *in vivo* redox state of *Drosophila* by over-expression of catalase extends lifespan and healthspan in females through redox-regulation of autophagy via a key redox-responsive cysteine in Atg4a. Our findings further emphasise the importance of fine-tuning autophagy in health and disease, and demonstrate how manipulation of redox signalling *in vivo* can ameliorate the effects of ageing. Furthermore, our data are consistent with a growing view in the ageing field that many effects of ROS on longevity are likely to be through alterations in redox signalling rather than through diminished oxidative damage[9,10,53].

Methods

Fly strains and husbandry

The *white Dahomey* (w^{Dah}) strain of *Drosophila melanogaster* was used as the WT background. The *Dahomey* stock was collected in 1970 in Dahomey (presently the Republic of Benin), and maintained since then as large population cages, ensuring outbreeding and overlapping generations. The w^{Dah} stock was derived by incorporation of the w^{1118} mutation into the outbred *Dahomey* background by back-crossing. Flies were either negative (w^{Dah}) or positive (w^{Dah+}) for the bacterial cytoplasmic endosymbiont *Wolbachia*, with infection status confirmed by PCR using published primers against *wsp*[18]. The w^{Dah} stock was originally achieved by tetracycline treatment of the w^{Dah+} stock[18]. The following transgenic lines were used: UAS-cat (BDSC 24621), UAS-catRNAi (VDRC 6283), gstD-GFP[37], UAS-Atg5RNAi[32], UAS-mito-cat[54], UAS-mito-QC[31]. The following driver lines were used: da-GAL4[55], da-GS[56], act5c-GAL4 (BDSC 4414), Uro-GAL4[57], dilp2-GAL4[58], elav-GS (BDSC 43642), S$_1$106-GS (BDSC 8151), TIGS-2[59], and MHC-GS[60]. All transgenic lines were back-crossed into the appropriate w^{Dah} or w^{Dah+} background for at least 6-10 generations. Experimental flies were incubated at 25ºC on a 12 h light:12 h dark cycle with 65% humidity.

Generation of the Atg4a lines

The Atg4a-C102S mutant fly line was generated by a fully transgenic CRISPR/Cas9 approach using two guide RNAs (gRNAs)[61] targeting the Atg4a (CG4428) gene. The C102S mutation together with an amino-terminal FLAG-tag were introduced using a donor construct for homology directed repair.

As a control, a fly line carrying only the FLAG-tag, referred to as Atg4a-WT, was generated using the same approach. The gRNA construct was generated by PCR with Phusion polymerase (NEB) and primers SOL897 and SOL898 (see Supplementary Table 4 for primer sequences) and the plasmid pCDF4 as template[61]. The resulting PCR product was cloned into the pCDF4 vector using the Gibson Assembly kit (NEB) and the resulting plasmid was then used to generate ATG4-gRNA transgenic flies via injection into the *y¹, sc¹, v¹, P{y[+t7.7]=nos-phiC31\int.NLS}X; P{y[+t7.7]=CaryP}attP2* fly line (BDSC 25710). Atg4a donor constructs were cloned in a two-step approach. First, a 2792 bp part encompassing the Atg4a gene locus was amplified by PCR using SOL926 and SOL927 on BAC clone CH321-39B1 (BACPAC Resources) as a template and cloned into the pBluescript II KS vector via *NotI* and *KpnI* restriction mediated ligation. In the next step, gene synthesis (Eurofins Genomics) was used to synthetise 806 bp long fragments encoding an amino-terminal FLAG-tag, the C102S mutation and mutations within the PAM sites to introduce the Atg4a-C102S mutation, or an amino-terminal FLAG-tag and mutations within the PAM sites to generate the Atg4a-WT construct, respectively. These fragments were then used to replace the wild-type sequence in the 2792 bp construct using the endogenous restriction sites *BstBI* and *MfeI*. In order to introduce the mutation, Atg4a-gRNA transgenic flies were crossed with flies expressing Cas9 under the ubiquitous actin promotor, and their progeny was injected with the Atg4a-C102S and Atg4a-WT donor constructs. PCR screening with primers SOL955 and SOL929 that specifically target the FLAG-tag sequence was used to identify positive CRISPR events and the presence of the C102S mutation was confirmed by sequencing. All injections were performed by the transgenic fly facility of the Max Planck Institute for Biology of Ageing.

Fly media

Flies were raised on standard sugar-yeast-agar medium (SYA) consisting of: 5% w/v sucrose (granulated sugar, Silver Spoon), 10% w/v brewer's yeast (#903312, MP Biomedicals), 1.5% w/v agar (A7002, Sigma), supplemented with nipagin (H5501, Sigma; 30 mL/L of 10% w/v nipagin in 95% EtOH) and propionic acid (P1386, Sigma; 0.3% v/v) as mould inhibitors, added once the food had cooled down to ~60ºC[23]. Expression via the inducible GeneSwitch system was achieved by addition of the drug RU (RU486/mifepristone; M8048, Sigma) to standard SYA once cooled down to ~60ºC, typically at 200 μM from a 0.1 M stock in EtOH. For dietary restriction (DR) experiments, the yeast content was varied to give 1% (0.1x), 5% (0.5x), 7.5% (0.75x), 10% (1x = SYA) or 15% (1.5 SYA) w/v yeast[62].

Experimental flies

For all experiments, eggs were collected over a defined period (<24 h) to ensure a synchronous population and reared at constant density in 200 mL bottles with SYA[62]. Eclosing adults of a defined age were kept as a mixed population for ~48 h to allow mating, then separated into males and females under mild CO_2 anaesthesia, and maintained as separate sexes from then onwards.

Lifespan & stress assays

Lifespan assays were set up as above, typically with n~10-15 flies per vial and a total of n~100-250 flies per condition. Flies were transferred to fresh food without gassing every ~2-3 days, with deaths and censors recorded. Stress assays were performed on d7 flies (typically n>100 per condition in groups of ~15-20 flies per vial), with deaths scored regularly following initiation of treatment. See Supplementary Table 1 for full survival assay information. For H_2O_2 resistance, flies were transferred onto medium containing 5% v/v H_2O_2 (H1009, Sigma), 5% w/v sucrose, 1.5% w/v agar. For paraquat stress, flies were either transferred onto standard SYA food supplemented with 20 mM paraquat (856177, Sigma), or injected with 75 nL of 1 mg/mL paraquat in Ringers buffer (3 mM $CaCl_2$, 182 mM KCl, 46 mM NaCl, 10 mM Tris base, pH 7.2 HCl) and maintained on standard SYA[63]. Starvation stress was assayed by transferring flies to 1.5% w/v agar medium, which lacks nutrients but allows hydration. Chloroquine (C6628, Sigma) was prepared as 10 mM in SYA or 5% w/v sugar, 1.5% w/v agar. Hyperoxia was performed by incubating flies on standard SYA vials in a glove box chamber set at 90% O_2 using a ProOx controller (BioSpherix). The majority of lifespans (Figs. 1a,b,e, 2a,b,h,i, 4e,f and Supplementary Figs. 1j-l) and stress assays (Figs. 1f,g, 2c, 4d and Supplementary Figs. 2f,h, 4d) were repeated at least twice as independent biological experiments, except Figs. 1h,i and Supplementary Figs. 1e,f,h,i,n-s which were performed once.

Development time

Eggs were collected from flies in cages onto grape juice agar plates over a defined time window (~4 h). After ~24 h, the resulting L1 larvae were picked onto SYA food at a density of 50 per vial (n=500 total per genotype), and the time to adult eclosion was monitored.

Climbing assay

Climbing ability (negative geotaxis) was assayed essentially as described[63]. Briefly, groups of 15 flies were transferred to a sawn-open 25 mL serological pipette (35 cm long, 1.5 cm diameter), with the base sealed by parafilm. The flies were tapped down within the column and observed during 45 s, after which their location was recorded. The column was separated into three sections: top 10 cm, middle, bottom 3 cm. Each cohort was evaluated 3 times, using 5 groups per genotype. The climbing performance index was calculated as: $1/2 \, (n^{total} + n^{top} - n^{bottom}/n^{total})$.

Metabolic and molecular assays

Flies for molecular experiments were rapidly transferred to pre-chilled microtubes via a small plastic funnel and snap frozen in liquid nitrogen, then stored at –80°C until required. Flies were always frozen at approximately the same time of day to minimise any circadian variation. For some assays, frozen flies were separated into body segments using forceful manual impact[64].

Quantitative PCR (QPCR)

Total RNA was extracted using Trizol (Ambion) according to the manufacturer's instructions (typically n=10 flies per sample, with n=4-5 replicates per condition). For some experiments, RNA was extracted from head and thorax tissue (i.e. abdomens removed), to avoid interference from the ovaries as the UAS/GAL4 system does not express in the germline. The concentration of total RNA purified for each sample was measured spectrophotometrically. 3 µg of total RNA was then subjected to DNA digestion using DNAse I (Ambion), immediately followed by reverse transcription using the Superscript II system (Invitrogen) with oligo dT primers. QPCR was performed using Fast SYBR Green Master Mix (Applied Biosystems) in a 7900HT Fast Real-Time PCR machine, and results processed by SDS2.4 software (Applied Biosystems), or using SYBR Green Supermix (1725124, Biorad) analysed with the QuantStudio 7 Flex Real-Time PCR System (Thermo Scientific). The primers were designed using Primer BLAST[65] and are listed in Supplementary Table 5.

Western blotting

Frozen fly samples were homogenised directly into 2X Laemmli loading buffer (Bio-Rad) supplemented with 5% v/v β-mercaptoethanol (Sigma) using a pellet pestle and motor (usually 5/10 females into 100/200 µL) and separated by standard SDS-PAGE. The following primary antibodies were used at the indicated dilutions: anti-actin (Ab1801, Ab8224, or Ab8227, AbCam; 1:1,000), anti-Atg8 (a generous gift from K. Köhler[66]; 1:1,000), anti-catalase (C0979, Sigma; 1:10,000), anti-GFP (#2955, Cell Signaling Technology; 1:1,000). The following secondary antibodies were used: anti-mouse IgG (#7076, Cell Signaling Technology, or A4416, Sigma; 1:5,000), anti-rabbit IgG (#7074, Cell Signaling Technology; 1:5,000). Blots were developed using standard ECL, followed by analysis with FIJI software[67].

Catalase activity

Catalase activity was measured according to a published protocol[68], adapted for *Drosophila* samples. Briefly, 5 whole flies were homogenised in 200 µL PBS supplemented with protease inhibitor (HALT 1861284, ThermoScientific) using a TissueLyser (Precellys, 30sec), and centrifuged

(16,000 x g, 4°C, 10 min). 50 µL supernatants (1:10 diluted in PBS for the da-GAL4>UAS-cat

genotype) were transferred to flow cytometry tubes (Falcon 5 mL polystyrene, round bottom, 12 x

75 mm), and combined with 50 µL 1% Triton-X100, followed by 50 µL 30% H_2O_2 solution (Sigma

H1009). After incubating at RT for 10 min, the height of the stable foam was marked, the tubes

photographed and analysed using FIJI. Enzyme activity was calculated by linear regression against

a standard curve of purified catalase (C40, Sigma) dissolved in PBS.

Glutathione assay

Total glutathione levels were measured as described[69], adapted for *Drosophila* samples. Briefly,

cohorts of 10 female flies (separated into head+thorax and abdomen fractions[64]) were homogenised

in 200 µL assay buffer (0.1 M potassium phosphate (pH 7.5), 5 mM EDTA), supplemented with

0.1% v/v Triton X-100 and 0.6% sulfosalicylic acid. Samples were centrifuged (16,000 x g, 4°C,

10 min), and diluted 1:10 in assay buffer. 20 µL samples and glutathione standards (G6529, Sigma)

were transferred to a 96-well plate and overlayed with 120 µL of assay buffer containing DTNB

(D8130, Sigma; 2 mg/ 3mL) and 10 U glutathione reductase (G3664, Sigma; 10 U/3 mL), followed

by the addition of 60 µL NADPH (N7505, Sigma; 2 mg/3 mL assay buffer). The formation of TNB

was monitored kinetically at 412 nm over 5 min in a spectrophotometer (Fluostar Omega, BMG

Labtech). Glutathione levels were determined from the slope relative to the standard curve.

Energy storage assays

Whole body triacylglyceride (TAG) and glycogen levels were measured in d7 and d28 females (n=5

flies per sample, n=6-12 replicates per genotype) under control (fed) conditions and in response to

starvation. For the TAG assay, flies were homogenised in 0.05% v/v Tween-20 and assayed using

the Triglyceride Infinity Reagent (TR22421, ThermoScientific) in a 96-well plate measuring

absorbance at 540 nm. For the glycogen assay, d7 flies were homogenised in saturated sodium

sulphate, then the subsequent pellet was resuspended in anthrone reagent (319899, Sigma) and

assayed in a 96-well plate measuring absorbance at 620 nm[70].

OxICAT sample preparation and LC-MS/MS analysis

To measure the bulk redox state of protein cysteine residues, we performed unbiased redox

proteomics using OxICAT. Protein isolation, cysteine-residue labelling, peptide preparation and

LC-MS/MS analysis were performed exactly as described previously[6]. Briefly, cohorts of n=10

females flies were rapidly frozen in liquid N_2, and the frozen head and thorax tissue was separated

from the abdomens[64] to avoid signal from the ovaries. A total of n=5 biological replicates were

processed per condition. Samples were homogenised in ice-cold 100% w/v TCA to stabilise thiols

and solubilise proteins. Homogenates were centrifuged to pellet the chitin exoskeleton and other insoluble components, and the resulting supernatant transferred to a fresh tube. Protein samples were precipitated by decreasing the TCA concentration to 20% w/v with addition of H_2O, then washed successively with 10% and 5% w/v TCA. This protein precipitation was shown to occur with minimal protein losses or distortion of the protein complement[6].

Protein samples (~30-40 μg) were resuspended in denaturing alkylating buffer (DAB; 6 M urea, 2% w/v SDS, 200 mM Tris-HCl, 10 mM EDTA, 100 μM DTPA, 10 μM neocuprine). Reduced Cys residues were labelled with light ICAT reagent (4339036, AB Sciex) for 2 h at 37°C, 1400 rpm in an Eppendorf Thermomixer. Proteins were precipitated and washed with ice-cold acetone, solubilised in DAB with 1 mM TCEP (tris(2-carboxyethyl)phosphine) to reduce previously reversibly oxidised Cys residues, which were then labelled with heavy ICAT reagent (4339036, AB Sciex) as above, before again precipitating and washing in ice-cold acetone. ICAT-labelled protein samples were resuspended in DAB and digested with trypsin overnight at 37°C. Digested samples were enriched for Cys-containing peptides first on a cation exchange cartridge, and then subsequently on an avidin affinity cartridge (both provided with the ICAT kit; 4339036, AB Sciex). The eluted peptides were dried down overnight in a SpeedVac, and the biotin moiety of the ICAT label cleaved.

Liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis of the OxICAT-labelled peptides was performed exactly as published[6], using an Orbitrap LTQ XL (Thermo) after chromatography on a nanoscale reverse-phase column. Raw files for each LC-MS/MS run were analysed using MaxQuant software to determine the ratio of heavy over light OxICAT-labelled peptides. Each biological sample (n=5 per condition) was run as 2 technical replicates, with the raw files from both LC-MS/MS runs grouped into a combined dataset. Besides light or heavy ICAT labelling of Cys residues, methionine oxidation was included as a possible modification, and up to 2 possible missed cleavages were allowed in the MaxQuant search parameters[6]. As a reference sequence database, a FASTA file containing all protein sequences associated with *D. melanogaster* was generated from UniProt, and used to identify peptides. In addition, the reference sequence database was used to create an *in silico* tryptic digest of the *D. melanogaster* proteome, listing all Cys-containing peptides (with 0, 1 or 2 missed cleavages), the Cys residue number, and the UniProt protein ID[6]. Ratios of heavy over light ICAT-labelled peptides obtained from MaxQuant were converted to % of the Cys residue reversibly oxidised (combining intensities from all peptide signals containing the Cys residue of interest (e.g. miscleaved, methionine oxidation, different z values), generating a mean % reversible oxidation for each unique Cys residue identified within the biological replicates (Supplementary Table 2). Functional annotations were obtained from DAVID[71] (https://david.ncifcrf.gov/tools.jsp; Supplementary Table 3).

Confocal microscopy

Autophagy and autophagic flux were assessed by LysoTracker Red and CytoID Green staining, respectively[30]. Briefly, intact guts were dissected into 1x PBS (70011-036, Gibco) and stained in the dark for 30 min with CytoID Green (ENZ-51036-K025, Enzo Life Sciences; 1:1,000 in 1x PBS), followed by a second incubation for 3 min in PBS with LysoTracker Red (L7528, Invitrogen; 1:2,000 in 1x PBS) and HOECHST (H1399, Invitrogen; 1:1,000 from a 1 mg/mL stock in ultrapure water) followed by three washes with PBS. Guts were mounted in mounting buffer (50% glycerol in PBS) and immediately imaged. The number of LysoTracker punctae and HOECHST-stained nuclei were quantified using using FIJI. As a positive control for CytoID staining, d7 female flies were fed for 48 h on standard SYA medium supplemented with 30 mM choloroquine prior to gut dissection and staining as described above. Quantification of punctae per cell was perfomed in FIJI.

Cellular ROS levels were assessed using CellROX Deep Red (C10422, Invitrogen; 1:500 in PBS). Guts were dissected with minimal light exposure, and stained for 30 min at 25ºC in the dark, then washed 3 times with 1x PBS prior to mounting (50% glycerol in PBS, with HOECHST 1:1,000 from a 1 mg/mL stock). Mean fluorescence intensity (MFI) was analysed using FIJI. As a positive control, dissected guts were treated *ex vivo* with 10 μM H_2O_2 for 10 min in 1x PBS at RT prior to CellRox staining as described above.

For catalase staining, whole guts were dissected in PBS and fixed in 4% paraformaldehyde (30450002-2, 2B Scientific Ltd) for 20 min. After 3x 5 min washes with PBST (0.1% v/v Triton X-100 in PBS), guts were blocked with 5% w/v BSA (A7906, Sigma) in PBST for 1 h. Guts were incubated overnight at 4ºC with a primary anti-catalase antibody (C0979, Sigma; 1:250 in 5% w/v BSA in PBS). After three washes with PBS, guts were incubated with an anti-mouse secondary antibody (A10037, Invitrogen; Alexa568) for 1 h at RT, washed and mounted in VectaShield containing DAPI (H1200-10, Vector Laboratories). Images were acquired on a Leica SP5 confocal microscope (zoom: 20x optical plus 5x digital; resolution: 512 x 512 px at 700 Hz). Laser power and optical settings were kept constant between images.

Structural modelling

The structure of *Drosophila* ATG4a was modelled using Phyre2[73] and visualised in Chimera X[74], superimposed on the crystal structure of human ATG4A (PDB 2P82). Distances between the amino acids belonging to the catalytic triad, as calculated in Chimera X[74], were comparable between human and *Drosophila*:

Human: His281_NE2⇄SG_Cys77 3.286 Å, His281_ND1⇄OD1_Asp279 2.616 Å;

Drosophila: Hs273_NE2⇄SG_Cys98 3.413 Å, His273_ND1⇄OD1_Asp271 2.620 Å;

MitoB injections

To assay *in vivo* mitochondrial H_2O_2 levels, d7 females were injected with the ratiometric mass spectrometry probe MitoB under mild CO_2 anaesthesia as described [36,64]. Flies were then returned to standard food (cohorts of 10 flies, n=6-7 replicates per condition) and incubated for 4 h on SYA medium at 25°C, before snap freezing. A t=0 h control (n=3 replicates) was snap frozen immediately to determine the background level of MitoP. Flies were homogenised, spiked with deuterated internal standards, and extracted to quantify the MitoP/MitoB ratio[36,64].

Statistical analysis

Lifespan and stress assays were plotted as cumulative survival curves, and statistical analysis was performed by Log-Rank test. Full details of survival data (n numbers, p values) are provided in Supplementary Table 1. Other data were analysed by Student's *t*-test or ANOVA as appropriate in GraphPad Prism v8-10.

Data availability

The proteomics dataset has been deposited to the ProteomeXchange Consortium[72] via the PRIDE partner repository (accession number #PXD002195).

Acknowledgements

This project has received funding from the Medical Research Council UK (MC-A654-5QB90) to HMC, and from the Max Planck Society to LP. MPM is supported by the Medical Research Council UK (MC-U105663142) and by a Wellcome Trust Investigator Award (110159/Z/15/Z). FC was supported by the Wellcome Trust/Royal Society (Sir Henry Dale Fellowship 102532/Z/12/Z and 102531/Z/13/A) and the DFG, German Research Foundation (EXC 2030 -390661388). We are grateful to Pirrko Salmiheimo and Mumtaz Ahmad for help with preparation of fly media, Jigna Patel for early technical support, Mike Harbour and Ian Fearnley for assistance with redox proteomics, and Yu-Xuan Lu for advice on confocal imaging. We thank members of the Partridge and Cochemé labs for valuable discussions throughout this project. We acknowledge the Bloomington *Drosophila* Stock Center and the Vienna *Drosophila* Resource Center (VRDC) for fly strains, and the BACPAC Resource Center for BAC clones. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

Author contributions

Conceived the project: LP and HMC. Performed experiments and data analysis: CL, IB, JICQ, LAGvL, AF, MB, JA, AM, HBK, PVS, AL, FC and HMC. Generated the CRISPR mutants: SG. Conducted and analysed the redox proteomics: KEM and AMJ. Supervised the work: MPM, LP and HMC. Wrote the manuscript: CL, MPM, LP and HMC, with input from all the authors.

Competing interests

The authors declare no competing interests.

References

Paulsen, C. E. & Carroll, K. S. Cysteine-mediated redox signaling: chemistry, biology, and tools for discovery. *Chem. Rev.* **113**, 4633-4679 (2013). https://doi.org/10.1021/cr300163e

Holmstrom, K. M. & Finkel, T. Cellular mechanisms and physiological consequences of redox-dependent signalling. *Nat. Rev. Mol. Cell Biol.* **15**, 411-421 (2014). https://doi.org/10.1038/nrm3801

Sies, H. & Jones, D. P. Reactive oxygen species (ROS) as pleiotropic physiological signalling agents. *Nat. Rev. Mol. Cell. Biol.* **21**, 363-383 (2020). https://doi.org/10.1038/s41580-020-0230-3

Lennicke, C. & Cochemé, H. M. Redox metabolism: ROS as specific molecular regulators of cell signaling and function. *Mol. Cell* **81**, 3691-3707 (2021). https://doi.org/10.1016/j.molcel.2021.08.018

Brandes, N. *et al.* Time line of redox events in aging postmitotic cells. *eLife* **2**, e00306 (2013). https://doi.org/10.7554/eLife.00306

Menger, K. E. *et al.* Fasting, but Not Aging, Dramatically Alters the Redox Status of Cysteine Residues on Proteins in *Drosophila melanogaster. Cell Rep.* **11**, 1856-1865 (2015). https://doi.org/10.1016/j.celrep.2015.05.033

Xiao, H. *et al.* A Quantitative Tissue-Specific Landscape of Protein Redox Regulation during Aging. *Cell* **180**, 968-983 e924 (2020). https://doi.org/10.1016/j.cell.2020.02.012

Meng, J. *et al.* Global profiling of distinct cysteine redox forms reveals wide-ranging redox regulation in *C. elegans. Nat. Commun.* **12**, 1415 (2021). https://doi.org/10.1038/s41467-021-21686-3

Jones, D. P. Redox theory of aging. *Redox Biol.* **5**, 71-79 (2015). https://doi.org/10.1016/j.redox.2015.03.004

Lennicke, C. & Cochemé, H. M. Redox signalling and ageing: insights from *Drosophila. Biochem. Soc. Trans.* **48**, 367-377 (2020). https://doi.org/10.1042/BST20190052

Stocker, S., Van Laer, K., Mijuskovic, A. & Dick, T. P. The Conundrum of Hydrogen Peroxide Signaling and the Emerging Role of Peroxiredoxins as Redox Relay Hubs. *Antioxid. Redox Signal.* **28**, 558-573 (2018). https://doi.org/10.1089/ars.2017.7162

van Dam, L. *et al.* The Human 2-Cys Peroxiredoxins form Widespread, Cysteine-Dependent- and Isoform-Specific Protein-Protein Interactions. *Antioxidants (Basel)* **10** (2021). https://doi.org/10.3390/antiox10040627

Lennicke, C. & Cochemé, H. M. Redox regulation of the insulin signalling pathway. *Redox Biol.* **42**, 101964 (2021). https://doi.org/10.1016/j.redox.2021.101964

14 Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. Hallmarks of
aging: An expanding universe. *Cell* **186**, 243-278 (2023).
https://doi.org/10.1016/j.cell.2022.11.001

15 Piper, M. D. W. & Partridge, L. *Drosophila* as a model for ageing. *Biochim. Biophys. Acta*
Mol. Basis Dis. **1864**, 2707-2717 (2018). https://doi.org/10.1016/j.bbadis.2017.09.016

16 Austad, S. N. & Fischer, K. E. Sex Differences in Lifespan. *Cell Metab.* **23**, 1022-1033
(2016). https://doi.org/10.1016/j.cmet.2016.05.019

17 Scialo, F. *et al.* Mitochondrial ROS Produced via Reverse Electron Transport Extend
Animal Lifespan. *Cell Metab.* **23**, 725-734 (2016).
https://doi.org/10.1016/j.cmet.2016.03.009

18 Toivonen, J. M. *et al.* No influence of Indy on lifespan in *Drosophila* after correction for
genetic and cytoplasmic background effects. *PLOS Genet.* **3**, e95 (2007).
https://doi.org/10.1371/journal.pgen.0030095

19 Ikeya, T., Broughton, S., Alic, N., Grandison, R. & Partridge, L. The endosymbiont
Wolbachia increases insulin/IGF-like signalling in *Drosophila*. *Proc. Biol. Sci.* **276**, 3799-
3807 (2009). https://doi.org/10.1098/rspb.2009.0778

20 Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The hallmarks of
aging. *Cell* **153**, 1194-1217 (2013). https://doi.org/10.1016/j.cell.2013.05.039

21 Mair, W., Goymer, P., Pletcher, S. D. & Partridge, L. Demography of dietary restriction
and death in *Drosophila*. *Science* **301**, 1731-1733 (2003).
https://doi.org/10.1126/science.1086016

22 Fontana, L. & Partridge, L. Promoting health and longevity through diet: from model
organisms to humans. *Cell* **161**, 106-118 (2015). https://doi.org/10.1016/j.cell.2015.02.020

23 Grandison, R. C., Wong, R., Bass, T. M., Partridge, L. & Piper, M. D. Effect of a
standardised dietary restriction protocol on multiple laboratory strains of *Drosophila*
melanogaster. *PLOS One* **4**, e4067 (2009). https://doi.org/10.1371/journal.pone.0004067

24 Vilchez, D., Saez, I. & Dillin, A. The role of protein clearance mechanisms in organismal
ageing and age-related diseases. *Nat. Commun.* **5**, 5659 (2014).
https://doi.org/10.1038/ncomms6659

25 Kaushik, S. & Cuervo, A. M. Proteostasis and aging. *Nat. Med.* **21**, 1406-1415 (2015).
https://doi.org/10.1038/nm.4001

26 Madeo, F., Zimmermann, A., Maiuri, M. C. & Kroemer, G. Essential role for autophagy in
life span extension. *J. Clin. Invest.* **125**, 85-93 (2015). https://doi.org/10.1172/JCI73946

27 Hansen, M., Rubinsztein, D. C. & Walker, D. W. Autophagy as a promoter of longevity:
insights from model organisms. *Nat. Rev. Mol. Cell Biol.* **19**, 579-593 (2018).
https://doi.org/10.1038/s41580-018-0033-y

28 Aman, Y. *et al.* Autophagy in healthy aging and disease. *Nat Aging* **1**, 634-650 (2021).
https://doi.org/10.1038/s43587-021-00098-4

29 Scherz-Shouval, R. & Elazar, Z. Regulation of autophagy by ROS: physiology and
pathology. *Trends Biochem. Sci.* **36**, 30-38 (2011).
https://doi.org/10.1016/j.tibs.2010.07.007

30 Lu, Y. X. *et al.* A TORC1-histone axis regulates chromatin organisation and non-canonical
induction of autophagy to ameliorate ageing. *eLife* **10** (2021).
https://doi.org/10.7554/eLife.62233

31 Lee, J. J. *et al.* Basal mitophagy is widespread in *Drosophila* but minimally affected by
loss of Pink1 or parkin. *J. Cell Biol.* **217**, 1613-1622 (2018).
https://doi.org/10.1083/jcb.201801044

32   Scott, R. C., Schuldiner, O. & Neufeld, T. P. Role and regulation of starvation-induced
autophagy in the *Drosophila* fat body. *Dev. Cell* **7**, 167-178 (2004).
https://doi.org/10.1016/j.devcel.2004.07.009

33   Kang, C., You, Y. J. & Avery, L. Dual roles of autophagy in the survival of
Caenorhabditis elegans during starvation. *Genes & development* **21**, 2161-2171 (2007).
https://doi.org/10.1101/gad.1573107

34   Bjedov, I. *et al.* Fine-tuning autophagy maximises lifespan and is associated with changes
in mitochondrial gene expression in Drosophila. *PLOS Genet.* **16**, e1009083 (2020).
https://doi.org/10.1371/journal.pgen.1009083

35   Scherz-Shouval, R. *et al.* Reactive oxygen species are essential for autophagy and
specifically regulate the activity of Atg4. *EMBO J.* **26**, 1749-1760 (2007).
https://doi.org/10.1038/sj.emboj.7601623

36   Cochemé, H. M. *et al.* Measurement of H_2O_2 within living *Drosophila* during aging using a
ratiometric mass spectrometry probe targeted to the mitochondrial matrix. *Cell Metab.* **13**,
340-350 (2011). https://doi.org/10.1016/j.cmet.2011.02.003

37   Sykiotis, G. P. & Bohmann, D. Keap1/Nrf2 signaling regulates oxidative stress tolerance
and lifespan in *Drosophila*. *Dev. Cell* **14**, 76-85 (2008).
https://doi.org/10.1016/j.devcel.2007.12.002

38   Orr, W. C. & Sohal, R. S. The effects of catalase gene overexpression on life span and
resistance to oxidative stress in transgenic *Drosophila melanogaster*. *Arch. Biochem.*
Biophys. **297**, 35-41 (1992). https://doi.org/10.1016/0003-9861(92)90637-c

39   Orr, W. C., Mockett, R. J., Benes, J. J. & Sohal, R. S. Effects of overexpression of copper-
zinc and manganese superoxide dismutases, catalase, and thioredoxin reductase genes on
longevity in *Drosophila melanogaster*. *J. Biol. Chem.* **278**, 26418-26422 (2003).
https://doi.org/10.1074/jbc.M303095200

40   Doonan, R. *et al.* Against the oxidative damage theory of aging: superoxide dismutases
protect against oxidative stress but have little or no effect on life span in *Caenorhabditis*
elegans. *Genes & development* **22**, 3236-3241 (2008). https://doi.org/10.1101/gad.504808

41   Perez, V. I. *et al.* The overexpression of major antioxidant enzymes does not extend the
lifespan of mice. *Aging Cell* **8**, 73-75 (2009). https://doi.org/10.1111/j.1474-
9726.2008.00449.x

42   Gems, D. & Doonan, R. Antioxidant defense and aging in *C. elegans*: is the oxidative
damage theory of aging wrong? *Cell Cycle* **8**, 1681-1687 (2009).
https://doi.org/10.4161/cc.8.11.8595

43   Regan, J. C. *et al.* Sexual identity of enterocytes regulates autophagy to determine
intestinal health, lifespan and responses to rapamycin. *Nat Aging* **2**, 1145-1158 (2022).
https://doi.org/10.1038/s43587-022-00308-7

44   Gusarov, I. *et al.* Dietary thiols accelerate aging of C. elegans. *Nat. Commun.* **12**, 4336
(2021). https://doi.org/10.1038/s41467-021-24634-3

45   Filomeni, G., De Zio, D. & Cecconi, F. Oxidative stress and autophagy: the clash between
damage and metabolic needs. *Cell Death Differ.* **22**, 377-388 (2015).
https://doi.org/10.1038/cdd.2014.150

46   Toth, M. L. *et al.* Longevity pathways converge on autophagy genes to regulate life span in
Caenorhabditis elegans. *Autophagy* **4**, 330-338 (2008). https://doi.org/10.4161/auto.5618

47   Bjedov, I. *et al.* Mechanisms of life span extension by rapamycin in the fruit fly
Drosophila melanogaster. *Cell Metab.* **11**, 35-46 (2010).
https://doi.org/10.1016/j.cmet.2009.11.010

48 Simonsen, A. *et al.* Promoting basal levels of autophagy in the nervous system enhances
longevity and oxidant resistance in adult *Drosophila*. *Autophagy* **4**, 176-184 (2008).
https://doi.org/10.4161/auto.5269

49 Lapierre, L. R. *et al.* The TFEB orthologue HLH-30 regulates autophagy and modulates
longevity in *Caenorhabditis elegans*. *Nat. Commun.* **4**, 2267 (2013).
https://doi.org/10.1038/ncomms3267

50 Pyo, J. O. *et al.* Overexpression of Atg5 in mice activates autophagy and extends lifespan.
Nat. Commun. **4**, 2300 (2013). https://doi.org/10.1038/ncomms3300

51 Frudd, K., Burgoyne, T. & Burgoyne, J. R. Oxidation of Atg3 and Atg7 mediates
inhibition of autophagy. *Nat. Commun.* **9**, 95 (2018). https://doi.org/10.1038/s41467-017-
02352-z

52 Ligeon, L. A. *et al.* Oxidation inhibits autophagy protein deconjugation from phagosomes
to sustain MHC class II restricted antigen presentation. *Nat. Commun.* **12**, 1508 (2021).
https://doi.org/10.1038/s41467-021-21829-6

53 Orr, W. C., Radyuk, S. N. & Sohal, R. S. Involvement of redox state in the aging of
Drosophila melanogaster. *Antioxid. Redox Signal.* **19**, 788-803 (2013).
https://doi.org/10.1089/ars.2012.5002

54 Radyuk, S. N. *et al.* Mitochondrial peroxiredoxins are critical for the maintenance of redox
state and the survival of adult Drosophila. *Free Radic. Biol. Med.* **49**, 1892-1902 (2010).
https://doi.org/10.1016/j.freeradbiomed.2010.09.014

55 Wodarz, A., Hinz, U., Engelbert, M. & Knust, E. Expression of crumbs confers apical
character on plasma membrane domains of ectodermal epithelia of *Drosophila*. *Cell* **82**,
67-76 (1995). https://doi.org/10.1016/0092-8674(95)90053-5

56 Tricoire, H. *et al.* The steroid hormone receptor EcR finely modulates *Drosophila* lifespan
during adulthood in a sex-specific manner. *Mech. Ageing Dev.* **130**, 547-552 (2009).
https://doi.org/10.1016/j.mad.2009.05.004

57 Terhzaz, S. *et al.* Cell-specific inositol 1,4,5 trisphosphate 3-kinase mediates epithelial cell
apoptosis in response to oxidative stress in *Drosophila*. *Cell Signal.* **22**, 737-748 (2010).
https://doi.org/10.1016/j.cellsig.2009.12.009

58 Ikeya, T., Galic, M., Belawat, P., Nairz, K. & Hafen, E. Nutrient-dependent expression of
insulin-like peptides from neuroendocrine cells in the CNS contributes to growth
regulation in *Drosophila*. *Curr. Biol.* **12**, 1293-1300 (2002). https://doi.org/10.1016/S0960-
9822(02)01043-6

59 Poirier, L., Shane, A., Zheng, J. & Seroude, L. Characterization of the *Drosophila* gene-
switch system in aging studies: a cautionary tale. *Aging Cell* **7**, 758-770 (2008).
https://doi.org/10.1111/j.1474-9726.2008.00421.x

60 Osterwalder, T., Yoon, K. S., White, B. H. & Keshishian, H. A conditional tissue-specific
transgene expression system using inducible GAL4. *Proc. Natl. Acad. Sci. U. S. A.* **98**,
12596-12601 (2001). https://doi.org/10.1073/pnas.221303298

61 Port, F., Chen, H. M., Lee, T. & Bullock, S. L. Optimized CRISPR/Cas tools for efficient
germline and somatic genome engineering in *Drosophila*. *Proc. Natl. Acad. Sci. U. S. A.*
111, E2967-2976 (2014). https://doi.org/10.1073/pnas.1405500111

62 Bass, T. M. *et al.* Optimization of dietary restriction protocols in *Drosophila*. *J. Gerontol.*
A Biol. Sci. Med. Sci. **62**, 1071-1081 (2007). https://doi.org/10.1093/gerona/62.10.1071

63 Kerr, F. *et al.* Direct Keap1-Nrf2 disruption as a potential therapeutic target for
Alzheimer's disease. *PLOS Genet.* **13**, e1006593 (2017).
https://doi.org/10.1371/journal.pgen.1006593

Cochemé, H. M. *et al.* Using the mitochondria-targeted ratiometric mass spectrometry probe MitoB to measure H_2O_2 in living *Drosophila*. *Nat. Protoc.* **7**, 946-958 (2012). https://doi.org/10.1038/nprot.2012.035

Ye, J. *et al.* Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC Bioinform.* **13**, 134 (2012). https://doi.org/10.1186/1471-2105-13-134

Barth, J. M., Szabad, J., Hafen, E. & Kohler, K. Autophagy in *Drosophila* ovaries is induced by starvation and is required for oogenesis. *Cell Death Differ.* **18**, 915-924 (2011). https://doi.org/10.1038/cdd.2010.157

Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676-682 (2012). https://doi.org/10.1038/nmeth.2019

Iwase, T. *et al.* A simple assay for measuring catalase activity: a visual approach. *Sci. Rep.* **3**, 3081 (2013). https://doi.org/10.1038/srep03081

Rahman, I., Kode, A. & Biswas, S. K. Assay for quantitative determination of glutathione and glutathione disulfide levels using enzymatic recycling method. *Nat. Protoc.* **1**, 3159-3165 (2006). https://doi.org/10.1038/nprot.2006.378

Castillo-Quan, J. I. *et al.* Lithium Promotes Longevity through GSK3/NRF2-Dependent Hormesis. *Cell Rep.* **15**, 638-650 (2016). https://doi.org/10.1016/j.celrep.2016.03.041

Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update). *Nucleic Acids Res.* **50**, W216-W221 (2022). https://doi.org/10.1093/nar/gkac194

Vizcaino, J. A. *et al.* ProteomeXchange provides globally coordinated proteomics data submission and dissemination. *Nat. Biotechnol.* **32**, 223-226 (2014). https://doi.org/10.1038/nbt.2839

Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. The Phyre2 web portal for protein modeling, prediction and analysis. *Nat. Protoc.* **10**, 845-858 (2015). https://doi.org/10.1038/nprot.2015.053

Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci.* **30**, 70-82 (2021). https://doi.org/10.1002/pro.3943

Figure 1

Fig. 1 | Ubiquitous catalase up-regulation extends lifespan in flies, independently of oxidative stress resistance. a, Constitutive, ubiquitous catalase over-expression (da-GAL4>UAS-cat) extended the survival of female flies in a w^{Dah} (*white Dahomey*) WT background relative to the UAS-cat/+ (p=5.7x10^{-7}) and da-GAL4/+ (p=3.3x10^{-6}) controls. The control lines were not significantly different from each other (p=0.7504). No effect was observed in males (p>0.05 for all comparisons). Lifespans were performed with n=200 flies per condition. **b,** Inducible catalase over-expression from early adulthood (d2) using the GeneSwitch system extended the lifespan of female flies (da-GS>UAS-cat ±RU, p=1.1x10^{-16}). RU had no effect on the da-GS/+ control line (p=0.7161). Lifespans were performed with n=225-300 flies per condition. Inset: catalase over-expression assessed by Western blotting in whole d9 females (=d7 of RU induction), with actin as a loading control. **c,** Healthspan, inferred from climbing performance, was improved in catalase over-expressing females. Climbing was assayed on da-GS>UAS-cat females ±RU to control for effects of eye colour on this behaviour. Data are presented as box-and-whisker plots (min/max error bars) of n=5 replicates per condition, each with n=15 flies per sample, analysed by unpaired Student's *t*-test. **d,** Mortality trajectories of the da-GS>UAS-cat ±RU survival curves from (**b**), fitted with a linear regression trendline (dotted line). **e,** Late onset over-expression of catalase using the inducible GeneSwitch system from either middle-age (d28 and d42) and old-age (d56) extended the lifespan of female flies (p=7.2x10^{-8}, p=1.0x10^{-7} and p=1.4x10^{-3}, respectively against the –RU control). Lifespans were performed with n=270 flies per condition, and were plotted from point of RU induction relative to the remaining –RU control flies at that age (see Supplementary Fig. 1l for the full survival data). **f,** Catalase over-expressing flies were strongly resistant to exogenous H_2O_2 stress relative to controls (da-GAL4>UAS-cat *v.* UAS-cat/+; p=5.0x10^{-61} females, p=3.2x10^{-33} males). H_2O_2 treatment (5% v/v in sucrose/agar medium) was initiated at d7, with n=105 males (n=75 for da-GS>UAS-cat) and n=120 females per condition. **g,** Catalase over-expressing flies were resistant to chronic dietary paraquat stress relative to control flies (da-GAL4>UAS-cat *v.* UAS-cat/+; p=6.93x10^{-31} females, p=7.99x10^{-16} males). Paraquat treatment (20 mM in SYA food) was initiated at d7, with n=100 flies per condition. **h,** Catalase over-expressing flies (da-GAL4>UAS-cat) were resistant to acute paraquat stress relative to controls (UAS-cat/+). d7 females were injected with 75 nL of 1 mg/mL paraquat in Ringers buffer (+PQ, n=150 flies) or mock injected with buffer alone (–PQ, n=120 flies). **i,** Catalase over-expressing flies were resistant to environmental hyperoxia stress relative to controls (da-GAL4>UAS-cat *v.* UAS-cat/+; p=1.7x10^{-8} females, p=8.1x10^{-3} males). Incubation at 90% O_2 was initiated at d7, with n=120 flies per condition (except n=90 for UAS-cat/+ females). All survival assays (**a,b,e,f,g,i**) were analysed by Log-Rank test (see Supplementary Table 1 for full n numbers and p values). n/s, p>0.05; *, p<0.05; **, p<0.01; ***, p<0.001.

Figure 2

Fig. 2 | Catalase lifespan extension requires autophagy. a, Catalase over-expressor (da-
GAL4>UAS-cat) and control (UAS-cat/+) females displayed a normal response to dietary restriction
(DR). Median lifespan is plotted against the yeast content of the diet, with 1x corresponding to
standard SYA food. Data are the means ±range of 2 independent lifespan experiments, each set up
with n=150-160 flies per genotype, analysed by two-way ANOVA (Fisher's LSD). **b,** Survival curve
on 0.1x-fold yeast from (**a**). The catalase over-expressor females (da-GAL4>UAS-cat) were longer-
lived than controls (UAS-cat/+) under low yeast nutritional conditions ($p=3.6 \times 10^{-10}$, n=160 flies per
genotype). **c,** Catalase over-expressor females (da-GAL4>UAS-cat) were sensitive to starvation
stress relative to UAS-cat/+ and da-GAL4/+ controls ($p=5.6 \times 10^{-5}$ and $p=7.6 \times 10^{-7}$, respectively). No
difference was observed in males (p>0.05 for all comparisons). Assays were performed at d7 with
n=120 flies per condition (except n=80 for UAS-cat/+ females). **d-e,** Triacylglyceride (TAG, **d**) and
glycogen (**e**) levels in whole females assayed at d7 (t=0) and depletion in response to starvation
treatment. Data are presented as box-and-whisker plots (min/max error bars) of n=4-6 replicates per
genotype, each with n=5 females per sample, analysed by unpaired Student's *t*-test (p>0.05). **f-g,**
Catalase over-expressor females (da-GAL4>UAS-cat) displayed enhanced autophagy induction
compared to UAS-cat/+ controls at d7. **f,** LysoTracker Red staining of midguts quantified as the
number of punctae relative to HOECHST-stained nuclei (scale bar = 10 μm). Data are means ±SD
of n=8 biological replicates, analysed by unpaired Student's *t*-test (***, p<0.001). **g,** Western
blotting against Atg8, normalised to actin. Data are means ±SD of n=6 biological replicates, each
with n=10 abdomens per sample, analysed by paired Student's *t*-test (*, p<0.05). Right, typical
bands probed against Atg8 with actin as a loading control (see Supplementary Fig. 4e for the full
blot). **h,** Global Atg5 knock-down did not decrease lifespan in a WT background under control
conditions (da-GS>UAS-Atg5RNAi ±RU; p=0.4177). Lifespan assays were performed on n=225-
240 females. **i,** Lifespan extension upon catalase over-expression (da-GS>UAS-cat ±RU; $p=7.2 \times 10^{-15}$)
was abolished in an autophagy-deficient background (da-GS>UAS-cat+UAS-Atg5RNAi ±RU;
p=0.1701). Survival assays (**b,c,h,i**) were analysed by Log-Rank test (see Supplementary Table 1
for full n numbers and p values). n/s, p>0.05; ***, p<0.001.

Figure 3

Fig. 3 | Catalase flies undergo an oxidising shift in global thiol redox state. a, Scheme showing the dual function of Atg4 in autophagy: 1) initial redox-independent cleavage of Atg8, to expose a C-terminal glycine residue enabling lipidation by PE (phosphatidyl-ethanolamine) via the E1-like enzyme Atg7 and the E2-like enzyme Atg3. The conjugated Atg8-PE is involved in autophagosome elongation/closure; 2) redox-dependent de-lipidation of Atg8-PE, allowing interaction and fusion of the autophagosome with the endosomal-lysosomal compartments, and recycling of cleaved Atg8. Oxidation of a redox-regulatory cysteine (C102 in *Drosophila* Atg4a) selectively inactivates the Atg8-PE de-conjugation activity of Atg4, promoting autophagosome biogenesis and therefore enhancing Atg4-mediated autophagy. **b,** Redox proteomic (OxICAT) analysis of d7, d28 and d56 catalase over-expressing females (da-GAL4>UAS-cat) compared to control (UAS-cat/+). Distribution of total cysteine residue oxidation levels, plotted as the proportion of the total number of peptides containing unique cysteine residues in each 5% quantile of percentage oxidation. Data are means ±SEM of n=5 biological replicates. **c-d,** Oxidation state of cysteine residues present, comparing control versus catalase over-expressor females at d7 (**c**) and d56 (**d**). Data points above the diagonal dotted line (slope=1) indicate cysteine residues more oxidised upon catalase up-regulation, with red symbols designating significance (p<0.05), assessed by unpaired two-tailed Student's *t*-test. The total number of unique Cys-containing peptides is indicated in black. **e,** Levels of Nrf2 signalling inferred using a gstD-GFP reporter in da-GS>UAS-cat females treated ±RU for d7 and d28. Western blotting against GFP, catalase and actin, showing 2 biological replicates per condition. **f**, Levels of total glutathione measured in d7 control (UAS-cat/+) and catalase over-expressor females. Data are means ±SEM of n=6 biological replicates, each with n=10 females per sample, analysed by unpaired Student's *t*-test (*, p>0.01 head+thorax; ***, p>0.001 abdomens).

Figure 4

Fig. 4 | Redox regulation of autophagy via Atg4a Cys102 extends lifespan. a, Structural representation of *Drosophila* Atg4a (brown, modelled with Phyre2[73]), superimposed on the crystal structure of human ATG4A (pink, PDB 2P82). Conservation of the catalytic triad and the regulatory cysteine (Cys102/81 in *Drosophila* and human, respectively) are highlighted. Structures were visualised in Chimera X[74]. **b,** Multiple sequence alignment of the Atg4a protein from *Drosophila* with the mouse and human orthologues, showing the catalytic cysteine residue (Cys98 in *Drosophila*) and the redox-regulatory cysteine (Cys102 in *Drosophila*). See Supplementary Fig. 4a for the full sequence. **c-d,** Autophagy induction in response to catalase over-expression was maintained in d7 Atg4a-WT female flies, but abolished in the Atg4a-C102S mutant background. **c,** Autophagy levels assessed by Western blotting against Atg8 (see Supplementary Fig. 4e for the full blot). Quantification of Atg8 levels by densitometry, normalised to actin as a loading control. Data are means ±range of n=2 biological replicates, analysed by two-way ANOVA (Fisher LSD). **d,** LysoTracker Red staining of midguts quantified as the number of punctae relative to HOECHST-stained nuclei (scale bar = 10 µm). Data are means ±SD of n=8 biological replicates, analysed by ANOVA (Tukey; n/s, $p > 0.05$; **, $p < 0.01$; ***, $p < 0.001$). **e-f,** Catalase over-expression extended lifespan in the Atg4a-WT control (**e**), but not in the redox-insensitive Atg4a-C102S knock-in background (**f**). Survival assays (**e**,**f**) were analysed by Log-Rank test (see Supplementary Table 1 for full n numbers and p values). n/s, $p > 0.05$; ***, $p < 0.001$. **g,** Graphical summary: Catalase up-regulation leads to the quenching of H_2O_2 as a second messenger molecule, which results in altered physiological redox signalling. Redox-responsive pathways, such as Nrf2, are not induced, causing an oxiding shift in bulk cysteine thiol redox state, which mimics an internal state of starvation. Autophagy is induced via redox-regulation of Cys102 in Atg4 as a protective response mechanism, which provides health and survival benefits.

Enhancing autophagy by redox regulation extends lifespan in *Drosophila*

Supplementary Data

Supplementary Figures 1-4

Supplementary Tables 1-5

- Supplementary Table 1: Summary of survival data
- Supplementary Table 2: List of cysteine residues identified by OxICAT redox proteomics
- Supplementary Table 3: Molecular function and pathway analysis of proteins identified in this study by OxICAT redox proteomics
- Supplementary Table 4: Oligonucleotide primers used for cloning and genotyping of the Atg4a-WT and Atg4a-C102S flies
- Supplementary Table 5: QPCR primers

Supplementary Figure 1

Supplementary Fig. 1 | Ubiquitous catalase up-regulation extends lifespan in flies, independently of oxidative stress resistance. a, Quantification of catalase over-expression by QPCR. Analysis of catalase mRNA levels normalised to actin in constitutive da-GAL4>UAS-cat females and males at d7, and inducible da-GS>UAS-cat +RU females at d9 (=d7 of RU treatment) compared to controls. Data are means ±SD of n=3-4 replicates, each with n=10 whole flies per sample, analysed by one-way ANOVA (Tukey). **b,** Representative Western Blot showing catalase protein abundance in female and male flies at d7. Data are means ±range of n=2 biological replicates, each with n=10 whole flies per sample, analysed by ANOVA (Tukey). **c,** Confocal microscopy of midgut sections showing catalase protein abundance by immunofluorescence (magenta) and nuclei (DAPI, blue) in control (UAS-cat/+) and catalase over-expressor (daGAL4>UAS-cat) females at d7 (scale bar = 100 μm). **d,** Catalase enzyme activity in whole female and male flies at d7. Data are means ±SEM of n=4 replicates per condition. n/s (p<0.05) by ANOVA (Tukey). **e-f,** Ubiquitous over-expression of mitochondria-targeted catalase (da-GAL>UAS-mito-cat) did not extend lifespan in females (p>0.05 *v.* UAS-mito-cat/+ and da-GAL4 controls) or males (p>0.05 for all comparisons). Survival assays were performed with n=150 flies per condition. **g,** Development time course. Constitutive ubiquitous catalase over-expression (da-GAL4>UAS-cat) causes a mild developmental delay relative to the UAS-cat/+ and da-GAL4/+ controls. Synchronised L1 larvae (n=500 per genotype) were transferred to SYA vials, and the subsequent eclosion time of adults was recorded. **h,** Dose response of da-GS>UAS-cat lifespan to a range of RU concentrations (0, 50, 200 and 400 μM) in both females and males. Survival assays were performed with n=240-255 flies per condition. **i,** Over-expression of catalase with a different ubiquitous driver, actin5c-GAL4, extended lifespan (act5c-GAL4>UAS-cat *v.* UAS-cat/+, p=3.6x10^{-12}). Survival assays were performed with n=200 females per condition. **j-k,** Lifespan extension by catalase was independent of *Wolbachia* status. Constitutive catalase over-expression (da-GAL4>UAS-cat) extended the lifespan of female flies in a w^{Dah+} background (c), but not males (d) relative to all controls (+/+, UAS-cat/+ and da-GAL4/+). Survival assays were performed with n=200 flies per condition. **l,** Late onset over-expression of catalase using the inducible GeneSwitch system from either middle-age (d28 and d42) and old-age (d56) extended the lifespan of female flies. Lifespans were performed with n=270 flies per condition. Original survival curves corresponding to Fig. 1e. **m,** Quantification of catalase protein levels by Western blotting with age and upon late-onset induction (see Fig. 1e). Flies (da-GS>UAS-cat) were sampled 7 d after the indicated age of RU treatment initiation. Left, typical Western blot probed against catalase with actin as a loading control. Right, quantification by densitometry. Data are means ±SD of n=3 biological replicates per time point, each with n=5 flies per sample, analysed by one-way ANOVA (Tukey). **n-s,** Tissue-specific over-expression of catalase: **n,** in the Malphigian tubules (the fly equivalent of the kidney) driven by Uro-GAL4 (n=300

females per condition, except n=285 for UAS-cat/+); **o,** in the insulin producing cells (IPCs) driven

by dilp2-GAL4 (n=300 females per condition, except n=285 for UAS-cat/+); **p,** pan-neuronal driven

by elav-GS (n=150 females per condition); **q,** in the intestine and fat body (the fly equivalent of the

liver/adipose tissue) driven by $S_1$106-GS (n=225 females per condition); **r,** in the gut driven by

TIGS2 (n=150 females per condition); **s,** pan-muscular driven by the myosin heavy chain driver

MHC-GS (n=150 females per condition). **t,** Catalase RNAi flies were sensitive to hyperoxia stress

relative to controls. Incubation at 90% O_2 was initiated at d9 (= d7 of RU treatment), and the fraction

alive scored after 4 d. Data are presented as box-and-whisker plots (min/max error bars) with n=7

vials per condition (each with n=15 flies), analysed by one-way ANOVA (Tukey) (da-GS/+ ±RU,

p=0.939; da-GS>UAS-cat ±RU, $p<1x10^{-15}$; da-GS>UAS-catRNAi ±RU, $p=7.01x10^{-6}$). All survival

assays (**e,f,h,i,j,k,l,m,n,o,p,q,r,s**) were analysed by Log-Rank test (see Supplementary Table 1 for

full n numbers and p values). n/s, p>0.05; ***, p<0.001.

Supplementary Figure 2

Supplementary Fig. 2 | Catalase lifespan extension requires autophagy. a, Response of egg laying to DR. Cumulative fecundity (egg laying at d7 + d14 + d21) for each genotype at a range of dietary yeast concentrations, with 1x corresponding to standard SYA food. Data are means ±SD of n=16 vials per genotype, each set up with n=10 females per vial, analysed by one-way ANOVA (Tukey, p<0.001 for all comparisons within each genotype). **b,** Whole body TAG levels in females assayed at d28 (t=0) and in response to a starvation timecourse. Data are presented as box-and-whisker plots (min/max error bars) of n=4-12 replicates per genotype, each with n=5 females per sample, analysed by ANOVA (n/s, p>0.05). **c-d,** Confocal imaging of d7 female midguts stained with LysoTracker Red and CytoID, as indicators of autophagolysosome formation and autophagic flux respectively (scale bar = 10 µm). **c,** Starvation (48 h) increased LytoTracker punctae in control (UAS-cat/+) females, while treatment with the inhibitor chloroquine (10 mM in SYA, 48 h) blocked autophagic flux, leading to increased CytoID signal. **d,** Autophagy is induced in catalase over-expressors (da-GAL4>UAS-cat) compared to controls (UAS-cat/+), without affecting flux. Images of individual channels, corresponding to Fig. 2f. **e,** Catalase up-regulation does not impact mitophagy, as inferred from the mito-QC reporter. Confocal images of dissected midguts showing mitolysosome punctae (scale bar = 5 µm). Data are the means ±SD of n=8-10 counts per genotype, analysed by unpaired Student's t-test (p>0.05). **f,** Catalase over-expressor females (da-GAL4>UAS-cat) were sensitive to treatment with the autophagy inhibitor chloroquine (10 mM in sucrose/agar medium) relative to UAS-cat/+ and da-GAL4/+ controls (p=2.6×10^{-5} and p=3.5×10^{-9}, respectively). Assays were performed at d7 with n=160 flies per condition. **g,** QPCR quantification of Atg5 and catalase levels upon Atg5 RNAi and catalase over-expression, normalised to actin. Female flies were sampled at d9 (=d7 of RU treatment), using RNA extracted from head and thorax tissue. Data are means ±SD of n=4-5 replicates per condition, each with n=10 flies per sample, analysed by one-way ANOVA (Tukey). **h,** Ubiquitous Atg5 RNAi results in sensitivity to starvation stress. The assay was set up at d7 with n=20-140 flies per genotype, using the GAL4 driver (rather than GeneSwitch) to maintain Atg5-RNAi during starvation stress where RU is not present (p<0.001 for da-GAL4>UAS-Atg5RNAi compared to all controls). All survival assays (**f,g**) were analysed by Log-Rank test (see Supplementary Table 1 for full n numbers and p values). n/s, p>0.05; ***, p<0.001.

Supplementary Figure 3

Supplementary Fig. 3 | Catalase flies undergo an oxidising shift in global thiol redox state. a, Scheme of the OxICAT workflow for unbiased redox proteomics. Cysteines are differentially labelled according to redox status using light and heavy ICAT probes. **b-e,** OxICAT analysis. Oxidation state of Cys residues present, comparing control (UAS-cat/+) *v.* catalase over-expressor (da-GAL4>UAS-cat) females at d28 (**a**), d7 *v.* d28 catalase over-expressors (**b**), d7 *v.* d56 catalase over-expressors (**c**), and d28 *v.* d56 catalase over-expressors (**d**). Data points above or below the diagonal dotted line (slope=1) indicate a change in Cys redox state, with red symbols designating low stringency significance (p<0.05, assessed by unpaired two-tailed Student's *t*-test) and blue symbols high stringency significance (Benjamini-Hochberg test). The total number of unique Cys-containing peptides is indicated in black. **f-g,** H_2O_2 levels inferred from the fluorescent redox-reactive probe CellROX assessed by confocal microscopy of dissected midguts (scale bar = 50 µm). Data are means ±SD of n=3 samples per condition. **f,** Response of control tissue incubated ±H_2O_2 as a positive control for the assay (p=0.0044). **g,** CellROX staining of catalase up-regulated guts, showing decreased signal compared to control (p=0.003564). **h,** Mitochondrial H_2O_2 levels quantified using the MitoB probe, showing similar levels in control and catalase up-regulated flies. Data are means ±SEM of n=6-7 samples per genotype (p=0.5045). Data (**f-h**) are for d7 females, analysed by unpaired 2-tailed Student's *t*-test (n/s, p>0.05; **, p<0.01).

Supplementary Fig. 4 | Redox regulation of autophagy via Atg4 Cys102 extends lifespan. a, Multiple sequence alignment of Atg4a from *Drosophila* with the mouse and human orthologues (UniProt IDs # Q9VPW2, Q8C9S8 and Q8WYN0, respectively) performed in T-Coffee (www.tcoffee.org). **b,** Atg4a expression levels were not affected by the C102S mutation or catalase over-expression, as assessed by QPRC, normalised to tubulin. Data are the means ±SD of n=4 replicates, analysed by ANOVA (Tukey, $p > 0.05$ for all comparisons). **c,** The Atg4a-C102S mutation did not change levels of catalase up-regulation, as measured by enzyme activity. Data are the means ±SD of n=3 replicates, analysed by ANOVA (Tukey, *** $p < 0.001$, n/s $p > 0.05$). **d,** H_2O_2 stress resistance was comparable between Atg4a-WT flies and Atg4a-C102S knock-in mutants, both under control conditions and upon catalase up-regulation. H_2O_2 treatment (5% v/v in sucrose/agar medium) was initiated at d7, with n=160 females per genotype. Survival assays were analysed by Log-Rank test (see Supplementary Table 1 for full n numbers and p values). n/s, $p > 0.05$, ***, $p < 0.001$. **e,** Full Western blots corresponding to data in Figs. 2g and 4c. After transfer, the upper portion of the membrane was probed for actin as a loading control, while the lower portion of the membrane was probed against Atg8. **f,** Autophagy induction in response to catalase over-expression was maintained in d7 Atg4a-WT female flies, but abolished in the Atg4a-C102S mutant background, as assessed by LysoTracker Red staining of midguts (scale bar = 10 μm). Images of individual channels, corresponding to Fig. 4d. **g,** Climbing assay of control (da-GAL/+) and catalase over-expressors (da-GAL>UAS-cat) in Atg4a-WT and Atg4a-C102S backgrounds. Data are presented as box-and-whisker plots (min/max error bars) of n=5 replicates per condition, each with n=15 flies per sample, analysed by ANOVA (n/s, $p > 0.05$; *, $p < 0.05$; **, $p < 0.01$). **h,** The Atg4a-C102S mutation does not alter egg laying upon catalase up-regulation compared to the control Atg4a-WT background in d7 females. Data are the means ±SEM of n=7 vials per genotype, each set up with n=15 females per vial, analysed by one-way ANOVA (Tukey, n/s, $p > 0.05$; ***, $p < 0.001$).