Classification: BIOLOGICAL SCIENCES, cell biology

Title: Interplay between Desmoglein2 and hypoxia controls metastasis in breast cancer

Author Affiliation: Po-Hao Chang[1], Min-Che Chen[2], Ya-Ping Tsai[2], Pang-Hung Hsu[3], Yung-Ming Jeng[4], Yi-Fang Tsai[5], Muh-Hwa Yang[6], Wendy W. Hwang-Verslues[1,*]

[1]Genomics Research Center, Academia Sinica, No. 128, Sec. 2, Academia Road, Taipei 115, Taiwan

[2]Asclepiumm Taiwan Co., Ltd, Mackay Memorial Hospital Innovation & Incubation Center, No. 45, Minsheng Rd, Tamsui, New Taipei City 25160, Taiwan

[3]Department of Bioscience and Biotechnology, National Taiwan Ocean University, No.2, Beining Rd., Jhongjheng District, Keelung City 20224, Taiwan

[4]Department of Pathology, National Taiwan University Hospital, No.7, Chung Shan S. Rd., Zhongzheng Dist., Taipei 100, Taiwan

[5]Department of Surgery, Taipei-Veterans General Hospital, No. 201, Section 2, Shipai Road, Beitou District, Taipei City 112, Taiwan

[6]Institute of Clinical Medicine, National Yang-Ming University, No.155, Sec.2, Linong Street, Taipei 112, Taiwan

Corresponding Author*: Wendy W. Hwang-Verslues, Ph.D., Genomics Research Center, Academia Sinica, No. 128, Sec. 2, Academia Road, Taipei 115, Taiwan.   **Email:**

wendyhv@gate.sinica.edu.tw, **Phone:** +886-2-2787-1246, **Fax:** +886-2-2789-9924, **ORCID:** 0000-

0002-0383-1710

Keywords: Desmoglein2 (DSG2), breast cancer, hypoxia, HIF1α, circulating tumor cells (CTCs),

metastasis

Abstract

Metastasis is the major cause of cancer death. Metastatic cancer cells that have intravasated into the circulatory system and formed clusters of circulating tumor cells (CTCs) are particularly associated with colonization of distant organs and poor prognosis. However, the key factors required for tumor cell dissemination and CTC clustering remain elusive. We found that high expression of Desmoglein2 (DSG2), a cell adhesion molecule involved in desmosome mediated intercellular adhesion, promoted tumor growth, increased the prevalence of CTC clusters and facilitated distant organ colonization. The dynamic regulation of DSG2 by hypoxia was key to this process as downregulation of DSG2 in hypoxic regions of primary tumors led to elevated epithelial-mesenchymal transition (EMT) gene expression allowing cells to detach from the primary tumor and undergo intravasation. Subsequent derepression of DSG2 after intravasation and release of hypoxic stress allowed CTCs to colonize distant organs. This dynamic regulation of DSG2 was mediated by Hypoxia-Induced Factor1α (HIF1α). In contrast to its more widely observed function to promote expression of hypoxia-inducible genes, HIF1α repressed DSG2 by recruitment of the Polycomb Repressive Complex 2 components, EZH2 and SUZ12, to the DSG2 promoter in hypoxic cells. Consistent with our experimental data, DSG2 expression level correlated with poor prognosis and recurrence risk in breast cancer patients. Together, these results demonstrated the importance of DSG2 expression in metastasis and revealed a new mechanism by which hypoxia drives metastasis.

Significant Statement

During metastasis, a hypoxic microenvironment is a major force driving primary tumor cells to disseminate into the circulatory system. The majority of these circulating tumor cells (CTCs) cannot survive when traveling alone. However, collective movement as CTC clusters enables them to avoid immune surveillance and increases the probability that they will successfully metastasize to distal organs. Dynamic down regulation of DSG2 in hypoxic tumor tissue and reactivation of expression in CTCs allowed high rates of tumor cell dissemination, CTC cluster formation and metastasis that are characteristics of the most aggressive breast cancers. The findings highlight both the potential, and potential pitfalls, of using DSG2 as a therapeutic target.

Introduction

Breast cancer is the most common cancer of women worldwide. With early detection and advance in therapeutic strategies, the 5-year relative survival rate for all stages combined is higher than 90% (according to the SEER database maintained by the U.S. National Cancer Institute). However, metastasis is now the major cause of breast cancer death (1) as the survival rate for women with metastasized breast cancers remains below 30%. Thus, identification of key factors in breast tumorigenesis and metastasis is important to identify therapeutic targets and strategies to improve prognosis.

Metastasis is a complex process involving tumor cell intrinsic alterations and extrinsic interaction with the microenvironment to select for highly aggressive cancer cells. Hypoxia, a key microenvironmental factor in solid tumors, activates hypoxic signaling to increase plasticity and promote epithelial-mesenchymal transition (EMT) to drive the first step of metastasis (2). High plasticity allows cancer cells to disseminate from the primary site and intravasate into the circulatory or lymphatic system. Most of these circulating tumor cells (CTCs) die in circulation and only a small fraction of CTCs are able to survive and eventually colonize distant organs (3, 4). Recent evidence has shown that CTC numbers can be used as an independent predictor for survival in patients with metastatic cancers (5-7). Further improvement in CTC detection methods led to identification of

CTC clusters and the finding that clustered CTCs exhibit epithelial/mesenchymal hybrid (partial

EMT) phenotype which allows them to move collectively (8). Collective movement makes these

cancer cells more apoptosis-resistant, more capable of avoiding immune surveillance and, better able

to colonize distant organs. Importantly, CTC clustering ability has been positively correlated with

poor clinical outcome (9-12). While a few factors that promote CTC cluster formation have been

identified (9, 13-15), the key mechanisms that allow CTC clusters to survive in the vascular system

and allow them to more efficiently metastasis than unclustered CTCs remain elusive.

   Cell adhesion proteins play critical roles in intercellular contacts and epithelial tissue dynamics.

Deregulation of cell adhesion molecules contributes to tumor metastasis (16, 17). Among cell

adhesion molecules, desmosomes are of particular interest for cancer biology. Desmosomes form

patch-like adhesion structures that mark the intercellular midline and connect to the intermediate

filament cytoskeleton to maintain cell-cell adhesion and tissue integrity (18). The desmosome is a

protein complex containing two transmembrane proteins, Desmocollin (DSC1-DSC3) and

Desmoglein (DSG1-DSG4), as well as adaptor proteins, Plakoglobin and Desmoplakin, that bind

intermediate filaments (19). Among the DSGs, DSG2 is the most ubiquitously expressed isoform,

including mammary tissue, and is a key factor for cell aggregation and oncogenic function in lung

and prostate cancers (20-23). However, whether DSG2 is involved in CTC clustering and metastasis

remains unknown.

We found that tumor growth and colonization were promoted by DSG2 expression at both the primary site and distant organ. However, HIF1α−mediated suppression of DSG2 under hypoxia was required for cancer cell invasion. Once in the vascular system, the cancer cells were released from hypoxic stress and DSG2 expression was de-repressed. This DSG2 reactivation was associated with increased prevalence of CTC clusters and was essential for CTCs to colonize distant organs. Consistent with these experimental observations, clinical data indicated that breast cancer patients whose tumors expressed DSG2 (DSG2-positive) had worse prognosis and higher recurrence risk than those with DSG2-negative tumors. Together, these results show that dynamic changes of DSG2 expression are required for breast tumorigenesis, invasion, CTC clustering and metastasis. Our data also identify regulatory mechanisms underlying DSG2 repression and de-repression during specific stages of breast cancer progression.

Results

High expression of DSG2 in breast cancer is associated with poor prognosis and high recurrence

To identify key factors driving CTC clustering and metastasis, we analyzed a transcriptome dataset for metastatic and non-metastatic breast cancers from the Gene Expression Omnibus (GEO)

database using Gene Set Enrichment Analysis (GSEA). Forty-six genes associated with metastatic

disease were enriched in metastatic breast cancers. Among these, DSG2 was the most highly

expressed (Fig. 1A, 1B). Because formation of CTC clusters (groups of two or more aggregated

CTCs) promotes metastasis and because cell-cell interaction is critical for CTC clustering (9), we

also analyzed genes associated with cell-cell junctions and found DSG2 as the second most highly

expressed gene among 56 genes enriched in the metastatic group (*SI Appendix*, Fig. S1A, S1B).

Overlap of these two gene sets (metastasis and cell-cell junction) identified three genes (*SI Appendix*,

Fig. S1C). Among these, DSG2 was the most highly upregulated in the metastasis and cell-cell

junction gene sets that was also enriched in metastatic breast cancer patients.

We then evaluated clinical correlation of DSG2 expression with prognosis using

immunohistochemistry (IHC) analysis of 164 biopsy specimens (Fig. 1C). Based on the IHC

analysis, patients were grouped into "DSG2-positive" and "DSG2-negative" groups for Kaplan–

Meier survival analyses. The disease-free survival was significantly lower in the DSG2-positive

group (Fig. 1D, $p = 0.004$). Importantly, DSG2 expression was highly correlated with cancer

recurrence (Fig. 1E, $p = 0.03$). To examine whether high DSG2 expression promoted metastasis, we

analyzed a breast cancer cohort from the UCSC Xena public hub (Yau 2010 dataset)(24) and found

significantly lower distant metastasis-free survival in DSG2-high breast cancer patients (Fig. 1F, $p =$

0.008). The results from these three independent cohorts indicated that DSG2 possesses a metastatic

role in breast cancer and could be used as a prognostic marker.

DSG2 expression is required for CTC clustering and metastasis

To determine DSG2 function in breast cancer progression, we first examined endogenous DSG2

levels in a panel of cell lines (*SI Appendix*, Fig. S2A). MB231, a model cell line for *in vivo*

metastasis studies due to its high metastatic potential (25), had the highest level of DSG2 out of eight

cell lines assayed. Suppression of DSG2 expression in MB231 using a lentiviral-shRNA system (*SI*

Appendix, Fig. S2B) led to depletion of DSG2 from the cell membrane, as confirmed by fluorescence

activated cell sorting (FACS) (*SI Appendix*, Fig. S2C). These DSG2-depleted cells were then used to

determine the role of DSG2 in CTC clustering and metastasis by orthotopic xenograft assays where

EGFP expressing MB231 cells, without (shCtrl) or with (shDSG2) DSG2 depletion, were injected

into the 4^{th} mammary fat-pad of NOD/SCIDγ mice. Nine weeks after mammary fat pad injection,

tumor size, CTC counts and lung nodules were evaluated (Fig. 2A). Mice bearing shCtrl cell-derived

tumors had more metastatic nodules in the lungs compared to the mice bearing shDSG2 cell-derived

tumors (Fig. 2B, right panel). Moreover, IHC staining confirmed that the cells which colonized and

formed nodules in the lungs expressed DSG2 (Fig. 2B, left panel, *SI Appendix*, Fig. S2D).

Interestingly, we observed a 5-fold higher prevalence of CTC clusters in the shCtrl group (Fig. 2C,

2D, black bars) than the shDSG2 group (278 clusters per gram tumor in the shCtrl group versus 59

clusters per gram tumor in the shDSG2 group). No significant difference in single CTC numbers

were found between shCtrl and shDSG2 group (Fig. 2C, 2D, white bars).

A syngeneic tumor mouse model was used to confirm that similar DSG2-driven tumorigenic

phenotypes also occurred in immune-competent mice. Mouse mammary tumor cells labeled with

GFP-luciferase (4T1-GFP/LUC) with or without DSG2 depletion were injected into the 4th

mammary fat-pad of BALB/c mice (Fig. 2E). Consistent with the MB231 xenograft model, shCtrl-

injected syngeneic mice had more lung metastasis than shDSG2-injected mice (Fig. 2F). Importantly,

blood of shCtrl-injected mice also had significantly higher levels of both single CTCs and CTC

clusters than blood from the shDSG2 group (Fig. 2G, 2H). This occurred despite the fact that there

was no significant difference in primary tumor size between these two groups (*SI Appendix*, Fig.

S2E). Together, these data indicated that DSG2 facilitates breast cancer metastasis and also promotes

CTC clustering.

DSG2 expression promotes metastatic colonization and tumor growth

As colonization is a crucial step to determine whether cancer cells can survive in distant organs

(26, 27), we injected SKBR3 cells, which also had high levels of endogenous DSG2 expression (*SI*

Appendix, Fig. S2A), with or without DSG2 depletion into mouse tail vein to examine whether

DSG2 contributes to colonization ability. Fixation and hematoxylin and eosin (H&E) staining of lung tissue two months after the initial tail vein injection showed that mice injected with DSG2-depleted SKBR3 cells had significantly lower numbers of lung nodules compared to mice injected with control cells (Fig. 3A). Conversely, DSG2 overexpression in MB157 cells, which had low levels of endogenous DSG2 expression (*SI Appendix*, Fig. S2A), significantly increased lung colonization (Fig. 3B).

In addition, DSG2 promoted tumor growth at the primary site. DSG2 depletion led to significant reduction of colony forming efficiency in soft agar colony forming (SACF) assays (Fig. 3C). Conversely, DSG2 overexpression in MB157 and MB468 cells, which had low endogenous DSG2 levels (*SI Appendix*, Fig. S2A-2C), resulted in higher colony numbers (Fig. 3D). Consistent with the tumorigenic role of DSG2 in promoting colony formation, orthotopic xenografts showed that tumors derived from DSG2-depleted cancer cells were significantly smaller than the control while cells overexpressing DSG2 formed significantly larger tumors (Fig. 3E, 3F, and *SI Appendix*, Fig. S2F, S2G). Together, these results demonstrated that DSG2 expression promotes breast tumor growth at the primary site and increases colonization in the distal organs.

Repression of DSG2 promotes cell invasion, migration and expression of EMT genes

Initiation of cancer metastasis, including invasion, migration and intravasation, relies heavily on

tumor cell dissemination (1, 28). Since DSG2 expression increased CTC clusters and tumor growth

in the mammary fat pad and also promoted colonization in the lung, we next evaluated its role in

dissemination. Seemingly in contrast to its tumorigenic role, DSG2 knockdown in MB231 cells

significantly increased migration ability (*SI Appendix*, Fig. S3A) and DSG2 overexpression in

MB157 cells inhibited migration (*SI Appendix*, Fig. S3B). Similarly, depletion of DSG2 increased

cell invasion; whereas overexpression of DSG2 inhibited this ability (*SI Appendix*, Fig. S3C, S3D).

Consistent with these observations, DSG2 depletion resulted in an upregulation of EMT genes,

including *SNAIL*, *SLUG* and *VIMENTIN (VIM)*, all of which are critical for cancer cell mobility and

dissemination (29, 30). Conversely, DSG2 overexpression downregulated these genes (*SI Appendix*,

Fig. S3E, S3F). Thus, DSG2 suppression may facilitate invasion and migration not only by

disrupting cell-cell adhesion but also by upregulating EMT genes to allow cancer cells to detach

from the primary tumor and intravasate into the circulatory system as single cells.

Hypoxia-induced HIF1α suppresses DSG2 expression

Since both single CTCs and CTC clusters were detected in mice with DSG2-expressing tumors

(Fig. 2), DSG2 must be suppressed in some parts of the primary tumors to allow cancer cells to

invade and disseminate into the blood. To examine this hypothesis, we performed IHC using breast

cancer patient samples. A negative correlation of expression pattern between DSG2 and CA9, a

hypoxic stress marker, across different tumor regions was observed (Fig. 4; *SI Appendix*, Fig. S4).

DSG2 expression was low in hypoxic regions with high CAIX expression (Fig. 4, red box), but high

in regions without CAIX staining (Fig. 4, blue box). These results indicated that DSG2 expression

may be downregulated under hypoxia.

Hypoxia plays a critical role in metastasis (2) but it is not known whether hypoxia affects DSG2

expression. Consistent with the pattern of DSG2 expression in tumor specimens, breast cancer cells

subjected to hypoxic stress had decreased DSG2 gene expression and protein level (Fig. 5A, 5B).

Similarly, DSG2 expression decreased in cells treated with different doses or durations of the

hypoxia-mimetic agent $CoCl_2$ (Fig. 5C; *SI Appendix*, Fig. S5A). Since HIF1α is a critical mediator

of hypoxia-associated cancer progression (31), we determined the effect of HIF1α on DSG2

expression. Transient transfection of a non-degradable HIF1α (P564A) in 293T cells resulted in

downregulation of DSG2 (Fig. 5D). Importantly, depletion of HIF1α in 293T treated with $CoCl_2$

abolished hypoxia-induced DSG2 suppression (Fig. 5E).   These results indicated that HIF1α is

responsible for DSG2 suppression under hypoxic stress.

To further elucidate whether HIF1α directly suppressed DSG2 transcription, we used Integrated

Motif Activity Response Analysis (ISMAR) (32) to identify putative HIF1α binding sites and found

that mutation of one of these sites (-946 to -943) abolished HIF1α inhibition of DSG2 expression

(Fig. 5F). Chromatin-immunoprecipitation (ChIP) further confirmed that hypoxia-induced HIF1α

bound to this region and this HIF1α binding was associated with increased H3K27-trimethylation

(H3K27me3), indicative of heterochromatin formation, during hypoxia (Fig. 5G, 5H; *SI Appendix*,

Fig. S5B).

This observation of HIF1α-mediated repression of DSG2 contrasted with the more typical

function of HIF1α to promote expression of hypoxia-inducible genes (33). To understand how

HIF1α carried out this transcriptional suppression, we performed Co-IP assays and found stabilized

HIF1α interaction with polycomb repressive complex 2 components, EZH2 and SUZ12, but not

HDAC1 and HDAC2, in breast cancer cells under hypoxia (Fig. 6A, *SI Appendix*, Fig. S5C). This

interaction was also confirmed using 293T cells ectopically expressing the non-degradable HIF1α

(P564A) (Fig. 6B). Importantly, depletion of EZH2 or SUZ12 significantly abolished hypoxia-

induced DSG2 suppression (Fig. 6C) indicating that these PRC2 components are required for DSG2

downregulation under hypoxic stress. Together, these results suggested that HIF1α may recruit

EZH2 and SUZ12 to the DSG2 promoter to downregulate DSG2 expression under hypoxia.

To further investigate whether EZH2 and SUZ12 were recruited to the DSG2 promoter by

HIF1α under hypoxic stress, we performed ChIP assays using cells with or without HIF1α depletion

(Fig. 6D). Upon $CoCl_2$ treatment, HIF1α was induced and recruited these co-repressors to the

HIF1α-binding region on the DSG2 promoter in the shCtrl cells (Fig. 6E, gray bar). However, the

recruitment of EZH2 or SUZ12 was abolished in HIF1α depleted cells under hypoxic stress (Fig. 6E,

blue bar). These results demonstrated that stabilized HIF1α is required to recruit PRC2 complex to

the DSG2 promoter to suppress DSG2 transcription under hypoxia.

Since DSG2 is critical for distant organ colonization and is highly expressed in metastatic

nodules (Fig. 2 and Fig. 3), DSG2 suppression that occurs in hypoxic conditions must be released

before colonization can occur. Consistent with this idea, de-repression of DSG2 expression was

observed within one hour after breast cancer cells were released from hypoxia *in vitro* (Fig. 6F). To

confirm that this reactivation is important *in vivo*, we injected EGFP expressing MB231 cells into

mammary fat pads and evaluated DSG2 expression in CTCs after injection. Before injection, nearly

all cells expressed DSG2 (98.6%, Fig. 6G, left panel). At nine weeks after injection, more than 90%

of CTCs disseminated from the primary tumor site expressed DSG2 while 9% had low/non-

detectable DSG2. Given our other observations, it seems likely that the cells with low or non-

detectable DSG2 had just been released from the primary tumor and undergone intravasation (Fig.

6G, right panel). Taken together, our *in vitro* and *in vivo* observations are all consistent with dynamic

regulation of DSG2 where hypoxia-mediated DSG2 suppression followed by de-repression of DSG2

when hypoxic stress is released in the circulatory system is required for DSG2-promoted breast

cancer metastasis.

Discussion

Our study demonstrated that dynamic changes in DSG2 expression are important for breast

tumorigenesis and malignancy (Fig. 7). Expression of DSG2 not only facilitated breast tumor growth

in mammary tissue and colonization in lung, it also promoted CTC collective migration in the

circulatory system (Fig. 2, 3). In hypoxic tumor regions, HIF1α-mediated DSG2 suppression was

critical for cancer cells to gain mobility, detach from the primary tumor and intravasate into the

blood (Fig. 5; *SI Appendix*, Fig. S3).

Recent research has proposed two mechanisms describing how CTCs form clusters to

collectively migrate in the circulatory system. One is that tumor cells intravasate as multicellular

clusters via plakoglobin-dependent intercellular adhesion into the blood (9). The other is that

individual tumor cells aggregate via CD44-PAK2 mediated FAK signaling after intravasation (15).

Our observations of dynamic, hypoxia-dependent regulation of DSG2 and its effects on CTC

prevalence suggest that DSG2 could participate in both mechanisms. Since DSG2 is an integral

component of desmosomes and its cytoplasmic tail binds to plakoglobin and plakophilins (34),

DSG2 may facilitate CTC clustering prior to intravasation through its cell-cell adhesion function.

Therefore, non-hypoxic tumor regions where DSG2 expression is high may shed CTC clusters at low

frequency. In contrast, hypoxic tumor regions where DSG2 is downregulated may more readily

generate single CTCs and rapid de-repression of DSG2 after intravasation (Fig. 6) allows these

single CTCs to aggregate. Once CTC clusters are in the circulatory system, high DSG2 expression

may allow CTC clusters to remain intact and persist longer in the circulatory system. This itself may

give more chance for colonization. DSG2 may also more directly promote colonization and

metastatic tumor growth via its roles in cell adhesion.

CTCs that can survive in the circulatory system and colonize in distant organs are thought to be

cancer stem-like cells and are highly associated with metastasis (35, 36). We also observed that

DSG2 expression promoted mammosphere formation and survival in a low-attached culture system

(*SI Appendix*, Fig. S6), suggesting that DSG2 may contribute to CTC survival during circulation and

may help maintain cancer stemness. Consistent with this notion, DSG2 has recently been identified

as a marker for pluripotent stem cells (PSCs) and is critical for PSC self-renewal and reprogramming

(37). How DSG2 may contribute to cancer stemness is a topic of interest for further experiments.

In several cancers, HIF1α acts as a transcription activator to upregulate genes required for

survival, EMT, angiogenesis and metastasis upon hypoxic stress (31, 33). HIF1α upregulates these

target genes by recruiting coactivators such as p300/CBP acetyl-transferase, CDK8-mediator, Potin

and SWI/SNF chromatin remodelers (38). In contrast, there is relatively little evidence that HIF1α

can also act as a transcription repressor. Under hypoxia, stabilized HIF1α was found to displace the

transcription activator Myc from Sp1 binding to repress MutSα expression to contribute to genomic

instability in colon cancer cells (39). However, what corepressors HIF1α recruited to the target

promoter upon hypoxia was not known. Our observation that HIF1α recruited EZH2 and SUZ12 to

suppress DSG2 transcription (Fig. 5, 6) provides a new regulatory mechanism and expands the

oncogenic role of HIF1α in promoting breast cancer progression. Whether this HIF1α-EZH2-SUZ12

complex serves as a common gene suppression mechanism to facilitate cancer malignancy in other

cancer types is worthy of further investigation.

Our clinical data and animal models clearly demonstrated that DSG2 can be used as a

prognostic marker for breast cancer. Interestingly, DSG2 expression is also associated with poor

prognosis in other cancers including cervical, head and neck and lung cancers (*SI Appendix*, Fig. S7).

This suggests a widespread role of DSG2 in cancer malignancy. As a transmembrane protein, DSG2

may be a drugable target. However, the strategy has to be carefully designed. Using a monoclonal

DSG2 specific antibody to target the DSG2 to inhibit anchorage independent growth may be

possible. However, antibody-DSG2 internalization may occur and result in enhancement of

intravasation, as downregulation of DSG2 promotes invasion. In this case, a well-designed antibody-

cytotoxic drug conjugate (ADC) (40) may be a more effective therapeutic method to target DSG2

expressing cancer cells. In addition, since DSG2 expressing CTCs tend to form clusters which

facilitate distant organ colonization, dialysis for DSG2 expressing cancer cells in the blood may be

another way to inhibit metastasis. Alternatively, DSG2 may be used as a specific marker for CTC

cluster detection and isolation. Overall, the underlying molecular mechanisms of DSG2-mediated

tumorigenesis and the possibility of using DSG2 as a cancer therapeutic target are promising areas

for further work.

Materials and Methods

Breast cancer specimens for IHC analysis were collected from National Taiwan University Hospital. All specimens were encoded to protect patients under protocols approved by the Institutional Review Board of Human Subjects Research Ethics Committee of Academia Sinica (AS-IRB01-16031) and National Taiwan University (201605057RINA), Taipei, Taiwan. Animal care and experiments were approved by the Institutional Animal Care and Utilization Committee of Academia Sinica (IACUC# 15-11-885). NOD/SCID mice were kindly provided by Dr. Michael Hsiao (Genomics Research Center, Academia Sinica, Taiwan) and BALB/c mice were purchased from the National Laboratory Animal Center (Taiwan). Cancer cell lines including MB231, MB157, MB468, SKBR3 and 4T1 were obtained from the American Type Culture Collection. Human breast cancer cell lines were maintained in DMEM and mouse breast cancer cell line 4T1 was maintained in RPMI supplemented with 10% fetal bovine serum and antibiotics and cultured at 37°C in a humidified incubator supplemented with 5% CO_2. For hypoxia experiments, medium was first incubated in HypoxyCOOL system (Baker) for 8 hours to reduce oxygen level to 1%. This medium was then used to culture the cells in INVIVO 400 hypoxia chamber (Baker) supplemented with 5% CO_2 and 1% O_2 at 37°C.

Single CTC and clusters were identified using a CTC platform from MiCareo, Inc. A spike-in control experiment using 10^3 4T1-GFP/LUC cells mixed with 2ml blood from wildtype BALB/c mice was

performed to determine whether the centrifugation step of CTC isolation increases cell aggregation

(Fig S8). Further details of experimental methods are given in the *SI Appendix*, Materials and

Methods.

Author contributions

P.-H.C. and W.W.H.-V. conceived the study. P.-H.C. performed all experiments not attributed to

other authors. P.-H.H. assisted with the LC-MS/MS analysis in the original manuscript and Y.-M.J.

assisted with immunohistochemistry analysis, P.-H.C. and P.-H.H. analyzed the data, P.-H.C., M.-

C.C. and Y.-P.T. generated the DSG2 antibody, Y.-M.J., Y.-F.T. and M.-H.Y. provided clinical

samples. P.-H.C. and W.W.H.-V. wrote the paper.

Author competing interest

P.-H.H., Y.-M.J., Y.-F. T., M.-H. Y. and W.W.H.-V. declare no competing interests. P.-H.C. and M.-

C.C. are co-inventors on a DSG2 monoclonal antibody patent owned by Asclepiumm Taiwan Co.,

Ltd.   M.-C.C. has equity ownership in and serves on the board of director of Asclepiumm Taiwan

Co., Ltd.

Acknowledgements

This work was supported by Academia Sinica [AS-SUMMIT-108, AS-SUMMIT-109] and the

Taiwan Ministry of Science and Technology [MOST 109-0210-01-18-02, MOST 108-3114-Y-001-

002, MOST 107-0210-01-19-01, MOST 105-2628-B-001-008-MY3 and MOST-108-2311-B-001-

005-MY3]. The authors would like to thank Dr. Michael Hsiao (Genomics Research Center,

Academia Sinica, Taiwan) for providing NOD/SCID mice, and Dr Yi-Cheng Chang (Graduate

Institute of Medical Genomics and Proteomics, National Taiwan University) for discussion.

Reference

1.   Chaffer CL & Weinberg RA (2011) A perspective on cancer cell metastasis. *Science* 331(6024):1559-1564.

2.   Rankin EB & Giaccia AJ (2016) Hypoxic control of metastasis. *Science* 352(6282):175-180.

3.   Twomey JD & Zhang B (2019) Circulating Tumor Cells Develop Resistance to TRAIL-Induced Apoptosis Through Autophagic Removal of Death Receptor 5: Evidence from an In Vitro Model. *Cancers (Basel)* 11(1).

4.   Theodoropoulos PA*, et al.* (2010) Circulating tumor cells with a putative stem cell phenotype in peripheral blood of patients with breast cancer. *Cancer Lett* 288(1):99-106.

5.   Cohen SJ*, et al.* (2008) Relationship of circulating tumor cells to tumor response, progression-free survival, and overall survival in patients with metastatic colorectal cancer. *J Clin Oncol* 26(19):3213-3221.

6.   Cristofanilli M*, et al.* (2004) Circulating tumor cells, disease progression, and survival in metastatic breast cancer. *N Engl J Med* 351(8):781-791.

7.   de Bono JS*, et al.* (2008) Circulating tumor cells predict survival benefit from treatment in metastatic castration-resistant prostate cancer. *Clin Cancer Res* 14(19):6302-6309.

8.   Liu X*, et al.* (2019) Epithelial-type systemic breast carcinoma cells with a restricted mesenchymal transition are a major source of metastasis. *Sci Adv* 5(6):eaav4275.

9.   Aceto N*, et al.* (2014) Circulating tumor cell clusters are oligoclonal precursors of breast cancer metastasis. *Cell* 158(5):1110-1122.

10.   Cho EH*, et al.* (2012) Characterization of circulating tumor cell aggregates identified in patients with epithelial tumors. *Phys Biol* 9(1):016001.

11.   Hou JM*, et al.* (2012) Clinical significance and molecular characteristics of circulating tumor cells and circulating tumor microemboli in patients with small-cell lung cancer. *J Clin Oncol* 30(5):525-532.

12.   Mu Z*, et al.* (2015) Prospective assessment of the prognostic value of circulating tumor cells and their clusters in patients with advanced-stage breast cancer. *Breast Cancer Res Treat* 154(3):563-571.

13.   Szczerba BM*, et al.* (2019) Neutrophils escort circulating tumour cells to enable cell cycle progression. *Nature* 566(7745):553-557.

14.   Gkountela S*, et al.* (2019) Circulating Tumor Cell Clustering Shapes DNA Methylation to Enable Metastasis Seeding. *Cell* 176(1-2):98-112 e114.

15.   Liu X*, et al.* (2019) Homophilic CD44 Interactions Mediate Tumor Cell Aggregation and Polyclonal Metastasis in Patient-Derived Breast Cancer Models. *Cancer Discov* 9(1):96-113.

16. Eger A, Stockinger A, Schaffhauser B, Beug H, & Foisner R (2000) Epithelial mesenchymal transition by c-Fos estrogen receptor activation involves nuclear translocation of beta-catenin and upregulation of beta-catenin/lymphoid enhancer binding factor-1 transcriptional activity. *J Cell Biol* 148(1):173-188.

17. Onder TT*, et al.* (2008) Loss of E-cadherin promotes metastasis via multiple downstream transcriptional pathways. *Cancer Res* 68(10):3645-3654.

18. Garrod D & Chidgey M (2008) Desmosome structure, composition and function. *Biochim Biophys Acta* 1778(3):572-587.

19. Delva E, Tucker DK, & Kowalczyk AP (2009) The desmosome. *Cold Spring Harb Perspect Biol* 1(2):a002543.

20. Schafer S, Koch PJ, & Franke WW (1994) Identification of the ubiquitous human desmoglein, Dsg2, and the expression catalogue of the desmoglein subfamily of desmosomal cadherins. *Exp Cell Res* 211(2):391-399.

21. Cai F*, et al.* (2017) Desmoglein-2 is overexpressed in non-small cell lung cancer tissues and its knockdown suppresses NSCLC growth by regulation of p27 and CDK2. *J Cancer Res Clin Oncol* 143(1):59-69.

22. Barber AG*, et al.* (2014) Characterization of desmoglein expression in the normal prostatic gland. Desmoglein 2 is an independent prognostic factor for aggressive prostate cancer. *PLoS One* 9(6):e98786.

23. Saias L, Gomes A, Cazales M, Ducommun B, & Lobjois V (2015) Cell-Cell Adhesion and Cytoskeleton Tension Oppose Each Other in Regulating Tumor Cell Aggregation. *Cancer Res* 75(12):2426-2433.

24. Mary Goldman BC, Mim Hastie, Kristupas Repečka, Akhil Kamath, Fran McDade, Dave Rogers, Angela N Brooks, Jingchun Zhu, David Haussler (The UCSC Xena platform for public and private cancer genomics data visualization and interpretation. in *bioRxiv 326470*.

25. Price JE, Polyzos A, Zhang RD, & Daniels LM (1990) Tumorigenicity and metastasis of human breast carcinoma cell lines in nude mice. *Cancer Res* 50(3):717-721.

26. Massague J & Obenauf AC (2016) Metastatic colonization by circulating tumour cells. *Nature* 529(7586):298-306.

27. Nguyen DX, Bos PD, & Massague J (2009) Metastasis: from dissemination to organ-specific colonization. *Nat Rev Cancer* 9(4):274-284.

28. Welch DR & Hurst DR (2019) Defining the Hallmarks of Metastasis. *Cancer Res* 79(12):3011-3027.

29. Nieto MA (2017) Context-specific roles of EMT programmes in cancer cell dissemination. *Nat Cell Biol* 19(5):416-418.

30. Barrallo-Gimeno A & Nieto MA (2005) The Snail genes as inducers of cell movement and survival: implications in development and cancer. *Development* 132(14):3151-3161.

31. Harris AL (2002) Hypoxia--a key regulatory factor in tumour growth. *Nat Rev Cancer* 2(1):38-47.

32. Balwierz PJ*, et al.* (2014) ISMARA: automated modeling of genomic signals as a democracy of regulatory motifs. *Genome Res* 24(5):869-884.

33. Majmundar AJ, Wong WJ, & Simon MC (2010) Hypoxia-inducible factors and the response to hypoxic stress. *Mol Cell* 40(2):294-309.

34. Li J & Radice GL (2010) A new perspective on intercalated disc organization: implications for heart disease. *Dermatol Res Pract* 2010:207835.

35. Plaks V, Koopman CD, & Werb Z (2013) Cancer. Circulating tumor cells. *Science* 341(6151):1186-1188.

36. Joosse SA, Gorges TM, & Pantel K (2015) Biology, detection, and clinical implications of circulating tumor cells. *EMBO Mol Med* 7(1):1-11.

37. Park J*, et al.* (2018) DSG2 Is a Functional Cell Surface Marker for Identification and Isolation of Human Pluripotent Stem Cells. *Stem Cell Reports* 11(1):115-127.

38. Dengler VL, Galbraith M, & Espinosa JM (2014) Transcriptional regulation by hypoxia inducible factors. *Crit Rev Biochem Mol Biol* 49(1):1-15.

39. Koshiji M*, et al.* (2005) HIF-1alpha induces genetic instability by transcriptionally downregulating MutSalpha expression. *Mol Cell* 17(6):793-803.

40. Li JY*, et al.* (2016) A Biparatopic HER2-Targeting Antibody-Drug Conjugate Induces Tumor Regression in Primary Models Refractory to or Ineligible for HER2-Targeted Therapy. *Cancer Cell* 29(1):117-129.

Figures and Figure Legends

Figure 1. High DSG2 expression is positively correlated with metastasis and high recurrence

risk in breast cancer patients.

(**A, B**) GSEA-P enrichment plot (A) and heat map (B) for the top 30 up-regulated genes for metastasis in metastatic (n = 7) versus non-metastatic breast cancer patients (n = 15). Normalized enrichment score (NES) and FDR q are listed on the enrichment plots.

(**C**) Representative immunohistochemistry (IHC) of breast tumors with negative (upper panel) and positive (lower panel) DSG2 expression. Enlarged images are presented on the right. Scale bar, 30 μm.

(**D**) Kaplan-Meier disease free survival analysis of breast cancer patients grouped by DSG2 expression. DSG2-positive group is indicated by red line (n = 113); DSG2-negative group is indicated by black line (n = 51). $p = 0.004$. The p-value was determined by log-rank test.

(**E**) Comparison of recurrence rate between patients with DSG2-positive vs. DSG2-negative tumors using chi-square test. $p = 0.03$.

(**F**) Kaplan–Meier analysis of distant metastasis-free survival of patients with different levels of DSG2 using a breast cancer cohort (Yau 2010 dataset) from the UCSC Xena public hub. Receiver operating characteristic (ROC) curve analysis was used to determine the relative level of DSG2. $p = 0.008$. The p-value was determined by log-rank test.

Figure 2. DSG2 expression promoted CTC clustering and metastatic colonization.

(**A**) Diagram of the procedure used for orthotopic xenografts. MB231 cells expressing EGFP without

(MB231-EGFP shCtrl) or with DSG2 depletion (shDSG2) were injected into 4th mammary fat pads

of NOD/SCIDγ mice. CTCs, primary tumors and lung tissues were collected at 9 weeks after

injection.

(**B**) Representative hematoxylin and eosin (H&E) staining of lung sections from mice orthotopically injected with shCtrl or shDSG2 MB231-EGFP cells and quantification of lung nodule numbers in shCtrl and shDSG2 groups. Error bars indicate S.D., *p* value was determined by unpaired T-test.

(**C**) Representative images of CTCs from shCtrl or shDSG2 MB231-EGFP tumor bearing mice captured by MiCareo on-chip filtration and having positive immunofluorescence (IF) staining of DAPI, EGFR and GFP to identify MB231-EGFP cells.

(**D**) Numbers of single CTCs and CTC clusters collected from pooled blood of shCtrl or shDSG2 MB231-EGFP tumor bearing mice. Three mice per group were assayed. Data are CTC counts ± 95% confidence intervals.

(**E**) Diagram of the syngeneic mouse model procedure. 4Tl-GFP/LUC without (shCtrl) or with DSG2 depletion (shDSG2) were injected into 4th mammary fat pads of BALB/c mice. Primary tumors and lung tissues were collected at 3 weeks after injection.

(**F**) Bioluminescence imaging of lung metastasis and quantitation of bioluminescence between shCtrl and shDSG2 groups. Four mice were used for each group. Data are means ± S.D., significant difference determined by T-test.

(**G**) Representative images of CTCs from shCtrl or shDSG2 4T1-GFP/LUC tumor bearing mice captured by MiCareo on-chip filtration and having positive IF staining of DAPI, CD29 and GFP to

identify 4T1-GFP/LUC cells.

(**H**) Numbers of single CTCs and CTC clusters collected from shCtrl- or shDSG2-transduced 4T1-

GFP/LUC tumor bearing mice. Four mice were used for each group. Data are means ± SD with

significant differences based on unpaired T-test (* indicates $p < 0.05$, ** indicates $p < 0.01$).

Figure 3. DSG2 expression promoted metastatic colonization and tumor growth.

(**A**) Lung metastasis and lung nodule numbers after tail vein injection using control or DSG2-depleted SKBR3 cells. Data are means ± S.D. with *p* value based on unpaired T-test. Images show representative H&E staining of lung sections from mice after tail vein injection. Scale bar indicates 100 μm.

(**B**) Lung metastasis and lung nodule numbers after tail vein injection using control or DSG2 overexpressing MB157 cells. Data formatting is as described for A.

(**C**) Soft agar colony formation assays using DSG2-depleted MB231 and SKBR3 cells. Data are

shown as mean ± SD with *p* value based on unpaired T-test. The experiments were repeated three

times.

(**D**) Soft agar colony formation assays using DSG2-overexpressing MB157 and MB468 cells.

Replication and data formatting are as described for C. **$p < 0.01$.

(**E**) Xenograft models in NOD/SCIDγ mice using control or DSG2-depleted MB231 and SKBR3

cells. Five mice were used for each group. Data are means ±S.D., significant difference is based on

unpaired T-test of the tumor size at the last time point.

(**F**) Xenograph models in NOD/SCIDγ mice using control or DSG2-overexpressing MB157 and

MB468 cells. Replication and data analysis are as described for E.

Figure 4. Negative correlation between hypoxia and DSG2 expression observed in clinical

specimens. Representative images of DSG2 and CA9 IHC staining using serial tumor sections from

clinical breast cancer patients. Red boxes show enlarged images of low DSG2 but high CA9

expression regions. Blue boxes show enlarged images of high DSG2 but low CA9 expression

regions. Scale bars indicate 2 mm or 100 μm as shown in the images. Examples shown are

representative of 151 patient samples examined. Among the 151 samples, 66 slides contained

DSG2^{high}/CA9^{low} cancer cells only, 40 slides contained DSG2^{low}/CA9^{low} cancer cells only and 45

slides contains both DSG2^{high}/CA9^{low} and DSG2^{low}/CA9^{high} cells in different regions.

Figure 5. HIF1α suppressed DSG2 expression under hypoxic stress.

(**A**) qRT-PCR analysis of DSG2 level in SKBR3 and MB231 cells under normoxia or hypoxia for 16 hours. Cells under hypoxic stress are indicated by elevated HIF1α. HIF1α expression was detected by immuoblot with GAPDH as a loading control. Data are means ±S.D. (n =3). Significant differences are based on unpaired T-test. The experiments were repeated three times.

(**B**) Time-course immunoblot of DSG2 and HIF1α in SKBR3 cells under normoxia or hypoxia.

GAPDH was used as a loading control. Blots shown are representative of three replicate

experiments.

(**C**) qRT-PCR analysis of DSG2 in SKBR3 and MB231 cells treated with 200 μM CoCl2 for 2 or

24hr. Data are means ± S.D. (n =3). Significant differences are based on unpaired T-test. The

experiments were repeated three times.

(**D**) qRT-PCR analysis of DSG2 level in 293T cells transiently transfected with pcDNA vector or

pcDNA-HIF1α-P564A plasmid. HIF1α expression was detected by immuoblot with GAPDH as a

loading control. Three independent experiments were performed and data are means ± SD from one

representative experiment (n =3). Significant differences are based on unpaired T-test.

(**E**) qRT-PCR of DSG2 level in 293T cells transduced with shHIF1α lentiviral vectors (clone # 675

and # 677) and treated with 100 μM CoCl2 for 24 hours. Immunoblot of HIF1α shows that hypoxia

induced accumulation of HIF1α was effectively blocked by shHIF1α. The experiments were

repeated three times.

(**F**) Diagram shows three putative HIF1α binding sites on the DSG2 promoter predicted using

ISMAR and the mutated promoter sequences used in luciferase reporter assays. Luciferase reporter

assays were conducted using 293T cells co-transfected with pcDNA.HIF1α and the wild type (Wt) or

mutant DSG2 promoter constructs. Three replicate experiments were performed and data are means

± SD from one representative experiment and significant differences detected using T-test.

(**G**) ChIP-qPCR analysis of HIF1α occupancy on the DSG2 promoter region containing putative

HIF1α binding site (-946~-943 nt) in SKBR3 cells under normoxia or hypoxia.

(**H**) ChIP-qPCR analysis of H3K27me3 on the same DSG2 promoter region as in G. Three

independent experiments were performed and data are means ± SD from one representative

experiment with significant differences detected by un-paired T-test. (* indicates $p < 0.05$, **

indicates $p < 0.01$ and *** indicates $p < 0.001$).

Figure 6. Hypoxia suppresses DSG2 expression via formation of a HIF1α-PRC2 complex.

(**A**) Co-IP of HIF1α, EZH2 and SUZ12 in SKBR3 cells without (normoxia) or with CoCl₂ treatment.

IgG was used as a negative control. The experiments were repeated three times.

(**B**) Co-IP of SUZ12, EZH2 and HIF1α in 293T cells transfected with HIF1α.P564A. IgG was used

as a negative control. The experiments were repeated three times.

(**C**) IB of EZH2, SUZ12 and HIF1α (top) and qRT-PCR analysis of DSG2 level (bottom) in shCtrl

(shC), shEZH2 or shSUZ12-transduced SKBR3 cells treated with $CoCl_2$. GAPDH was used as a

control. qRT-PCR data are means ±S.D. (n = 3). Significant differences were detected by T-test.

(**D**) IB of HIF1α in SKBR3 cells transduced with shHIF1α lentiviral vectors (clone # 674 and # 677)

and treated with $CoCl_2$ for 6 hours. The experiments were repeated three times.

(**E**) ChIP-qPCR analysis of EZH2 and SUZ12 on the P2 promoter region (-1062 to-552) in HIF1α-

depleted SKBR3 cells with or without $CoCl_2$ treatment. Data are mean ± SD with significant

differences base on unpaired T-test (*$p < 0.05$, **$p < 0.01$. ***$p < 0.001$). The experiments were

repeated three times.

(**F**) qRT-PCR analysis of DSG2 in SKBR3 cells kept under normoxia throughout the experiment

(black line), or under hypoxia for 2 hr (blue line) or 3 hr (red line) and then released from the stress.

Data are means ±S.D (n = 3). Significant differences are based on T-test of data at each time point.

The experiments were repeated three times.

(**G**) FACS analysis of DSG2 levels in shCtrl EGFP-MB231 cells before injection (left panel) or in

CTCs collected from pooled blood of four mice at 9 weeks after injection (right panel).

Figure 7. Dynamic changes of DSG2 expression contribute to breast tumorigenesis and metastasis. High DSG2 expression promotes breast tumor growth in mammary tissue. In non-hypoxic tumor regions with high DSG2, CTC clusters may be shed and DSG2-mediated cell adhesion may facilitate CTC clustering prior to intravasation. Alternatively, when the tumor is large enough to induce hypoxia, HIF1α is stabilized and recruits PRC2 complex (EZH2 and SUZ12) to the DSG2 promoter to suppress its transcription. DSG2 suppression allows cancer cells to undergo EMT and single CTCs are released from the primary tumor. In circulatory system, hypoxic stress is released, DSG2 expression reactivated and DSG2-mediated adhesion allows CTC aggregation in blood and colonization in lung.