ANCESTRAL MITOCHONDRIAL PROTEIN SECRETION MACHINERY

Lenka Horváthová[1*], Vojtěch Žárský[1*], Tomáš Pánek[2*], Romain Derelle[3], Jan Pyrih[4], Alžběta Motyčková[1], Veronika Klápšťová[1], Vladimír Klimeš[2], Markéta Petrů[1], Zuzana Vaitová[1], Ivan Čepička[5], Karel Harant[6], Michael W. Gray[7], Ingrid Guilvout[8], Olivera Francetic[8], B. Franz Lang[9], Čestmír Vlček[10], Anastasios D. Tsaousis[4], Marek Eliáš[2#], Pavel Doležal[1#]

[1]Department of Parasitology, Faculty of Science, Charles University, BIOCEV, Průmyslová 595, Vestec, 252 42, Czech Republic
[2]Department of Biology and Ecology, Faculty of Science, University of Ostrava, Chittussiho 10, 710 00 Ostrava, Czech Republic
[3]School of Biosciences, University of Birmingham, Edgbaston, B15 2TT, UK
[4]Laboratory of Molecular & Evolutionary Parasitology, RAPID group, School of Biosciences, University of Kent, Canterbury, CT2 7NZ, UK
[5]Department of Zoology, Faculty of Science, Charles University, Viničná 7, Prague 2, 128 44, Czech Republic
[6]Proteomic core facility, Faculty of Science, Charles University , BIOCEV, Průmyslová 595, Vestec, 252 42, Czech Republic
[7]Department of Biochemistry and Molecular Biology and Centre for Comparative Genomics and Evolutionary Bioinformatics, Dalhousie University, Halifax, NS B3H 4R2, Canada
[8]Institut Pasteur, Biochemistry of Macromolecular Interactions Unit, Department of Structural Biology and Chemistry, CNRS UMR3528, 75015 Paris, France
[9]Robert Cedergren Centre for Bioinformatics and Genomics, Département de Biochimie, Université de Montréal, Montreal, QC, Canada H3T 1J4
[10]Institute of Molecular Genetics, Czech Academy of Sciences, 142 20 Prague 4, Czech Republic

*these authors equally contributed to the study
#corresponding authors: Marek Eliáš (marek.elias@osu.cz), Pavel Doležal (pavel.dolezal@natur.cuni.cz)

Abstract

Modern mitochondria have preserved few traits of the original bacterial endosymbiont. Unexpectedly, we find that certain representatives of heteroloboseans, jakobids and malawimonads possess homologues of four core components of the type 2 secretion system (T2SS) so far restricted to eubacteria. We show that these components are localized to the mitochondrion, and their behaviour in functional assays is consistent with the formation of a mitochondrial T2SS-derived protein secretion system. We additionally identified 23 protein families exactly co-occurring in eukaryotes with the T2SS. Seven of these proteins could be directly linked to the core T2SS by functional data and/or sequence features, whereas others may represent different parts of a broader functional pathway, possibly linking the mitochondrion with the peroxisome. Its distribution in eukaryotes and phylogenetic evidence indicate that the whole mitochondrial T2SS-centred pathway is an ancestral eukaryotic trait. Our findings thus have direct implications for the functional properties of the early mitochondrion.

Introduction

Mitochondria of all eukaryotes arose from the same Alphaproteobacteria-related endosymbiotic bacterium[1,2]. New functions have been incorporated into the bacterial blueprint during mitochondrial evolution, while many ancestral traits have been lost. Importantly, in some cases, these losses occurred independently in different lineages of eukaryotes, resulting in a patchy distribution of the respective ancestral mitochondrial traits in extant eukaryotes. A good example of this is the ancestral mitochondrial division apparatus (including homologues of bacterial Min proteins) retained in several distantly related protist lineages[3,4]. It is likely that additional pieces of the ancestral bacterial cell physiology will be discovered in mitochondria of poorly studied eukaryotes.

An apparent significant difference between the mitochondrion and bacteria (including those living as endosymbionts of eukaryotes) lies in the directionality of protein transport across their envelope. All bacteria export specific proteins from the cell via the plasma membrane using the Sec or Tat machineries[5], and many diderm (Gram-negative) bacteria exhibit specialized systems mediating further protein translocation across the outer membrane (OM)[6]. In contrast, the mitochondrion depends on a newly evolved protein import system spanning both envelope membranes and enabling import of proteins encoded by the nuclear genome[7]. The capacity of mitochondria to secrete proteins seems to be limited. Mitochondrial homologues of Tat translocase subunits occur in some eukaryotic taxa, but their role in protein secretion has not been established[8]. A mitochondrial homologue of the SecY protein (a Sec translocase subunit) has been described only in jakobids[9,10], but its function remains elusive[11]. No dedicated machinery for protein export from the mitochondrion across the outer mitochondrial membrane has been described.

One of the best characterized bacterial protein translocation machineries is the so-called type 2 secretion system (T2SS)[12,13]. The T2SS belongs to a large bacterial superfamily of type 4 pili (T4P)-related molecular machines, most of which

secrete long extracellular filaments (pili) for motility, adhesion, or DNA uptake[14–16].
Using building blocks homologous to components of the other members of the T4P
superfamily, the T2SS constitutes a specialized secretion apparatus, whose filament
(pseudopilus) remains in the periplasm[12,13]. It is composed of 12-15 conserved
components, commonly referred to as general secretion pathway (Gsp) proteins,
which assemble into four main subcomplexes (Fig. 1A). The OM pore is formed by
an oligomer of 15-16 molecules of the GspD protein[17]. The subcomplex in the inner
membrane (IM) is called the assembly platform and consists of the central
multispanning membrane protein GspF surrounded by single-pass membrane
proteins GspC, GspL, and GspM. GspC links the assembly platform to the OM pore by
interacting with the periplasmic N-terminal domain of GspD[18]. The third
subcomplex, called the pseudopilus, is a helical filament formed mainly of GspG
subunits, with minor pseudopilins (GspH, GspI, GspJ and GspK) assembled at its tip[15].
The pseudopilus is assembled at the assembly platform and its growth is believed to
push the periplasmic T2SS substrate through the OM pore. The energy for
pseudopilus assembly is provided by the fourth subcomplex, the hexameric ATPase
GspE, interacting with the assembly platform from the cytoplasmic side[16].
Substrates for T2SS-mediated secretion are first transported by the Tat (as
folded proteins) or the Sec (in an unfolded form) system across the IM into the
periplasm, where they undergo maturation and/or folding. The folded substrates
are finally loaded onto the pseudopilus for the release outside the cell via the OM
pore. The known T2SS substrates differ between taxa and no common sequence
features have been identified for them. Proteins transported by the T2SS in different
species include catabolic enzymes (such as lipases, proteases or phosphatases) and,
in the case of bacterial pathogens, toxins[12]. A recent survey of bacterial genomes
showed that the T2SS is mainly present in Proteobacteria[19]. Crucially, neither the
T2SS nor other systems of the T4P superfamily have been reported from
eukaryotes[6,12,20].
Here we show that certain distantly related eukaryotes unexpectedly contain
homologues of key T2SS subunits representing all four functional T2SS
subcomplexes. We provide evidence for mitochondrial localization of these
eukaryotic Gsp homologues and describe experimental results supporting the idea
that they constitute a system similar to the bacterial T2SS. Furthermore, we point to
the existence of 23 proteins with a perfect taxonomic co-occurrence with the
eukaryotic Gsp homologues. Some of these co-occurring proteins seem to be
additional components of the mitochondrial T2SS-related machinery, whereas
others are candidates for components of a broader functional pathway linking the
mitochondrion with other parts of the cell. Given its phylogenetic distribution we
propose that the newly discovered pathway was ancestrally present in eukaryotes.
Its further characterization may provide fundamental new insights into the
evolutionary conversion of the protomitochondrion into the mitochondrial
organelle.
135
Results
Certain protist lineages code for a conserved set of homologues of T2SS core
components

While searching the genome of the heterolobosean *Naegleria gruberi* for proteins of
bacterial origin with a possible mitochondrial role, we surprisingly discovered
homologues of four core subunits of the bacterial T2SS, specifically GspD, GspE,
GspF, and GspG (Fig. 1A; Supplementary Table 1). Using genomic and transcriptomic
data from public repositories and our on-going sequencing projects for several
protist species of key evolutionary interest, we mapped the distribution of these
four components in eukaryotes. All four genes were found in the following
characteristic set of taxa (Fig. 1B, Supplementary Table 1): three additional
heteroloboseans (*Naegleria fowleri*, *Neovahlkampfia damariscottae*, *Pharyngomonas*
kirbyi), two jakobids (*R. americana* and *Andalucia godoyi*), and two malawimonads
(*Malawimonas jakobiformis* and *Gefionella okellyi*). In addition, three separate
representatives of the heterolobosean genus *Percolomonas* (Supplementary Fig. 1)
each exhibited a homologue of GspD, but not of the remaining Gsp proteins, in the
available transcriptomic data. In contrast, all four genes were missing in sequence
data from all other eukaryotes investigated, including the genome and
transcriptome of another malawimonad ("Malawimonas californiana") and deeply-
sequenced transcriptomes of a third jakobid (*Stygiella incarcerata*) and four
additional heteroloboseans (*Creneis carolina*, "Dactylomonas venusta", *Harpagon*
schusteri, and the undescribed strain Heterolobosea sp. BB2).
Probing *N. gruberi* nuclei with fluorescent *in situ* hybridization ruled out an
unidentified bacterial endosymbiont as the source of the Gsp genes (Supplementary
Fig. 2). Moreover, the eukaryotic Gsp genes usually have introns and constitute
robustly supported monophyletic groups well separated from bacterial homologues
(Fig. 1C; Supplementary Fig. 3), ruling out bacterial contamination in all cases. In an
attempt to illuminate the origin of the eukaryotic Gsp proteins we carried out
systematic phylogenetic analyses based on progressively expanded datasets of
prokaryotic homologues and for each tree inferred the taxonomic identity of the
bacterial ancestor of the eukaryotic branch (see Methods for details on the
procedure). The results, summarized in Supplementary Fig. 3, showed that the
inference is highly unstable depending on the dataset analysed, and no specific
bacterial group can be identified as an obvious donor of the eukaryotic Gsp genes.
This result probably stems from a combination of factors, including the long
branches separating the eukaryotic and bacterial Gsp sequences, the length of Gsp
proteins restricting the amount of the phylogenetic signal retained, and perhaps
also rampant horizontal gene transfer of the T2SS system genes between bacterial
taxa. The eukaryotic Gsp genes are in fact so divergent that some of them could not
be unambiguously classified as specific homologs of T2SS components (as opposed
to the related machineries of the T4P superfamily) when analysed using models
developed for the bacterial genomes[19] (Supplementary Fig. 3).
Heteroloboseans, jakobids and malawimonads have been classified in the
hypothetical supergroup Excavata[21]. However, recent phylogenomic analyses
indicate that excavates are non-monophyletic and even suggest that malawimonads
are separated from heteroloboseans and jakobids by the root of the eukaryote
phylogeny[22–25]. Hence, the current phylogenetic distribution of the Gsp homologues
in eukaryotes may reflect their presence in the last eukaryotic common ancestor
(LECA) followed by multiple independent losses (Fig. 1C). Heteroloboseans and

malawimonads have two GspG paralogues, but the phylogenetic analyses did not
resolve whether this is due to multiple independent GspG gene duplications or one
ancestral eukaryotic duplication followed by loss of one of the paralogues in
jakobids (Supplementary Fig. 3D; Supplementary Table 1).
189
The eukaryotic Gsp proteins localize to the mitochondrion
We hypothesized that the eukaryotic homologues of the four Gsp proteins are parts
of a functional T2SS-related system localized to the mitochondrion. This notion was
supported by the presence of predicted N-terminal mitochondrial targeting
sequences (MTSs) in some of the eukaryotic Gsp proteins (Supplementary Table 1).
The prediction algorithms identified putative N-terminal MTSs for proteins from
jakobids and malawimonads but failed to recognize them in the orthologues from
heteroloboseans, which, however, carry the longest N-terminal extensions
(Supplementary Fig. 4). We assumed that these extensions might still function as
MTSs in heteroloboseans. Indeed, labelling of *N. gruberi* cells using specific
polyclonal antibodies showed that GspD, GspF and GspG1 are present in
mitochondria (Fig. 2A). Moreover, the atypical MTSs of *N. gruberi* Gsp proteins were
efficiently recognized by the yeast mitochondrial import machinery (Supplementary
Fig. 5). Analogously, three Gsp proteins from *G. okellyi* were all localized to
mitochondria when expressed in yeast (Fig. 2B).
In order to further confirm the mitochondrial localization of the Gsp proteins
in *N. gruberi*, we analysed the mitochondrial proteome of this species by partial
purification of the organelle and identification of resident proteins by mass
spectrometry. A mitochondria-enriched fraction was obtained from a cellular lysate
by several steps of differential centrifugation and further separated by OptiPrep
gradient centrifugation. Three sub-fractions of different densities were collected
(Supplementary Fig. 6A) and subjected to proteomic analysis. The relative amount
of each protein in the gradient was determined by label-free quantification and the
proteins were grouped by a multicomponent analysis (for details see Methods)
according to their distributions across the gradient (Fig. 3). A set of marker proteins
(homologs of well characterized typical mitochondrial proteins from other species)
was used to identify a cluster of mitochondrial proteins. Due to the partial co-
purification of peroxisomes with mitochondria, a peroxisome-specific cluster was
defined analogously. As a result, 946 putative mitochondrial and 78 putative
peroxisomal proteins were identified among the total of 4,198 proteins detected.
Encouragingly, the putative mitochondrial proteome of *N. gruberi* is dominated by
proteins expected to be mitochondrial or whose mitochondrial localization is not
unlikely (Supplementary Fig. 6B, Supplementary Table 2A). On the other hand, the
putative peroxisomal proteome seems to be contaminated by mitochondrial
proteins (owing to the presence of several mitochondrial ribosomal proteins;
Supplementary Table 2B). Importantly, all five Gsp proteins (including both GspG
paralogs) were identified in the putative mitochondrial but not peroxisomal
proteome of *N. gruberi*.
228
229 ***The properties of the eukaryotic Gsp proteins support the existence of a***
230 ***mitochondrial T2SS-related machinery***

The foregoing experiments support the idea that all four eukaryotic Gsp homologues localize to and function in the mitochondrion. However, direct *in vivo* demonstration of the existence of a functional mitochondrial T2SS-related machinery is currently not feasible, because none of the Gsp homologue-carrying eukaryotes represents a tractable genetic system. We thus used *in vitro* approaches and heterologous expression systems to test the key properties of the eukaryotic Gsp proteins.

Crucial for the T2SS function is the formation of the OM pore, which is a β-barrel formed by the oligomerization of the C-domain of the GspD protein[26]. The actual assembly of the bacterial pore requires the interaction of the very C-terminal domain of GspD (S-domain) with the outer membrane lipoprotein GspS[27]. In addition, the bacterial GspD carries four short N-terminal domains exposed to the periplasm, called N0 to N3, of which N1 to N3 share a similar fold[28] (Fig. 4A). While the N3 domain has been shown to participate in the pore assembly, N0 interacts with GspC of the assembly platform[18]. However, sequence analysis of the mitochondrial GspD homologue revealed that it, in fact, corresponds to only a C-terminal part of the bacterial GspD β -barrel C-domain, whereas the N-terminal domains N0 to N3, the N-terminal part of the C-domain, and the S-domain are missing (Fig. 4A). This finding raised a question whether the mitochondrial GspD homologue has retained the ability to form a membrane pore. Nevertheless, homology modelling of GspD from *G. okellyi* (*Go*GspD) using *Vibrio cholerae* GspD[29] as a template indicated that the protein could be fitted into solved structure of the pentadecameric pore complex with the conserved amphipathic helical loop (AHL)(Fig. 4B).

Testing the function of *Go*GspD in bacteria was impossible due to its high toxicity leading to rapid cell death upon induction of protein expression (Fig. 4C), which is a typical behaviour of pore-forming proteins. The protein toxicity was less pronounced in the yeast two-hybrid (Y2H) system, which indicated strong self-interaction of *Go*GspD (Fig. 4D), and hence its ability to oligomerize. Indeed, radioactively labelled *Go*GspD assembled into a high-molecular-weight complex in an experimental membrane in an *in vitro* translation assay (Fig. 4E). The formation of the complex was dependent on the presence of the membrane and the complex was resistant to 2M urea treatment, which would remove nonspecific protein aggregates. These results showed that the mitochondrial GspD, despite being significantly truncated when compared to its bacterial homologues, has retained the capability to form membrane pores, characteristic for the secretins of the T2SS [30]. Compared to the bacterial GspD, the predicted *Go*GspD structure suggests a unique biogenesis pathway, where the secretin pore-forming domain may be directly inserted in the mitochondrial outer membrane, bypassing the membrane transport essential for its bacterial counterparts.

The secretion mechanism of the T2SS relies on assembly of pseudopilus made up of GspG subunits[15]. A possible assembly of mitochondrial GspG from *G. okellyi* (*Go*GspG1) into the pseudopilus structure was indicated by modelling the protein sequence into the recently obtained cryoEM reconstruction of the PulG complex from *Klebsiella oxytoca*[20] (Supplementary Fig. 7). The actual interaction properties of *Go*GspG1 were followed by the bacterial two-hybrid assay (B2H). When expressed in bacteria (in a truncated form with the MTS region removed, see

Fig. 5A), the mitochondrial *Go*GspG1 interacted with itself (Fig. 5B), which is a
prerequisite for pseudopilus formation. An analogous B2H assays of *N. gruberi* Gsp
proteins also showed GspG1 self-interaction (data not shown). In addition, *Go*GspG1
showed positive interaction with *Go*GspF, the IM component believed to participate
in transfer of energy for the pseudopilus assembly from GspE (Fig. 1A). Moreover,
the mitochondrial *Go*GspF and *Go*GspE each self-interacted in the B2H assay (Fig.
5B). These interactions are in agreement with the role of both proteins as T2SS
components, as GspF forms dimers within the IM complex and GspE assembles into
an active hexameric ATPase. Furthermore, B2H assay has identified the same
interactions between the GspG and GspF homologues in the bacterial T2SS [31]. Tests
of all other possible interactions of *G. okellyi* Gsp proteins were negative.
The *in silico* analyses and experiments described above are consistent with
the hypothesized existence of a functional mitochondrial secretion machinery
derived from the bacterial T2SS. However, the mitochondrial subunits identified
would assemble only a minimalist version of the secretion system, reduced to the
functional core of the four subcomplexes of the bacterial T2SS, i.e. the luminal
ATPase (GspE), the IM pseudopilus assembly platform (GspF), the intermembrane
space pseudopilus (GspG), and the OM pore (truncated GspD). Despite using
sensitive HMM-based searches, we did not detect homologues of other conserved
T2SS subunits in any of the eukaryotes possessing GspD to GspG proteins. One of the
missing subunits is GspC, which connects the assembly platform with the N0 domain
of GspD pore[18,32]. Thus, the absence of GspC in eukaryotes correlates with the lack of
the N0 domain in the eukaryotic GspD. Analogously, the absence of the C-terminal S-
domain in the mitochondrial GspD (Fig. 4A), known to be missing also from some
bacterial GspD proteins, rationalizes the lack of a eukaryotic homologue of the
bacterial OM component GspS that binds to GspD via the S-domain during the pore
assembly[27].
The mitochondrial system also apparently lacks a homologue of GspO, a
bifunctional enzyme that is essential for GspG maturation. Despite this absence,
eukaryotic GspG homologues have conserved all the characteristic sequence
features required for GspG maturation (the polar anchor and the trans-membrane
domain with a conserved glutamate residue at the +5 position relative to the
processing site) (Fig. 5A, Supplementary Fig. 4D). Notably, all the *Ng*GspG1 and
*Ng*GspG2-derived peptides detected in our proteomic analysis come from the region
of the protein downstream of the conserved processing site (Fig.5C), and an anti-
*Ng*GspG1 antibody detected a specific band of a much smaller size than expected for
the full-length protein (around 44 kDa) on a western blot of electrophoretically
separated *N. gruberi* proteins (Fig.5D). However, the theoretical Mw of the *Ng*GspG1
processed at the conserved site is 25.5 kDa, whereas the protein detected by the
immunoblot is even smaller, with a size similar to that of bacterial pseudopilins.
Hence, the actual nature of the mitochondrial GspG maturation needs to be studied
further.
319
New putative components of the mitochondrial T2SS-based functional pathway
identified by phylogenetic profiling

Since none of the eukaryotes with the Gsp homologues is currently amenable to
functional studies, we tried to further illuminate the role of the mitochondrial T2SS
system using a comparative genomic approach. Specifically, we reasoned that
possible additional components of the machinery, as well as its actual substrate(s),
might show the same phylogenetic distribution as the originally identified four
subunits. Using a combination of an automated identification of candidate protein
families and subsequent manual scrutiny by exhaustive searches of available
eukaryote sequence data (for details of the procedure see Methods), we identified
23 proteins (more precisely, groups of orthologues) that proved to exhibit precisely
the same phylogenetic distribution in eukaryotes as the four core T2SS components.
Specifically, all 23 proteins were represented in each of the heterolobosean, jakobid,
and malawimonad species possessing all four core Gsp proteins, whereas only seven
of them were found in the transcriptomic data from the *Percolomonas* lineage that
possesses only GspD (Fig. 1B; Supplementary Table 3). Except for two presumably
Gsp-positive jakobids represented by incomplete EST surveys and a case of a likely
contamination (Supplementary Table 4), no orthologues of any of these proteins
were found in any other eukaryote (including the Gsp-lacking members of
heteroloboseans, jakobids and malawimonads). The sequences of these 23 proteins
were analysed by various *in silico* approaches, including sensitive homology-
detection methods (HMM-HMM comparisons with HHpred[33] and protein modelling
using the Phyre2 server[34]) to assess their possible function (Fig. 6A; Supplementary
Table 3).
These analyses revealed that seven of the families have a direct link to the
T2SS suggested by discerned homology to known T2SS components. One of them
represents an additional, more divergent homologue of the C-terminal part of the
bacterial GspD. Hence, the protein has been marked as GspDL (GspD-like). Three
other families, referred to as GspDN1 to GspDN3, proved to be homologous to the
Secretin_N domain (Pfam family PF03958), present in the bacterial GspD protein in
three copies as the domains N1, N2, and N3 (Fig. 4A). The N1-N3 array protrudes
into the periplasmic space, where it oligomerizes to form three stacked rings[35]. As
mentioned above, the initially identified eukaryotic GspD homologues lack the N-
terminal region, suggesting that the gene was split into multiple parts in eukaryotes.
Unfortunately, high sequence divergence makes it impossible to identify potential
specific correspondence between the N1 to N3 domains of the bacterial GspD and
the eukaryotic GspDN1 to GspDN3 proteins. Importantly, an initial Y2H assay
indicated that the two separate polypeptides GspD and GspDN1 of *N. gruberi* may
interact *in vivo* (Fig. 4F), perhaps forming a larger mitochondrial complex. In
addition, we identified most of the newly discovered GspD-related proteins (GspDL
and GspDN) in the *N. gruberi* mitochondrial proteome (the exception being GspDN1,
which was not detected in a sufficient number of replicates to be included in the
downstream analysis; Supplementary Table 2A).
The final three proteins linked to the T2SS based on their sequence features
represent three divergent paralogues of the GspE subunit (GspE-like) here denoted
GspEL1 to GspEL3. However, abrogation of ATPase-specific motifs in these
paralogues suggests the loss of the ATPase activity (Supplementary Fig. 4B). GspEL2
and GspEL3 were identified among *N. gruberi* mitochondrial proteins in the

proteomic analysis, whereas GspEL1 was found in the cluster of putative
peroxisomal proteins.
The remaining sixteen proteins co-occurring with the core eukaryotic T2SS
subunits, hereafter referred to as Gcp (Gsp-co-occurring proteins), were divided
into three categories. The first comprises four proteins that constitute novel
paralogues within broader common eukaryotic (super)families (Fig. 6B). Three of
them (Gcp1 to Gcp3) belong to the WD40 superfamily, in which they form a single
clade together with the peroxisomal protein import co-receptor Pex7 (Fig. 6B;
Supplementary Fig. 8). None of these proteins has any putative N-terminal targeting
sequence, but interestingly, the peroxisomal targeting signal 1 (PTS1) could be
predicted on most Gcp1 and some Gcp2 proteins (Supplementary Table 3). However,
these predictions are not fully consistent with the results of our proteomic analysis:
*Ng*Gcp1 was found among the mitochondrial proteins and *Ng*Gcp2 in the cluster of
putative peroxisomal proteins (Supplementary Table 2), but PTS1 is predicted to be
present in the *Ng*Gcp1 protein (Supplementary Table 3). The fourth Gcp protein
(Gcp4) is a novel paralogue of the ubiquitin-like superfamily, distinctly different
from the previously characterized members including ubiquitin, SUMO, NEDD8 and
others (Supplementary Fig. 9).
The second Gcp category comprises eleven proteins (Gcp5 to Gcp15) well
conserved at the sequence level among the Gsp-containing eukaryotes, yet lacking
any discernible homologues in other eukaryotes or in prokaryotes. Two of these
proteins (Gcp8, Gcp15) were not identified in the proteomic analysis of *N. gruberi*
(Supplementary Table 3). Of those identified, several (Gcp5, Gcp6, Gcp13) were
found among the mitochondrial proteins, whereas some others (Gcp9, Gcp10,
Gcp11) clustered with peroxisomal markers. Specific localization of the three
remaining proteins (Gcp7, Gcp12, and Gcp14) could not be determined due to their
presence at the boundaries of the mitochondrial or peroxisomal clusters.
No homology to other proteins or domains could be discerned for the Gsp5 to Gsp15
proteins even when sensitive homology-detection algorithms were employed.
However, four of them are predicted as single-pass membrane proteins, with the
transmembrane segment in the N- (Gcp7, Gcp11, Gcp15) or C-terminus (Gcp5) (Fig.
6A; Supplementary Fig. 10). Interestingly, Gcp6 and Gcp12 proteins contain multiple
absolutely conserved cysteine or histidine residues (Fig. 6A; Supplementary Fig. 11).
Finally, Gcp16 constitutes a category of its own. It typifies a family of
predicted membrane proteins with non-eukaryotic representatives restricted to
bacteria of the PVC superphylum (Supplementary Fig. 12), some of which are known
to have the T2SS[36]. Interestingly, Gcp16 proteins from *Neochlamydia* spp. are fused
to the N-terminus of a protein from the Lactamase_B_2 (PF12706) family that
generally occurs as an independent protein widely conserved in various bacteria.
Phylogenetic analyses confirmed that the eukaryotic members of the family are of
the same origin rather than acquisitions by independent HGT events into different
lineages of eukaryotes (Supplementary Fig. 13). Most eukaryotic Gcp16 proteins
exhibit an N-terminal extension compared to the bacterial homologues
(Supplementary Fig. 12), but only some of these extensions are recognized as
putative MTSs and the *N. gruberi* Gcp16 was not identified either in putative
mitochondrial or peroxisomal proteome.

Discussion

Our analyses revealed that a subset of species belonging to three eukaryotic lineages share a set of at least 27 proteins (or families of orthologues) absent from other eukaryotes for which genomic or transcriptomic data are currently available (Fig. 1C). At least eleven of these proteins (the Gsp proteins) are evolutionarily related to components of the bacterial T2SS, although seven of them are so divergent that their evolutionary connection to the T2SS could be recognized only retrospectively after their identification based on their characteristic phylogenetic profile. For the sixteen remaining proteins (Gcp1 to Gcp16) no other evolutionary or functional link to the T2SS is evident apart from the same phyletic pattern as exhibited by the T2SS subunit homologues. Nevertheless, similar phylogenetic profiles are generally a strong indication for proteins being parts of the same functional system or pathway, and have enabled identification of new components of different cellular structures or pathways (e.g. refs[37,38]). Is it, therefore, possible that the 27 Gsp/Gcp proteins similarly belong to a single functional pathway?

The phylogenetic profile shared by the eukaryotic Gsp and Gcp proteins is not trivial, as it implies independent gene losses in a specific set of multiple eukaryotic branches (Fig. 1B). The likelihood of a chance emergence of the same taxonomic distribution of these proteins is thus low. Nevertheless, false positives cannot be completely excluded among the Gcp proteins and their list may be revised when a more comprehensive sampling of eukaryote genomes or transcriptomes becomes available. It is also possible that the currently inferred phylogenetic profile of some of the Gsp/Gcp proteins is inaccurate due to incomplete sampling of the actual gene repertoire of species represented by transcriptome assemblies only. An interesting case in point is the heterolobosean *Percolomonas* lineage. Transcriptomic data from three different members revealed only the presence of GspD, GspDL, the three GspDN variants, and four Gcp proteins (Fig. 1B, Supplementary Tables 1 and 3), which may reflect incomplete data. However, the relatively coherent pattern of Gsp/Gcp protein occurrence in the three independently sequenced transcriptomes and the fact that in other Gsp/Gcp - containing eukaryotes all 27 families are always represented in the respective transcriptome assembly (Supplementary Tables 1 and 3) suggest that the *Percolomonas* lineage has preserved only a subset of Gsp/Gcp families. Genome sequencing is required to test this possibility.

All uncertainties notwithstanding, our data favour the idea that a hitherto unknown complex functional pathway exists in some eukaryotic cells, underpinned by most, if not all, of the 27 Gsp/Gcp proteins and possibly others yet to be discovered. Direct biochemical and cell biological investigations are required for testing its very existence and the actual cellular role. Nevertheless, we integrated the experimental data gathered so far with the insights from bioinformatic analyses to propose a hypothetical working model (Fig. 7).

Our main proposition is that the eukaryotic homologues of the bacterial Gsp proteins assemble a functional transport system, here denoted miT2SS, that spans the mitochondrial OM and mediates the export of specific substrate proteins from the mitochondrion. Although the actual architecture of the miT2SS needs to be

determined, the available data suggest that it departs in detail from the canonical
bacterial T2SS organization, as homologues of some of the important bacterial T2SS
components are apparently missing. Most notable is the absence of GspC,
presumably related to the modified structure of its interacting partner GspD, which
in eukaryotes is split into multiple polypeptides and seems to completely lack the
N0 domain involved in GspC binding. It thus remains unclear whether and how the
IM assembly platform and the OM pore interact in mitochondria. One possible
explanation is that GspC has been replaced by an unrelated protein. It is notable that
three Gcp proteins (Gcp7, Gcp11, and Gcp15) have the same general architecture as
GspC: they possess a transmembrane segment at the N-terminus and a (predicted)
globular domain at the C-terminus (Fig. 6A). Testing possible interactions between
these proteins and T2SS core subunits (particularly GspF and GspDN) using B2H or
Y2H assays will be of future interest.
Future investigations also must address the question of whether the
mitochondrial GspG is processed analogously to the bacterial homologues and how
such processing occurs in the absence of discernible homologues of GspO (see
above). The mitochondrial GspG is presumably inserted into the IM by the Tim22 or
Tim23 complex, resulting in a GspG precursor with the N-terminus, including the
MTS, protruding into the matrix. It is possible that N-terminal cleavage by matrix
processing peptidase serves not only to remove the transit peptide, but at the same
time to generate the mature N-terminus of the processed GspG form, ready for
recruitment into the pseudopilus.
In parallel with its apparent simplification, the miT2SS may have been
specifically elaborated compared to the ancestral bacterial machinery. This
possibility is suggested by the existence of the three divergent, possibly ATPase
activity-deficient GspE paralogues (GspEL1 to GspEL3) that we discovered in all
miT2SS-containing eukaryotes but not elsewhere. We can only speculate as to the
function of these proteins, but they may interact with and regulate the catalytically
active GspE protein. The fact that the bacterial GspE assembles into a homohexamer
raises the possibility that in eukaryotes GspEL proteins are included in a
heterooligomer with GspE, a situation analogous to the presence of catalytically
active and inactive paralogous subunits in some well known protein complexes (e.g.
refs[39,40]). The co-occurrence of two different paralogues of the GspD C-domain, one
(GspDL) being particularly divergent, suggests a eukaryote-specific elaboration of
the putative pore in the mitochondrial OM.
An unanswered key question is what is the actual substrate (or substrates)
exported from the mitochondrion by the miT2SS. No bioinformatic tool for T2SS
substrate prediction is available due to the enigmatic nature of the mechanism of
substrate recognition by the pathway[12], so at the moment we can only speculate. It
is notable that no protein encoded by the mitochondrial genomes of jakobids,
heteroloboseans and malawimonads stands out as an obvious candidate for the
miT2SS substrate, since they either have well-established roles in the
mitochondrion or are hypothetical proteins with a restricted (genus-specific)
distribution. Therefore, we hypothesize that the substrate is encoded by the nuclear
genome and imported into the mitochondrion to undergo a specific processing step.
This may include addition of a prosthetic group – a scenario modelled on the

process of cytochrome *c* or Rieske protein maturation[41,42]. Interestingly, the
proteins Gcp6 and Gcp12, each exhibiting an array of absolutely conserved cysteine
and histidine residues (Supplementary Fig. 11), are good candidates for proteins
that are loaded with a specific prosthetic group, so any of them may well be the
sought-after miT2SS substrate. Some of the other Gcp proteins may then represent
components of the hypothetical machinery responsible for the substrate
modification. The putative functionalization step may occur either in the
mitochondrial matrix or in the intermembrane space (IMS), but we note that the
former localization would necessitate a mechanism of protein translocation across
the mitochondrial IM in the direction from the matrix to the IMS, which has not been
demonstrated yet. Regardless, the modified protein would eventually be
translocated across the mitochondrial OM by the T2SS system to the cytoplasm.
However, this may not be the end of the journey, since there are hints of a
link between the miT2SS-associated pathway and peroxisomes. First, three Gcp
proteins, namely Gcp1 to Gcp3, are specifically related to Pex7, a protein mediating
import of peroxisomal proteins characterized by the peroxisomal targeting signal 2
(PTS2)[43]. Second, some of the Gcp proteins (Gcp1, Gcp2, Gcp13) have at the C-
terminus a predicted PTS1 signal (at least in some species; Supplementary Table 3).
Third, several Gcp proteins (Gcp2, Gcp9, Gcp10, and Gcp11) and GspEL1 were
assigned to the putative peroxisomal proteome in our proteomic analysis
(Supplementary Table 2B). We note the discrepancy between the PTS1 signal
predictions and the actual set of experimentally defined peroxisomal proteins,
which might be due to an incomplete separation of peroxisome and mitochondria by
our purification procedure, but may also reflect protein shuttling between the two
organelles.
We thus hypothesize that upon its export from the mitochondrion, the
miT2SS substrate is eventually delivered to the peroxisome. This is possibly
mediated by the Gcp1/2/3 trio, but other Gcp proteins might participate as well.
One such protein might be the ubiquitin-related protein Gcp4. Ubiquitination and
deubiquitination of several components of the peroxisome protein import
machinery is a critical part of the import mechanism[43] and Gcp4 could serve as an
analogous peptide modifier in the hypothetical novel peroxisome import pathway
functionally linked to the miT2SS.
Altogether, our data suggest the existence of a novel elaborate functional
pathway combining components of bacterial origin with newly evolved eukaryote-
specific proteins. The modern phylogenetic distribution of the pathway is sparse,
but our current understanding of eukaryote phylogeny suggests that it was
ancestrally present in eukaryotes and for some reason dispensed with, multiple
times during evolution. Although we could not define a specific bacterial group as
the actual source of the eukaryotic Gsp genes, it is tempting to speculate that the
T2SS was introduced into eukaryotes by the bacterial progenitor of mitochondria
and that it was involved in delivering specific proteins from the endosymbiont into
the host cell, as is known in the case of current intracellular bacteria[36]. Elucidating
the actual role of this communication route in establishing the endosymbiont as a
fully integrated organelle requires understanding the cellular function of the
modern miT2SS-associated pathways, which is a challenge for future research.

Methods

Sequence data and homology searches

Homologues of relevant genes/proteins were searched in sequence databases accessible via the National Center for Biotechnology Information BLAST server (https://blast.ncbi.nlm.nih.gov/Blast.cgi), including the nucleotide and protein non-redundant (nr) databases, whole-genome shotgun assemblies (WGAs), expressed sequence tags (ESTs), and transcriptome shotgun assemblies (TSAs). Additional public databases searched included the data provided by the Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP[44]) comprising TSAs from hundreds of diverse protists (https://www.imicrobe.us/#/projects/104), the OneKP project[45] (https://sites.google.com/a/ualberta.ca/onekp/) comprising TSAs from hundreds of plants and algae, and individual WGAs and TSAs deposited at various on-line repositories (Supplementary Table 5). Non-public sequence data analysed included genome and/or transcriptome assemblies from several heteroloboseans, jakobids and malawimonads generated in our laboratories using standard sequencing technologies (454 and or Illumina) and sequence assembly programs (Supplementary Table 5). Details on the sequencing and assembly and full analyses of these genomes and transcriptomes will be published elsewhere.

Homology searches were done using BLAST[46] (blastp or tblastn, depending on the database queried) and HMMER[47] using profile HMMs built from sequence alignments of proteins of interest. Hits were evaluated by BLAST (blastp or blastx) searches against the nr protein dataset at NCBI to distinguish orthologues of Gsp and Gcp proteins from paralogous proteins or non-specific matches. This was facilitated by a high degree of conservation of individual eukaryotic Gsp/Gcp proteins among different species (see also Supplementary Figs 4 and 10-12) and in most cases by the lack of other close homologues in eukaryotic genomes (the exceptions being members of broader protein families, including the ATPase GspE, the WD40 superfamily proteins Gcp1 to Gcp3, and the ubiquitin related protein Gcp4). All identified eukaryotic Gsp and Gcp sequences were carefully manually curated to ensure maximal accuracy and completeness of the data, which included correction of existing gene models, extension of truncated sequences by manual analysis of raw sequencing reads, and correction of assembly errors (for details see Supplementary Methods). All newly predicted or curated Gsp and Gcp sequences are provided in Supplementary Tables 1 and 3, respectively; additional Gsp and Gcp sequences from non-target species are listed in Supplementary Table 4.

Phylogenetic profiling

In order to identify genes with the same phylogenetic distribution as the eukaryotic homologues of the four core T2SS components, we carried out two partially overlapping analyses based on defining groups of putative orthologous genes in select Gsp-positive species and phylogenetically diverse Gsp-negative eukaryotic species. The list of taxa included is provided in Supplementary Table 6. The first analysis was based on 18 species, including three Gsp-positive ones (*N. gruberi*, *A. godoyi* and *M. jakobiformis*), for the second analysis the set was expanded by adding

one additional Gsp-positive species (*G. okellyi*) and one Gsp-negative species
(*Monocercomonoides* sp. PA203). Briefly, the protein sequences of a given species
were compared to those of all other species using blastp followed by fast
phylogenetic analyses and orthologous relationships between proteins were then
inferred from this set of phylogenetic trees using a reference-species-tree-
independent approach. This procedure was repeated for each species and all
resulting sets of orthologous relationships, also known as phylomes[48], were
combined in a dense network of orthologous relationships. This network was finally
trimmed in several successive steps to remove week or spurious connections and to
account for (genuine or artificial) gene fusions, with the first analysis being less
restrictive than the second. Details of this pipeline are provided in Supplementary
Methods.
For each of the two analyses, the final set of defined groups of orthologs
(orthogroups) was parsed to identify those comprising genes from at least two Gsp-
positive species yet lacking genes from any Gsp-negative species. The orthogroups
passing this criterion were further analysed manually by blastp and tblastn searches
against various public and private sequence repositories (see the section "Sequence
data and homology searches") to exclude those orthogroups with obvious orthologs
in Gsp-negative species. *Percolomonas* spp. exhibiting only GspD and jakobids
represented by incomplete EST surveys (these species are likely to possess the
miT2SS system) were not considered as Gsp-negative. The orthogroups that
remained were then evaluated for their conservation in Gsp-positive species and
those that proved to have a representative in all these species (*N. gruberi*, *N. fowleri*,
N. damariscottae, *P. kirbyi*, *A. godoyi*, *R. americana*, *M. jakobiformis*, *G. okellyi*) were
considered as bona fide Gcp (Gsp-co-occurring protein) candidates. It is of note that
some of these proteins are short and were missed by the automated annotation of
some of the genomes, so using relaxed criteria for the initial consideration of
candidate orthogroups (i.e. allowing for their absence from some of the Gsp-positive
species) proved critical for decreasing the number of false-negative identifications.
627
Sequence analyses and phylogenetic inference
The presence of N-terminal mitochondrial transit peptides and peroxisomal
targeting signal 1 (PTS1) in the Gsp and Gcp proteins was evaluated using
MitoFates[49] (http://mitf.cbrc.jp/MitoFates/cgi-bin/top.cgi) and PTS1 predictor[50]
(http://mendel.imp.ac.at/pts1/), respectively. Transmembrane domains were
predicted using TMHMM[51] (http://www.cbs.dtu.dk/services/TMHMM/). Homology
of Gsp and Gcp protein families to other proteins was evaluated by searches against
Pfam v. 31 (ref.[52]; http://pfam.xfam.org/) and Superfamily 1.75 database[53]
(http://supfam.org/SUPERFAMILY/index.html) and by using HHpred[33]
(https://toolkit.tuebingen.mpg.de/#/tools/hhpred) and the Phyre2 server[34]
(http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index). The relative
position of the Gcp4 family among Ubiquitin-like proteins was analysed by a cluster
analysis using CLANS[54] (https://www.eb.tuebingen.mpg.de/protein-
evolution/software/clans/); for the analysis the Gcp4 family was combined with all
59 defined families included in the clan Ubiquitin (CL0072) as defined in the Pfam
database (each family was represented by sequences from the respective seed

alignments stored in the Pfam database). For further details on the procedure see
the legend of Supplementary Fig. 9A. Multiple sequence alignments used for
presentation of the conservation and specific sequence features of Gsp and Gcp
families were built using MUSCLE[55] and shaded using BioEdit
(http://www.mbio.ncsu.edu/BioEdit/bioedit.html)
In order to obtain datasets for the phylogenetic analyses of eukaryotic GspD
to GspG proteins, the protein sequences were aligned using MAFFT[56] and trimmed
manually. Profile hidden Markov models (HMMs) built on the basis of the respective
alignments were used as queries to search the UniProt database using HMMER. All
recovered sequences were assigned to components of the T4P superfamily
machineries using HMMER searches against a collection of profile HMMs reported
by Abby et al. (ref.[19]). For each GspD to GspG proteins, a series of alignments was
built by progressively expanding the sequence set by including more distant
homologues (as retrieved by the HMMER searches). Specifically, the different sets of
sequences were defined by the HMMER score based on the formula $score_{cutoff} =$
$c * score_{best\ prokaryotic\ hit}$, with the coefficient c decreasing from 0.99 to 0.70
incrementally by 0.01. The sequences were then aligned using MAFFT, trimmed
with BMGE[57] and the phylogenies were computed with IQ-TREE[58] using the best-fit
model (selected by the program from standard protein evolution models and the
mixture models[59] offered). The topologies were tested using 10,000 ultra-fast
bootstraps. The resulting trees were systematically analyzed for support of the
monophyly of eukaryotic sequences and for the taxonomic assignment of the
parental prokaryotic node of the eukaryotic subtree. The assignment was done
using the following procedure. The tree was artificially rooted between the
eukaryotic and prokaryotic sequences. From sub-leaf nodes to the deepest node of
the prokaryotic subtree, the taxonomic affiliation of each node was assigned by
proportionally considering the known or inferred taxonomic affiliations (at the
phylum or class level) of the descending nodes. See the legend to Supplementary Fig.
3 for further details.
The phylogenetic analysis of the WD40 superfamily including Gcp1 to Gcp3
proteins was performed as follows. The starting dataset was prepared by a
combination of two different approaches: 1) each identified sequence of Gcp1 to
Gcp3 proteins was used as a query in a blastp search against the non-redundant (nr)
NCBI protein database and the 500 best hits for each sequence were kept; 2) protein
sequences of each the Gcp1 to Gcp3 family were aligned using MAFFT and the
multiple alignment was used as a query in a HMMER3 search
(https://toolkit.tuebingen.mpg.de/#/tools/hmmer) against the UniProt database.
Best hits (E-value cutoff 1e-50) from all three searches were pooled and de-
duplicated, and the resulting sequence set (including Gcp1 to Gcp3 sequences) was
aligned using MAFFT and trimmed manually to remove poorly conserved regions.
Because WD40 proteins are extremely diversified, sequences that were too
divergent were removed from the starting dataset during three subsequent rounds
of sequence removal, based on a manual inspection of the alignment and
phylogenetic trees computed by IQ-TREE (using the best-fit model as described
above). The final dataset was enriched by adding PEX7 and WDR24 orthologues
from eukaryotes known to possess miT2SS components. The final phylogenetic tree

was computed using IQ-TEE as described in the legend to Supplementary Fig. 8. IQ-
TREE was used also for inferring trees of the heterolobosean 18S rRNA gene
sequences (Supplementary Fig. 1), ubiquitin-related proteins (Supplementary Fig.
9B) and the Gcp16 family (Supplementary Fig. 13); details on the analyses are
provided in legends to the respective figures.
695
Homology modelling
The PDB database was searched by the SWISS-MODEL server[60] for structural
homologues of *Go*GspD and *Go*GspG1. *V. cholerae* GspD[35] (PDB entry 5Wq9) and
K. oxytoca PulG[20] pseudopilus (PDB entry 5wda) were selected as the top matches,
respectively. Models were built based on the target-template alignment using
ProMod3[60]. Coordinates that were conserved between the target and the template
were copied from the template to the model. Insertions and deletions were
remodelled using a fragment library, followed by rebuilding side chains. Finally, the
geometry of the resulting model was regularized by using a force field. In the case of
loop modelling with ProMod3 fails, an alternative model was built with PROMOD-
II[61]. The quaternary structure annotation of the template was used to model the
target sequence in its oligomeric form[62].
708
Cultivation and fractionation of *N. gruberi* and proteomic analysis
Naegleria gruberi str. NEG-M was axenically cultured in M7 medium with PenStrep
(100 U/mL of penicillin and 100 µg/mL of streptomycin) at 27°C in vented tissue
culture flasks. Mitochondria of *N. gruberi* were isolated in seven independent
experiments, which were analyzed individually (see below). Each time ~1x10^9 *N.*
gruberi cells were resuspended in 2 mL of SM buffer (250 mM sucrose, 20 mM MOPS,
pH 7.4) supplemented with DNase I (40 ug/mL) and Roche cOmplete™ EDTA-free
Protease Inhibitor Cocktail and homogenized by eight passages through a 33-gauge
hypodermic needle (Sigma Aldrich). The resulting cell homogenate was then
cleaned of cellular debris using differential centrifugation and separated by a 2-hr
centrifugation in a discontinuous density OptiPrep gradient (10%, 15%, 20%, 30%
and 50%) as described previously[63]. Three visually identifiable fractions
corresponding to 10-15% (OPT-1015), 15-20% (OPT-1520) and 20-30% (OPT-
2023) OptiPrep densities were collected (each in five biological replicates) and
washed with SM buffer.
Proteins extracted from these samples were then digested with trypsin and
peptides were separated by nanoflow liquid chromatography and analyzed by
tandem mass spectrometry (nLC-MS2) on a Thermo Orbitrap Fusion (q-OT-IT)
instrument as described elsewhere[64]. The quantification of mass spectrometry data
in the MaxQuant software[65] provided normalized intensity values for 4,198
proteins in all samples and all three fractions. These values were further processed
using the Perseus software[66]. Data were filtered and only proteins with at least two
valid values in one fraction were kept. Imputation of missing values, which
represent low-abundance measurements, was performed with random distribution
around the value of instrument sensitivity using default settings of Perseus
software[66].

The data were analyzed by principle component analysis (PCA). The first two
loadings of the PCA were used to plot a two-dimensional graph. Based on a set of
marker proteins (376 mitochondrial and 26 peroxisomal, Supplementary Table 2),
clusters of proteins co-fractionating with mitochondria and peroxisomes were
defined and the proteins within the clusters were further analyzed. This workflow
was set up on the basis of the LOPIT protocol[67]. As a result, out of the 4,198 proteins
detected, 946 putative mitochondrial and 78 putative peroxisomal proteins were
defined. All proteins were subjected to *in silico* predictions concerning their function
(BLAST, HHpred[33]) and subcellular localization (Psort II,
https://psort.hgc.jp/form2.html; TargetP,
http://www.cbs.dtu.dk/services/TargetP/; MultiLoc2, https://abi.inf.uni-
tuebingen.de/Services/MultiLoc2). The mass spectrometry proteomics data have
been deposited in the ProteomeXchange Consortium via the PRIDE[68] partner
repository with the dataset identifier PXD007764.
749
Fluorescence *in situ* hybridization (FISH)
The PCR products of the *Ng*GspE and *Ng*GspF genes were labelled by alkali-stable
digoxigenin-11-dUTP (Roche) using DecaLabel DNA Labeling Kit (Thermo
Scientific). Labelled probes were purified on columns of QIAquick Gel Extraction Kit
(Qiagen, 28704) in a final volume of 50 µL. Labelling efficiencies were tested by dot
blotting with anti-digoxigenin alkaline phosphatase conjugate and CSPD
chemiluminescence substrate for alkaline phosphatase from DIG High Prime DNA
Labeling and Detection Starter Kit II (Roche) according to the manufacturer`s
protocol. FISH with digoxigenin-labelled probes was performed essentially
according to the procedure described in Zubacova at al. (ref.[69]) with some
modifications. *N. gruberi* cells were pelleted by centrifugation for 10 min at 2,000 x
g at 4°C. Cells were placed in hypotonic solution, fixed twice with a freshly prepared
mixture of methanol and acetic acid (3:1) and dropped on superfrost microscope
slides (ThermoScientific). Preparations for hybridizations were treated with RNase
A, 20 µg in 100 µL 2 x SSC, for 1 hr at 37°C, washed twice in 2 x SSC for 5 min,
dehydrated in a methanol series and air-dried. Slides were treated with 50% acetic
acid followed by pepsin treatment and postfixation with 2% paraformaldehyde.
Endogenous peroxidase activity of the cell remnants (undesirable for tyramide
signal amplification) was inactivated by incubation in 1% hydrogen peroxide,
followed by dehydration in a graded methanol series. All slides were denatured
together with 2 µL (25 ng) of the probe in 50 µL of hybridization mixture containing
50% deionised formamide (Sigma) in 2 x SSC for 5 min at 82°C. Hybridizations
were carried out overnight. Slides were incubated with tyramide reagent for 7 min.
Preparations were counterstained with DAPI in VectaShield and observed under an
Olympus IX81 microscope equipped with a Hamamatsu Orca-AG digital camera
using the Cell^R imaging software.
776
Heterologous gene expression, preparation of antibodies, and
immunodetection of Gsp proteins
The selected Gsp genes from *G. okellyi* and *N. gruberi* were amplified from
commercially synthesized templates (Genscript) (for primers used for PCR

amplification of the coding sequences see Supplementary Table 7) and cloned into
the pUG35 vector. The constructs were introduced into *S. cerevisiae* strain YPH499
by lithium acetate/PEG method. The positive colonies grown on SD-URA plates were
incubated with MitoTracker Red CMXRos (Thermo Fisher Scientific) and observed
for GFP and MitoTracker fluorescence (using the same equipment as used for FISH,
see above). For bacterial protein expression, *N. gruberi* GspD, GspE, GspF and GspG
genes were amplified from commercially synthesized templates and cloned into
pET42b vector (for primers used for PCR amplification of the coding sequences, see
Supplementary Table 6). The constructs were introduced into chemically-competent
E. coli strain BL21(DE3) and their expression induced by 1 mM IPTG. The
recombinant proteins were purified under denaturing conditions on Ni-NTA
agarose (Qiagen). The purified proteins were used for rat immunization in an in-
house animal facility at Charles University.
The sera obtained were used for immunodetection of Gsp proteins in *N.*
gruberi cells. Briefly, cells were fixed for 5 min in methanol (-20°C) and
permeabilized for 5 min by acetone (-20°C). The slides were incubated in blocking
buffer (BB) (PBS supplemented by 0.25% BSA, 0.05% TWEEN® 20 and 0.25%
gelatin) for 1 hr at room temperature. The slides were incubated overnight at 4°C
with primary antibodies diluted in BB and washed three times in PBS for 10 min.
Slides were then incubated for 1 hr with an anti-rat antibody conjugated with
Alexa488 (Thermo Fisher Scientific) diluted in. After washing three times for 10 min
in PBS, the slides were mounted in VectaShield DAPI solution and observed as above.
For mitochondrial labelling, the cells were incubated with MitoTracker Red CMXRos
for 30 min before fixation.
805
***In vitro* protein translation**
The *Go*GspD gene was amplified from the commercially synthesized template (for
primers used for PCR amplification of the coding sequences, see Supplementary
Table 6) and cloned into pDHFR vector provided in the PURExpress *In Vitro* Protein
Synthesis Kit (NEB). The 25 µl translation reaction contained 10 µL of solution A, 7.5
µL of solution B, 250 ng of pDHFR plasmid carrying *Go*GspD gene, 1 µL of an RNase
inhibitor (RNAsin, Promega), radioactively labelled ^{35}S-methionine, and 50 µg of
lecithin liposomes. The liposomes were prepared from a stock solution of soybean
L-α-lecithin in chloroform by evaporating the chloroform under a nitrogen flow,
resuspending the lipid film in dH_2O, and subsequent sonication in a waterbath
sonicator. The translation reaction was incubated for 2 hr at 37°C and then
centrifuged for 45 min at 13,000 x *g*. The pellet was resuspended in 50 mM sodium
phosphate buffer (pH = 8) with 2 M urea, centrifuged, and then washed in clear 50
mM sodium phosphate buffer. The output was analyzed by Blue Native PAGE using
2% digitonin and NativePAGE Novex 4-16% Bis-Tris Protein Gel (Thermo Fisher
Scientific).
822
Testing protein interactions using two-hybrid systems
Bacterial two-hybrid system (B2H) analysis was performed as described in ref.[70].
Gsp genes were amplified for commercially synthesized DNA and cloned into pKT25
and pUT18c plasmids. *E. coli* strain DHT1 competent cells were co-transformed with

two plasmids with different combinations of Gsp genes. Co-transformants were
selected on LB plates with ampicillin (100 μg/mL) and kanamycin (25 μg/mL).
Colonies were grown at 30°C for 48 to 96 hr. From each plate three colonies were
picked, transferred to 1 mL of LB medium with ampicillin and kanamycin, and
grown overnight at 30°C with shaking. Next day precultures (0.25 mL) were
inoculated to 5 mL of LB medium with ampicillin, kanamycin and 1 mM IPTG.
Cultures were grown with shaking at 30°C to OD_{600} of about 1-1.5. Bacteria (0.5 mL)
were mixed with 0.5 mL of Z buffer and subjected to the β-galactosidase assay[71].
The yeast two-hybrid system (Y2H) was employed as described in ref.[72]. Cells
of *S. cerevisiae* strain AH109 were co-transformed with two plasmids (pGADT7,
pGBKT7) with different combinations of Gsp genes. Co-transformants were selected
on double-dropout SD-Leu/-Trp and triple-dropout SD-Leu/-Trp/-His plates. The
colonies were grown for a few days. Positive colonies from the triple dropout were
grown overnight at 30°C with shaking and then the serial dilution test was
performed on double- and triple-dropout plates.
842
843 **Data availability**
844 All newly reported sequences of Gsp and Gcp proteins are provided in
845 Supplementary Table 1 and were deposited at GenBank with accession numbers
846 ######. Other relevant data (e.g. multiple sequence alignments used for
847 phylogenetic analyses) are available from the authors upon request.
848
849 **References**
850

1.   Roger, A. J., Muñoz-Gómez, S. A. & Kamikawa, R. The origin and diversification
of mitochondria. *Curr. Biol.* **27,** R1177–R1192 (2017).
2.   Martijn, J., Vosseberg, J., Guy, L., Offre, P. & Ettema, T. J. G. Deep mitochondrial
origin outside the sampled alphaproteobacteria. *Nature* **557,** 101–105 (2018).
3.   Leger, M. M. *et al.* An ancestral bacterial division system is widespread in
eukaryotic mitochondria. *Proc. Natl. Acad. Sci. U. S. A.* **112,** 10239–46 (2015).
4.   Beech, P. L. Mitochondrial FtsZ in a chromophyte alga. *Science (80-.).* **287,**
1276–1279 (2000).
5.   Natale, P., Brüser, T. & Driessen, A. J. M. Sec- and Tat-mediated protein
secretion across the bacterial cytoplasmic membrane—Distinct translocases
and mechanisms. *Biochim. Biophys. Acta - Biomembr.* **1778,** 1735–1756
(2008).
6.   Costa, T. R. D. *et al.* Secretion systems in Gram-negative bacteria: structural
and mechanistic insights. *Nat. Rev. Microbiol.* **13,** 343–359 (2015).
7.   Dolezal, P., Likic, V., Tachezy, J. & Lithgow, T. Evolution of the molecular
machines for protein import into mitochondria. *Science* **313,** 314–8 (2006).
8.   Palmer, T. & Berks, B. C. The twin-arginine translocation (Tat) protein export
pathway. *Nat. Rev. Microbiol.* **10,** 483–96 (2012).
9.   Lang, B. F. *et al.* An ancestral mitochondrial DNA resembling a eubacterial
genome in miniature. *Nature* **387,** 493–7 (1997).
10.   Burger, G., Gray, M. W., Forget, L. & Lang, B. F. Strikingly bacteria-like and
gene-rich mitochondrial genomes throughout jakobid protists. *Genome Biol.*

Evol. **5,** 418–38 (2013).

11. Tong, J. *et al.* Ancestral and derived protein import pathways in the
mitochondrion of *Reclinomonas america. Mol. Biol. Evol.* **28,** 1581–91 (2011).

12. Korotkov, K. V., Sandkvist, M. & Hol, W. G. J. The type II secretion system:
biogenesis, molecular architecture and mechanism. *Nat. Rev. Microbiol.* **10,**
336–51 (2012).

13. Thomassin, J.-L., Santos Moreno, J., Guilvout, I., Tran Van Nhieu, G. & Francetic,
O. The trans-envelope architecture and function of the type 2 secretion
system: new insights raising new questions. *Mol. Microbiol.* **105,** 211–226
(2017).

14. Berry, J.-L. & Pelicic, V. Exceptionally widespread nanomachines composed of
type IV pilins: the prokaryotic Swiss Army knives. *FEMS Microbiol. Rev.* **39,**
134–54 (2015).

15. Nivaskumar, M. & Francetic, O. Type II secretion system: a magic beanstalk or
a protein escalator. *Biochim. Biophys. Acta* **1843,** 1568–77 (2014).

16. Peabody, C. R. *et al.* Type II protein secretion and its relationship to bacterial
type IV pili and archaeal flagella. *Microbiology* **149,** 3051–72 (2003).

17. d'Enfert, C., Reyss, I., Wandersman, C. & Pugsley, A. P. Protein secretion by
gram-negative bacteria. Characterization of two membrane proteins required
for pullulanase secretion by *Escherichia coli* K-12. *J. Biol. Chem.* **264,** 17462–8
(1989).

18. Wang, X. *et al.* Cysteine scanning mutagenesis and disulfide mapping analysis
of arrangement of GspC and GspD protomers within the type 2 secretion
system. *J. Biol. Chem.* **287,** 19082–93 (2012).

19. Abby, S. S. *et al.* Identification of protein secretion systems in bacterial
genomes. *Sci. Rep.* **6,** 23080 (2016).

20. López-Castilla, A. *et al.* Structure of the calcium-dependent type 2 secretion
pseudopilus. *Nat. Microbiol.* **2,** 1686–1695 (2017).

21. Adl, S. M. *et al.* The revised classification of eukaryotes. *J. Eukaryot. Microbiol.*
59, 429–514 (2012).

22. Derelle, R. *et al.* Bacterial proteins pinpoint a single eukaryotic root. *Proc. Natl.*
Acad. Sci. U. S. A. **112,** E693-9 (2015).

23. Karnkowska, A. *et al.* A eukaryote without a mitochondrial organelle. *Curr.*
Biol. **26,** 1274–84 (2016).

24. Heiss, A. A. *et al.* Combined morphological and phylogenomic re-examination
of malawimonads, a critical taxon for inferring the evolutionary history of
eukaryotes. *R. Soc. Open Sci.* **5,** 171707 (2018).

25. Brown, M. W. *et al.* Phylogenomics places orphan protistan lineages in a novel
eukaryotic super-group. *Genome Biol. Evol.* **10,** 427–433 (2018).

26. Nouwen, N. *et al.* Secretin PulD: Association with pilot PulS, structure, and
ion-conducting channel formation. *Proc. Natl. Acad. Sci.* **96,** 8173–8177
(1999).

27. Hardie, K. R., Lory, S. & Pugsley, A. P. Insertion of an outer membrane protein
in *Escherichia coli* requires a chaperone-like protein. *EMBO J.* **15,** 978–88
(1996).

28. Korotkov, K. V, Pardon, E., Steyaert, J. & Hol, W. G. J. Crystal structure of the N-

terminal domain of the secretin GspD from ETEC determined with the
assistance of a nanobody. *Structure* **17,** 255–265 (2009).

29.   Yin, M., Yan, Z. & Li, X. Structural insight into the assembly of the Type II
secretion system pilotin-Secretin complex from enterotoxigenic *Escherichia*
coli. *Nat Microbiol* (2018). doi:10.2210/PDB5ZDH/PDB

30.   Guilvout, I. *et al.* In vitro multimerization and membrane insertion of bacterial
outer membrane secretin PulD. *J. Mol. Biol.* **382,** 13–23 (2008).

31.   Nivaskumar, M. *et al.* Pseudopilin residue E5 is essential for recruitment by
the type 2 secretion system assembly platform. *Mol. Microbiol.* **101,** 924–41
(2016).

32.   Korotkov, K. V. *et al.* Structural and functional studies on the interaction of
GspC and GspD in the type II secretion system. *PLoS Pathog.* **7,** e1002228
(2011).

33.   Alva, V., Nam, S.-Z., Söding, J. & Lupas, A. N. The MPI bioinformatics Toolkit as
an integrative platform for advanced protein sequence and structure analysis.
Nucleic Acids Res. **44,** W410–W415 (2016).

34.   Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. E. The
Phyre2 web portal for protein modeling, prediction and analysis. *Nat. Protoc.*
10, 845–858 (2015).

35.   Yan, Z., Yin, M., Xu, D., Zhu, Y. & Li, X. Structural insights into the secretin
translocation channel in the type II secretion system. *Nat. Struct. Mol. Biol.* **24,**
177–183 (2017).

36.   Nguyen, B. D. & Valdivia, R. H. Virulence determinants in the obligate
intracellular pathogen *Chlamydia trachomatis* revealed by forward genetic
approaches. *Proc. Natl. Acad. Sci.* **109,** 1263–1268 (2012).

37.   Tabach, Y. *et al.* Identification of small RNA pathway genes using patterns of
phylogenetic conservation and divergence. *Nature* **493,** 694–698 (2012).

38.   Nevers, Y. *et al.* Insights into ciliary genes and evolution from multi-level
phylogenetic profiling. *Mol. Biol. Evol.* **34,** 2016–2034 (2017).

39.   Okuno, D., Iino, R. & Noji, H. Rotation and structure of FoF1-ATP synthase. *J.*
Biochem. **149,** 655–664 (2011).

40.   Tomko, R. J. & Hochstrasser, M. Molecular architecture and assembly of the
eukaryotic proteasome. *Annu. Rev. Biochem.* **82,** 415–445 (2013).

41.   Babbitt, S. E., Sutherland, M. C., San Francisco, B., Mendez, D. L. & Kranz, R. G.
Mitochondrial cytochrome c biogenesis: no longer an enigma. *Trends Biochem.*
Sci. **40,** 446–55 (2015).

42.   Hartl, F. U., Schmidt, B., Wachter, E., Weiss, H. & Neupert, W. Transport into
mitochondria and intramitochondrial sorting of the Fe/S protein of ubiquinol-
cytochrome c reductase. *Cell* **47,** 939–51 (1986).

43.   Francisco, T. *et al.* Protein transport into peroxisomes: Knowns and
unknowns. *BioEssays* **39,** 1700047 (2017).

44.   Keeling, P. J. *et al.* The marine microbial eukaryote transcriptome sequencing
project (MMETSP): illuminating the functional diversity of eukaryotic life in
the oceans through transcriptome sequencing. *PLoS Biol.* **12,** e1001889
(2014).

45.   Matasci, N. *et al.* Data access for the 1,000 Plants (1KP) project. *Gigascience* **3,**

17 (2014).

46.   Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Res.* **25,** 3389–402 (1997).

47.   Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence
similarity searching. *Nucleic Acids Res.* **39,** W29-37 (2011).

48.   Huerta-Cepas, J., Dopazo, H., Dopazo, J. & Gabaldón, T. The human phylome.
Genome Biol. **8,** R109 (2007).

49.   Fukasawa, Y. *et al.* MitoFates: improved prediction of mitochondrial targeting
sequences and their cleavage sites. *Mol. Cell. Proteomics* **14,** 1113–26 (2015).

50.   Neuberger, G., Maurer-Stroh, S., Eisenhaber, B., Hartig, A. & Eisenhaber, F.
Prediction of peroxisomal targeting signal 1 containing proteins from amino
acid sequence. *J. Mol. Biol.* **328,** 581–92 (2003).

51.   Käll, L., Krogh, A. & Sonnhammer, E. L. L. Advantages of combined
transmembrane topology and signal peptide prediction--the Phobius web
server. *Nucleic Acids Res.* **35,** W429-32 (2007).

52.   Finn, R. D. *et al.* The Pfam protein families database: towards a more
sustainable future. *Nucleic Acids Res.* **44,** D279-85 (2016).

53.   de Lima Morais, D. A. *et al.* SUPERFAMILY 1.75 including a domain-centric
gene ontology method. *Nucleic Acids Res.* **39,** D427-34 (2011).

54.   Frickey, T. & Lupas, A. CLANS: a Java application for visualizing protein
families based on pairwise similarity. *Bioinformatics* **20,** 3702–3704 (2004).

55.   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and
high throughput. *Nucleic Acids Res.* **32,** 1792–7 (2004).

56.   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30,**
772–780 (2013).

57.   Criscuolo, A. & Gribaldo, S. BMGE (Block Mapping and Gathering with
Entropy): a new software for selection of phylogenetic informative regions
from multiple sequence alignments. *BMC Evol. Biol.* **10,** 210 (2010).

58.   Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies.
Mol. Biol. Evol. **32,** 268–74 (2015).

59.   Lartillot, N. & Philippe, H. A Bayesian mixture model for across-site
heterogeneities in the amino-acid replacement process. *Mol. Biol. Evol.* **21,**
1095–109 (2004).

60.   Bienert, S. *et al.* The SWISS-MODEL Repository-new features and functionality.
Nucleic Acids Res. **45,** D313–D319 (2017).

61.   Guex, N., Peitsch, M. C. & Schwede, T. Automated comparative protein
structure modeling with SWISS-MODEL and Swiss-PdbViewer: A historical
perspective. *Electrophoresis* **30,** S162–S173 (2009).

62.   Biasini, M. *et al.* SWISS-MODEL: modelling protein tertiary and quaternary
structure using evolutionary information. *Nucleic Acids Res.* **42,** W252-8
(2014).

63.   Jedelský, P. L. *et al.* The minimal proteome in the reduced mitochondrion of
the parasitic protist *Giardia intestinalis*. *PLoS One* **6,** e17285 (2011).

64.   Černá, M., Kuntová, B., Talacko, P., Stopková, R. & Stopka, P. Differential

regulation of vaginal lipocalins (OBP, MUP) during the estrous cycle of the
house mouse. *Sci. Rep.* **7,** 11674 (2017).
65.   Cox, J. *et al.* Accurate proteome-wide label-free quantification by delayed
normalization and maximal peptide ratio extraction, termed MaxLFQ. *Mol. Cell.*
Proteomics **13,** 2513–2526 (2014).
66.   Tyanova, S. *et al.* The Perseus computational platform for comprehensive
analysis of (prote)omics data. *Nat. Methods* **13,** 731–740 (2016).
67.   Dunkley, T. P. J., Watson, R., Griffin, J. L., Dupree, P. & Lilley, K. S. Localization
of organelle proteins by isotope tagging (LOPIT). *Mol. Cell. Proteomics* **3,**
1128–1134 (2004).
68.   Vizcaíno, J. A. *et al.* 2016 update of the PRIDE database and its related tools.
Nucleic Acids Res. **44,** 11033–11033 (2016).
69.   Zubáčová, Z., Krylov, V. & Tachezy, J. Fluorescence in situ hybridization (FISH)
mapping of single copy genes on *Trichomonas vaginalis* chromosomes. *Mol.*
Biochem. Parasitol. **176,** 135–137 (2011).
70.   Battesti, A. & Bouveret, E. The bacterial two-hybrid system based on
adenylate cyclase reconstitution in *Escherichia coli*. *Methods* **58,** 325–334
(2012).
71.   Miller, J. H. *Experiments in molecular genetics*. (Cold Spring Harbor Laboratory,
1972).
72.   Fields, S. & Song, O. A novel genetic system to detect protein-protein
interactions. *Nature* **340,** 245–246 (1989).

1033
1034

Acknowledgements

We would like to thank Michelle Leger and Alastair Simpson for granting us an
access, prior to publication, to their transcriptomic data from *A. godoyi* and *G. okellyi*,
respectively, which were instrumental in annotating the Gsp and Gcp genes in our
genome assemblies. This work was supported by Czech Science Foundation grants
13-29423S to PD and 18-18699S to ME and the KONTAKT II grant LH15253
provided by Ministry of Education, Youth and Sports of CR (MEYS) to PD. This work
was also supported by MEYS within the National Sustainability Program II (Project
BIOCEV-FAR, LQ1604) the project BIOCEV (CZ.1.05/1.1.00/02.0109), and the
project "Centre for research of pathogenicity and virulence of parasites" (No.
CZ.02.1.01/0.0/0.0/16_019/0000759) funded by European Regional Development
Fund (ERDF) and MEYS. The work in OF laboratory was funded by the ANR-14-
CE09-0004 grant. JP was supported by a grant from the Gordon and Betty Moore
Foundation to ADT. This work was supported by The Ministry of Education, Youth
and Sports from the Large Infrastructures for Research, Experimental Development
and Innovations project "IT4Innovations National Supercomputing Center –
LM2015070".

1052
1053

Author information

1055

Affiliations

Department of Parasitology, Faculty of Science, Charles University, BIOCEV,
Průmyslová 595, Vestec, 252 42, Czech Republic
L. Horváthová, V. Žárský, A. Krupičková, V. Klápšťová, M. Petrů, Z. Vaitová & P. Doležal

Department of Biology and Ecology, Faculty of Science, University of Ostrava,
Chittussiho 10, 710 00 Ostrava, Czech Republic
T. Pánek, V. Klimeš & M. Eliáš

School of Biosciences, University of Birmingham, Edgbaston, B15 2TT, UK
R. Derelle

Laboratory of Molecular & Evolutionary Parasitology, RAPID group, School of
Biosciences, University of Kent, Canterbury, CT2 7NZ, UK
J. Pyrih & A. D. Tsaousis

Department of Zoology, Faculty of Science, Charles University, Viničná 7, Prague 2, 128
44, Czech Republic
I. Čepička

Proteomic core facility, Faculty of Science, Charles University , BIOCEV, Průmyslová
595, Vestec, 252 42, Czech Republic
K. Harant

Department of Biochemistry and Molecular Biology and Centre for Comparative
Genomics and Evolutionary Bioinformatics, Dalhousie University, Halifax, NS B3H 4R2,
Canada
M. W. Gray

Institut Pasteur, Biochemistry of Macromolecular Interactions Unit, Department of
Structural Biology and Chemistry, CNRS UMR3528, 75015 Paris, France
I. Guilvout & O. Francetic

Robert Cedergren Centre for Bioinformatics and Genomics, Département de Biochimie,
Université de Montréal, Montreal, QC, Canada H3T 1J4
B. F. Lang

Institute of Molecular Genetics, Czech Academy of Sciences, 142 20 Prague 4, Czech
Republic
Č. Vlček

Author contributions
L.H. planned and carried out the experiments, V.Ž. conceived the original idea and carried out the bioinformatics analyses, T.P. carried out the genome and bioinformatic analyses, R.D. designed and carried out comparative genomic analyses, J.P. planned and carried out the experiments on *N. gruberi* mitochondrial proteome

and analysed the data, A.M. carried out the experiments, Ve.K. carried out the
experiments, Vl.K. participated in genome sequencing and analysis, M.P. planned
carried out the experiments, I.Č. participated in genome data acquisition, Z.V.
carried out the experiments, K.H. analysed the proteome of *N. gruberi* mitochondria,
M.W.G. contributed to the interpretation of the results and manuscript preparation,
I.G. designed and planned the experiments, O.F. designed, planned and carried out
the experiments and analyzed the data, B.F.L. provided the genome data and
analyses and contributed to manuscript preparation, Č.V. participated in genome
data acquisition, A.D.T. designed and planned the experiments, M.E. conceived the
idea, performed genomic analyses and wrote the manuscript, P.D. conceived the
idea, designed and performed experiments and wrote the manuscript.
1114
1115
1116 **Competing interests**
1117 None declared.
1118
1119 **Corresponding author**
1120 Correspondence to M. Eliáš or P. Doležal.
1121
1122
1123
1124
Figure Legends
Fig. 1 Some eukaryotes harbour homologues of core components of the bacterial
T2SS machinery. (A) Schematic representation of the complete bacterial T2SS;
subunits having identified eukaryotic homologues are highlighted in colour. (B)
Phylogenetic distribution of eukaryotic homologues of bacterial T2SS subunits (Gsp
proteins) and co-occurring proteins (Gcp). Core T2SS components (cyan),
eukaryote-specific T2SS components (dark blue), Gcp proteins carrying protein
domains found in eukaryotes (magenta), and Gcp proteins without discernible
homologues or with homologues only in prokaryotes (orange). Coloured sections
indicate proteins found to be present in genome or transcriptome data; white
sections, proteins absent from complete genome data; grey sections, proteins absent
from transcriptome data. The asterisk indicates the presence of the particular
protein in at least two of three species of *Percolomonas* analyzed. The two species
names in parentheses have not been yet been formally published. Sequence IDs and
additional details on the eukaryotic Gsp and Gcp proteins are provided in
Supplementary Table 1. (C) Maximum likelihood (ML) phylogenetic tree of
eukaryotic and selected bacterial GspF proteins demonstrating the monophyletic
origin of the eukaryotic GspF proteins and their separation from bacterial
homologues by a long branch (the tree inferred using IQ-TREE). Branch support
(bootstrap / posterior probability values) was assessed by ML ultrafast
bootstrapping and is shown only for branches where > 50.
1146
Fig. 2 Eukaryotic T2SS components are localized in mitochondria. (A) *N. gruberi*
cells labelled with specific polyclonal antibodies raised against GspD, GspF and

GspG1, and co-stained with MitoTracker red CMX ROS show mitochondrial
localization of the proteins; scale bar, 10 µm. (B) *S. cerevisiae* expressing *G. okellyi*
T2SS components as C-terminal GFP fusions co-stained with MitoTracker red CMX
ROS; scale bar, 10 µm.
1153
1154 **Fig. 3** Analysis of the *N. gruberi* mitochondrial proteome. PCA analysis of 4198
1155 proteins identified in the proteomic analysis of *N. gruberi* mitochondria. The cluster
1156 of mitochondrial proteins was defined on the basis of 376 mitochondrial markers.
1157 The boundaries of the cluster of co-purified peroxisomal proteins were defined by
1158 26 peroxisomal markers.
1159
Fig. 4. Mitochondrial GspD oligomerizes towards the formation of membrane pores.
(A) Domain architecture of the canonical bacterial GspD protein and eukaryotic
proteins homologous to its different parts. (B) Structural model of *Go*GspD built by
ProMod3 on the *Vibrio cholerae* GspD template. Top and side view of a cartoon and a
transparent surface representation of the *Go*GspD pentadecamer model is shown in
blue. The amphipathic helical loop (AHL), the signature of the secretin family, is
highlighted and coloured according to the secondary structure with strands in
magenta, helices in cyan and loops in light brown. The C-terminal GpsD residues are
highlighted as spheres. The detailed view of the AHL region shows the essential
residues V162 and F166 pointing towards the membrane surface. (C) Expression of
the mitochondrial *Go*GspD quickly induces cell death in bacteria. (D) Y2H assay
shows the self-interaction of the mitochondrial *Go*GspD. (E) *In vitro* translation and
assembly of mitochondrial *Go*GspD into a high-molecular-weight complex; lipo –
liposomes added, urea – extraction by 2M urea. (F) Y2H assay suggests the
interaction of *Ng*GspDN1 with itself and with *Ng*GspD.
1175
Fig. 5 Structure, maturation, and interactions of the mitochondrial GspG. (A)
Domain architecture of the bacterial and the mitochondrial pseudopilin GspG. The
arrow indicates the processing site of the bacterial GspG during protein maturation.
MTS – mitochondria targeting sequence, + – polar anchor, TMD – transmembrane
domain. (B) Positive interactions between the mitochondrial GspG protein and other
T2SS subunits were determined by the B2H assays. (C) Peptides specific to *Ng*GspG1
retrieved from the proteomic analysis of *N. gruberi* mitochondria. The arrow
indicates the position of the processing site of bacterial GspG proteins. (D)
Immunodetection of *Ng*GspG1 in *N. gruberi* cellular fractions. The arrow marks the
*Ng*GspG1-specific band.
1186
Fig. 6 Proteins with the same phylogenetic profile as the originally identified
mitochondrial Gsp homologues. (A) Schematic domain representation of 23 proteins
occurring in heteroloboseans, jakobids and malawimonads with the core T2SS
subunits but not in other eukaryotes analyzed. Proteins with a functional link to the
T2SS suggested by sequence homology are shown in royal blue, proteins
representing novel paralogues within broader (super)families are shown in red, and
proteins without discernible homologues or with homologues only in prokaryotes
are shown in yellow. The presence of conserved protein domains or characteristic

structural motifs is shown if detected in the given protein. Grey block – predicted
transmembrane domain (see also Supplementary Fig. 10); "C H C H" – the presence
of absolutely conserved cysteine and histidine residues (see also Supplementary Fig.
11) that may mediate binding of a prosthetic group. The length of the rectangles
corresponds to the relative size of the proteins. (B) Evolutionary relationships
among Gcp1 to Gcp3 proteins and other members of the WD40 superfamily. The
schematic phylogenetic tree was drawn on the basis of a ML phylogenetic tree
available as Supplementary Fig. 8.

Fig. 7 A hypothetical novel eukaryotic functional pathway including a mitochondrial
version of the T2SS (miT2SS) and connecting the mitochondrion with the
peroxisome. A nucleus-encoded protein (magenta) is imported via the TOM complex
into the mitochondrial inner membrane space, where it is modified by addition of a
specific prosthetic group catalysed by certain Gcp proteins. After folding it becomes
a substrate of the miT2SS machinery and is exported from the mitochondrion.
Finally it is imported into the peroxisome by the action of a dedicated import system
including other Gcp proteins. OMM – outer mitochondrial membrane, IMS –
intermembrane space, IMM – inner mitochondrial membrane, MM – mitochondrial
matrix.

Figure 1

A

B

Figure 2

Figure 3

Figure 4

A

bacteria + TMD pseudopilin

mitochondria *MTS* + TMD pseudopilin

B

β-galactosidase (Miller units)

neg.
GoGspG+GoGspG
GoGspG+GoGspF
GoGspF+GoGspF
GoGspE+GoGspE

C

MMHNSKTQPFSMMGHTPLSSERRKYRRMEQIIKHLIPFATIKSSSSIEENNYCTTSVNNNRKCENNNQMM *70*

EILQFLIRHILVTKRNNKDSINQLNIILQKYIQQTNGDFKKYLPIGIIAGLICANKYNQNNSTISNRFNI *140*

                                        VSNALLDLAK

VKNIFSNSAGILKDNNIAVTIHHLSMIEILIAISIIVTFSGLTGAVLAQVYEESRVSNALLDLAKLQEGL *210*

    YPLSLEDLLEGGELNK
    YPLSLEDLLEGGELNKVPK   ILLTTTTSNSGNNSTSQQQLTY
HGKYPLSLEDLLEGGELNKVPK DPWGTDYLYVPHLDWNR ILLTTTTSNSGNNSTSQQQLTY
VLYFTRHGKYPLSLEDLLEGGELNKVPKDPWGTDYLYVPHLDWNRLNRILLTTTTSNSGNNSTSQQQLTY *280*

FNEVLK   IDVDVSR
FNEVLKR FPSKIDVDVSR
FNEVLKRMETVLITLPGGVTPMSLLAIANEQPFCICVGTKIPRFPSKIDVDVSRERIRYVANLMKMSMER *350*

                                    IKNILNVTR
SSVAGSVQSNNQQGVTSIMNQITELNAR
SSVAGSVQSNNQQGVTSIMNQITELNARIKNILNVTRQASGGGE *394*

D

kDa L C M

35
25

Figure 5

A

B

Figure 6

Figure 7