Title: A simple model for the evolution of temperature-dependent sex determination explains the

temperature sensitivity of embryonic mortality in imperiled reptiles.

Authors: Lauren Lawson[*,**], Njal Rollinson[*,**]

Corresponding author: Njal Rollinson, njal.rollinson@utoronto.ca

Acknowledgements: We thank Anthony Rajkumar, Sami Troendle for constructive comments

on the original manuscript, and Melanie Massey for help compiling data. We also thank Daniel

Noble and two anonymous reviewers for their generous, constructive, and extensive feedback

that greatly improved the quality of this work. Funding was provided by an NSERC Discovery

Grant Number 2016-06469 to NR.

Lay summary: We analyze data on embryonic mortality under constant-temperature incubation

for 15 species of chelonians with temperature-dependent sex determination (TSD). Mortality is

lowest near species-specific pivotal temperatures (Tpiv) but increases rapidly above TPiv,

consistent with a theory that explains the adaptive significance of TSD. Conservation managers

should incubate embryos near TPiv.

Word count: 5,364

lauren.lawson@mail.utoronto.ca, njal.rollinson@utoronto.ca

[*]University of Toronto, Department of Ecology and Evolutionary Biology. 25 Willcocks Street,

Room 3055, Toronto, Ontario, Canada M5S 3B2.

[**]University of Toronto, School of the Environment. 33 Willcocks Street, Suite 1016V, Toronto

ON Canada M5S 3E8.

Title: A simple model for the evolution of temperature-dependent sex determination explains the

temperature sensitivity of embryonic stress in imperiled reptiles.

Authors:

Lauren Lawson, LL. University of Toronto, Department of Ecology and Evolutionary Biology.

25 Willcocks Street, Room 3055, Toronto, Ontario, Canada M5S 3B2.

Lauren.lawson@mail.utoronto.ca. 647-504-5515.

Njal Rollinson, NR. [1]University of Toronto, Department of Ecology and Evolutionary Biology.

25 Willcocks Street, Room 3055, Toronto, Ontario, Canada M5S 3B2. [2]University of Toronto,

School of the Environment. 33 Willcocks Street, Suite 1016V, Toronto ON Canada M5S 3E8.

Abstract

A common reptile conservation strategy involves artificial incubation of embryos and release of hatchlings or juveniles into wild populations. Temperature-dependent sex determination (TSD) occurs in most chelonians, permitting conservation managers to bias sex ratios towards females by incubating embryos at high temperatures, ultimately allowing the introduction of more egg-bearing individuals into populations. Here, we revisit classic sex allocation theory and hypothesize that TSD evolved in some reptile groups (specifically, chelonians and crocodilians) because male fitness is more sensitive to condition (general health, vigor) than female fitness. It follows that males benefit more than females from incubation environments that confer high-quality phenotypes, and hence high-condition individuals. We predict that female-producing temperatures, which comprise relatively high incubation temperatures in chelonians and crocodilians, are relatively stressful for embryos and subsequent life stages. We synthesize data from 28 studies to investigate how constant temperature incubation affects embryonic mortality in chelonians with TSD. We find several lines of evidence suggesting that female-producing temperatures, especially relatively warm temperatures, are more stressful than male-producing temperatures, and we find some evidence that pivotal temperatures (TPiv, the temperature that produces a 1:1 sex ratio) exhibit a correlated evolution with embryonic thermal tolerance. If patterns of temperature-sensitive embryonic mortality are also indicative of chronic thermal stress that occurs post hatching, then conservation programs may benefit from incubating eggs close to species-specific TPivs, thus avoiding high-temperature incubation. Indeed, our models predict that, on average, a sex ratio of more than 75% females can generally be achieved by incubating eggs only 1°C above TPiv. Of equal importance, we provide insight into the enigmatic evolution of TSD in chelonians, by providing support to the hypothesis that TSD

evolution is related to the quality of the phenotype conferred by incubation temperature, with

males produced in high-quality incubation environments.

Introduction

Despite recent and widespread interest in reptile conservation (Roll *et al.*, 2017), reptile

populations are declining globally (Todd *et al.*, 2010). Turtles, for example, are among the most

imperiled group of vertebrates in the world (Rhodin *et al.*, 2018; Gibbons and Lovich, 2019;

Stanford *et al.*, 2020). Natural rates of replacement and population growth are low in most turtle

species, and because of high juvenile mortality, slow life histories (e.g., Kondo, Morimoto, Sato,

& Suganuma, 2017), and low genetic diversity (Romiguier *et al.*, 2014), populations are unable

to adapt to environmental change on the time scale of anthropogenic impacts (Hawkes *et al.*,

2009). Elevated adult mortality of turtles arising from road collisions (Steen and Gibbs, 2004),

fishing gear entrapment (Lewison *et al.*, 2004; Bolten *et al.*, 2011), predation (Bolten *et al.*,

2011), and direct consumption as a food source (Conway-Gómez, 2007; Hancock *et al.*, 2017),

has therefore resulted in dramatic population declines. At present, 56.3% of data sufficient

species (51.9% of all recognized species) are considered critically endangered, endangered, or

vulnerable by the International Union for Conservation of Nature (Rhodin *et al.*, 2018).

Common turtle conservation initiatives include protecting nesting turtles from natural

predation and poachers (Eckert *et al.*, 1999), rehabilitation programs at trauma centers (Feck

and Hamann, 2013), education initiatives (Hassan *et al.*, 2017), and captive breeding (Bowkett,

2009). Another common initiative undertaken by zoos, parks, conservation authorities, and

even private individuals, involves incubation of reptile eggs in an ex-situ setting (Eckert *et al.*,

1999), followed by the release of hatchlings or juveniles. Growth of wild populations may

then be enhanced by protecting eggs and embryos, which reduces embryonic and hatchling

mortality, and/or rearing individuals to larger body sizes before release (Carstairs *et al.*, 2019;

Tetzlaff *et al.*, 2019), which may increase survival rates in the wild (Rollinson and Rowe,

2015).

Artificial incubation is an integral part of many conservation initiatives for reptiles.

Artificial incubation typically follows the removal of eggs from the wild (e.g., natural nests,

or gravid females that died on a road), or egg production after captive breeding. Embryos are

placed in incubators, hatched under a specified incubation regime, then released into the wild

(Páez *et al.*, 2015). Although the physical incubation of reptilian embryos may seem to

represent a small or insignificant portion of conservation programs, it is widely recognized

incubation temperature has a profound influence on morphology, performance, and fitness

components (Bobyn and Brooks, 1994a, 1994b; Booth, 2006; Noble *et al.*, 2018b). Further,

some lizards, all crocodilians, the tuatara, and most turtles exhibit temperature-dependent sex

determination (TSD), where sex is permanently affected by incubation temperature

(Valenzuela and Lance, 2004). The evolution and maintenance of TSD may in fact be related

to the effect of incubation temperature on fitness components (West, 2009), and as such,

exploring artificial incubation regimes for reptiles through the lens of TSD evolution may

provide insight into best practices for conservation programs.

In reptiles with TSD, the incubation temperatures experienced by an embryo during

the thermosensitive period influences sex, where the thermosensitive period comprises

specific anatomical stages that occur roughly during the middle third of embryonic

development (Yntema, 1968, 1979; Girondot *et al.*, 2018). Under constant temperature, the

temperature-sex reaction norm is known to take three forms (reviewed in Valenzuela and

Lance, 2004). The FMF pattern occurs when males are produced at intermediate temperatures

and females are produced at extreme temperatures. FMF is hypothesized to be ancestral, and it

is found in some turtles, lizards, and all crocodilians. The MF pattern occurs when males are

produced at cool temperatures and females at hot temperatures. Although turtles are the only

taxon to exhibit MF, turtles happen to comprise the majority of extant TSD species, and

therefore MF is the most common pattern of TSD. Finally, the FM pattern is when females are

produced at cool temperatures and males at hot temperatures; but this pattern is rare and found

only in a few lizards and the tuatara. Evolutionary explanations for TSD in lizards and the

tuatara (FM, and FMF) may be different than evolutionary explanations for TSD in turtles and

crocodilians (MF, and FMF) (e.g., Janzen and Phillips 2006). For instance, squamates are a

sister group to the tuatara (Rest *et al.*, 2003), whereas turtles are sister to the Archosaurs,

which includes crocodilians (Crawford *et al.*, 2012), underlining that fundamentally different

patterns of TSD are found in divergent evolutionary lineages.

In general, the evolution of TSD is hypothesized to arise by virtue of a sex-by-

temperature interaction for fitness (Charnov and Bull 1977). In short-lived squamates with

rapid maturation, the FM pattern of TSD may represent an adaptation to ensure females hatch

under the cool conditions that occur relatively early in the season, as to maximize time for

growth and achieve high fecundity during their short lives (Pen *et al.*, 2010, see also Conover,

1984). In contrast, turtles and crocodilians mature late and are very long lived, so TSD may be

an adaptation to an effect of incubation temperature on adult fitness that occurs over their

relatively longer lives (Figure 1). Specifically, males generally experience competition for

mates, and a small proportion of males often capture a large portion of mating opportunities

(Bateman 1948), with males in high "condition" (i.e., vigorous males in good health) securing

most mates (Rowe and Houle, 1996). Male fitness is therefore more sensitive to condition than

female fitness (Trivers and Willard, 1983), such that TSD evolves in age- and size-structured

populations to ensure that males are produced under temperatures that represent a high-quality

incubation environment, conferring relatively high post-hatching condition, ultimately

enhancing male fitness under strong sexual selection (Trivers and Willard, 1973; Deeming and

Ferguson, 1989; West, 2009). This hypothesis is an extension of Trivers-Willard (1973).

The FMF pattern can easily be reconciled with our hypothesis, where females are

produced in extreme environments (i.e., hot and cold incubation environments), and males at

intermediate, or "optimal", temperatures (Fig 1a). The MF pattern can, however, also be

reconciled with our hypothesis: because sex is determined by the amount of development that

occurs above vs below TPiv (Georges *et al.*, 1994), lower temperatures must have relatively

little effect on sexual differentiation as they have little developmental leverage (Rollinson *et*

al., 2018; Massey *et al.*, 2019), and therefore natural selection to maintain the lower TPiv is

weak in FMF species; this ultimately gives rise to MF species (Fig 1). Thus, a critical

prediction of our hypothesis is that fitness should decline with temperature at a relatively

faster rate above TPiv vs below the TPiv for MF species, and above vs below the upper TPiv

for FMF species. The reasoning is that male-producing temperatures should reflect the peak of

the performance curve, where a unit change in temperature has little influence on fitness,

whereas female-producing temperatures should represent the shoulder of the curve, where

fitness changes rapidly with temperature (Fig. 1). Therefore, performance metrics should be

relatively invariant with respect to temperature in the range of male-producing temperatures

(i.e, slightly cooler than TPiv), although male-producing temperatures that are far cooler than

TPiv may result in a decrease in performance.

We have outlined a theoretical ground to suggest that incubation temperature has

predictable and long-term effects on phenotypes and fitness components, especially in turtles

and crocodilians. Yet, incubation regimes in conservation programs for these taxa are broadly

unstandardized. For instance, some head-starting programs allow natural egg incubation

(Bona *et al.*, 2012), some allow temperature to fluctuate with above-ground ambient

temperature (Tetzlaff *et al.*, 2019), and others incubate different sets of eggs at different

constant temperatures (Carstairs *et al.*, 2019). A broad and theoretically informed exploration

of how incubation temperature affects fitness in embryos and hatchlings may therefore be

useful in developing best practices for artificial incubation in conservation programs.

Our goal is to better understand the evolution of TSD, and to inform best practices for

artificial incubation programs geared towards the conservation of reptiles, particularly for

chelonians. In the present study, we synthesize existing data to explore the fitness consequences

of artificial incubation regimes through the lens of TSD theory. We focus on patterns of

temperature-sensitive embryonic mortality in chelonians, as chelonians are relatively data rich,

and mortality is both a fitness component and is commonly measured. We generalize our

mortality findings by centering study-specific incubation temperatures with respect to species-

specific pivotal temperatures (TPiv), which is the constant temperature at which the sex ratio is

1:1 (Bull, 1980; Ewert *et al.*, 1994). We focus on a simple explanation for TSD that is specific to

chelonians and crocodilians (Figure 1), which is that males are produced in incubation

environments that confer high condition (Charnov and Bull, 1977; Deeming and Ferguson, 1989;

West, 2009). Several predictions arise from this hypothesis. The first is that embryonic thermal

tolerance exhibits a positive correlated evolution with TPiv, where thermal tolerance and TPiv

values evolve together. This pattern occurs because embryonic physiology is locally adapted to

incubation regimes (Ewert 1985), and TPiv evolution ensures that extreme temperatures within

local regimes produce female hatchlings, consistent with the adaptive explanation for TSD.

Therefore, we predict that (1) embryonic mortality will be best explained by the difference

between observed incubation temperature and population-specific TPivs, compared to the

difference between observed incubation temperature and mean incubation temperature, or simply

the observation incubation temperature. We also predict that (2) mortality will be positively

associated with the difference between incubation temperature and TPiv. Next, under the

adaptive explanation for TSD explored herein, temperature affects individual quality and males

are produced in high-quality thermal environments, but the association between sex and

individual quality arises through temperature. We therefore predict that (3) sex ratio (percent

male) will be negatively associated with embryonic mortality, but after accounting for

temperature, mortality will not be associated with sex. Finally, if TPiv evolves to delineate high-

quality thermal environments (for male production), from low quality thermal environments (for

female production), then (4) embryonic mortality in chelonians should increase with temperature

more rapidly above the TPiv, compared to below the TPiv. Mechanistically, fitness should

depreciate rapidly above the TPiv because TPiv evolves to be closer to the upper end of thermal

tolerance, rather than the lower end, and reaction rates for most or all processes changes more

rapidly near the upper tolerance limit (Sharpe and DeMichele, 1977; Kingsolver, 2009).

Methods

Data collection

Data was collected from previously published studies collated in the Reptile

Development Database (Noble *et al.*, 2018a). Only chelonians were considered in this study, as

they are of extreme conservation concern (Rhodin *et al.*, 2018; Gibbons and Lovich, 2019;

Stanford *et al.*, 2020), they represent the overwhelming majority of species in the world that

exhibit TSD, they feature a pattern of TSD different from squamates and the tuatara (see

below), and very limited data was available on crocodilians, especially for TPivs. Thus,

studies from the Reptile Development Database were included in the present study if they

featured (1) a turtle species with TSD (Sabath *et al.*, 2016), (2) artificial incubation under

constant temperature, (3) embryonic mortality (or survival) specific to a temperature treatment,

and (4) the number of embryos comprising the sample. Data collected from studies fitting

inclusion criteria included species, incubation temperature, TPiv, percent mortality per

incubation regime, number of eggs incubated, and geographic location of egg collection.

Exclusion factors included genotypic sex determined species (GSD), in situ incubation, or

fluctuating temperature trials. In studies including both TSD and GSD species, and studies

including both constant and fluctuating incubation trials, only TSD and constant trial results were

extracted. Trials involving temperature shifts were also excluded. Several studies report

mortality results in figures and do not specify results textually; in these cases we extracted data

from figures using WebPlotDigitzer (Rohatgi, 2019). We also extracted all available information

on substrate moisture; however, moisture was only measured directly in ten studies. Notably, all

studies either reported moisture values or maintained moisture at approximately constant levels

across the incubation regime; further, the few studies that manipulated moisture and temperature

did so in a factorial manner. It is therefore unlikely that moisture was confounded with

temperature in the present study.

Ultimately, 28 studies in the Reptile Development Database fit inclusion criteria for the

current study (Supplementary Table 1). In total, 199 data points were collected on a total of 15

different species in 6 different families of turtle, representing 43% of known taxonomic families.

Pivotal temperatures

Some papers that were used in the present study did not explicitly state the population-

specific TPiv(s). These values were supplemented with TPivs from other papers studying the

exact same population, or the most proximal population of the same species with a TPiv

estimate, as TPiv can vary geographically and among species (Bull, 1980; Ewert *et al.*, 1994;

Carter *et al.*, 2019). To accomplish this, we performed literature searches to find TPivs of species

using the species name, 'pivotal temperature', and 'sex ratios' as search terms. Species-specific

TPivs were found, and the closest geographically available population TPiv found was used as

the TPiv. In total, thirteen studies required supplementation of TPiv from previously published

literature (Supplementary Table 1).

There was only one species with FMF TSD (Pattern II), the snapping turtle (*Chelydra*

serpentina) in which both cool and hot temperatures produce females (Figure 1). For this species,

we use the upper TPiv in our analyses, since high temperature incubation is the focus of this

study; however, we perform all analyses with and without this species to ensure patterns are

robust regardless of the type of TSD.

Data Analysis
We reasoned that the relationship between temperature and embryonic mortality is likely

non-linear (e.g., Amarasekare and Savage, 2012). We therefore allowed flexibility in fixed

effects in all models (described below) by using basis splines within the package *splines* (R Core

Team, 2020). We chose basis splines over natural splines as basis splines are not constrained to

linearity at the tails, which is important as the tails of our functions are associated with few data

points, and basis splines therefore emphasize the uncertainty, rather than minimize it. Except

where noted, embryonic mortality was always the dependent variable, and it was weighted by

sample size of embryos in each temperature treatment. Given that many species were represented

by only one study, we chose to use only StudyID as a random intercept. We explored the

possibility of including basis splines as random slopes within StudyID, but these models

generally exhibited signatures of overfitting, and results were broadly robust whether or not

random slopes were included. Having selected our lone random effect to be applied in all

models, our models were subsequently fit using maximum likelihood. Model comparisons

outlined below were performed using second order Akaike Information Criterion (AIC_c) and

Akaike weights (w_i) within the package 'AICmodavg' in R (Fabozzi *et al.*, 2014; Mazzerole,

2019); basis spline regression was applied to fixed effects using the *bs* function in *splines* and

lmer in lme4 (Bates *et al.*, 2015).

We expect that TPiv exhibits a correlated evolution with embryonic thermal tolerance. To

address this expectation, we generated three plausible hypotheses for how incubation

temperature affects mortality, and we expressed these hypotheses as statistical models (Burnham

and Anderson 2002). The first and simplest hypothesis is that mortality is a function of

incubation temperature. In other words, the value of incubation temperature reported in the study

from which the data were collected best explains mortality. For clarity we refer to this variable as

"observed incubation temperature", or T_{OBS}. Second, we reasoned that TPiv and embryonic

thermal tolerance exhibit a correlated evolution, in which case embryonic mortality would be

best explained by the difference between TPiv and T_{OBS}. To create this new variable, we

subtracted T_{OBS} from species-specific TPivs, and we refer to this new variable as "deviation from

TPiv", or ΔT_{TPIV}. Finally, we reasoned that if the mean incubation temperature chosen by

researchers is generally similar to the mean temperature of the natural incubation environment,

then the mean temperature of a given study provides a good reference point for the average

incubation conditions. Mortality may then be related to the extremity of the incubation treatment

and would be best explained by the difference between the mean incubation temperature of the

study and T_{OBS}. To create this new variable, we subtracted each value of T_{OBS} from the mean

temperature of its respective study, and we refer to this new variable as "deviation from mean

temperature", or $\Delta T_{T\mu}$. We fit the three mixed linear models using maximum likelihood, where

the sole fixed-effect predictor of embryonic mortality was T_{OBS} (Model A), ΔT_{TPIV} (Model B), or

$\Delta T_{T\mu}$ (Model C). Models were compared using AICc. We ran these analyses twice, once with all

available data, which includes all FMF and MF species (n = 199), and again after excluding FMF

species, thereby including only MF species (n = 159).

We also investigated whether there is an association between sex ratio and embryonic

mortality, and if so, whether this association is realized over-and-above any temperature effect.

This analysis leverages the subset of data for which both temperature and sex ratio are measured

at a given incubation temperature. We fit the model that best-explained variation in embryonic

mortality (which was Model B, in which temperature is expressed as ΔT_{TPIV}, see Results), and

compared this model to two new models: Model D, in which mortality was expressed as a linear

effect of sex ratio (proportion of clutch male, or *PropMale*), and Model E, in which mortality

was expressed as a linear effect of *PropMale* and ΔT_{TPIV} (as in Model B). We ran separate

analyses for MF species alone, then for both FMF and MF species together (FM species: 19

studies, 13 species, 130 observations; FMF & MF species: 21 studies, 14 species, 158

observations). Finally, we also fit *PropMale* as a function of ΔT_{TPIV} , with StudyID as a random

effect, simply to visualize how *PropMale* changes with temperature across broad species.

Theory suggests mortality should increase more rapidly with temperature under female-

producing temperatures, compared to male producing temperatures (Figure 1). We therefore

tested whether the relationship between mortality and temperature is different above the TPiv

compared to below the TPiv. We took the absolute value of ΔT_{TPIV}, thereby creating a new

variable which we refer to as "absolute deviation from TPiv", or $|\Delta T_{TPIV}|$. We also created a

categorical variable "direction of deviation", which identifies the direction of the temperature

difference with respect to TPiv (i.e., above TPiv or below TPiv). We used model selection to

compare three models, and continuous fixed effects were fit as basis splines. First, we fit Model

F, in which mortality was expressed simply as $|\Delta T_{TPIV}|$. Next, we fit Model G, in which we fit

$|\Delta T_{TPIV}|$ and the additional term "direction of deviation", which explores whether average

mortality is different above vs below the TPiv. Finally, we fit Model H, which included an

interaction between "direction of deviation" and $|\Delta T_{TPIV}|$, which explores whether shape of the

relationship between embryonic mortality and temperature differs above vs below TPiv. We ran

these analyses twice, once with all available data, which includes all FMF and MF species (n =

199), and again after excluding FMF species, thereby including only MF species (n = 159).

Results

We tested whether embryonic mortality is best explained by incubation temperature per

se (T_{OBS}, Model A), the difference between incubation temperature and TPiv (ΔT_{TPIV}, Model B),

or the difference between incubation temperature and the mean incubation temperature of the

study ($\Delta T_{T\mu}$, Model C). We found no support for Model A (k = 6, LogLik = -356.4, ΔAIC = 43.3,

$w_i = 0$, n = 199) or Model C (k = 6, LogLik = -353.3, ΔAIC = 37.1, $w_i = 0$, n = 199). Instead,

Model B was strongly supported (k = 6, LogLik = -334.7, $w_i = 1$, n = 199). Visualization of

Model B indicates that the relationship between ΔT_{TPIV} and embryonic mortality is concave-

upward, centered below TPiv (Table S1, Figure 2a). We reran this analysis excluding all FMF

species, and Model B remained highly supported (k = 6, LogLik = -258.7, w_i = 1, n = 159, Figure

2b), whereas Model A (k = 6, LogLik = -273.0, ΔAIC = 28.5, w_i = 0, n = 159) and Model C (k =

6, LogLik = -278.6, ΔAIC = 39.6, w_i = 0, n = 159) remained unsupported.

Visualization of sex ratios (*PropMale*) as a function of ΔT_{TPIV} revealed the expected

patterns of variation: when data from FMF and MF species are combined there is a well-defined

peak of male producing temperatures below species-specific (upper) TPivs (Figure 3A), and

when MF species are considered alone there was relatively little variation in sex ratios below

species-specific TPiv (Figure 3B). Next, we investigated whether embryonic mortality is best

predicted by temperature or sex, and we did so by modelling whether mortality was associated

with *PropMale*, or ΔT_{TPIV}, or both *PropMale* and ΔT_{TPIV}. Of the three models compared, we

found that the model featuring *PropMale* as the sole predictor of embryonic mortality, Model D,

was poorly supported (k = 4, LogLik = 270.3, ΔAIC = 44.3, w_i = 0, n = 158). In fact, Model D

was poorly supported even though there was a linear association between *PropMale* and

embryonic mortality in this model ($b_{SexRatio} \pm SE$ = -1.02 $\pm$ 0.210, Figure 4a). The most strongly

supported model, Model B, featured ΔT_{TPIV} as the sole predictor of embryonic mortality (k = 6,

LogLik = -246.0, w_i = 0.75, n = 158). Finally, the model featuring both *PropMale* and ΔT_{TPIV},

Model E , was not supported: it differed from the best model (Model B) by one parameter and

likelihoods were nearly identical (see Burnham and Anderson, 2010, pg 131) (k = 7, LogLik = -

246.0, ΔAIC = 2.19, w_i = 0.25, n = 158). Importantly, the marginal association between

PropMale and embryonic mortality was weak in Model E, i.e., when controlling for ΔT_{TPIV}

($b_{SexRatio}$ = 0.0187 $\pm$ 0.322), indicating that temperature, not sex, influences mortality. Next, we

excluded all FMF data and reran the above analyses, revealing qualitatively identical results. We

found that Model D was unsupported (k = 4, LogLik = -216.9, ΔAIC = 36.2, w_i = 0, n = 130)

even though there was a non-trivial association between *PropMale* and embryonic mortality

($b_{SexRatio}$ = -0.638 ± 0.237, Figure 4B). Model B was again supported (k = 6, LogLik = -196.6, w_i

= 0.72, n = 130), and the marginal association between *PropMale* and embryonic mortality was

weak when controlling for ΔT_{TPIV} ($b_{SexRatio}$ = -0.200 ± 0.330).

We tested whether patterns of temperature-dependent mortality differed above vs below

species-specific TPivs. We compared three models, and we found that the model featuring an

interaction between $|\Delta T_{TPIV}|$ and 'Direction of Deviation', Model H, was strongly supported (k =

10, LogLik = -330.7, w_i = 1, n = 199). The nature of the interaction in Model H supported the

prediction that mortality increases relatively rapidly above the TPiv, compared to below the

TPiv. Other models were poorly supported, namely Model F (k = 6, Loglik = -346.3, ΔAIC =

22.5, w_i = 0, n = 199) and Model G (k = 7, LogLik = -340.0, ΔAIC = 12.1, w_i = 0, n = 199).

Next, we excluded FMF species and included only MF species in our analyses. Results were

similar, as the only model without the term 'Direction of Deviation' was poorly supported,

namely Model F (k = 6, Loglik = -261.3, ΔAIC = 6.92, w_i = 0.02, n = 159). Support was

otherwise similar between Model G (k = 7, LogLik = -257.2, ΔAIC = 0.93, w_i = 0.38, n = 159)

and Model H (k = 10, LogLik = -253.4, w_i = 0.60, n = 159, Figure 5B). Visualization again

supports the interpretation that mortality increases relatively rapidly above the TPiv, compared to

below the TPiv, at least for the first 5-6^{o}C above TPiv (Figure 5B).

Discussion

The present study presents a simple theory of TSD evolution (Figure 1) to explore how

constant temperature incubation is associated with embryonic mortality in chelonians. We

communicate four main findings. First, we show that embryonic morality is best explained by the

temperature difference between species-specific TPiv and incubation temperature (ΔT_{TPIV}).

Second, and similarly, we find mortality is lowest near TPiv, and highest when the difference

between TPiv and the incubation temperature is large. We interpret these first two patterns as

consistent with the prediction that TPiv exhibits correlated evolution with embryonic thermal

tolerance. Third, we found that embryonic mortality was negatively correlated with sex ratio

(proportion male) for both MF species alone and for FMF and MF species combined, but that sex

ratio was not associated with mortality after accounting for ΔT_{TPIV}. This is consistent with the

notion that temperature, not sex, exhibits an overarching effect on performance, and that female-

producing thermal environments are relatively stressful for embryos. Finally, and relatedly, we

demonstrate that embryonic mortality increases relatively rapidly with temperature above the

TPiv, compared to below the TPiv, which is also consistent with the hypothesis that female-

producing temperatures are relatively stressful for embryos. Below we explore these results and

emphasize their significance in evolution and conservation.

The relatively rapid increase in mortality with temperature above the TPiv has important

evolutionary implications. Mechanistically, fitness may depreciate rapidly above the TPiv

because TPiv evolves to be closer to the upper end of thermal tolerance, rather than the lower

end, as reaction rates for most or all processes changes more rapidly near the upper tolerance

limit (Sharpe and DeMichele, 1977; Kingsolver, 2009). Indeed, the lowest level of mortality

observed in the present study was just below TPiv, suggesting TPiv often falls toward the upper

limit of an intermediate thermal range that is optimal for development (Stubbs and Mitchell,

2018). One interpretation of our results, therefore, is that female-producing temperatures

coincide, at least roughly, with a departure from optimal incubation temperatures. Indeed, we

also found that sex ratio (percent male) *per se* was negatively correlated with mortality.

Although this pattern could arise from differential mortality of male embryos when overall

clutch mortality is relatively high, there is no obvious mechanism that would explain this

phenomenon. We therefore suggest that the negative correlation between sex ratio (percent male)

per se and embryonic mortality arises because female-producing temperatures are relatively

stressful. Our findings are therefore important as they are consistent with the simple explanation

for TSD explored herein (Figure 1). Specifically, classic theory suggests females in age-

structured populations should produce more sons when sons are expected to become high-

condition adults (Trivers and Willard, 1973), where condition refers to general health, energy

reserves, vigor, and overall quality. Therefore, if incubation temperature has, on average, an

effect on the general health and vigor of individuals at adulthood, then a sex by incubation

temperature interaction for fitness (Charnov and Bull, 1977) is inexorable when males are

produced under favourable incubation temperatures and females under unfavourable incubation

temperatures. We suggest that this explanation for TSD should be limited to the long-lived

chelonians and crocodilians, and not necessarily extended to squamates and the tuatara. This is

because chelonians and crocodilians share an evolutionary history, because neither group

features the FM pattern of TSD, and because the longevity of both chelonians and crocodilians

results in populations that are highly age- and size- structured, where variation in male quality

can be maintained.

Pivotal temperatures appear to be heritable and may be under weak selection

(Beukeboom and Perrin, 2014), and there is evidence, albeit mixed, that TPivs follow broad

environmental gradients in temperature (Ewert *et al.*, 2004, 2005; but see Carter *et al.*, 2019).

Evolution of TPivs is therefore possible, and another prediction from the simple theory of TSD

(Figure 1) is that TPiv and embryonic thermal tolerance experience a correlated evolution, as

would be necessary to ensure males are produced in relatively high-quality environments.

Admittedly, we define 'tolerance' fairly broadly, in that tolerance encompasses the range of

thermal environments that support normal embryonic development, and we assume that

embryonic mortality and tolerance are strongly associated.. Nevertheless, we found mortality

was best explained by ΔT_{TPIV} (Model B), rather than the difference between observed temperature

and the mean temperature used in the study ($\Delta T_{T\mu}$, Model C), or the observed incubation

temperature (T_{OBS}, Model A). One interpretation of this pattern is that TPiv and embryonic

thermal tolerance undergo a correlated evolution, although the extent to which this particular

finding supports our simple theory of TSD is debatable. We suggest that good evidence of a

correlated evolution of TPiv and embryonic thermal tolerance would indeed arise if ΔT_{TPIV}

(Model B) better-explained mortality patterns than deviation from the long-term average

environmental temperature, but our variable $\Delta T_{T\mu}$ (Model C) is the average temperature used in a

particular study, which is not necessarily the long-term average temperature of the local

environment. In sum, it is unclear if our study provides evidence for correlated evolution

between TPiv and embryonic thermal tolerance, consistent with TSD theory (Figure 1), or

whether relatively strong support for our Model B vs Model C arose because researchers choose

values of $\Delta T_{T\mu}$ that do not reflect the typical incubation environments of their focal populations.

In any event, it is necessary to study further whether TPivs are strongly linked with thermal

tolerance and physiology of the embryo, as this may provide insight into the adaptive

significance of TSD.

The overarching message communicated by the present study is that a constant

temperature used to produce females is associated with thermal stress, measured here as

mortality. Although we did not measure post-hatching phenotypes (but see below), we argue that

negative effects of thermal stress are likely to persist beyond the embryo stage. Indeed, persistent

phenotypic effects of thermal stress would have ramifications for incubation techniques used in

conservation programs, and such effects would be consistent with our simple model for TSD

evolution. Stress is a physiological state in which normal function is disrupted, and stress may

have a negative fitness impact (Walker *et al.*, 2005; Klockmann *et al.*, 2017). Our proxy of stress

(mortality) represents the most extreme fitness outcome that is possible, as mortality represents

an inability to persist, let alone reproduce (Kingsolver, 2009). Thermal stress can be induced at

temperatures which do not necessarily induce mortality but may induce phenotypic morbidity

(Cavallo *et al.*, 2015; Refsnider *et al.*, 2015; Kingsolver and Woods, 2016). Even within the

present study, for example, several studies reported abnormalities in hatchlings reared in high

temperature incubation trials (Packard *et al.*, 1987; Zhu *et al.*, 2006), and previous studies

suggest extreme temperatures lead to developmental instability (Noble *et al.*, 2018b). We

therefore suggest that our findings are probably not restricted to the effect of temperature on

embryos: the relationship between temperature and fitness observed herein (i.e., embryonic

mortality is lowest just below TPiv, and increases more rapidly above TPiv than below TPiv)

may persist into later life by virtue of the thermal stress survivors experienced as embryos.

The form in which thermal stress might persist in post-hatching phenotypes is unclear.

However, thirteen studies included in the present analysis included a measure of post-hatching

growth, performance metrics, or mortality, encompassing a total of twenty-three discrete

measurements (Table 1). Fifty-two percent of the measurements indicated a significant

temperature effect. Due to the heterogeneity of the papers and fitness metrics, and the multi-level

comparisons within some studies, a formal meta-analysis was not possible. However, our

inclusion criteria for the present study likely resulted in the exclusion of many studies examining

post-hatching growth or performance, such that it may prove fruitful for future research to

formally examine post-hatching growth and performance under different incubation regimes, as

evidence suggests the impacts of temperature stress may be observed at a variety of life stages

(Jonsson *et al.*, 2014; Noble *et al.*, 2018b).

If artificial incubation is geared toward conservation, then presumably artificial

incubation regimes fall within a normal range of incubation temperatures experienced by the

focal population. However, conservation managers face at least two reasons that incentivize

high-temperature incubation of turtle embryos. The first is that most turtles exhibit TSD, and for

turtles, relatively warm temperatures result in an over-production of female hatchlings (Ewert

et al., 1994). Given that a major goal of conservation programs is ultimately to increase the

number of wild births in the population, and births are more strongly related to female than male

abundance (Girondot *et al.*, 2008), TSD may represent an incentive to incubate at warm

temperatures. Second, high temperatures result in faster development rate and hence shorter

incubation time, potentially reducing the expense associated with husbandry, further

incentivizing high temperature incubation (Du *et al.*, 2006). Despite the incentives associated

with high-temperature incubation, our study suggests constant temperature incubation near TPiv

results in the highest hatching success and perhaps the least thermal stress. Notably, female

production does not require extreme temperatures, as female-biased sex ratios can be produced at

temperatures that are only slightly above the TPiv. For example, or analysis suggests that by

incubating embryos only 1ºC above TPiv (Figure 3), a sex ratio of over 75% females can be

achieved, on average, with a trivial increase in embryonic mortality (and presumably, a trivial

increase in post-hatching stress), although we note that the actual sex ratio achieved will depend

on the shape of the temperature-sex reaction norm at the population level (Bentley *et al.*, 2017;

Carter *et al.*, 2019). The notion of incubating embryos as near as possible to the pivotal

temperature may be especially useful for head starting programs, where incubation protocols can

be reviewed to prevent the release of hatchlings or juveniles that underwent significant thermal

stress during development.

While some studies suggest head-starting can be beneficial (Cunnington and Brooks,

1996; Crowder and Heppell, 2011; Mullin, 2019), the long-term benefits of head starting are

often unclear or negligible (Páez et al., 2015; Spencer, Van Dyke, & Thompson, 2017), as

factors such as predation and adverse environmental conditions result in low natural or even

negligible survival of hatchlings and juveniles in wild environments (Heppell, 1998; Enneson

and Litzgus, 2008; Altobelli, 2017). Although the present study does not definitively link

incubation environments with long-term survival as we do not examine post-hatching fates,

we suggest that incubation of embryos near TPiv may ultimately improve survival prospects

of individuals, especially females, released into the wild. As we have pointed out, incubation

near TPiv is compatible with sex ratio manipulation for many species and populations with

TSD. Indeed, it is desirable when increases in adult recruitment translate into increases in

juvenile recruitment, and this requires that careful attention is paid to the sex of released

individuals, as population size is not indicative of extinction risk when sex ratios are skewed

(Dale, 2001; Clout *et al.*, 2002; Wedekind, 2002; López-Sepulcre *et al.*, 2009).

Data syntheses and analyses are able to identify large-scale patterns, and ultimately help

strengthen evidence-based decision-making (Stewart, 2010). Indeed, it is useful for conservation

managers to periodically review methodologies used for species recovery, and with the

substantial increase in reptile conservation organizations since the 1990s, there are likely many

different initiatives that leverage artificial incubation (Gibbons and Lovich, 2019). The present

study underlines how constant temperature incubation in artificial egg incubation regimes affects

embryonic mortality, and possibly post-hatching thermal stress. We emphasize that we studied

chelonians in particular, but similar studies of crocodilians, and especially squamates and the

tuatara where different patterns of TSD can be found (Valenzuela and Lance 2004), would be

useful to uncover how TPiv relates to mortality in those groups. More broadly, we suggest

chelonian conservation programs leveraging artificial incubation may benefit from avoiding high

temperature incubation environments, and incubating embryos near species-specific TPivs.

Literature Cited

Altobelli JT (2017) Nest-Site Selection and Neonate Survival of Eastern Box Turtles (*Terrapene Carolina Carolina*) in Michigan's Northern Lower Peninsula (Masters of Science). Grand Valley State University.

Amarasekare P, Savage V (2012) A Framework for Elucidating the Temperature Dependence of Fitness. *The American Naturalist* 179: 178–191.

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. *J Stat Soft* 67. doi:10.18637/jss.v067.i01

Bentley BP, Haas BJ, Tedeschi JN, Berry O (2017) Loggerhead sea turtle embryos (*Caretta caretta)* regulate expression of stress response and developmental genes when exposed to a biologically realistic heat stress. *Mol Ecol* 26: 2978–2992.

Beukeboom LW, Perrin N (2014) The Evolution of Sex Determination, First edition. Edition. Oxford University Press, Oxford, United Kingdom□; New York, NY, United States of America.

Bobyn ML, Brooks RJ (1994a) Interclutch and interpopulation variation in the effects of incubation conditions on sex, survival and growth of hatchling turtles (*Chelydra serpentina*). *Journal of Zoology* 233: 233–257.

Bobyn ML, Brooks RJ (1994b) Incubation conditions as potential factors limiting the northern distribution of snapping turtles, *Chelydra serpentina*. *Can J Zool* 72: 28–37.

Bolten AB, Crowder LB, Dodd MG, MacPherson SL, Musick JA, Schroeder BA, Witherington BE, Long KJ, Snover ML (2011) Quantifying multiple threats to endangered species: an example from loggerhead sea turtles. *Frontiers in Ecology and the Environment* 9: 295–301.

Bona M, Novotny M, Danko S, Buresova A (2012) Headstarting in a small population of european pond turtles (*Emys orbicularis*) in Central European conditions: First results. *Hereptology Notes* 5.

Booth DT (2006) Influence of Incubation Temperature on Hatchling Phenotype in Reptiles. *Physiological and Biochemical Zoology* 79: 274–281.

Bowkett AE (2009) Recent Captive-Breeding Proposals and the Return of the Ark Concept to Global Species Conservation. *Conservation Biology* 23: 773–776.

Bull JJ (1980) Sex Determination in Reptiles. *The Quarterly Review of Biology* 55: 3–21.

Burnham KP, Anderson DR (2010) Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach, 2. ed. Edition. Springer, New York, NY.

Carstairs S, Paterson JE, Jager KL, Gasbarrini D, Mui AB, Davy CM (2019) Population
reinforcement accelerates subadult recruitment rates in an endangered freshwater turtle.
Anim Conserv 22: 589–599.

Carter AL, Bodensteiner BL, Iverson JB, Milne☐Zelman CL, Mitchell TS, Refsnider JM,
Warner DA, Janzen FJ (2019) Breadth of the thermal response captures individual and
geographic variation in temperature☐dependent sex determination. *Funct Ecol* 33: 1928–
1939.

Cavallo C, Dempster T, Kearney MR, Kelly E, Booth D, Hadden KM, Jessop TS (2015)
Predicting climate warming effects on green turtle hatchling viability and dispersal
performance. *Funct Ecol* 29: 768–778.

Charnov E, Bull J (1977) When is sex environmentally determined? *Nature* 266: 828–830.

Clout MN, Elliott GP, Robertson BC (2002) Effects of supplementary feeding on the offspring
sex ratio of kakapo: a dilemma for the conservation of a polygynous parrot. *Biological
Conservation* 107: 13–18.

Conover DO (1984) Adaptive Significance of Temperature-Dependent Sex Determination in a
Fish. *The American Naturalist* 123: 297–313.

Conway-Gómez K (2007) Effects of Human Settlements on Abundance of *Podocnemis unifilis*
and *P. expansa* Turtles in Northeastern Bolivia. *Chelonian Conserv Biol* 6: 199.

Crawford NG, Faircloth BC, McCormack JE, Brumfield RT, Winker K, Glenn TC (2012) More
than 1000 ultraconserved elements provide evidence that turtles are the sister group of
archosaurs. *Biol Lett* 8: 783–786.

Crowder L, Heppell S (2011) The Decline and Rise of a Sea Turtle: How Kemp's Ridleys Are
Recovering in the Gulf of Mexico. *Solutions* 2: 67–73.

Cunnington DC, Brooks RJ (1996) Bet-hedging theory and eigenelasticity: a comparison of the
life histories of loggerhead sea turtles (*Caretta caretta*) and snapping turtles (*Chelydra
serpentina*). *Can J Zool* 74: 291–296.

Dale S (2001) Female-biased dispersal, low female recruitment, unpaired males, and the
extinction of small and isolated bird populations. *Oikos* 92: 344–356.

Deeming DC, Ferguson MWJ (1989) The mechanism of temperature dependent sex
determination in crocodilians: a hypothesis. *American Zoologist* 29: 973–985.

Du W-G, Zheng R-Q, Shu L (2006) The Influence of Incubation Temperature on Morphology,
Locomotor Performance, and Cold Tolerance of Hatchling Chinese Three-Keeled Pond
Turtles, *Chinemys reevesii*. *Chelonian Conservation and Biology* 5: 294–299.

Eckert KL, IUCN/SSC Marine Turtle Specialist Group, World Wildlife Fund (U.S.), Center for
Marine Conservation, United States, National Oceanic and Atmospheric Administration,

International Union for Conservation of Nature and Natural Resources, Species Survival
Commission (1999) Research and Management Techniques for the Conservation of Sea
Turtles. IUCN/SSC Marine Turtle Specialist Group, Washington, D.C. (1725 De Sales
Street, NW #600, Washington 20036).

Enneson JJ, Litzgus JD (2008) Using long-term data and a stage-classified matrix to assess
conservation strategies for an endangered turtle (*Clemmys guttata*). *Biological*
Conservation 141: 1560–1568.

Ewert MA, Etchberger CR, Nelson CE (2004) Turtle sex-determining modes and TSD patterns,
and some TSD pattern correlates. In: Temperature Dependent Sex Determination in
Vertebrates (Eds. Valenzuela, N. & Lance, V). Smithsonian Institute, Washington, D.C,
pp 21–32.

Ewert MA, Jackson DR, Nelson CE (1994) Patterns of temperature-dependent sex determination
in turtles. *J Exp Zool* 270: 3–15.

Ewert MA, Lang JW, Nelson CE (2005) Geographic variation in the pattern of temperature-
dependent sex determination in the American snapping turtle (*Chelydra serpentina*). *J*
Zoology 265: 81–95.

Fabozzi FJ, Focardi SM, Rachev ST, Arshanapalli BG (2014) Appendix E Model Selection
Criterion: AIC and BIC. In: The Basis of Financial Econometrics: Tools, Concepts, and
Asset Management Applications. John Wiley & Sons Inc., pp 399–403.

Feck A, Hamann M (2013) Effect of sea turtle rehabilitation centres in Queensland, Australia, on
people's perceptions of conservation. *Endang Species Res* 20: 153–165.

Georges A, Limpus C, Stoutjesdijk R (1994) Hatchling sex in the marine turtle *Caretta caretta* is
determined by proportion of development at a temperature, not daily duration of
exposure. *J Exp Zool* 270: 432–444.

Gibbons JW, Lovich JE (2019) Where Has Turtle Ecology Been, and Where Is It Going?
Herpetologica 75: 4.

Girondot M, Fouillet H, Thinsp CP (2008) Feminizing Turtle Embryos as a Conservation Tool.
Conservation Biology 12: 353–362.

Girondot M, Monsinjon J, Guillon J-M (2018) Delimitation of the embryonic thermosensitive
period for sex determination using an embryo growth model reveals a potential bias for
sex ratio prediction in turtles. *Journal of Thermal Biology* 73: 32–40.

Hancock JM, Furtado S, Merino S, Godley BJ, Nuno A (2017) Exploring drivers and deterrents
of the illegal consumption and trade of marine turtle products in Cape Verde, and
implications for conservation planning. *Oryx* 51: 428–436.

Hassan R, Yahya NK, Ong LM, Kheng LK, Abidin ZZ, Ayob A, Jainal AM (2017) Public
Awareness Program and Develoment of Education Toolkit for Green Sea Turtle
Conservation in Sarawak, MALAYSIA. *ISESE* 12: 463–474.

Hawkes L, Broderick A, Godfrey M, Godley B (2009) Climate change and marine turtles.
Endang Species Res 7: 137–154.

Heppell SS (1998) Application of Life-History Theory and Population Model Analysis to Turtle
Conservation. *Copeia* 1998: 367.

Jonsson B, Jonsson N, Finstad A (2014) Linking embryonic temperature with adult reproductive
investment in Atlantic salmon Salmo salar. *Mar Ecol Prog Ser* 515: 217–226.

Kingsolver JG (2009) The Well☐Temperatured Biologist: (American Society of Naturalists
Presidential Address). *The American Naturalist* 174: 755–768.

Kingsolver JG, Woods HA (2016) Beyond Thermal Performance Curves: Modeling Time-
Dependent Effects of Thermal Stress on Ectotherm Growth Rates. *The American
Naturalist* 187: 283–294.

Klockmann M, Kleinschmidt F, Fischer K (2017) Carried over: Heat stress in the egg stage
reduces subsequent performance in a butterfly. *PLoS ONE* 12: e0180968.

Kondo S, Morimoto Y, Sato T, Suganuma H (2017) Factors Affecting the Long-Term Population
Dynamics of Green Turtles (*Chelonia mydas*) in Ogasawara, Japan: Influence of Natural
and Artificial Production of Hatchlings and Harvest Pressure. *Chelonian Conservation
and Biology* 16: 83–92.

Lewison RL, Freeman SA, Crowder LB (2004) Quantifying the effects of fisheries on threatened
species: the impact of pelagic longlines on loggerhead and leatherback sea turtles:
Fisheries effects on sea turtles. *Ecology Letters* 7: 221–231.

López-Sepulcre A, Norris K, Kokko H (2009) Reproductive conflict delays the recovery of an
endangered social species. *Journal of Animal Ecology* 78: 219–225.

Massey MD, Holt SM, Brooks RJ, Rollinson N (2019) Measurement and modelling of primary
sex ratios for species with temperature-dependent sex determination. *J Exp Biol* 222:
jeb190215.

Mazzerole MJ (2019) AICcmodavg: Model Selection and Multimodel Inference Based on
(Q)AIC.

Mullin DI (2019) Evaluating the Effectiveness of Headstarting for Wood Turtle (*Glyptemys
Insculpta*) Population Recovery (Masters of Science). Laurentian University, The Faculty
of Graduate Studies.

Noble DWA, Stenhouse V, Riley JL, Warner DA, While GM, Du W-G, Uller T, Schwanz LE
(2018a) A comprehensive database of thermal developmental plasticity in reptiles. *Sci*
Data 5: 180138.

Noble DWA, Stenhouse V, Schwanz LE (2018b) Developmental temperatures and phenotypic
plasticity in reptiles: a systematic review and meta-analysis: Incubation temperature and
plasticity. *Biol Rev* 93: 72–97.

Packard GC, Packard MJ, Miller K, Boardman TJ (1987) Influence of Moisture, Temperature,
and Substrate on Snapping Turtle Eggs and Embryos. *Ecology* 68: 983–993.

Páez VP, Lipman A, Bock BC, Heppell SS (2015) A Plea to Redirect and Evaluate Conservation
Programs for South America's Podocnemidid River Turtles. *Chelonian Conservation and*
Biology 14: 205–216.

Pen I, Uller T, Feldmeyer B, Harts A, While GM, Wapstra E (2010) Climate-driven population
divergence in sex-determining systems. *Nature* 468: 436–438.

R Core Team (2020) Splines. *R Foundation for Statistical Computing, Vienna, Austria*.

Refsnider JM, Palacios MG, Reding DM, Bronikowski AM (2015) Effects of a novel climate on
stress response and immune function in painted turtles (*Chrysemys picta*): NOVEL
CLIMATE EFFECTS ON TURTLE PHYSIOLOGY. *J Exp Zool* 323: 160–168.

Rest JS, Ast JC, Austin CC, Waddell PJ, Tibbetts EA, Hay JM, Mindell DP (2003) Molecular
systematics of primary reptilian lineages and the tuatara mitochondrial genome.
Molecular Phylogenetics and Evolution 29: 289–297.

Rhodin AGJ, Stanford CB, Dijk PPV, Eisemberg C, Luiselli L, Mittermeier RA, Hudson R,
Horne BD, Goode EV, Kuchling G, *et al.* (2018) Global Conservation Status of Turtles
and Tortoises (Order Testudines). *Chelonian Conservation and Biology* 17: 135.

Rohatgi A (2019) WebPlotDigitizer.

Roll U, Feldman A, Novosolov M, Allison A, Bauer AM, Bernard R, Böhm M, Castro-Herrera
F, Chirio L, Collen B, *et al.* (2017) The global distribution of tetrapods reveals a need for
targeted reptile conservation. *Nature Ecology & Evolution* 1: 1677–1682.

Rollinson N, Holt SM, Massey MD, Holt RC, Nancekivell EG, Brooks RJ (2018) A new method
of estimating thermal performance of embryonic development rate yields accurate
prediction of embryonic age in wild reptile nests. *Journal of Thermal Biology* 74: 187–
194.

Rollinson N, Rowe L (2015) Persistent directional selection on body size and a resolution to the
paradox of stasis: DIRECTIONAL SELECTION ON BODY SIZE. *Evolution* 69: 2441–
2451.

Romiguier J, Gayral P, Ballenghien M, Bernard A, Cahais V, Chenuil A, Chiari Y, Dernat R,
Duret L, Faivre N, *et al.* (2014) Comparative population genomics in animals uncovers
the determinants of genetic diversity. *Nature* 515: 261–263.

Rowe L, Houle D (1996) The lek paradox and the capture of genetic variance by condition
dependent traits. *Proc R Soc Lond B* 263: 1415–1421.

Sabath N, Itescu Y, Feldman A, Meiri S, Mayrose I, Valenzuela N (2016) Sex determination,
longevity, and the birth and death of reptilian species. *Ecol Evol* 6: 5207–5220.

Schwarzkopf L, Brooks RJ (1985) Sex determination in northern painted turtles: effect of
incubation at constant and fluctuating temperatures. *Can J Zool* 63: 2543–2547.

Sharpe PJH, DeMichele DW (1977) Reaction kinetics of poikilotherm development. *Journal of
Theoretical Biology* 64: 649–670.

Stanford CB, Iverson JB, Rhodin AGJ, Paul van Dijk P, Mittermeier RA, Kuchling G, Berry KH,
Bertolero A, Bjorndal KA, Blanck TEG, *et al.* (2020) Turtles and Tortoises Are in
Trouble. *Current Biology* 30: R721–R735.

Steen DA, Gibbs JP (2004) Effects of Roads on the Structure of Freshwater Turtle Populations.
Conservation Biology 18: 1143–1148.

Stewart G (2010) Meta-analysis in applied ecology. *Biol Lett* 6: 78–81.

Stubbs JL, Mitchell NJ (2018) The Influence of Temperature on Embryonic Respiration,
Growth, and Sex Determination in a Western Australian Population of Green Turtles
(*Chelonia mydas*). *Physiological and Biochemical Zoology* 91: 1102–1114.

Tetzlaff SJ, Sperry JH, Kingsbury BA, DeGregorio BA (2019) Captive-rearing duration may be
more important than environmental enrichment for enhancing turtle head-starting
success. *Global Ecology and Conservation* 20: e00797.

Todd B, Willson J, Gibbons J (2010) The Global Status of Reptiles and Causes of Their Decline.
In: Sparling D, Linder G, Bishop C, Krest S, eds. Ecotoxicology of Amphibians and
Reptiles, Second Edition. CRC Press, pp 47–67.

Trivers RL, Willard DE (1973) Natural Selection of Parental Ability to Vary the Sex Ratio of
Offspring. *Science* 179: 90–92.

Valenzuela N, Lance VA (2004) Temperature-Dependent Sex Determination in Vertebrates.
Smithsonian Institution.

Walker BG, Boersma PD, Wingfield JC (2005) Field endocrinology and conservation biology
45: 12–18.

Wedekind C (2002) Manipulating sex ratios for conservation: short-term risks and long-term
benefits. *Animal Conservation* 5: 13–20.

West SA (2009) Sex Allocation. Princeton University Press, Princeton.

Yntema CL (1968) A series of stages in the embryonic development of *Chelydra serpentina*. *J
733   Morphol* 125: 219–251.

734   Yntema CL (1979) Temperature levels and periods of sex determination during incubation of
735   eggs of *Chelydra serpentina*. *J Morphol* 159: 17–27.

736   Zhu X-P, Wei C-Q, Zhao W-H, Du H-J, Chen Y-L, Gui J-F (2006) Effects of incubation
737   temperatures on embryonic development in the Asian yellow pond turtle. *Aquaculture*
738   259: 243–248.

 Tables

758   Table 1: Post-hatching performance and growth temperature effect for papers included in the

759   current study.

Species	Trait	Effect	Is cooler better?	Study
Chinemys reevesii	Condition - body condition	No	-	Du et al. (2009)
Chinemys reevesii	Condition - body shape	No	-	Du et al. (2009)
Maureymys mutica	Condition - size	No	-	Zhao et al. (2015)
Maureymys mutica	Condition - size	Yes	No	Zhao et al. (2015)
Malaclemys terrapin	Growth	Yes/No*	-	Roosenburg and Kelley (1996)
Chrysemys picta	Growth	No	-	Les et al. (2009)
Trachemys scripta	Growth	No	-	Les et al. (2009)
Emys marmorata	Growth	No	-	Geist et al. (2015)
Ocadia sinensis	Growth	No	-	Du et al. (2010)
Chinemys reevesii	Growth	Yes**	-	Du et al. (2009)
Chinemys reevesii	Growth	Yes**	-	Du et al. (2007)
Gopherus agassizii	Growth	Yes	Yes	Spotila et al.(1994)
Mauremys mutica	Growth	Yes	Yes	Du et al. (2010)
Chelonia mydas	Growth	Yes	Yes	Booth et al. (2004)
Gopherus Polyphemus	Growth	Yes***	-	Demuth (2004)
Chelydra serptenina	Growth - mass	Yes	Yes/No****	Bobyn and Brooks (1994)
Podocnemis lewyana	Growth - straight line carapace	Yes	Yes	Paez et al. (2009)
Gopherus Polyphemus	Performance - crawl speed	No	-	Demuth (2001)
Maureymys mutica	Performance - righting response	No	-	Zhao et al. (2015)
Mauerymys	Performance - righting response	Yes	No	Zhao et al.

mutica				(2015)
Caretta caretta	Performance - righting response, crawl test, swim test	Yes	No	Fisher et al. (2014)
Chinemys reevesii	Performance - swimming ability	No	-	Du et al. (2009)
Chelonia mydas	Performance - swimming trials	Yes	No	Booth et al. (2004)

761   *Not able to determine if due to temperature or due to sex, ** Temperature effect gone after three months,

762   ***Temperature effect gone after 271 days, ****May be due to sexual dimorphism, intermediate (male temperature)

763   led to larger

Figure 1: Conceptual model for the evolution of TSD in turtles (and crocodilians), with explicit links between pivotal temperatures (TPiv), expected fitness, and reaction norms for sex. Types of TSD are (a) an "FMF" pattern (or Type II TSD) or (c) an "MF" pattern (or Type Ia TSD). An optimal incubation temperature produces high-quality phenotypes as in FMF species (panel a), or lower temperatures have low developmental leverage and have little influence on sex as in MF species (panel c). In either case, males should be produced under the conditions that maximize adult fitness for both sexes, as males stand to gain more fitness from good incubation environments, and to lose more fitness from bad incubation environments. This is equivalent to a sex-by-environment interaction for fitness, which gives rise to (b) the FMF reaction norm, or a special case of the FMF reaction norm, which is (d) the MF reaction norm. In panel (c) the lower TPiv (light grey) indicates females can be produced under extremely low temperatures in MF species.

Figure 2: The relationship between proportion clutch mortality and incubation temperature deviation from TPiv (ΔT_{TPIV}), which is the difference between incubation temperature and species-specific TPivs. In panel (a) all MF & FMF species are included, and in panel (b) FMF species are excluded. Points represent raw mortality data collected from published literature, the size of the point reflects relative sample size of each data point, and color reflects study ID. The 95% confidence intervals are in shaded grey.

Figure 3: Visualization of sex ratio (proportion of clutch male) as a function of the difference between observed incubation temperature and species-specific TPivs (ΔT_{TPIV}). Fitted line for sex is in blue. In grey, we overlaid the relationship between clutch mortality and ΔT_{TPIV}, estimated

from Model B (see Figure 2). (A) Both MF and FMF species are included in the model, and in (B) FMF species are excluded. Color of points indicate different studies and size of points indicates relative sample size. The single MF species in panel B with a prevalence of males below TPiv is the painted turtle (Schwarzkopf and Brooks, 1985).

Figure 4: Patterns of clutch mortality as a function of proportion of clutch male for (A) FMF and MF species, and (B) MF species only. Color of points indicate different studies. Size of points reflects relative sample size of raw data points.

Figure 5: Clutch mortality as a function of the absolute difference between incubation temperature and species-specific TPivs ($|\Delta T_{TPIV}|$) and the direction of the difference (above vs below TPiv) for (A) FMF and MF species, and (B) MF species only. Blue indicates raw data points below TPiv and red indicates raw data points above TPiv. Size of points reflects relative sample size of raw data points.