Sensitization of knee-innervating sensory neurons by tumor necrosis factor-α

activated fibroblast-like synoviocytes: an in vitro, co-culture model of inflammatory

pain.

Sampurna Chakrabarti, Luke A. Pattison, Charity N. Bhebhe, Gerard Callejo, David C. Bulmer,

Ewan St. John Smith.

Department of Pharmacology, University of Cambridge.

Corresponding author: Ewan St. John Smith

Department of Pharmacology, University of Cambridge

Tennis Court Road, Cambridge CB2 1PD.

Email: es336@cam.ac.uk

Keywords: synoviocytes, sensory neurons, pain, inflammation, tumor necrosis factor, complete

Freund's adjuvant, knee

Abbreviations:

FB = Fast Blue, FLS = Fibroblast-like synoviocyte, IL = Interleukin, PG = Prostaglandin, COX =

Cyclooxygenase, knee neurons = knee-innervating dorsal root ganglion neurons, CFA = Complete

Freund's adjuvant, CDH = Cadherin, CD = Cluster of differentiation, P = Passage, RT-PCR =

Reverse transcription – polymerase chain reaction, DRG = Dorsal root ganglion, RA =

Rheumatoid arthritis, OA = Osteoarthritis, MLS = Macrophage-like synoviocyte, MMP = Matrix

metalloprotease, VCAM = Vascular cell adhesion molecule, TNF = Tumor necrosis factor, AIA =

Antigen-induced arthritis, TRP = Transient receptor potential, ASIC = Acid-sensing ion channel,

CGRP = Calcitonin gene-related peptide, ECS = Extracellular solution, PBS = Phosphate-buffered

saline, Ipsi = CFA-injected side, Contra = Non-injected knee, RNA = Ribonucleic acid, DNA =

Deoxyribonucleic acid, L = Lumbar, KC = Keratinocyte chemoattractant, ANOVA = Analysis of

variance, AP = Action potential, RMP = Resting membrane potential, AHP =

Afterhyperpolarization, HPD = Half peak duration, LIX = Lipopolysaccharide induced

chemokine, MCP = Monocyte chemoattractant protein, MCSF = Macrophage colony stimulating

factor, MIG = Monokine induced by gamma interferon, MIP = Macrophage inflammatory protein,

RANTES = Regulated on activation, normal T-cell expressed and secreted, SDF= Stromal cell

derived factor, GM-CSF = Granulocyte-macrophage colony stimulating factor, GPCR = G protein-

coupled receptors.

Abstract (350 words)

Background:

Pain is a principle contributor to the global burden of arthritis with peripheral sensitization being

a major cause of arthritis-related pain. Within the knee joint, distal endings of dorsal root ganglion

neurons (knee neurons) interact with fibroblast-like synoviocytes (FLS) and the inflammatory

mediators they secrete, which are thought to promote peripheral sensitization. Therefore, we

investigated the communication between knee neurons and FLS in a co-culture system.

Methods:

Inflammation in FLS (isolated from mouse patella) was induced by stimulation with tumor

necrosis factor-α (TNF-FLS). Expression of relevant genes, secretion of cytokines and functional

acid response was assessed using polymerase chain reactions, inflammatory antibody array blots

and Ca^{2+} imaging respectively. Electrophysiology was performed on dissociated knee neurons in

mono-culture, co-culture with control FLS or TNF-FLS, or supernatant derived from TNF-FLS

cultures to determine electrical excitability and sensitivity to transient receptor potential (TRP)

agonists. Two group comparisons were conducted using Student's t-tests (distributed variable) or

chi-sq tests (categorical variable). ANOVA followed by Tukey's post-hoc test was performed for

comparing more than two groups.

Results:

Compared to control, TNF-FLS showed increased expression of interleukin-6 (*Il-6*, $p < 0.01$,

unpaired t-test) and *Il-1r1* ($p < 0.05$, unpaired t-test), and enhanced cytokine secretion. FLS also

responded to acidic stimuli and the percentage of TNF-FLS acid responders increased in the pH

range 6.0-5.0 ($p < 0.001$, chi-sq test). Co-culture of DRG neurons with TNF-FLS or supernatant

derived from TNF-FLS depolarized the resting membrane potential ($p < 0.01$, ANOVA), increased

spontaneous action potential firing ($p < 0.05$, ANOVA) and enhanced TRPV1 function ($p < 0.05$,

ANOVA) of knee neurons consistent with a role for FLS in the sensitization of pain sensing nerves

in arthritis.

Conclusions:

TNF-FLS have a pro-nociceptive phenotype, displaying enhanced acid sensitivity and increased

secretion of pro-inflammatory cytokines. Co-culture with TNF-FLS or supernatant from TNF-

FLS, induces hyperexcitability of knee neurons and enhances TRPV1 function. Data from this

study demonstrate the ability of FLS activated by TNF-α to promote neuronal sensitization, results

that highlight the importance of both non-neuronal and neuronal cells to the development of pain

in arthritis.

Background

Joint inflammation and pain are the major clinical symptoms of arthritis. Inflammation is part of

the body's immune response to tissue damage and involves multiple cell types, including

lymphocytes and synoviocytes. These non-neuronal cells are either in direct contact with, or in

close proximity to, the distal endings of dorsal root ganglion (DRG) sensory neurons that innervate

the joints. The physical interaction of sensory neurons with non-neuronal cells and the mediators

they secrete, is thought to cause peripheral sensitization of knee-innervating nerves, which is a key

pathology in the development of inflammatory pain.

The knee is a richly innervated, diarthrodial joint that is lined by the synovial membrane, which is

composed of synoviocytes in its intimal lining layer and lymphocytes in its sub-lining layer (1).

During inflammatory diseases such as rheumatoid arthritis (RA), the joint undergoes hyperplasia

due to T lymphocyte infiltration and fibroblast-like synoviocyte (FLS) proliferation. FLS are key

effector cells in RA as they become active upon stimulation by inflammatory cytokines (released

by macrophage-like synoviocytes, MLS, and T-cells) and secrete matrix metalloproteases (MMP)

that cause joint destruction (2). In addition, FLS can also support and maintain the ongoing

inflammation in arthritic joints by themselves secreting pro-inflammatory mediators (2). Building

on this, recent single-cell transcriptional analysis of patient-derived RA-FLS, and their subsequent

transfer into mouse knee joints, has identified distinct FLS subgroups: "destructive" and

"inflammatory" (3). Furthermore, deletion of cadherin-11 (CDH-11), a key surface marker of FLS,

causes hypoplasticity in murine joints and makes them resistant to inflammatory arthritis (4),

evidence further supporting a critical role of FLS in arthritic pathology.

FLS derived from rodent and human joints proliferate in vitro, unlike MLS, enabling their

extensive characterization (1). Interestingly, human RA-FLS are shown to be "imprinted

aggressors" that retain inflammatory phenotype over several passages, although the effect of these

activated FLS on sensory neurons remains largely unexplored (1,5). In rodents, it has been reported

that FLS derived from K/BxN mice (a spontaneous, chronic arthritis model) (5) and supernatants

from cultured FLS derived 3-days after induction of antigen-induced arthritis (AIA) in rats (6)

have increased levels of secreted cytokines compared to control FLS. Several other studies have

utilized interleukin 1β (IL-1β) or tumor necrosis factor-α (TNF-α) stimulation to robustly induce

an inflammatory phenotype in FLS (7–10). In this study we tested two modes of FLS activation to

understand their inflammatory phenotype. Firstly, we tested for pro-inflammatory phenotype in

FLS derived from mice undergoing the commonly used acute, unilateral complete Freund's

adjuvant (CFA) model, which, as previously reported, can produce robust knee inflammation, knee

neuron hyperexcitability and a concomitant decrease in digging behavior (11). Secondly, we tested

TNF-α (a cytokine that is upregulated in tissues derived CFA-injected mice (12)) stimulated FLS

derived from mouse knee, and determined their inflammatory phenotype to understand the

potential of activated FLS to induce sensitization of sensory neurons.

Since arthritic FLS reside in a pro-inflammatory environment which is often (13) (but not always

(14)) acidic, it is perhaps unsurprising that they express algogen sensing ion channels including

proton sensors (15,16) and various transient receptor potential (TRP) channels (17), such as

TRPV1 (transducer of noxious heat and low pH) and TRPA1 and TRPM8 (cold transducers).

Experiments conducted on SW982, a human tumor derived synoviocyte cell line, and primary FLS

from an RA patient, have demonstrated that a Ca^{2+} influx occurs upon application of the TRPV1

agonist capsaicin in an acidic (pH 7.1) environment and upon application of the TRPA1 and

TRPM8 agonist icilin (18). In addition to *Trpv1*, the human FLS cell lines SW982 and MH7A also

express *Trpv4* (warmth and tonicity transducer), and both *Trpv1* and *Trpv4* are upregulated by

TNF-α stimulation (19,20). Since most of these experiments were conducted on immortalized cell

lines, which thus might not be physiologically relevant, there is a need to investigate TRP channel

function in primary FLS. By contrast, acid sensors have been better studied in FLS derived from

mouse joints, leading to the identification of acid-sensing ion channel 3 (ASIC3) as a major proton

transducer. FLS from wild-type mice cultured in IL-1β show increased intracellular Ca^{2+} influx

and cell death in response to acid compared to FLS derived from ASIC3$^{-/-}$ mice, suggesting a

physiological role of ASIC3 and acid-sensing in general in FLS (16,21,22). Therefore, in this study

we sought to compare the algogen sensitivity of FLS before and after inducing an inflammatory

phenotype using Ca^{2+} imaging.

Given the complex interaction of cells in the joint environment, co-culture has proved to be an

important tool for dissecting FLS biology. Co-culture of human RA-FLS with T-cells (23–26) or

macrophages/monocytes (27–29) increased the concentration of prostaglandin (PG) E_2, IL-6, IL-

8, MMP-1 and MMP-3 in the culture medium, along with increased expression of adhesion

molecules such as vascular cell-adhesion molecule (*Vcam-1*) on FLS in a cell-mediated and

soluble-factor mediated manner. Furthermore, upregulation of these pro-inflammatory mediators

in culture supernatants was inhibited by anti-TNF-α (26,30) and anti-IL-6 antibodies (30).

Additionally, IL-15 secreted from human RA-FLS promotes survival of B-cells (31) and natural

killer cells (32) in co-culture, further demonstrating the importance of FLS in maintaining the

proinflammatory environment of arthritic joints. Although co-culture techniques have revealed

multiple immune interactions within the joint environment, the communication between neurons

and non-neuronal cells is less well understood in the context of knee inflammation. Culturing DRG

neurons with FLS derived from chronic AIA rats increased expression of receptors associated with

nociception, namely neurokinin 1, bradykinin 2 and TRPV1 in neurons (6), validating the premise

of co-culture studies to understand neuron-synoviocyte communication within the arthritic joint.

However, these studies did not functionally assess modulation of DRG neuron excitability, which

is a key mechanism of peripheral sensitization and hence pain. We recently showed that human

osteoarthritic (OA) synovial fluid (a lubricating fluid largely secreted by FLS (1)) can cause

hyperexcitability of murine sensory neurons and increase TRPV1 function (33). Therefore, in this

study, we hypothesized that after induction of inflammation, mouse knee-derived FLS will

increase excitability and TRP function of knee-innervating DRG neurons (knee neurons) in an

FLS-DRG neuron co-culture system.

Methods

Animals

All mice used in this study were 6-12 week old C57BL/6J mice (Envigo). Unless otherwise stated,

female mice were used since in humans, females are at a higher risk for inflammatory pain (34).

Mice were housed in groups of up to 5 in a temperature controlled (21 °C) room with appropriate

bedding materials, a red shelter and enrichment. They were on a 12-hour/light dark cycle with food

and water available *ad libitum*. Experiments in this study were regulated under the Animals

(Scientific Procedure) Act 1986, Amendment Regulations 2012. All protocols were approved by

a UK Home Office project license granted to Dr Ewan St. John Smith (P7EBFC1B1) and reviewed

by the University of Cambridge Animal Welfare and Ethical Review Body.

Knee injections

Under anesthesia (100 mg/kg ketamine and 10 mg/kg xylazine, intra-peritoneally) mice were

injected intra-articularly through the patellar tendon into each knee with the retrograde tracer, fast

blue (FB, 1.5 µl 2% in 0.9% saline, Polysciences) or into the left knee with 7.5 µl CFA (10 mg/ml,

Chondrex). Vernier's calipers were used to measure knee width (as before (11)) pre- and 24-hour

post-CFA injection.

Isolation and culture of FLS

24-hour after CFA injection into the knee, mice were killed by cervical dislocation and decapitation. Knee joints were exposed by removing the skin, then the quadriceps muscles were resected in the middle and pulled distally to expose the patellae. Patellae were then collected by cutting through the surrounding ligaments, as described before (35), in phosphate-buffered saline (PBS) and then transferred into one well of a 24-well plate with FLS culture media containing: Dulbecco's Modified Eagle Medium F-12 Nutrient Mixture (Ham) (Life Technologies), 25% fetal bovine serum (Sigma), 2 mM glutamine (Sigma) and 100 mg/ml penicillin/streptomycin (Life Technologies). Cells took approximately 10 days to grow to 70% confluency; medium was changed every other day. For P1, FLS were trypsinized with 1% trypsin (Sigma), re-suspended in FLS culture media and transferred into two wells of a 6-well plate. FLS from two animals were combined at P2. For subsequent passages, FLS were transferred into 60 mm dishes. Contralateral (Contra) and CFA-injected knees/cells (Ipsi) were kept separate at all stages. The cells were maintained in a humidified, 37 °C, 5% CO_2 incubator. FLS were also cultured until P5 from mice that had not undergone any knee CFA injection (control). A random selection of these dishes, from three separate biological replicates, was incubated for 24 to 48-hours (as per experimental design) in culture medium with recombinant mouse TNF-α (10 ng/ml from a stock solution of 100 μg/ml made up in 0.2 % bovine serum albumin and sterile PBS, R&D systems, aa-80325) to stimulate release of inflammatory mediators (TNF-FLS) (5).

Culture of Raw 264.7 cells

Raw 264.7 cells (EACC) were cultured in medium containing Dulbecco's Modified Eagle Medium

F-12 Nutrient Mixture (Ham) (Life Technologies), 10% fetal bovine serum (Sigma), 2 mM

glutamine (Sigma) and 100 mg/ml penicillin/streptomycin (Life Technologies). Cells were

maintained in a humidified 37 °C, 5% CO_2 incubator.

RNA extraction and reverse transcriptase – quantitative/ polymerase chain reaction (RT-q/PCR)

For all conditions, RNA was extracted from two 60 mm dishes (Thermo Fisher) of FLS (various

passages) and from one T-25 flask (Greiner Bio-one) of Raw 264.7 cells at P3 using the RNeasy

Mini Kit (Qiagen). 500 ng of the extracted RNA was used to synthesize cDNA using a High

Capacity cDNA RT kit (Applied Biosystems) following the manufacturer's guidelines, using a

T100 Thermal Cycler (Bio-Rad). The resultant cDNA was diluted to a 1:5 ratio with nuclease free

water and quantitative PCR (qPCR) was performed using a StepOnePlus Real Time PCR system,

following the manufacturer's guidelines on settings (Applied Biosystems) using TaqMan probes

(Thermo Fisher) (See Supplementary Table 1 in Additional File 1). The fluorescence intensity of

samples was captured during the last minute of each cycle. All reactions were run in triplicate with

appropriate negative controls with water containing no cDNA.

For qPCR reactions, data were obtained as Ct values (the cycle number at which fluorescent signals

emitted by the TaqMan probe crossed a threshold value). Only Ct values below 35 were analyzed

to determine ΔCt values by subtracting the Ct of *18S* ribosomal RNA from the Ct of target gene

(36). ΔCt values of target genes were subtracted from average ΔCt values of their controls to

calculate ΔΔCt, followed by 2^(- ΔΔCt) to calculate fold change (37).

FLS gene expression was also assessed by RT-PCR. DreamTaq Polymerase (Thermo Fisher) was

used to amplify a section within the open reading frames of various genes from 5 ng template

cDNA. The sequences of designed oligonucleotides (Sigma) are listed in Supplementary Table 1

in Additional File 1. Negative controls (using water/RNA as the template) were performed for

each biological sample with a randomly selected primer pairing. PCR products were resolved on

2% agarose containing 1X GelRed Nucleic Acid Stain (Biotum) and imaged with a GeneFlash Gel

Documentation System (Syngene). A randomly selected subset of reactions was repeated with 2.5

ng template cDNA to ensure reproducibility. Densitometry analyses were performed using ImageJ

Software (NIH) where relative expression was determined by dividing the band intensity of each

gene by that of the housekeeping gene, *18S* ribosomal RNA, for each biological replicate.

Cytokine antibody array

Before RNA extraction, 2 ml of culture media from P5 Contra, Ipsi, control and TNF-FLS (48-

hour) were collected and stored at -80 °C until use. Culture medium was pooled from three cultures

for each of the four conditions and assayed (undiluted) for the presence of 40 inflammatory

mediators using Mouse Inflammatory Antibody Array Membranes (ab133999, Abcam) according

to the manufacturer's instructions. Chemiluminescence was imaged using a BioSpectrum 810

imaging system (UVP) with 3 min exposure. Location of the 40 cytokines detected by the array

membranes are shown in Supplementary Table 2 in Additional file 1.

Densitometry of the spots in the array membranes was performed using ImageJ software (NIH).

Briefly, the mean gray value of each spot was measured from all membranes using the same

circular region of interest. The spots of interests were then background (average of all negative

control spots) subtracted and normalized to the positive control spots of the reference membrane

(control FLS media). Fold change was obtained by dividing normalized intensities of the

membrane of interest and the control membrane, analyte-by-analyte.

DRG neuron isolation and culture

Lumbar DRG (L2-L5, those that primarily innervate the knee) were collected from FB labelled

mice 7-10 days after knee injections in ice cold dissociation media containing L-15 Medium (1X)

+ GlutaMAX-l (Life Technologies), supplemented with 24 mM $NaHCO_3$. DRG were then

enzymatically digested in type 1A collagenase (Sigma) and trypsin solution (Sigma) at 37 °C,

followed by mechanical trituration as described before (11). Dissociated DRG neurons were plated

onto poly-D-lysine and laminin coated glass bottomed dishes (MatTek, P35GC-1.5-14-C) and

cultured either on their own (mono-culture), on a layer of FLS (co-culture, see below) or in 48-

hour conditioned media from TNF-FLS for 24-hours. The DRG culture medium contained L-15

Medium (1X) + GlutaMAX-l, 10% (v/v) fetal bovine serum, 24 mM $NaHCO_3$, 38 mM glucose,

2% (v/v) penicillin/streptomycin.

DRG neuron/FLS co-culture

For co-culture studies, FLS were plated onto MatTek dishes and cultured for 24-hours with FLS

culture media (with or without TNF-α stimulation). The next day medium was removed from FLS

plates, then DRG neurons were isolated as described above and plated on top of the FLS. Co-

culture plates were then kept in DRG culture medium for up to 24-hours for electrophysiogical

recording.

Cell staining

FLS immunocytochemistry: FLS were plated overnight in wells of a 24-well plate, fixed with

Zamboni's fixative (38) for 10 min, permeabilized with 0.05 % TritonX-100 and blocked with

antibody diluent (0.2% (v/v) Triton X-100, 5% (v/v) donkey serum and 1% (v/v) bovine serum

albumin in PBS) for 30 min. The cells were then incubated overnight at 4 °C in 1:100 (in antibody

diluent) anti-cadherin-11 antibody (CDH-11, rabbit polyclonal, Thermo Fisher, 71-7600). Cells

were washed three times with PBS-tween and incubated in the conjugated secondary antibody,

anti-rabbit Alexa-568 (1:1000 in PBS, Thermo Fisher, A10042) for 1-hour at room temperature

(21 °C). The secondary antibody was washed off three times with PBS-tween and the cells were

incubated in the nuclear dye DAPI (1:1000 in PBS, Sigma, D9452) for 10 min. Cells were further

washed with PBS-tween once and imaged in PBS using an EVOS FLoid Cell Imaging Station

(Thermo Fisher) at 598 nm (for CDH-11) and 350 nm (for DAPI) wavelength of light. Cells

without primary antibody did not show fluorescence.

DRG/FLS co-culture live cell stain: To visualize neuron/FLS co-culture, live cell imaging was

performed. FLS were plated on MatTek dishes with 1:1000 (diluted in FLS culture medium)

CellTracker Deep Red Dye (Thermo Fisher, C34565) and incubated for 24-hours in a humidified

37 °C, 5% CO_2 incubator. Dissociated DRG neurons (see above) were incubated in CellTracker

Green Dye (1: 1000 diluted in DRG culture media, Thermo Fisher, C7025) for 15 min at room

temperature (21 °C), centrifuged (16000 g, 3 min, 5415R, Eppendorf) and re-suspended in fresh

medium. The FLS dishes were then washed twice with PBS and the neuronal suspension was

plated on top of the FLS monolayer and incubated overnight in the incubator. The co-culture dishes

were washed with PBS and imaged the following day using an Olympus BX51 microscope and

QImaging camera at 650 nm (for deep red dye) and 488 nm (for green dye) wavelength of light.

Whole-cell patch-clamp electrophysiology

Neurons were bathed in extracellular solution containing (ECS, in mM): NaCl (140), KCl (4),

$MgCl_2$ (1), $CaCl_2$ (2), glucose (4) and HEPES (10) adjusted to pH 7.4 with NaOH. Only FB

labelled neurons identified by their fluorescence upon excitation with a 365 nm LED (Cairn

Research) were recorded. Patch pipettes of 5–10 MΩ were pulled with a P-97 Flaming/Brown

puller (Sutter Instruments) from borosilicate glass capillaries and the intracellular solution used

contained (in mM): KCl (110), NaCl (10), $MgCl_2$ (1), EGTA (1), HEPES (10), Na_2ATP (2),

Na_2GTP (0.5) adjusted to pH 7.3 with KOH.

Action potentials (AP) were recorded in current clamp mode without current injection (to

investigate spontaneous AP firing) or after step-wise injection of 80 ms current pulses from 0 –

1050 pA in 50 pA steps using a HEKA EPC-10 amplifier (Lambrecht) and the corresponding

Patchmaster software. AP properties were analyzed using Fitmaster software (HEKA) or IgorPro

software (Wavemetrics) as described before (11) and shown in Figure 3D (inset). Neurons were

excluded from analysis if they did not fire an AP in response to current injection. For recording

whole-cell voltage-clamp currents in response to TRP agonists capsaicin (1 µM from a 1 mM stock

in ethanol, Sigma-Aldrich), cinnamaldehyde (100 μM from a 1 M stock in ethanol, Alfa Aesar)

and menthol (100 μM from a 1 M stock in ethanol, Alfa Aesar), a 5 s baseline was established

with ECS, followed by a 5 s randomized drug application. Peak drug response was measured in

Fitmaster by subtracting the average of 2 s baseline immediately preceding the drug application

and the maximum peak response reached during the 5 s of drug application. Peak current density

is represented in graphs by dividing this peak response by the capacitance of the neuron. Data from

at least four mice were used in all conditions (with each mouse being used for at least two

conditions) and at least three neurons were recorded from each mouse in each category.

Ca^{2+} imaging

FLS or neurons were incubated with the Ca^{2+} indicator, Fluo-4 AM (10 μM diluted in ECS from

a 10 mM stock solution in DMSO, Invitrogen) for 30 min at room temperature (21 °C). Culture

dishes were then washed and imaged using an inverted Nikon Eclipse T*i* microscope. Fluo-4

fluorescence was excited using a 470 nm LED (Cairn Research) and captured with a camera (Zyla

cSMOS, Andor) at 1 Hz with 50 ms (for neurons) 250 ms (for FLS) exposure time using Micro-

Manager software (v1.4; NIH). Solutions were perfused in this system through a gravity-driven 12

barrel perfusion system (39). During imaging of neurons, MIP-1γ (10 and 100 ng/ml diluted in

ECS from a stock concentration of 0.1 mg/ml in 0.1% bovine serum albumin/sterile PBS, Sigma)

was applied for 20 s after establishing a baseline with ECS for 10 s. Neurons were allowed to

recover for 5 minutes between drug applications. 50 mM KCl was used as a positive control.

During imaging of FLS, a 10 s baseline was established with ECS and then pH 4-7, and TRP

agonists, 1 μM capsaicin, 100 μM cinnamaldehyde and 100 μM menthol were applied for 10 s

before a wash out period of 4 min between drug applications. TRP agonist sensitivity and acid

sensitivity was assessed on separate dishes. Three biological repeats were conducted on separate

days for each condition. 10 μM ionomycin was used as a positive control after all FLS Ca^{2+}

imaging experiments.

Analysis was conducted as described before (33). Briefly, mean grey values were extracted by

manually drawing around FLS or neurons in the ImageJ software. These values were then fed into

a custom-made R-toolbox (https://github.com/amapruns/Calcium-Imaging-Analysis-with-R.git)

to compute the proportion of cells responding to each drug and their corresponding magnitude of

response (normalized to their peak ionomycin (FLS) or KCl (neuron) response, (ΔF/Fmax); cells

not crossing threshold for positive controls were excluded from the analysis).

Statistics

Comparisons between two groups with distributed variables were performed using two-sided

Student's t-tests (paired if comparing two conditions of the same sample, unpaired otherwise) with

suitable corrections and amongst three groups using one-way analysis of variance (ANOVA)

followed by Tukey's post-hoc tests. Proportions were compared for categorical data using chi-sq

tests. Data are shown as mean ± SEM.

Results

1. TNF-FLS, but not FLS derived from CFA-induced inflamed knees, have a pro-

inflammatory phenotype and secrete cytokines that activate DRG neurons.

Key genes have been described that act as identifiers of cultured FLS isolated by enzymatic

digestion of mouse joints (5), as well as for establishing their inflammatory phenotype. In this

study, we performed RT-qPCR on adherent cells (P2-P5) originating from mouse patellae that

proliferated in culture (Figure 1A, B). From P2 to P3, the expression of the macrophage marker

(1,5) cluster of differentiation 68 (*Cd-68*, relative to the macrophage cell line RAW 267.4) was

significantly reduced, whereas the expression of FLS markers *Cdh-11* and *Cd-248* (1,5) remained

consistent from P2 to P5 (Supplementary Figure 1 in Additional file 1); the endothelial marker *Cd-*

31 was not detected (data not shown). These results suggest that from P3, cells cultured from

mouse patellae were predominantly FLS and hence subsequent studies were conducted on FLS

from P3-P6.

Concurrently, to establish the pro-inflammatory phenotype of FLS, we used RT-qPCR to

determine the expression of the inflammatory genes *Il-6*, *Il-1r1* and *Cox-2*, as well as the

constitutively expressed gene *Cox-1*. When control FLS (P5) were stimulated with 10 ng/ml TNF-

α for 48-hours, there was an upregulation of *Il-6* (fold changes: Control, 1.5 ± 0.8 vs. TNF-FLS,

135.1 ± 31.1, n = 3, p = 0.006, unpaired t-test) and *Il-1r1* (fold changes: Control, 1.1 ± 0.2 vs.

TNF-FLS, 3.4 ± 1.0, n = 3, p = 0.04, unpaired t-test) expression levels, but not that of *Cox-2* (fold

changes: Control, 1.0 ± 0.2 vs. TNF-FLS, 0.8 ± 0.2, n = 3, p = 0.2, unpaired t-test) or *Cox-1* (fold

changes: Control, 1.0 ± 0.2 vs. TNF-FLS, 0.8 ± 0.2, n = 3, p = 0.18, unpaired t-test) (Figure 1C).

However, when FLS derived from the inflamed knee (Ipsi, knee width, pre-CFA, 3.1 ± 0.09 mm

vs. post-CFA, 4.1 ± 0.08 mm, n = 6, p = 0.0001, paired t-test, Supplementary Figure 1 in Additional

File 1) of CFA injected mice was compared to those of the matched contralateral knee (Contra),

we did not find any changes in expression levels of the genes between Contra and Ipsi FLS across

P2-P5 (Figure 1C, Supplementary Figure 1 in Additional File 1).

We next tested the media (n = 3, each) isolated from control, Contra, Ipsi and TNF-FLS against a

mouse inflammation antibody array (Figure 1E) to determine the levels of different secreted pro-

inflammatory mediators. This demonstrated that when compared to control FLS media, TNF-FLS

media showed presence of Regulated on activation, normal T-cell expressed and secreted

(RANTES) and granulocyte-macrophage colony stimulating factor (GM-CSF). Additionally,

TNF-FLS contained higher levels of IL-6 (p < 0.0001), keratinocyte chemoattractant (KC, mouse

homolog of IL-8, p < 0.0001), lipopolysaccharide induced chemokine (LIX, p < 0.0001), stromal

cell derived factor 1 (SDF-1, p < 0.0001), fractalkine (p < 0.0001), monocyte chemoattractant

protein 1 (MCP-1, p < 0.0001) and macrophage inhibitory protein 1γ (MIP-1γ, p = 0.002) and

lower levels of macrophage colony stimulating factor (MCSF, p = 0.003) compared to control FLS

media (multiple t-tests with Holm-Sidak corrections, Figure 1Dii). TNF-α was not detected in

TNF-FLS media as previously shown in TNF-α stimulated airway epithelial cells (40). Mirroring

the absence of pro-inflammatory gene upregulation in Ipsi vs. Contra FLS, the spot intensity values

of inflammatory cytokines between Ipsi and Contra FLS were similar (multiple t-tests with Holm-

Sidak corrections, graph not shown), suggesting similar levels of secreted cytokines. However,

when spot intensities were normalized to control FLS, TNF-FLS media showed increased levels

of secreted IL-6 (p < 0.0001, ANOVA followed by Tukey's multiple comparison test) and KC (p

= 0.001, ANOVA followed by Tukey's multiple comparison test) compared to Ipsi and Contra

FLS (Figure 1Diii).

Interestingly, MIP-1γ was elevated in both Ipsi and Contra FLS when compared to control and

TNF-FLS (Figure 1Diii), perhaps indicating its role in acute, systemic inflammatory pathways.

MIP-1γ has been described as having a role in hyperalgesia induced by diabetic neuropathy (41)

and OA-related pain (42), and we find that MIP-1γ directly activates mouse DRG neurons by

producing a dose-dependent Ca^{2+} influx (Supplementary Figure 1 in Additional File 1), which thus

might be a potential mechanism for the *in vivo* effects described above. However, because MIP-

1γ does not have a human homolog (NCBI gene database entry #20308), and thus has limited

clinical potential, we did not explore this further.

Taken together, our data suggest that TNF-α stimulation of FLS upregulates expression of

canonical inflammatory markers, whereas the *in vivo* pathology observed 24-hours after CFA-

induced knee inflammation does not correlate with a sustained pro-inflammatory phenotype of

FLS isolated from patellae of CFA mice. We also identify other soluble mediators secreted by

TNF-FLS that might be involved in FLS-DRG neuron communication. Therefore, given the pro-

inflammatory phenotype of TNF-FLS (and the lack of such phenotype in the CFA model), we

focused solely on the TNF-FLS for the rest of the study.

2.　TNF-FLS have reduced sensitivity to mild acidosis, but enhanced sensitivity to more acidic

pH.

During arthritis, FLS are located in an environment with abundant algogens that have been shown

to signal through the modulation of TRP channels (2). Therefore we tested FLS sensitivity to a

range of acidic pH stimuli, as well as the prototypic TRP channel agonists, capsaicin (TRPV1),

cinnamaldehyde (TRPA1) and menthol (TRPM8) using Ca^{2+} imaging. Confirming results from

other studies on acid sensing in FLS (22), we found that both control and TNF-FLS respond to a

range of pH solutions (pH 4.0 – 7.0) with an increase in intracellular Ca^{2+} (Figure 2A, B). An

increased percentage of control FLS (24.1 %) respond to pH 7.0 compared to TNF-FLS (11.0 %,

$p = 0.0004$, chi-sq test), but in the more acidic range an increased percentage of TNF-FLS

responded to acid than did control FLS, i.e. pH 6.0 (control, 7.2 % vs. TNF-FLS, 26.3 %, p <

0.0001, chi-sq test) and pH 5.0 (control, 14.4 % vs. TNF-FLS, 30.5 %, $p < 0.0001$, chi-sq test).

At the very acidic pH of 4.0, ~ 40% ($p = 0.4$, chi-sq test) of FLS responded in both groups possibly

due to ceiling effect. Although the proportion of responding neurons varied, the magnitude of peak

normalized response was similar across the pH range in control (pH 7.0: 0.4 ± 0.04 F/Fmax (n =

50), pH 6.0: 0.4 ± 0.08 F/Fmax (n = 27), pH 5.0: 0.3 ± 0.03 F/Fmax (n = 46), pH 4.0: 0.3 ± 0.02

F/Fmax (n = 129)) and TNF-FLS (pH 7.0: 0.3 ± 0.06 F/Fmax (n = 21), pH 6.0: 0.4 ± 0.03 F/Fmax

(n = 50), pH 5.0: 0.3 ± 0.02 F/Fmax (n = 58), pH 4.0: 0.3 ± 0.02 F/Fmax (n = 72), Figure 2Bii).

To identify the proton sensors expressed by FLS that likely mediate the responses measured, we

conducted RT-PCR of control and TNF-FLS to determine expression of known proton-sensing G

protein-coupled receptors (GPCRs), ASICs and TRPV1. Our data show that FLS express the

proton sensing GPCRs – *Gpr4*, *Gpr65*, *Gpr68* and *Gpr132*, as well as *Asic1*, Asic*3* and *Trpv1*

(Figure 2Ci). When relative expression was assessed compared to the housekeeping gene *18S*,

Gpr132 was found to be increased ($p = 0.005$, Figure 2Cii) in TNF-FLS implicating it as a potential

contributor to the enhanced proportion of acid responders seen in TNF-FLS (Figure 2Bi). The pro-

inflammatory phenotype of TNF-FLS was confirmed by the increased *Il-6* band intensity (p <

0.0001), as observed previously using qPCR (multiple t-tests with Holm-Sidak corrections, Figure

2Cii).

We also tested FLS sensitivity to capsaicin (control, 3/201, TNF-FLS, 13/252), cinnamaldehyde

(control, 1/201, TNF-FLS, 11/252) and menthol (control, 0/201, TNF-FLS, 1/194) to find that only

a few cells responded to each compound (data not shown). This led us to conclude that there are

very few mouse primary FLS with functional TRPV1 (also corroborated by the observation of low

Trpv1 gene expression, Figure 2C), TRPA1 or TRPM8 ion channels and hence we did not explore

this further.

3.   TNF-FLS increase knee-innervating DRG neuron excitability in co-culture.

Changes is primary sensory neuron excitability underlie peripheral sensitization which drives

arthritis-related pain (43). Non-neuronal cells like FLS, which are in close proximity with distal

terminals of knee neurons, can play instrumental roles in modulating sensory neuron excitability

by direct cell contact and/or secretion of pro-inflammatory cytokines. To determine neuron/FLS

communication in health and inflammation we compared the DRG neuronal excitability of four

groups: 1) knee neuron mono-culture, 2) knee neurons co-cultured with control FLS (Figure 3A,

B), 3) knee neurons with FLS that have been exposed to TNF media for 24-hour followed by DRG

culture media for 24-hours and 4) knee neurons with conditioned media from FLS that have been

exposed to TNF media for 48-hours to understand the role of soluble mediators in peripheral

sensitization. In current-clamp mode, we observed many neurons spontaneously firing AP in

groups 3 and 4. In order to statistically compare the proportions, we converted our four group data

into binary categories: neuron mono-culture and neuron/control FLS co-culture were assigned to

the class "healthy", and neuron/TNF-FLS and neuron/TNF-FLS media were assigned to the class

"inflamed". 19.5% of inflamed neurons (neuron/TNF-FLS, 5/21; neuron/TNF-FLS media, 3/19)

fired spontaneous AP compared to 2.7% (neuron mono-culture, 0/19; neuron/control FLS, 1/18)

of healthy neurons (p = 0.02, chi-sq test) suggesting that there is a general increase in excitability

of knee neurons when exposed to an FLS-mediated inflammatory environment (Figure 3C, D).

Upon measuring the AP properties (Figure 3D inset) we found that the resting membrane potential

(RMP) was more depolarized in the neuron/TNF-FLS media group compared to both neuronal

mono-culture (p = 0.0004) and neuron/control FLS co-culture (p = 0.012), which highlights that

secreted pro-inflammatory factors from TNF-FLS likely act upon knee neurons to increase their

excitability (ANOVA followed by Tukey's post hoc comparison, Figure 3Ei). The other measured

properties were unchanged among the four groups (ANOVA followed by Tukey's post hoc

comparison, Table 1, Figure 3Eii-iv).

4. TNF-FLS increase TRPV1 function and decrease TRPA1/TRPM8 function of DRG

neurons in co-culture.

DRG neurons in co-culture with inflamed FLS derived from AIA rats reportedly have increased

expression of TRPV1 (6). Here we sought to investigate whether FLS can modulate knee neuron

responses to TRP agonists using whole-cell patch-clamp recordings. Knee neurons in mono-

culture and when co-cultured with control FLS displayed very similar responses, i.e. a mean

capsaicin peak current density response of 3.5 ± 1.0 pA/pF (n = 7) and 3.2 ± 0.9 pA/pF (n = 7)

respectively. By contrast, knee neurons in neuron/TNF-FLS co-culture showed a trend for larger

magnitude of capsaicin responses (27.7 ± 9.3 pA/pF, n = 8), and those cultured with TNF-FLS

media had a capsaicin response of 65.8 ± 25.7 pA/pF (n = 7) which was significantly larger in

magnitude than neuronal mono-culture (p = 0.02) and neuron/control FLS co-culture (p = 0.02)

(ANOVA followed by Tukey's post-hoc comparison, Figure 4Ai,ii). No difference was observed

between the percentage of capsaicin responders in healthy (40.5%) vs. inflamed groups (37.5%, p

= 0.8, chi-sq test, Figure 4Aiii).

With regards to menthol (mono-culture, 2.4 ± 1.2 pA/pF, n = 7; neuron/healthy FLS, 4.8 ± 2.1

pA/pF, n = 6; neuron/TNF-FLS, 1.4 ± 1.2 pA/pF, n = 2; neuron/TNF-FLS conditioned media, 4.0

± 1.9 pA/pF, n = 4; Figure 4Bi) and cinnamaldehyde (mono-culture, 1.9 ± 0.7 pA/pF, n = 10;

neuron/healthy FLS, 2.8 ± 1.1 pA/pF, n = 4; neuron/TNF-FLS, 0.6 ± 0.3 pA/pF, n = 3,

neuron/TNF-FLS conditioned media, 2.3 ± 1.5 pA/pF, n = 3; Figure 4Ci) responses, all four groups

showed similar mean peak current density values (Figure 4Bii, Cii). However, in response to both

the TRPA1 and TRPM8 agonists, the percentage of responding neurons was significantly less in

the inflamed group compared to healthy (menthol: healthy vs. inflamed, 37.8% vs. 15%, p = 0.02,

chi-sq test; cinnamaldehyde: healthy vs. inflamed, 43.2% vs. 15%, p = 0.006, chi-sq test, Figure

4Biii, Ciii). Taken together our data suggests that "inflamed" FLS can alter TRP agonist response

of knee neurons in co-culture.

Discussion

In this study we demonstrate that FLS can be obtained from mouse patella via cell-outgrowth and maintained in culture. When stimulated with TNF-α, these FLS show an increase in: pro-inflammatory gene expression, secretion of pro-inflammatory cytokines and acid sensitivity in the pH 6.0-5.0 range consistent with a pro-nociceptive activated phenotype. Finally, we establish a co-culture system, which shows that FLS and knee neurons can interact, specifically, factors secreted by TNF-FLS increase knee neuron excitability and the magnitude of the response to capsaicin, whereas the proportions of cinnamaldehyde and menthol responding neurons are diminished. To the best of our knowledge, this is the first report to demonstrate FLS-mediated changes in neuronal excitability in co-culture and hence directly demonstrate how FLS can regulate articular neurons and in turn arthritis-related pain.

FLS from mouse are generally cultured by enzymatically digesting and combining excised joints of fore- and hind-limbs (5,21,22,44), which assumes similarity of FLS derived from all joints. However, a genome-wide study on DNA-methylation has shown that important differences exist between knee and hip FLS, including of genes involved in IL-6 signaling (45). By using a cell-outgrowth method to culture mouse FLS, as previously described in humans (46) and rats (6), we have avoided the biological ambiguity introduced by joint-to-joint variability. Using these FLS, we investigated the expression of inflammatory genes *Il-6*, *Il-1r1* and *Cox-2*, all of which have been linked to the inflammatory phenotype of FLS. In brief: stimulating human-derived FLS with

IL-1β increases *Cox-2* and *Il-6* expression (46); increased *Il-6* expression is seen in FLS derived

from K/BxN mice and following 24-hour stimulation of healthy mouse FLS with TNF-α (5).

Lastly, supernatants from FLS derived 3-days into a rat AIA model show increased IL-6 and PGE$_2$

(6). Here we observed that neither the expression level of *Il-6, Il-1r1, Cox-2*, nor the level of

secreted cytokines were upregulated in FLS derived from CFA-injected knee. This is possibly

because the model used here is too brief to influence FLS gene expression. Indeed, evidence to

support this is that FLS derived from rats with longer (3-28 days) AIA-induced knee inflammation

did have higher PGE$_2$ and IL-6 concentrations in culture supernatants compared to control FLS

(6). Alternatively, the affected area of the synovium might be too small for sufficient proliferation

of "inflamed" FLS and thus they are lost over time in culture. This result suggests a limited role

of FLS in the acute CFA model of knee inflammation with regard to modulation of neuronal

excitability.

We reasoned that FLS need more direct and prolonged exposure to inflammatory mediators to

show a robust inflammatory phenotype. TNF-α is one such cytokine that is locally upregulated

within 3-hours of intra-plantar CFA injection in mice (12). It is also present in high concentrations

in the synovial fluid (47) and tissue of OA and RA patients (48,49); along with anti-TNF-α agents

being the leading treatment of RA (50). Indeed, we show that TNF-α stimulated FLS display

increased expression of *Il-6* and *Il-1r1* mRNA, with a concomitant increase in secretion of many

pro-inflammatory cytokines, including IL-6 as has been reported previously in human FLS (51).

This observation is consistent with the "passive responder and imprinted aggressor model" of

inflammatory FLS behavior (2), i.e. these cells can respond to pro-inflammatory environment and

then themselves become effectors to drive disease pathology. The functional repercussions of this

model are not well-explored and here we investigate two such possibilities: change in FLS

functionality after induction of inflammation and "inflamed" FLS-induced functional changes in

nerves supplying the knee joint to drive pain.

Comparing the control and TNF-FLS response to acidic stimuli, we found a decreased percentage

of TNF-FLS responding to pH 7.0, but in the pH 6.0-5.0 range the percentage of TNF-FLS

responding to acidic stimulation increased. Multiple proton-sensors with sensitivity to varied pH

range might underlie the apparent incongruity of TNF-FLS acid sensitivity in mild and strongly

acidic environment (52). For example, increased expression of proton sensors programmed to

detect highly acidic environments can come at the cost of sensors active at mildly acidic range.

This concept has been alluded to previously with mouse primary FLS showing increased *Asic3*,

but decreased *Asic1*, expression after IL-1β stimulation (22). The present study shows that ASIC1

and 3 may underlie acid sensitivity in FLS, as has been shown previously (21,22) and here we also

show that multiple proton sensing GPCRs are expressed in FLS. Additionally, an increased level

of *Gpr132* in TNF-FLS lends support to the hypothesized role of GPR132 (15) in Ca^{2+}

mobilization of FLS in response to acidification. Furthermore, TRPV1 is activated at pH below 6,

which can contribute to the enhanced acid response of TNF-FLS in that range. Although we report

very few FLS respond to capsaicin at physiological pH of 7.4 (as has been shown before (18)),

more cells responded in the TNF condition compared to control.

In order to understand the effector role of FLS in driving nociception through peripheral

sensitization, we set up a co-culture system combining FLS and knee neurons. Studying co-culture

of rat FLS and DRG neurons, von Banchet *et al* showed using immunohistochemistry that

bradykinin 2 receptor labelling (but not that of neurokinin 1 or TRPV1) was increased when DRG

neurons were cultured with healthy rat FLS (6), i.e. co-culture of DRG neurons with FLS can alter

expression of genes associated with nociception. Therefore, we first verified that knee neurons in

co-culture with control FLS do not show dysregulation of excitability or TRP agonist response.

Then we asked whether TNF-FLS modulates knee neuron function and found using whole-cell

patch clamp that 23% and 16% of knee neurons in neuron/TNF-FLS co-culture and neuron/TNF-

FLS media (inflamed conditions) respectively evoked spontaneous AP compared to 6% in

neuron/control FLS and 0% in neuron mono-culture (healthy conditions). This suggests that TNF-

FLS increases excitability of knee neurons. von Banchet *et al* (6) also showed that compared to

mono-culture, FLS derived from acute and chronic AIA rats induced an increase in TRPV1 protein

expression in DRG neurons. However, we did not see increase in proportion of neurons responding

to capsaicin between healthy and inflamed conditions, which might reflect a species difference

and/or difference in knee-specific neuronal population (unlike in this study, von Banchet *et al* did

not discriminate between knee-innervating neurons and non-knee innervating neurons); however,

we did observe that capsaicin responsive neurons produced larger magnitude responses when

incubated with TNF-FLS medium, see below. We also observed that cinnamaldehyde and

menthol-evoked responses were decreased in the inflamed condition, suggesting functional

downregulation of TRPA1 and TRPM8. Functional downregulation of TRPA1 might be explained

through desensitization via increased TRPV1 function (see below) (53) or via an increase in

intracellular Ca^{2+} (due to increased excitability in the inflamed condition) (54). The latter reason

can also be applied to explain decrease of menthol-evoked responses (55).

We also observed a tendency of a more depolarized RMP and an enhanced magnitude of capsaicin-

evoked peak current density of knee neurons in inflamed conditions, albeit both only reached

statistical significance in the TNF-FLS media incubated group when compared to mono-culture

and neuron/control FLS. These results suggest that soluble mediators released by FLS are key

players in modulating knee neuron excitability. We posit that because our experimental design

involved a media change 24-hours after TNF incubation of FLS (so as not to directly stimulate

neurons with residual TNF-α (56)) in the neuron/TNF-FLS co-culture condition, the accumulated

soluble mediator concentrations were lower compared to when knee neurons were incubated in

48-hours TNF-FLS conditioned media (no remaining TNF-α was measured in the media at this

point). Indeed we have recently established the role of soluble mediators present in OA synovial

fluid in increasing neuronal excitability and TRPV1 function (33). These soluble mediators mainly

consist of cytokines/chemokines which form complex signaling pathways with neurons (reviewed

in (57,58)). In this study, we have identified IL-6 (59), KC (60), RANTES (61), GM-CSF (62),

LIX (63), SDF-1 (61), MCP-1 (64), MIP-1γ (present study) to be upregulated in TNF-FLS media

which can also directly signal to neurons; and in the cases of IL-6, MCP-1 and GM-CSF increase

TRPV1 function (59,62,64). Although fractalkine does not directly stimulate neurons (57), a recent

study showed a correlation between levels of this cytokine in protein extracted from human OA

synovia and brain derived neurotrophic factor - tropomyosin receptor kinase B mediated signaling

in joint nerves to induce mechanical hyperalgesia (65).

Therefore, this study demonstrates an increase in knee neuron excitability and TRPV1 function

upon interaction with non-neuronal TNF-FLS.

630   Conclusion

Knee pain in arthritis is a major cause of disability and peripheral sensitization mediated by

enhanced excitability of nociceptive neurons play a key role in pain (43,66). Those neurons

supplying the knee joint are exposed to a pro-inflammatory environment due to cell-contact and

soluble mediator release from immune cells. One such key effector in the joint is FLS which can

respond to inflammatory mediators and also themselves secrete inflammatory soluble factors (2).

In this study we show that TNF-FLS increase mRNA expression of inflammation-linked genes *Il-*

6 and *Il-1r1* along with enhanced secretion of cytokines compared to control. Additionally, TNF-

FLS can increase excitability of knee neurons and alter TRP channel function in these neurons in

a co-culture system. Therefore, we establish a direct inflammation-pain axis through FLS and DRG

neurons. This system can be adapted to investigate peripheral sensitization using FLS activated by

other pro-inflammatory cytokine or arthritic synovial fluid.

Declarations

Ethics approval: Mice used in this study were regulated under the Animals (Scientific Procedure)

Act 1986, Amendment Regulations 2012. All protocols were reviewed by the University of

Cambridge Animal Welfare and Ethical Review Body.

Consent for publication: Not applicable.

Availability of data and materials: The datasets supporting the conclusions of this article will be

available in University of Cambridge Apollo Repository (https://doi.org/10.17863/CAM.44367).

Competing interests: The authors declare no competing interests.

Funding: This study was supported by Versus Arthritis Project Grants (RG 20930 and RG 21973)

and Rosetrees grant (M818) to E. St. J. S. and G.C., and Crohn's and Colitis UK grant to D.C.B.

(PC2019/1-Bulmer). S.C. and C.N.B were supported by Gates Cambridge Trust scholarships.

L.A.P. was supported by the University of Cambridge BBSRC Doctoral Training Programme

(BB/M011194/1).

Authors' contributions:

S.C. designed experiments, conducted experiments, analyzed data and drafted the manuscript.

L.A.P. and C.N.B. conducted and analyzed experiments and revised manuscript. G.C. designed

experiments and revised the manuscript. D.C.B revised the manuscript. E. St. J. S. designed

experiments and drafted the manuscript.

Acknowledgements:

The authors thank Dr Dora Lopresto (Department of Pharmacology, University of Cambridge) for

technical help with cell staining.

Figure Legends:

Figure 1: Induction of inflammation in FLS derived from mouse knee. A) Representative picture

showing exposed inside of the patella (black triangle) and the surrounding ligament and fat pad

(white triangle) after midline resection and distal pull of the quadriceps muscles. Scale bar = 2

mm. B) Representative picture of FLS in culture. Scale bar = 50 μm. C) Bars represent fold change

of the genes *Il-6*, *Il-1r1*, *Cox-2* and *Cox-1* from either Contra (vs. Ipsi) or control (vs. TNF). Black

hatched bars = control FLS, red bars = TNF-FLS, grey bar = Contra, white bar = Ipsi. All FLS at

P5. D) Images of mouse inflammatory array membranes probed against FLS conditioned medium

(i) which were quantified by densitometry and represented as bar graphs showing fold change of

various cytokines between control and TNF-FLS (ii, multiple t-test with Holm-Sidak correction)

and among Contra, Ipsi and TNF from control FLS media (iii, ANOVA with Tukey's post-hoc

test) and spot intensity differences. Dotted rectangles highlight IL-6 spots (Di) and corresponding

quantifications (Dii, iii). Only cytokines that were present in all of the compared groups are shown

in graphs. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ and **** $p < 0.0001$. Error bars = SEM.

Figure 2: Acid sensitivity in FLS. A) Representative Ca^{2+} imaging trace from an FLS responding

to pH 5 and ionomycin. Bi) Bars representing proportions and magnitudes (ii) of control (black

hatched) and TNF-FLS (red) responding to pH 7, 6, 5 and 4. Comparison between control vs. TNF-

FLS made using chi-sq test. Data from three biological replicates in each category. Ci) Gel image

of control and TNF-FLS expression of proton sensors along with the densitometry analysis of the

relative band intensity compared to the 18S band (ii, multiple t-test with Holm-Sidak correction).

** $p < 0.01$, *** $p < 0.001$ and **** $p < 0.0001$. Error bars = SEM.

Figure 3: TNF-FLS mediated increase in excitability of knee neurons. A) Representative live-cell

imaging picture showing FLS (magenta) and neuron (cyan) in co-culture. Scale bar = 50 μm. B)

Representative FB labelled knee neuron (white arrow) being recorded using a patch pipette

(triangular shadow), surrounded by FLS (yellow arrow). Scale bar = 50 μm. C) Pie-chart showing

proportion of knee neurons that fired AP without current stimulation in healthy (neuron mono-

culture + neuron/control FLS, n = 37, black) and inflamed (neuron/TNF-FLS + neuron/TNF-FLS

media, n = 41, red) condition. D) Representative knee neuron incubated with TNF-FLS media

firing spontaneous AP along with schematic diagram of AP properties measured (inset). E) Bar

graphs showing measured RMP (i), threshold (ii), AHP peak (iii) and HPD (iv) from knee neurons

in mono-culture (n = 19, white bar/black open circle), in co-culture with control FLS (n = 18, grey

bar/black dotted circle), in co-culture with TNF-FLS (n = 20, light red bar/red dotted circle) and

incubated in TNF-FLS media (n = 21, white bar/red open circle). * p < 0.05 and *** p < 0.001,

ANOVA followed by Tukey's post hoc test. Data from 4-5 female mice in each group. Error bars

= SEM.

Figure 4: TNF-FLS mediated modulation of TRP agonist response in knee neurons. Representative

traces showing capsaicin (TRPV1 agonist, Ai), cinnamaldehyde (TRPA1 agonist, Bi) and

menthol- (TRPM8 agonist, Ci) evoked responses in knee neurons. Black traces obtained from knee

neuron in mono-culture, red trace obtained from knee neuron incubated in TNF-FLS media. White

boxes represent perfusion of extracellular solution. Bar graphs showing peak current densities of

capsaicin- (Aii), cinnamaldehyde- (Bii) or menthol-evoked (Cii) currents from knee neurons in

mono-culture (white bar/black open circle), in co-culture with control FLS (grey bar/black dotted

circle), in co-culture with TNF-FLS (light red bar/red dotted circle) and incubated in TNF-FLS

media (white bar/red open circle) Comparison between groups made using ANOVA with Tukey's

post hoc test. Bar graphs showing percent of knee neurons that responded to capsaicin (Aiii),

cinnamaldehyde (Biii) and menthol (Ciii) in healthy (neuron mono-culture + neuron/control FLS)

and inflamed (neuron/TNF-FLS + neuron/TNF-FLS media) condition. Comparison made using

chi-sq test. * $p < 0.05$. Data from 4-5 female mice in each group. Error bars = SEM.

Additional materials:

• File name: Additional file 1

• File format including the correct file extension: .docx

• Title of data: Table 1, 2 and Figure 1

• Description of data: Table 1 contains list of genes assessed in this study using qPCR and

RT-PCR, Table 2 contains location of cytokines detected by the mouse inflammatory

antibody array membrane. Figure 1 shows characterization of mouse primary FLS and

neuronal activation of FLS secreted MIP-1γ.

References:

1. Bartok B, Firestein GS. Fibroblast-like synoviocytes: key effector cells in rheumatoid arthritis. Immunological reviews. 2010 Jan;233(1):233–255.

2. Bottini N, Firestein GS. Duality of fibroblast-like synoviocytes in RA: passive responders and imprinted aggressors. Nat Rev Rheumatol [Internet]. 2013;9. Available from: http://dx.doi.org/10.1038/nrrheum.2012.190

3. Croft AP, Campos J, Jansen K, Turner JD, Marshall J, Attar M, et al. Distinct fibroblast subsets drive inflammation and damage in arthritis. Nature. 2019 Jun 1;570(7760):246–51.

4. Lee DM, Kiener HP, Agarwal SK, Noss EH, Watts GFM, Chisaka O, et al. Cadherin-11 in Synovial Lining Formation and Pathology in Arthritis. Science. 2007 Feb 16;315(5814):1006.

5. Hardy RS, Hülso C, Liu Y, Gasparini SJ, Fong-Yee C, Tu J, et al. Characterisation of fibroblast-like synoviocytes from a murine model of joint inflammation. Arthritis Research & Therapy. 2013;15(1):R24–R24.

6. von Banchet GS, Richter J, Hückel M, Rose C, Bräuer R, Schaible H-G. Fibroblast-like synovial cells from normal and inflamed knee joints differently affect the expression of pain-related receptors in sensory neurones: a co-culture study. Arthritis Research & Therapy. 2007;9(1):R6–R6.

7. Llop-Guevara A, Porras M, Cendón C, Di Ceglie I, Siracusa F, Madarena F, et al. Simultaneous inhibition of JAK and SYK kinases ameliorates chronic and destructive arthritis in mice. Arthritis Research & Therapy. 2015 Dec 10;17(1):356.

8. Guo W, Yu D, Wang X, Luo C, Chen Y, Lei W, et al. Anti-inflammatory effects of interleukin-23 receptor cytokine-binding homology region rebalance T cell distribution in rodent collagen-induced arthritis. Oncotarget. 2016 May 31;7(22):31800–13.

9. Chen C-X, Chen J-Y, Kou J-Q, Xu Y-L, Wang S-Z, Zhu Q, et al. Suppression of Inflammation and Arthritis by Orally Administrated Cardiotoxin from Naja naja atra. Evid Based Complement Alternat Med. 2015;2015:387094.

10. Hong R, Sur B, Yeom M, Lee B, Kim KS, Rodriguez JP, et al. Anti-inflammatory and anti-arthritic effects of the ethanolic extract of Aralia continentalis Kitag. in IL-1β-stimulated human fibroblast-like synoviocytes and rodent models of polyarthritis and nociception. Phytomedicine. 2018 Jan 1;38:45–56.

11. Chakrabarti S, Pattison LA, Singhal K, Hockley JRF, Callejo G, Smith ESJ. Acute
inflammation sensitizes knee-innervating sensory neurons and decreases mouse digging
behavior in a TRPV1-dependent manner. Neuropharmacology. 2018 Dec 1;143:49–62.

12. Woolf CJ, Allchorne A, Safieh-Garabedian B, Poole S. Cytokines, nerve growth factor and
inflammatory hyperalgesia: the contribution of tumour necrosis factor α. British Journal of
Pharmacology. 1997 May 1;121(3):417–24.

13. Andersson SE, Lexmuller K, Johansson A, Ekstrom GM. Tissue and intracellular pH in
normal periarticular soft tissue and during different phases of antigen induced arthritis in
the rat. J Rheumatol. 1999 Sep;26(9):2018–24.

14. Wright AJ, Husson ZMA, Hu D-E, Callejo G, Brindle KM, Smith ESJ. Increased
hyperpolarized [1-13C] lactate production in a model of joint inflammation is not
accompanied by tissue acidosis as assessed using hyperpolarized 13C-labelled bicarbonate.
NMR in Biomedicine. 2018;e3892-n/a.

15. Christensen BN, Kochukov M, McNearney TA, Taglialatela G, Westlund KN. Proton-
sensing G protein-coupled receptor mobilizes calcium in human synovial cells. Am J
Physiol Cell Physiol [Internet]. 2005;289. Available from:
http://dx.doi.org/10.1152/ajpcell.00039.2005

16. Kolker SJ, Walder RY, Usachev Y, Hillman J, Boyle DL, Firestein GS, et al. Acid-sensing
ion channel 3 expressed in type B synoviocytes and chondrocytes modulates hyaluronan
expression and release. ANNALS OF THE RHEUMATIC DISEASES. 2010
May;69(5):903–9.

17. Galindo T, Reyna J, Weyer A. Evidence for Transient Receptor Potential (TRP) Channel
Contribution to Arthritis Pain and Pathogenesis. Pharmaceuticals (Basel, Switzerland).
2018 Oct 15;11(4):105.

18. Kochukov MY, McNearney TA, Fu Y, Westlund KN. Thermosensitive TRP ion channels
mediate cytosolic calcium response in human synoviocytes. Am J Physiol Cell Physiol
[Internet]. 2006;291. Available from: http://dx.doi.org/10.1152/ajpcell.00553.2005

19. Itoh Y, Hatano N, Hayashi H, Onozaki K, Miyazawa K, Muraki K. An environmental
sensor, TRPV4 is a novel regulator of intracellular Ca2+ in human synoviocytes. American
Journal of Physiology-Cell Physiology. 2009 Sep 16;297(5):C1082–90.

20. Kochukov MY, McNearney TA, Yin H, Zhang L, Ma F, Ponomareva L, et al. Tumor
Necrosis Factor-Alpha (TNF-α) Enhances Functional Thermal and Chemical Responses of
TRP Cation Channels in Human Synoviocytes. Mol Pain. 2009 Jan 1;5:1744-8069-5–49.

21. Sluka KA, Rasmussen LA, Edgar MM, O'Donnell JM, Walder RY, Kolker SJ, et al. Acid-
sensing ion channel 3 deficiency increases inflammation but decreases pain behavior in
murine arthritis. 2013;(1529-0131 (Electronic)).

22.　Gong W, Kolker SJ, Usachev Y, Walder RY, Boyle DL, Firestein GS, et al. Acid-sensing
ion channel 3 decreases phosphorylation of extracellular signal-regulated kinases and
induces synoviocyte cell death by increasing intracellular calcium. Arthritis Research &
Therapy. 2014;16(3):R121.

23.　Lebre MC, Vieira PL, Tang MW, Aarrass S, Helder B, Newsom-Davis T, et al. Synovial
IL-21/TNF-producing CD4+ T cells induce joint destruction in rheumatoid arthritis by
inducing matrix metalloproteinase production by fibroblast-like synoviocytes. Journal of
Leukocyte Biology. 2017 Mar 1;101(3):775–83.

24.　Bombara MR, Webb DL, Conrad P, Marior CW, Sarr T, Ranges GE, et al. Cell contact
between T cells and synovial fibroblasts causes induction of adhesion molecules and
cytokines. Journal of Leukocyte Biology. 1993 Nov 1;54(5):399–406.

25.　Yamamura Y, Gupta R, Morita Y, He X, Pai R, Endres J, et al. Effector Function of Resting
T Cells: Activation of Synovial Fibroblasts. J Immunol. 2001 Feb 15;166(4):2270.

26.　Burger D, Rezzonico R, Li J-M, Modoux C, Pierce RA, Welgus HG, et al. Imbalance
between interstitial collagenase and tissue inhibitor of metalloproteinases 1 in synoviocytes
and fibroblasts upon direct contact with stimulated T lymphocytes: Involvement of
membrane-associated cytokines. Arthritis & Rheumatism. 1998 Oct 1;41(10):1748–59.

27.　Bondeson J, Wainwright SD, Lauder S, Amos N, Hughes CE. The role of synovial
macrophages and macrophage-produced cytokines in driving aggrecanases, matrix
metalloproteinases, and other destructive and inflammatory responses in osteoarthritis.
Arthritis Research & Therapy. 2006 Dec 19;8(6):R187.

28.　Blue ML, Conrad P, Webb DL, Sarr T, Macaro M. Interacting monocytes and synoviocytes
induce adhesion molecules by a cytokine-regulated process. Lymphokine Cytokine Res.
1993 Aug;12(4):213–8.

29.　Chomarat P, Rissoan MC, Pin JJ, Banchereau J, Miossec P. Contribution of IL-1, CD14,
and CD13 in the increased IL-6 production induced by in vitro monocyte-synoviocyte
interactions. J Immunol. 1995 Oct 1;155(7):3645.

30.　Scott BB, Weisbrot LM, Greenwood JD, Bogoch ER, Paige CJ, Keystone EC. Rheumatoid
arthritis synovial fibroblast and U937 macrophage/monocyte cell line interaction in
cartilage degradation. Arthritis & Rheumatism. 1997 Mar 1;40(3):490–8.

31.　Benito-Miguel M, García-Carmona Y, Balsa A, Bautista-Caro M-B, Arroyo-Villa I, Cobo-
Ibáñez T, et al. IL-15 Expression on RA Synovial Fibroblasts Promotes B Cell Survival.
PLOS ONE. 2012 Jul 9;7(7):e40620.

32.　Chan A, Filer A, Parsonage G, Kollnberger S, Gundle R, Buckley CD, et al. Mediation of
the proinflammatory cytokine response in rheumatoid arthritis and spondylarthritis by
interactions between fibroblast-like synoviocytes and natural killer cells. Arthritis &
Rheumatism. 2008 Mar 1;58(3):707–17.

33.   Chakrabarti S, Jadon DR, Bulmer DC, Smith ESJ. Human osteoarthritic synovial fluid
increases excitability of mouse dorsal root ganglion sensory neurons: an in-vitro
translational model to study arthritic pain. Rheumatology [Internet]. 2019 Aug 13 [cited
2019 Aug 14]; Available from: https://doi.org/10.1093/rheumatology/kez331

34.   Hannan MT. Epidemiologic perspectives on women and arthritis: An overview. Arthritis &
Rheumatism. 1996 Dec 1;9(6):424–34.

35.   Futami I, Ishijima M, Kaneko H, Tsuji K, Ichikawa-Tomikawa N, Sadatsuki R, et al.
Isolation and Characterization of Multipotential Mesenchymal Cells from the Mouse
Synovium. PLoS ONE. 2012 Sep 18;7(9):e45517.

36.   McCall MN, McMurray HR, Land H, Almudevar A. On non-detects in qPCR data.
Bioinformatics. 2014 Aug 15;30(16):2310–6.

37.   Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time
Quantitative PCR and the 2−ΔΔCT Method. Methods. 2001 Dec 1;25(4):402–8.

38.   Stefanini M, Martino C de, Zamboni L. Fixation of Ejaculated Spermatozoa for Electron
Microscopy. Nature. 1967 Oct 1;216(5111):173–4.

39.   Dittert I, Benedikt J, Vyklický L, Zimmermann K, Reeh PW, Vlachová V. Improved
superfusion technique for rapid cooling or heating of cultured cells under patch-clamp
conditions. Journal of Neuroscience Methods. 2006 Mar 15;151(2):178–85.

40.   Elizur A, Adair-Kirk TL, Kelley DG, Griffin GL, Demello DE, Senior RM. Tumor necrosis
factor-alpha from macrophages enhances LPS-induced clara cell expression of
keratinocyte-derived chemokine. Am J Respir Cell Mol Biol. 2008 Jan;38(1):8–15.

41.   Rojewska E, Zychowska M, Piotrowska A, Kreiner G, Nalepa I, Mika J. Involvement of
Macrophage Inflammatory Protein-1 Family Members in the Development of Diabetic
Neuropathy and Their Contribution to Effectiveness of Morphine. Front Immunol. 2018
Mar 12;9:494–494.

42.   Dawes JM, Kiesewetter H, Perkins JR, Bennett DLH, McMahon SB. Chemokine
expression in peripheral tissues from the monosodium iodoacetate model of chronic joint
pain. Mol Pain. 2013 Nov 8;9:57–57.

43.   Syx D, Tran PB, Miller RE, Malfait A-M. Peripheral Mechanisms Contributing to
Osteoarthritis Pain. Current Rheumatology Reports. 2018 Feb 26;20(2):9.

44.   Rosengren S, Boyle DL, Firestein GS. Acquisition, culture, and phenotyping of synovial
fibroblasts. Methods in molecular medicine. 2007;135:365–375.

45.   Ai R, Hammaker D, Boyle DL, Morgan R, Walsh AM, Fan S, et al. Joint-specific DNA
methylation and transcriptome signatures in rheumatoid arthritis identify distinct
pathogenic processes. Nat Commun. 2016 Jun 10;7:11849–11849.

46. Kawashima M, Ogura N, Akutsu M, Ito K, Kondoh T. The anti-inflammatory effect of
cyclooxygenase inhibitors in fibroblast-like synoviocytes from the human
temporomandibular joint results from the suppression of PGE2 production. Journal of oral
pathology & medicine : official publication of the International Association of Oral
Pathologists and the American Academy of Oral Pathology. 2013 Jul;42(6):499–506.

47. Larsson S, Englund M, Struglics A, Lohmander LS. Interleukin-6 and tumor necrosis factor
alpha in synovial fluid are associated with progression of radiographic knee osteoarthritis in
subjects with previous meniscectomy. Osteoarthritis and Cartilage. 2015 Nov
1;23(11):1906–14.

48. Tetta C, Camussi G, Modena V, Di Vittorio C, Baglioni C. Tumour necrosis factor in serum
and synovial fluid of patients with active and severe rheumatoid arthritis. Ann Rheum Dis.
1990 Sep;49(9):665–7.

49. Parsonage G, Falciani F, Burman A, Filer A, Ross E, Bofill M, et al. Global gene
expression profiles in fibroblasts from synovial, skin and lymphoid tissue reveals distinct
cytokine and chemokine expression patterns. Thromb Haemost. 2017 May 12;90(10):688–
97.

50. Taylor PC, Feldmann M. Anti-TNF biologic agents: still the therapy of choice for
rheumatoid arthritis. Nature Reviews Rheumatology. 2009 Oct 1;5:578.

51. Jones DS, Jenney AP, Swantek JL, Burke JM, Lauffenburger DA, Sorger PK. Profiling
drugs for rheumatoid arthritis that inhibit synovial fibroblast activation. Nature Chemical
Biology. 2016 Oct 31;13:38.

52. Pattison LA, Callejo G, St John Smith E. Evolution of acid nociception: ion channels and
receptors for detecting acid. Philosophical Transactions of the Royal Society B: Biological
Sciences. 2019 Nov 11;374(1785):20190291.

53. Akopian AN, Ruparel NB, Jeske NA, Hargreaves KM. Transient receptor potential TRPA1
channel desensitization in sensory neurons is agonist dependent and regulated by TRPV1-
directed internalization. J Physiol. 2007 Aug 15;583(Pt 1):175–93.

54. Wang YY, Chang RB, Waters HN, McKemy DD, Liman ER. The nociceptor ion channel
TRPA1 is potentiated and inactivated by permeating calcium ions. J Biol Chem. 2008 Nov
21;283(47):32691–703.

55. Sarria I, Ling J, Zhu MX, Gu JG. TRPM8 acute desensitization is mediated by calmodulin
and requires PIP(2): distinction from tachyphylaxis. J Neurophysiol. 2011
Dec;106(6):3056–66.

56. Czeschik JC, Hagenacker T, Schäfers M, Büsselberg D. TNF-α differentially modulates ion
channels of nociceptive neurons. Neuroscience Letters. 2008 Apr 4;434(3):293–8.

57. Miller RJ, Jung H, Bhangoo SK, White FA. Cytokine and chemokine regulation of sensory
neuron function. Handb Exp Pharmacol. 2009;(194):417–49.

58. Schaible H-G. Nociceptive neurons detect cytokines in arthritis. ARTHRITIS RESEARCH
& THERAPY. 2014;16(5).

59. Fang D, Kong L-Y, Cai J, Li S, Liu X-D, Han J-S, et al. Interleukin-6-mediated functional
upregulation of TRPV1 receptors in dorsal root ganglion neurons through the activation of
JAK/PI3K signaling pathway: roles in the development of bone cancer pain in a rat model.
Pain. 2015 Jun;156(6):1124–44.

60. Brandolini L, Benedetti E, Ruffini PA, Russo R, Cristiano L, Antonosante A, et al.
CXCR1/2 pathways in paclitaxel-induced neuropathic pain. Oncotarget. 2017 Apr
4;8(14):23188–201.

61. Oh SB, Tran PB, Gillard SE, Hurley RW, Hammond DL, Miller RJ. Chemokines and
Glycoprotein120 Produce Pain Hypersensitivity by Directly Exciting Primary Nociceptive
Neurons. J Neurosci. 2001 Jul 15;21(14):5027.

62. Donatien P, Anand U, Yiangou Y, Sinisi M, Fox M, MacQuillan A, et al. Granulocyte-
macrophage colony-stimulating factor receptor expression in clinical pain disorder tissues
and role in neuronal sensitization. Pain Rep. 2018 Sep 17;3(5):e676–e676.

63. Merabova N, Kaminski R, Krynska B, Amini S, Khalili K, Darbinyan A. JCV agnoprotein-
induced reduction in CXCL5/LIX secretion by oligodendrocytes is associated with
activation of apoptotic signaling in neurons. J Cell Physiol. 2012 Aug;227(8):3119–27.

64. Jung H, Toth PT, White FA, Miller RJ. Monocyte chemoattractant protein-1 functions as a
neuromodulator in dorsal root ganglia neurons. J Neurochem. 2008 Jan;104(1):254–63.

65. Gowler PRW, Li L, Woodhams SG, Bennett AJ, Suzuki R, Walsh DA, et al. Peripheral
brain derived neurotrophic factor contributes to chronic osteoarthritis joint pain. PAIN
[Internet]. 2019;Articles in Press. Available from:
https://journals.lww.com/pain/Fulltext/publishahead/Peripheral_brain_derived_neurotrophi
c_factor.98598.aspx

66. Neogi T. The epidemiology and impact of pain in osteoarthritis. Osteoarthritis and
Cartilage. 2013 Sep 1;21(9):1145–53.

Table 1: AP properties of knee neurons. *** represents p < 0.001 when compared to knee neuron mono-culture, $ represents p < 0.05 when compared to neuron/control FLS co-culture.

	Knee neuron mono-culture (n = 19)		Knee neuron/control FLS (n = 18)		Knee neuron/TNF-FLS (n = 21)		Knee neuron/TNF-FLS media (n = 20)	
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM.
RMP (mV)	-46.6	2.0	-42.8	2.9	-39.7	2.3	-32.0***/$	2.3
Threshold (pA)	447.4	71.5	319.4	56.3	245.2	49.4	315.0	70.0
Half peak duration (HPD, ms)	2.2	0.4	2.3	0.2	2.4	0.5	2.4	0.6
Afterhyperpolarization amplitude (AHP peak, mV)	13.7	2.6	19.6	2.7	16.1	2.9	21.6	3.3

Figure 1

Figure 2

1000   Supplementary Table 1: Genes of interest analyzed in this study.

Gene	TaqMan Assay ID/sequence	Role
Taqman Probes		
18S	Mm03928990_g1	Housekeeping
Cd-68	Mm03047343_m1	Macrophage marker
Cdh-11	Mm00515466_m1	Intimal synovial fibroblast marker
Cd-248	Mm00547485_s1	Synovial fibroblast marker
Cd-31	Mm01242576_m1	Endothelial marker
Il-6	Mm00446190_m1	Inflammation
Il-1r1	Mm00434237_m1	Inflammation
Cox-1	Mm04225243_g1	Constitutively active gene
Cox-2	Mm03294838_g1	Inflammation
RT-PCR primers		
18S	Fwd: CCGGTACAGTGAAACTGCGA Rev: ATCTAGAGTCACCAAGCCGC Product size: 230 bp	Housekeeping
Gpr4	Fwd: ATTCAGCACCGCTCTTCCAT Rev: CAGGGCCAGACGTTTGATCT Product size: 236 bp	Proton-sensing GPCR
Gpr65	Fwd: CAACATCGGATCTTTATGCG Rev: ATGTAGGTGAAGAAAACGCT Product size: 195 bp	Proton-sensing GPCR
Gpr68	Fwd: TTCTCCCTCCTCCTCACCAG Rev: GGCTGAGTGGAGCTTGGTTA Product size: 234 bp	Proton-sensing GPCR
Gpr132	Fwd: CCACTACCTGCGTTTCACCT Rev: CCAGGAAGATGGTGACGACC Product size: 161 bp	Proton-sensing GPCR
Asic1	Fwd: ACACATTCAACTCGGGCCAA Rev: TGCTCCTGGCAAGACACAAA Product size: 250 bp	Acid-sensing ion channel

Asic2	Fwd: TGCTGCCTTACTTGGTGACA Rev: CGGAGTGGTTTGGCATTGTG Product size: 194 bp	Acid-sensing ion channel
Asic3	Fwd: AGAAGGAGCTCTCAAAGGCG Rev: AGGTAACAGGTACGGTGGGA Product size: 158 bp	Acid-sensing ion channel
Asic4	Fwd: AGCGGCTAACTTATCTGCCC Rev: CAAGGGAGTCCAGTGTGTGG Product size: 234 bp	Acid-sensing ion channel
Trpv1	Fwd: GACACCATTGCTCTGCTCCT Rev: GCCTGGACATCTGCTCCATT Product size: 176 bp	Heat and proton transducing ion channel
Il6	Fwd: AGCCAGAGTCCTTCAGAGAGA Rev: TGGTCTTGGTCCTTAGCCAC Product size: 226 bp	Inflammation

1001

1002   Fwd = Forward primer sequence, Rev = reverse primer sequence.

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013 Supplementary Table 2: Location of cytokines detected by the mouse inflammatory antibody

1014 array membrane.

	A	B	C	D	E	F	G	H	I	J	K	L
1	Pos	Pos	Neg	Neg	Blank	BLC	CD30 L	Eotaxin	Eotaxin-2	Fas Ligand	Fractalkine	GCSF
2	Pos	Pos	Neg	Neg	Blank	BLC	CD30 L	Eotaxin	Eotaxin-2	Fas Ligand	Fractalkine	GCSF
3	GM-CSF	IFNγ	IL-1α	IL-1β	IL-2	IL-7	IL-4	IL-6	IL-9	IL-10	IL-12 p40/p70	IL-12 p70
4	GM-CSF	IFNγ	IL-1α	IL-1β	IL-2	IL-7	IL-4	IL-6	IL-9	IL-10	IL-12 p40/p70	IL-12 p70
5	IL-13	IL-17	I-TAC	KC	Leptin	LIX	Lymphotactin	MCP-1	MCSF	MIG	MIP-1α	MIP-1γ
6	IL-13	IL-17	I-TAC	KC	Leptin	LIX	Lymphotactin	MCP-1	MCSF	MIG	MIP-1α	MIP-1γ
7	RANTES	SDF-1	TCA-3	TECK	TIMP-1	TIMP-2	TNF-α	sTNF RI	sTNF R II	Blank	Blank	Pos
8	RANTES	SDF-1	TCA-3	TECK	TIMP-1	TIMP-2	TNF-α	sTNF RI	sTNF R II	Blank	Blank	Pos

1015

BLC = B lymphocyte chemoattractant, GCSF = granulocyte colony stimulating factor, GM-CSF

= granulocyte-macrophage colony stimulating factor, IFN = interferon, IL= interleukin, I-TAC =

interferon inducible T-cell alpha chemoattractant, KC = Keratinocyte chemoattractant (chemokine

ligand 1), LIX = lipopolysaccharide induced chemokine, MCP = monocyte chemoattractant

protein, MCSF = macrophage colony stimulating factor, MIG = monokine induced by gamma

interferon, MIP = macrophage inflammatory protein, RANTES = Regulated on activation, normal

T-cell expressed and secreted, SDF= stromal cell derived factor, TCA = T-cell activation gene,

TECK = thymus expressed chemokine, TIMP = Tissue inhibitor of metalloproteinase, TNF(R) =

tumor necrosis factor (receptor), Pos = Positive spot, Neg = negative spot.

1025

Characterization of FLS

Expression of inflammation-related genes in FLS derived from CFA-injected knee joint

MIP-1γ activation of DRG neurons

1028

1029

Supplementary Figure 1: Characterization of mouse primary FLS and neuronal activation of FLS
secreted MIP-1γ. A) Bars showing reduction in mRNA expression of the macrophage marker *Cd-*
68 from P2 to P3 expressed as fold change from macrophage cell line Raw 264.7 cells. ** p <0.01,
ratio paired t-test. Bi) Expression of FLS marker *Cdh-11* across P2-P5 in ipsi and contra FLS
normalized to housekeeping gene *18S* (left) and a representative image of CDH-11 (red) and DAPI
(blue) stained FLS in culture. ii) Expression of FLS marker *Cd-68* across P2-P5 in ipsi and contra
FLS normalized to housekeeping gene *18S*. Ci) Mouse knee width in Contra (grey circles) and Ipsi
(open circles) limbs before and after injection of CFA (n = 6, paired t-test. ii) Bars represent fold
change of the genes *Il-6*, *Il-1r1*, *Cox-2* and *Cox-1* from Contra vs. Ipsi. Black bars = P2, grey bars
= P3. D) Intracellular Ca^{2+} influx in lumbar DRG neurons in response to 20 s application of 100
ng/ml MIP-1γ, followed by percentage of neurons responding to 10 and 100 ng/ml of MIP-1γ and
their respective peak response. > 500 neurons imaged from three male mice. *** $p < 0.001$, chi-
sq test. Error bars represent SEM.

1043

1044

1045