REPURPOSING THE *STREPTOCOCCUS MUTANS* CRISPR-CAS9 SYSTEM TO UNDERSTAND ESSENTIAL GENE FUNCTION

Shields RC[1], Walker AR[1], Maricic N[1], Underhill SAM[2], Burne RA[1]

[1]Department of Oral Biology, College of Dentistry, University of Florida, Gainesville, Florida, USA;

[2]Department of Physics, University of Florida, 2001 Museum Road, Gainesville, Florida, USA

ORCiDs:

Shields: http://orcid.org/0000-0002-8823-9504

Underhill: https://orcid.org/0000-0002-5214-4562

Burne: https://orcid.org/0000-0002-4234-0316

Keywords: Gene silencing, gene functional genomics, exopolysaccharides, cell division, dental caries

Corresponding Author Information:

Robert C. Shields

Department of Oral Biology

University of Florida College of Dentistry

1395 Center Drive

Gainesville, FL 32610

Phone: (352) 273-8850

Fax: (352) 273-8829

E-mail: rshields@dental.ufl.edu

Abstract

A recent genome-wide screen identified ~300 essential or growth-supporting genes in the dental caries pathogen *Streptococcus mutans*. To be able to study these genes, we built a CRISPR interference tool around the Cas9 nuclease (Cas9$_{Smu}$) encoded in the *S. mutans* UA159 genome. Using a xylose-inducible dead Cas9$_{Smu}$ with a constitutively active single-guide RNA (sgRNA), we observed titratable repression of GFP fluorescence that compared favorably to that of *Streptococcus pyogenes* dCas9 (Cas9$_{Spy}$). We then investigated sgRNA specificity and proto-spacer adjacent motif (PAM) requirements. Interference by sgRNAs did not occur with double or triple base-pair mutations, or if single base-pair mutations were in the 3' end of the sgRNA. Bioinformatic analysis of >450 *S. mutans* genomes allied with *in vivo* assays revealed a similar PAM recognition sequence as the Cas9$_{Spy}$. Next, we created a comprehensive library of sgRNA plasmids that were directed at essential and growth-supporting genes. We discovered growth defects for 77% of the CRISPRi strains expressing sgRNAs. Phenotypes of CRISPRi strains, across several biological pathways, were assessed using fluorescence microscopy. A variety of cell structure anomalies were observed, including segregational instability of the chromosome, enlarged cells, and ovococci-to-rod shape transitions. CRISPRi was also employed to observe how silencing of cell wall glycopolysaccharide biosynthesis (rhamnose-glucose polysaccharide, RGP) affected both cell division and pathogenesis in a wax worm model. The CRISPRi tool and sgRNA library are valuable resources for characterizing essential genes in *S. mutans*, some of which could prove to be promising therapeutic targets.

Introduction

Streptococcus species are natural inhabitants of humans and animals. Under the right conditions many streptococci can cause or exacerbate diverse infections in humans, including life-threatening illnesses that include meningitis or pneumonia [1]. The transition between commensalism and pathogenicity is an important feature of the oral streptococci, which is the most abundant bacterial genus in the human mouth [2]. Oral streptococci have important roles in the initiation and maturation of dental biofilms (plaque), and many are associated with oral and dental health. However, oral streptococci that are especially acidogenic and acid-tolerant are implicated as primary etiologic agents in the pathogenesis of dental caries (tooth decay) [3]. Dental caries is a common cause of pain, impaired quality of life, and tooth loss in children and adults [3]. The organism most often linked to caries development is *Streptococcus mutans*, which has been shown to have multiple virulence traits, some unique, that imbue the organism with the capacity to initiate and worsen the disease [4]. Extraorally, *S. mutans* and other oral streptococci can cause infective endocarditis [5]. More recently, epidemiologic and mechanistic studies point to a role for certain *S. mutans* strains in cardiovascular diseases, including stroke [6]. Investigating the biology of this organism is important for developing therapies that combat dental caries and other diseases associated with *S. mutans* and related organisms. More broadly, expanding the understanding of streptococcal biology will positively impact efforts to prevent or treat many other diseases in humans and animals.

Antimicrobial therapies usually target biological processes that are essential for bacterial growth and persistence. These pathways include protein synthesis, transcription, DNA replication, cell division, peptidoglycan biosynthesis, or folic acid metabolism. The most traditional approach for discovering new antibiotics has been screening the ability of natural and synthetic compounds to inhibit growth or processes essential for pathogenesis [7]. Alternatively, antibacterial targets could be identified by combining genome sequence data with genome-wide techniques that probe gene essentiality, followed by the identification or design of molecules that target the factors involved in the process of interest [8]. Global identification of essential genes is possible with a technique known as transposon sequencing (Tn-seq) [9]. Using Tn-seq, we identified a panel of genes that appears essential for the growth of *S. mutans* UA159, together with genes that are required for growth *in vitro* in rich or defined medium [10]. Although Tn-seq can be used to discover essential genes, it does not yield conditional mutations, and therefore does not allow for the functional study of these genes unless culture conditions can be found that allow for growth. Before moving toward studies that attempt to block or inhibit these pathways in *S. mutans*, or other streptococci, there is a need for further validation and investigation of these apparently essential genes. However, functional studies of essential genes in bacteria have traditionally been technically challenging and time-consuming, because mutants of these genes cannot be isolated and studied. One approach involves the generation of a conditional knockout, usually by mutating an essential gene and simultaneously expressing a copy of the gene under the control of a promoter that can be activated by an exogenously provided compound [11]. Two major drawbacks to this approach are that it is not easily adapted to a comprehensive study of the essential genome of an organism in a time-efficient manner (low throughput) and differences in gene expression levels between the conditional expresser and the wild type can generate misleading results.

Reprogramming of the bacterial CRISPR (clustered regularly interspaced short palindromic repeats) - Cas system into a gene-silencing tool has made it more practical to perform high-throughput studies of essential genes [12,13]. This technique, known as CRISPR interference (CRISPRi), combines an inducible, catalytically inactive *Streptococcus pyogenes* Cas9 nuclease, dead Cas9 (dCas9), with a programmable single-guide RNA (sgRNA). The sgRNA encodes a 20-nt sequence that is complementary to the target DNA. When expressed together in a cell, the sgRNA-dCas9$_{Spy}$ complex represses transcription initiation or elongation of the target gene. CRISPRi libraries that silence all putatively essential genes have been generated for both *Streptococcus pneumoniae* D39 [14] and *Bacillus subtilis* 168 [15]. A complicating factor for developing a CRISPRi system in *S. mutans* UA159 is

the presence of an endogenous Cas9 nuclease (SMu.1405c) [16]. Introduction of a CRISPRi system
into *S. mutans* built around dCas9$_{Spy}$ would likely not function without the deletion of the endogenous
Cas9, because of the potential for Cas9$_{Smu}$-sgRNA complexes to cause lethal double-strand breaks in
the chromosome. In fact, Cas9$_{Smu}$ and Cas9$_{Spy}$ can cleave DNA in the presence of a *S. pyogenes* dual-
RNA, and vice versa [17]. The Cas9 target recognition sequence, known as the proto-spacer adjacent
motif (PAM), is also predicted to be 5'-NGG'-3' in *S. mutans*, as is the case in *S. pyogenes* [18]. Given
the similarities, one possible approach to CRISPRi in *S. mutans* could be to repurpose the endogenous
Cas9$_{Smu}$ nuclease into a nuclease-dead version, while also using elements of the original interference
system (*e.g.* sgRNA). A CRISPRi system built around Cas9$_{Smu}$ could function more effectively in *S.*
mutans than one based on Cas9$_{Spy}$. Heterologous expression of *S. pyogenes* dCas9 can be toxic to
certain bacteria [19,20], although problems of this nature were not observed in a recent CRISPRi study
in *S. pneumoniae* [14]. Irrespective of *S. pyogenes* dCas9 toxicity, the dCas9$_{Spy}$ could perform
suboptimally compared to dCas9$_{Smu}$ simply due to codon-bias. Lastly, developing an *S. mutans*
CRISPRi system has the potential to add to the number of tools for genome editing that are currently
available.
111
Motivated by a desire to functionally study essential genes in *S. mutans,* we created a CRISPRi tool
assembled around dCas9$_{Smu}$. We explored sgRNA specificity and PAM requirements, and compared
interference activity with dCas9$_{Spy}$. Having demonstrated good CRISPRi activity, we then constructed a
library of sgRNAs against all putative essential and growth-supporting genes [10]. With this library we
evaluated growth and certain phenotypes in a subset of strains with CRISPRi-silenced essential genes.
Finally, we used the technique to systematically study the impact that depletion of L-rhamnose-
containing cell wall polysaccharides has on cell division and pathogenicity in an insect model.
119
Results
Robust gene knockdown with a titratable dCas9$_{Smu}$ system
More than 95% of the 477 *S. mutans* strains for which whole or draft genome sequences are presently
available carry at least one CRISPR-Cas system (Fig 1A). The Type II-A CRISPR-Cas system, which
can be repurposed for CRISPRi, is the most prevalent across the species with 339/477 strains carrying
a complete system (Fig 1A). For reprogramming of *S. mutans* UA159 Cas9 (SMu.1405c) into a
CRISPRi system, we began by mutating two amino acid residues (D10 and H840) known to be required
for the nuclease activity of *S. pyogenes* Cas9 [21] (Fig 1B). We were able to confirm the DNA-binding
and repressing activity of dCas9$_{Smu}$ by expressing a sgRNA against a constitutively expressed *gfp* gene
[22] (Fig. 1C). Next, the *S. mutans dcas9* gene was cloned into pZX9 [23], such that expression could
be activated by xylose. Prior to this, we discovered, using green fluorescent protein (GFP) as a readout
for xylose-induced gene expression, that the P$_{xyl}$ promoter was less active when glucose was the sole
carbohydrate source compared to maltose (Fig S1). We suspect the different expression levels on
glucose versus maltose could be due to carbohydrate catabolite repression and/or effects of certain
phosphoenolpyruvate:sugar phosphotransferase system (PTS) components on P$_{xyl}$ activity. Thus, we
used the defined media 'FMC' containing maltose for all CRISPRi related experiments. To make the
CRISPRi strain (Fig 1D), the sgRNA plasmid was integrated in single copy into the *S. mutans* genome
in a strain that contained i) a *gfp* gene driven by the constitutively active P23 promoter and integrated
into the chromosome, ii) the P$_{xyl}$-*dcas9*$_{Smu}$ plasmid and iii) point mutations that prevent translation of
SMu.1405c (*cas9*; start codon mutated to stop codon). *S. mutans* Cas9 must be mutated, otherwise
endogenous *cas9* expression in combination with sgRNAs would likely cause lethal chromosome
damage; *S. mutans* is unable to repair the double-strand cleavage catalyzed by a functional Cas9.
142
Having developed the system, we moved to assessing its efficiency. Using the *S. mutans* CRISPRi
technique, we observed titratable repression of GFP fluorescence (Fig 1E) and GFP protein levels (Fig
1F). Importantly, there were no obvious effects on the growth *S. mutans* as increasing amounts of
xylose were added (up to 0.025% w/v). Also of note, slightly higher amounts of xylose (0.025% vs.

0.00625%) were required to achieve similar levels of repression of GFP production in a strain
containing a xylose-inducible *S. pyogenes dcas9* (Figs 1E and F). We also observed slight growth
impairments as increasing amounts of dCas9$_{Spy}$ were produced (Fig 1E). Overall, the differences in
performance of the two Cas9 constructs were minor. Lastly, we measured gene-silencing (absence of
GFP fluorescence) at the single-cell level (Fig S2). The number of cells escaping CRISPRi-mediated
gene knockdown (GFP positive) was very small, indicating that, at saturating amounts of xylose, the
effects of the *S. mutans*-derived CRISPRi system are uniform across the population when the dCas9
protein is delivered on a multi-copy plasmid.
155
Guide RNA design principles for the sgRNA-Cas9$_{Smu}$ system
To begin to optimize the *S. mutans* CRISPRi system, we examined parameters that are important for
functionality; the first of these was sgRNA specificity. sgRNA-Cas9 systems that operate with lower
sequence stringency could be prone to off-target effects that occur when guide RNA and dCas9 bind to
a site(s) in the genome that was not intended to be affected. To test this, we cloned an sgRNA with a
complementary sequence for the gene *atpB*, which encodes the beta subunit of the F_1F_O proton-
translocating ATPase. We modified the sgRNA-*atpB* sequence by introducing substitutions along the
length of guide RNA (Fig 2A). The sgRNA plasmids were added to competent *S. mutans* and plated
onto selective agar. We expected that transformation into *S. mutans* of the sgRNA-*atpB* plasmid lacking
sequence modifications would not be tolerated, leading to a very low transformation efficiency. As
expected, transformation of the sgRNA-*atpB*^{T} (T; targeting the template strand) occurred at very low
efficiency, 1 x 10^{-5} (Fig 2B). Conversely, transformation efficiency in a Δ*cas9* (deletion of Smu.1405c)
strain, the dead *cas9* strain, or in a strain carrying a deletion of 20-bp of the *atpB*^{T} sequence from the
sgRNA (sgRNAneg) was 100- to 1000-fold more efficient than with the intact *atpB* sgRNA (Fig 2B).
170
We next evaluated the effects of expressing sgRNAs with one, two or three base mutations within the
atpB^{T} sequence. Single base-pair mismatches were tolerated at the 5' end of the sgRNA (*atpB*_3,
*atpB*_6, *atpB*_9 and *atpB*_12). Plasmids with mismatches closer to the PAM (*atpB*_15 and *atpB*_18)
transformed with higher efficiency into *S. mutans*, indicative of a loss of sgRNA activity. All of the
double and triple base mismatches reduced the activity of the sgRNAs, as these plasmids transformed
S. mutans at levels comparable to the *cas9*-deficient strain (Fig 2B). These results suggest that near
perfect sgRNA sequence complementarity is required to silence genes, and thus it should be possible
to minimize off-target effects with careful design of sgRNAs.
179
Another important feature of CRISPR-Cas9 systems, both as a natural bacterial immune system and
for CRISPRi, is the proto-spacer adjacent motif (PAM). This sequence is required for the Cas9 protein
to recognize interference targets and is also critical during the acquisition of CRISPR spacers (for Type
II-A systems) [21,24]. For a given bacterial strain, Type II-A PAM sequences are generated by aligning
CRISPR spacers to phage genomes, extracting the sequence downstream of the spacer target, and
then building a conserved sequence from these. The difficulty with taking this approach with *S. mutans*
UA159 is that there are too few CRISPR spacers from which to build a reliable PAM sequence.
Consequently, we compiled PAM sequences from a large (>450) collection of *S. mutans* genomes.
After narrowing the data to strains that had Type II-A CRISPR-Cas systems and ≥5 spacers, PAM
sequences were generated for 100 individual *S. mutans* strains using WebLogo (Fig S3). Per this
bioinformatics analysis, a highly conserved PAM for Cas9$_{Smu}$ was not evident. Rather, a significant
number of *S. mutans* strains with Cas9 contained spacers that might require the 5'-NGG-3' recognition
sequence for phage DNA interference, and these spacer arrays grouped together by alignment (Fig 3A,
yellow color). For other *S. mutans* strains with Cas9, PAM sequences were more random, but at least
two other groups, 5'-TTN-3' and 5'-NAAA-3', existed (Fig S3). We observed no obvious grouping for *S.*
mutans of the predicted PAM sequences by Cas9 protein identity (Fig 3A). We reasoned that PAM
recognition might be protein sequence specific and thus expected that the different PAM groups
predicted by bioinformatics would also group together. A lack of coherence between the predicted PAM

sequences and Cas9 protein sequences suggests that the PAM sequences assembled via
bioinformatics may not be the best indicator of the requirements for recognition for all *S. mutans* strains.
So, to investigate DNA recognition in greater detail we developed an interference assay with GFP
fluorescence as the readout. Using site-directed mutagenesis we incorporated assorted PAMs into a
construct in which the P23 promoter could drive the expression of a *gfp* gene (Fig 3B). This assay
confirmed that *S. mutans* dCas9 recognizes a 5'-NGG-3' PAM (Fig 3C). With 5'-NAG-3' we observed
moderate CRISPRi activity with dCas9$_{Smu}$, although the results varied as a function of the DNA base
present in the first position, with up to an 80% loss of GFP fluorescence. Weak binding of *S. pyogenes*
Cas9 to a 5'-NAG-3' PAM has been reported before [25–27]. Nuclease activity via 5'-NGA-3'
recognition has previously been reported for Cas9$_{Spy}$ [26,27] but we were unable to observe CRISPRi
activity with either dCas9 protein with a 5'-AGA-3' PAM. No interference was observed with any of the
other PAM sequence combinations tested.
210
Creation of a comprehensive library of essential gene targeting sgRNAs
We constructed a library of sgRNAs targeting 253 potentially essential and growth supporting genes
(Fig 4A). These genes (Table S1) were selected from previously conducted Tn-seq experiments [10],
Tn-seq performed as part of this study, and genes shown to be essential in previous studies (e.g.
vicR/walR). We were unable to design sgRNAs for 21 putative essential or growth supporting genes, of
which over half (12) were ribosomal proteins. These genes were small (average size of 275-bp) and
generally lacked PAM sites that would give desirable sgRNA sequences (*e.g.* RNA folding, off-target
effects, location in 5' region of gene). During the first round of cloning (see Materials and Methods) we
obtained correct sgRNA clones for 181 of 282 (64%) of the desired strains. After three rounds of
cloning 259/282 (92%) of the sgRNAs were successfully cloned. We were unable to obtain sgRNA
clones for 23 (8%) genes. After plasmid cloning in *E. coli* 10-beta, sequence-verified sgRNA plasmids
were transformed into *S. mutans* Δ*cas9* P$_{xyl}$-*dcas9*$_{Smu}$. While it would be possible to clone the sgRNA
plasmids directly into *S. mutans*, we wanted to minimize the number of passages in *S. mutans* as
CRISPRi strains are known to acquire suppressors when guide RNAs are expressed in their hosts [14].
225
Growth phenotypes of strains with silenced essential genes
We measured the growth of each sgRNA strain in FMC-maltose either in the absence or presence of
0.1% xylose for 16 h using a Bioscreen C Automated Microbiology Growth Curve System. Afterwards,
the area under the curve (AUC) was calculated for both conditions. Quantification of growth by AUC is
useful as it combines growth phases (lag, exponential and stationary phases). For each strain, the
AUC$_{xylose}$ was divided by AUC$_{uninduced}$. Growth phenotypes were grouped per these ratios: >0.5, no
phenotype; 0.25-0.49, moderate growth defect; >0.25, strong growth defect). Growth defects were
observed in 194/253 (77%) of the strains expressing sgRNAs that were designed to target essential or
growth supporting genes (Fig 4B; S1 Table). Overall, we observed a higher percentage of growth
defects for strains with silenced essential genes versus growth-supporting genes, 82% and 62%,
respectively (Fig 4B). When comparing between different Clusters of Orthologous Groups (COGs), the
lowest median growth defects were observed for silenced genes in the translation (J) and lipid
metabolism and transport pathways (I) (Fig 4C). The highest median growth defects were observed
when sgRNAs targeted genes related to coenzyme transport and metabolism (H) and signal
transduction mechanisms (T) (Fig 4C). Growth defects were not observed for the six sgRNAs that
targeted non-essential genes. There was no significant relationship between the growth defect we
observed and the position of the sgRNA within the gene or the size of the target gene (Fig. S4).
243
Assessing cell biology phenotypes of CRISPRi strains with microscopy
Next we evaluated the morphological phenotypes of eighty CRISPRi strains placed in the following
functional categories: fatty acid biosynthesis/membrane biogenesis, peptidoglycan synthesis,
glycolysis, DNA replication, transcription, translation, cell division, and ATP synthase activity. Using
DAPI staining, we were able to visualize the bacterial chromosome, and with nile red we could detect

the cell membrane. The cell morphology defects of essential gene-targeted strains were diverse (Fig 5).
Defects included increased cell chaining, abnormally large cells, heterogeneously sized cells, cell lysis
and anucleate cells (Figs S5-S12). The most commonly observed morphological phenotypes across all
the pathways investigated were increased cell chaining (*e.g. pfk*, *accD*, *dnaE*, *rs3*, *atpC*) and
heterogeneously sized cells (*e.g. pgi*, *glmM*, *dnaG*, *eftS*, *atpF*) (Fig. 5A). It is important to note that
there were strains without obvious growth defects that displayed major cell morphology defects (*e.g.*
ftsX and *glmS*). Repression of genes involved in the glycolysis pathway led to significant growth defects
and larger/misshapen cells (Fig S5). When *pgm*, encoding phosphoglucomutase, was repressed, cells
appeared to be somewhat rod-shaped, probably associated with altered or deficient biosynthesis of cell
wall polysaccharides. When genes involved in peptidoglycan biosynthesis were repressed, the most
common phenotype was the formation of singular large round cells (Fig S6). The percentage of cells
displaying this phenotype varied, with it being most common in a strain with silenced *murM*, which in *S.*
pneumoniae encodes a protein that incorporates amino acids into the side-chain of peptidoglycan [28].
Repression of genes involved in the conversion of acetyl-CoA to malonyl-CoA (*accACD* and *bccP*), an
important step in fatty acid biosynthesis, led to excessive chaining of cells (Fig S7). Repression of fatty
acid biosynthetic (*fab*) genes yielded minor phenotypic changes, whereas silencing of genes
downstream of the FAS II pathway (*plsX* and *cdsA*) led to more obvious morphological defects. As
anticipated, when genes involved in DNA replication were repressed, not all cells contained nucleoids
(Fig S8). In addition to anucleate cells, certain CRISPRi strains with repressed chromosome replication
also showed severe morphological defects (e.g. *dnaE*, *dnaG*, *priA*). CRISPRi repression of translation-
related genes increased cell chaining (Fig S9). When the ribosomal proteins *rs3* (small ribosomal
subunit) and *rl11* (large ribosomal subunit) were repressed, we observed anucleate cells that lacked a
well-defined cell envelope. CRISPRi silencing of genes encoding subunits of the F_1F_O-ATPase led to
strong growth defects, increased cell chaining and variability in nile red staining (Fig S10). Repression
of transcription-related genes also led to increased cell chaining and attenuated growth (Fig S11). The
cell shape and size of *rpoB*-, *rpoC*-, and *rpoZ*-silenced CRISPRi strains were severely impacted. When
genes involved in cell division were repressed, strains displayed heterogeneity in cell size and shape
(Fig S12). Notably, with repression of *divIC*, *ftsA*, *ftsL* and *ftsZ* we observed similar morphological
changes. Cells appeared to be rod-shaped, except that they bulged at the cell septum, probably
consistent with an inability of these cells to complete septation. FtsZ is a tubulin homologue that forms
a Z-ring at the midcell that is important for scaffolding and cytokinesis [29]. FtsA, DivIC and FtsL also
localize to the midcell (at different stages) during cell division [29]. Repression of *gbpB* (glucan-binding
protein B), an ortholog of the *S. pneumoniae* peptidoglycan hydrolase gene, *pcsB*, led to cell division
defects. Although specific peptidoglycan hydrolase activity has not been observed for GbpB, the cell
division defects we observed, along with the observation that GbpB can be localized to areas of active
peptidoglycan synthesis [30], provide support for the idea that it may act enzymatically to affect the
structure of the cell wall. Taken together, microscopic analysis of CRISPRi strains from different
pathways has revealed generalized phenotypes of essential gene depletion, but also clear functional
links between genes in established pathways.
288
L-rhamnose glycopolymers are critical for *S. mutans* cell division and pathogenicity
Having shown that CRISPRi allows for the functional study of essential genes, we next focused our
attention on a pathway that is thought to be of major importance to *S. mutans*, rhamnose-glucose
polysaccharide (RGP) biosynthesis. Certain streptococcal species lack cell wall teichoic acids (WTAs)
and instead produce RGPs that are covalently linked to the peptidoglycan [31]. RGP biosynthesis
begins with a four-step pathway encoded by the genes *rmlABCD* that converts glucose-1-phosphate
into dTDP-L-rhamnose. The last gene in the pathway, *rmlD*, resides in an operon (SMu.824-835) that
contains genes required for additional RGP biosynthesis steps and transport of the carbohydrates
through the cell membrane. By our Tn-seq analysis, most of the genes involved in producing RGP are
either essential or required for the growth of *S. mutans* UA159 (Fig 6A) [10]. L-rhamnose biosynthetic
pathways are also essential in *S. pyogenes* [32] and *Mycobacterium smegmatis* [33]. Several of these

genes have, however, been mutated in *S. mutans*, including *rmlBC* [34], *rmlD* [32], *rgpE* [30,35], *rgpF*
[35,36], and *rgpG* [37]. Inconsistencies between our observations and others could be related to the
nature of our Tn-seq selection and/or the presence of undetected suppressor mutations that bypass
essentiality in strains that carry RGP gene deletions in other studies. Use of CRISPRi to study RGP
function facilitates the rapid evaluation of depleting each gene product under conditions that allow for
selective pressure to be applied only when xylose is provided in the medium. By CRISPRi, down-
regulation of every gene directly involved in RGP biosynthesis displayed severe growth defects when
silenced with sgRNAs (Fig 6B; Fig S13). In the invertebrate infection model organism *Galleria*
mellonella, strains with silenced *rmlB*, *rgpC*, and *rgpF* genes displayed attenuated virulence compared
to wild-type *S. mutans* UA159 and a d*cas9* strain (Fig 6C). For this proof-of-concept study, sgRNAs
were expressed in an *S. mutans* strain where the endogenous Cas9 had been converted to dCas9. A
strain with silenced *rmlD* was not attenuated. However, we suspect this strain acquired suppressor
mutation(s) as it originally grew very poorly, but after some passage it was able to grow to a final OD_{600}
similar to the parental strain. Having shown that RGP biosynthesis genes are required for growth in
laboratory media and during infection, we next investigated their importance to *S. mutans* cell biology.
Consistent with previous observations [30,37], cellular morphologies were exceptionally aberrant,
compared to the wild type (Fig S13). Cells were large, not consistently ovococcoid, and formed longer
chains than normal. Although resolutions were limiting, it also appeared that certain large cells
contained multiple nucleoids by DAPI strain (Fig S14). To gain greater ultrastructural insights, we
investigated cell biology phenotypes of *rmlB*, *rmlD*, *rgpC*, and *rgpF* silenced strains with transmission
electron microscopy (TEM). In these strains, division site positioning was irregular and not always in the
mid-point of cells (Fig 6D). There were instances where the division septum was unable to traverse the
cell (*rmlB*, *rmlD, rgpF*) or cells had double septa (*rgpF*) (Fig 6D). Accordingly, large cells with
incomplete septation appeared to have multiple unequally sized compartments, which likely has a
deleterious effect on chromosome segregation, and might explain the unusual patterns of DAPI staining
observed during fluorescence microscopy. In *rgpC*-silenced cells, the equatorial ring, an outgrowth of
cell wall material that is normally duplicated and separates as cells divide, was abnormally thick and in
some cases was 'peeling' back from the cell (Fig 6D). Measurements from the CRISPRi strain
micrographs using ImageJ confirmed that they were wider than wild-type cells (Fig 6E).
329
Discussion
Essential genes represent promising targets for antimicrobials. The essential genome of *S. mutans* is
comprised of ~300 genes and, like other bacteria, is enriched for genes that function in processing
genetic information, energy production, and maintenance of the cell envelope. However, until the recent
development of CRISPRi technologies, high-throughput functional studies of these genes has not been
possible. Here we demonstrate the conversion of the *S. mutans* Cas9 nuclease into a gene silencing
tool for the systematic interrogation of the essential genome. Our system is relatively streamlined as it
only requires a constitutively expressed sgRNA and a xylose-inducible dCas9. Importantly, it allowed us
to study all the known essential genes in a relatively short time. Most of the genes predicted to be
essential by Tn-seq displayed strong growth defects once silenced by CRISPRi. Furthermore, CRISPRi
strains with sgRNAs targeting essential genes displayed morphological defects when analyzed by
microscopy. This work confirms the importance of these genes to the integrity of *S. mutans* cells, and of
their potential as interesting therapeutic targets for the control of *S. mutans* or certain other pathogenic
streptococci.
344
One interesting finding of this work is that $Cas9_{Smu}$ recognition of target DNA shares similarities with
$Cas9_{Spy}$. First, guide RNA specificity of $Cas9_{Smu}$ is similar to $Cas9_{Spy}$ as single base-pair mismatches
are not tolerated in the 5' region of sgRNA (the seed region) [38,39]. $Cas9_{Smu}$ was also unable to cause
DNA cleavage *in vivo* when double or triple base-pair mutations were introduced into the sgRNA
sequence, again consistent with findings in $Cas9_{Spy}$ [39]. Second, PAM recognition relied on a 5'-NGG-
3' sequence. We also observed moderate CRISPRi activity with a 5'-NAG-3' PAM. $Cas9_{Spy}$ activity also

relies on a 5'-NGG-3' PAM with weak DNA recognition reported for the non-canonical PAMs 5'-NAG-3'
and 5'-NGA-3' [25–27], although we were unable to measure this with our assay. Although target
recognition also relies on other factors [40], we are confident that off-target effects should be minimized
when using a 5'-NGG-3' PAM and a guide RNA designed to ensure a lack of complementarity (two or
more mismatches) to other regions of the genome. The similarities of $Cas9_{Smu}$ and $Cas9_{Spy}$ are
interesting given that other streptococcal Cas9 nucleases require PAMs that differ to 5'-NGG-3' [20,41].
For example, the Cas9 of *Streptococcus canis*, which shares ~90% sequence similarity to $Cas9_{Spy}$,
recognizes a 5'-NNG-3' PAM [41]. Bioinformatics analysis of the species-wide PAM requirements were
cryptic, but it is possible that other *S. mutans* Cas9 proteins recognize alternative PAMs *e.g.* 5'-NAAA-
3'. Intriguingly the Cas9 of *Streptococcus macacae* NCTC 11558 is presumed to recognize an adenine
enriched PAM [42]. There is an arms race between the bacteria that generate Cas9 proteins
recognizing new PAMs and the phage that undergo mutations to avoid CRISPR interference, with the
latter being a confounding issue with PAMs that are generated solely using bioinformatic tools. Despite
these limitations, it is possible there is species-wide PAM diversity, similar to what has been observed
with *Neisseria meningitidis* [43]. This is an interesting area of study as it should aid in the engineering of
new variants of Cas9 that recognize different PAMs, thus expanding the genetic engineering
capabilities of the Cas9 proteins.
368
The CRISPRi system described here was able to perturb the expression of over 250 putative essential
and growth-supporting genes. The severe growth defects exhibited by most of these strains
emphasizes their critical functions in *S. mutans*. The percentage of silenced genes that displayed
significant growth defects was comparable with what was observed for *S. pneumoniae* [14]. As noted
above, CRISPRi has many advantages over more traditional approaches for studying essential genes,
but the technique has its limitations. For example, failure to display an aberrant growth phenotype could
be caused by sub-optimal sgRNA design or selection for suppressor mutations that bypass essentiality.
Likewise, certain essential genes could still function with a low number of protein molecules per cell if
the target gene is no completely silenced. In some cases, strains that displayed relatively minor growth
defects when the target genes were depleted had significant alterations in cell morphology. Tn-seq,
which was used to identify to candidate genes for this study (REF), is a competitive assay, so even
mutants with minor growth phenotypes could be significantly underrepresented at the chosen end-point.
In this study, we supplemented our previous Tn-seq essential gene analysis [10] with a different Tn-seq
approach, which involved plating the original library on BHI instead of blood agar, then passaging the
libraries in BHI or FMC for ~10 generations, as compared with ~30 generations in our previous study
(REF). The differences between the older and the newer Tn-seq analysis were minor (Table S1).
Overall, the CRISPRi growth study paired with comprehensive Tn-seq enquiries provide a solid
foundation for determining which genes are essential in *S. mutans* UA159. It is reasonable to predict
that some of these gene products could be targeted for the therapeutic control of *S. mutans* and
possibly some other pathogenic streptococci.
389
In addition to growth studies, we were able to observe morphological phenotypes of CRISPRi strains
with light and electron microscopy. Several morphology phenotypes were evident that were consistent
with other high-throughput microscopic studies of essential gene depletions [14,15]. The changes in
morphology could represent more generalized responses to an inability to grow or, in some cases, may
prove useful in linking gene products that function in concert. For example, *ftsA*, *ftsZ*, *ftsL* and *divIC*
shared similar phenotypes, consistent with their products being required for correct cell septum
formation. Perturbation of peptidoglycan synthesis led in some cases to the formation of large round
cells. Blocking or deleting peptidoglycan synthesis genes in other bacteria can lead to the generation of
giant cells [44] or L-forms [45], although the phenotypes in this study appear distinct from those
previously described variants.
400

The rhamnose-glucose polysaccharide (RGP) of mutans streptococci were used in the past to
differentiate serotypes of strains that were previously all designated as *S. mutans.* Today, it is
recognized that the species *S. mutans* consists primarily of Bratthall serotypes *c, e,* and *f*, along with
the unusual serotype *k* strains [46,47]. In contrast, what was formerly designated serotype *b S. mutans*
is now classified as *Streptococcus ratti*. It is also now well recognized that RGP is an important
constituent of the envelope of mutans streptococci. Clear impacts on growth, pathogenicity and cell
division were observed in strains that had silenced genes contributing to RGP biogenesis. The cell
morphology phenotypes were stark, with depletion strains clearly exhibiting aberrant septum formation,
as has been noted before in *S. mutans* and other related species [30,48,49]. An inability to divide
normally has been linked to aberrant localization of the peptidoglycan hydrolase PcsB (GbpB in *S.*
mutans) [30,49]. When *gbpB* was targeted with CRISPRi, cells appeared to be unable to divide
correctly (Fig S12), but the phenotype did not match that associated with *rgp* gene silencing. Given the
vital role and essentiality of the RGP biosynthetic pathway, coupled with the decreased virulence of
certain *rgp* knockdowns reported here, RGP biogenesis may be an attractive antimicrobial target.
Further, if RGP and peptidoglycan synthesis are linked, it is conceivable that targeting RGP
biosynthesis could potentiate the efficacy of β-lactam and other cell wall-active antibiotics. Other
attractive aspects of the RGP pathway as an antimicrobial target include the species or strain-specificity
of the glycopolymers and the absence of the pathway in humans [31].
419
In summary, this is the first study to utilize CRISPRi in *S. mutans* as a genetic tool. We did so by
repurposing the Cas9 that is widely distributed among the species, opening new avenues for
investigating the diversity of CRISPR-Cas in *S. mutans* and other oral streptococci, and the possible
development of next-generation CRISPR-Cas tools. We demonstrate that the technique provides a
powerful approach for studying essential genes and gene function, in agreement with studies using
other organisms [14,15]. We anticipate that these tools can be used to investigate new aspects of *S.*
mutans cell biology and to identify promising therapeutic candidates.
427

Materials and methods
Bacterial strains and growth conditions. Strains of *Streptococcus mutans* were cultured in Brain
Heart Infusion (BHI) broth (Difco) or chemically defined medium (FMC) [50] supplemented with 0.5%
maltose. Streptococci were grown in a 5% CO_2 aerobic environment at 37°C, unless stated otherwise.
Strains of *Escherichia coli* were routinely grown in lysogeny broth (LB) with slight modifications (Lennox
LB; 10 g/L tryptone, 5 g/L yeast extract and 5 g/L NaCl) at 37°C with aeration. Antibiotics were added to
growth media at the following concentrations: kanamycin (1.0 mg/ml for *S. mutans*, 50 μg/ml for *E. coli*),
erythromycin (10 μg/ml for *S. mutans*, 300 μg/ml for *E. coli*), spectinomycin (1.0 mg/ml for *S. mutans*,
50 μg/ml for *E. coli*) and ampicillin (100 μg/ml for *E. coli*). Strains and plasmids are listed in S2 Table.
437

CRISPR-Cas bioinformatics. The 477 *S. mutans* genomes available in the National Center for
Biotechnology Information (NCBI) GenBank in June 2018 were mined using CRISPRone [51] to
determine the occurrence and diversity of CRISPR-Cas systems in this species. CRISPR arrays were
also identified using CRISPRone and these were used to predict the PAM sequence based on spacer-
protospacer matches using CRISPRTarget [52]. Afterwards, consensus PAM sequences for each strain
were built using the WebLogo server [53].
444

Construction of the xylose-inducible CRISPRi system. Xylose-inducible plasmids were cloned by
following a protocol developed by Xie *et al* [23]. Three plasmids were generated that contained *gfp*,
dcas9$_{Smu}$ and *dcas9*$_{Spy}$ inserts cloned into the pZX9 plasmid. For each plasmid, two linear fragments
were created by PCR using pZX9 or *S. mutans* UA159 genomic DNA as templates. *dcas9*$_{Spy}$ was
amplified from a plasmid obtained from Addgene (Addgene #44249). *dcas9*$_{Smu}$ was amplified from the
strain described in the supplemental material. The *gfp* gene was amplified from previously constructed
gfp-reporter strains [22]. The PCR fragments are designed so that they have overlapping

complementary regions that facilitate concatemer formation during a 2nd PCR. The products generated
by this 2nd PCR are transformed directly into *S. mutans*. Primers pZX9seqF and pZX9seqR were used
to amplify gene inserts followed by Sanger sequencing. For cloning of the sgRNA plasmid
(pPM::sgRNA), a synthetic DNA fragment was generated (Integrated DNA Technologies) that
contained the following components sequentially: 1) a P$_{veg}$ promoter [22], 2) a 20-nt sgRNA targeting
gfp, 3) a 42-nt dCas9 binding handle sequence, and 4) an *S. pyogenes* terminator. Restriction enzyme
recognition sequences for *Sac*I and *Sph*I were included at the 5' and 3' ends of the sgRNA construct
respectively. The synthetic DNA fragment was subsequently cloned into the pPMZ plasmid which
integrates at the *phnA-mtlA* locus of *S. mutans* UA159. This plasmid was also used to generate
plasmids containing sgRNAs that target other areas of the *S. mutans* genome. Protocols for cloning of
additional strains are provided in S1 File. Oligonucleotides used in this study are listed in S3 Table.
463
Green fluorescent protein CRISPRi assay. To test the CRISPRi system and compare the activity of
dCas9$_{Smu}$ to dCas9$_{Spy}$ we used a GFP based assay. CRISPRi strains were cultured overnight, grown to
an OD$_{600}$ = 0.5, and then diluted 1:100 into FMC-maltose with or without xylose. For these assays the
concentration of xylose was serially diluted. In triplicates, 200 µL samples were added to clear-
bottomed, black-sided, 96-well microtiter plates. After loading of the cultures, the 96-well plate was
placed in a Synergy HT microtiter plate reader (BioTek). The fluorescence and optical density
(absorbance at 600 nm) were measured at 30-min intervals using Gen5 software (BioTek). The
fluorescence settings were excitation at 485 nm, emission at 525 nm, and a sensitivity of 65. Data
readings were collected, and the background fluorescence or OD600 was subtracted prior to data
visualization using GraphPad Prism (v7) software (GraphPad Software). A slightly modified version of
this assay was used for PAM comparisons (S1 File). The protocol for Western blotting of GFP and
dCas9$_{Spy}$ is available in S1 File.
476
sgRNA library design and cloning. sgRNA sequences (20-nt) were selected with CRISPy-web [54].
With this tool we uploaded the *S. mutans* UA159 GenBank file (NC_004350.2) and selected essential
genes. For each gene, the tool designs sgRNAs with predicted off-target matches (13-bp upstream of
the PAM; off-targets containing 0, 1, 2, or 3-bp mismatches). To minimize off-target effects we only
selected sgRNAs with two or more mismatches, which we have shown to diminish sgRNA activity (Fig
2B). We also aimed to select sgRNA sequences as close to the start of each gene as possible. For
each sgRNA we predicted the secondary structure (20-nt sequence and 42-nt dCas9 binding handle)
with the RNAfold program provided by ViennaRNA [55]. Only sgRNAs that maintained the hairpin
structure of the dCas9 binding handle were selected for cloning. sgRNA sequences were cloned using
the Q5® Site-Directed Mutagenesis Kit (New England Biolabs). Primers were designed using the
NEBaseChanger™ tool (v.1.2.9), substituting the sgRNA-*gfp* 20-nt sequence for whichever sgRNA we
were cloning. Using the designed primers, the pPM::sgRNA plasmid was amplified and then added to a
mix of Kinase-Ligase-*Dpn*I for 5 min at room temperature. This mixture (5 µL) was then transformed
into chemically competent *E. coli* 10-beta with selection on LB-kanamycin. For each sgRNA plasmid
transformation, two independent clones were selected and cultured overnight. With each culture we
made a glycerol stock for storage at -80°C, and purified plasmid with the QIAprep Spin Miniprep Kit
(Qiagen). Each plasmid was Sanger sequenced using the pRCS1seqR primer (S3 Table).
494
sgRNA plasmid interference assays. sgRNA sequence specificity was tested using a plasmid
interference assay. A 20-nt sequence complementary to the template strand of the *atpB* gene
(SMu.1528) was introduced into the pPM::sgRNA plasmid with the Q5® Site-Directed Mutagenesis Kit.
Using the same kit, single, double, or triple base-pair mismatches were introduced into the 20-nt *atpB*
sgRNA sequence. These plasmids were transformed into competent *S. mutans* and cells were serially
diluted onto BHI agar, or BHI agar containing kanamycin. The percentage of cells that were
transformed was calculated from the colony forming units present on both types of agar.
502

CRISPRi growth assays. Strains of *S. mutans* carrying sgRNA cassettes were grown overnight in BHI with the appropriate antibiotics in a microaerophilic environment at 37°C. The following day the bacterial cultures were washed once in FMC-maltose, and then diluted 1:1000 into FMC-maltose or FMC-maltose containing 0.025% xylose. Triplicates of 300 µL were added to a Bioscreen C 10 x 10 well plate for each condition. OD_{600} was measured every 30 min for 16 h with a Bioscreen C Automated Microbiology Growth Curve Analysis System (Growth Curves USA).

Fluorescence microscopy of CRISPRi strains. Overnight cultures of CRISPRi strains were diluted 1:100 into 200 µL FMC-maltose containing 0.025% xylose, and then incubated at 37°C in a 5% CO_2 incubator for 16 h. After incubation, 1 µL of 1 mg/mL nile red (ThermoFisher Scientific) was added to each sample and incubated for 5 min at room temperature (RT) in darkness. Next, 1 µL of 1 mg/mL DAPI (4′,6-diamidino-2-phenylindole; ThermoFisher Scientific) was added to each sample and incubated at RT in darkness for a further 5 min. Samples were centrifuged for 5 min at 4,000 x *g* and suspended in ProLong™ Live Antifade Reagent (ThermoFisher Scientific). For imaging, 2.5 µL of stained cells were placed on PBS (pH 7.4; 137 mM NaCl, 2.7 mM KCl, 8 mM Na_2HPO_4, 2 mM KH_2PO_4) agarose pads [56] before sealing with a coverslip. Images were captured with a Zeiss Axiovert 200M microscope equipped with a Zeiss AxioCam MRm camera using a 100X oil objective.

Transmission electron microscopy. CRISPRi strains were grown as described for fluorescence microscopy. Afterwards cells were rinsed with 0.1M sodium cacodylate buffer and then fixed in 3% glutaraldehyde overnight at 4°C. The following day, cells were treated with 1.5% osmium tetroxide in the dark for 1 h at 4°C. Next the cells were mixed in equal parts with 5% agarose in PBS, collected by centrifugation at 2000 x *g*, and cooled to 4°C. Small chunks of the bacterial pellet plus agarose were then dehydrated in ethanol via the following steps: 30%, 50%, 70%, 80%, 90% each for 15 min, 99% 10 min, and then absolute ethanol 2 x 10 min. The dehydrated cells were embedded in an epoxy resin, sectioned and stained with uranyl acetate and lead citrate. Microscopy was conducted using a Hitachi H7600 transmission electron microscope.

Waxworm experiments. *Galleria mellonella* killing assays were performed as described previously [57]. Insects in the final instar larval stage were purchased from Vanderhorst, Inc. (St. Marys, OH), stored at 4°C in the dark and used within 7 days of shipment. Groups of 15 larvae, ranging in weight from 200 to 300 mg and with no signs of melanization, were selected at random for subsequent infection. Insects were injected with $1x10^7$ CFU of bacterial inoculum, prepared from overnight cultures that had been washed and resuspended in 0.9% saline solution, into the hemocoel of each larva via the last left proleg using a 25-µl Hamilton syringe. Fifteen larvae were injected for each condition. The number of viable cells injected into the larvae was verified by colony counts on BHI plates. As a negative control, one group of *G. mellonella* were injected with UA159 that was heat killed at 80°C for 30 minutes. After injection, insects were kept at 37°C and survival at different time intervals was recorded. Larvae were scored as dead when thigmotaxis was absent. Kaplan-Meier killing curves were plotted, and estimations of differences in survival were compared using a log rank test. All data were analyzed with GraphPad Prism, version 7.0c software. Experiments were repeated three times.

Data availability. The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgements

We thank Dr. Sharon Matthews and Chao Chen at the Electron Microscopy core at the University of Florida for help with TEM imaging and analysis; Dr. Vincent Richards for sending us the *S. mutans* genomes; Dr. Justin Merritt for providing us with the pZX plasmids for xylose-inducible gene expression in *S. mutans*; and Dr. Edward Chan and John Calise for use of, and technical assistance with,

fluorescence microscopy. This work was supported by National Institutes of Health grant R01DE013239 (to R.A.B.).

References

1. Krzyściak W, Pluskwa KK, Jurczak A, Kościelniak D. The pathogenicity of the Streptococcus genus. Eur J Clin Microbiol Infect Dis. 2013;32: 1361–1376. doi:10.1007/s10096-013-1914-9

2. Dewhirst FE, Chen T, Izard J, Paster BJ, Tanner ACR, Yu W-H, et al. The human oral microbiome. J Bacteriol. 2010;192: 5002–17. doi:10.1128/JB.00542-10

3. Pitts NB, Zero DT, Marsh PD, Ekstrand K, Weintraub JA, Ramos-Gomez F, et al. Dental caries. Nat Rev Dis Primer. 2017;3: 17030. doi:10.1038/nrdp.2017.30

4. Lemos JA, Quivey RG, Koo H, Abranches J. *Streptococcus mutans*: a new Gram-positive paradigm? Microbiology. 2013;159: 436–45. doi:10.1099/mic.0.066134-0

5. Nomura R, Nakano K, Nemoto H, Fujita K, Inagaki S, Takahashi T, et al. Isolation and characterization of *Streptococcus mutans* in heart valve and dental plaque specimens from a patient with infective endocarditis. J Med Microbiol. 2006;55: 1135–1140. doi:10.1099/jmm.0.46609-0

6. Nakano K, Hokamura K, Taniguchi N, Wada K, Kudo C, Nomura R, et al. The collagen-binding protein of *Streptococcus mutans*is involved in haemorrhagic stroke. Nat Commun. 2011;2: 485. doi:10.1038/ncomms1491

7. Lewis K. Platforms for antibiotic discovery. Nat Rev Drug Discov. 2013;12: 371–387. doi:10.1038/nrd3975

8. Juhas M, Eberl L, Church GM. Essential genes as antimicrobial targets and cornerstones of synthetic biology. Trends Biotechnol. 2012;30: 601–607. doi:10.1016/j.tibtech.2012.08.002

9. van Opijnen T, Bodi KL, Camilli A. Tn-seq: high-throughput parallel sequencing for fitness and genetic interaction studies in microorganisms. Nat Methods. 2009;6: 767–72. doi:10.1038/nmeth.1377

10. Shields RC, Zeng L, Culp DJ, Burne RA. Genomewide Identification of Essential Genes and Fitness Determinants of *Streptococcus mutans* UA159. mSphere. 2018;3: e00031-18. doi:10.1128/mSphere.00031-18

11. Fan F, Lunsford RD, Sylvester D, Fan J, Celesnik H, Iordanescu S, et al. Regulated Ectopic Expression and Allelic-Replacement Mutagenesis as a Method for Gene Essentiality Testing in *Staphylococcus aureus*. Plasmid. 2001;46: 71–75. doi:10.1006/plas.2001.1526

12. Bikard D, Jiang W, Samai P, Hochschild A, Zhang F, Marraffini LA. Programmable repression and activation of bacterial gene expression using an engineered CRISPR-Cas system. Nucleic Acids Res. 2013;41: 7429–7437. doi:10.1093/nar/gkt520

13. Qi LS, Larson MH, Gilbert LA, Doudna JA, Weissman JS, Arkin AP, et al. Repurposing CRISPR as an RNA-Guided Platform for Sequence-Specific Control of Gene Expression. Cell. 2013;152: 1173–1183. doi:10.1016/j.cell.2013.02.022

14. Liu X, Gallay C, Kjos M, Domenech A, Slager J, van Kessel SP, et al. High-throughput CRISPRi phenotyping
identifies new essential genes in *Streptococcus pneumoniae*. Mol Syst Biol. 2017;13: 931–931.
doi:10.15252/MSB.20167449

15. Peters JM, Colavin A, Shi H, Czarny TL, Larson MH, Wong S, et al. A Comprehensive, CRISPR-based Functional
Analysis of Essential Genes in Bacteria. Cell. 2016;165: 1493–1506. doi:10.1016/j.cell.2016.05.003

16. Serbanescu MA, Cordova M, Krastel K, Flick R, Beloglazova N, Latos A, et al. Role of the *Streptococcus*
mutans CRISPR-Cas Systems in Immunity and Cell Physiology. J Bacteriol. 2015;197: 749–761.
doi:10.1128/JB.02333-14

17. Fonfara I, Le Rhun A, Chylinski K, Makarova KS, Lécrivain A-L, Bzdrenga J, et al. Phylogeny of Cas9
determines functional exchangeability of dual-RNA and Cas9 among orthologous type II CRISPR-Cas
systems. Nucleic Acids Res. 2014;42: 2577–2590. doi:10.1093/nar/gkt1074

18. van der Ploeg JR. Analysis of CRISPR in *Streptococcus mutans* suggests frequent occurrence of acquired
immunity against infection by M102-like bacteriophages. Microbiol Read Engl. 2009;155: 1966–1976.
doi:10.1099/mic.0.027508-0

19. Cho S, Choe D, Lee E, Kim SC, Palsson B, Cho B-K. High-Level dCas9 Expression Induces Abnormal Cell
Morphology in *Escherichia coli*. ACS Synth Biol. 2018;7: 1085–1094. doi:10.1021/acssynbio.7b00462

20. Rock JM, Hopkins FF, Chavez A, Diallo M, Chase MR, Gerrick ER, et al. Programmable transcriptional
repression in mycobacteria using an orthogonal CRISPR interference platform. Nat Microbiol. 2017;2:
16274. doi:10.1038/nmicrobiol.2016.274

21. Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E. A Programmable Dual-RNA-Guided DNA
Endonuclease in Adaptive Bacterial Immunity. Science. 2012;337: 816–821. doi:10.1126/science.1225829

22. Shields RC, Kaspar JR, Lee K, Underhill S a. M, Burne RA. Fluorescence tools adapted for real-time monitoring
of the behaviors of *Streptococcus* species. Appl Environ Microbiol. 2019. doi:10.1128/AEM.00620-19

23. Xie Z, Qi F, Merritt J. Development of a tunable wide-range gene induction system useful for the study of
streptococcal toxin-antitoxin systems. Appl Environ Microbiol. 2013;79: 6375–84. doi:10.1128/AEM.02320-
13

24. Heler R, Samai P, Modell JW, Weiner C, Goldberg GW, Bikard D, et al. Cas9 specifies functional viral targets
during CRISPR–Cas adaptation. Nature. 2015;519: 199–202. doi:10.1038/nature14245

25. Leenay RT, Maksimchuk KR, Slotkowski RA, Agrawal RN, Gomaa AA, Briner AE, et al. Identifying and
Visualizing Functional PAM Diversity across CRISPR-Cas Systems. Mol Cell. 2016;62: 137–147.
doi:10.1016/j.molcel.2016.02.031

26. Zhang Y, Ge X, Yang F, Zhang L, Zheng J, Tan X, et al. Comparison of non-canonical PAMs for CRISPR/Cas9-
mediated DNA cleavage in human cells. Sci Rep. 2014;4. doi:10.1038/srep05405

27. Kleinstiver BP, Prew MS, Tsai SQ, Topkar V, Nguyen NT, Zheng Z, et al. Engineered CRISPR-Cas9 nucleases
with altered PAM specificities. Nature. 2015;523: 481–485. doi:10.1038/nature14592

28. Filipe SR, Tomasz A. Inhibition of the expression of penicillin resistance in *Streptococcus pneumoniae* by
inactivation of cell wall muropeptide branching genes. Proc Natl Acad Sci U S A. 2000;97: 4891–4896.
doi:10.1073/pnas.080067697

29. Massidda O, Nováková L, Vollmer W. From models to pathogens: how much have we learned about
Streptococcus pneumoniae cell division? Environ Microbiol. 2013;15: 3133–3157. doi:10.1111/1462-
2920.12189

30. Kovacs CJ, Faustoferri RC, Bischer AP, Quivey RG. *Streptococcus mutans* requires mature rhamnose-glucose
polysaccharides for proper pathophysiology, morphogenesis and cellular division. In: Molecular
Microbiology [Internet]. 12 Jul 2019 [cited 14 Aug 2019]. doi:10.1111/mmi.14330

31. Mistou M-Y, Sutcliffe IC, van Sorge NM. Bacterial glycobiology: rhamnose-containing cell wall
polysaccharides in Gram-positive bacteria. FEMS Microbiol Rev. 2016;40: 464–79.
doi:10.1093/femsre/fuw006

32. Beek SL van der, Breton YL, Ferenbach AT, Chapman RN, Aalten DMF van, Navratilova I, et al. GacA is
essential for Group A *Streptococcus* and defines a new class of monomeric dTDP-4-dehydrorhamnose
reductases (RmlD). Mol Microbiol. 2015;98: 946–962. doi:10.1111/mmi.13169

33. Ma Y, Pan F, McNeil M. Formation of dTDP-Rhamnose Is Essential for Growth of Mycobacteria. J Bacteriol.
2002;184: 3392–3395. doi:10.1128/JB.184.12.3392-3395.2002

34. Beek SL van der, Zorzoli A, Çanak E, Chapman RN, Lucas K, Meyer BH, et al. Streptococcal dTDP-L-rhamnose
biosynthesis enzymes: functional characterization and lead compound identification. Mol Microbiol.
2019;111: 951–964. doi:10.1111/mmi.14197

35. Quivey RG, Grayhack EJ, Faustoferri RC, Hubbard CJ, Baldeck JD, Wolf AS, et al. Functional profiling in
Streptococcus mutans: construction and examination of a genomic collection of gene deletion mutants.
Mol Oral Microbiol. 2015;30: 474–495. doi:10.1111/omi.12107

36. Kovacs CJ, Faustoferri RC, Quivey RG. RgpF Is Required for Maintenance of Stress Tolerance and Virulence in
Streptococcus mutans. J Bacteriol. 2017;199. doi:10.1128/JB.00497-17

37. De A, Liao S, Bitoun JP, Roth R, Beatty WL, Wu H, et al. Deficiency of RgpG Causes Major Defects in Cell
Division and Biofilm Formation, and Deficiency of LytR-CpsA-Psr Family Proteins Leads to Accumulation of
Cell Wall Antigens in Culture Medium by *Streptococcus mutans*. Appl Environ Microbiol. 2017;83.
doi:10.1128/AEM.00928-17

38. Boyle EA, Andreasson JOL, Chircus LM, Sternberg SH, Wu MJ, Guegler CK, et al. High-throughput biochemical
profiling reveals sequence determinants of dCas9 off-target binding and unbinding. Proc Natl Acad Sci.
2017;114: 5461–5466. doi:10.1073/pnas.1700557114

39. Anderson EM, Haupt A, Schiel JA, Chou E, Machado HB, Strezoska Ž, et al. Systematic analysis of CRISPR–
Cas9 mismatch tolerance reveals low levels of off-target activity. J Biotechnol. 2015;211: 56–65.
doi:10.1016/j.jbiotec.2015.06.427

40. Klein M, Eslami-Mossallam B, Arroyo DG, Depken M. Hybridization Kinetics Explains CRISPR-Cas Off-
Targeting Rules. Cell Rep. 2018;22: 1413–1423. doi:10.1016/j.celrep.2018.01.045

41. Chatterjee P, Jakimo N, Jacobson JM. Minimal PAM specificity of a highly similar SpCas9 ortholog. Sci Adv.
2018;4: eaau0766. doi:10.1126/sciadv.aau0766

42. Jakimo N, Chatterjee P, Nip L, Jacobson JM. A Cas9 with Complete PAM Recognition for Adenine
Dinucleotides. bioRxiv. 2018; 429654. doi:10.1101/429654

43. Edraki A, Mir A, Ibraheim R, Gainetdinov I, Yoon Y, Song C-Q, et al. A Compact, High-Accuracy Cas9 with a
Dinucleotide PAM for In Vivo Genome Editing. Mol Cell. 2019;73: 714-726.e4.
doi:10.1016/j.molcel.2018.12.003

44. Bailey J, Cass J, Gasper J, Ngo N-D, Wiggins P, Manoil C. Essential gene deletions producing gigantic bacteria.
PLOS Genet. 2019;15: e1008195. doi:10.1371/journal.pgen.1008195

45. Kawai Y, Mercier R, Mickiewicz K, Serafini A, Carvalho LPS de, Errington J. Crucial role for central carbon
metabolism in the bacterial L-form switch and killing by β-lactam antibiotics. Nat Microbiol. 2019; 1–11.
doi:10.1038/s41564-019-0497-3

46. Nakano K, Nomura R, Nakagawa I, Hamada S, Ooshima T. Demonstration of *Streptococcus mutans* with a
Cell Wall Polysaccharide Specific to a New Serotype, k, in the Human Oral Cavity. J Clin Microbiol. 2004;42:
198–202. doi:10.1128/JCM.42.1.198-202.2004

47. Coykendall AL. Classification and identification of the viridans streptococci. Clin Microbiol Rev. 1989;2: 315–
328. doi:10.1128/CMR.2.3.315

48. Beaussart A, Péchoux C, Trieu-Cuot P, Hols P, Mistou M-Y, F. Dufrêne Y. Molecular mapping of the cell wall
polysaccharides of the human pathogen *Streptococcus agalactiae*. Nanoscale. 2014;6: 14820–14827.
doi:10.1039/C4NR05280C

49. Caliot É, Dramsi S, Chapot-Chartier M-P, Courtin P, Kulakauskas S, Péchoux C, et al. Role of the Group B
antigen of *Streptococcus agalactiae*: a peptidoglycan-anchored polysaccharide involved in cell wall
biogenesis. PLoS Pathog. 2012;8: e1002756. doi:10.1371/journal.ppat.1002756

50. Terleckyj B, Willett NP, Shockman GD. Growth of several cariogenic strains of oral streptococci in a
chemically defined medium. Infect Immun. 1975;11: 649–655.

51. Zhang Q, Ye Y. Not all predicted CRISPR–Cas systems are equal: isolated cas genes and classes of CRISPR like
elements. BMC Bioinformatics. 2017;18: 92. doi:10.1186/s12859-017-1512-4

52. Biswas A, Gagnon JN, Brouns SJJ, Fineran PC, Brown CM. CRISPRTarget. RNA Biol. 2013;10: 817–827.
doi:10.4161/rna.24046

53. Crooks GE, Hon G, Chandonia J-M, Brenner SE. WebLogo: a sequence logo generator. Genome Res. 2004;14:
1188–90. doi:10.1101/gr.849004

54. Blin K, Pedersen LE, Weber T, Lee SY. CRISPy-web: An online resource to design sgRNAs for CRISPR
applications. Synth Syst Biotechnol. 2016;1: 118–121. doi:10.1016/j.synbio.2016.01.003

55. Lorenz R, Bernhart SH, Höner zu Siederdissen C, Tafer H, Flamm C, Stadler PF, et al. ViennaRNA Package 2.0.
Algorithms Mol Biol. 2011;6: 26. doi:10.1186/1748-7188-6-26

56. Jong IG de, Beilharz K, Kuipers OP, Veening J-W. Live Cell Imaging of *Bacillus subtilis* and *Streptococcus*
698   *pneumoniae* using Automated Time-lapse Microscopy. JoVE J Vis Exp. 2011; e3145. doi:10.3791/3145

699   57. Abranches J, Miller JH, Martinez AR, Simpson-Haidaris PJ, Burne RA, Lemos JA. The collagen-binding protein
700   Cnm is required for Streptococcus mutans adherence to and intracellular invasion of human coronary
701   artery endothelial cells. Infect Immun. 2011;79: 2277–2284. doi:10.1128/IAI.00767-10

702
703

704
Figure 1. Design of an inducible CRISPRi gene-silencing system for *S. mutans*. (A) Distribution of
CRISPR-Cas systems in *S. mutans* by type (top half) and subtype (bottom half). For example, 67 *S.*
mutans strains contain Type I systems (blue), and the subtypes of this system are B, C, and E (I-C is
the most common). The numbers on the exterior of the plot denote the number of *S. mutans* strains that
possess a certain CRISPR-Cas system. (B) Diagram of the Cas9 protein of *S. mutans* UA159 showing
domains. The two conserved amino acids required for nuclease activity in the *S. pyogenes* Cas9
protein that were mutated to create the *S. mutans* dCas9 protein are indicated. Each domain was
compared with *S. pyogenes* Cas9 for percentage amino acid identity. (C) Gene-silencing activity of
endogenous dCas9$_{Smu}$ was assessed by expressing an sgRNA with complementarity to a
chromosomally integrated *gfp* gene and comparing GFP protein levels using Western blotting (sgRNA-
gfp). Also shown is a negative control, d*cas9* without *gfp*, and a positive control strain that constitutively
expreses the *gfp* gene from the P23 promoter (see text for details). (D) Overview of the CRISPRi
system engineered for *S. mutans*, which consists of two primary elements, a chromosomally located
sgRNA and a plasmid-borne inducible dead Cas9. Combined, these two elements can block gene
transcription as shown on the right. (E) Comparison of dCas9$_{Smu}$ (left) and dCas9$_{Spy}$ (right) *gfp* gene
silencing activity using real-time fluorescence monitoring. Fluorescence readings are shown as dots,
and cell density (OD$_{600}$) readings as lines. As the xylose concentration was decreased, data points
change from a dark to a light blue color. (F) Comparison of dCas9$_{Smu}$ (top panel) and dCas9$_{Spy}$ (middle
panel) *gfp* gene silencing activity with Western blotting as as a function of xylose concentration. The
bottom panel is a Western blot against the xylose-inducible dCas9$_{Spy}$; as this antibody detected
dCas9$_{Spy}$, but not dCas9$_{Smu}$.
726
727

728
Figure 2. Determining guide RNA specificity of Cas9$_{Smu}$. (A) Diagram of the sgRNA for *atpB* and the
730 mutagenesis strategy. Positions 3, 6, 9, 12, 15 and 18 of the sgRNA were mutated (including in pairs or
731 triplets) as a way of investigating sgRNA specificity (see text for additional details). (B) Plasmid
732 interference assay of wild-type and mutated sgRNA-*atpB* plasmids, with Δ*cas9*, d*cas9* and sgRNAneg
733 serving as controls (for no plasmid interference). If the plasmid interfered with *S. mutans atpB*
734 expression, a lower transformation efficiency would be expected.
735
736

737
Figure 3. Investigating the proto-spacer adjacent motif (PAM) requirements of Cas9$_{Smu}$. (A)
Phylogenetic relatedness of extracted CRISPR-Cas9 spacer sequences (left) and Cas9 protein
sequences (right). Consensus PAM sequences were built from spacer sequences and are available in
Fig. S3. A key describing the colors chosen for consensus PAM sequences is shown on the far right.
(B) To determine the optimal PAM sequence for dCas9$_{Smu}$, different PAM sequences were inserted into
between the P23 promoter and *gfp* gene. (C) The functionality of various PAM sequences were
compared by measuring CRISPRi repression of GFP fluorescence for both *S. pyogenes* and *S. mutans*
Cas9. Repression was calculated as a percentage of maximal fluorescence without exogenous xylose
compared to fluorescence with xylose added. Bars show the average and standard deviation, and light
blue dots are the individual values, obtained using the dCas9$_{Smu}$ construct, whereas the dark blue dots
show data obtained using dCas9$_{Spy}$. Maximal repression was measured with a 5'-NGG-3' PAM, with
moderate repression for some 5'-NAG-3' PAMs.
750
751

752
Figure 4. High-throughput growth analysis of a comprehensive library of sgRNAs targeting
essential and growth supporting genes. (A) Overview of the CRISPRi growth study that included
generating sgRNAs against >250 essential or growth-supporting genes. Growth analysis of CRISPRi
strains (B) whether putative essential or required for growth and (C) sorted for physiological functions
by Clusters of Orthologous Groups. CRISPRi strains were cultured in the presence or absence of
xylose and the area under the curve (AUC) was calculated for each condition. Growth phenotypes were
categorized according to the ratio of AUC_{xylose} to $AUC_{uninduced}$ (the lower the number the stronger the
growth defect).

763
Figure 5. Microscopic analysis of the morphological defects caused by CRISPRi essential gene
depletion. (A) CRISPRi strains, by functional pathway, were grouped according to cell morphology
phenotypes. Strains with depleted essential genes can have more than one morphological defect (*e.g.*
increased cell chaining and heterogeneous sizes). (B) Representative micrographs for several of the
cell morphology phenotypes that were observed are shown. Additional microscopy across all the
pathways investigated can be found in Figs. S5-S12. For sgRNA-*atpA* the arrow shows an area of
increased nile red staining compared to other cells. The large round cells observed when silencing
peptidoglycan biosynthesis is depicted with the sgRNA-*murN* micrograph. Arrows indicate potential
anucleate cells when targeting the DNA replication-related gene *dnaC*.
773

774
Figure 6. Rhamnose-glucose polysaccharide biosynthesis genes are required for growth,
pathogenesis and correct cell division of *S. mutans*. (A) Distribution of Tn-seq reads within genes
that function in RGP biosynthesis (*rmlA-rmlC*, *rgpG*, and *rmlD-rgpF*; SMu.831-835 reside in the operon
that starts with *rmlD*). Each peak corresponds to a transposon insertion with a lack of peaks within a
gene representative of an essential gene (*e.g. rmlA* or *rgpE*). (B) Growth phenotypes of silenced RGP
biosynthesis genes compared to a control strain (sgRNA-*lacG*). (C) Pathogenesis phenotypes of
silenced RGP biosynthesis genes (*rmlB*, *rmlD*, *rgpC*, and *rgpF*) in a *Galleria mellonella* infection model.
Heat killed UA159 served as a negative control, and wild-type UA159 and d*cas9* absent sgRNA as
positive controls. (D) Electron micrographs of CRISPRi strains targeting the same genes investigated in
the pathogenesis model. Arrowheads point to putative division septa. Fluorescence microscopy images
and growth curves for CRISPRi strains for all of the genes involved in this pathway can be found in Fig.
S13. Cell length (E) and width (F) measurements from TEM micrographs using ImageJ.
787

Supplemental Materials and Methods

789
Strain cloning. Site-directed mutagenesis, using the splice overlap extension method (1), was
employed to mutate aspartic acid residue 10 to alanine and histidine residue 840 to alanine within *S.*
mutans cas9 (SMu.1405c). For each mutation, two PCR products were generated using primers that
contained the desired mutations, and primers which annealed 0.6 kbp upstream or downstream of the
region of interest. The two PCR products, with 25-bp homology that included the mutation, were
subjected to PCR for 1 cycle in the absence of added primers. Next, the outer primers were added and
a second PCR of 30 cycles was used to amplify the 1.2 kbp fragment with the desired mutation. Both
PCR products (for each mutation) were purified and transformed into competent *S. mutans*. Colonies
were selected on the basis of erythromycin resistance using a suicide plasmid carrying an internal
fragment of the *lacG* gene. Potential clones containing the desired mutations were screened using PCR
and DNA Sanger sequencing. Afterwards, strains that had been selected using the *lacG* plasmid were
grown in the presence of lactose, to identify strains that had lost the suicide plasmid and the sensitivity
of the strains to erythromycin was used as further confirmation of the plasmid. A $\Delta cas9$ (SMu.1405c)
strain was generated in a similar manner, except here the ATG start codon was mutated to a stop
codon (TAG; AT>TA). For marking *S. mutans* with GFP, a fragment of DNA was synthesized that
contained the P23 promoter and a synthetic RBS followed by a superfolder *gfp* sequence (2). This
fragment was subsequently cloned into the pBGE plasmid that integrates at the *gtfA* locus of *S. mutans*
(3). The restriction enzymes *Xba*I and *Bsr*GI were used to clone the DNA fragment into the pBGE
plasmid.

809
Green fluorescent protein PAM comparison assay. PAM sequences were introduced into a P23
promoter driving *gfp* gene expression (pBGE::P23-*gfp*) using the Q5® Site-Directed Mutagenesis Kit.
An sgRNA with complementarity adjacent to this PAM was introduced into the pPM::sgRNA plasmid
also using the Q5® Site-Directed Mutagenesis Kit. pBGE::P23-*gfp* plasmids (containing PAMs) were
introduced into an *S. mutans* strain with a $\Delta cas9$ mutation and carrying pPM::sgRNA-P23 and P$_{xyl}$-
d*cas9*$_{Smu}$. GFP fluorescence and microbial growth were measured with a Synergy HT, as detailed in the
Materials and Methods section.

817
Western blotting. Bacterial strains were grown in FMC-maltose (with or without xylose) at 37°C to an
OD$_{600}$ of 0.5. Cells were harvested by centrifugation at 3,500 × *g* for 10 min, spent medium was
discarded, and cell pellets were stored overnight at −20°C. On the following day, the pellets were
resuspended in 100 µl of lysis buffer (60 mM Tris, pH 6.8, 2% sodium dodecyl sulfate [SDS]) and
transferred to a screw-cap microcentrifuge tube that contained 100 µl 0.1-mm ice-cold glass beads.
Samples were homogenized in a bead beater for 30 s three times with 5-min intervals on ice. The
samples were then centrifuged at 8,000 × *g* for 10 min at 4°C. The supernates, which contained the *S.*
mutans proteins, were carefully removed and placed into a fresh 1.5-ml microcentrifuge tube. The
protein concentration was determined using a bicinchoninic acid (BCA) assay following the supplier's
protocol (Pierce), with purified bovine serum albumin used as the standard. Ten µg of each lysate was
diluted in 4X SDS loading buffer and the mixture was boiled for 10 min. Proteins were separated by
SDS-polyacrylamide gel electrophoresis (PAGE) and then transferred to a polyvinylidene difluoride
(PVDF) membrane using a Trans-Blot Turbo transfer system (Bio-Rad). Green fluorescent protein was
detected using a polyclonal anti-GFP antibody (1:5,000 dilution; Millipore Sigma) and a goat-anti-rabbit
immunoglobulin G (IgG) antibody (1:5,000 dilution; SeraCare Life Sciences, USA). dCas9$_{Spy}$ was
detected using a mouse monoclonal anti-CRISPR-Cas9 antibody (1:5,000 dilution; abcam #ab191468)
and goat-anti-mouse IgG antibody (1:5,000 dilution; SeraCare Life Sciences, USA). Western blot
signals were detected using a SuperSignalWest Pico chemiluminescent substrate kit (Thermo Fisher
Scientific) and visualized with a FluorChem 8900 imaging system (Alpha Innotech, USA).

837
Essential and growth-supporting genes of *S. mutans* UA159

Essential genes in *S. mutans* UA159 were identified with Tn-seq performed as described before (4, 5),
with slight modifications to growth conditions. Transposon mutants were selected on BHI agar, whereas
previous libraries were initially selected on blood agar. After libraries were created and stored in frozen
aliquots, they were grown in BHI or FMC for 7-10 generations; previous libraries were passaged for ~30
generations. The results of these experiments were used to determine which genes to target with
CRISPRi in parallel with previous Tn-seq in *S. mutans* UA159. In addition, for poorly annotated
essential or growth-supporting genes we ran the protein sequence through BLASTP.

846
847 **Figure S1. Determining the optimal conditions for xylose induced gene expression in *S. mutans*.**
848 The graphs show (glucose left, maltose right) that P_{xyl}-*gfp* expression was higher in FMC-maltose than
849 FMC-glucose for comparable amounts of xylose. Fluorescence readings are shown as dots, and cell
850 density readings as lines. As the xylose concentration is diluted data points change from a dark to a
851 light blue color.

852
Figure S2. Single-cell analysis of P$_{xyl}$-*gfp* expression and CRISPRi knockdown. Addition of xylose
to FMC-maltose induced *gfp* expression in virtually all imaged cells (first versus second panel) although
gfp expression was not as high as observed with a constitutive P$_{veg}$ promoter (third panel). In a
CRISPRi strain containing a chromosomally integrated *gfp* reporter all cells were GFP positive without
xylose addition (fourth panel). Virtually no cells expressed *gfp* when xylose was added to the CRISPRi
strain as median GFP fluorescence was barely above background (fifth panel).

859
860 **Figure S3. Cas9 PAM predictions for *Streptococcus mutans* strains.** Spacer sequences found
861 within Type II CRISPR cassettes of *S. mutans* strains were aligned to the genome of the *Streptococcus*
862 phage M102AD. After alignment protospacer adjacent motifs (PAMs) were extracted and consensus
863 PAMs were generated with WebLogo. Consensus PAMs broadly fit into five groups, 5'-NGG-3', T-rich,
864 A-rich, no distinct PAM and a fifth group of PAMs that didn't fit into the first four groups.

865
866 **Figure S4. Investigating the impact of sgRNA position within gene and overall gene length on**
867 **CRISPRi growth defects.** (A) Overall, there was no discernable impact on CRISPRi mediated growth
868 defects according to the placement of the sgRNA within the gene. (B) Gene length did not impact
869 CRISPRi-mediated growth defects across all measured strains (*e.g.* larger genes showing stronger
870 growth defects versus smaller genes).

871
Figure S5. Growth phenotypes and microscopy of CRISPRi strains with silenced genes related to carbohydrate metabolism. For growth assays, light blue dots indicate the no xylose condition and dark blue dots represent the addition of 0.1% xylose. Cells were stained with DAPI (DNA stain) and nile red (membrane stain) and imaged with a x100 oil objective. For each strain a representative microscopic image is shown.

877
878 **Figure S6. Growth phenotypes and microscopy of CRISPRi strains with silenced genes related**
879 **to peptidoglycan biosynthesis.** For growth assays, light blue dots indicate the no xylose condition
880 and dark blue dots represent the addition of 0.1% xylose. Cells were stained with DAPI (DNA stain) and
881 nile red (membrane stain) and imaged with a x100 oil objective. For each strain a representative
882 microscopic image is shown.

883
884 **Figure S7. Growth phenotypes and microscopy of CRISPRi strains with silenced genes related**
885 **to lipid metabolism.** For growth assays, light blue dots indicate the no xylose condition and dark blue
886 dots represent the addition of 0.1% xylose. Cells were stained with DAPI (DNA stain) and nile red
887 (membrane stain) and imaged with a x100 oil objective. For each strain a representative microscopic
888 image is shown.

889
890 **Figure S8. Growth phenotypes and microscopy of CRISPRi strains with silenced genes related**
891 **to DNA replication.** For growth assays, light blue dots indicate the no xylose condition and dark blue
892 dots represent the addition of 0.1% xylose. Cells were stained with DAPI (DNA stain) and nile red
893 (membrane stain) and imaged with a x100 oil objective. For each strain a representative microscopic
894 image is shown.

895
**Figure S9. Growth phenotypes and microscopy of CRISPRi strains with silenced genes related
to translation and ribosome biogenesis.** For growth assays, light blue dots indicate the no xylose
condition and dark blue dots represent the addition of 0.1% xylose. Cells were stained with DAPI (DNA
stain) and nile red (membrane stain) and imaged with a x100 oil objective. For each strain a
representative microscopic image is shown.

901
902 **Figure S10. Growth phenotypes and microscopy of CRISPRi strains with silenced genes related**
903 **to ATP synthase activity.** For growth assays, light blue dots indicate the no xylose condition and dark
904 blue dots represent the addition of 0.1% xylose. Cells were stained with DAPI (DNA stain) and nile red
905 (membrane stain) and imaged with a x100 oil objective. For each strain a representative microscopic
906 image is shown.

907
Figure S11. Growth phenotypes and microscopy of CRISPRi strains with silenced genes related
908
to transcription. For growth assays, light blue dots indicate the no xylose condition and dark blue dots
909
represent the addition of 0.1% xylose. Cells were stained with DAPI (DNA stain) and nile red
910
(membrane stain) and imaged with a x100 oil objective. For each strain a representative microscopic
911
image is shown.
912

913
914 **Figure S12. Growth phenotypes and microscopy of CRISPRi strains with silenced genes related**
915 **to cell division.** For growth assays, light blue dots indicate the no xylose condition and dark blue dots
916 represent the addition of 0.1% xylose. Cells were stained with DAPI (DNA stain) and nile red
917 (membrane stain) and imaged with a x100 oil objective. For each strain a representative microscopic
918 image is shown.

919
920 **Figure S13. Growth phenotypes and microscopy of CRISPRi strains with silenced genes related**
921 **to rhamnose-glucose polysaccharide biosynthesis and transport.** For growth assays, light blue
922 dots indicate the no xylose condition and dark blue dots represent the addition of 0.1% xylose. Cells
923 were stained with DAPI (DNA stain) and nile red (membrane stain) and imaged with a x100 oil
924 objective. For each strain a representative microscopic image is shown.
925

926
927 **Figure S14. Multiple nucleoids per cell for RGP biosynthesis knockdowns.** Cell borders are shown
928 with dotted orange lines. Inside outlined cells multiple nucleoids are visible as punctate bright dots, also
929 highlighted with white arrowheads.

Supplemental Text References

930
1. Ho SN, Hunt HD, Horton RM, Pullen JK, Pease LR. 1989. Site-directed mutagenesis by overlap
extension using the polymerase chain reaction. Gene 77:51–59.
2. Shields RC, Kaspar JR, Lee K, Underhill S a. M, Burne RA. 2019. Fluorescence tools adapted for
real-time monitoring of the behaviors of Streptococcus species. Appl Environ Microbiol.
3. Zeng L, Burne RA. 2009. Transcriptional Regulation of the Cellobiose Operon of Streptococcus
mutans. J Bacteriol 191:2153–2162.
4. Shields RC, Zeng L, Culp DJ, Burne RA. 2018. Genomewide Identification of Essential Genes and
Fitness Determinants of *Streptococcus mutans* UA159. mSphere 3:e00031-18.
5. Shields RC, O'Brien G, Maricic N, Kesterson A, Grace M, Hagen SJ, Burne RA. 2018. Genome-
wide screens reveal new gene products that influence genetic competence in *Streptococcus*
mutans. Journal of Bacteriology 200:16 e00508-17.
942
943