First fossil skull of an anteater (Vermilingua, Myrmecophagidae) from northern South America, a taxonomic reassessment of *Neotamandua* and a discussion of the myrmecophagid diversification

Kevin Jiménez-Lara[a,b*], Jhon González

[a] División de Paleontología de Vertebrados, Museo de La Plata, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque s/n, B1900FWA La Plata, Argentina.

[b] CONICET, Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina.

*Corresponding author. E-mail: kjimenezlara@fcnym.unlp.edu.ar

The evolutionary history of the South American anteaters, Vermilingua, is incompletely known as consequence of the fragmentary and geographically biased nature of the fossil record of this group. The only record of a nominal extinct species for northern South America is *Neotamandua borealis*, from the Middle Miocene of La Venta area, southwestern Colombia (Hirschfeld 1976). A new genus and species of myrmecophagid for La Venta is described here from a new partial skull. Additionally, the taxonomic status of the genus to which was referred the co-occurrent species of Gen. et sp. nov., i.e. *Neotamandua*, is revised. The morphological and taxonomic analyses of these taxa indicate that Gen. et sp. nov. may be related to *Tamandua* and that the justification of the generic assignments of the referred species to *Neotamandua* is weak, with high probability of

reassigning some of them to a new genus. Two species previously referred to *Neotamandua*

(*N. magna* and *N.? australis*) were designated as *species inquirendae* and new diagnostic

information for the redefined genus and its type species, *N. conspicua*, is provided.

Together, these results suggest that the diversification of Myrmecophagidae was

taxonomically and biogeographically more complex than that what has been proposed so

far. Considering the new evidence, it is proposed a synthetic model on the diversification of

these xenartrans during the late Cenozoic based on the probable relationship between their

intrinsic ecological constraints and some major abiotic changes in the Americas.

Key words: Vermilingua, Myrmecophagidae, diversification, *Neotamandua*, La Venta

area, Neogene.

Introduction

The anteaters of the suborder Vermilingua are part of Xenarthra, one of the more inclusive

clades in the evolutionary tree of the placental mammals (Eutheria) and a characteristic

group in the land mammal assemblages of the middle-late Cenozoic of the Americas

(McDonald et al. 2008; Foley et al. 2016; Halliday et al. 2016; Feijoo & Parada 2017).

Within Xenarthra, Vermilingua belongs to Pilosa, a clade that also includes the sloths, i.e.

Tardigrada. Today, Vemilingua comprises the genera *Cyclopes* (pygmy anteaters),

Tamandua (collared anteaters) and *Myrmecophaga* (giant anteaters). These genera groups

ten extant species, the most of them (seven) belonging to *Cyclopes*, according to the most

recent exhaustive taxonomic revision (Miranda et al. 2017). The classic phylogenetic

hypothesis reunites *Tamandua* and *Myrmecophaga* in the family Myrmecophagidae, while

Cyclopes is located in a basal position with respect to Myrmecophagidae as the only recent

form of the family Cyclopedidae (Engelmann 1985). With the connotation of a superior

taxonomic hierarchy (i.e. at the family level; Barros et al. 2008; Gibb et al. 2015) by an

early evolutionary divergence (Hirschfeld 1976; Delsuc et al. 2001; Gibb et al. 2016) and in

acknowledgment of a more extended use in the scientific literature, the names

Myrmecophagidae and Cyclopedidae are used here, instead of Myrmecophaginae and

Cyclopinae *sensu* Gaudin & Branham (1998), respectively. However, the taxonomic

content of Myrmecophaginae and Cyclopinae, including extinct forms, is considered as

transferable to their counterparties (McDonald et al. 2008).

The living anteaters, whose mean body mass ranges from ~0.4 and 30 kilograms (Gaudin et

al. 2018), are highly, morphologically specialized mammals by exhibiting remarkable

skeleton and soft-anatomy modifications, which are closely linked to their

myrmecophagous diets, i.e. diets consisting of at least 90% of ants/termites (Redford 1987;

McDonald et al. 2008). Many of these adaptations, anatomically located in the skull and

jaws, are associated between them in several ways by being part of the architecture of an

integrated functional system of food apprehension and ingestion. Among these features, the

following are some of the most noteworthy: rostral elongation and narrowing, basicranial-

basifacial axis curvature, complete loss of teeth, gracile jaw, reduction of the adductor jaw

muscles, unfused jaw symphysis and protrusible long tongue (Reiss 2001; Gaudin &

McDonald 2008; McDonald et al. 2008). Several of these morphological specializations are

convergent with those described for other myrmecophagous mammals such as the

pangolins (Pholidota) and the aardvarks (Tubulidentata), so it is not surprising that early

systematic researchers erroneously proposed close common ancestry of Vermilingua with

these Old world groups from their superficial similarities (e.g. Engelmann 1978; Norman &

Ashley 1994).

Despite their unique biology and ecology, at least in the context of the land mammals of the

Americas, the evolutionary history of the anteaters is largely obscured by their poor,

fragmentary and geographically biased fossil record (Hirschfeld 1976; Gaudin & Branham

1998; McDonald et al. 2008). Generally, five valid genera and nine species are recognized

in the fossil record of Vermilingua, of which two genera and two species have extant

representatives, i.e. *Myrmecophaga tridactyla* and *Tamandua tetradactyla*.

Myrmecophagidae groups nearly all these fossil taxa (only one genus and one species for

Cyclopedidae) in a general biochron beginning c. 18 million years before present, most of

them distributed throughout the Neogene (McDonald et al. 2008). But while the record of

this family for the latter period is taxonomically more diverse than that for the Quaternary,

it also poses more difficulties in the systematic framework of the implicated taxa. The

oldest member of Myrmecophagidae is *Protamandua rothi*, from the late Early Miocene of

the Santa Cruz Province, southern Argentina (Ameghino 1904). This species has been well

validated from a pair of incomplete skulls and several postcranial bones, but the validity of

other co-occurrent putative vermilinguan (myrmecophagid?) taxa is, at least, questionable

(Hirschfeld 1976; McDonald et al. 2008). For the early Middle Miocene has been reported

a myrmecophagid doubtfully assigned to *Neotamandua*, and yet used to create a new

species from isolated humeral remains (*N.? australis*; Scillato-Yané & Carlini 1998). In the

latter genus was also allocated, with some uncertainty, postcranial material of a middle-to-

large sized anteater recorded in the late Middle Miocene of La Venta area, southwestern

Colombia, whose description includes the only nominal extinct species for northern South

America, i.e. *N. borealis* (Hirschfeld 1976). *Neotamandua* chronologically extends to the

Late Miocene and Early Pliocene with the species *N. magna* (Ameghino 1919), *N.*

greslebini (Kraglievich 1940) and *N. conspicua* (type species; Rovereto 1914), all of them

from northwestern Argentina (provinces of Catamarca and Tucumán). This genus is

typically recognized as morphologically similar (even directly ancestral) to *Myrmecophaga*,

although smaller in body size (Hirschfeld 1976; Gaudin & Branham 1998). Considering the

very few anatomically correlatable elements in which are based the different species

referred to *Neotamandua*, Hirschfeld (1976) and Scillato-Yané & Carlini (1998) have

suggested that this genus could be paraphyletic. Furthermore, the latter authors proposed

the hypothesis that *Neotamandua* is composed by two distinct evolutionary lineages: one

more closely related to *Myrmecophaga* and other one to *Tamandua*. In turn, these two

lineages would have diverged in allopatry in South America, in such a way that the

geographical origin of *Myrmecophaga* is located in northern South America, while that of

Tamandua is in southern South America.

In this article, we describe the first fossil skull of a myrmecophagid (and vermilinguan)

from northern South America. This specimen was collected in the Middle Miocene La

Victoria Formation of La Venta area, Colombia. Additionally, in coming to analysis

because it is the only nominal taxon reported for the same region and geological unit, the

taxonomic status of *Neotamandua* is revised. The results prompt the development of a

discussion on a model of diversification for Myrmecophagidae in which new and previous

hypothesis about this evolutionary event are synthesized. This contribution is intended to

revaluate, expand and integrate biotic and abiotic evidence related to the diversification of

this fascinating mammal group, with emphasis on the biogeographic role of tropical, low

latitude regions of the Americas.

Material and methods

The cranial specimen described for the first time here for Colombia (VPPLT 975) comes

from a light-brown mudstone layer in the Llano Largo field, around 2 Km northeast of La

Victoria town, Municipality of Villavieja, Department of Huila (Fig. 1A-C). Strata of the

La Victoria Formation outcrop there, within the palaeontologically relevant area of La

Venta. The La Victoria Formation is a geological unit of ~500 meters in thickness which is

mainly composed by bioturbated mudstones (Anderson et al. 2016). These sedimentites are

interrupted by very continuous, coarse-to-fine grained sandstones with crossbedding and

erosive bases. According to the lithostratigraphic scheme of Guerrero (1997; Fig. 1D), the

new skull is from a level stratigraphically close (<20 m) and below the Chunchullo

sandstone beds, i.e. the lower part of the La Victoria Formation. This corresponds to the

unit referred as "Unit between the Cerro Gordo and Chunchullo sandstone beds". As

described by the same author, this unit, whose thickness ranges from ~80 to 160 m, is

predominantly composed of mudstones and some interlayers of sandstones. This

sedimentary body bears abundant plutonic and volcanic fragments from the lower Jurassic

basement of the Honda Group (Saldaña Formation), as well as clasts of volcanic rocks

formed in the magmatic arc of the Cordillera Central of Colombia during the Middle

Miocene (Anderson et al. 2016).

The general palaeoenvironment inferred for the La Victoria Formation is a meandering

fluvial system (except for the Cerbatana conglomerate, associated to an anastomosed

system) with significant soil development in flood plain zones (Guerrero 1997). The ages

calculated by Guerrero (1997) and Flynn et al. (1997) using magnetic polarity stratigraphy

and geochronology indicate sedimentary deposition during the interval 13.8□12.5 million

years ago (mya). These results have recently been reinforced by the U-Pb geochronology of

detrital zircons recovered in this formation (Anderson et al. 2016). The age range obtained

is 14.4 ± 1.9 □ 13.2 ± 1.3 mya. This interval coincides approximately with the early

Serravalian, sub-stage of the Middle Miocene [Insert Fig. 1 here].

Cranial measurements, with some nomenclatural modifications, are based on those of

Hossotani et al. (2017) (Fig. 2; see Anatomical Abbreviations). All these measurements are

in millimetres (mm). The description of the new skull of La Venta includes a rough body

mass estimation of the respective individual from a traditional allometric approach. All

these data and analyses are compiled in the Supplementary material (Appendices S1 and

S3). For the taxonomic analysis of the genus *Neotamandua* were revised the justifications

of generic allocations for the referred species (at least doubtfully) in all the relevant

scientific literature. These species are: *Neotamandua conspicua* Rovereto 1914 (type

species); *Neotamandua magna* Ameghino 1919; *Neotamandua greslebini* Kraglievich

1940; *Neotamandua borealis* Hirschfeld 1976; *Neotamandua*? *australis* Scillato-Yané &

Carlini 1998. Additionally, some observations were made on the holotypes of *N. conspicua*

(MACN 8097) and *N. borealis* (UCMP 39847) to reexamine the described characteristics

for these species in the original publications (Rovereto 1914 and Hirschfeld 1976,

respectively). The conceptual model of Plotcnick & Warner (2006) to recognize taxonomic

wastebaskets was applied to *Neotamandua*. From the foregoing and the designation of the

specimen FMNH P14419 as epitype of *N. conspicua*, it was proposed a diagnosis for

Neotamandua. See a list of all the studied fossil specimens in the Appendix S1 of the

Supplementary material [Insert Fig. 2 here].

On other hand, considering that Hirschfeld (1976), in her description of *N. borealis*, did not

include morphological comparisons from postcranial bones of this species and homologous

elements of the species referred to *Neotamandua* for southern South America, we

performed this task and a preliminary character distribution analysis from postcrania of

these taxa to explore the hypothesis that they are closely related. Forcibly, *N. magna* and

N.? australis are excluded from the comparisons since they do not have osteological

elements correlated with those of *N. borealis* (or any other species referred to

Neotamandua). Additionally, as a result of loss of its holotype (McDonald et al. 2008),

comparisons with *N. greslebini* are based exclusively on the non-illustrated description by

Kraglievich (1940). Other comparisons include postcranium collected by Juan Méndez in

1911 in the upper Miocene of the Andalhuala locality, Catamarca Province, Argentina. This

material was assigned to *Neotamandua* (*Neotamandua* sp.) without a reference publication.

McDonald et al. (2008) manifested doubt about this taxonomic assignment

(*Neotamandua*?), but these authors simultaneously speculated that it might be the lost

holotype of *N. greslebini*.

Following to McKenna & Bell (1997), the genus *Nunezia* is considered a junior synonym

of *Myrmecophaga*. Myological inferences are based on Hirschfeld (1976) and Gambaryan

et al. (2009).

Institutional abbreviations

CAC: Cátedra de Anatomía Comparada, Facultad de Ciencias Naturales y Museo,

Universidad Nacional de La Plata; **FMNH:** Field Museum, Chicago, IL., USA; **ICN:**

Instituto de Ciencias Naturales, Facultad de Ciencias, Universidad Nacional, Bogotá,

Colombia; **MACN:** Museo Argentino de Ciencias Naturales 'Bernardino Rivadavia',

Buenos Aires, Argentina; **MLP:** Museo de La Plata, Facultad de Ciencias Naturales y

Museo, Universidad Nacional de La Plata, La Plata, Argentina; **MPT:** Museo Provincial de

Tucumán, Tucumán, Argentina; **UCMP:** University of California Museum of

Paleontology, Berkeley, CA., USA; **VPPLT:** Museo de Historia Natural La Tatacoa, La

Victoria, Huila, Colombia; **YPM:** Peabody Museum, Yale University, New Haven, CT,

USA.

Anatomical abbreviations

Abbreviations of equivalent measurements by Hossotani et al. (2017) in parenthesis. **FL**,

frontal length; **GSL (SL)**, greatest skull length; **MBW (NC)**, maximum braincase width;

ML, maxilla length; **NL**, nasal length; **NW (NB)**, nasal width; **PL**, parietal length.

SYSTEMATIC PALAEONTOLOGY

Xenarthra Cope, 1889

Pilosa Flower, 1883

Vermilingua Illiger, 1811

 Myrmecophagidae Gray, 1825

 Gen. nov.

 Etymology. [intentionally in blank].

 Diagnosis. Same as that of the type and only known species.

 Gen. et sp. nov.

 (Fig. 3)

 Etymology. [intentionally in blank]

 Holotype. VPPLT 975, anterior portion of a skull, without jugals nor premaxillae.

 Diagnosis. Middle sized myrmecophagid, slightly smaller than *Tamandua* and even more

 than *Neotamandua*. It can be differentiated from other genera/species of anteaters by the

 following combination of cranial features: relatively width rostrum, similar to *Tamandua*;

 narrow and strongly tapered nasals toward their anterior end; anteroposterior length of the

 pre-orbital section of frontals equal to more than two thirds of the anteroposterior length of

 nasals; jugals inserted from the same level of the most anterior border of the lacrimal;

 anterior portion of the orbit more laterally extended in the superior wall in the inferior one,

 without forming a conspicuous dome as in *Neotamandua conspicua*.

 Comparative description of the holotype of Gen. et sp. nov. The specimen VPPLT 975

 consists of a partial skull that preserves nearly all the anatomical elements from the anterior

 section of the frontals (at the mid anteroposterior level of the orbit) to the anterior end of

 the rostrum. See cranial measurements taken for this new taxon and other myrmecophagids

in the Table 1. The estimated body mass for this individual is around 3.9 Kg (Appendix S3

of the Supplementary material). As consequence of the preservation, some sutures are

distinguishable in dorsal and lateral views, but virtually no suture is clearly detectable in

ventral view. The rostrum is proportionally shorter and more robust than those in

Myrmecophaga and *N. conspicua* (see below), but less than in *Tamandua*. In dorsal view, it

is very similar to the skull of *Tamandua*, with at least four characteristics remarkably

different with respect this extant genus: (1) lower rostrum; (2) rostrum more regularly

tapered; (3) narrower and more anteriorly tapered nasals; (4) pre-orbital section of the

frontals more anteroposteriorly elongated. In dorsal view, the rostrum shows a slight bulge

in its middle part, similar to that in *Tamandua* and *Myrmecophaga*. However, in VPPLT

975 this bulge is even subtler than in the living myrmecophagids. Apparently, the nasals are

shorter than frontals and are poorly exposed in lateral view. The jugals are absent by

preservation, but it is possible to recognize their insertion location. This is more anterior

than in *Myrmecophaga*, but more posterior than in *Tamandua*. Associated to the insertion

of the jugal, there is a reduced posterolateral process of the maxilla in comparison with that

of *Myrmecophaga*, similar in *Tamandua*. The right side of the skull preserves better the

lacrimal zone, but it is simultaneously more deformed around the fronto-maxillary suture

than in the left side. The lacrimal is longer in its anteroposterior axis than in that

dorsoventral. The same bone is proportionally smaller than in *Tamandua* and even more

than in *N. conspicua*. It has a triangular outline (at least anteriorly), similar to

Myrmecophaga and unlike *Tamandua* (irregularly rounded, ovated, or, infrequently, sub-

triangular lacrimal). The maxilla is not part of the orbit. The superior wall of the orbit is

more laterally expanded than the inferior wall, without forming a conspicuous dome as in

N. conspicua. This is similar to the condition observed in *Myrmecophaga* and differs from

that in *Tamandua*, in which the inferior wall is prominent given that it is more laterally

expanded. It is not possible to recognize lacrimal nor orbital foramina. In ventral view, the

dorsal border of the orbit is regularly concave. The palatines are less laterally extended than

in *Tamandua* and apparently there are no palatine "wings" (noticeable lateral expansions of

the palatines), unlike *N. conspicua* [Insert Fig. 3 here].

Taxonomic reanalysis of the genus *Neotamandua*

Taxonomic history and discussion on the taxonomic status of *Neotamandua*

The genus *Neotamandua* was proposed by Rovereto (1914) from a posterior portion of a

skull (MACN 8097), which was collected in upper Miocene-to-Pliocene strata of the

Catamarca Province, Argentina. The name *Neotamandua*, literally meaning 'new

tamandua', was coined by Rovereto in allusion to the cranial similarity of the type species,

N. conspicua, with the extant genus *Tamandua*, rather than with *Myrmecophaga*. This

detail would be historically paradoxical, as will be shown below. It is important to note that

Rovereto did not provide a diagnosis for *Neotamandua*, but he just briefly described the

holotype of *N. conspicua*, emphasizing its elongated parietals. However, this feature, more

comparable with that in *Myrmecophaga* than that in any other myrmecophagid, was

correlated with the anteroposterior length of the parietals in *Tamandua*. A few years after

the Rovereto's work, Carlos Ameghino (Ameghino 1919) used a pelvis (MPT 58)

recovered in contemporary strata of the Tucuman Province, Argentina, to create a new

species, *N. magna*. Despite the taxonomic assignment of this pelvis to *Neotamandua*,

Ameghino discussed that, alternatively, this species could belong to other genus of larger

body size, as Kraglievich (1940) also held. Formally, *N. magna* has not been reevaluated,

but McDonald et al. (2008) suggested that, given that this species was transferred to

Nunezia by Kraglievich (1934), and *Nunezia* is considered a junior synonym of

Myrmecophaga (Hirschfeld 1976; McKenna & Bell 1997), then *N. magna* should be

included in the latter genus, i.e. *Myrmecophaga magna* comb. nov. (unpublished). Indeed,

the morphological differences cited by Ameghino (1919) and Kraglievich (1940) between

the pelvis of *N. magna* and that of *M. tridactyla* (e.g. greater width and ventral flattening of

the intermediate sacral vertebrae) do not seem sufficient to consider a generic distinction

between these species.

Two decades later, Kraglievich (1940) proposed a new species from postcranium collected

in the upper Miocene of the Catamarca Province. This was initially assigned to *N.*

conspicua. According to Kraglievich, the then new species, *N. greslebini*, is easily

identifiable by its large size, intermediate between those of *N. conspicua* and *N. magna*.

Like Rovereto (1914), this author also correlated his generic assignment of *N. greslebini* to

Neotamandua from the similarity between the fossil specimens of this species and

homologous elements of *Tamandua* (Kraglievich, p. 633). The holotype of *N. greslebini* is

missing or mixed up with material labelled with generic names of extinct anteaters (i.e.

Neotamandua and *Palaeomyrmidon*) in the Museo Argentino de Ciencias Naturales

(MACN), in Buenos Aires, Argentina (McDonald et al. 2008).

Already in the second half of the XX century, a controversy about the possible synonymy

between *Neotamandua* and *Myrmecophaga* arose. This means that there was a radical

paradigmatic shift in myrmecophagid systematics, from that in early XX century, in which

Neotamandua was considered closely related to *Tamandua*, to that in late XX century, in

which *Neotamandua* was even a serious candidate to be a junior synonym of

Myrmecophaga. This historical change began with the non-cladistic systematic analysis of

Hirschfeld (1976), in which *Neotamandua* was originally proposed as the direct ancestor

(anagenetic form) of *Myrmecophaga*. In the same work, Hirschfeld created the first and,

until now, only nominal extinct species of Vermilingua and Myrmecophagidae for northern

South America, *N. borealis* (Middle Miocene of Colombia). Given the scarcity and

fragmentation of the specimens referred to *Neotamandua*, Hirschfeld recognized the need

to revise the taxonomic validity of *N. conspicua*, *N. magna* and *N. greslebini*. Indeed, she

went beyond and stated that *Neotamandua* species could be representatives of more than

one single genus. However, her assignment of *N. borealis* to *Neotamandua* was based

primarily on the idea that the fossils she studied are 'considerably more advanced than

those known from the Santacruzian [late Early Miocene], closer to the Araucanian [Late

Miocene-Pliocene] species and…to the line leading to *Myrmecophaga* than *Tamandua*'

(Hirschfeld, p. 421). For this author, many postcranial traits of *N. borealis* are intermediate

between *Tamandua* and *Myrmecophaga*. As a questionable methodological aspect, it is

important to note that Hirschfeld did not make osteological comparisons with the southern

species of *Neotamandua*, only with postcranium of *Protamandua*, *Tamandua* and

Myrmecophaga (extant species of the two latter genera).

In implicit reply to Hirschfeld (1976), Patterson et al. (1992) highlighted the morphological

similarities between the unpublished skull FMNH P14419, catalogued as *N. conspicua* in

the Field Museum, and the modern skulls of *Myrmecophaga*. For these authors, FMNH

P14419 only differs from skulls of the living giant anteater in its smaller size.

Consequently, Patterson et al. (1992) suggested synonymize *Neotamandua* and

Myrmecophaga, with nomenclatural priority for the latter. Nevertheless, Gaudin &

Branham (1998) provided (weak) support for the validity of *Neotamandua* through a

comprehensive phylogenetic analysis of Vermilingua. Their results indicate that

Neotamandua is an independent taxon based on two autapomorphies, being one of them

ambiguous and the other one unambiguous. The latter is the horizontal inclination of the

glenoid. In the only most parsimonious tree recovered by Gaudin & Branham (1998),

Neotamandua is closely related to *Myrmecophaga*, not *Tamandua*, as opposed to Rovereto

(1914) and Kraglievich (1940).

Finally, the last species referred, with doubt, to the genus was *N.? australis* (Scillato-Yané

& Carlini 1998). The holotype of this species consists only of a humerus (MLP 91-IX-6-5)

collected in the lower Middle Miocene of the Río Negro Province, Argentina. Scillato-Yané

& Carlini (1998) highlighted some similarities of this material with the humerus of

Tamandua. They also expressed considerably uncertainty in assigning it to *Neotamandua*,

not only by its fragmentary nature, but from the idea of Hirschfeld about the non-natural

(i.e. non-monophyletic) status of this genus. Without performing a phylogenetic analysis,

these authors proposed a hypothesis that *N. borealis* is closely related to *Myrmecophaga*,

while *N. conspicua* and *N.? australis* are closer to *Tamandua*. If this hypothesis is correct,

N. borealis does not belong to *Neotamandua* as consequence of the application of the

nomenclatural priority principle.

In summary, multiple historical factors, including the lack of a diagnosis, insufficient

number of anatomically correlatable/highly diagnostic postcranial elements and, especially,

absence of cranial-postcranial associations, aroused the relatively arbitrary use of

Neotamandua as a wastebasket taxon, i.e. a residual genus deriving from weak and/or

inadequate systematic analysis. According to the conceptual model of Plotnick & Warner

(2006), *Neotamandua* has five (from a total of seven) properties of a genus potentially

classifiable as wastebasket: (1) it is an old name (i.e. more than one century to the present);

(2) it is [relatively] rich in species (five species, i.e. the most speciose extinct genus of

Vermilingua); (3) it has a [relatively] high number of occurrences; (4) it has wide temporal

and geographical distributions; (5) it [primarily] groups together specimens poorly

preserved and/or difficult to identify. To these five properties we may add the lack of a

diagnosis, which is related in some way to the property number two of the Plotnick-Warner

model, i.e. genera diagnosed from generalized characters, probably plesiomophies or easily

recognizable characters.

As it was shown, *Neotamandua* has been invoked as a directly ancestral form of

Tamandua, or, more recently, of *Myrmecophaga*, from its morphological characteristics in

common with these two extant genera. But precisely because of this character mosaic, the

generic allocation of isolated postcranial remains of myrmecophagids potentially referable

to *Neotamandua* should not be reduced or exclusively focused on their comparison with the

crown-group, but should also consider the effect of possible homoplasies (e.g. those related

to ecological convergences), plesiomorphies and limitations of the fossil record (Plotnick &

Warner 2006). In other words, the apparent affinity between isolated postcranial elements

of any Neogene anteater and their homologous in *Myrmecophaga* is not enough to make a

reliable generic allocation in *Neotamandua*; diagnostic information of the latter genus is

needed, preferably autapomorphies, which allow it to be individually identified and not

simply as a set of forms similar to *Myrmecophaga*.

359   **Comparisons between northern and southern species referred to *Neotamandua***

360   ***N. borealis* and *N. greslebini.*** See the Table 2 for comparison of postcranial measurements

361   between these species.

362   **Radius.** Both *N. borealis* and *N. greslebini* show two longitudinal, parallel radial

363   ridges, of which the lateral ridge is higher and reaches a more distal level than the

364   cranial one. This is similar to the condition observed in *Myrmecophaga* and differs

365   from the distally convergent radial ridges of *Tamandua*. In *N. borealis*, the lateral

366   ridge is even more distally extended than in *N. greslebini,* in such a way that the

367   flanks of this structure contact the lateral border of the styloid process. According to

368   Kraglievich (1940), in the Argentinean species this ridge ends at an intermediate level

369   between the distal end of the cranial ridge and the styloid process.

370   **Tibia.** The type material of *N. borealis* includes a proximal epiphysis and part of the

371   diaphysis of a right tibia. According to Kraglievich (1940), the holotype of *N.*

372   *greslebini* includes two fragments of a tibia, one of them proximal and the other one

373   distal. Both Hirschfeld and Kraglievich claimed greater overall similarity between the

374   tibial fragments of these species and the homologous parts of *Tamandua*, rather than

375   *Myrmecophaga*. This way, the mid-section of the tibias both of *N. borealis* and *N.*

376   *greslebini* is not as strongly triangular as in *Myrmecophaga*. Rather, this bone

377   segment is from gently triangular to sub-rounded in these two species referred to

378   *Neotamandua*, without being rounded as in *Tamandua*.

379   **Astragalus.** Hirschfeld (1976) described the astragalus of *N. borealis* (Fig. 4A-B, 4E-

380   F) as intermediate in morphology and size between those in *Tamandua* and

Myrmecophaga. In contrast, Kraglievich (1940) stated that the astragalus of *N.*

greslebini closely resembles that of *Tamandua*. New observations allow to

determinate that, in dorsal view, the astragalus of *N. borealis* is more similar to that in

Tamandua than *Myrmecophaga* as consequence of a lateral side of the trochlea larger

than the medial one (trochlear asymmetry). Like *N. greslebini*, the regular concavity

in which is inserted the *flexor digitorum fibularis* tendon extends posteroventrally

like a well-defined wedge ("pointed shape" in Kraglievich's words) and it contacts

the calcaneal facets across the entire width of the latter. In ventral view, the

arrangement of the calcaneal facets in *N. borealis* is a kind of 'transition' between

that in *Myrmecophaga* and *Tamandua*. In *N. borealis*, the ectal and sustentacular are

largely separated by a wide and deep sulcus, but there is an incipient connection. This

condition is approximately comparable to that described by Kraglievich (1940) for *N.*

greslebini and differs from the fully separated calcaneal facets in *Protamandua* and

Tamandua. In this sense, Kraglievich was not very explicit in pointing out the degree

of development of the connection between these facets, but it is inferred that it is not

exactly wide as in *Myrmecophaga* when he wrote that '…these calcaneal articulations

are, *apparently*, posteriorly fused…' (italics are ours; Kraglievich, p. 635).

Calcaneum. Like *N. greslebini*, *N. borealis* has a narrow fibular calcaneal facet,

which is located laterally and in a slightly different plane with respect to that of the

ectal facet (Fig. 4C-D, 4G-H). In both of the former species, the *sustentaculum* is less

medially projected than in *Myrmecophaga*. They also show an accessory facet in the

anterior end of the calcaneum that articulates with the astragalar head, similarly to

Tamandua. In all the aforementioned taxa, this facet is closer (even in contact) to the

cuboid facet. In *N. borealis* and *N. greslebini*, the cuboid facet is transversely ovate

and concave. A unique feature in common for them is the presence of a short

tendinous groove (shorter than in *Myrmecophaga*) and strongly concave (Fig. 4H). It

is the continuation of the longitudinal and conspicuous ridge that runs the calcaneum

in its lateral side. The latter separates tendons of the *fibularis longus* and *accesorius*

muscles (Hirschfeld 1976; Gambaryan et al. 2009). In *N. borealis*, this ridge is more

conspicuous than in *Tamandua* and less than in *Myrmecophaga* [Insert Fig. 4 here].

N. borealis and **Neotamandua sp.** See the Table 3 for comparison of postcranial

measurements between these species.

Radius. The distal epiphysis of the radius in *Neotamandua* sp. (MACN 2408) is more

massive than that in *N. borealis*. In the latter species, the distal end of the radius is

relatively stylized, like *Tamandua*. However, the morphologies of *N. borealis* and

Neotamandua sp. are more comparable between them. In distal view, the styloid

process of these species is more elongated and posteriorly oriented than in

Tamandua. In the latter extant genus, the transverse axis (longer axis) of the facet for

distal articulation is forming an angle close to 45° with respect to the plane of the

anterior side of the radius, while this axis is nearly parallel with respect that plane in

N. borealis and *Neotamandua* sp. This difference gives to the distal radius of the

compared *Neotamandua* species a non-rotated appearance, unlike the same epiphysis

in *Tamandua*. In anterior view, the distal articulation facet of *N. borealis* and

Neotamandua sp. is visible in wedge shape pointing towards the medial border.

Additionally, in the same view, this facet exhibits comparable exposures in both of

the latter species, considerably more than in *Tamandua*. The posterior side of the

distal epiphysis is from flat to slightly concave in *N. borealis* and *Neotamandua* sp.,

unlike the convex posterior side in *N. greslebini* (this observation could suggest that

the material of *Neotamandua* sp. is not the holotype of *N. greslebini*, as speculated by

McDonald et al. 2008) and *Tamandua*. The distal extension of the lateral ridge in *N.*

borealis and *Neotamandua* sp. is similar.

Astragalus. The astragalus of *Neotamandua* sp. (MACN 2406) is only represented

by the astragalar body. The medial trochlea is smaller than the lateral trochlea, but

this asymmetry is less than in *N. borealis*. In addition, these sections of the trochlea

are proportionally less separated in the latter species than in *Neotamandua* sp.

Calcaneum. The calcaneum is fragmentary in *Neotamandua* sp. (MACN 2411). As

in the case of the astragalus, the preserved portion is the bone body. The ectal facet is

sub-triangular in shape in *Neotamandua* sp., while it is approximately sub-oval in *N.*

borealis. The sustentacular facet is more medially extended in the latter species than

in *Neotamandua* sp. In both species, the cuboid facet is partially visible in dorsal

view, particularly in *Neotamandua* sp. In the same view, the lateral ridge is slightly

exposed in *N. borealis*, but not so in *Neotamandua* sp.

Discussion. The former comparisons allow to recognize a few morphological similarities

and differences between homologous postcranial elements of *N. borealis*, *N. greslebini* and

Neotamandua sp. It is considered that some similarities in these species are potentially

diagnostic at the genus level, namely the sub-rounded to gently triangular shape of the tibial

mid-section; ectal and sustentacular facets incipiently connected in the astragalus; and a

short tendinous groove in the lateral side of the calcaneum (Table 4). These similarities

seem to provide support to the hypothesis that these northern and southern South American

species referred to *Neotamandua* are closely related and, consequently, that they are

correctly included in the same genus. Alternatively, these common features could be

symplesiomorphies of a hypothetical lineage of myrmecophagids more late diverging than

Protamandua and apparently closer to *Myrmecophaga* than *Tamandua*. Provisionally, from

the analysis presented, it is proposed to circumscribe the genus *Neotamandua* to the

nominal species *N. conspicua* (type species), *N. greslebini* and *N. borealis*. Since *N. magna*

and *N.? australis* are doubtfully assigned to *Neotamandua* or its allocation in this genus has

been seriously questioned (McDonald et al. 2008; this work), they are considered *species*

inquirendae, following the International Code of Zoological Nomenclature (Ride et al.

1999). To denote the questionable generic allocation of *N. magna* is suggested the use of

inverted commas, i.e. *'N.' magna*. The material referred to *Neotamandua* sp. seems

correctly referred to this genus, but it should be further tested. It is possible that these

specimens correspond to a new species.

The diagnosis for *Neotamandua* proposed below is largely based on the designation of the

specimen FMNH P14419 as epitype for the type species, *N. conspicua*, after considering

the fragmentary nature of the holotype of this taxon (MACN 8097; Rovereto 1914), and,

consequently, its ambiguity or lack of some taxonomically relevant features, particularly in

the rostrum. In addition, the potentially diagnostic postcranial features for *Neotamandua*

that has been identified above are also incorporated in the new diagnosis until cranial-

postcranial associations are found and studied.

472   *Neotamandua* Rovereto, 1914

LSID. urn:lsid:zoobank.org:act:4EC0ABE1-C013-4113-9956-5DBD6E79FCEA

Type species. *N. conspicua* Rovereto, 1914.

Other referred species. *N. greslebini* Kraglievich 1940; *N. borealis* Hirschfeld 1976.

Related *species inquirendae*. '*N.*' *magna* Ameghino 1919 (*Myrmecophaga magna* new

combination? See McDonald et al. 2008); *N.*? *australis* Scillato-Yané & Carlini 1998.

Diagnosis. Middle-to-large sized myrmecophagid, larger than *Tamandua* but smaller than

Myrmecophaga. It can be differentiated from other vermilinguans by the following

combination of characteristics: in dorsal view, rostrum strongly tapered towards its anterior

end (more than in any other myrmecophagid), with a regular transition in width from the

anterior portion of frontals to the anterior end of nasals; reduced lacrimal which is not part

of the orbit; jugal inserted in posteroventral position with respect to the lacrimal and

slightly projected in posterodorsal direction; frontal forming a dorsal dome at the orbit

level; hard palate well extended towards the posterior end of the skull, close to the ventral

border of the occipital condyles; squamosal (= posterior) zygomatic process dorsally

inclined; presence of palatine "wings"; horizontal inclination of the glenoid (Gaudin &

Branham 1998); sub-oval to gently triangular shape of the tibial mid-section; ectal and

sustentacular facets incipiently connected in the astragalus; short tendinous groove in the

lateral side of the calcaneum.

492   *Neotamandua conspicua* Rovereto, 1914

493   (Fig. 5)

LSID. urn:lsid:zoobank.org:act:C4DC62D5-6470-4A04-B152-D42ED3BA332C

Holotype. MACN 8097, posterior portion of a skull.

Epitype. FMNH P14419, nearly complete skull but with fractured rostrum and partially

eroded frontals and parietals.

Geographical and stratigraphic provenance. MACN 8097 is from an indeterminate

locality in the Santa María Valley, Catamarca Province, Argentina (Rovereto 1914).

Probably Andalhuala Formation. Upper Miocene (McDonald et al. 2008; Bonini 2014;

Esteban et al. 2014).

FMNH P 14419 is from the Corral Quemado area, Catamarca Province, Argentina. Corral

Quemado Formation. Lower Pliocene (Bonini 2014; Esteban et al. 2014). This specimen

was collected by Robert Thorne and Felipe Méndez during the Second Captain Marshall

Field Palaeontological Expedition, which was led by Elmer S. Riggs and developed in

Argentina and Bolivia in 1926☐1927 (Simpson, pers. comm.; Riggs 1928). In the Field

Museum, where it is deposited, has been catalogued as *N. conspicua*. No known reference

publication exists as support for the taxonomic assignation to this species, except in Gaudin

& Branham (1998) and, now, in this work from direct comparison with the holotype.

Diagnosis. See the diagnosis for *Neotamandua* above. The postcranial diagnostic features

included there do not belong to material known for this species.

Comparative description of the epitype. The skull FMNH P14419 is anteroposteriorly

elongated, with a general architecture more similar to that in *Myrmecophaga* than

Tamandua. The cranial measurements taken for this specimen are shown in the Table 1. In

dorsal view, both the rostrum, in general, as well as the nasals, in particular, are anteriorly

tapered. The pre-orbital section of the frontals is proportionally less elongated than in

Myrmecophaga. The lacrimal has a sub-triangular outline and its anteroposterior and

dorsoventral lengths are similar, unlike *Myrmecophaga*, in which the lacrimal is triangular

and more anteroposteriorly elongated. The insertion of the jugals is more ventral and

posterior than in *Myrmecophaga* and even more than *Tamandua*. Each jugal is slightly

tapered by mediolateral compression in its posterior end and it is posterodorsally projected,

instead of posteroventrally like *Myrmecophaga*. The posterolateral process of the maxilla

contacts the entire anterior and ventral borders of the lacrimal. The orbital ridge is less

prominent than in *Myrmecophaga*. The superior orbital wall is laterally expanded, forming

a roof more developed than in *Myrmecophaga*. At the orbit level, the palatines are also

laterally expanded, forming palatine "wings". These structures make the anterior hard

palate look wider than the posterior palate. The posterior end of the hard palate is less

ventrally projected, unlike *Tamandua* and *Myrmecophaga*. In lateral view, the squamosal

zygomatic processes are dorsally inclined, unlike the ventral inclination of the same bone

projection in *Tamandua* and *Myrmecophaga*. This feature would be a convergence with

Cyclopes. The braincase is proportionally larger than in *Myrmecophaga*, but smaller than in

Tamandua. The tympanic bulla is less developed than in *Tamandua*. The external auditory

meatus has subcircular to circular shape, like *Myrmecophaga* (ovated in *Tamandua*). In *N.*

conspicua the same opening is located in a posterodorsal position, like *Myrmecophaga* and

in contrast with *Tamandua*, in which it has an anterodorsal position. Despite the

palatopterygoid suture is not well preserved, appears to be more similar to the irregular

suture in *Myrmecophaga*, with a posteriorly opened, asymmetrical "V" shape, than the

regular suture in *Tamandua*, with an anteriorly opened, symmetrical "V" shape. There is no

interpterygoid vacuity in which it could be established a soft palate, like *Myrmecophaga*.

The occipital condyles are proportionally larger than in *Myrmecophaga* [Insert Fig. 5 here].

Discussion

Systematic implications

This works includes the first description of a new, valid extinct genus for

Myrmecophagidae in the last century, i.e. *Myrmedioctes* gen. nov. Likewise, it constitutes a

novel taxonomic comprehensive reassessment for *Neotamandua* from Hirschfeld (1976).

The results suggest that there are still critical gaps in our knowledge on the composition

and diversity of the Neogene assemblages of these xenartrans, particularly in the tropical

region of South America. With the inclusion of Gen. et sp. nov. (Fig. 6), Myrmecophagidae

now comprises at least five genera (three of them fully extinct) and 11 nominal species

(eight extinct species), namely [the dagger means extinct species]: *Protamandua rothi*[†];

Neotamandua? australis[†]; *Neotamandua borealis*[†]; Gen. et sp. nov.[†]; '*Neotamandua*'

magna[†]; *Neotamandua greslebini*[†]; *Neotamandua conspicua*[†]; *Myrmecophaga*

caroloameghinoi[†]; *Myrmecophaga tridactyla*; *Tamandua tetradactyla*; and *Tamandua*

mexicana. Of these taxa, only two genera and two species have fossil occurrence in

northern South America: *N. borealis* (Middle Miocene of Colombia; Hirschfeld 1976) and

Gen. et sp. nov. (Middle Miocene of Colombia; this work) (Fig. 7). The latter taxon is a

small-to-middle sized myrmecophagid, comparable but slightly smaller than *Tamandua*.

The general morphology of the skull of this new anteater resembles more to that of

Tamandua than any other known taxon. It shows remarkable features such as: (1) strongly

tapered nasals toward its anterior rostrum; (2) relatively low rostrum and anterior section of

frontals; (3) large pre-orbital section of frontals; and (4) strongly triangulated (anterior)

lacrimal. The tapering of nasals is a characteristic in common with *N. conspicua*, but in the

latter species the entire rostrum is tapered, not only the nasals, like Gen. et sp. nov. The

relatively low rostrum and anterior section of frontals seems to indicate a plesiomorphy,

given that this feature is apparently present in *P. rothi*. A large pre-orbital section of

frontals is shared, in (nearly) extreme condition, by *N. conspicua* and, especially,

Myrmecophaga, but it should be noted that in Gen. et sp. nov. there is no such as elongated

skull. And, finally, the strongly triangulated (anterior) lacrimal in the latter species is

superficially similar to that in *Myrmecophaga*. Estimates of cranial measurements and

features (rostrum length, exposure of the maxilla in the orbit and curvature of the

basicranial-basifacial axis) used for coding the characters with numbers 4, 8, 9 and 42 of

the character list by Gaudin & Branham (1998), allow tentatively infer the phylogenetic

position of Gen. et sp. nov. as a taxon included within the clade *Tamandua + Neotamandua*

+ Myrmecophaga and located in a polytomy with *Tamandua*. Under this preliminary

phylogenetic analysis, which is not presented in the results section because there is no

enough information for coding the new taxon, *Protamandua* is well supported as the most

basal myrmecophagid as consequence of sharing several character states with non-

Myrmecophagidae Vermilingua (i.e. *Cyclopes* and *Palaeomyrmidon*; for more details, see

Gaudin & Branham 1998). For future studies, it is tentatively suggested the subfamilial

name "Myrmecophaginae" for all the Myrmecophagidae more late diverging than

Protamandua, including possibly Gen. et sp. nov. In this sense, new and more complete

material referable to the latter taxon is required to shed light on its phylogenetic position

[Insert Fig. 6 here].

On other hand, the taxonomic analysis of *Neotamandua* and its referred species indicates

that these taxa were based on a poorly supported taxonomy. Other case of extinct

vermilinguans with flawed systematics in low levels of the taxonomic hierarchy was noted

by McDonald et al. (2008) with regard to genera and species proposed from isolated

postcranial elements of putative myrmecophagids or even members of new, distinct

families from the Early Miocene of Santa Cruz, southern Argentina. These authors,

partially based on comparisons by Hirschfeld (1976), argued that the number of taxa

claimed for that area and interval (seven genera and nine species; e.g. *Promyrmephagus*,

Adiastaltus; Ameghino 1894) has been artificially inflated, even though it is still possible to

revalidate taxa other than the well validated species *P. rothi* (McDonald et al. 2008). All

these research problems in systematics imply the need to regularly reevaluate the taxonomy

of extinct anteaters through reexamination, when possible, of previously described material

and the study of new specimens. While it is true that the fossil record of Vermilingua is

poor and fragmentary in comparison, for instance, with that of other xenartrans such as

Tardigrada, the sampling effort should be increased in order to have greater recovery of

fossil material for this group, especially in areas known for their preservation potential (e.g.

southern and northwestern Argentina, southwestern Colombia).

The reevaluation of the taxonomic status of *Neotamandua* found that there was no

diagnosis for this genus. The newly proposed diagnosis includes multiple cranial and

potential postcranial characteristics, which uphold that *Neotamandua*, independently

whether it is a natural group or not, certainly contains species that do not belong to

Myrmecophaga, despite their great resemblance with the latter. This outcome is congruent

with the taxonomic opinion of Gaudin & Branham (1998) and is at odds with Patterson et

al. (1992). Now, can we confidently say that *Neotamandua* is monophyletic from current

evidence? *Neotamandua*, as previously defined by other workers, may be composed of

successive basal species or genera in relation to the hypothetical clade of *Myrmecophaga*

(i.e. *My. tridactyla + My. caroloameghinoi*). If that is correct, *Neotamandua* would be

paraphyletic by definition, since it excludes some of its descendants (Sereno et al. 1991).

This possible pattern of basal paraphyly is consequence of a taxonomy not defined by

clades, but grades (Huxley 1958; Wood & Lonergan 2008). The monophyly of

Neotamandua, as was redefined here (i.e. *N. conspicua + N. greslebini + N. borealis*), is

tentatively supported by three potential synapomorphies shared by two of its species whose

postcranium is known (*N. greslebini* and *N. borealis*): (1) sub-oval to gently triangular mid-

section of the tibia; (2) ectal and sustentacular facets incipiently connected in the

astragalus; (3) short tendinous groove in the lateral side of the calcaneum. However, the

synapomorphic condition of these features for *Neotamandua* need to be further tested from

systematic analysis of new, more complete and/or associated material of Gen. et sp. nov.

and species referred to *Neotamandua*. That would allow to assess more adequately the

global morphological variability and character distribution in Miocene myrmecophagids

more late diverging than *Protamandua*. In turn, knowing this distribution better, it is more

likely to disentangle the taxonomic identities and affinities of the *Neotamandua* species in

order to corroborate the monophyly of this genus. For the moment, the hypothesis of

Hirschfeld (1976) that *Neotamandua* is not monophyletic is, in principle, less probable if

the *species inquirendae* '*N.*' *magna* and *N.? australis* are excluded from the genus, as it

was decided here, than if they are retained within it. The exclusion of the *species*

inquirendae does not affect the hypothesis that *Neotamandua* is closer to *Myrmecophaga*

than any other known nominal genus. Consequently, the type species of *Neotamandua*, *N.*

conspicua, is reiterated as closer to *Myrmecophaga* than *Tamandua*, in line with the

phylogeny of Gaudin & Branham (1998) and unlike the hypothesis of Carlini & Scillato-

Yané (1998).

Finally, the material referred to *Neotamandua* sp. and used in this study to make

comparisons with *N. borealis*, seems correctly allocated in that genus, but it might

eventually be assigned to a new species with very large body size, larger than *N. greslebini*.

This is partially conditioned to the clarification of the taxonomic status of '*N.*' *magna*,

which is a species comparable in body size to *Neotamandua* sp., so they could be (or not)

the same taxon.

The diversification of Myrmecophagidae

McDonald et al. (2008) pointed out that since the highly incomplete fossil record of

Vermilingua, several fundamental questions on the evolution of this group, including

morphological trends and the acquisition of ecological preferences in its distinct taxa, are

largely unknown. Likewise, they highlighted some uncertainty related to the divergence

times of possible sub-clades. However, several inferences and hypotheses about the

evolutionary history of anteaters and, particularly, the myrmecophagids, can be outlined

from the current evidence, including that presented in this work. Following to Pascual &

Ortiz-Jaureguizar (1990), McDonald et al. (2008) and Toledo et al. (2017), the next

discussion is based on multiple palaeobiological, ecological and biogeographical aspects as

major constraints and/or consequences of the myrmecophagid evolution.

The diversification of Myrmecophagidae was an macroevolutionary event that occurred

through the Neogene, at least as early as the Burdigalian (Early Miocene), according to the

minimal age estimated for the most basal genus, i.e. *Protamandua*. The beginning of this

diversification is approximately overlapped in time with the onset or development of

similar events in other higher taxa in South America, such as the xenartrans

Megatherioidea, Mylodontidae, Glyptodontidae and Dasypodini (Croft et al. 2007;

McDonald & De Iuliis 2008; Bargo et al. 2012; Carlini et al. 2014; Boscaini et al. 2019), or

the South American native ungulates Pachyrukhinae, Mesotheriinae and Toxodontidae

related to *Pericotoxodon* and *Mixotoxodon* (Seoane et al. 2017; Armella et al. 2018a;

Armella et al. 2018b). This pattern shows the importance of the Early Miocene, particularly

the Burdigalian, as a critical interval for the diversification of multiple South American

land mammal lineages. In light of the geographical provenance of *Protamandua*, the most

probable ancestral area for Myrmecophagidae is southern South America (Fig. 7). The

palaeonvironmental conditions inferred for the Early Miocene of this area are considerably

warmer and more humid (1000□1500 mm/year) than today, with presence of a subtropical

dry forest (Iglesias et al. 2011; Quattrocchio et al. 2011; Kay et al. 2012; Brea et al. 2017;

Raigenborm et al. 2018). In line with this reconstruction, Palazzesi et al. (2014), using a

rarefied richness analysis from palynological evidence, reported that southern Argentina

harboured in the Early Miocene a plant richness comparable to that documented today for

the Brazilian Atlantic Forest, in southeastern Brazil. Similar to *Tamandua*, *Protamandua*

would have preferred forested habitats and would have had semiarboreal habits (Gaudin &

Branham 1998; McDonald et al. 2008; Kay et al. 2012). Whether the ancestral condition of

substrate use in Myrmecophagidae is arboreal, as held by Gaudin & Branham (1998), the

preference for open biomes (e.g. savannah) and terrestriality in *Myrmecophaga* (and

possibly in *Neotamandua*) is a derived condition (McDonald et al. 2008; Toledo et al.

2017). The semiarboreal habits of *Tamandua* are explained from niche conservatism or,

alternatively, from convergence with *Protamandua* if the ancestor of *Tamandua* was

hypothetically terrestrial [Insert Fig. 7 here].

Since their particular, low basal metabolic rates and myrmecophagous diets (McNab 1984,

1985), it is likely that the global warm recovery during the early Neogene (Early Miocene

to early Middle Miocene; including the Middle Miocene Climatic Optimum or MMCO;

Fig. 8), linked to latitudinal temperature gradient reduction and expansion of the tropical

(warm) forest belt towards higher latitudes in the continents (including South America; see

Anderson 2009; Herold et al. 2011; Morley 2011; Palazzesi et al. 2014), has influenced on

the evolutionary differentiation of the myrmecophagids, maybe predominantly *in situ* as in

the climatically-induced evolution of other small Cenozoic mammals (Fortelius et al.

2014), such as *Protamandua*. This differentiation would have been triggered by increase in

suitable area in terms of preferred biomes (warm forests in this case) and, especially,

temporarily sustained availability of social insects for their feeding (McDonald et al. 2008;

Kay et al. 2012; Toledo et al. 2017). Indeed, extant termites and ants (Termitidae and

Formicidae, respectively) concentrate the vast majority of their biomass (and species

richness) in the tropics and warm subtropical regions (Hölldobler & Wilson 1990; Tobin

1995; Davidson & Patrell-Kim 1996; Eggleton et al. 1996; Davidson et al. 2003; Ellwood

& Foster 2004; Keller & Gordon 2009). This ecogeographical pattern is consistent with the

fossil record of the former higher taxa, which shows a strong tropical niche conservatism

from their respective evolutionary origins in the Late Cretaceous/Early Paleogene (see

below; Archibald et al. 2011; LaPolla et al. 2013; Guénard et al. 2015; Bourguignon et al.

2017).

According to Blois & Hadly (2009), the responses of mammalian taxa to climate change

throughout the Cenozoic are causally interconnected. These responses at the level of

individual taxa may include changes in abundance, genetics, morphology and/or

distributional range, and they may instigate multitaxa responses such as diversification

events comparable to that placed on the root of the evolutionary tree of Myrmecophagidae.

This case of a cladogenetic event possibly induced by climate contrasts in kind of biome

with those that have been repeatedly documented for intervals of grassland expansion (e.g.

Equidae, Bovidae, Cervidae, Ochotonidae, Hippopotaminae; MacFadden 2000;

Bouchenak-Khelladi et al. 2009; Boisserie & Merceron, 2011; Ge et al. 2013).

In the Middle Miocene, *N.? australis*, *N. borealis* and Gen. et sp. nov. exhibit a mosaic of

morphological features in common with *Tamandua* and/or *Myrmecophaga*, as well as some

exclusive characteristics, which suggest an early, important increase in morphological

disparity in Myrmecophagidae and possibly the evolutionary divergence of those lineages

comprising its crown-group. This coincides with the interpretation of Hirschfeld (1976),

according to which the lineages including the extant genera of Myrmecophagidae

differentiated morphologically at least from the Friasian (Middle Miocene). Same way, it is

compatible with the results of the molecular phylogenies by Delsuc et al. (2001, 2012) and

Gibb et al. (2016), which estimated that the evolutionary divergence of *Tamandua* and

Myrmecophaga occurred in the late Middle Miocene, c. 13 mya. On the other hand, relative

body sizes inferred for the Middle Miocene taxa show an apparent trend towards increase in

body size in comparison with the basal taxon *Protamandua*. During this interval, the

myrmecophagids have a wide geographical distribution in South America (Fig. 7), from

low to medium-high latitudes. This is in line with the evolution of larger body sizes since

when this attribute increases, the foraging area also increases and, with it, the distributional

range, according the general foraging strategy of the extant myrmecophagids (Naples 1999;

Toledo et al. 2017; Gaudin et al. 2018). The co-occurrence pattern of *N. borealis* and Gen.

et sp. nov. in La Venta area in Colombia constitutes the earliest pattern of this kind for

Myrmecophagidae until pending systematic revisions for putative taxa from the Early

Miocene of Santa Cruz, Argentina, are carried out. These revisions would allow to

determine whether there are two or more co-occurrent myrmecophagid taxa in the latter

area. Given that *N. borealis* and Gen. et sp. nov. probably are not sister taxa, it would imply

a non-sympatric diversification followed by dispersal of at least one of the involved taxa.

The habitat preference of Gen. et sp. nov. in the palaeoenvironmental mosaic of La Venta

area (Kay & Madden 1997; Spradley et al. 2019) is speculated as tropical forest

(semiarboreal?) by analogy with *Protamandua*, while it is proposed a more generalized

habitat selection for *N. borealis* in line with the palaeobiological inference of

predominantly terrestrial locomotion for the latter taxon by Hirschfeld (1976). If this hold

true, opens the possibility that *N. borealis* is the oldest myrmecophagid inhabiting zones

with semi-open or even open vegetation (see below).

The morphological and probably taxonomic diversification of Myrmecophagidae continued

in the Late Miocene. Inferred body sizes range from larger than *Tamandua* and nearly

comparable to *Myrmecophaga*. Considering the wide geographical distribution during the

Middle Miocene, there is probably a geographical bias in the fossil record of the

myrmecophagids during the Late Miocene as the only known occurrences are

Myrmecophaga-like forms from northwestern Argentina (Fig. 7). If *N. borealis* and *N.*

greslebini are sister taxa, as it seems, that means there was a biogeographical connection

for Myrmecophagidae between northern and southern South America in the late

Middle/early Late Miocene. This inference is congruent with the palaeobiogeographical

analyses of Cozzuol (2006) and Carrillo et al. (2015), according to which the affinities

between several Late Miocene, northern and southern South American land mammal

assemblages are strong or, at least, not so distant as those between Middle Miocene

assemblages from the same regions. This pattern might be explained from the geographical

shrinks of the Pebas Mega-Wetland System and the Paranean Sea in the Middle-Late

Miocene transition (Aceñolaza & Sprechmann 2002; Cozzuol 2006; Salas-Gismondi et al.

2015). It is also possible that the expansion of open biomes in South America during the

Late Miocene has facilitated this biotic connection, as has been acknowledged in the case

of other mammal taxa (e.g. Glyptodontinae, a xenartran group like Myrmecophagidae;

Ortiz-Jaureguizar & Cladera 2008; Oliva et al. 2010). Indeed, from a palaeoenvironmental

viewpoint, the (partial?) co-occurrence of '*N.*' *magna*, *N. greslebini* and *N. conspicua* in

northwestern Argentina is important inasmuch as this pattern is related, for the first time in

the evolutionary history of Myrmecophagidae, to savannahs well developed with respect to

other kinds of vegetation cover (Latorre et al. 1997; Brandoni et al. 2012; Cotton et al.

2014; Amidon et al. 2017; Zimicz et al. 2018). On the basis of the foregoing and by

generalization of morphological and ecological features of the living vermilinguans, e.g.

less dependence on trees related to greater taxonomic and/or ecological diversity of

consumed insects (Hirschfeld 1976; Montgomery 1985a; Rodrigues et al. 2008; Toledo et

al. 2017; Table 5), it is hypothesized that, as early as the late Middle Miocene, with the

triggering of a global cooling (Fig. 8), *Neotamandua* was involved in a niche evolution

process within Myrmecophagidae which implied a significative increase in dietary diversity

as myrmecophagous and expansion of substrate use and biome selection. Probably the

species of this genus preferred the frequent use of the ground by biomechanical constraints

and made inroads into largely open environments as humid savannahs, without excluding

use of forested environments, like *Myrmecophaga* (Fuster et al. 2018; Gaudin et al. 2018).

The former model is further supported from the evolutionary response pattern to major

climatic-vegetational changes documented by Badgley et al. (2008) in a faunal sequence of

mammals from the Late Miocene of southern Asia, according to which the trophic niche

evolution and, particularly the expansion of this attribute, in conjunction with habitat

changes, is related to increase in the probabilities of local and regional survivorship in the

studied lineages [Insert Fig. 8 here].

On other hand, the fossil record of the crown-group genera, *Tamandua* and *Myrmecophaga*,

is confined to the Pliocene-Pleistocene, but the evolutionary (morphological) divergence of

Myrmecophaga would date back at least to the late Middle Miocene according the first

appearance of *Neotamandua*, i.e. *N. borealis*. Under this assumption, the hypothesis of 'N.'

magna as a species of *Myrmecophaga* is perfectly feasible. In any case, the biogeographical

dynamics of the two extant myrmecophagid genera may have been constrained by their

respective ecological tolerances and, they, in turn, by the rapidly changing habitat and

biome distribution in the Americas during at least the last five or six million years (de Vivo

& Carmignotto 2004; Salzmann et al. 2011; Sniderman et al. 2016; Amidon et al. 2017;

Roberts et al. 2018; Grimmer et al. 2018). This applies especially to the case of *Tamandua*

since this taxon is less generalist in relation to habitat selection than *Myrmecophaga*

(McDonald 2005). Considering the hypothesis of niche expansion for *Neotamandua*, the

differentiation of *Myrmecophaga* would have accentuated this putative evolutionary trend

through stronger preference for open environments, which is consistent with the general

palaeoenvironment of savannah in the Early Pliocene of the area where occurs the oldest

species of the latter genus, i.e. *My. caroloameghinoi* (Zavala & Navarro 1993; McDonald et

al. 2008).

The myrmecophagid evolution has a late episode with the complete formation of the

Panama Land Bridge (PLB) in the terminal Neogene (Coates & Stallard 2013; O'dea et al.

2016; Jaramillo 2018). *Myrmecophaga tridactyla* invaded and colonized Central- and

southern North America (northern Mexico) at least as early as the Early Pleistocene (Shaw

& McDonald 1987; Fig. 7). This dispersal event is part of the Great American Biotic

Interchange (GABI), specifically the episode referred as GABI 2 (Woodburne 2010).

Today, the northern boundary of this species is located in northern Central America, over

3000 Km to the south of the northernmost fossil record (Gaudin et al. 2018). This

distributional difference was interpreted by Shaw & McDonald (1987) from the occurrence

of warmer and more humid conditions in the Early Pleistocene of southern North America

(southern United States-northern Mexico) than today in the same area. These conditions

would have allowed that *Myrmecophaga* colonize subtropical savannahs with permanent

availability of insects included in its diet (Croxen III et al. 2007; McDonald 2005), but

subsequent climatic-vegetational shifts (desertification) during the Late Pleistocene would

have forced from extirpation a range shrinkage of this taxon towards lower latitudes

(McDonald 2005; Ferrusquía-Villafranca et al. 2017). The distributional range pattern of

tropical taxa expanded towards southern North America during some intervals of the

Pleistocene has been well supported from the records of multiple taxa other than

Myrmecophaga, including mammals and sauropsids (Shaw & McDonald 1987; Moscato &

Jasinski 2016; Ferrusquía-Villafranca et al. 2017).

Like *Myrmecophaga*, *Tamandua* also colonized (or evolved in) northern continental

territories outside South America. This is supported from the occurrence of *Tamandua* sp.

in the terminal Pleistocene of Central Mexico (Arroyo-Cabrales et al. 2004; Ferrusquía-

Villafranca et al. 2010; Fig. 7). In its northern zone, the current distributional area of *T.*

mexicana includes latitudes comparable with that of the referred fossil record for this

species (Navarrete & Ortega 2011). Central Mexico is part of the transitional area between

the current Neotropical and Neartic regions, called Mexican Transition Zone (MTZ;

Halffter & Morrone 2017). All these observations, in conjunction with the above

interpretation of the Neogene biogeographical and environmental patterns, suggest that

Myrmecophagidae kept throughout its evolutionary history a niche conservatism associated

with tropical (warm) habitats (a case of phylogenetic niche conservatism or PNC; see

Cooper et al. 2011; Fig. 8), possibly in parallel with the same pattern in species groups of

its prey insects (Thompson 1994). Even more, the fact that Myrmecophagidae currently

accumulates its highest species richness in the warmest and wettest belt of the Americas

(Hayssen 2011; Navarrete & Ortega 2011; Miranda et al. 2017; Gaudin et al. 2018) is

further interpreted as evidence that this higher taxon represents support for the tropical

niche conservatism hypothesis (TCH; Wiens & Donoghue 2004; Wiens & Graham 2005).

However, in line with the discussion above, this major ecological constraint in

Myrmecophagidae is not only related to environmental thermal tolerance (see McNab

[1985] for an analysis on the thermophysiological constraints of the Xenarthra; McNab

[1984] also discussed the same issue for myrmecophagous mammals), as emphasized by

TCH, but it is also driven by food availability, at least by limiting or preventing historical

colonization of low-productivity regions far from the tropics (Shaw & McDonald 1987;

McDonald 2005; Šímová & Storch 2017; Fig. 8).

Conclusion

The systematic evidence presented here suggests that probably the diversification of

Myrmecophagidae is taxonomically and biogeographically more complex than previously

thought. This insight is based on the description of the new taxon Gen. et sp. nov. for the

Middle Miocene of Colombia (co-occurrent species of *N. borealis*) and the determination

of *Neotamandua*, as previously defined, as a wastebasket taxon which is probably formed

by species belonging to more than one single genus. While Gen. et sp. nov. possibly has

affinities with *Tamandua*, more information is needed to test its phylogenetic position

within Myrmecophagidae. On the other hand, *N. borealis*, *N. greslebini* and *Neotamandua*

sp. share postcranial features (potential synapomorphies) that imply some grade of kinship

between them. Therefore, the two nominal species among the former ones are provisionally

kept within *Neotamandua*. Alternatively, these features also may constitute

symplesiomorphies of a hypothetical lineage which is apparently close to *Myrmecophaga*.

The remaining nominal species referred to *Neotamandua*, i.e. '*N.*' *magna* and *N.?* *australis*

were designated as *species inquirendae*. Overall, it is necessary to develop new systematic

revisions, including new phylogenetic analyses similar to that of Gaudin & Branham

(1998), from new material referable to Gen. et sp. nov. and the referred species to

Neotamandua, so as to obtain enough evidence to solidly determine the phylogenetic

position of the new species from La Venta and corroborate the putative monophyletic status

of *Neotamandua*. In line with the foregoing considerations, the paleontological exploration

of Neogene sedimentary units in northern South America and northern Argentina is crucial

to improve our understanding of the diversification of Myrmecophagidae.

Acknowledgements

A lot of people and institutions have contributed in distinct ways to the development of this

research. We would like to specially thank Andrés Vanegas and his family; the group of

defenders of the palaeontological heritage *Vigías del Patrimonio Paleontológico La*

Tatacoa; Carlos Jaramillo (Smithsonian Tropical Research Institute); and all the staff of the

Museo de Historia Natural La Tatacoa for assistance and providing access to the holotype

of the new species described for La Venta. Alejandro Kramarz (MACN), Kenneth

Angielczyk (FMNH), Bill Simpson (FMNH) and Patricia Holroyd (UCMP), Eduardo

Flórez (ICN), Alfredo Carlini (CAC) and Itatí Olivares (MLP) gave permission to study

extinct and extant anteater specimens in collections under their care. Sergio Vizcaíno and

Susana Bargo (MLP - CONICET) provided photos of postcranial specimens labelled as

Neotamandua sp in the MACN. Daniel Brinkman and Jamie Henderson (YPM) shared

photos of a skull of *Protamandua rothi* (YPM-15267). Matthew Pound (Northumbria

University) and Camila Hossotani (Universidade Federal de Mato Grosso do Sul, Brazil)

kindly helped by facilitating their postgraduate theses. James Zachos (University of

California, Santa Cruz) gave permission to reuse part of the data for his Cenozoic global

temperature curve. Jorge Gonzalez was the palaeoartist who exquisitely recreated the new

anteater from La Venta. K.J.L. received financial support from the following institutions:

CONICET (Internal Doctoral Fellowship); Field Museum (Science Visiting Scholarship);

University of California Museum of Paleontology (Welles Fund).

Statement of data archiving

The nomenclatural acts contained in this work are registered in Zoobank:

[identifier Gen. et sp. nov.]

LSID. urn:lsid:zoobank.org:act:4EC0ABE1-C013-4113-9956-5DBD6E79FCEA

LSID. urn:lsid:zoobank.org:act:C4DC62D5-6470-4A04-B152-D42ED3BA332C

References

Aceñolaza, F. & **Sprechmann, P.** 2002. The Miocene marine transgression in the

meridional Atlantic of South America. *Neues Jahrbuch für Geologie und Paläontologie-*

Abhandlungen, **225**, 75–84. doi:10.1127/njgpa/225/2002/75

Ameghino, F. 1894. Enumération synoptique des espèces de mammifères fossiles des

formations éocènes de Patagonie. *Boletín de la Academia Nacional de Ciencias en*

Córdoba, **13**, 259–455.

Ameghino, F. 1904. Nuevas especies de mamíferos, cretáceos y terciarios de la República

Argentina. *Anales de la Sociedad Científica Argentina*, **56–58**, 1–142.

Ameghino, C. 1919. Sobre mamíferos fósiles del piso araucanense de Catamarca y

Tucumán. Primera Reunión Nacional de la Sociedad Argentina de Ciencias Naturales, 1,

150–153.

Amidon, W. H., Fisher, G. B., Burbank, D. W., Ciccioli, P. L., Alonso, R. N., Gorin, A.

L. & Christoffersen, M. S. 2017. Mio-Pliocene aridity in the southcentral Andes

associated with Southern Hemisphere cold periods. *Proceedings of the National Academy*

of Sciences, **114**(25), 6474–6479. doi:10.1073/ pnas.1700327114.

Andersson, C. 2009. Neogene climates. Pp. 609–612 in V. Gornitz, (ed) *Encyclopedia of*

Paleoclimatology and Ancient Environments. Dordrecht, Springer.

Anderson, V. J., Horton, B. K., Saylor, J. E., Mora, A., Tesón, E., Breecker, D. O. &

Ketcham, R. A. 2016. Andean topographic growth and basement uplift in southern

Colombia: Implications for the evolution of the Magdalena, Orinoco, and Amazon river

systems. *Geosphere*, **12**(4), 1235–1256. doi:10.1130/GES01294.1

Archibald, S. B., Johnson, K. R., Mathewes, R. W. & Greenwood, D. R. 2011.

Intercontinental dispersal of giant thermophilic ants across the Arctic during early Eocene

hyperthermals. *Proceedings of the Royal Society B: Biological Sciences*, **278**(1725), 3679–

3686. doi:10.1098/rspb.2011.0729

Armella, M. A., García-López, D. A. & Domínguez, L. 2018a. A new species of *Xotodon*

(Notoungulata, Toxodontidae) from northwestern Argentina. *Journal of Vertebrate*

Paleontology, **38**(1), e1425882. doi:10.1080/02724634.2017.1425882

Armella, M. A., Nasif, N. L. & Cerdeño, E. 2018b. Small-sized mesotheriines

(Mesotheriidae, Notoungulata) from Northwestern Argentina: Systematic, chronological,

and paleobiogeographic implications. *Journal of South American Earth Sciences*, **83**, 14–

26. doi:10.1016/j.jsames.2018.02.002

Arroyo-Cabrales, J., Polaco, O. J. & **Johnson, E.** 2004. Quaternary mammals from

Mexico. Pp. 69–70 in L. C. Maul & R. D. Kahlke (eds) *Late Neogene and Quaternary*

Biodiversity and Evolution: Regional developments and interregional correlations (Vol. 2).

Weimar (Germany).

Badgley, C., Barry, J. C., Morgan, M. E., Nelson, S. V., Behrensmeyer, A. K., Cerling,

T. E. & **Pilbeam, D.** 2008. Ecological changes in Miocene mammalian record show impact

of prolonged climatic forcing. *Proceedings of the National Academy of Sciences*, **105**(34),

12145–12149. doi: 10.1073/pnas.0805592105

Bargo, M. S., Toledo, N. & **Vizcaíno, S. F.** 2012. Paleobiology of the Santacrucian soths

and anteaters (Xenarthra, Pilosa). Pp. 216–242 in S. F. Vizcaíno, R. F. Kay & M. S. Bargo,

(eds) *Early Miocene paleobiology in Patagonia: High-latitude paleocommunities of the*

Santa Cruz formation. Cambridge University Press, New York.

Barros, M. C., Sampaio, I. & **Schneider, H.** 2008. Novel 12S mtDNA findings in sloths

(Pilosa, Folivora) and anteaters (Pilosa, Vermilingua) suggest a true case of long branch

attraction. *Genetic and Molecular Biology*, **31**(3), 793–799.

doi:10.1016/j.jsames.2018.02.002

Blois, J. L. & **Hadly, E. A.** 2009. Mammalian response to Cenozoic climatic change.

Annual *Review of Earth and Planetary Sciences*, **37**, 181–208.

doi:10.1146/annurev.earth.031208.100055

Boisserie, J. R. & **Merceron, G.** 2011. Correlating the success of Hippopotaminae with

the C4 grass expansion in Africa: relationship and diet of early Pliocene hippopotamids

from Langebaanweg, South Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology*,

308(3–4), 350–361. doi:10.1016/j.palaeo.2011.05.040

Bonini, R. A. 2014. *Bioestratigrafía y diversidad de los mamíferos del Neógeno de San*

Fernando y Puerta de Corral Quemado (Catamarca, Argentina). PhD thesis, Universidad

Nacional de La Plata, Argentina, 366 pp. doi:10.35537/10915/34382

Boscaini A., Pujos F. & **Gaudin T.** 2019. A reppraisal of the phylogeny of Mylodontidae

(Mammalia, Xenarthra) and the divergence of mylodontine and lestodontine sloths.

Zoologica Scripta. doi:10.1111/zsc.12376

Bouchenak☐Khelladi, Y., Anthony Verboom, G., Hodkinson, T. R., Salamin, N.,

Francois, O., Ni Chonghaile, G. & **Savolainen, V.** 2009. The origins and diversification

of C4 grasses and savanna☐adapted ungulates. *Global Change Biology*, **15**(10), 2397–

2417. doi:10.1111/j.1365-2486.2009.01860.x

Bourguignon, T., Lo, N., Šobotník, J., Ho, S.Y., Iqbal, N., Coissac, E., Lee, M.,

Jendryka, M.M., Sillam–Dussès, D., Krížková, B., Roisin, Y. & **Evans, T.A.** 2017.

Mitochondrial phylogenomics resolves the global spread of higher termites, ecosystem

engineers of the tropics. *Molecular Biology and Evolution*, **34**(3), 589–597.

doi:10.1016/j.ympev.2017.09.025

Brandoni, D., Schmidt, G. I., Candela, A. M., Noriega, J. I., Brunetto, E. & **Fiorelli, L.**

E. 2012. Mammals from the Salicas Formation (late Miocene), La Rioja Province,

northwestern Argentina: Paleobiogeography, age and paleoenvironment. *Ameghiniana*,

49(3), 375–388. doi:10.5710/AMGH.v49i3(467)

Brea, M., Zucol, A.F., Bargo, M.S., Fernicola, J.C. & **Vizcaíno, S.F.** 2017. First

Miocene record of Akaniaceae in Patagonia (Argentina): a fossil wood from the early

Miocene Santa Cruz Formation and its palaeobiogeographical implications. *Botanical of*

the Journal Linnean Society, **183**, 334–347. doi:10.1093/botlinnean/bow014

Carlini, A. A., Castro, M. C., Madden, R. H. & **Scillato-Yané, G. J.** 2014. A new

species of Dasypodidae (Xenarthra: Cingulata) from the late Miocene of northwestern

South America: Implications in the Dasypodini phylogeny and diversity. *Historical*

Biology, **26**(6), 728–736. doi:10.1080/08912963.2013.840832

Carrillo, J. D., Forasiepi, A., Jaramillo, C. & **Sánchez-Villagra, M. R. 2015.**

Neotropical mammal diversity and the Great American Biotic Interchange: Spatial and

temporal variation in South America's fossil record. *Frontiers in Genetics*, **5**, 451.

http://doi.org/10.3389/fgene.2014.00451.

Chan, W. L., Abe-Ouchi, A. & **Ohgaito, R.** 2011. Simulating the mid-Pliocene climate

with the MIROC general circulation model: experimental design and initial results.

Geoscientific Model Development, **4**(4), 1035–1049.

Coates, A. G. & **Stallard, R. F.** 2013. How old is the Isthmus of Panama? *Bulletin of*

Marine Science, **89**(4), 801–813. doi:10.5343/bms.2012.1076

Cooper, N., Freckleton, R. P. & **Jetz, W**. 2011. Phylogenetic conservatism of

environmental niches in mammals. *Proceedings of the Royal Society B: Biological*

Sciences, **278**(1716), 2384–2391. doi:10.1098/rspb.2010.2207

Cope, E. D. 1889. The Edentata of North America. *The American Naturalist*, **23**, 657–664.

Cotton, J. M., Hyland, E. G. & **Sheldon, N. D.** 2014. Multiproxy evidence for tectonic

control on the expansion of C4 grasses in northwest Argentina. *Earth and Planetary*

Science Letters, **395**, 41–50. doi:10.1016/j.epsl.2014.03.014

Cozzuol, M. A. 2006. The Acre vertebrate fauna: age, diversity, and geography. *Journal of*

South American Earth Sciences, **21**(3), 185–203. doi:10.1016/j.jsames.2006.03.005

Croft, D. A., Flynn, J. J. & **Wyss, A. R. 2007.** A new basal glyptodontid and other

Xenarthra of the early Miocene Chucal Fauna, northern Chile. *Journal of Vertebrate*

Paleontology, **27**(4), 781–797. doi:10.1671/0272-4634(2007)27[781:ANBGAO]2.0.CO;2

Croxen III, F. W., Shaw, C. A. & **Sussman, D. R.** 2007. Pleistocene geology and

paleontology of the Colorado River Delta at Golfo de Santa Clara, Sonora, Mexico. Pp. 84–

89 in: R. E. Reynolds (ed.), *Wild, Scenic and Rapid: A Trip Down the Colorado River*

Trough. The 2007 Desert Symposium, Field Guide and Abstracts from Proceedings.

California State University, Desert Studies Consortium, Fullerton.

Davidson, D. W. & **Patrell-Kim, L.** 1996. Tropical arboreal ants: Why so abundant? Pp.

127–140 in A. C. Gibson (ed) *Neotropical biodiversity and conservation*. UCLA Botanical

Garden, Los Angeles, CA.

Davidson, D. W., Cook, S. C., Snelling, R. R. & **Chua, T. H.** 2003. Explaining the

abundance of ants in lowland tropical rainforest canopies. *Science*, **300**(5621), 969–972.

doi:10.1126/science.1082074

de Vivo, M. & Carmignotto, A. P. 2004. Holocene vegetation change and the mammal faunas of South America and Africa. *Journal of Biogeography*, **31**(6), 943–957. doi:10.1111/j.1365-2699.2004.01068.x

Delsuc, F., Ctzeflis, F. M., Stanhope, M. J. & Douzery, E. J. 2001. The evolution of armadillos, anteaters and sloths depicted by nuclear and mitochondrial phylogenies: Implications for the status of the enigmatic fossil *Eurotamandua. Proceedings of the Royal Society of London. Series B: Biological Sciences*, **268**(1476), 1605–1615. doi:10.1098/rspb.2001.1702

Delsuc, F., Superina, M., Tilak, M. K., Douzery E. J. P. & Hassanin, A. 2012 Molecular phylogenetics unveils the ancient evolutionary origins of the enigmatic fairy armadillos. *Molecular Phylogenetics and Evolution*, **62**, 673–680. doi:10.1016/j.ympev.2011.11.008

Dowsett, H., Dolan, A., Rowley, D., Moucha, R., Forte, A. M., Mitrovica, J. X., Pound, M., Salzmann, U., Robinson, M., Chandler, M. & Foley, K. 2016. The PRISM4 (mid–Piacenzian) palaeoenvironmental reconstruction. *Climate of the Past Discussions*, **12**, 1519–1538. doi:10.5194/cp-2016-33

Eggleton, P., Bignell, D. E., Sands, W. A., Mawdsley, N. A., Lawton, J. H., Wood, T. G. & Bignell, N. C. 1996. The diversity, abundance and biomass of termites under differing levels of disturbance in the Mbalmayo Forest Reserve, southern Cameroon. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, **351**(1335), 51–68. doi:10.1098/rstb.1996.0004

Ellwood, M. D. F. & Foster, W. A. 2004. Doubling the estimate of invertebrate biomass in a rainforest canopy. *Nature*, **429**, 549–551. doi:10.1038/nature02560.

Engelmann, G. F. 1978. *The Logic of Phylogenetic Analysis and the Phylogeny of the*

Xenarthra. Unpublished Ph.D. thesis dissertation, Columbia University, New York, 329 pp.

Engelmann, G. F. 1985. The phylogeny of the Xenarthra. Pp. 51–64 in G. G. Montgomery

(ed) *The Evolution and Ecology of Armadillos, Sloths, and Vermilinguas*. Smithsonian

Institution Press, Washington, DC.

Esteban, G. I., Nasif, N., & **Georgieff, S. M.** 2014. Cronobioestratigrafía del Mioceno

tardío-Plioceno temprano, Puerta de Corral Quemado y Villavil, Provincia de Catamarca,

Argentina. *Acta Geológica Lilloana*, **26**, 165–192. doi:10.1206/0003-0090-404.1.1

Feijoo, M. & **Parada, A.** 2017. Macrosystematics of eutherian mammals combining HTS

data to expand taxon coverage. *Molecular phylogenetics and evolution*, **113**, 76–83.

doi:10.1016/j.ympev.2017.05.004

Ferrusquía-Villafranca, I., Arroyo-Cabrales, J., Martínez-Hernández, E., Gama-

Castro, J., Ruiz-González, J., Polaco, O. J. & **Johnson, E.** 2010. Pleistocene mammals of

Mexico: A critical review of regional chronofaunas, climate change response and

biogeographic provinciality. *Quaternary International*, **217**(1–2), 53–104.

doi:10.1016/j.quaint.2009.11.036

Ferrusquía-Villafranca, I., Arroyo-Cabrales, J., Johnson, E., Ruiz-González, J.,

Martínez-Hernández, E., Gama-Castro, J., de Anda-Hurtado, P. & **Polaco, O. J.** 2017.

Quaternary mammals, people, and climate change: A view from southern North America.

Pp. 27–67 in G. Monks (ed.) *Climate change and human responses*. Springer, Dordrecht.

doi:10.1007/978-94-024-1106-5

Flynn, J. J., Guerrero, J. & **Swisher III, C. C.** 1997. Geochronology of the Honda

Group. Pp. 44–60 in R. F. Kay, R. H. Madden, R. L. Cifelli & J. J. Flynn (eds) *Vertebrate*

Paleontology in the Neotropics: The Miocene Fauna of La Venta, Colombia. Smithsonian

Institution Press, Washington.

Flower, W. H. 1883. On the arrangement of the Orders and Families of existing

Mammalia. *Proceedings of the Zoological Society of London*, **1883**, 178–186.

Foley, N. M., Springer, M. S. & **Teeling, E. C.** 2016. Mammal madness: Is the mammal

tree of life not yet resolved? *Philosophical Transactions of the Royal Society B: Biological*

Sciences, **371**(1699), 20150140. doi:10.1098/rstb.2015.0140

Forrest, M., Eronen, J. T., Utescher, T., Knorr, G., Stepanek, C., Lohmann, G. &

Hickler, T. 2015. Climate-vegetation modelling and fossil plant data suggest low

atmospheric CO2 in the late Miocene. *Climate of the Past*, **11**, 1701–1732. doi:10.5194/cp-

11-1701-2015

Fortelius, M., Eronen, J. T., Kaya, F., Tang, H., Raia, P. & **Puolamäki, K.** 2014.

Evolution of Neogene mammals in Eurasia: Environmental forcing and biotic interactions.

Annual Review of Earth and Planetary Sciences, **42**, 579–604. doi:10.1146/annurev-earth-

050212-124030

Frigola, A., Prange, M. & **Schulz, M.** 2018. Boundary conditions for the Middle Miocene

Climate Transition (MMCT v1. 0). *Geoscientific Model Development*, **11**(4), 1607–1626.

doi:10.5194/gmd-11-1607-2018

Fuster, A., Diodato, L. & **Contreras, J. H.** 2018. Dieta de *Myrmecophaga tridactyla*

(Pilosa: Myrmecophagidae) en pastizales y bosques del noreste de Santiago del Estero y su

relación con las hormigas. *Acta zoológica lilloana*, **62**(1), 1–9.

Gallo, J. A., Abba, A. M., Elizalde, L., Di Nucci, D., Ríos, T. A. & **Ezquiaga, M. C.**

2017. First study on food habits of anteaters, *Myrmecophaga tridactyla* and *Tamandua*

tetradactyla at the southern limit of their distribution. *Mammalia*, **81**(6), 601–604. doi:

10.1515/mammalia-2016-0117

Gambaryan, P. P., Zherebtsova, O. V., Perepelova, A. A. & **Platonov, V. V.** 2009. Pes

muscles and their action in giant anteater *Myrmecophaga tridactyla* (Myrmecophagidae,

Pilosa) compared with other plantigrade mammals. *Russian Journal of Theriology*, **8**(1), 1–

15. doi:10.15298/rusjtheriol.8.1.01

Gaudin, T. J. & **Branham, D. G.** 1998. The phylogeny of the Myrmecophagidae

(Mammalia, Xenarthra, Vermilingua) and the relationship of *Eurotamandua* to the

Vermilingua. *Journal of Mammalian Evolution*, **5**, 237–265.

doi:10.1023/A:1020512529767

Gaudin, T. J. & **McDonald, H. G.** 2008. Morphology-based investigations of the

phylogenetic relationships among extant and fossil xenarthrans. Pp. 24–36 in S. F. Vizcaíno

& W. J. Loughry. (eds) *The biology of the Xenarthra*. University of Florida Press,

Gainesville.

Gaudin, T. J., Hicks, P. & **Di Blanco, Y.** 2018. *Myrmecophaga tridactyla* (Pilosa:

Myrmecophagidae). *Mammalian Species*, **50**(956), 1–13. doi: 10.1093/mspecies/sey001

Ge, D., Wen, Z., Xia, L., Zhang, Z., Erbajeva, M., Huang, C. & **Yang, Q.** 2013.

Evolutionary history of lagomorphs in response to global environmental change. *PLoS One*,

8(4), e59668. doi:10.1371/journal.pone.0059668

Gibb, G. C., Condamine, F. L., Kuch, M., Enk, J., Moraes-Barros, N., Superina, M.,

Poinar, H. N. & **Delsuc, F.** 2015. Shotgun mitogenomics provides a reference

phylogenetic framework and timescale for living xenarthrans. *Molecular Biology and*

Evolution, **33**(3), 621–642. doi:10.1093/molbev/msv250

Gray, J. E. 1825. An outline of an attempt at the disposition of Mammalia into tribes and

families, with a list of genera apparently pertaining to each tribe. *Annals of Philosophy*, **10**,

337–344.

Grimmer, F., Dupont, L., Lamy, F., Jung, G., González, C. & **Wefer, G.** 2018. Early

Pliocene vegetation and hydrology changes in western equatorial South America. *Climate*

of the Past, **14**(11), 1739–1754. doi:10.5194/cp-14-1739-2018

Guénard, B., Perrichot, V. & **Economo, E. P.** 2015. Integration of global fossil and

modern biodiversity data reveals dynamism and stasis in ant macroecological patterns.

Journal of Biogeography, **42**(12), 2302–2312. doi:10.1111/jbi.12614

Guerrero, J. 1997. Stratigraphy, sedimentary environments, and the Miocene uplift of the

Colombian Andes. Pp. 15–43 in R. F. Kay, R. H. Madden, R. L. Cifelli & J. J. Flynn (eds)

Vertebrate Paleontology in the Neotropics: The Miocene Fauna of La Venta, Colombia.

Smithsonian Institution Press, Washington, DC.

Halffter, G. & **Morrone, J. J.** 2017. An analytical review of Halffter's Mexican transition

Zone, and its relevance for evolutionary biogeography, ecology and biogeographical

regionalization. *Zootaxa*, **4226**(1), 1–46. doi:10.1007/s13127-013-0148-0

Halliday, T. J. D., Upchurch, P. & **Goswami, A.** 2016. Eutherians experienced elevated

evolutionary rates in the immediate aftermath of the Cretaceous-Palaeogene mass

extinction. *Proceedings of the Royal Society B: Biological Sciences*, **283**(1833), 20153026.

doi:10.1098/rspb.2015.3026

Hayssen, V. 2011. *Tamandua tetradactyla* (Pilosa: Myrmecophagidae). *Mammalian*

Species, 43(875), 64–74. doi:10.1644/875.1

Hayssen, V., Miranda, F. & **Pasch, B.** 2012. *Cyclopes didactylus* (Pilosa: Cyclopedidae).

Mammalian Species, 44(895), 51–58. doi: 10.1644/895.1

Henrot, A. J., Utescher, T., Erdei, B., Dury, M., Hamon, N., Ramstein, G., Krapp, M.,

Herold, N., Goldner, A., Favre, E. & **Munhoven, G.** 2017. Middle Miocene climate and

vegetation models and their validation with proxy data. *Palaeogeography,*

Palaeoclimatology, Palaeoecology, **467**, 95–119. doi:10.1016/j.palaeo.2016.05.026

Herold, N., Huber, M., Greenwood, D. R., Müller, R. D. & **Seton, M.** 2011. Early to

middle Miocene monsoon climate in Australia. *Geology*, **39**(1), 3–6. doi:10.1130/G31208.1

Hirschfeld, S. E. 1976. A new fossil anteater (Edentata, Mammalia) from Colombia, SA

and evolution of the Vermilingua. *Journal of Paleontology*, **50**, 419–432.

Hölldobler, B. & **Wilson EO.** 1990. *The Ants*. Cambridge, MA: Harvard University Press.

732 pp.

Hossotani C. M. S., Ragusa-Netto J. & **Luna HS.** 2017. Skull morphometry and vault

sutures of *Myrmecophaga tridactyla* and *Tamandua tetradactyla*. *Iheringia, Série Zoologia*,

107, e2017038. doi:10.1590/1678-4766e2017038.

Huntley, B. & **Webb III, T.** 1988. *Vegetation History* (Vol. 7). Springer Science &

Business Media. Kluwer, Dordrecht, 803 pp.

Huxley, J. S. 1958. Evolutionary processes and taxonomy with special reference to grades.

Uppsala Universitets Arsskrift, 6, 21–39.

Iglesias A., Artabe, A. E. & **Morel, E. M.** 2011. The evolution of Patagonian climate and

vegetation from the Mesozoic to the present. Biological *Journal of the Linnean Society*,

103, 409–422. doi: 10.1111/j.1095-8312.2011.01657.x

Illiger, J. K. W. 1811. *Prodromus systematis mammalium et avium additis terminis*

zoographicis triusque classis, eorumque versione germanica. Berlin, 301 pp.

Jaramillo, C. 2018. Evolution of the Isthmus of Panama: Biological, paleoceanographic,

and paleoclimatological implications. Pp. 323–338 in C. Hoorn & A. Antonelli (eds)

Mountains, climate and biodiversity. John Wiley & Sons, Oxford.

Jiménez, N. L., Di Blanco, Y. E. & **Calcaterra, L. A.** 2018. Ant diversity in the diet of

giant anteaters, *Myrmecophaga tridactyla* (Pilosa: Myrmecophagidae), in the Iberá Nature

Reserve, Argentina. *Mastozoologia Neotropical*, **25**(2) 1–14.

doi:10.31687/saremMN.18.25.2.0.16

Kay, R. F. & **Madden, R. H.** 1997. Mammals and rainfall: Paleoecology of the Middle

Miocene at La Venta (Colombia, South America). *Journal of Human Evolution*, **32**(2–3),

161–199. doi: 10.1006/jhev.1996.0104

Kay R. F., Vizcaíno S. F. & **Bargo M, S.** 2012. A review of the paleoenvironment and

paleoecology of the Miocene Santa Cruz Formation. Pp. 331–365 in S. F. Vizcaíno, R. F.

Kay & M. S. Bargo (eds) *Early Miocene Paleobiology in Patagonia: High-Latitude*

Paleocommunities of the Santa Cruz Formation. Cambridge University Press, New York.

Keller, L. & **E. Gordon.** 2009. *The Lives of Ants*. Oxford University Press, New York, 273

pp.

Kraglievich, L. 1934. *La antigüedad pliocena de las faunas de Monte Hermoso y*

Chapadmalal, deducidas de su comparación con las que le procedieron y sucedieron.

Imprenta El Siglo Ilustrado, Montevideo, 136 pp.

Kraglievich, L. 1940. Dos nuevas especies fósiles de oso hormigueros. Pp. 631–636 in A.

J. Torcelli and C. A. Marelli (eds) *Obras de geología y paleontología* (Vol. 3). Ministerio

de Obras Públicas de la Provincia de Buenos Aires, La Plata.

LaPolla, J. S., Dlussky, G. M. & **Perrichot, V.** 2013. Ants and the fossil record. *Annual*

review of entomology, **58**, 609–630. doi:10.1111/syen.12090

Latorre, C., Quade, J. & **McIntosh, W. C.** 1997. The expansion of C4 grasses and global

change in the late Miocene: Stable isotope evidence from the Americas. *Earth and*

Planetary Science Letters, **146**(1–2), 83–96. doi:10.1016/S0012-821X(96)00231-2

Lohmann, G., Butzin, M. & **Bickert, T.** 2015. Effect of vegetation on the Late Miocene

ocean circulation. *Journal of Marine Science and Engineering*, **3**(4), 1311–1333.

doi:10.3390/jmse3041311

Lubin, Y. D. & **Montgomery, G. G.** 1981. Defenses of *Nasutitermes* termites (Isoptera,

Termitidae) against *Tamandua* anteaters (Edentata, Myrmecophagidae). *Biotropica*, **13**, 66–

76. doi: 10.2307/2387872

MacFadden, B. J. 2000. Cenozoic mammalian herbivores from the Americas:

Reconstructing ancient diets and terrestrial communities. *Annual Review of Ecology and*

Systematics, **31**(1), 33–59. doi:10.1146/annurev.ecolsys.31.1.33

McDonald, H. G. 2005. Paleoecology of extinct xenarthrans and the Great American

Biotic Interchange. *Bulletin of the Florida State Museum of Natural History*, **45**, 313–333.

McDonald, H. G. & **De Iuliis, G.** 2008. Fossil history of sloths. Pp. 39–55 in S. F.

Vizcaíno & W. J. Loughry (eds) *The biology of the Xenarthra*. University Press of Florida,

Gainesville.

McDonald, H. G., Vizcaíno, S. F., & **Bargo, M. S.** 2008. Skeletal anatomy and the fossil

history of the Vermilingua. Pp. 257–268 in S. F. Vizcaíno & W. J. Loughry (eds*) The*

biology of the Xenarthra. University Press of Florida, Gainesville.

McKenna, M. C. & **Bell S. K.** 1997. *Classification of mammals above the species level*

(Vol. XIII). Columbia University Press, New York, 631 pp.

McNab, B. K. 1984. Physiological convergence amongst ant-eating and termite-eating

mammals. *Journal of Zoology*, **203**(4), 485–510. doi:10.1111/j.1469-7998.1984.tb02345.x

McNab, B. K. 1985. Energetics, population biology, and distribution of xenarthrans, living

and extinct. Pp. 219–232 in G. G. Montgomery (ed.) *The evolution and ecology of*

armadillos, sloths, and vermilinguas. Smithsonian Institution Press, Washington, DC.

Medri, I. M., Mourao, G. D. M. & **Harada, A. Y.** 2003. Dieta de tamanduá-bandeira

(*Myrmecophaga tridactyla*) no Pantanal da Nhecolândia, Brasil. *Edentata*, **5**, 29–34.

Miranda, F., Veloso, R., Superina, M. & **Zara, F. J.** 2009. Food habits of wild silky

anteaters (*Cyclopes didactylus*) of São Luis do Maranhão, Brazil. *Edentata*, **2009**(10), 1–6.

doi:10.1896/020.010.0109

Miranda, F. R., Casali, D. M., Perini, F. A., Machado, F. A. & **Santos, F. R.** 2017.

Taxonomic review of the genus *Cyclopes* Gray, 1821 (Xenarthra: Pilosa), with the

revalidation and description of new species. *Zoological Journal of the Linnean Society*,

183(3), 687–721. doi:10.1093/zoolinnean/zlx079

Montgomery, G.G. 1985a. Impact of vermilinguas (*Cyclopes*, *Tamandua*: Xenarthra =

Edentata) on arboreal ant populations. Pp. 351–363 in G.G. Montgomery (ed) *Ecology and*

evolution of sloths, anteaters and armadillos (Mammalia, Xenarthra = Edentata).

Smithsonian Institution Press, Washington, D.C.

Montgomery, G. G. 1985b. Movements, foraging and food habits of the four extant

species of neotropical vermilinguas (Mammalia; Myrmecophagidae). Pp. 365–367 in G.G.

Montgomery (ed) *The Evolution and Ecology of Armadillos, Sloth, and Vermilinguas*.

Smithsonian Institution Press, Washington, D.C.

Morales-Sandoval, V. H. 2010. Caracterización nutricional de la dieta de Tamandua

mexicana en el Zoológico "Miguel Álvarez del Toro" (ZooMAT) Chiapas, México.

Edentata, **11**(1), 44–49. doi:10.1896/020.011.0108

Morley, R. J. 2011. Cretaceous and Tertiary climate change and the past distribution of megathermal rainforests. Pp. 1–34 in M. B. Bush & J. R. Flenley (eds) *Tropical rainforest responses to climatic change*. Springer, Berlin, Heidelberg.

Moscato, D. A. & **Jasinski, S. E.** 2016. First record of fossil chelydrid and trionychid turtles (testudines) from the Pleistocene of Sonora, Mexico. Pp. 163–168 in R. M. Sullivan & G. L. Spencer (eds*) Fossil Record 5: Bulletin*. New Mexico Museum of Natural History & Science, Albuquerque.

Naples, V. L. 1999. Morphology, evolution and function of feeding in the giant anteater (*Myrmecophaga tridactyla*). *Journal of Zoology*, **249**(1), 19–41. doi:10.1111/j.1469–7998.1999.tb01057.x

Navarrete, D. & **Ortega, J.** 2011. *Tamandua mexicana* (Pilosa: Myrmecophagidae). *Mammalian Species*, **43**(874), 56–63. doi:10.1644/874.1

Norman, J. E. & **Ashley, M. V.** 1994. Molecular systematics of the xenarthrans. American Society of Mammalogists, 75th Anniversary Meeting, Abstract 271A, Washington, DC.

Oliva, C., Zurita, A. E., Dondas, A. & **Scillato-Yané, G. J.** 2010. Los Glyptodontinae (Xenarthra, Glyptodontidae) del Piso/Edad Chapadmalalense (Plioceno tardío): Revisión y aportes a su conocimiento. *Revista mexicana de ciencias geológicas*, **27**(1), 112–120.

O'Dea, A., Lessios, H. A., Coates, A. G., Eytan, R. I., Restrepo-Moreno, S. A., Cione, A. L., Collins, S. L., Queiroz, A., Farris, D. W., Norris, R. D., Stallard, R. F., Woodburne, M. O., Aguilera, O., Aubry, M. P., Berggren, W. A., Budd, A. F., Cozzuol, M. A., Coppard, S. E., Duque-Caro. H., Finnegan. S., Gasparini, G. M., Grossman, E. L., Johnson, K. G., Keigwin, L. D., Knowlton, N., Leigh, E. G., Pingel, J.

S., Marko, P. B., Pyenson, B. M., Rachello-Dolmen, P. G., Soibelzon, E., Soilbezon, L.,

Todd, J. A. Vermeij, G. J. & **Stallard, R. F.** 2016. Formation of the Isthmus of Panama.

Science advances, **2**(8), e1600883. doi:10.1126/sciadv.1600883

Ortiz-Jaureguizar, E. & **Cladera, G. A.** 2006. Paleoenvironmental evolution of southern

South America during the Cenozoic. *Journal of Arid Environments*, **66**(3), 498–532.

doi:10.1016/j.jaridenv.2006.01.007

Palazzesi, L., Barreda, V. D., Cuitiño, J. I., Guler, M. V., Tellería, M. C. & **Santos, R.**

V. 2014. Fossil pollen records indicate that Patagonian desertification was not solely a

consequence of Andean uplift. *Nature Communications*, **5**, 3558. doi:10.1038/ncomms4558

Pascual, R. & **Ortiz-Jaureguizar, E. 1990.** Evolving climates and mammal faunas in

Cenozoic South America. *Journal of Human Evolution*, **19**, 23–60. doi:10.1016/B978-0-12-

260345–7.50006-0

Patterson, B., Segall, W., Turnbull, W. D. & **Gaudin, T. J.** 1992. The ear region in

xenarthrans (=Edentata: Mammalia). Part II. Pilosa (sloths and anteaters), palaeanodonts,

and a miscellany. *Fieldiana Geology*, *n.s. 24*, 1–79.

Plotnick, R. E. & **Wagner, P. J.** 2006. Round up the usual suspects: Common genera in

the fossil record and the nature of wastebasket taxa. *Paleobiology*, **32**(1), 126–146.

doi:10.1666/0094–8373(2006)032[0126:RUTUSC]2.0.CO;2

Pound, M. J. 2012. *Middle to Late Miocene terrestrial biota and climate*. Unpublished

Ph.D. thesis, University of Leeds, Leeds, 416 pp.

Pound, M. J., Haywood, A. M., Salzmann, U. & **Riding, J.** B. 2012. Global vegetation

dynamics and latitudinal temperature gradients during the Mid to Late Miocene (15.97–

5.33 Ma). *Earth-Science Reviews*, **112**(1–2), 1–22. doi:10.1016/j.earscirev.2012.02.005

Quattrocchio M. E., Volkheimer, W., Borromei, A. M. & **Martínez, M. A.** 2011.

Changes of the palynobiotas in the Mesozoic and Cenozoic of Patagonia: A review.

Biological Journal of the Linnean Society, **103**, 380–396. doi: 10.1111/j.1095-

8312.2011.01652.x

Raigemborn, M. S., Krapovickas, V., Beilinson, E., Peral, L. E. G., Zucol, A. F.,

Zapata, L. & **Sial, A. N.** 2018. Multiproxy studies of Early Miocene pedogenic calcretes in

the Santa Cruz Formation of southern Patagonia, Argentina indicate the existence of a

temperate warm vegetation adapted to a fluctuating water table. *Palaeogeography,*

Palaeoclimatology, Palaeoecology, 500:1–23. doi:10.1016/j.palaeo.2018.03.037

Redford, K. H. 1985. Feeding and food preference in captive and wild giant anteaters

(*Myrmecophaga tridactyla*). *Journal of Zoology*, **205**(4), 559–572. doi:10.1111/j.1469–

7998.1985.tb03544.x

Redford, K. H. 1987. Ants and termites as food: Patterns of mammalian myrmecophagy.

Pp. 349–399 in H. H. Genoways (ed.) *Current mammalogy*. Plenum Press, New York.

Reiss, K. Z. 2001. Using phylogenies to study convergence: the case of the anteating

mammals. *American Zoologist*, **41**(3), 507–525. doi:10.1093/icb/41.3.507

Ride, W. D. L., Cogger, H. G., Dupuis, C., Kraus, O., Minelli, A., Thompson, F. C. &

Tubbs, P. K. 1999. *International Code of Zoological Nomenclature*. 4th ed. International

Trust for Zoological Nomenclature and the Natural History Museum, London, 106 pp.

Riggs, E. S. 1928. Work accomplished by the Field Museum Paleontological Expeditions

to South America. *Science*, **67**(1745), 585–587.

Roberts, N. J., Barendregt, R. W. & **Clague, J. J.** 2018. Pliocene and Pleistocene

chronostratigraphy of continental sediments underlying the Altiplano at La Paz, Bolivia.

Quaternary Science Reviews, **189**, 105–126. doi:10.1016/j.quascirev.2018.03.008

Rodrigues, F. H., Medri, I. M., De Miranda, G. H., Alves, C. & **Mourão, G.** 2008.

Anteater behavior and ecology. Pp. 257–268 in S. F. Vizcaíno & W. J. Loughry (eds) *The*

biology of the Xenarthra. University Press of Florida, Gainseville.

Rovereto, C. 1914. Los estratos araucanos y sus fósiles. *Anales del Museo Nacional de*

Historia Natural de Buenos Aires, **25**,1–247.

Salas-Gismondi, R., Flynn, J. J., Baby, P., Tejada-Lara, J. V., Wesselingh, F. P. &

Antoine, P. O. 2015. A Miocene hyperdiverse crocodylian community reveals peculiar

trophic dynamics in proto-Amazonian mega-wetlands. *Proceedings of the Royal Society B:*

Biological Sciences, **282**(1804), 20142490. doi:10.1098/rspb.2014.2490

Salzmann, U., Williams, M., Haywood, A. M., Johnson, A. L., Kender, S. &

Zalasiewicz, J. 2011. Climate and environment of a Pliocene warm world.

Palaeogeography, Palaeoclimatology, Palaeoecology, **309**(1–2), 1–8.

doi:10.1016/j.palaeo.2011.05.044

Sandoval-Gómez, V. E., Ramírez-Chaves, H. E. & **Marín, D.** 2012. Registros de

hormigas y termitas presentes en la dieta de osos hormigueros (Mammalia:

Myrmecophagidae) en tres localidades de Colombia. *Edentata*, **13**(1), 1–10.

doi:10.5537/020.013.0104

Scillato-Yané, G. J. & **Carlini, C. C.** 1998. Nuevos Xenarthra del Friasense (Mioceno medio) de Argentina. *Studia Geológica Salmanticensia*, **34**: 43–67.

Sereno, P. C. 1991. Basal archosaurs: Phylogenetic relationships and functional implications. *Journal of Vertebrate Paleontology*, **11**(S4), 1–53. doi:10.1080/02724634.1991.10011426

Seoane, F. D., Roig Juñent, S. & **Cerdeño, E.** 2017. Phylogeny and paleobiogeography of Hegetotheriidae (Mammalia, Notoungulata). *Journal of Vertebrate Paleontology*, **37**(1), e1278547. doi:10.1080/02724634.2017.1278547

Spradley, J. P., Glazer, B. J. & **Kay, R. F.** 2019. Mammalian faunas, ecological indices, and machine-learning regression for the purpose of paleoenvironment reconstruction in the Miocene of South America. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **518**, 155–171. doi:10.1016/j.palaeo.2019.01.014

Shaw C. A. & **McDonald H. G.** 1987. First record of giant anteater (Xenarthra: Myrmecophagidae) in North America. *Science*, **236**, 186–188. doi:10.1126/science.236.4798.186

Šímová, I. & **Storch, D.** 2017. The enigma of terrestrial primary productivity: Measurements, models, scales and the diversity-productivity relationship. *Ecography*, **40**(2), 239–252. doi:10.1111/ecog.02482

Sniderman, J. K., Woodhead, J. D., Hellstrom, J., Jordan, G. J., Drysdale, R. N., Tyler, J. J. & **Porch, N.** 2016. Pliocene reversal of late Neogene aridification. *Proceedings of the National Academy of Sciences*, **113**(8), 1999–2004. doi:10.1073/pnas.1520188113

Tobin, J. E. 1995. Ecology and diversity of tropical forest canopy ants. Pp. 129–147. In M.

D. Lowman & N. M. Nadkarni (eds.) *Forest Canopies*. Academic Press, New York.

Toby Pennington, R., Prado, D. E. & **Pendry, C. A.** 2000. Neotropical seasonally dry

forests and Quaternary vegetation changes. *Journal of Biogeography*, **27**(2), 261–273.

doi:10.1046/j.1365–2699.2000.00397.x

Toledo, N., Bargo, M. S., Vizcaíno, S. F., De Iuliis, G. & **Pujos, F.** 2017. Evolution of

body size in anteaters and sloths (Xenarthra, Pilosa): Phylogeny, metabolism, diet and

substrate preferences. *Earth and Environmental Science Transactions of the Royal Society*

of Edinburgh, **106**(4), 289–301. doi:10.1017/S1755691016000177

Thompson, J. N. 1994. *The Coevolutionary Process*. University of Chicago Press, Chicago

IL., 360 pp.

Wiens, J. J. & **Donoghue, M. J.** 2004. Historical biogeography, ecology and species

richness. *Trends in Ecology & Evolution*, **19**(12), 639–644. doi:10.1016/j.tree.2004.09.011

Wiens, J. J. & **Graham, C. H.** 2005. Niche conservatism: Integrating evolution, ecology,

and conservation biology. *Annual Reviews in Ecology, Evolution and Systematics*, **36**, 519–

539. doi: 10.1146/annurev.ecolsys.36.102803.095431

Williams, J. W., Shuman, B. N., Webb III, T., Bartlein, P. J. & **Leduc, P. L.** 2004.

Late☐Quaternary vegetation dynamics in North America: Scaling from taxa to biomes.

Ecological Monographs, **74**(2), 309–334. doi:10.1890/02–4045

Williams, J. 2009. Quaternary vegetation distributions. Pp. 856–862 in V. Gornitz (ed.)

Encyclopedia of Paleoclimatology and Ancient Environments. Springer, Dordrecht.

Wood, B. & **Lonergan, N.** 2008. The hominin fossil record: Taxa, grades and clades.

Journal of Anatomy, **212**(4), 354–376. doi:10.1111/j.1469–7580.2008.00871.x

Woodburne, M. O. 2010. The Great American Biotic Interchange: Dispersals, tectonics,

climate, sea level and holding pens. *Journal of Mammalian Evolution*, **17**(4), 245–264.

doi:10.1007/s10914-010-9144-8

Zavala, C. & **Navarro, E.** 1993. Depósitos fluviales en la Formación Monte Hermoso

(Plioceno inferior-medio), Provincia de Buenos Aires. *Actas*, **5**(2), 236–244.

Zachos, J., Pagani, M., Sloan, L., Thomas, E. & **Billups, K.** 2001. Trends, rhythms, and

aberrations in global climate 65 Ma to present. *Science*, **292**(5517), 686–693.

doi:10.1126/science.1059412

Zachos, J. C., Dickens, G. R. & **Zeebe, R. E.** 2008. An early Cenozoic perspective on

greenhouse warming and carbon-cycle dynamics. *Nature*, **451**(7176), 279.

Zimicz, N., Payrola, P. & **del Papa, C.** 2018. New, Late Miocene mammalian assemblage

from the Palo Pintado Formation (Northwestern Argentina). *Journal of South American*

Earth Sciences, **81**, 31–44. doi:10.1016/j.jsames.2017.11.003

1358

Figure captions

Figure 1. Geographical and stratigraphic provenance of the skull VPPLT 975 of the new

taxon described here and the holotype of *Neotamandua borealis* (Hirschfeld 1976). **A**,

location of the Department of Huila in Colombia; **B**, location of the fossil area of interest,

i.e. northern of La Venta area, in the Department of Huila (small rectangle); **C**, location of

the fossil site (black star), near the La Victoria town; **D**, stratigraphic scheme of Guerrero

(1997) for La Venta area, with approximate stratigraphic provenance of VPPLT 975 and

the holotype of *N. borealis*.

Figure 2. Cranial measurements used in this work. All are based on Hossotani et al. (2017).

Nomenclatural modifications from these measurements are shown in the section of

Anatomical Abbreviations. **A**, skull of *Tamandua* in dorsal view; **B**, the same skull in

ventral view. **Abbreviations: FL**, frontal length; **GSL**, greatest skull length; **MBW**,

maximum braincase width; **ML**, maxilla length; **NL**, nasal length; **NW**, nasal width; **PL**,

parietal length.

Figure 3. Holotypic skull (VPPLT 975) of Gen. et sp. nov. **A**, dorsal view; **B**, right lateral

view; **C**, ventral view; **D**, left lateral view; **E**, anatomical drawing in dorsal view; **F**,

anatomical drawing in right lateral view. **Abbreviations: fr,** frontals; **ji,** jugal insertion; **la,**

lacrimal; **mx,** maxilla; **na,** nasals; **or,** orbit. Scale bar equal to 30 mm.

Figure 4. Two very informative postcranial bones of the holotype (UCMP 39847) of

Neotamandua borealis (Hirschfeld 1976). **A**, right astragalus, dorsal view; **B**, right

astragalus, ventral view; **C**, left calcaneum, dorsal view; **D**, left calcaneum, lateral view; **E**,

anatomical drawing of the astragalus in dorsal view; **F**, anatomical drawing of the

astragalus in ventral view; **G**, anatomical drawing of the calcaneum in dorsal view; **H**,

anatomical drawing of the calcaneum in lateral view. **Abbreviations: af**, calcaneal

accessory facet; **ct**, calcaneal tuber; **ef**, ectal facet; **ff**, fibular facet; **h**, astragalar head; **lr**,

lateral ridge; **lt**, lateral trochlea; **mt**, medial trochlea; **sf**, sustentacular facet. Scale bar equal

to 20 mm.

Figure 5. Epitype (FMNH P14419) of *Neotamandua conspicua*. **A**, dorsal view; **B**, right

lateral view; **C**, ventral view; **D**, left lateral view; **E**, right hemimandible; **F**, left

hemimandible; **G**, anatomical drawing in dorsal view; **H**, anatomical drawing in ventral

view; **I**, anatomical drawing in left lateral view. **Abbreviations: ab**, auditory bullae; **fr**,

frontals; **j**, jugal; **mx**, maxilla; **na**, nasals; **oc**, occipital condyles; **pal**, palatines; **ptb**,

pterygoid bullae; **pte**, pterygoids; **szp**, squamosal zygomatic process. Scale bar equal to 80

mm.

Figure 6. Reconstruction the external appearance in life of Gen. et sp. nov. (close-up view).

In the background, individuals of the macraucheniid *Theosodon* (left) and the alouattine

Stirtonia (upper right corner) in the tropical forest of La Venta, late Middle Miocene of

Colombia.

Figure 7. Geographic and chronological distribution of the myrmecophagid fossil records

during the Late Cenozoic. Note the only two fossil records of these xenartrans outside

South America in the Pleistocene of southern and northern Mexico (*Tamandua* sp. and

Myrmecophaga tridactyla, respectively). Based on information compiled by McDonald et

al. (2008). Original references in the same work and, largely, in the main text here.

Figure 8. Chronological collation of data on: **A**, biochrons of the myrmecophagid genera or

questionable grouping (horizontal solid bars and dashed line); **B**, distribution of the highest

latitudinal fossil records (northern and/or southern) of myrmecophagids (horizontal solid

bars) and approximate, chronologically discrete latitudinal ranges of tropical rainforest plus

tropical and subtropical dry broadleaf forest (i.e. frost-free areas [mean annual temperatures

higher than 15°C] with significant rainfall, at least seasonally; large vertical rectangles); **C**,

general trend curve of global temperature and climatic episodes during the Late Cenozoic:

a, early Neogene warm recovery, including the thermal peak in the late Early-early Middle

Miocene known as Middle Miocene Climatic Optimum or MMCO; **b**, Middle Miocene

climatic transition; **c**, late Middle-Late Miocene cooling; **d**, Early Pliocene warming; **e**,

Late Pliocene-Pleistocene cooling and glaciations. The vertical dashed line indicates the

time of complete formation of the Panama Land Bridge, which represented thereafter a

fundamentally continuous physical connection between South- and North America.

Palaeocological data used for the plot in 'B' is from the following references: Huntley &

Webb III (1988); Toby Pennington et al. (2000); Williams et al. (2004); Williams (2009);

Chan et al. (2011); Morley (2011); Kay et al. (2012); Pound (2012); Pound et al. (2012);

Forrest et al. (2015); Lohmann et al. (2015); Dowsett et al. (2016); Sniderman et al. (2016);

Henrot et al. (2017); Frigola et al. (2018). The temperature curve in 'C' is based on Zachos

et al. (2001, 2008) and it is reproduced with permission.

1421

1422

1423

1424

1425

1426

1427

1428

1429

TABLES

Table 1. Cranial measurements (in mm) for the holotype of Gen. et sp. nov. and other myrmecophagid species

Measurement	Gen. et sp. nov. [a]	*P. rothi* [b]	*N. conspicua* [c]	*T. tetradactyla* [d]	*M. tridactyla* [e]
GSL	80.7*	77*	210**	125.7	327.5
NL	30.9	☐	~110	38.2	127.5
NW	4.8	☐	11.1	7.7	14.2
FL	☐	~27	☐	53.1	143.9
MBW	☐	36	~51	42.2	60.4
PL	☐	~20	☐	20.1	26

a. VPPLT 975 (holotype)
b. YPM-15267
c. FMNH P14419
d. Mean of a sample of (sub) adults, n = 8 (Appendix 2, Supplementary material)
e. Mean of a sample of (sub) adults, n = 10 (Appendix 2, Supplementary material)

*Incomplete skull. Rough estimate of GSL for Gen. et sp. nov. from a simple linear model based on some skull measurements for *Tamandua* (see Supplementary Material) is equal to 118.6 mm.

**Fractured rostrum

Table 2. Comparison of some postcranial measurements (in mm) between *N. borealis* and *N. greslebini*.

Measurement	*N. borealis*	*N. greslebini*
Maximum distal width of the radius	16.5	29
Maximum proximal width of the tibia	29.5	46
Proximo-distal length of the astragalus	41.5	55
Maximum width of the astragalus	16.4	20.5

1430

1431

Table 3. Comparison of some postcranial measurements (in mm) between *N. borealis* and

Neotamandua sp.

Measurement	*N. borealis*	*Neotamandua* sp.
Maximum distal width of the radius	16.5	35
Maximum width of the tibial articulation of the astragalus	20	24
Maximum width of the calcaneum	20	~21

1432

1433

Table 4. Distribution of some postcranial characters of species referred to *Neotamandua* and other myrmecophagid taxa. The

	N. borealis	*N. greslebini*	*Neotamandua* sp.	*Protamandua*	*Tamandua*	*Myrmecophaga*
Relative body size	Medium to large	Large	Very large	Small	Medium	Very large
Character						
1. Arrangement of the radial ridges	Parallel ridges	Parallel ridges	Parallel ridges	NA	Distally convergent ridges	Parallel ridges
2. Rotated appearance of the distal radius	Absent	Absent	Absent	Absent	Present	Absent
3. Tibial mid-section*	Sub-rounded to gently triangular	Sub-rounded to gently triangular	NA	NA	Rounded	Strongly triangular
4. Arrangement of the ectal and sustentacular facets in the astragalus*	Largely separated, but with an incipient connection	Largely separated, but with an incipient connection	NA	Fully separated	Fully separated	Widely connected
5. Tendinous groove in the lateral side of the calcaneum*	Short	Short	NA	Absent	Absent	Long

characters marked with asterisk contain potentially diagnostic character states (synapomorphies?) for *Neotamandua*.

Table 5. Taxonomic breadth in diet (genus level) of extant genera of Vermilingua and habitat preference of their MDCs (genera or species groups considered main dietary components). Key for the references: (1) Best & Harada (1985); (2) Fuster et al. 2018; (3) Gallo et al. (2017); (4) Gaudin et al. (2018); (5) Gómez et al. (2012); (6) Hayssen (2011); (7) Hayssen et al. (2012); (8) Jiménez et al. (2018); (9) Lubin & Montgomery (1981); (10) Medri et al. (2003); (11) Miranda et al. (2009); (12) Montgomery (1981); (13) Montgomery (1985a); (14) Montgomery (1985b); (15) Morales-Sandoval (2010); (16) Navarrete & Ortega (2011); (17) Redford (1985); (18) Rodrigues et al. (2008); (19) Sandoval-Gómez et al. (2012).

Extant anteater genera	Termite genera	Ant genera	Some MDCs	Habitat preference of MDCs	References
Cyclopes	0	8	*Camponotus* *Dolichoderus* *Solenopsis*	Rainforest, seasonally dry tropical forest; trees	1, 7, 11, 13, 19
Tamandua	7	17	*Nasutitermes* *Camponotus* *Crematogaster* *Solenopsis*	Seasonally dry tropical forest, rainforest; trees and ground	3, 6, 9, 12, 14, 15, 16
Myrmecophaga	8	31	*Cornitermes* *Syntermes* *Atta* *Solenopsis* *Camponotus* *Acromyrmex* *Pheidole*	savannah, grassland; ground	2, 3, 4, 5, 8, 10, 14, 17, 18, 19

Figure 1

Lithology	Unit
	Cerbatana cglm.
	Unit between
	Tatacoa ssb.
	Unit between
	Chunchullo ssb.
	Unit between
	Cerro Gordo ssb.
	Unit below

← *Neotamandua borealis*

← VPPLT 975

ssb.   sandstone beds
cglm.   conglomerate

Fine grained sedimentites (mudstones, claystones)

Massive and geographically extended sandstones

Clast-supported conglomerate

★ Fossil locality (specimen VPPLT 975)

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Early Miocene - *Protamandua*

Middle Miocene - *Neotamandua*

Middle Miocene - Gen. nov.

Middle Miocene - *Neotamandua*?

Late Miocene - *Neotamandua*

Late Miocene - '*Neotamandua*'

Early Pliocene - *Neotamandua*

Early Pliocene - *Myrmecophaga*

Early Pleistocene - *Myrmecophaga*

(Late?) Pleistocene - *Myrmecophaga*

Late Pleistocene - *Tamandua*

Figure 8