Ingested insecticide to control *Aedes aegypti*: Developing a novel dried attractive toxic sugar bait device for intra-domiciliary control

Rachel Sippy[1,2,†], Galo E. Rivera[3,a,†], Valeria Sanchez[1,4], Froilán Heras Heras[1,4], Bianca Morejón[3,b], Efraín Beltrán Ayala[4], Robert S. Hikida[5], María A. López-Latorre[3,c], Alex Aguirre[3], Anna M. Stewart-Ibarra[1,5,d,*], David A. Larsen[6,*], Marco Neira[3,*]

[1] Institute for Global Health & Translational Science, SUNY-Upstate Medical University, Syracuse, NY, USA

[2] Department of Geography, University of Florida, Gainesville, FL, USA

[3] Center for Research on Health in Latin America, Pontificia Universidad Católica del Ecuador, Quito, Ecuador

[4] Unidad Académica de Ciencias Químicas y de la Salud, Universidad Técnica de Machala, Machala, Ecuador

[5] Department of Medicine and Department of Public Health and Preventative Medicine, SUNY-Upstate Medical University, Syracuse, NY, USA

[6] Department of Public Health, Syracuse University, Syracuse, NY, USA

[a] Current affiliation: Vector Biology Group, Max Plank Institute for Infection Biology, Berlin, Germany

[b] Current affiliation: Biology Division, College of Arts and Sciences, Kansas State University, Manhattan, KS, USA

[c] Current affiliation: Medical School, College of Health Sciences, Universidad de las Américas, Quito, Ecuador

[d] Current address: InterAmerican Institute for Global Change Research (IAI), Montevideo, Uruguay

[†] These authors contributed equally to the work

*Corresponding authors:
 Anna Stewart: astewart@dir.iai.int
 David Larsen: dalarsen@syr.edu
 Marco Neira: mvneira@puce.edu.ec

Abstract (350 words)

Background: Illnesses transmitted by *Aedes aegypti* (Linnaeus, 1762) such as dengue, chikungunya and Zika comprise a considerable global burden; mosquito control is the primary public health tool to reduce disease transmission. Current interventions are inadequate and insecticide resistance threatens the effectiveness of these options. Dried attractive bait stations (DABS) are a novel mechanism to deliver insecticide to *Ae. aegypti*. The DABS are a high-contrast 28 inch^2 surface coated with dried sugar-boric acid solution. *Ae. aegypti* are attracted to DABS by visual cues only, and the dried sugar solution elicits an ingestion response from *Ae. aegypti* landing on the surface. The study presents the development of the DABS and tests of their impact on *Ae. aegypti* mortality in the laboratory and a series of semi-field trials. *Methods*: We conducted multiple series of laboratory and semi-field trials to assess the survivability of *Ae. aegypti* mosquitoes exposed to the DABS. For laboratory experiments we assessed the lethality, the killing mechanism, and the shelf life of the device through controlled experiments. In the semi-field trials, we released laboratory-reared female *Ae. aegypti* into experimental houses typical of peri-urban tropical communities in South America in three trial series with six replicates each. Laboratory experiments were conducted in Quito, Ecuador, and semi-field experiments were conducted in Machala, Ecuador – an area with abundant wild populations of *Ae. aegypti* and endemic arboviral transmission. *Results*: In the laboratory, complete lethality was observed after 48 hours regardless of physiological status of the mosquito. The killing mechanism was determined to be through ingestion, as the boric acid disrupted the gut of the mosquito. In experimental houses, total mosquito mortality was greater in the treatment house for all series of experiments (p<0.0001). *Conclusions*: The DABS devices were effective at killing female *Ae. aegypti* under a variety of laboratory and semi-field conditions. DABS are a promising intervention for interdomiciliary control of *Ae. aegypti* and arboviral disease prevention.

Keywords: *Aedes aegypti*, vector control, toxic sugar bait, attractive bait, semi-field, dengue, arbovirus, ATSB

Background

Arboviral illnesses, including dengue, chikungunya, yellow fever, and Zika, are major contributors to morbidity and mortality in the tropics and subtropics. The burden is particularly apparent in Central and South America: between 2010—2018, the estimated annual number of dengue cases in the region ranged from 500,000 to 2,400,000 [1], and since 2013 PAHO has estimated that there have been more than 2.5 million suspected and confirmed cases of chikungunya and 800,000 cases of Zika. The viruses causing these diseases are spread mainly by the mosquitoes *Aedes aegypti* (Linnaeus, 1762; *Ae. aegypti*) and *Aedes albopictus* (Skuse, 1894; *Ae. albopictus*), with *Ae. aegypti* serving as the principal vector in many South American countries, including Ecuador [2]. Due to the lack of commercially-available vaccines for most human arboviral diseases, prevention efforts focus on vector surveillance and control methods [3].

Vector control relies heavily on contact-based insecticides, which are available in four main classes: organophosphates, pyrethroids, carbamates, and organochlorines. Indoor residual spraying is a common approach to vector control, for which twelve insecticides are available and approved for human use [4]. This small number of approved insecticides constitutes an impediment for the implementation of effective vector control strategies (such as pesticide rotation cycles) aimed at decreasing the development of resistance to any single insecticide [5]. As a result, pesticide resistance has become a major limitation for current vector control strategies, and is widespread in South American countries [6–8]. Our current reliance on a few chemical molecules to control *Ae. aegypti* is an increasingly flawed strategy, as evidenced by the proliferation of this disease vector across the globe and increasing arbovirus epidemics [9].

In contrast to the contact-based insecticide approach of the public health sector, the agricultural industry has focused on ingested insecticides for pest control. The use of ingested insecticides could be applied in disease control programs and interventions if disease vectors are successfully led to ingest the insecticide. One solution, attractive toxic sugar baits (ATSB), exploits

the nectar-feeding behavior of mosquitoes [10,11] to deliver the insecticide. An ATSB uses a mixture

of a lethal agent with sugar water and an additional attractant [12]. ATSBs have been tested for

Anopheles spp. [13–17], *Culex* spp. [15,16,18,19], *Ae. albopictus* [20–23], and other vector or

nuisance species [16] with a variety of attractants, baits, active ingredients, designs, and placement

strategies. Although laboratory bioassays demonstrate that ATSBs are toxic to *Ae. aegypti* [16,24,25],

semi-field and field evaluations have had poor results in reducing *Ae. aegypti* populations [26,27],

indicating that ATSB devices must be carefully designed and tested for each target species [12].

Compared to other mosquito species, *Ae. aegypti* appear to have a lower propensity for sugar-

feeding, preferring human blood meals instead [11]. Despite this, *Ae. aegypti* females will readily

feed on sugar in the laboratory, and often feed on plant sugars in the wild [28–31]. However,

traditional attractive sugar bait strategies that rely only on fruit volatiles as an attractant are likely

insufficient to "lure" highly anthropophilic female *Ae. aegypti* in the natural environment.

Herein we present the development of dried attractive bait stations (DABS) (Figure 1), and

show results from laboratory and semi-field experiments. In the laboratory we first identified the

lethality of DABS (Series 1.1), aimed to identify the killing mechanism of the DABS (Series 1.2),

assessed how the physiological status altered the effectiveness of DABS (Series 1.3), and assessed

the shelf life of the DABS (Series 1.4). In the semi-field trials, we sought to determine the timing of

mosquito mortality (Series 2.1), assess the relationship between DABS exposure time and mosquito

mortality (Series 2.2), and to demonstrate these effects in the presence of competing attractants (Series

2.3).

112   Results

113   Laboratory experiments

Series 1.1: Effects of DABS exposure on mosquito survival

We measured survival in mosquitoes exposed to DABS and compared to mosquitoes exposed

to control DABS in 20x20x20cm cages during in four independent replicates. An average of 13.5

(N=4, std = 3.88, SE= 1.94) out of 30 mosquitoes exposed to DABS survived at the first 24 hours

post-exposure. All mosquitoes had died by the end of the experiments at 48h post-exposure (Figure

2). In contrast, an average 29.75 (N=4, std = 0.50, SE=0.25) out of 30 mosquitoes exposed to control

devices survived after the first 24h. After the second 24h period, by the end of the experiment an

average of 29. 25 (N=4, std= 0.96, SE=0.48) had survived. Differences between toxic and control

treatments were highly significant at 24h (df=7, t=8.32, p<0.001) and 48h (df=7, t=61.1, p<0.001)

post-exposure.

Series 1.2: Characterization of the biological mode of action of the device.

125   We disrupted the feeding parts of mosquitoes and examined survival in those exposed to

DABS compared to those exposed to control DABS. After 48 hours, all mosquitoes that could still

feed (*i.e*. mosquitoes with an intact proboscis), died when exposed to the toxic device, and out of 20

19.33 (N=3, std= 0.58, SE=0.29) survived when exposed to the non-toxic control device. On average

12.33 mosquitoes that could not feed (those with ablated proboscis), survived regardless of inclusion

of the device's toxic condition (N=3, std=1.52, SE=0.87) or control non-toxic condition (N=3,

std=2.86, SE=1.65). Significant differences were found between the four treatments (df=11, F=70.55,

p<0.001). Post-hoc pairwise comparison determined that (a) the mortality of ablated mosquitoes

exposed to toxic devices was not significantly different to the mortality of ablated mosquitoes exposed

to control devices, and (b) the mortality of ablated mosquitoes was significantly different from the

mortality of whole mosquitoes exposed to toxic devices and whole mosquitoes exposed to control

devices (Figure 3).

137   Mosquitoes that had ingested toxic sugar solution presented histological abnormalities in the

posterior midgut. Electron micrographs revealed disruptions in the continuity of the gut epithelium

(Figures 4A, 4C). In addition, we found abnormal adipocytes that we hypothesize are undergoing a

process of necrosis (Figures 4E, 4F), and an increase in both the size and number of basal infolds in

the gut epithelial cells. We hypothesize that boric acid ingestion is the cause of these pathological

changes, which contributed to the mortality observed in specimens exposed to toxic devices.

Microscopic images of individuals exposed to control devices presented none of these pathologies on

the posterior midgut (Figures 4B, 4D).

Series 1.3: assessment of mosquito physiological status on DABS effectiveness

We measured survival in mosquitoes at blood fed and parous life stages in mosquitoes

exposed to DABS and mosquitoes exposed to control DABS. Mosquitoes with both physiological

statuses evaluated (blood fed and parous) presented lower survival when exposed to toxic devices

than when exposed to non-toxic control devices. In toxic conditions, an average of 19.33 (N= 3, std=

1.73, SE=0.99) blood fed females' out of 30 had survived at 48h. 2.67 (N= 3, std= 3.05, SE=1.76)

blood fed mosquitoes had survived at 72h, by the end of the experiment. 27 (N= 3, std= 1.73,

SE=0.99) out of 30 blood fed mosquitoes had survived to non-toxic DABS at the end of 72 hours

(Figure 5). Differences between control and toxic treatment survival were significant at 48h (df=5,

t=5.75, p<0.01) and 72h (df=5, t=12, p<0.001) post-exposure.

10.33 (N= 3, std= 3.51, SE=2.02) parous female mosquitoes at the first 24 hours and a 0 at

48 hours of exposure had survived to the toxic DABS devices (Figure 5B). In the non-toxic control

group 29.33 (N= 3, std= 0.58,) at 48 hours of exposure. Differences between control and toxic

treatment survival curves were significant at 24h (df=5, t=9.25, p<0.001) and 48h (df=5, t=87.99,

p<0.001) post-exposure.

Series 1.4: Assessment of shelf-life of the DABS device

We tested the shelf life of DABS by measuring survival of mosquitoes exposed to DABS

compared to those not exposed to DABS, using DABS that had been constructed at various time

points in the past. 30 out of 30 mosquitoes exposed to toxic devices stored for 38 days had survived

at 24 hours; 28.67 (N= 3, std= 0.58, SE=0.33) mosquitoes exposed to control conditions survived at

48 hours (Fig. 6A). Differences in survival between conditions were highly significant at 48h post-

exposure (df=5, t=86, p < 0.001).

5 (N= 3, std= 1.0, SE=0.58) mosquitoes exposed to toxic devices stored for 80 had survived

at the first 24 hour period and 0 after 48 hours; 29.33 (N=3, std=1.16, SE= 0.67) mosquitoes exposed

to control conditions survived at 48 hours (Fig. 6B). Differences in survival between conditions were

highly significant at 48h post-exposure (df=5, t=44, p < 0.001).

On average 28.33 (N=3, std= 0.58, SE=0.33), 10.66 (N=3, std=4.62, SE=2.67), and 0

mosquitoes exposed to toxic devices stored for 118 had survived at 24h, 48h, and 72h, respectively

(Fig. 6C). Differences in survival between conditions were highly significant at 48h (df=5, t=6.95,

p<0.01) and 72h (df=5, t=87.99, p < 0.001) post-exposure.

Semi-field experiments

We assessed the attractiveness of DABS by measuring mortality in mosquitoes exposed to

DABS compared to mosquitoes not exposed to DABS in experimental houses. When exposed to

DABS in semi-field trials (Series 2.1, Figure 7), mosquito mortality was 0.0—6.0% (mean: 2.0%,

standard error: 0.9%) in the control and 17.0—57.1% (mean: 36.7%, standard error: 5.3%) in the

treatment house after 24 hours (df=5, t=-7.0, p<0.001). At 48 hours, mortality was 0.0—18.0% (mean:

5.4%, standard error: 2.4%) in the control and 22.0—51.1% (mean: 38.9%, standard error: 3.9%) in

the treatment house (df=5, t=-5.36, p<0.01). At 72 hours, mortality was 0.0—4.1% (mean: 0.7%,

standard error: 0.6%) in the control and 0.0—4.0% (mean:1.4%, standard error: 0.6%) in the treatment

house (df=5, t=-0.80, p>0.05). The cumulative mortality of the control was 4.1—18.0% (mean: 8.2%,

standard error: 1.9%) and 54.0—98.0% (mean: 76.9%, standard error: 6.2%) in the treatment house

(df=5, t=-8.37, p<0.001). Most mosquito mortality was observed within the first 48 hours of the

experiment, with no difference in mosquito mortality after this time period.

When exposed to DABS for 48 hours (Series 2.2, Figure 8), mosquito mortality was 2.0—

22.9% (mean: 11.7%, standard error: 2.8%) in the control and 77.3—100.0% (mean: 91.5%, standard

error: 3.8%) in the treatment house (df=5, t=-17.0, p<0.001), indicating high mortality from 48 hours

of exposure to DABS in the treatment houses.

When alternative attractants were included in the experimental houses (Series 2.3, Figure 9),

mosquito mortality was 2.0—32.7% (mean: 14.1%, standard error: 4.1%) in the control and 68.0—

100.0% (mean: 89.6%, standard error: 4.5%) in the treatment house (df=5, t=-12.90, p<0.001),

indicating that DABS results in high mortality even in the presence of a competing attractant.

When comparing the results of 24 hours (Series 2.1) to 48 hours of exposure (Series 2.2), 48 hours of

exposure results in higher mortality at 48 hours (df=10, t=-8.78, p<0.001) in the treatment group

(Supplemental Table 1), with no difference in the control groups (df=10, t=-1.55, p>0.05).

When comparing 48 hours of exposure to DABS only (Series 2) and 48 hours of exposure to

DABS in the presence of a competing attractant (Series 2.3), there is no effect of a competing

attractant on the effect of DABS on mosquito mortality (df=10, t=0.28, p>0.05) in the treatment group

(Table 1). High mortality from 48 hours of DABS exposure was observed despite the presence of a

competing attractant.

Discussion

These experiments demonstrate that DABS can strongly impact the mortality of female *Ae.*

aegypti under laboratory and semi-field conditions. In these settings, we show that mortality occurs

within the first 48 hours of exposure to our devices. In addition, DABS attract and kill *Ae. aegypti*

even in the presence of an alternative sugar source. To the best of our knowledge, this device is the

only known "dry" ATSB. The simple and economic design lends itself to in-home use in resource

limited settings where *Ae. aegypti* target human hosts and transmit dangerous arboviruses.

Our assessment of the biological action of the devices provides an insight into the mechanism

by which low concentrations of boric acid affect *Ae. aegypti*. We determined that boric acid enters

the insect´s body by ingestion, further supporting the notion that this inorganic pesticide acts as a

stomach poison, as previously suggested [32,33]. Based on our electron microscopy analysis, we

hypothesize that the ingestion of boric acid disrupts the integrity of the gut epithelium.

Considering that the proposed mechanism by which boric acid exerts its toxic effect (gut

disruption) is markedly different from the neurotoxic mechanism by which most traditional pesticides

cause mortality, we propose that our devices have the potential to act as efficient complementary tools

to combat the spread of resistance to traditional pesticides. By combining the use of DABS with

traditional pesticides in the same areas, it would be possible to target two different and crucial systems

(namely, the nervous and digestive systems) in the insect´s body simultaneously, thereby reducing

the mosquito´s probability of survival and decreasing the probability of the development of

insecticide resistance.

We observed significant mortality of blood fed female *Ae. aegypti* exposed to the DABS

device, albeit at lower rate than starved females. Interestingly, the largest drop in survival probability

in blood fed females is observed between 48h and 72h post-exposure to the device (Fig 5), suggesting

that after 48h, females have already used imbibed blood for the development of eggs and are keen to

search for further meals. Based on this evidence, it is plausible to suggest that if employed in the field,

DABS devices may be efficient in killing female mosquitoes of various physiological states, including

females that have already ingested blood – a particularly important group for disease transmission.

Novel vector control methods have the potential to serve as critical tools in the public health

effort to control persistent and emerging vector borne diseases. Various designs of ATSBs have had

promising field trials for potential control of *Aedes albopictus* Skuse, 1894, *Anopheles* spp. and *Culex*

spp. [15–17,20,21,24]. Previous research shows that several formulations of ATSBs can achieve *Ae.*

aegypti mortalities above 80% in laboratory settings [16,25], but results from ATSBs in semi-field or

field settings have been mixed. Early field trials did not show a positive effect of ATSBs on *Ae.*

aegypti [26,27], however a recent field trial in Bamako, Mali showed promising success [31]. The

principle barrier to field trial success appears to be the ability to attract *Ae. aegypti* to ATSBs and

mixed results have been achieved when using floral-based attractants.

We hypothesize that our device attracts *Ae. aegypti* with strong visual cues (as opposed to a

chemical) as an attractant. *Ae. aegypti* are container breeders [34,35], that utilized tree holes in their

natural forested habitat before adapting to life in human civilization. The DABS device has a high-

contrast (black and white) 28-inch^2 surface to simulate a refuge for *Ae. aegypti* [36]. High contrast

coloring has similarly been integrated into prior trap designs and has been shown to improve capture

rates of *Ae. aegypti* [37]. We believe the high-contrast coloring of DABS draws *Ae. aegypti* to land

on the device.

These experiments have demonstrated the effectiveness of DABS on *Ae. aegypti* in laboratory

and semi-field experimental conditions. Our approach differs from most ATSB approaches in two

important ways: firstly, we use a device with a dried sugar solution to elicit an ingestion response

while other ATSBs typically use liquid sprayed on vegetation [12,15,17,26]. We hypothesize that the

device is a key element in the effectiveness of DABS. Similar to other diptera [38], *Ae. aegypti* are

able to evaluate surfaces with their feet, and the "taste" of a landing surface can either lead the

mosquito to feed and ingest, or reject, the surface [39]. Additionally, the device provides two

operational advantages over spraying liquid solutions: 1) liquid solutions are more difficult to

manufacture, ship, and distribute than devices, and 2) the device can be smaller and more easily

deployed. Secondly, we use a visual rather than chemical attractant to lure *Ae. aegypti* to the device.

Chemical attractants add to the cost and decrease the shelf life of any device. Previous research has

questioned the ability of sugar solutions alone to attract mosquitoes [26,33], leading to research on

chemical attractant additives for ATSBs, but the use of chemical attractants in ATSBs targeting *Ae.*

aegypti have been unsuccessful [26,27]. We demonstrate that a simple black-and-white visual

attractant is a sufficient motivator for female *Ae. aegypti* to land on the surface of DABS even in the

presence of a competing oasis. Taken together, we hypothesize that the visual cues attract the *Ae.*

aegypti to land on the device, upon which the presence of the dry sugar on the device's surface entices

the insect to ingest it. When this sugar solution is mixed with boric acid, ingestion results in insect

mortality.

We propose that these encouraging results justify larger field trials of DABS in open-air

environments. We show that 48 hours of DABS exposure leads to high mosquito mortality when used

in the laboratory and in experimental houses reminiscent of peri-urban tropical housing. Furthermore,

we have established that the effectiveness of DABS for killing *Ae. aegypti* is maintained even after

prolonged storage periods, a characteristic that would facilitate their use in semi-field and field

conditions.

Semi-field trials are a crucial step to bring a scalable, marketable product to intra domiciliary

field testing. An in-home approach is ideal for control of *Ae. aegypti*, as the vector has an extremely

limited flight range, often spending its entire life within a single household [5,35,40]. Other research

with ATSBs has shown that end-users of these products prefer to have them placed indoors [14]. The

successful design and placement strategy of DABS used in our experiments indicate that the device

is ideal for in-home field testing.

279   Limitations

These experiments were conducted under laboratory and semi-field conditions, which can

only moderately emulate real-world/field conditions. Semi-field experiments were limited to

nulliparous females and we cannot be certain how DABS will affect gravid or blood-fed females or

males in an open-air environment—though it should be noted that DABS were equally effective in

attracting and killing blood-fed and nulliparous females under laboratory conditions. It is also unclear

if DABS would impact non-target insect species, such as butterflies or other pollinators, though if

DABS are limited to use inside the home, it is unlikely to affect these species. Although DABS

performed well in the presence of a competing attractant (100 grams of apples), it is unlikely that the

attractant used in our experiments are a realistic substitute for open-air field conditions. An actual

home will contain many competing attractants, including human hosts. It is difficult to know if the

success of DABS in semi-field conditions will be replicated in occupied homes in the field; the

number and placement of DABS may need to be modified. In addition, it is unclear how end users

will react to placement of DABS in their homes, although our preliminary examinations (unpublished)

suggest residents are receptive of DABS and there is evidence that residents in areas of high *Ae.*

aegypti burden are willing to utilize numerous home-based mosquito control products [41].

Conclusions

With careful design and device placement consideration, we have created a promising vector

control device ready for large-scale trials to test its ability to control *Ae. aegypti* in natural conditions.

We demonstrated that DABS are capable of attracting and killing female *Ae. aegypti* in experimental

houses, and that 48 hours in the presence of DABS leads to high mortality among female *Ae. aegypti*.

Importantly, DABS were efficient at killing female mosquitoes of diverse physiological statuses, and

can attract and kill female *Ae. aegypti* even in the presence of a competing attractant.

Materials and methods

Aim and design

The general aim of our study was to assess the effectiveness of DABS at killing female *Ae.*

aegypti in numerous laboratory and semi-field experimental conditions. In the laboratory we first

measured a killing effect of DABS (Series 1.1), identified the killing mechanism (Series 1.2), and

measured how mosquito physiological status affected DABS lethality (Series 1.3). In the semi-field

experiments our specific aims were (a) to determine the timing of mosquito mortality (Series 2.1), (b)

to assess the relationship between DABS exposure time and mosquito mortality (Series 2.2), and (c)

to demonstrate these effects in the presence of competing attractants (Series 2.3).

In the laboratory we first identified the lethality of DABS (Series 1.1), aimed to identify the

killing mechanism of the DABS (Series 1.2), assessed how the physiological status altered the

effectiveness of DABS (Series 1.3), and assessed the shelf life of the DABS (Series 1.4). In the semi-

field trials, we sought to determine the timing of mosquito mortality (Series 2.1), assess the

relationship between DABS exposure time and mosquito mortality (Series 2.2), and to demonstrate

these effects in the presence of competing attractants (Series 2.3).

Study setting

Laboratory experiments

Laboratory experiments were conducted at the Center for Research on Health in Latin America (CISeAL, by its Spanish acronym), where they were reared and maintained under standard insectary conditions: $28 \pm 1°C$ temperature, $80 \pm 10\%$ relative humidity, and a 12h:12h (L:D) light cycle. Larvae were fed finely ground fish food. When required, mosquitoes were sexed during the pupal stage. Adults were kept in 20x20x20 cm cages. For maintenance, adult mosquitoes were fed 10% sucrose solution *ad libitum*. For blood feeding, female adult mosquitoes were offered access to a restrained female mouse. All mosquitoes were maintained under insectary conditions after adult emergence before they were used for experiments. Mosquitoes referred to as "starved" in this manuscript were deprived of access to sugar or blood (but not water) for 48h prior to their use in experiments.

Semi-field trials

Assays were performed in experimental houses meant to emulate typical housing found in areas with active dengue transmission. Photographs of the houses are available in an additional file (see Additional file 1). The houses are constructed of wood and cane and are raised on a 1-meter platform with walkways to improve structural integrity and facilitate window access; one window on each house is equipped with window escape traps with sleeves to monitor escape behavior. The dimensions of the houses are 3.85 meters (m) wide x 4.85 m long x 3 m high. Each house has three windows (0.9 m wide × 0.6 m high) and one door (1.03 m wide × 3 m high). The house frames are made of wood, untreated wooden plank flooring, walls of untreated cane, and a roof of zinc panels. The window traps are 0.45 m long x 0.66 m wide x 0.45 m high. The houses are located on campus at the Universidad Técnica de Machala in the city of Machala, Ecuador (3°15' S, 79°57' W), a region with abundant wild populations of *Ae. aegypti* and endemic arbovirus transmission. Experiments were conducted under ambient climate conditions (temperature range: 23.1—35.6° C, mean temperature:

342   28.4° C, relative humidity range: 43.9—95.0%, mean relative humidity: 75.8%). Each trial replicate

343   was conducted with one control and one experimental house; the specific house used as the

344   experimental or control house was alternated upon each replicate.

Biological material

Aedes aegypti Linnaeus, 1762 eggs were provided by the Center for Research on Health in

Latin America (CISeAL, by its Spanish acronym) at the Pontificia Universidad Católica del Ecuador.

All strains used in this work originated from Ecuador, and had been maintained in laboratory

conditions since 2015. Laboratory work was performed with strains originally collected in Ecuador

from the cities of Guayaquil and Puerto Francisco de Orellana. Semi-filed work was performed with

a strain originally collected in the city of Machala.

Semi-field experiments

Hatching and rearing of *Ae. aegypti* for semi-field experiments were performed at the

Laboratory of Entomology at the Universidad Técnica de Machala. Considering this laboratory is

located in a region where *Ae. aegypti* actively reproduces and thrives, environmental conditions

(temperature: 28-32 °C; relative humidity: 60-80%) were not artificially controlled in our mosquito-

rearing facilities. A vacuum pressure system was used to synchronize egg hatching (one hour

exposure to obtain first stage larvae). Larvae were fed with finely ground fish food. At the pupal

stage, males and females were separated. Adults were kept in 20x20x20 cm cages. Adults were fed

on 10% sugar solution *ad libitum*. Each experimental semi-field experiment series used nulliparous

females aged 1-5 days and starved for 24 hours prior to experimental release.

Dried Attractive Bait Stations (DABS)

The DABS device consists of two concentric foam disks (an inner white disk one centimeter

in diameter, and an outer black disk eight centimeters in diameter). Experimental DABS were

impregnated with a 10% sucrose solution containing 1% boric acid as a lethal agent. Control DABS

were impregnated with 10% sucrose solution without boric acid. Photographs of the devices are

available in an additional file (see Supplementary Figure 2, Additional file 2), US Patent App.

15/990,931, 2018 .

Laboratory experiments

Series 1.1: Survival assessment of mosquitoes exposed to the device

371   To determine whether exposure to the DABS devices has an influence on adult mosquito

372 survival probability, we conducted an experiment in which groups of 30 adult female mosquitoes,

373 placed in a 15x15x15cm cage, were exposed during 48 hours to either a DABS device or a control

374 device (sugar solution but no boric acid). We replicated each experiment four times. The assessment

375 was repeated using each of the two laboratory strains described previously.

Series 1.2: Appraisal of biological mode of action of the device

To establish whether the toxic component of DABS needs to be ingested by the mosquitoes

in order to exert its effect, we presented the devices to cohorts of adult females aged 1 to 7 days,

which were unable to ingest food due to the surgical ablation of their mouthparts. To establish these

cohorts, individual mosquitoes were first anesthetized by placing them at 4°C during 10-15 minutes.

Anesthetized specimens were individually placed under a dissection microscope and, using a human

hair, we tied a knot at the proboscis' proximal end in order to create a constriction that would impede

the flow of food. Subsequently, the part of the proboscis anterior to the knot was removed using

micro-dissection scissor (Supplemental Figure 3). Following the surgery, mosquitoes were left to rest

for 24 hours before being used in any experiment. To control for the potential negative effect of the

anesthetizing procedure in mosquito survival, non-ablated mosquitoes used in control groups were

also placed at 4°C during 10-15 minutes, and allowed to recover during 24 hours before experimental

set-up.

We conducted the experiment with four separate cages, each with 20 starved mosquitoes

(Table 2 shows different treatments). We treated cage 1 with toxic DABS devices and used 20 ablated

mosquitoes; cage 2 held non-toxic control devices and 20 ablated mosquitoes. We treated cage 3 with

toxic DABS devices and non-ablated mosquitoes; cage 4 held a non-toxic control device and non-

ablated mosquitoes. We assessed mortality in all groups at 24 and 48 hours of exposure to the devices.

We replicated the experiment three times.

We then conducted an experiment wherein 30 adult starved female mosquitoes aged 1 to 7

days were introduced to a cage with a DABS device, and 30 adult starved female mosquitoes of

similar age were introduced to a cage with a non-toxic control device. We monitored cages for 24

hours, and removed dead mosquitoes by aspiration every hour from the cages. Using a dissection

microscope, we removed the legs, head, and wings of every dead specimen onto a drop of 70%

ethanol. Through this process we gently disrupted the abdominal cuticle to permit the exposure of

internal tissues to the fixative. Afterwards we fixed individual mosquitoes in a solution containing

2.5% glutaraldehyde, 2.5% paraformaldehyde in 0.1M cacodylate buffer (pH 7.4), and stored them at

4°C for 72 hours. We then washed specimens in cacodylate buffer with 0.1M sucrose overnight. Post-

fixing was achieved by leaving the specimens for two hours at 4°C in 2% osmium tetroxide in 0.1

cacodylate buffer (pH 7.4). Subsequently, individuals were stained using 2% uranyl acetate and left

to rest for three hours in the dark at room temperature. Tissues were later dehydrated through a series

of ethanol baths (50%, 70%, 95%, 100%). Afterwards, they were placed in propylene oxide for 30

minutes, then in a 1:1 volume propylene oxide resin mixture (Epon 812, Araldite 502, dodecenyl

succinic anhydride, benzyl dimethylamine) for one hour and later, one more volume of resin was

added and left on a rotator overnight. Finally, mosquitoes were embedded in resin and incubated at

60°C for 24 hours. Resin were stained using 2% uranyl acetate. We then utilized a transmission

electron microscope to observe specimens and obtain micrographs of interesting tissues.

Series 1.3: Effects of the physiological status of the mosquitoes on the performance of DABS

We examined two different physiological statuses using mated starved female adult

mosquitoes aged 1 to 7 days, namely blood fed and parous. We established females deemed as "blood

fed" by selecting blood-engorged individuals immediately after a blood meal. We established females

deemed as "parous" by first blood feeding and subsequently maintaining mosquitoes for 7 days under

insectary conditions in order to ensure that they had oviposited before being used for experimentation.

We set up two cages for each of the defined physiological statuses, each with 30 mosquitoes. One

cage exposed the mosquitoes to an ATSB device, and the other held a control non-toxic device. We

gathered survival data at 24 and 48 hours following introduction to the cages, and replicated these

experiments three times.

Series 1.4: Shelf-life of the device

In order to determine the shelf life of ATSB devices, toxicity tests were performed using

devices which had been stored for 38, 80, and 118 days after their production. For storage, devices

were individually wrapped inside a sealed plastic bag and placed in an incubator at $28 \pm 2°C$ and 80

$\pm 10\%$ relative humidity. We conducted three replicates of previously described experiments for each

storage time.

Semi-field trials

Series 2.1: 24 hours of DABS exposure in experimental houses

Each house contained four DABS devices (control or treatment DABS as appropriate)

suspended on strings attached to the roof of the house at a height of 30—50cm above ground and

approximately 30cm from the nearest wall. Diagrams of the experimental layout are available in an

additional file (see Supplementary Figure 3, Additional file 2). For each trial replicate, 50 female *Ae.*

aegypti were released into each house through the escape window sleeve (release time 11:00a.m.—

2:00p.m.). Twenty-four hours after release, dead mosquitos were collected from the floor and window

escape traps in each house, and the remaining live mosquitoes were captured with a hand-held

aspirator (Prokopack, John W. Hock Company, Gainesville, USA). All live mosquitoes were labeled

by experimental group and observed for 48 additional hours in laboratory cages (under laboratory

conditions with food available). Mortality was calculated for 24 hours, 48 hours, and 72 hours. Six

trial replicates were performed for Series 1.

Series 2.2: 48 hours of DABS exposure in experimental houses

Each house contained four DABS devices (control or treatment DABS as appropriate) and

two sources of water (wet cotton in a black plastic bucket). Photographs of the water source are

available in an additional file (see Supplementary Figure 3, Additional file 2). For each trial replicate,

50 female *Ae. aegypti* were released into each house through the escape window sleeve (release time

8:00a.m.—11:00a.m.). Forty-eight hours after release, dead mosquitos were collected in each house

and remaining live mosquitoes were captured with an aspirator. Mortality was calculated for 48 hours.

Six replicates were performed for Series 2.

Series 2.3: 48 hours of DABS exposure in experimental houses with competing attractant

Each house contained four DABS devices (control or treatment DABS as appropriate), two

sources of water (wet cotton in a black plastic bucket), and 100 grams of peeled, cut apples in a dish

placed on a chair in the center of the house as a competing attractant. Photographs of the competing

attractant are available in an additional file (see Supplementary Figure 3, Additional file 2). Recently

emerged female *Ae. aegypti* rely on sugar meals for energy, these meals may include aging fruit and

female *Ae. aegypti* will feed on fructose (as is found in apples). For each trial replicate, 50 female *Ae.*

aegypti were released into each house through the escape window sleeve (release time 9:00a.m.—

12:00p.m.). Forty-eight hours after release, dead mosquitos were collected in each house and

remaining live mosquitoes were captured with an aspirator. Mortality was calculated for 48 hours.

Six replicates were performed for Series 3.

Statistical analyses

For the Series 1 experiments, data was processed, plotted, and analyzed using Python v2.7.13.

For data processing we used Pandas v0.22.0 module. Plots were generated using the Plotly v3.10.0

module. We examined the normal distribution of the data with Kolmogorov-Smirnov and Shapiro-

Wilk tests. In experiments in Series 1.1, 1.3, and 1.4 Student's T-test comparisons were performed

using the Scipy v1.0.0 module. In Series 1.2, one-way ANOVA was performed using Scipy v1.0.0

module with four experimental groups. Tukey's range test, using Statsmodels v.0.10.0 module, was

performed after ANOVA for determining ranges for each group. All data and codes used for this data

have been stored in a private online git repository and will be provided upon request. In Series 2.1—

2.3, mosquito mortality data from each series were compared using a two-tailed paired t-test (paired

by replicate). Mean mosquito mortality was compared across series using a two-tailed t-test. Data

were analyzed using Excel (Microsoft, Redmond, USA).

List of abbreviations

Ae. aegypti=*Aedes aegypti* Linnaeus, 1762

Ae. albopictus=Aedes albopictus Skuse 1894

ATSB=attractive toxic sugar bait

DABS=dried attractive bait stations

cm =centimeters

m=meters

Declarations

Ethics approval:
These study protocols were found to be exempt from Institutional Review Board review by

Syracuse University Institutional Review Board.

Consent for publication:
Not applicable

Availability of data and material:
The datasets used and/or analysed during the current study are available from the

corresponding author on reasonable request.

Competing interests:

ASI, DL, and MN hold a patent on the DABS technology.

Funding:

Laboratory work was funded by Pontificia Universidad Católica del Ecuador´s Internal Research Grant L13234, awarded to MN. Semi-field work was funded by a seed grant from Syracuse University, awarded to DL.

Author contributions:

MN, ASI, and DL designed the study.

MN, ASI, DL, and EBA provided study materials.

GR, ML, AA, BM VS and FHH performed the experiments.

RH performed electron microscopy work

RS and GR conducted data analysis.

RS, GR, MN, VS, and FHH drafted the manuscript.

All authors reviewed and approved the final draft of the manuscript.

Acknowledgements:

The authors would like to thank Cinthya Cueva Aponte for her assistance with study documentation, Chad Ward and Matt Kovacs for assistance with mosquito rearing, Mario Grijalva for his logistical support for the performance of electron microscopy, Felipe Andrade and Phillip Huegli for providing useful recommendations on programming, and Santiago Cadena for his support in insectary maintenance.

Additional files

Additional file 1 (pdf)

Supplementary Figure 1: Experimental Houses

Additional file 2 (pdf)

Supplementary Figure 2: Experimental Layout

516   Additional file 3 (pdf)

517   Supplemental Figure 2: Feeding disruption procedure. A) Anesthetized individual with whole

518   proboscis. B) Human hair tied at the proximal end of the proboscis. C) Micro-dissection scissor

519   removal of proboscis' segment anterior to the knot. D) Feeding disrupted individual.

References

1. PAHO, WHO. Epidemiological Update: Dengue. 2019;1–14.
2. Boyer S, Calvez E, Chouin-Carneiro T, Diallo D, Failloux AB. An overview of mosquito vectors
of Zika virus. Microbes Infect. 2018;20:646–60.
3. Olliaro P, Fouque F, Kroeger A, Bowman L, Velayudhan R, Santelli AC, et al. Improved tools and
strategies for the prevention and control of arboviral diseases: A research-to-policy forum. PLoS
Negl Trop Dis. 2018;12:e0005967.
4. Oxborough RM. Trends in US President's Malaria Initiative-funded indoor residual spray coverage
and insecticide choice in sub-Saharan Africa (2008-2015): urgent need for affordable, long-lasting
insecticides. Malar J. 2016;15:146.
5. Harrington LC, Scott TW, Lerdthusnee K, Coleman RC, Costero A, Clark GG, et al. Dispersal of
the dengue vector Aedes aegypti within and between rural communities. Am J Trop Med Hyg.
2005;72:209–20.
6. Hemingway J, Ranson H. Insecticide Resistance in Insect Vectors of Human Disease. Annu Rev
Entomol. 2000;45:371–91.
7. Chanda E, Kamuliwo M, Sikaala CH, Thomsen EK, Ranson H, Hemingway J, et al. An operational
framework for insecticide resistance management planning. Emerg Infect Dis. 2016;22:773–9.
8. Ryan SJ, Mundis SJ, Aguirre A, Lippi CA, Beltrán E, Heras F, et al. Seasonal and geographic
variation in insecticide resistance in Aedes aegypti in southern Ecuador. PLoS Negl Trop Dis.
2019;13:e0007448.
9. Kraemer MUG, Sinka ME, Duda KA, Mylne A, Shearer FM, Barker CM, et al. The global
distribution of the arbovirus vectors Aedes aegypti and Ae. albopictus. Elife. 2015;4:e08347.
10. Yuval B. The other habit: sugar feeding by mosquitoes. Bull Soc Vector Ecol. Society for Vector
Ecology; 1992;17:150–6.
11. Foster WA. Mosquito sugar feeding and reproductive energetics. Annu Rev Entomol.
1995;40:443–74.
12. Fiorenzano JM, Koehler PG, Xue R De. Attractive toxic sugar bait (ATSB) for control of
mosquitoes and its impact on non-target organisms: A review. Int J Environ Res Public Health.
2017;14.
13. Tenywa FC, Kambagha A, Saddler A, Maia MF. The development of an ivermectin-based
attractive toxic sugar bait (ATSB) to target Anopheles arabiensis. Malar J. 2017;16:1–10.
14. Maia MF, Tenywa FC, Nelson H, Kambagha A, Ashura A, Bakari I, et al. Attractive toxic sugar
baits for controlling mosquitoes: A qualitative study in Bagamoyo, Tanzania. Malar J. 2018;17:1–
6.
15. Stewart ZP, Oxborough RM, Tungu PK, Kirby MJ, Rowland MW, Irish SR. Indoor application
of attractive toxic sugar bait (ATSB) in combination with mosquito nets for control of pyrethroid-
resistant mosquitoes. PLoS One. 2013;8.
16. Qualls WA, Müller GC, Revay EE, Allan SA, Arheart KL, Beier JC, et al. Evaluation of attractive
toxic sugar bait (ATSB)-Barrier for control of vector and nuisance mosquitoes and its effect on
non-target organisms in sub-tropical environments in Florida. Acta Trop. 2014;131:104–10.
17. Qualls WA, Müller GC, Traore SF, Traore MM, Arheart KL, Doumbia S, et al. Indoor use of
attractive toxic sugar bait (ATSB) to effectively control malaria vectors in Mali, West Africa.

Malar J. Malaria Journal; 2015;14:301.

18. Qualls WA, Scott-Fiorenzano J, Müller GC, Arheart KL, Beier JC, Xue R-D. Evaluation and
Adaptation of Attractive Toxic Sugar Baits For Culex tarsalis and Culex quinquefasciatus Control
In The Coachella Valley, Southern California . J Am Mosq Control Assoc. 2017;32:292–9.

19. Müller GC, Junnila A, Schlein Y, Ller GCM. Effective Control of Adult Culex pipiens by
Spraying an Attractive Toxic Sugar Bait Solution in the Vegetation Near Larval Habitats. Source
J Med Entomol J Med Entomol. 2010;47:63–6.

20. Revay EE, Müller GC, Qualls WA, Kline DL, Naranjo DP, Arheart KL, et al. Control of Aedes
albopictus with attractive toxic sugar baits (ATSB) and potential impact on non-target organisms
in St. Augustine, Florida. Parasitol Res. 2014;113:73–9.

21. Junnila A, Revay EE, Müller GC, Kravchenko V, Qualls WA, Xue R de, et al. Efficacy of
attractive toxic sugar baits (ATSB) against Aedes albopictus with garlic oil encapsulated in beta-
cyclodextrin as the active ingredient. Acta Trop. 2015;152:195–200.

22. Scott-Fiorenzano JM, Fulcher AP, Seeger KE, Allan SA, Kline DL, Koehler PG, et al. Evaluations
of dual attractant toxic sugar baits for surveillance and control of Aedes aegypti and Aedes
albopictus in Florida. Parasites and Vectors. 2017;10:1–9.

23. Scott JM, Seeger KE, Gibson-Corrado J, Muller GC, Xue R De. Attractive Toxic Sugar Bait
(ATSB) Mixed With Pyriproxyfen for Control of Larval Aedes albopictus (Diptera: Culicidae)
Through Fecal Deposits of Adult Mosquitoes. J Med Entomol. 2017;54:236–8.

24. Khallaayoune K, Qualls WA, Revay EE, Allan SA, Arheart KL, Kravchenko VD, et al. Attractive
Toxic Sugar Baits: Control of Mosquitoes With the Low-Risk Active Ingredient Dinotefuran and
Potential Impacts on Nontarget Organisms in Morocco. Environ Entomol. 2013;42:1040–5.

25. Lea AO. Sugar-baited insecticide residues against mosquitoes. Mosq News. 1965;25:65.

26. Fikrig K, Johnson BJ, Fish D, Ritchie SA. Assessment of synthetic floral-based attractants and
sugar baits to capture male and female Aedes aegypti (Diptera: Culicidae). Parasites and Vectors.
2017;10:1–9.

27. Xue R-D, Ali A, Kline DL, Barnard DR. Field Evaluation of Boric Acid- and Fipronil-Based Bait
Stations Against Adult Mosquitoes. J Am Mosq Control Assoc. 2008;24:415–8.

28. Martinez-Ibarra JA, Rodriguez MH, Arredondo-Jiménez JI, Yuval B, Arredondo-Jimenez JI,
Yuval B. Influence of plant abundance on nectar feeding by Aedes aegypti (Diptera: Culicidae) in
southern Mexico. J Med Entomol. 1997;34:589–93.

29. Spencer CY, Pendergast TH, Harrington LC. Fructose variation in the dengue vector, Aedes
aegypti, during high and low transmission seasons in the Mae Sot region of Thailand. J Am Mosq
Control Assoc. 2005;21:177–81.

30. van Handel E, Edman JD, DAY JF, Scott TW, Clark GG, Reiter P, et al. Plant-sugar, glycogen,
and lipid assay of Aedes aegypti collected in urban Puerto Rico and rural Florida. J Am Mosq
Control Assoc. 1994;10:149–53.

31. Sissoko F, Junnila A, Traore MM, Traore SF, Doumbia S, Dembele SM, et al. Frequent sugar
feeding behavior by Aedes aegypti in Bamako, Mali makes them ideal candidates for control with
attractive toxic sugar baits (ATSB). PLoS One. 2019;14:1–21.

32. Xue R-D, Kline DL, Ali A, Barnard DR. Application of boric acid baits to plant foliage for adult
mosquito control. J Am Mosq Control Assoc. American Mosquito Control Association;
2006;22:497–500.

33. Xue R-D, Barnard DR. Boric acid bait kills adult mosquitoes (Diptera: Culicidae). J Econ
Entomol. 2003;96:1559–62.

34. Garcia-Rejon J, Loroño-Pino MA, Farfan-Ale JA, Flores-Flores L, Del Pilar Rosado-Paredes E,
Rivero-Cardenas N, et al. Dengue virus-infected Aedes aegypti in the home environment. Am J
Trop Med Hyg. 2008;79:940–50.

35. Kuno G. Review of the factors modulating dengue transmission. Epidemiol Rev. 1995;17:321–
35.

36. Fay RW, Prince WH. A modified visual trap for Aedes aegypti. Mosq News. 1970;

37. Barrera R, Mackay AJ, Amador MA. An improved trap to capture adult container-inhabiting
mosquitoes. J Am Mosq Control Assoc. 2013;29:358–68.
38. Meunier N, Marion-Poll F, Rospars JP, Tanimura T. Peripheral coding of bitter taste in
Drosophila. J Neurobiol. 2003;56:139–52.
39. Dennis EJ, Goldman O V., Vosshall LB. Aedes aegypti Mosquitoes Use Their Legs to Sense
DEET on Contact. Curr Biol. 2019;29:1551-1556.e5.
40. Hemme RR, Thomas CL, Chadee DD, Severson DW. Influence of urban landscapes on population
dynamics in a short-distance migrant mosquito: evidence for the dengue vector Aedes aegypti.
PLoS Negl Trop Dis. 2010;4:e634.
41. Heydari N, Larsen D, Neira M, Beltrán Ayala E, Fernandez P, Adrian J, et al. Household Dengue
Prevention Interventions, Expenditures, and Barriers to Aedes aegypti Control in Machala,
Ecuador. Int J Environ Res Public Health. 2017;14:196.
625
626

627 **<u>Figures</u>**

629   Figure 1. Attractive toxic sugar bait devices. A) 3D model of the DABS devices. Diameters

630   for each of the foam sheet circles are shown.B) Photography of a DABS device used in the study.

Survival of Mosquitoes Exposed to DABS

631
632   Figure 2. Survival assessment of mosquitoes exposed to the device. All mosquitoes (n=30)

633   exposed to toxic devices die after 48 hours of exposure. When presented with non-toxic device almost

634   all survive. Box plot indicating median 25% and 75% quartiles. Error bars indicate max and min

635   values. Each dot indicates an independent count of mosquitoes (y axis) exposed to control devices

636   (blue) or toxic devices (red) at different time points (x axis).

639
640   Figure 3. Uptake mechanism of the toxic component. Toxic effect is dependent on the ability

of mosquitoes to ingest the toxic component. When mosquitoes are able to ingest the toxic component

all mosquitoes (n =20) die after 48 hours (red). Mosquitoes with ablated mouthparts die equally

regardless of the toxic or non-toxic condition of the device (green and yellow). Almost all mosquitoes

with no ablation or toxic devices survive. Box plot indicating median 25% and 75% quartiles. Error

bars indicate max and min values. Each dot indicates an independent count of mosquitoes (y axis)

exposed to control devices (blue), control devices + mosquito mouthparts ablation (yellow), toxic

devices + mosquito mouthparts ablation, or toxic devices (red) at different time points (x axis).

Figure 4. Histopathological effects on the midgut. Longitudinal sections of Ae. aegypti posterior midgut. Panels A, C, D: Mosquitoes exposed to toxic devices. Panel B: Mosquito exposed to non-toxic device. Specimens exposed to toxic devices showed disruptions in the gut integrity (ED, panel A). Because of the even distribution of adjacent bacterial cells in the gut lumen, this disruption is unlikely to be the result of sample processing for electron microscopy. Abbreviations: AD: Adipocite; BC: Bacterial cells in gut lumen; ED: Epithelial disruption; GL: Gut lumen; ME: Midgut epithelium. Magnifications: A: 15,000x; B: 10,000x; C: 3,000x; D: 5,000x.

Figure 5. Effects of the physiological status of the mosquitos on the performance of the device. DABS

are efficient at killing mosquitoes through different physiological stages. (A) A significant majority

of blood fed mosquitoes exposed to toxic devices die after 72 hours of exposure. When presented

with non-toxic device almost all survive. (B) All parous mosquitoes exposed to toxic devices die after

48 hours of exposure. When presented with non-toxic device almost all survive. Box plot indicating

663   median 25% and 75% quartiles. Error bars indicate max and min values. Each dot indicates an

664   independent count of mosquitoes (y axis) exposed to control devices (blue) or toxic devices (red) at

665   different time points (x axis).

Figure 6. Shelf life of the device. DABS are still effective after (A) 38 days, (B) 80 days and (C) 118

days of storage in insectary conditions. Box plot indicating median 25% and 75% quartiles. Error bars

indicate max and min values. Each dot indicates an independent count of mosquitoes (y axis) exposed

to control devices (blue) or toxic devices (red) at different time points (x axis).

Figure 7: Mortality of mosquitoes when exposed to DABS over time (Series 2.1). Mosquitos

were exposed to DABS for 24 hours and observed afterwards; mosquito mortality was calculated

immediately after the exposure period, and during laboratory observation at 48 hours and 72 hours.

Mean control and experimental house mortalities are shown as bars, and standard deviation as error

lines. Points indicating the mortality from each individual replicate are overlaid on each experimental

condition.

Figure 8: Mortality of mosquitoes when exposed to DABS for 48 hours (Series 2.2).

Mosquitos were exposed to DABS for 48 hours; mosquito mortality was calculated immediately after

the exposure period. Mean control and experimental house mortalities are shown as bars, and standard

deviation as error lines. Points indicating the mortality from each individual replicate are overlaid on

each experimental condition.

Figure 9: Mortality of mosquitoes when exposed to DABS for 48 hours in the presence of

alternative sugar source (Series 2.3). Mosquitos were exposed to DABS for 48 hours in the presence

of a competing attractant; mosquito mortality was calculated immediately after the exposure period.

Mean control and experimental house mortalities are shown as bars, and standard deviation as error

lines. Points indicating the mortality from each individual replicate are overlaid on each experimental

condition and time point.

<u>Supplemental table</u>

Table 1: Comparison of Results Across Semi-field Experimental Series Mean and (standard error) of 48-hour mosquito mortality from 6-replicate series are given. Mean mortality was compared across trial series using a t-test.			
Experimental Conditions		48-hour Mortality	p-value
Control	24-hour DABS exposure (Series 1)	5.5% (2.4)	0.151
	48-hour DABS exposure (Series 2)	11.7% (2.8)	
Treatment	24-hour DABS exposure (Series 1)	38.9% (3.9)	<0.001
	48-hour DABS exposure (Series 2)	91.5% (3.8)	
Control	48-hour DABS exposure (Series 2)	11.7% (2.8)	0.671
	48-hour DABS exposure with competing attractant (Series 3)	14.1% (4.1)	
Treatment	48-hour DABS exposure (Series 2)	91.5% (3.8)	0.784
	48-hour DABS exposure with competing attractant (Series 3)	89.6% (4.5)	

697