Ingested insecticide to control *Aedes aegypti*: Developing a novel dried attractive toxic sugar bait device for intra-domiciliary control

Rachel Sippy[1,2,†], Galo Rivera[3,a,†], Valeria Sanchez[1,4], Froilán Heras Heras[1,4], Bianca Morejón[3,b], Efraín Beltrán Ayala[4], Robert S. Hikida[5], María A. López-Latorre[3,c], Alex Aguirre[3], Anna M. Stewart-Ibarra[1,5,d,*], David A. Larsen[6,*], Marco Neira[3,*]

[1] Institute for Global Health & Translational Science, SUNY-Upstate Medical University, Syracuse, NY, USA

[2] Department of Geography, University of Florida, Gainesville, FL, USA

[3] Center for Research on Health in Latin America, Pontificia Universidad Católica del Ecuador, Quito, Ecuador

[4] Unidad Académica de Ciencias Químicas y de la Salud, Universidad Técnica de Machala, Machala, Ecuador

[5] Department of Medicine and Department of Public Health and Preventative Medicine, SUNY-Upstate Medical University, Syracuse, NY, USA

[6] Department of Public Health, Syracuse University, Syracuse, NY, USA

[a] Current affiliation: Vector Biology Group, Max Plank Institute for Infection Biology, Berlin, Germany

[b] Current affiliation: Biology Division, College of Arts and Sciences, Kansas State University, Manhattan, KS, USA

[c] Current affiliation: Medical School, College of Health Sciences, Universidad de las Américas, Quito, Ecuador

[d] Current address: InterAmerican Institute for Global Change Research (IAI), Montevideo, Uruguay

[†] These authors contributed equally to the work

*Corresponding authors:
 Anna Stewart: astewart@dir.iai.int
 David Larsen: dalarsen@syr.edu
 Marco Neira: mvneira@puce.edu.ec

Abstract (350 words)

Background: Illnesses transmitted by *Aedes aegypti* (dengue, chikungunya and Zika) comprise a considerable global burden; mosquito control is the primary public health tool to reduce disease transmission. Current interventions are inadequate and insecticide resistance threatens the effectiveness of these options. Dried attractive bait stations (DABS) are a novel mechanism to deliver insecticide to *Ae. aegypti*. The DABS are a high-contrast 28 inch^2 surface coated with dried sugar-boric acid solution that elicits an ingestion response from *Ae. aegypti* landing on the surface. The study presents the development of the DABS and tests of their impact on *Ae. aegypti* mortality in the laboratory and a series of semi-field trials. *Methods*: We conducted multiple series of laboratory and semi-field trials to assess the survivability of *Ae. aegypti* mosquitoes exposed to the DABS. For laboratory experiments we assessed the lethality, the killing mechanism, and the shelf life of the device through controlled experiments. In the semi-field trials, we released laboratory-reared female *Ae. aegypti* into experimental houses typical of peri-urban tropical communities in South America in three trial series with six replicates each. Laboratory experiments were conducted in Quito, Ecuador, and semi-field experiments were conducted in Machala, Ecuador – an area with abundant wild populations of *Ae. aegypti* and endemic arboviral transmission. *Results*: In the laboratory, complete lethality was observed after 48 hours regardless of physiological status of the mosquito. The killing mechanism was determined to be through ingestion, as the boric acid disrupted the gut of the mosquito. In experimental houses, total mosquito mortality was greater in the treatment house for all series of experiments ($p<0.0001$). *Conclusions*: The DABS devices were effective at killing female *Ae. aegypti* under a variety of laboratory and semi-field conditions. DABS are a promising intervention for interdomiciliary control of *Ae. aegypti* and arboviral disease prevention.

66   **Keywords:** *Aedes aegypti*, vector control, toxic sugar bait, attractive bait, semi-field, dengue,

67   arbovirus, ATSB

68   Background

Arboviral illnesses, including dengue, chikungunya, yellow fever, and Zika, are major

contributors to morbidity and mortality in the tropics and subtropics. The burden is particularly

apparent in Central and South America: between 2010—2018, the estimated annual number of

dengue cases in the region ranged from 500,000 to 2,400,000 [1], and since 2013 it is estimated that

there have been a total of >1 million cases of chikungunya [2], and 3-4 million cases of Zika [3].

The viruses causing these diseases are spread mainly by the mosquitoes *Aedes* (*Ae.*) *aegypti* and

Ae. albopictus, with *Ae. aegypti* serving as the principal vector in many South American countries,

including Ecuador [3]. Due to the lack of commercially-available vaccines for most human

arboviral diseases, prevention efforts focus on vector surveillance and control methods [4].

Vector control relies heavily on contact-based insecticides, which are available in four main

classes: organophosphates, pyrethroids, carbamates, and organochlorines. Indoor residual spraying

is a common approach to vector control, for which twelve insecticides are available and approved

for human use. This small number of approved insecticides constitutes an impediment for the

implementation of effective vector control strategies (such as pesticide rotation cycles) aimed at

decreasing the development of resistance to any single insecticide [5]. As a result, pesticide

resistance has become a major limitation for current vector control strategies, and is widespread in

South American countries [6–8]. Our current reliance on a few chemical molecules to control *Ae.*

aegypti is an increasingly flawed strategy, as evidenced by the proliferation of this disease vector

across the globe and increasing arbovirus epidemics [9].

In contrast to the contact-based insecticide approach of the public health sector, the

agricultural industry has focused on ingested insecticides for pest control. The use of ingested

insecticides could be applied in disease control programs and interventions if disease vectors are

successfully led to ingest the insecticide. One solution, attractive toxic sugar baits (ATSB), exploits the nectar-feeding behavior of mosquitoes [10,11] to deliver the insecticide. An ATSB uses a mixture of a lethal agent with sugar water and sometimes an additional attractant [12]. ATSBs have been tested for *Anopheles* spp. [13–17], *Culex* spp. [15,16,18,19], *Ae. albopictus* [20–23], and other vector or nuisance species [16] with a variety of attractants, baits, active ingredients, designs, and placement strategies. Although laboratory bioassays demonstrate that ATSBs are toxic to *Ae. aegypti* [16,24,25], semi-field and field evaluations have had poor results in reducing *Ae. aegypti* populations [26,27], indicating that ATSB devices must be carefully designed and tested for each target species [12].

Compared to other mosquito species, *Ae. aegypti* appear to have a lower propensity for sugar-feeding, preferring human blood meals instead [11]. Despite this, *Ae. aegypti* females will readily feed on sugar in the laboratory, and often feed on plant sugars in the wild [28–30]. However, traditional attractive sugar bait strategies that rely only on sugar as an attractant are likely insufficient to "lure" female *Ae. aegypti* in the natural environment

Herein we present the development of dried attractive bait stations (DABS) (Figure 1), and show results from laboratory and semi-field experiments. In the laboratory we first identified the lethality of DABS (Series 1.1), aimed to identify the killing mechanism of the DABS (Series 1.2), assessed how the physiological status altered the effectiveness of DABS (Series 1.3), and assessed the shelf life of the DABS (Series 1.4). In the semi-field trials, we sought to determine the timing of mosquito mortality (Series 2.1), assess the relationship between DABS exposure time and mosquito mortality (Series 2.2), and to demonstrate these effects in the presence of competing attractants (Series 2.3).

Results

Laboratory experiments

116   *Series 1.1: Effects of DABS exposure on mosquito survival*
117

Mosquitoes exposed to DABS presented 55% survival probability reduction in the first 24

hours post-exposure, and an additional 45% reduction between 24h and 48h post-exposure,

resulting in 100% mortality by the end of the experiments (Figure 2). In contrast, mosquitoes

exposed to control devices presented only a 1% reduction in their survival probability during the

first 24h. This value increased to 2% during the second 24h period, resulting in 98% survival

probability by the end of the experiment. Differences between the survival curves of toxic and

control treatments were highly significant (p<0.001).

Series 1.2: Characterization of the biological mode of action of the device.
127

After 48 hours, mosquitoes that could still feed (*i.e.* mosquitoes with an intact proboscis),

presented 100% mortality when exposed to the toxic device, and 3% mortality when exposed to the

non-toxic control device. Mosquitoes that could not feed (those with ablated proboscis) presented

38% mortality regardless of inclusion in the ATSB-device condition or control non-toxic condition.

Significant differences were found between the four treatments (p<0.001). Post-hoc pairwise

comparison determined that (a) the mortality of ablated mosquitoes exposed to toxic devices was

not significantly different to the mortality of ablated mosquitoes exposed to control devices, and (b)

the mortality of ablated mosquitoes was significantly different from the mortality of whole

mosquitoes exposed to toxic devices and whole mosquitoes exposed to control devices (Figure 3).

Mosquitoes that had ingested toxic sugar solution presented histological abnormalities in

the posterior midgut. Electron micrographs revealed disruptions in the continuity of the gut

epithelium (Figures 4A, 4C). In addition, we found abnormal adipocytes that we hypothesize are

undergoing a process of necrosis (Figures 4E, 4F), and an increase in both the size and number of

basal infolds in the gut epithelial cells. We hypothesize that boric acid ingestion is the cause of

these pathological changes, which contributed to the mortality observed in specimens exposed to

toxic devices. Microscopic images of individuals exposed to control devices presented none of these

pathologies on the posterior midgut (Figures 4B, 4D).

Series 1.3: assessment of mosquito biological status on DABS effectiveness
147

Mosquitoes with both physiological statuses evaluated (blood fed and parous) presented a

lower survival probability when exposed to toxic devices than when exposed to non-toxic control

devices. Blood fed females' survival probability dropped 13% during the first 24h interval, 22%

during the second 24h interval, and 55.6% during the last interval, resulting in 8.89% survival

probability by the end of the experiment. Blood fed non-toxic mosquitoes exposed to control

conditions resulted in 90% survival probability at the end of 72 hours after having dropped 9%, 1%,

and 0% during the three 24-hour intervals, respectively (Figure 5A). Differences between control

and toxic treatment survival curves were significant ($p < 0.001$)

Parous female mosquitoes presented a 66% decline in survival probability during the first

24 hours and a 0% survival probability after 48 hours of exposure to the ATSB devices (Figure 5B).

These results are significantly different ($p < 0.001$) from the non-toxic control group, that showed a

drop in survival probability of 2% during the first 24h, resulting in 98% survival after 48 hours of

exposure.

Series 1.4: Assessment of shelf-life of the DABS device
163

Mosquitoes exposed to toxic devices stored for 38 days showed 0% survival probability

after the first 24 hour interval; mosquitoes exposed to control conditions showed 96% survival

through 48 hours (Fig. 6A). Differences in survival between conditions were highly significant ($p <$

0.001). Mosquitoes exposed to toxic devices stored for 80 days showed 16% survival probability

after the first 24 hour period and 0% survival probability after 48 hours; mosquitoes exposed to

control conditions showed 97% survival probability after 48 hours (Fig. 6B). Differences in survival

between conditions were highly significant (p < 0.001). Mosquitoes exposed to toxic devices stored

for 118 days showed 95% survival probability after the first 24-hour period, 64% survival during

the second 24-hour interval, and 35% survival probability after 72 hours; mosquitoes exposed to

control conditions showed 96% survival probability after 72 hours (Fig. 6C). Differences in survival

between conditions were highly significant (p = 0.001).

Semi-field experiments

The results of Series 2.1 (24 hours of exposure to DABS) are shown in Figure 7. Mosquito

mortality was 0—6% in the control and 17—57% in the treatment house after 24 hours (p<0.001).

At 48 hours, mortality was 0—18% in the control and 22—51% in the treatment house (p=0.003).

At 72 hours, mortality was 0—4% in the control and 0—2% in the treatment house (p=0.458). The

cumulative mortality of the control was 4—18% and 54—98% in the treatment house (p<0.001).

Following 24 hours of exposure to DABS in the house, most mosquito mortality was observed

within the first 48 hours of the experiment, with no difference in mosquito mortality after this time

period.

The results of Series 2.2 (48 hours of exposure to DABS) are shown in Figure 8. Mosquito

mortality was 2—23% in the control and 77—100% in the treatment house (p<0.001), indicating

high mortality from 48 hours of exposure to DABS in the treatment houses.

The results of Series 2.3 (48 hours of exposure to DABS in the presence of competing

attractant) are shown in Figure 9. Mosquito mortality was 2—33% in the control and 68—100% in

the treatment house (p=0.006), indicating that DABS results in high mortality even in the presence

of a competing attractant.

When comparing the results of 24 hours (Series 2.1) to 48 hours of exposure (Series 2.2), 48

hours of exposure results in higher mortality at 48 hours (p<0.001) in the treatment group (Table 1),

with no difference in the control groups (p=0.151). When comparing 48 hours of exposure to DABS

only (Series 2,2) and 48 hours of exposure to DABS in the presence of a competing attractant

(Series 2.3), there is no effect of a competing attractant on the effect of DABS on mosquito

mortality (p=0.784) in the treatment group (Table 1). High mortality from 48 hours of DABS

exposure was observed despite the presence of a competing attractant.

Discussion

These experiments demonstrate that DABS can strongly impact the mortality of female *Ae.*

aegypti under laboratory and semi-field conditions. In these settings, we show that mortality occurs

within the first 48 hours of exposure to our devices. In addition, DABS attract and kill *Ae. aegypti*

even in the presence of an alternative sugar source. To the best of our knowledge, this device is the

only known "dry" ATSB. The simple and economic design lends itself to in-home use in resource

limited settings where *Ae. aegypti* target human hosts and transmit dangerous arboviruses.

Our assessment of the biological action of the devices provides an insight into the

mechanism by which low concentrations of boric acid affect *Ae. aegypti*. We determined that boric

acid enters the insect´s body by ingestion, further supporting the notion that this inorganic pesticide

acts as a stomach poison, as previously suggested (Xue R-D, Kline DL, Ali A, Barnard DR.

Application of boric acid baits to plant foliage for adult mosquito control. J Am Mosq Control

Assoc. 2006;22: 497–500. Xue R-D, Barnard DR. Boric acid bait kills adult mosquitoes (Diptera:

Culicidae). J Econ Entomol. 2003;96: 1559–1562). Based on our electron microscopy analysis, we

hypothesize that the ingestion of boric acid disrupts the integrity of the gut epithelium.

Considering that the proposed mechanism by which boric acid exerts its toxic effect (gut

disruption) is markedly different from the neurotoxic mechanism by which most traditional

pesticides cause mortality, we propose that our devices have the potential to act as efficient

complementary tools to combat the spread of resistance to traditional pesticides. By combining the

use of DABS with traditional pesticides in the same areas, it would be possible to target two

different and crucial systems (namely, the nervous and digestive systems) in the insect´s body

simultaneously, thereby reducing the mosquito´s probability of survival.

We observed significant mortality of blood fed female *Ae. aegypti* exposed to the DABS

device, albeit at lower rate than starved females. Interestingly, the largest drop in survival

probability in blood fed females is observed between 48h and 72h post-exposure to the device (Fig

5), suggesting that after 48h, females have already used imbibed blood for the development of eggs

and are keen to search for further meals. Based on this evidence, it is plausible to suggest that if

employed in the field, DABS devices may be efficient in killing female mosquitoes of various

physiological statuses, including females that have already ingested blood – a particularly important

group for disease transmission.

Novel vector control methods have the potential to serve as critical tools in the public health

effort to control persistent and emerging vector borne diseases. Various designs of ATSBs have had

promising field trials for potential control of *Aedes albopictus*, *Anopheles* spp. and *Culex* spp. [15–

17,20,21,24]. Unfortunately, this success has not yet been achieved in efforts to control *Ae. aegypti*,

a vector of major public health importance. Previous research shows that several formulations of

ATSBs can achieve *Ae. aegypti* mortalities above 80% in laboratory settings [16,25], but these

ATSBs are ineffective in semi-field or field settings [26,27]. The principle barrier to their success

was an inability to sufficiently attract *Ae. aegypti* to ATSBs, rendering them ineffectual.

Particularly, ATSBs using synthetic floral-based attractants, while attractive to male *Ae. aegypti* in a

laboratory setting, do not function well when targeting *Ae. aegypti* in field evaluations [26]. An

ATSB with a host-based attractant (*i.e*. CO_2) was unable to meaningfully reduce adult populations

[27].

We hypothesize that our device attracts *Ae. aegypti* with strong visual cues (as opposed to a

chemical) as an attractant. *Ae. aegypti* are container breeders [31,32], that utilized tree holes in their

natural forested habitat before adapting to life in human civilization. The DABS device has a high-

contrast (black and white) 28-inch^2 surface to simulate a refuge for *Ae. aegypti* [33]. High contrast

coloring has similarly been integrated into prior trap designs and has been shown to improve

capture rates of *Ae. aegypti* [34]. We believe the high-contrast coloring of DABS draws *Ae. aegypti*

to land on the device.

These experiments have demonstrated the effectiveness of DABS on *Ae. aegypti* in

laboratory and semi-field experimental conditions. Our approach differs from most ATSB

approaches in two important ways: firstly, we use a dried sugar solution to elicit an ingestion

response while other ATSBs typically use liquid nectar-simulants [12,15,17,26]. We hypothesize

that the dried sugar solution is a key element in the effectiveness of DABS. Similar to other diptera

[35], *Ae. aegypti* are able to evaluate surfaces with their feet, and the "taste" of a landing surface

can either lead the mosquito to feed and ingest, or reject, the surface [36]. Additionally, the dried

sugar solution provides two operational advantages over liquid solutions: liquid solutions are more

difficult to manufacture, ship, and distribute than dried solutions, and the dried device can be

smaller and more easily deployed. Secondly, we use a visual rather than chemical attractant to lure

Ae. aegypti to the device. Chemical attractants add to the cost and decrease the shelf life of any

device. Previous research has questioned the ability of sugar solutions alone to attract mosquitoes

[26,37], leading to research on chemical attractant additives for ATSBs, but the use of chemical

attractants in ATSBs targeting *Ae. aegypti* have been unsuccessful [26,27]. We demonstrate that a

simple black-and-white visual attractant is a sufficient motivator for female *Ae. aegypti* to land on

the surface of DABS even in the presence of a competing oasis and alternative sugary food. Taken

together, we hypothesize that the visual cues attract the *Ae. aegypti* to land on the device, upon

which the presence of the dry sugar on the device's surface entices the insect to ingest it. When this

sugar solution is mixed with boric acid, ingestion results in insect mortality.

We propose that these encouraging results justify larger field trials of DABS in open-air

environments. We show that 48 hours of DABS exposure leads to high mosquito mortality when

used in the laboratory and in experimental houses reminiscent of peri-urban tropical housing.

Furthermore, we have established that the effectiveness of DABS for killing *Ae. aegypti* is

maintained even after prolonged storage periods, a characteristic that would facilitate their use in
semi-field and field conditions.

Semi-field trials are a crucial step to bring a scalable, marketable product to in-home field
testing. An in-home approach is ideal for control of *Ae. aegypti*, as the vector has an extremely
limited flight range, often spending its entire life within a single household [5,32,38]. Other
research with ATSBs has shown that end-users of these products prefer to have them placed indoors
[14]. The successful design and placement strategy of DABS used in our experiments indicate that
the device is ideal for in-home field testing.

Limitations
282

These experiments were conducted under laboratory and semi-field conditions, which can
only moderately emulate real-world/field conditions. Semi-field experiments were limited to
nulliparous females and we cannot be certain how DABS will affect gravid or blood-fed females or
males in an open-air environment—though it should be noted that DABS were equally effective in
attracting and killing blood-fed and nulliparous females under laboratory conditions. It is also
unclear if DABS would impact non-target insect species, such as butterflies or other pollinators,
though if DABS are limited to use inside the home, it is unlikely to affect these species. Although
DABS performed well in the presence of a competing attractant (100 grams of apples), it is unlikely
that the attractant used in our experiments are a realistic substitute for open-air field conditions. An
actual home will contain many competing attractants, including human hosts. It is difficult to know
if the success of DABS in semi-field conditions will be replicated in occupied homes in the field;
the number and placement of DABS may need to be modified. In addition, it is unclear how end
users will react to placement of DABS in their homes, although there is evidence that residents in
areas of high *Ae. aegypti* burden are willing to utilize numerous home-based mosquito control
products [39].

Conclusions

With careful design and device placement consideration, we have created a promising vector control device ready for large-scale trials to test its ability to control *Ae. aegypti* in natural conditions. We demonstrated that DABS are capable of attracting and killing female *Ae. aegypti* in experimental houses, and that 48 hours in the presence of DABS leads to high mortality among female *Ae. aegypti*. Importantly, DABS were efficient at killing female mosquitoes of diverse physiological statuses, and can attract and kill female *Ae. aegypti* even in the presence of a competing attractant. These devices have promising effectiveness and this semi-field evaluation of DABS is an important step forward in the development of novel devices for the control of *Ae. aegypti*.

<u>Materials and methods</u>

Aim and design

The general aim of our study was to assess the effectiveness of DABS at killing female *Ae. aegypti* under laboratory and semi-field experimental conditions. Our specific aims were (a) to determine the timing of mosquito mortality (Series 1), (b) to assess the relationship between DABS exposure time and mosquito mortality (Series 2), and (c) to demonstrate these effects in the presence of competing attractants (Series 3). These aims were tested in three series of six trial replicates to compare control and treatment experimental conditions.

Study setting

Laboratory experiments

Laboratory experiments were conducted at the Center for Research on Health in Latin

America (CISeAL, by its Spanish acronym), where they were reared and maintained under standard

insectary conditions: $28 \pm 1°C$ temperature, $80 \pm 10\%$ relative humidity, and a 12h:12h (L:D) light

cycle. Larvae were fed finely ground fish food. When required, mosquitoes were sexed during the

pupal stage. Adults were kept in 15x15x15 cm cages. For maintenance, adult mosquitoes were fed

10% sucrose solution *ad libitum*. For blood feeding, female adult mosquitoes were offered access to

a restrained female mouse. All mosquitoes were maintained under insectary conditions between 0

and 14 days after adult emergence before they were used for experiments. Mosquitoes referred to as

"starved" in this manuscript were deprived of access to sugar or blood (but not water) for 48h prior

to their use in experiments.

Semi-field trials

Assays were performed in experimental houses meant to emulate typical housing found in

areas with active dengue transmission. Photographs of the houses are available in an additional file

(see Additional file 1). The houses are constructed of wood and cane and are raised on a 1-meter

platform with walkways to improve structural integrity and facilitate window access; one window

on each house is equipped with window escape traps with sleeves to monitor escape behavior. The

dimensions of the houses are 3.85 meters (m) wide x 4.85 m long x 3 m high. Each house has three

windows (0.9 m wide $\times$ 0.6 m high) and one door (1.03 m wide $\times$ 3 m high). The house frames are

made of wood, untreated wooden plank flooring, walls of untreated cane, and a roof of zinc panels.

The window traps are 0.45 m long x 0.66 m wide x 0.45 m high. The houses are located on campus

at the Universidad Técnica de Machala in the city of Machala, Ecuador (3°15' S, 79°57' W), a

region with abundant wild populations of *Ae. aegypti* and endemic arbovirus transmission.

Experiments were conducted under ambient climate conditions (temperature range: 23.1—35.6° C,

mean temperature: 28.4° C, relative humidity range: 43.9—95.0%, mean relative humidity: 75.8%).

Each trial replicate was conducted with one control and one experimental house; the specific house
used as the experimental or control house was alternated upon each replicate.

Biological material

352
353

Aedes aegypti eggs were provided by the Center for Research on Health in Latin America
(CISeAL, by its Spanish acronym) at the Pontificia Universidad Católica del Ecuador. All strains
used in this work originated from Ecuador, and had been maintained in laboratory conditions since
2015. Laboratory work was performed with strains originally collected in Ecuador from the cities of
Guayaquil and Puerto Francisco de Orellana. Semi-filed work was performed with a strain
originally collected in the city of Machala.

Semi-field experiments
362

Hatching and rearing of *Ae. aegypti* for semi-field experiments were performed at the
Laboratory of Entomology at the Universidad Técnica de Machala. Considering this laboratory is
located in a region where *Ae. aegypti* actively reproduces and thrives, environmental conditions
(temperature: 28-32 °C; relative humidity: 60-80%) were not artificially controlled in our mosquito-
rearing facilities. A vacuum pressure system was used to synchronize egg hatching (one hour
exposure to obtain first stage larvae). Larvae were fed with finely ground fish food. At the pupal
stage, males and females were separated. Adults were kept in 15x15x15 cm cages. Adults were fed
on 10% sugar solution *ad libitum*. Each experimental semi-field experiment series used nulliparous
females aged 1—5 days and starved for 24 hours prior to experimental release.

Dried Attractive Bait Stations (DABS)

373
374

The DABS device consists of two concentric foam disks (an inner white disk one

centimeter in diameter, and an outer black disk eight centimeters in diameter). Experimental DABS

were impregnated with a 10% sucrose solution containing 1% boric acid as a lethal agent. Control

DABS were impregnated with 10% sucrose solution without boric acid. Photographs of the devices

are available in an additional file (see Supplementary Figure 2, Additional file 2).

Laboratory experiments
382

383   *Series 1.1: Survival assessment of mosquitoes exposed to the device*
384

To determine whether exposure to the DABS devices has an influence on adult mosquito

survival probability, we conducted an experiment in which groups of 30 adult female mosquitoes,

placed in a 15x15x15cm cage, were exposed during 48 hours to either an DABS device or a control

device (sugar solution but no boric acid). We replicated each experiment four times. The assessment

was repeated using each of the two laboratory strains described previously.

Series 1.2: Appraisal of biological mode of action of the device
392

To establish whether the toxic component of DABS needs to be ingested by the mosquitoes

in order to exert its effect, we presented the devices to cohorts of adult females that were unable to

ingest food due to the surgical ablation of their mouthparts. To establish these cohorts, individual

mosquitoes were first anesthetized by placing them at 4°C during 10-15 minutes. Anesthetized

specimens were individually placed under a dissection microscope and, using a human hair, we tied

a knot at the proboscis' proximal end in order to create a constriction that would impede the flow of

food. Subsequently, the part of the proboscis anterior to the knot was removed using micro-

dissection scissor (Supplemental Figure 3). Following the surgery, mosquitoes were left to rest for

24 hours before being used in any experiment. To control for the potential negative effect of the

anesthetizing procedure in mosquito survival, non-ablated mosquitoes used in control groups were

also placed at 4°C during 10-15 minutes, and allowed to recover during 24 hours before

experimental set-up.

We conducted the experiment with four separate cages, each with 20 starved mosquitoes

(Table 2 shows different treatments). We treated cage 1 with toxic DABS devices and used 20

ablated mosquitoes; cage 2 held non-toxic control devices and 20 ablated mosquitoes. We treated

cage 3 with toxic DABS devices and non-ablated mosquitoes; cage 4 held a non-toxic control

device and non-ablated mosquitoes. We assessed mortality in all groups at 24 and 48 hours of

exposure to the devices. We replicated the experiment three times.

We then conducted an experiment wherein 30 adult starved female mosquitoes were

introduced to a cage with a DABS device, and 30 adult starved female mosquitoes were introduced

to a cage with a non-toxic control device. We monitored cages for 24 hours, and removed dead

mosquitoes by aspiration every hour from the cages. Using a dissection microscope, we removed

the legs, head, and wings of every dead specimen onto a drop of 70% ethanol. Through this process

we gently disrupted the abdominal cuticle to permit the exposure of internal tissues to the fixative.

Afterwards we fixed individual mosquitoes in a solution containing 2.5% glutaraldehyde, 2.5%

paraformaldehyde in 0.1M cacodylate buffer (pH 7.4), and stored them at 4°C for 72 hours. We

then washed specimens in cacodylate buffer with 0.1M sucrose overnight. Post-fixing was achieved

by leaving the specimens for two hours at 4°C in 2% osmium tetroxide in 0.1 cacodylate buffer (pH

7.4). Subsequently, individuals were stained using 2% uranyl acetate and left to rest for three hours

in the dark at room temperature. Tissues were later dehydrated through a series of ethanol baths

(50%, 70%, 95%, 100%). Afterwards, they were placed in propylene oxide for 30 minutes, then in a

1:1 volume propylene oxide resin mixture (Epon 812, Araldite 502, dodecenyl succinic anhydride,

benzyl dimethylamine) for one hour and later, one more volume of resin was added and left on a

rotator overnight. Finally, mosquitoes were embedded in resin and incubated at 60°C for 24 hours.

Resin were stained using 2% uranyl acetate. We then utilized a transmission electron microscope to

observe specimens and obtain micrographs of interesting tissues.

Series 1.3: Effects of the physiological status of the mosquitoes on the performance of
DABS
432

We examined two different physiological statuses using mated starved female adult

mosquitoes, namely blood fed and parous. We established females deemed as "blood fed" by

selecting blood-engorged individuals immediately after a blood meal. We established females

deemed as "parous" by first blood feeding and subsequently maintaining mosquitoes for 7 days

under insectary conditions in order to ensure that they had oviposited before being used for

experimentation. We set up two cages for each of the defined physiological statuses, each with 30

mosquitoes. One cage exposed the mosquitoes to an ATSB device, and the other held a control non-

toxic device. We gathered survival data at 24 and 48 hours following introduction to the cages, and

replicated these experiments three times.

Series 1.4: Shelf-life of the device
444

445   In order to determine the shelf life of ATSB devices, toxicity tests were performed using

446 devices which had been stored for 38, 80, and 118 days after their production. For storage, devices

447 were individually wrapped inside a sealed plastic bag and placed in an incubator at $28 \pm 2°C$ and 80

448 $\pm 10\%$ relative humidity. We conducted three replicates of previously described experiments for

449 each storage time.

Semi-field trials
451
452

453 *Series 2.1: 24 hours of DABS exposure in experimental houses*
454

Each house contained four DABS devices (control or treatment DABS as appropriate)

suspended on strings attached to the roof of the house at a height of 30—50cm above ground and

approximately 30cm from the nearest wall. Diagrams of the experimental layout are available in an

additional file (see Supplementary Figure 3, Additional file 2). For each trial replicate, 50 female

Ae. aegypti were released into each house through the escape window sleeve (release time

11:00a.m.—2:00p.m.). Twenty-four hours after release, dead mosquitos were collected from the

floor and window escape traps in each house, and the remaining live mosquitoes were captured with

a hand-held aspirator (Prokopack, John W. Hock Company, Gainesville, USA). All live mosquitoes

were labeled by experimental group and observed for 48 additional hours in laboratory cages (under

laboratory conditions with food available). Mortality was calculated for 24 hours, 48 hours, and 72

hours. Six trial replicates were performed for Series 1.

Series 2.2: 48 hours of DABS exposure in experimental houses
468

Each house contained four DABS devices (control or treatment DABS as appropriate) and

two sources of water (wet cotton in a black plastic bucket). Photographs of the water source are

available in an additional file (see Supplementary Figure 3, Additional file 2). For each trial

replicate, 50 female *Ae. aegypti* were released into each house through the escape window sleeve

(release time 8:00a.m.—11:00a.m.). Forty-eight hours after release, dead mosquitos were collected

in each house and remaining live mosquitoes were captured with an aspirator. Mortality was

calculated for 48 hours. Six replicates were performed for Series 2.

Series 2.3: 48 hours of DABS exposure in experimental houses with competing attractant
478

Each house contained four DABS devices (control or treatment DABS as appropriate), two

sources of water (wet cotton in a black plastic bucket), and 100 grams of peeled, cut apples in a dish

placed on a chair in the center of the house as a competing attractant. Photographs of the competing

attractant are available in an additional file (see Supplementary Figure 3, Additional file 2).

Recently emerged female *Ae. aegypti* rely on sugar meals for energy, these meals may include aging

fruit and female *Ae. aegypti* will feed on fructose (as is found in apples). For each trial replicate, 50

female *Ae. aegypti* were released into each house through the escape window sleeve (release time

9:00a.m.—12:00p.m.). Forty-eight hours after release, dead mosquitos were collected in each house

and remaining live mosquitoes were captured with an aspirator. Mortality was calculated for 48

hours. Six replicates were performed for Series 3.

490 Statistical analyses
491

For each replication of the Series 1.1 experiments, we plotted and analyzed interval

censored survival data and non-parametric likelihood estimates. We then used a log-rank hypothesis

test to compare the survival distributions of the two treatments. In Series 1.2, we examined the

normal distribution of the data with Kolmogorov-Smirnov and Shapiro-Wilk tests. We assessed

treatment group using an analysis of variance and a post-hoc's test to determine ranks. For Series

1.3 and 1.4, we plotted and analyzed censored survival data with a logrank hypothesis test to

compare the survival distributions of experiment and control treatments. In Series 2.1—2.3,

mosquito mortality data from each series were compared using a two-tailed paired t-test (paired by

replicate). Mean mosquito mortality was compared across series using a two-tailed t-test. Data were

analyzed using Excel (Microsoft, Redmond, USA) and R (R Foundation for Statistical Computing,

Vienna, Austria) with the survival package [42,43].

504 <u>List of abbreviations</u>

505 Ae.=Aedes

506 ATSB=attractive toxic sugar bait

507 DABS=dried attractive bait stations

508 cm =centimeters

509 m=meters

Declarations

Ethics approval:

513
514   These study protocols were found to be exempt from Institutional Review Board review by

515 Syracuse University Institutional Review Board.

Consent for publication:

516
517   Not applicable

Availability of data and material:

518
519   The datasets used and/or analysed during the current study are available from the

520 corresponding author on reasonable request.

Competing interests:

521
522   ASI, DL, and MN hold a patent on the DABS technology.

Funding:

523
524   Laboratory work was funded by Pontificia Universidad Católica del Ecuador´s Internal

525 Research Grant L13234, awarded to MN. Semi-field work was funded by a seed grant from

526 Syracuse University, awarded to DL.

Author contributions:

527
528

MN, ASI, and DL designed the study.

MN, ASI, DL, and EBA provided study materials.

GR, ML, AA, BM VS and FHH performed the experiments.

RH performed electron microscopy work

RS and GR conducted data analysis.

RS, GR, MN, VS, and FHH drafted the manuscript.

All authors reviewed and approved the final draft of the manuscript.

Acknowledgements:
537

The authors would like to thank Cinthya Cueva Aponte for her assistance with study

documentation, Chad Ward and Matt Kovacs for assistance with mosquito rearing, Mario Grijalva

for his logistical support for the performance of electron microscopy, Felipe Andrade and Phillip

Huegli for providing useful recommendations on R programming, and Santiago Cadena for his

support in insectary maintenance.

Additional files

Additional file 1 (pdf)

Supplementary Figure 1: Experimental Houses

Additional file 2 (pdf)

Supplementary Figure 2: Experimental Layout

Additional file 3 (pdf)

Supplemental Figure 2: Feeding disruption procedure. A) Anesthetized individual with

whole proboscis. B) Human hair tied at the proximal end of the proboscis. C) Micro-dissection

scissor removal of proboscis' segment anterior to the knot. D) Feeding disrupted individual.

References

1. PAHO, WHO. Epidemiological Update: Dengue. 2019;1–14.

2. Leta S, Jibat Beyene T, De Clercq EM, Amenu K, Revie CW, Kraemer MUG. Global

risk mapping for major diseases transmitted by Aedes aegypti and Aedes albopictus HHS Public

Access. Int J Infect Dis. 2018;67:25–35.

3. Boyer S, Calvez E, Chouin-Carneiro T, Diallo D, Failloux AB. An overview of mosquito

vectors of Zika virus. Microbes Infect. 2018;20:646–60.

4. P. O, F. F, A. K, L. B, R. V, A.C. S, et al. Improved tools and strategies for the

prevention and control of arboviral diseases: A research-to-policy forum. PLoS Negl Trop Dis.

2018;12:e0005967.

5. Harrington LC, Scott TW, Lerdthusnee K, Coleman RC, Costero A, Clark GG, et al.

Dispersal of the dengue vector Aedes aegypti within and between rural communities. Am J Trop

Med Hyg. 2005;72:209–20.

6. Hemingway J, Ranson H. Insecticide Resistance in Insect Vectors of Human Disease.

Annu Rev Entomol. 2000;45:371–91.

7. Chanda E, Kamuliwo M, Sikaala CH, Thomsen EK, Ranson H, Hemingway J, et al. An

operational framework for insecticide resistance management planning. Emerg Infect Dis.

2016;22:773–9.

8. Ryan SJ, Mundis SJ, Aguirre A, Lippi CA, Beltrán E, Heras F, et al. Seasonal and

geographic variation in insecticide resistance in Aedes aegypti in southern Ecuador. PLoS Negl

Trop Dis. 2019;13:e0007448.

9. Kraemer MUG, Sinka ME, Duda KA, Mylne A, Shearer FM, Barker CM, et al. The

global distribution of the arbovirus vectors Aedes aegypti and Ae. albopictus. Elife. 2015;4:e08347.

10. Yuval B. The other habit: sugar feeding by mosquitoes. Bull Soc Vector Ecol [Internet].

Society for Vector Ecology; 1992;17:150–6.

11. Foster WA. Mosquito sugar feeding and reproductive energetics. Annu Rev Entomol.

1995;40:443–74.

12. Fiorenzano JM, Koehler PG, Xue R De. Attractive toxic sugar bait (ATSB) for control

of mosquitoes and its impact on non-target organisms: A review. Int J Environ Res Public Health.

2017;14.

13. Tenywa FC, Kambagha A, Saddler A, Maia MF. The development of an ivermectin-

based attractive toxic sugar bait (ATSB) to target Anopheles arabiensis. Malar J. BioMed Central;

2017;16:1–10.

14. Maia MF, Tenywa FC, Nelson H, Kambagha A, Ashura A, Bakari I, et al. Attractive

toxic sugar baits for controlling mosquitoes: A qualitative study in Bagamoyo, Tanzania. Malar J.

2018;17:1–6.

15. Stewart ZP, Oxborough RM, Tungu PK, Kirby MJ, Rowland MW, Irish SR. Indoor

application of attractive toxic sugar bait (ATSB) in combination with mosquito nets for control of

pyrethroid-resistant mosquitoes. PLoS One. 2013;8.

16. Qualls WA, Müller GC, Revay EE, Allan SA, Arheart KL, Beier JC, et al. Evaluation of

attractive toxic sugar bait (ATSB)-Barrier for control of vector and nuisance mosquitoes and its

effect on non-target organisms in sub-tropical environments in Florida. Acta Trop. 2014;131:104–

10.

17. Qualls WA, Müller GC, Traore SF, Traore MM, Arheart KL, Doumbia S, et al. Indoor

use of attractive toxic sugar bait (ATSB) to effectively control malaria vectors in Mali, West Africa.

Malar J. 2015;14:301.

18. Qualls WA, Scott-Fiorenzano J, Müller GC, Arheart KL, Beier JC, Xue R-D.

Evaluation and Adaptation of Attractive Toxic Sugar Baits For Culex tarsalis and Culex

quinquefasciatus Control In The Coachella Valley, Southern California . J Am Mosq Control Assoc.

2017;32:292–9.

19. Müller GC, Junnila A, Schlein Y, Ller GCM. Effective Control of Adult Culex pipiens

by Spraying an Attractive Toxic Sugar Bait Solution in the Vegetation Near Larval Habitats. Source

J Med Entomol J Med Entomol. 2010;47:63–6.

20. Revay EE, Müller GC, Qualls WA, Kline DL, Naranjo DP, Arheart KL, et al. Control of

Aedes albopictus with attractive toxic sugar baits (ATSB) and potential impact on non-target

organisms in St. Augustine, Florida. Parasitol Res. 2014;113:73–9.

21. Junnila A, Revay EE, Müller GC, Kravchenko V, Qualls WA, Xue R de, et al. Efficacy

of attractive toxic sugar baits (ATSB) against Aedes albopictus with garlic oil encapsulated in beta-

cyclodextrin as the active ingredient. Acta Trop. 2015;152:195–200.

22. Scott-Fiorenzano JM, Fulcher AP, Seeger KE, Allan SA, Kline DL, Koehler PG, et al.

Evaluations of dual attractant toxic sugar baits for surveillance and control of Aedes aegypti and

Aedes albopictus in Florida. Parasites and Vectors. Parasites & Vectors; 2017;10:1–9.

23. Scott JM, Seeger KE, Gibson-Corrado J, Muller GC, Xue R De. Attractive Toxic Sugar

Bait (ATSB) Mixed With Pyriproxyfen for Control of Larval Aedes albopictus (Diptera: Culicidae)

Through Fecal Deposits of Adult Mosquitoes. J Med Entomol. 2017;54:236–8.

24. Khallaayoune K, Qualls WA, Revay EE, Allan SA, Arheart KL, Kravchenko VD, et al.

Attractive Toxic Sugar Baits: Control of Mosquitoes With the Low-Risk Active Ingredient

Dinotefuran and Potential Impacts on Nontarget Organisms in Morocco. Environ Entomol.

2013;42:1040–5.

25. Lea AO. Sugar-baited insecticide residues against mosquitoes. Mosq News. 1965;25:65.

26. Fikrig K, Johnson BJ, Fish D, Ritchie SA. Assessment of synthetic floral-based

attractants and sugar baits to capture male and female Aedes aegypti (Diptera: Culicidae). Parasites

and Vectors. Parasites & Vectors; 2017;10:1–9.

27. Xue R-D, Ali A, Kline DL, Barnard DR. Field Evaluation of Boric Acid- and Fipronil-

Based Bait Stations Against Adult Mosquitoes. J Am Mosq Control Assoc. 2008;24:415–8.

28. Martinez-Ibarra JA, Rodriguez MH, Arredondo-Jiménez JI, Yuval B, Arredondo-

Jimenez JI, Yuval B. Influence of plant abundance on nectar feeding by Aedes aegypti (Diptera:

Culicidae) in southern Mexico. J Med Entomol. 1997;34:589–93.

29. Spencer CY, Pendergast TH, Harrington LC. Fructose variation in the dengue vector,

Aedes aegypti, during high and low transmission seasons in the Mae Sot region of Thailand. J Am

Mosq Control Assoc. 2005;21:177–81.

30. van Handel E, Edman JD, DAY JF, Scott TW, Clark GG, Reiter P, et al. Plant-sugar,

glycogen, and lipid assay of Aedes aegypti collected in urban Puerto Rico and rural Florida. J Am
Mosq Control Assoc. 1994;10:149–53.

31. Garcia-Rejon J, Loroño-Pino MA, Farfan-Ale JA, Flores-Flores L, Del Pilar Rosado-Paredes E, Rivero-Cardenas N, et al. Dengue virus-infected Aedes aegypti in the home environment. Am J Trop Med Hyg. 2008;79:940–50.

32. Kuno G. Review of the factors modulating dengue transmission. Epidemiol Rev. 1995;17:321–35.

33. Fay RW, Prince WH. A modified visual trap for Aedes aegypti. Mosq News. 1970.

34. Barrera R, Mackay AJ, Amador MA. An improved trap to capture adult container-inhabiting mosquitoes. J Am Mosq Control Assoc. 2013; 29:358–68.

35. Meunier N, Marion-Poll F, Rospars JP, Tanimura T. Peripheral coding of bitter taste in Drosophila. J Neurobiol. 2003;56:139–52.

36. Dennis EJ, Goldman O V., Vosshall LB. Aedes aegypti Mosquitoes Use Their Legs to Sense DEET on Contact. Curr Biol. 2019;29:1551-1556.e5.

37. Xue R-D, Barnard DR. Boric acid bait kills adult mosquitoes (Diptera: Culicidae). J Econ Entomol. 2003;96:1559–62.

38. Hemme RR, Thomas CL, Chadee DD, Severson DW. Influence of urban landscapes on population dynamics in a short-distance migrant mosquito: evidence for the dengue vector Aedes aegypti. PLoS Negl Trop Dis. 2010;4:e634.

39. Heydari N, Larsen D, Neira M, Beltrán Ayala E, Fernandez P, Adrian J, et al. Household Dengue Prevention Interventions, Expenditures, and Barriers to Aedes aegypti Control in Machala, Ecuador. Int J Environ Res Public Health. 2017;14:196.

Figures

Survival of Mosquitoes Exposed to DABS

676　*Figure 1. Attractive toxic sugar bait devices. A)* 3D model of the DABS devices. Diameters for
677　each of the foam sheet circles are shown.B) Photography of a DABS device used in the study.
Figure 2. Survival assessment of mosquitoes exposed to the device. All mosquitoes
(n=30) exposed to toxic devices die after 48 hours of exposure. When presented with non-
toxic device almost all survive. Box plot indicating median 25% and 75% quartiles. Error
bars indicate max and min values. Each dot indicates an independent count of mosquitoes
(y axis) exposed to control devices (blue) or toxic devices (red) at different time points (x
axis).
689
690
691

697
698
Figure 3. Uptake mechanism of the toxic component. Toxic effect is dependent on the ability of
699
mosquitoes to ingest the toxic component. When mosquitoes are able to ingest the toxic component
700
all mosquitoes (n =20) die after 48 hours (red). Mosquitoes with ablated mouthparts die equally
701
regardless of the toxic or non-toxic condition of the device (green and yellow). Almost all
702
mosquitoes with no ablation or toxic devices survive. Box plot indicating median 25% and 75%
703
quartiles. Error bars indicate max and min values. Each dot indicates an independent count of
704
mosquitoes (y axis) exposed to control devices (blue), control devices + mosquito mouthparts
705
ablation (yellow), toxic devices + mosquito mouthparts ablation, or toxic devices (red) at different
706
time points (x axis).
707
708
709

Figure 4. Histopathological effects on the midgut. Longitudinal sections of Ae. aegypti posterior midgut. Panels A, C, D: Mosquitoes exposed to toxic devices. Panel B: Mosquito exposed to non-toxic device. Specimens exposed to toxic devices showed disruptions in the gut integrity (ED, panel A). Because of the even distribution of adjacent bacterial cells in the gut lumen, this disruption is unlikely to be the result of sample processing for electron microscopy. Abbreviations: AD: Adipocite; BC: Bacterial cells in gut lumen; ED: Epithelial disruption; GL: Gut lumen; ME: Midgut epithelium. Magnifications: A: 15,000x; B: 10,000x; C: 3,000x; D: 5,000x.

Figure 5. Effects of the physiological status of the mosquitos on the performance of the device. DABS are efficient at killing mosquitoes through different physiological stages. (A) A significant majority of blood fed mosquitoes exposed to toxic devices die after 72 hours of exposure. When presented with non-toxic device almost all survive. (B) All parous mosquitoes exposed to toxic devices die after 48 hours of exposure. When presented with non-toxic device almost all survive. Box plot indicating median 25% and 75% quartiles. Error bars indicate max and min values. Each dot indicates an independent count of mosquitoes (y axis) exposed to control devices (blue) or toxic devices (red) at different time points (x axis).

Figure 6. Shelf life of the device. DABS are still effective after (A) 38 days, (B) 80 days and (C) 118 days of storage in insectary conditions. Box plot indicating median 25% and 75% quartiles. Error bars indicate max and min values. Each dot indicates an independent count of mosquitoes (y axis) exposed to control devices (blue) or toxic devices (red) at different time points (x axis).

Figure 7: Results of Series 2.1. Mosquitos were exposed to DABS for 24 hours and observed afterwards; mosquito mortality was calculated immediately after the exposure period, and during laboratory observation at 48 hours and 72 hours. Mean control and experimental house mortalities are shown as bars, and standard deviation as error lines. Points indicating the mortality from each individual replicate are overlaid on each experimental condition.

789 *Figure 8: Results of Series 2.2.* Mosquitos were exposed to DABS for 48 hours; mosquito
790 mortality was calculated immediately after the exposure period. Mean control and
791 experimental house mortalities are shown as bars, and standard deviation as error lines.
792 Points indicating the mortality from each individual replicate are overlaid on each
793 experimental condition.
794

799 *Figure 9: Results of Series 2.3.* Mosquitos were exposed to DABS for 48 hours in the
800 presence of a competing attractant; mosquito mortality was calculated immediately after the
801 exposure period. Mean control and experimental house mortalities are shown as bars, and
802 standard deviation as error lines. Points indicating the mortality from each individual
803 replicate are overlaid on each experimental condition and time point.
804