A Highly Conserved *Shh* Enhancer Coordinates Hypothalamic and Craniofacial Development

Zoe Crane-Smith[1], Jeffrey Schoenebeck[2], Katy A Graham[1], Paul S Devenney[1], Lorraine Rose[1], Mark
Ditzell[3], Eve Anderson[1,Ŧ], Joseph I Thomson[1], Natasha Klenin[4], Deborah M Kurrasch[4], Laura A
Lettice[1], and Robert E Hill[1,*]

[1] Medical Research Council Human Genetics Unit, Institute of Genetics and Molecular Medicine,
University of Edinburgh, Edinburgh EH4 2XU, United Kingdom

[2] The Roslin Institute and Royal (Dick) School for Veterinary Studies, University of Edinburgh, Easter
Bush, Midlothian, United Kingdom

[3] Cancer Research UK Edinburgh Centre, Institute of Genetics and Molecular Medicine, University
of Edinburgh, Edinburgh EH4 2XU, United Kingdom

[4] Department of Medical Genetics, Alberta Children's Hospital Research Institute & Hotchkiss Brain
Institute, University of Calgary, Calgary, AB T2N 4N1, Canada.

* Correspondence:
Robert Hill
bob.hill@igmm.ed.ac.uk

Keywords: *Shh* / enhancer / hypothalamic development / craniofacial formation / embryonic
development

[Ŧ] Present address – Cancer Research UK Beatson Institute, Garscube Estate, Switchback Rd,
Bearsden, Glasgow G61 1BD United Kingdom

Abstract

Enhancers that are conserved deep in evolutionary time regulate characteristics held in common
across taxonomic classes. Here, deletion of the highly conserved *Shh* enhancer SBE2 (*Shh* brain
enhancer 2) markedly reduced *Shh* expression within the embryonic brain specifically in the rostral
diencephalon; however, no abnormal anatomical phenotype was observed due to compensatory low
levels of expression mediated by a second enhancer. In contrast, a further reduction of *Shh* levels,
achieved by crossing the SBE2 deletion with the *Shh* null allele, disrupted brain and craniofacial
tissue; thus, linking SBE2 regulated *Shh* expression to multiple defects and further enabling the study
of the effects of differing levels of *Shh* on embryogenesis. Development of the hypothalamus,
derived from the rostral diencephalon, was disrupted along both the anterior-posterior (AP) and the
dorsal-ventral (DV) axes. Expression of DV patterning genes and subsequent neuronal population
induction were particularly sensitive to *Shh* expression levels, demonstrating a novel morphogenic
context for *Shh*. The role of SBE2, which is highlighted by DV gene expression, is to step-up
expression of *Shh* above the minimal activity of the second enhancer, ensuring the necessary levels of
Shh in a regional-specific manner. We also show that low *Shh* levels in the diencephalon disrupted
neighbouring craniofacial development, including mediolateral patterning of the bones along the
cranial floor and viscerocranium. Thus, SBE2 contributes to hypothalamic morphogenesis and
ensures there is coordination with the formation of the adjacent midline cranial bones that
subsequently protects the neural tissue.

Introduction

Long-distance enhancers that are conserved across multiple classes of vertebrates are
implicated in regulating early embryonic development (Nord et al., 2013). The *Shh* regulatory domain,
which extends 1Mb upstream of the gene contains a number of sequence elements identified by high
sequence conservation. These elements appear to be largely responsible for the typical pattern of *Shh*
expression observed at initial stages of organogenesis (Anderson et al., 2014; Jeong et al., 2008). The
deep conservation and early stages of expression mediated by these *Shh* enhancers suggest a
fundamental role in generating ancient, shared vertebrate characteristics. Studies to date show a
modular composition of the *Shh* regulatory domain and thus, for at least some of the *cis*-regulators,
there is a simple relationship of a single enhancer to a distinct spatial domain of expression (Jeong et
al., 2006; Lettice et al., 2014); although, in other instances, secondary enhancers provide compensatory
low levels of expression (Letelier et al., 2018; Tsukiji et al., 2014).

SHH is a signalling factor with a fundamental role in brain and craniofacial development. A
number of enhancers have been identified within the *Shh* regulatory domain that are responsible for
regulating expression in the midbrain and forebrain. Transgenic analysis indicates that the SBE2
enhancer (Jeong et al., 2008, 2006; Zhao et al., 2012) regulates *Shh* expression in the early developing
forebrain, from the zona limitans intrathalmica (ZLI) to the medial ganglionic eminence (MGE),
specifically designated the rostral diencephalon (Jeong et al., 2006; Zhao et al., 2012). Of particular
interest is a clinically relevant inactivating mutation in the SBE2 enhancer associated with features of
semilobar holoprosencephaly (Jeong et al., 2008). Holoprosencephaly (HPE) is a developmental
disorder of both the forebrain and midline facial structures that include microcephaly, midfacial
hypoplasia, cleft lip and palate, diabetes insipidus and moderate fusion of the hypothalamus
(Marcorelles and Laquerriere, 2010). More commonly, HPE is associated with heterozygous mutations
that inactivate the *Shh* gene (Chiang et al., 1996; Ming and Muenke, 2002). These data suggest that
disruption of SBE2 driven expression in the early developing forebrain may contribute to this disorder
(Jeong et al., 2008).

Here, we deleted the SBE2 enhancer to discern its overall contribution to *Shh* expression pattern
and to establish its requirement for mediating the embryonic phenotype. *Shh* expression is continuous
throughout the central nervous system, relying on a number of *cis*-regulators to generate the overall
pattern (Anderson et al., 2014). The deletion of SBE2 interrupted this spatial pattern by specifically
downregulating expression in the ventral rostral diencephalon; however, this reduction of SBE2 driven
Shh expression produced no overt phenotype. We confirmed the presence of a second enhancer (Sagai
et al., 2019) responsible for residual levels of *Shh* expression that compensated for the loss of SBE2 in
the rostral diencephalon. Further reduction in expression levels of *Shh* was attained by crossing the
SBE2 deletion mice to the line carrying the *Shh* null allele. The development of the hypothalamus and
of the surrounding tissue was disrupted in the compound mutants, linking diencephalic spatiotemporal
expression to multiple defects and enabling analysis of development at different expression levels. We
showed, firstly, that patterning in the hypothalamus along the AP axis was disrupted, but low levels of
Shh are sufficient for normal formation of the AP boundary. Further reductions in the levels of *Shh*,
however, led to misplaced spatial assignment of the hypophyseal lobes due to disruption of the AP
boundary. Secondly, gene expression along the DV axis was, in contrast, concentration sensitive and
consistent with SHH functioning as a morphogen along this developmental axis, with tissue patterning

and eventual neuronal fate reliant upon varying concentrations of *Shh* signalling. Thirdly, we found
that loss of SBE2 activity disrupts craniofacial development affecting medial lateral patterning of the
bones along the cranial floor and palatal shelf, revealing that SBE2 activity in the brain directly
regulates cranial and facial morphogenesis, providing further evidence of the ventral patterning role
played by *Shh* (Corman et al., 2018; Dworkin et al., 2016). These data confirm that signalling in the
brain has an influence upon craniofacial morphology and raises an interesting evolutionary relationship
that links patterning of a crucial region of the brain with the formation of the bony plate barrier
underlying this region in the skull. Hence, a single regulatory component coordinates the combined
function of directing brain development with the concomitant role of ensuring physical
neuroprotection.

Materials and methods

Embryo production and transgenic targeting
All animal studies were approved by the MRC IGMM 'Animal Care and Use Committee' and
according to the MRC 'Responsibility in the Use of Animals for Medical Research' (July 1993), EU
Directive 2010 and UK Home Office Project License no PPL 60/4424.
Embryos were harvested at various embryonic stages between E10.5-E17.5. For all
experiments described triplicate datasets were used, unless otherwise stated. Reduction in *Shh* levels
to 50%, in *Shh^{null/+}* embryos, is known to cause no phenotypic effects (Chiang et al., 1996).
Nevertheless, to allow for the use of littermate controls, differences between wild-type, *Shh^{ΔSBE2/+}* and
Shh^{null/+} embryos were assessed for each experiment described. As expected, no differences were
detected in between the three previously mentioned genotypes, as such all three genotypes were used
as control embryos.
The expression driven by potential enhancer region of interest was analysed in G0 transgenic
embryos using a lacZ construct which carried sequence of the region in question, a β-globin minimal
promoter and LacZ as has been previously described (Yee and Rigby, 1993) The DNA fragment of
interest was generated using PCR with the primers
GATCAT**GTCGAC**GCTCCAGGTACTGCTGTTCAG and
GATCAT**GCTGAC**ATGTGGATGGCAAGCATTGGC- the SalI restriction site used in the cloning
is highlighted in bold.
120
ATAC-seq and FAC sorting
Entire heads of GFP positive eGFP/Cre-*Shh* (Harfe et al., 2004) embryos were dissected and
single cell suspensions were made by incubating the dissected tissue in 1:5 dilution of trypsin:versene
at 37°C for 15 minutes. GFP positive cells were sorted using a fluorescence activated cell sorter and
ATAC-seq libraries were made from the positive cell population pooled across littermates as
previously described (Buenrostro et al., 2015). Subsequently libraries were subject to size selection to
exclude DNA fragments larger than 1 kb using SPRIselect magnetic beads (Beckman Coulter).
Samples were sequenced on a Illumina HiSeq 4000 platform to obtain 50 bp paired-end reads.
Resulting reads were trimmed using cutadapt paired-end trimming, and subsequently aligned to the
mm9 mouse genome using bowtie2 paired-end alignment. Unmapped reads and those mapping to the
mitochondrial genome were then removed and duplicate reads were filteres out using Picard
MarkDuplicates. Reads were shifted by +4bp for those mapping to the positive strand and -5bp for

those mapping to the negative strand. Broad peaks were then called, to identify open chromatin regions,
using MACS2 callpeak, using –g mm, -f BEDPE, -B –broads and cutoff 0.01 options.
135

SBE2 deletion line production by CRISPR

CRISPR guide oligo pairs were designed flanking the SBE2 enhancer region described by
Zhao, *et al* 2012. The pairs were annealed and cloned into the pX330 vector.
Guide sequences used were: Upstream of SBE2, 5' AAACACATTAAAGCCCTCCAGCG 3' and 5'
CACCGACGCTGGAGGGCTTTAATGT 3'; Downstream of SBE2, 5'
AAACACGAGCAAGCCAACCGGAGG 3' and 5' CACCGCTCCGGTTGGCTTGCTCGT 3'.
pX330-U6-Chimeric_BB-CBh-hSpCas9 was a gift from Feng Zhang (Addgene plasmid # 42230;
http://n2t.net/addgene:42230 ; RRID:Addgene_42230) (Cong et al., 2013).
Transgenic mice were made by pronuclear injection of plasmid DNA into C57Bl/6 x CBA F2
embryos. G0 pups were genotyped by PCR and sequencing. Pups carrying the correct deletions were
then used to establish the line by mating to C57BL/6J mice.
147

OPT analysis

OPT imaging was performed (Sharpe, 2003). Briefly, PFA-fixed embryos were embedded in
1% low-melting-point agarose and dehydrated by immersion in methanol for 24 h. The sampes were
cleared for 24 h in BABB (one part benzyl alcohol/two parts benzyl benzoate). before being scanned
using a Bioptonics 3001 scanner (www.biooptonics.com) Images taken every 0.9° (of a 360° rotation)
and were reconstructed using Bioptonics proprietary software with the outputs then being viewed with
Dataviewer (Bioptonics) and Bioptonics Viewer.

Morphometric analysis

MicroCTs of fifteen E17.5 embryos (7 controls, 8 mutants) collected from three litters were
scanned post mortem using a Bruker Skyscanner 1076. Individual skull scans were downsampled 2:1
using Bruker NRecon. Reconstructed images were converted to DICOM format using Bruker
DicomCT. Twenty-eight landmarks were placed on each isosurface using Stratovan CheckPoint
(v19.03.04.1102 Win64). In two mutants, dysmorphology of the vomer prevented placement of
corresponding landmarks and as such, the landmarks were indicated as "missing". Raw coordinates
data saved in Morphologika format were processed in R (v3.5.3) using Geomorph (v3.1.2), Rvcg
(v0.18), Morpho (v2.7), and Hotelling (v1.0-5) packages. Missing landmarks were estimated using a
multivariate regression method implemented in Geomorph. Following, a Generalised Procrustes
analysis (GPA) translated, rotated and rescaled raw landmark coordinate data to minimise the sum of
squared distances between specimens. The amount of rescaling required in the GPA is captured by
the centroid size, a variable which is used subsequently as a proxy for specimen size. A single outlier
(control, Litter 3) was removed as its Procrustes distance was >1 standard deviation from the mean.
Following removal of the outlier, the GPA was rerun to produce Procrustes shape variables. To test
for shape variation due to allometry, shape variables were regressed against log(centroid size) as well
as additional terms corresponding to genotype and litter. The best fit included log(centroid size)
interactions with both litter and genotype (ie coordinates ~ log(centroid size)*litter + log(centroid
size)*genotype).

In order to visualise shape differences, a randomly chosen ply surface file was warped using
thin plate splines using the mean coordinate values of all fourteen specimens. The resulting mesh
represented the dataset reference which was subsequently warped to using principal component
minimum and maximum shape variables.

1.1.1 Tissue processing

Embryos were dissected at the stages indicated and fixed overnight at 4°C in 4%
paraformaldehyde (PFA) in phosphate-buffered saline (PBS). Embryos were washed briefly in PBS,
and then passed through a sucrose gradient: 30 min in 5% sucrose in PBS, 30 min in 10% sucrose in
PBS before being placed in 20% sucrose in PBS overnight at 4°C. Embryos were embedded in OCT
compound and stored at -80°C. Sections were cut on a cryostat at varying thicknesses appropriate to
the embryonic stage: For E11.5 at 8 µm, for E13.5 at 10 µm and for E17.5 at 14 µm.

1.1.2 H&E staining

Cryosections were allowed to reach room temperature before the slides were rehydrated by
passage through 100% EtOH 3x 5 min then serial passage through 90%, 70%, 50% and lastly 30%
EtOH for a couple of min each. Slides were washed in H_2O and stained in haemotoxylin for 4 min.
They were rinsed in H_2O and subsequently differentiated in acid/alcohol for a few seconds. Slides were
again rinsed in H_2O then placed in saturated lithium carbonate solution for a few seconds. They were
rinsed in H_2O, stained in eosin for 2 min and again rinsed in H_2O. Slides were dehydrated through 4
changes of 100% ethanol and passaged 3x through xylene for 5 min each, subsequently mounted in
DPX (Cell Path, SEA-1304-00A), coverslipped and allowed to dry overnight.

lacZ expression analysis, Dawson staining and wholemount *in situ* hybridisation

Embryos were analysed for *lacZ* expression at E10.5 through staining for β-gal activity as
previously described (Mackenzie et al., 1997). Skeletal preparations from E17.5 foetuses were stained
simultaneously with Alizarin Red and Alcian Blue (Nagy et al., 2009a, 2009b). Whole-mount *in situ*
hybridisation on half heads was performed as previously described (Hecksher-Sørensen et al., 1998).
Whole-mount *in situ* hybridisation on E95 day embryos was performed as previously described (Pai et
al., 2015).

In situ hybridisation on cryosections

For in situ on cryosections, the protocol was adaped from (Lu et al., 2013) as described.
Briefly, *in situ* probes were placed at 37°C for 5 mins, added to hybridisation buffer (1x salt (2 M NaCl,
100 mM Tris, 65 mM NaH_2PO_4, 50 mM Na_2HPO_4 and 50 mM EDTA in ddH_2O, pH to 7.5), 50%
deionised formamide, 10% dextran sulfate, 1 mg/mL yeast RNA, 1x Denhardt's (2% BSA, 2% Ficcoll,
and 2% PVP made up to 100mL with ddH_2O)) pre-warmed to 65°C and subsequently denatured at
95°C for 2 mins. Probes were added to slides, that had been previously to reach room temperature, and
placed in a 65°C waterbath overnight in a chamber humidified with 5x SSC/50% formamide. Slides
were subject to 3x 30 min washes at 65°C in washing solution (50% formamide, 25 mL 1x SSC and
0.1% Tween 20 made up to 500 mL Milli-Q H_2O). Subsequently, slides were washed 2x 30 min in 1x
MABT (100 mM maleic acid, 167 mM NaCl in ddH_2O, pH to 7.5, then 0.1% Tween 20). Slides were
blocked in 1x MABT + 2% blocking reagent + 20% heat inactivated sheep serum. A 1/2500 dilution

of anti-DIG (Roche, 1109327490) in 1x MABT + 20% heat inactivated sheep serum was then added
and slides were placed in a humified chamber and incubated overnight at room temperature. Slides
were washed 5x for 20 min in 1x MABT. They were then washed with NTMT (100 mM NaCl, 50 mM
MgCl, 100 mM Tris pH 9.5 and 0.1% Tween 20 made up in ddH$_2$O) for 10 min followed by NTMTL
(NTMT + 5 mM Levamisole) for 10 min. Staining solution, NTMTL containing 3.75 µL/mL BCIP
(Roche, 11383221001) and 5 µL/mL NBT (Roche, 11383213001), was then added and slides were
placed at 37^{o}C to allow colour to develop. Slides were immersed in H$_2$O, air dried and mounted in
Prolong gold (Thermo Fisher, P39634).

1.1.3 Immunofluorescence
228   Cryosections were removed from the -80°C freezer and allowed to reach room temperature
before being subjected to antigen retrieval using a citric acid solution (10 mM citric acid dissolved in
ddH$_2$O, then pH to 6-6.5). Subsequently, slides were washed twice for 5 min in PBS then blocked using
10% goat serum in PBT (0.1% Triton-X100 in PBS) for 1 h at room temperature. For PITX2 staining,
donkey serum was used, Slides were incubated in primary antibodies diluted in blocking solution
overnight at 4°C as follows: rabbit anti-CASP3 (1:500, Abcam, ab49822); mouse anti-COL2A1 (1:400,
Santa Cruz, sc-52659); rabbit anti-ISL1 (1:200, Abcam, ab20670); rabbit anti-LHX3 (1:250, Abcam,
ab14555); rabbit anti-NKX2-1 (1:250, Abcam, ab76013); rabbit anti-NKX2-2 (1:250, Abcam,
ab191077); rabbit anti-NKX6-1 (1:250, Abcam, ab221549); rabbit anti-OLIG2 (1:200, Abcam,
ab109186); rabbit anti-oxytocin (1:500, Abcam, ab212193); rabbit anti-PAX6 (1:350, Abcam,
ab195045); sheep anti-PITX2 (1:200, R&D, AF7388); rabbit anti-SF1 (1:200, Abcam, ab65815);
rabbit anti-TBX3 (1:200, Abcam, ab99302); rabbit anti-TH1 (1:200, Abcam, ab137869); rabbit anti-
TCF4 (1:100, Cell Signalling; C48H11); rabbit anti-vasopressin (1:500, Abcam, ab213707). Slides
were then washed 3 times for 10 min each in PBT and incubated for 1 h at room temperature with
fluorescent secondary antibodies in PBT in the dark, secondaries as follows: goat anti-rabbit 488
(1:500, Thermo Fisher, A11034); goat anti-mouse 488 (1:500, Thermo Fisher, A11029); donkey anti-
sheep 594 (1:500, Thermo Fisher, A11016). Slides were then washed 3 times for 10 min each in PBT
and subsequently stained with 0.1 µg/ml DAPI in PBS for 10 min, then washed in PBS twice for 5
min each. Lastly, slides were mounted using Prolong Gold (Thermo Fisher, P39634) and allowed to
dry prior to imaging.

RESULTS

Loss of SBE2 enhancer activity leads to reduction in rostral diencephalic *Shh* expression

252   To examine the role of the SBE2 enhancer during development, this element was deleted using
CRISPR/Cas9 (Cong et al., 2013) to remove 1.2 kb (Fig. S1A, B) containing the conserved element.
The SBE2 deleted allele, henceforth referred to as *Shh^{ΔSBE2}*, was crossed to generate homozygous
embryos (*Shh$^{ΔSBE2/ΔSBE2}$*) and also to *Shh* null (*Shh^{null}*) mice to generate compound heterozygous
embryos (*Shh$^{ΔSBE2/null}$*). Loss of the SBE2 enhancer alone had no effect upon survival as the
Shh$^{ΔSBE2/ΔSBE2}$ mice were viable and fertile and at weaning no reduction in the number of mutant mice
was found. Conversely, no live *Shh$^{null/ΔSBE2}$* offspring were recorded by postnatal (P) 14 days (Fig. 1B),
these compound heterozygous embryos died perinatally failing to survive beyond P4. Normal ratios
were observed at embryonic (E) 17.5 (Fig. 1A).

The rostral diencephalon, extending from the ZLI to the MGE, is the target for SBE2 expression
as predicted by SBE2/reporter gene analysis in transgenic embryos (Jeong *et al.*, 2006). *Shh* mRNA
expression analysis at E10.5 in *Shh*$^{\Delta SBE2/\Delta SBE2}$ and *Shh*$^{null/\Delta SBE2}$ embryos revealed undetectable levels of
Shh expression in the ventral portion of the rostral diencephalon in both crosses (Fig. 1C-E), this region
of the diencephalon will give rise to the ventral portion of the tuberal and mammillary hypothalamus
(Xie and Dorsky, 2017), and for simplicity will henceforth be referred to as the ventral hypothalamus
(VH). The region lacking *Shh* expression in the SBE2 mutant embryos, however, was appreciably more
restricted than the region predicted by transgenic analysis, this is likely due to overlapping expression
caudally directed by other brain enhancers such as SBE3 and SBE4.

The differences in postnatal viability between *Shh*$^{\Delta SBE2/\Delta SBE2}$ and *Shh*$^{null/\Delta SBE2}$ suggested that there may
be residual, yet undetectable, *Shh* expression in the viable line. To investigate possible low levels of
Shh expression, patched-1 (*Ptch1*) expression, which is upregulated by hedgehog signalling and is a
sensitive readout of *Shh* expression, was examined at E10.5 (Casillas and Roelink, 2018). Similar to
the expression pattern observed for *Shh*, *Ptch1* was also downregulated in the VH upon loss of SBE2
activity (Fig. 1F-H). However, in this instance, levels differed between the compound heterozygous
and the homozygous embryos. While *Shh*$^{null/\Delta SBE2}$ embryos showed no detectable levels of *Ptch1*
expression in the VH (Fig. 1H, green arrow), the *Shh*$^{\Delta SBE2/\Delta SBE2}$ embryos presented residual levels (Fig.
1G, red arrow). It should be noted that while at E9.5 *Shh* levels are low in the ventral diencephalon in
the absence of SBE2 for both mutant conditions (Fig. 1I-K), that *Ptch1* expression pattern in
neighbouring tissue is comparable to wildtype (Fig. 1L-N). We suggest that *Ptch1* dependent
development is unaffected until after E9.5 and the defects occur after this stage. To determine when
VH *Shh* expression is downregulated, a mouse line containing the *LacZ* reporter gene inserted into the
Shh regulatory domain, SBLac526, was used (Anderson and Hill, 2014) as proxy for *Shh* expression.
VH *LacZ* expression was low but detectable up to E13.5 (Fig. S2A-C, red arrow), with expression
subsequently absent at E14.5 (Fig. S2D, black arrow). Hence, the consequential role of SBE2
encompasses a developmental window that begins sometime after E9.5 extending to E13.5 when all
regulatory activity in the VH is decreasing.

We suspected that presumed residual, low levels of *Shh* expression in the absence of SBE2
were capable of rescuing the postnatal lethality. Accordingly, an additional enhancer was identified in
Shh expressing neural cells (Harfe et al., 2004) from disected brains. ATAC-seq (Buenrostro et al.,
2015) revealed a vertebrate conserved sequence element (position chr5:29230287-29230750 [mm9]),
(Fig. S3A) and when used in reporter constructs in transgenic embryos at E10.5 (Fig. S3B-D) showed
broad expression activity extending anteriorly from the midbrain into the diencephalon (Fig. S3B-D).
Recently, an enhancer whose primary function is to drive expression initially in the prechordal plate,
and later throughout the ventral forebrain, called SBE7 (Sagai et al., 2019), was reported at the same
chromosomal position. Thus, SBE7 has a primary function in the prechordal plate and secondarily,
drives low but sufficient levels throughout the ventral forebrain.

Loss of SBE2 activity effects midline craniofacial deformities

SBE2 deletion embryos were analysed for craniofacial malformations. Unsurprisingly, given
the postnatal viability of *Shh*$^{\Delta SBE2/\Delta SBE2}$ embryos, no gross craniofacial malformations nor deformities
in the craniofacial skeletal elements (Fig. S4A-B) were observed. In contrast, at E17.5, *Shh*$^{null/\Delta SBE2}$

embryos exhibited external head deformities, in which the mutant embryos presented with a more
rounded head. To investigate these changes in head shape, control and $Shh^{null/\Delta SBE2}$ mutant embryos
were subject to geometric morphometric analysis to deconvolute the skull and test for differences
between control and mutant embryos. Centroid size, a proxy of skull scale, revealed that the compound
heterozygous mutant embryos tended to have an overall reduction in skull size, though this was not
significant (pairwise two-sample Wilcoxon, p=0.49) (Fig. 2A). Accordingly, allometric shape variation
due to size, genotype, and litter provided the best regression fit (R^2=0.83, ANOVA p<0.001). Principal
components of the non-allometric shape was also explored. We observed clear, genotype-specific
separation of control and mutant mice, with the latter occupying a relatively broader expanse of
morphospace (Fig. 2B). In mutants, Principal Component 1 (46.5%) (Fig. 2B) describes a reduction in
the anteroposterior axis while dorsoventral and mediolateral axes expanded (Fig. 2C-H). Procrustes
variances between control and mutant mice were not significant, however this was presumably due to
limitations in statistical power. Indeed, shape differences described by PC1 were significant (Welch
Two Sample t-test, p=1.5×10^{-4}). OPT analysis revealed that, in addition to presenting a perturbed
overall head structure, $Shh^{null/\Delta SBE2}$ embryos displayed affected nasal cavities (Fig. 2J,L), which were
reduced in length; however, the extent to which the nasal cavities were affected varied between mutant
embryos, with the example presented representing a moderate phenotype. Moreover, a region of brain
tissue, specifically the midline hypothalamic tissue, was absent in the mutant embryos (Fig. 2I,K).

$Shh^{null/\Delta SBE2}$ embryos displayed midline craniofacial bone deformities with a high degree of
variability, which, based on phenotypic severity, were classified into three categories: mild, moderate
and severe (Fig. 3A-D). All mutant embryos showed a reduction in the basisphenoid bone, ranging
from the presence of only a small remnant (Fig. 3B, pale green arrow) to complete loss (Fig. 3C, D)
and in the vomer bone, ranging from an excessive fusion (Fig. 3B, yellow arrow) to complete absence
(Fig. 3C, D). In addition, the moderate and severe embryos displayed defects in the pterygoid and
maxillary bones (Fig. 3C, D, white arrows). Notably, all embryos showed some midline craniofacial
element deformities, which indicates that Shh signalling directed by SBE2 is involved in the midline
development of the craniofacial bones, and parallels the role Shh plays in early midline brain
development (Choi et al., 2012).

Mislocalized pituitary and loss of SHH signalling disrupts craniofacial development

Closer examination of the heads of mutant compound heterozygous embryos revealed that the
basisphenoid bone, the most posterior defective cranial bone, was intercepted by the adenohypophyseal
pituitary lobe (labelled AP in Fig. 3E, F). This lobe of the pituitary was misplaced and formed in a
more anterior position, growing within the region where the basisphenoid will form (the basis for the
mislocalization is discussed below). Developmental timing of this event suggests that the
adenohypophysis disrupts this region before chondrogenesis initiates, which indicates that the
adenohypophyseal tissue likely presents a physical barrier to basisphenoid development. The
phenotypic variability may correlate with the extent to which the pituitary lobe intercepted the
basisphenoid bone in the mutant embryos.

Further investigations into the effect of SBE2 directed expression on craniofacial development
focused on mesenchymal $Ptch1$ expression in the vicinity of the forebrain. In control embryos at E12.5,
$Ptch1$ expression was detected within a narrow layer of mesenchyme located directly adjacent to the
floor of the diencephalon where Shh was expressed (Fig. 4A-B). COL2A1, a type 2 collagen, is
expressed during early cartilage formation (Hissnauer et al., 2010) and co-localised with $Ptch1$ in

control embryos (Fig. 4E-F), implicating that the underlying facial mesenchyme is under the influence
of SHH and is committed towards cartilage formation. At the same stage in the *Shh^{null/ΔSBE2}* embryos,
Ptch1 expressing cells were absent in the underlying mesenchyme (Fig. 4C-D) and only a thin lining
of cells expressing COL2A1 was seen (Fig. 4G-H). The loss of *Ptch1* and COL2A1 staining was found
to be independent of the physical disruptive effects caused by mislocalisation of the neurohypophysis,
as the absence of both markers was evident in regions of the cranial base which lie anterior to disrupted
neurohypophyseal tissue detected in the mutant embryos (Fig. 4C,G). Thus, SBE2 mediated *Shh*
expression is instrumental in ensuring adequate cell response and bone formation in the sub–
diencephalic mesenchyme along the midline of the palatal floor, and reduction of this cell population
is critical for the midline craniofacial phenotype.

Shh^{null/ΔSBE2} embryos display mislocalised and deformed pituitary lobes due to disruption of the AP hypothalamic boundary

Pituitary lobe development was analysed at stages E11.5, E13.5 and E15.5. In *Shh^{null/ΔSBE2}*
mutant embryos at E11.5, Rathke's pouch (RP), which will give rise to the adenohypophyseal lobe,
and the infundibulum, which will give rise to the neurohypophyseal pituitary lobe, were both
ectopically shifted ventrally (Fig. S5 A, B). These findings reflect the previous observations (Zhao et
al., 2012) in conditional *Shh* deletion mice, when recombination was driven by the SBE2 enhancer
(designated SBE2-Cre Δ*Shh*) an anterior shift in the pituitary lobes was observed. At E13.5, RP was
severely malformed in the *Shh^{null/ΔSBE2}* mutant embryos, while the infundibulum did not separate from
the neuroectodermal tissue to the same degree as control counterparts (Fig. 5A, C) and remained
ectopically located at E15.5 (Fig. S5C, D).

Interestingly, at E17.5 the neurophyphysis appeared absent in the *Shh^{null/ΔSBE2}* mutant
embryos (Fig. 3E, F), yet staining for markers of neurohypophyseal tissue revealed that this was due
to the lack of separation of the neurohypophysis from the neuroectodermal tissue of origin (Fig. 5L,
N). Moreover, caspase-3 (CASP3) staining revealed that this remnant neurohypophyseal tissue
appeared to be undergoing apoptosis (Fig. 5M, O) indicating that inadequate neurohypophyseal
partitioning likely led to tissue degeneration. Despite the fact that both hypophyseal lobes were
misspecified in the *Shh^{null/ΔSBE2}* embryos early hypophyseal patterning was unperturbed (Fig. S6C, F,
I, L). These findings agree with observations by Zhao et al., (2012) in SBE2-Cre Δ*Shh* mice, where
pituitary markers, such as LHX3, were unperturbed despite hypophyseal misplacement. However,
whilst SBE2-Cre Δ*Shh* mice displayed multiple invagination sites for RP, only one site was observed
in *Shh^{null/ΔSBE2}* embryos. To exclude the possibilty that the observed pituitary malformation and
mislocalisation, specifically that of the adenohypophysis, were due to a developmental delay,
expression of adenohypophyseal markers that display stage-specific restricted expression within the
lobe were analysed. ISL1 is expressed throughout the early developing adenohypophysis but later
becomes restricted to the ventral most portion of the lobe (Castinetti et al., 2015; Ericson et al., 1998).
This same restriction of expression was seen for TBX3, which was expressed broadly throught the
developing adenohypophysis, but by E12.5 expression was restricted to the ventral aspect of the lobe
(Pontecorvi et al., 2008). For both ISL1 (Fig, S7A-B) and TBX3 (Fig. S7C-D) expression in the
adenohypophysis at E12.5 in *Shh^{null/ΔSBE2}* embryos was found to reflect the pattern seen in control
embryos, with expression only detected in the ventral most portion, indicating that the
adenohypophyseal lobe was not developmentally delayed.

While patterning and early neuronal marker expression within the pituitary lobes appeared
normal, at later developmental stages, E17.5, neuronal populations within varied nuclei of the VH,
which establish connections with the pituitary lobes, were disrupted. Both the tyrosine hydroxylase
(TH1) (Fig. S8A-B) and the vasopressin neuronal population (Fig. S8E-F) of the paraventricular
nucleus (PVN) were lost in $Shh^{null/\Delta SBE2}$ embryos. Additionally, the oxytocin neuronal population of the
PVN was absent while the population found within the arcuate nucleus (ARC) was either misspecified
or mislocalised (Fig. S8C-D). To confirm that the loss of PVN and ARC neuronal populations did not
result from a developmental delay in hypothalamic neuronal specification, expression of oxytocin and
TH1 was analysed at E15.5. At E15.5 oxytocin expression was present in the PVN of the control
embryos (Fig. S9A, white arrow) but absent from $Shh^{null/\Delta SBE2}$ embryos (Fig. S9B). Similarly,
expression of TH1 was detected in the ARC of control embryos (Fig. S9C, yellow arrow), yet no
expression was detected in $Shh^{null/\Delta SBE2}$ embryos (Fig. S9D). This indicates that the lack of specific gene
expression in hypothalamic neuronal populations at E15.5 persists for at least two additional days up
to E17.5, arguing for a loss of the neuronal population and not a developmental delay in neuronal
specification. These results indicate that, in addition to the disruption of hypophyseal structure and
location in development, there is also disruption of hypothalamic neuronal populations, demonstrating
that SBE2 activity is vital for the role of *Shh* in hypophyseal and hypothalamic neuronal development.
Interestingly, the hypothalamic populations found to be affected are known to establish connections
with and regulate the hypophysis, indicating that *Shh* plays a role in coordinating hypophyseal
development and subsequent neuronal regulation. The disruption of these late neuronal populations in
$Shh^{null/\Delta SBE2}$ embryos revealed that early loss of SBE2 directed *Shh* activity in the VH has late effects
upon development in the hypothalamus, at timepoints at which *Shh* expression is no longer present in
the tissue in question (Fig. S2).

In $Shh^{\Delta SBE2/\Delta SBE2}$ mutant embryos, neither the adenohypophysis nor the neurohypophysis were
ectopically located and both lobes had separated from the tissue of origin E13.5 (Fig. S4). Additionally,
no defects were observed in early marker expression (Fig. S6B, E, H, K); however, mild deformities
were detected predominantly in the structure of the lateral portion of adenohypophyseal lobe in
$Shh^{\Delta SBE2/\Delta SBE2}$ mutant embryos (Fig. S4C-K).

Distinct anterior-posterior expression domains are established along the midline of the
hypothalamus by E9.5. The posterior hypothalamic domain expresses factors including *Bmp4* and
Fgf10 whilst the anterior domain expresses *Six6*, *Tcf4* and *Shh*. Mutual inhibition between these two
gene expression domains exists, whereby in the absence of one domain the opposing domain will
expand to occupy the hypothalamic space (Ericson *et al*., 1998; Takuma *et al*., 1998). The
infundibulum expresses the posterior markers *Bmp4* and *Fgf10* and is specified dorsally to the anterior
hypothalamic tissue, at the AP transition zone (Ferrand, 1972; Kawamura and Kikuyama, 1995;
Takuma *et al*., 1998; Treier *et al*., 1998). Inductive cues from the ventral hypothalamic tissue play a
role in orchestrating the formation of RP (Schwind, 1928; Takuma *et al*., 1998). As RP and the
infundibulum were both shifted ventrally in the $Shh^{null/\Delta SBE2}$ mutant embryos (Fig. 5C), verification of
any other associated defects in the establishment of the AP hypothalamic boundary were sought. The
posterior boundary of expression of TCF4 was shifted ventrally in the mutant embryos (Fig. 5I)
whereas the anterior boundary of expression of *Fgf10* was shifted ventrally, expanding expression (Fig.
5F). The shift observed for both the anterior and posterior markers was found to be significant in the
$Shh^{null/\Delta SBE2}$ mutant embryos (Fig. 5J, K). These results indicated that there is a shift in the AP domain
boundary with a reduction in anterior markers and gain of posterior markers. In the $Shh^{\Delta SBE2/\Delta SBE2}$
mutant embryos no shift in expression of TCF4 (Fig. 5H) or *Fgf10* (Fig. 5E) was observed, correlating
with correct positioning of the pituitary lobes (Fig. 5C). These results suggest that shift of the pituitary

lobes occurs as a result of the aberrant specification of the AP boundary, which does not appear to be
sensitive to reduced levels of SHH shown in $Shh^{\Delta SBE2/\Delta SBE2}$ embryos (Fig. S4C-F).

Loss of SBE2 enhancer activity leads to defects in hypothalamic patterning in a concentration
dependent manner

Much of our knowledge about DV hypothalamic patterning is informed by what is known
regarding DV neural tube patterning. In the developing neural tube, neural progenitors acquire distinct
transcriptional identities based on length of exposure to patterning signals and DV positioning
(Balaskas et al., 2012; Dessaud et al., 2007, 2010). Indeed it is the opposing action of the dorsal neural
tube signals *Wnts* and BMPs and the ventral neural tube signal *Shh* that drive patterning along this axis
(Burbridge et al., 2016). Subsequently, expression of distinct combinations of transcription factors
drives the differentiation of the neural progenitors towards specific neural fates (Dessaud et al., 2008).
In the developing hypothalamus *Shh* is initially expressed from the prechordal mesoderm; it is
subsequently expressed along the floor plate and plays a role in inducing ventral hypothalamic fate
(Burbridge et al., 2016; Corman et al., 2018). In contrast, BMPs are expressed from the roof of the
telencephalon, regulating alar hypothalamic development (Burbridge et al., 2016). Given that *Shh*
signalling from the hypothalamic floor plate was reduced in the $Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$ mutant
embryos, DV hypothalamic patterning was examined. Genes whose expression are responsive to SHH
levels in the neural tube were of particular interest. No differences were detected between wild-type,
$Shh^{\Delta SBE2/+}$ and $Shh^{null/+}$ embryos (Fig. S10), indicating that a 50% reduction in *Shh* levels has no effect
upon DV patterning marker expression.

Expression of PAX6, known to the expressed within the alar hypothalamus, and the opposing
basal hypothalamic marker NKX2-*2* (Croizier et al., 2015; Guillemot, 2007) were examined. NKX2-2
expression commences early in hypothalamic development, with expression detected in the basal
hypothalamic region which borders the alar zone. Later in development, this expression pattern alters
within the VH, whereby expression has forked into two bilateral stripes; one that occupies the transition
zone of the alar-basal hypothalamus extending into the alar hypothalamus itself, and a second stripe
that resides within the basal hypothalamic territory of the VH (Burbridge et al., 2016). At E13.5, this
bilateral stripe expression pattern was observed in control embryos (Fig. 6A). However, in
$Shh^{\Delta SBE2/\Delta SBE2}$ mutants the dorsal VH stripe, within the basal hypothalamus, presented patchy
expression (Fig. 6B). The $Shh^{null/\Delta SBE2}$ mutant embryos were more severely affected, with both the alar
and basal VH stripes absent (Fig. 6C). This data indicate that low levels of *Shh* expression can induce
some NKX2-2 expression in $Shh^{\Delta SBE2/\Delta SBE2}$, however, these low levels are not sufficient to either
specify or maintain the entirety of the NKX2-2 population.

PAX6 is a neural tube and hypothalamic marker, whose expression pattern is known to be
restricted to the alar hypothalamus (Burbridge et al., 2016) and, as expected, E13.5 control embryos
displayed PAX6 expression solely within the alar portion of the VH (Fig. 6D). In $Shh^{\Delta SBE2/\Delta SBE2}$ mutant
embryos there was a dorsal expansion of PAX6 expression into the basal hypothalamic region in the
embryos (Fig. 6E). In $Shh^{null/\Delta SBE2}$ embryos, this dorsal expansion of PAX6 expression was more
severe, with large patches of PAX6 expression detected in the basal hypothalamus of the mutant
embryos (Fig. 6F). These results indicate that *Shh* signalling from the basal hypothalamus is required
to restrict PAX6 expression solely to the alar region, and that low levels of *Shh* activity, those found
in basal hypothalamus of $Shh^{\Delta SBE2/\Delta SBE2}$ mutants, are not sufficient to exclude PAX6 activity from this
region.

481    These results for NKX2-2 and PAX6 within the VH, upon loss of SHH signalling activity,
482 mirror what was found by Corman and colleagues (2018). However, these data demonstrate a novel
483 dose dependent effect of SHH signalling whereby these transcription factors are not only able to
484 respond to the presence, or indeed absence, of *Shh* signalling, but are also sensitive to the amount of
485 *Shh*, with the levels of *Shh* activity being key for theestablishment of hypothalamic signaling domains.

486    Expression of the transcription factor OLIG2 is induced by SHH (Dessaud et al., 2008), and in
487 control embryos was detected in the alar hypothalamus in a narrow region along the edge of the third
488 ventricle (Fig. 6G). In *Shh$^{\Delta SBE2/\Delta SBE2}$* mutant embryos, however, the OLIG2 positive cells showed a
489 subtle dorsal shift away from the edge of the third ventricle and additional expressing cells were
490 detected in the basal hypothalamus (Fig. 6H). In *Shh$^{null/\Delta SBE2}$* mutant embryos, OLIG2 positive staining
491 cells had dramatically shifted into the basal portion of the hypothalamus, with large areas of positive
492 staining cells detected (Fig. 6I). These data indicated that *Shh* activity acts to ensure adequate OLIG2
493 expression within the alar hypothalamus, whereupon reduction of SHH allows for OLIG2 expression
494 within the basal hypothalamus. Unexpectedly, OLIG2 was found to be expanded into ectopic regions
495 of the VH in mutant embryos rather than gradually reduced with increasing loss of SHH activity.
496 However, Balaskas et al., 2012, demonstrated in the neural tube that, whilst NKX2-2 and OLIG2 are
497 both equally responsive to SHH activity, higher levels of signalling are required for NKX2-2
498 expression as it is repressed by both PAX6 and OLIG2, creating a gene regulatory network between
499 these three transcription factors. Thus, in this instance, the residual levels of *Shh* expression directed
500 by SBE7 in *Shh$^{null/\Delta SBE2}$* mutant embryos are likely sufficient to induce OLIG2 expression, yet not
501 NKX2-2 due to the repressive activity of OLIG2 and PAX6.

502    NKX6-1, another transcription factor induced by SHH in the neural tube (Dessaud et al., 2008),
503 was expressed in the alar hypothalamus in a narrow region along the edge of the third ventricle in
504 sagittal and parasagittal regions (Fig. 6J); but in contrast to OLIG2, these cells were only detected
505 along the basal lateral edge of the third ventricle. In *Shh$^{\Delta SBE2/\Delta SBE2}$* mutant embryos, NKX6-1 expression
506 along the basal lateral edge of the third ventricle was undetectable (Fig. 6K), however, isolated groups
507 of expressing cells were detected in the basal hypothalamus (Fig. 6K). *Shh$^{null/\Delta SBE2}$* mutants showed
508 complete loss of the region of staining along the basal lateral edge with no additional regions of
509 expression observed (Fig. 6L). NKX6-1 appears to require low levels of *Shh* expression to be induced,
510 levels which are normally found along the edge of the third ventricle. In *Shh$^{\Delta SBE2/\Delta SBE2}$* mutants the
511 levels of *Shh* expression are reduced within the basal hypothalamus to a level that allows for ectopic
512 expression of NKX6-1 within this region. In *Shh$^{null/\Delta SBE2}$* mutants the complete loss of the expressing
513 cells from the edge of the third ventricle indicate that the levels of *Shh* signalling activity within the
514 VH, directed by SBE7, are insufficient to induce expression of NKX6-1. These data demonstrate a
515 novel, previously undescribed role for SHH activity in the regulation of NKX6-1 expression within the
516 hypothalamus, which mirrors the role described in the neural tube (Sander et al., 2000). Thus, providing
517 further evidence that the signalling network which operates in the neural tube has been co-opted in
518 hypothalamic development.

519    For all hypothalamic markers analysed, the extent of changes in expression correlate to the
520 scale of alteration in levels of *Shh* expression, suggesting that *Shh* signalling is functioning in a
521 concentration dependent manner to induce ventral patterning in the hypothalamus. The data described
522 not only present novel examples of patterning genes regulated by SHH activity, as is the case for
523 NKX6-1, but also expand on data presented by Corman *et al.* (2018). We further demonstrate that, not
524 only do these genes require SHH activity to be adequately expressed, but they are capable of responding
525 to varying levels of SHH. This concentration dependent effect mirrors that seen for *Shh* in the neural

526   tube, and involves many of the same key players, indicating that *Shh* is acting as a morphogen within
527   the VH to regulate tissue establishment within this critical brain region.

529   SBE2 enhancer activity controls downstream effects upon hypothalamic neuronal populations

We next sought to assess the effects of disrupted DV patterning on neuronal populations. We
first verified that there were no detectable differences in marker expression between wild-type and
heterozygotes, *Shh*$^{\Delta SBE2/+}$ and *Shh*$^{null/+}$ embryos (Fig. S11), demonstrating that a 50% reduction in *Shh*
levels has no gross effect upon DV neuronal population specification.

To examine the effect of disrupted DV patterning upon the establishment of the different ventral
hypothalamic neuronal populations, we first analysed expression of the broad ventral hypothalamic
neuronal marker NKX2-1 (Ohyama et al., 2005). In control embryos at E13.5, NKX2-1 positive cells
were detected across the entirety of the basal hypothalamic domain (Fig. 7A). In *Shh*$^{\Delta SBE2/\Delta SBE2}$ mutant
embryos, the region of NKX2-1 expressing cells was reduced and embedded within smaller domains
intermingled with patches devoid of staining (Fig. 7B). In the *Shh*$^{null/\Delta SBE2}$ mutant the region of
expression was further reduced with larger and more numerous patches of negative cells observed (Fig.
7C). These results indicate that whilst SHH activity is required for proper induction of the NKX2-1
neuronal population another factor must also sustain the remnant expression seen in mutant embryos.

ISL1 is a melanocortinogenic neuronal marker, for which positive cells are detected throughout
the hypothalamus (Kim et al., 2015). In control embryos positive staining was detected throughout the
basal hypothalamic region, as expected (Fig. 7D). No differences in staining were detected between
control and *Shh*$^{\Delta SBE2/\Delta SBE2}$ mutant embryos (Fig. 7E), however, in *Shh*$^{null/\Delta SBE2}$ mutants there was
substantial loss of positive staining cells in the basal hypothalamus (Fig. 7F).

We additionally verified whether neurons that are restricted to specific nuclei of the basal
hypothalamus were affected upon removal of SBE2 activity. Steroidogenic factor 1 (SF1) positive
neurons are localised to the ventromedial hypothalamus (VMH) and are involved in the regulation of
glucose metabolism and energy balance (Choi et al., 2013). In control embryos, as expected, SF1
positive neurons were detected in the developing VMH of the basal hypothalamus (Fig. 7G). In
Shh$^{\Delta SBE2/\Delta SBE2}$ mutant embryos the region of the basal hypothalamus expressing SF1 was reduced (Fig.
7H), while in *Shh*$^{null/\Delta SBE2}$ mutant embryos there was a complete loss of SF1 positive neurons (Fig. 7I).
Somatostatin (SOM) expressing neurons are a subset of GABAergic inhibitory neurons that are
detected in varied nuclei of the hypothalamus, including the VMH and the arcuate nucleus, which serve
to regulate cell proliferation ultimately affecting growth (Urban-Ciecko and Barth, 2016). In control
embryos, SOM neurons were detected in the basal hypothalamic region corresponding to the arcuate
nucleus at E13.5 (Fig. 7J). No differences were detected in *Shh*$^{\Delta SBE2/\Delta SBE2}$ mutant embryos where
comparable staining was detected (Fig. 7K); however, in *Shh*$^{null/\Delta SBE2}$ mutant embryos no positive
staining for SOM was detected in the VH (Fig. 7L). These results indicated that low levels of *Shh* in
the VH were sufficient to retain specification of SOM neurons, but higher levels of *Shh* activity were
required to sustain and/or specify the SF1 neuronal population in the basal hypothalamus. In the
Shh$^{null/\Delta SBE2}$ mutant embryos both of these populations were absent which indicated that early *Shh*
activity was essential for the adequate establishment of both the SF1 and SOM neuronal populations
of the basal hypothalamus.

Corman *et al.* (2018), have previously demonstrated that loss of *Shh* signalling directed by
SBE2 leads to a loss or reduction of neuronal markers. Our data expand on these findings by

demonstrating a previously unknown concentration dependent requirement for *Shh* in VH neuronal
development whereby some neuronal populations, such as ISL1 and SOM, are only affected when the
levels of *Shh* are vastly reduced, as in *Shh$^{null/\Delta SBE2}$* mutants. In contrast, the NKX2-1 and SF1
populations are affected by moderate reductions in the levels of *Shh* and further affected by greater
loss of *Shh* signalling.

Discussion

Secondary enhancer activity rescues SBE2 inactivation

Loss of SBE2 in *Shh$^{\Delta SBE2/\Delta SBE2}$* homozygous mice did not cause lethality or overt phenotypic
effects in the brain and craniofacial regions due to additional enhancer activity most likely from the
broadly acting element SBE7. A multistep series of events generate the normal *Shh* expression levels
in the ventral diencephalon of which SBE2 activity plays a later role. SBE7 is responsible for early *Shh*
expression throughout the prechordal plate a tissue that underlies the developing brain, from as early
as E7.5 (Sagai et al., 2019). SBE7, subsequently, drives *Shh* expression at low levels in the ventral
midline of the developing forebrain throughout later developmental stages. By E9.5 SBE2 activation,
which is dependent on the earlier SBE7 activity (Sagai et al., 2019), contributes to a substantially
increased level of expression specifically in the ventral diencephalon.

SBE2 activity does not appear to be obligatory since in its absence the animals are viable and
fertile; raising questions about the value of this enhancer to the embryo. In contrast, the deep
conservation of this element strongly argues that SBE2 provides fitness to individuals and is crucial
across the vertebrate classes. Consequently, we argue that this is not an example of enhancer
redundancy that commonly occurs in the mammalian genome (Osterwalder et al., 2018). Our analysis
shows that expression of DV patterning genes is sensitive to level of SHH which is perturbed in the
absence of SBE2. SBE2 and SBE7 are components of a systematic temporal response required to reach
sufficient SHH levels to attain the precise pattern of gene expression. Since further reduction of *Shh*
levels (as in the *Shh$^{null/\Delta SBE2}$* mutant) causes greater disruption to the expression of hypothalamic marker
genes and perinatal lethality, it is reasonable to argue that, in the absence of SBE2, these animals are
not fit in the wild. Disruption of neuronal populations seen in *Shh$^{\Delta SBE2/\Delta SBE2}$* homozygous embryos,
such as the reduction in the NKX2-1 and SF1 positive population, may lead to phenotypic effects on
physiology, such as reproduction and energy balance. Indeed, specific depletion of the SF1 neuronal
population of the VMH is known to be viable, however the metabolic response of skeletal muscle are
attenuated, revealing an effect upon metabolism (Fujikawa et al., 2016).

SBE2 mediated expression links brain and craniofacial development

Heterozygous inactivating mutations in the *SHH* gene in patients cause HPE (Chiang et al.,
1996; Ming and Muenke, 2002) while in mice complete loss of *Shh* leads to multiple phenotypic
consequences including the absence of brain and craniofacial structures (Chiang et al., 1996). While
these observations highlight the links between brain and craniofacial development, the question as to
whether *Shh* signalling from the neuroectoderm is directly required for the development of the
craniofacial elements has remained largely unanswered. Physical interactions between the brain and

face begin early at the initial stages of facial formation (Marcucio et al., 2011). There are two
fundamentally different mechanisms by which development of the face and brain may be interrelated;
firstly, the developing brain serves as a dynamic architectural foundation upon which the face responds
and develops and secondly, signalling directly from the brain to the cranium and face regulates
morphogenesis.

In fish, *Shh* signalling from both the oral ectoderm and neural tissue is required for adequate
progression of craniofacial development (Wada et al., 2005). In mammalian development, *Shh*
signalling from the oral ectoderm, which underlies the cranial base and also neighbours the facial
skeletal elements, has been demonstrated to be key in adequate craniofacial development. Depletion
of signalling from the oral ectoderm leads to defects in the maxillary and mandibular bones, the palatine
bones and the basisphenoid bone among others (Jeong et al., 2004). The present study addressed the
role of SBE2 forebrain expression activity on the development of adjacent tissue which would be
predicted to form the midline of the cranial floor. Loss of SBE2 in the presence of the *Shh* null allele
leads to both lethality and craniofacial deformities. Defects were observed in multiple skeletal elements
of the cranial base including the vomer bone and the basisphenoid bone, which resides below the
pituitary glands and the hypothalamic tissue, demonstrating that *Shh* signalling from the rostral
diencephalon is directly required for adequate craniofacial develop.

SBE2 affects both AP and DV patterning in the hypothalamus

In addition to the craniofacial defects observed in the SBE2 mutant mouse lines, defects in
hypothalamic development were also observed. This is perhaps unsurprising given that the region of
Shh expression directed by SBE2 is the rostral diencephalon, the tissue from which the hypothalamus
originates. However, the mode in which *Shh* regulates hypothalamic patterning and establishment
appears to differ depending on the axis considered.

The AP axis was less sensitive to *Shh* levels of expression, whereby low, residual levels of
signalling activity in the $Shh^{\Delta SBE2/\Delta SBE2}$ mutants were sufficient to sustain normal patterning. In contrast,
further reduction of *Shh* expression in the $Shh^{null/\Delta SBE2}$ mutant embryos caused a loss of the anterior
hypothalamic identity, which is promoted by *Shh*, in favour of the posterior hypothalamic identity. A
shift in the anterior marker TCF4 was observed in the $Shh^{null/\Delta SBE2}$ mutant embryos, which contrasts
previous observations (Zhao *et al*., 2012), where a shift in all other AP hypothalamic markers with the
exception of TCF4 was observed. This difference likely resides in the different mouse lines used,
whereby the use of SBE2-Cre to conditionally knock-down *Shh* activity may lead, firstly, to a temporal
delay in the removal of *Shh* activity and, secondly, to a broader removal of *Shh* activity than that solely
directed by SBE2. Notably, *Shh* is also required for adequate development of neuronal populations
which connect to the hypophyseal lobes and in turn, regulate hormonal release. These results
demonstrate a dual role for *Shh* in ensuring hypophyseal function, directly in hypophyseal development
and secondarily in regulating innervation.

In contrast to AP patterning, the DV hypothalamic axis was affected by *Shh* activity in a
concentration dependent manner. This is apparent in the $Shh^{\Delta SBE2/\Delta SBE2}$ mutant embryos where defects
in DV patterning, and also subsequent neuronal induction, were observed. These defects were far less
severe than those observed in the $Shh^{null/\Delta SBE2}$ mutant embryos, where the disruption to DV
development resulted in a complete loss of hypothalamic neuronal populations. The results presented
regarding DV hypothalamic patterning are comparable to those described by Corman *et al*. (2018)

where the same key neural tube patterning genes and subsequence neuronal regulators are involved.
However, we have demonstrated a novel concentration dependent patterning activity of *Shh* in the
hypothalamus, which has previously only been described in the neural tube. Moreover, the results
demonstrate loss of neuronal populations beyond those of the dorsomedial, ventromedial and arcuate
nuclei, additionally implicating neurons of the paraventricular nucleus, which play a key role in
hypophyseal regulation. Corman *et al*. (2018) also propose that NKX2-2 is expressed in a domain
between dorsal PAX6 and ventral NKX2-1; however, our data suggest that NKX2-1 is expressed
broadly across the entire ventral hypothalamic region, with NKX2-2 expressed in a subdomain of the
ventral hypothalamus, in regions overlapping with NKX2-1. It is possible that the ventral hypothalamic
markers induced by *Shh*, including NKX2-2 are responsible for inducing expression of NKX2-1,
whereby ventral hypothalamic markers that are not completely absent in the $Shh^{null/\Delta SBE2}$ mutant
embryos, sustain remaining NKX2-1 expression.

Similar to the patterning process in the neural tube, the response of different DV patterning
transcription factors to *Shh* was concentration dependent. For example, NKX6-1 was ventrally
displaced in the $Shh^{\Delta SBE2/\Delta SBE2}$ mutants, much like OLIG2. However, it was lost altogether in
$Shh^{null/\Delta SBE2}$ mutant embryos, unlike OLIG2 which was further displaced; indicating that NKX6-1 is
more sensitive to loss of *Shh* signalling than OLIG2. Similarly, in the neural tube OLIG2 responds to
a wider range of *Shh* signalling levels than NKX6-1. This same variable sensitivity to *Shh* signalling
was seen for the neuronal populations, whereby only some of the neuronal population that were
disrupted in the $Shh^{null/\Delta SBE2}$ mutant embryos were also affected in the $Shh^{\Delta SBE2/\Delta SBE2}$ mutants; for
example, NKX2-1 expression required higher levels of *Shh* expression than those needed for ISL1.
Moreover, *Shh* was found to only be essential to maintain the SF1 and somatostatin neuronal
populations, which were completely absent in $Shh^{null/\Delta SBE2}$ mutants. Thus, early SBE2 directed *Shh*
signalling directs subsequent DV patterning in a concentration dependent manner acting as a
morphogen within ventral hypothalamic development, in which transcription factor expression patterns
are controlled by *Shh* levels which in turn instruct ventral hypothalamic neuronal populations.

SBE2 coordinates craniofacial development and patterning of the hypothalamus

SBE2 regulates the expression of *Shh* in a region of the diencephalon in which SHH signalling
coordinates development of both the diencephalon and the neighbouring tissue responsible for the
cranial and facial bones, skeletal elements which are in turn responsible for protecting the neural
structures from external exposure. Homologues of the craniofacial bones disrupted in mouse are found
in species of all vertebrate classes from fish to mammals (Maddin et al., 2016). In addition, the
hypothalamus is an ancient brain centre critical for the production and release of key hormones (Xie
and Dorsky, 2017). We suggest that deep in vertebrate evolution SBE2 activity arose as a means to
govern VH development and as a consequence of hypothalamus proximity to craniofacial mesenchyme
enabled *Shh* signalling to control formation of the bone to ensure its own protection. We have
demonstrated that disruptions to coordinated development of the VH and cranial vault in mice leads to
phenotypic defects which reflect those seen for HPE patients. These results reveal not only that the
interactions between the brain and the face display deeply conserved evolutionary roots, but also show
the key importance of mouse models for addressing human disease phenotypes.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

Author Contributions

Conceptualization, REH.; Methodology, LAL, NK and ZCS.; Validation, ZCS and DMK.;
Investigation, ZCS, JS, KAG, PSD, LR, MD, JIH, LAL Writing – Original Draft, ZCS and REH;
Writing – Review & Editing, DMK., JS and LAL; Visualization, ZCS, KAG, JS and LAL.;
Supervision, DMK, JS and REH; Funding Acquisition, DMK and REH

Funding

ZCS, KAG, PSD, LR, EA, JIT, LAL and REH were supported by a MRC core award to the
MRC Human Genetics Unit. NK, and DK were funded by a CIHR operating grant to DMK (MOP-
275053). JS was supported by aBBSRC Institute Strategic Programme grant BBS/E/D/10002071.

Acknowledgements

We would like to thank the MRC Central Services for providing DNA sequence support. We
would also like to thank the staff at the Evans Building for expert technical assistance. Collaborative
work was supported by the MRC and the University of Calgary

References

Anderson, E., and Hill, R.E. (2014). Long range regulation of the sonic hedgehog gene. Curr. Opin. Genet. Dev. *27*, 54–59.

Anderson, E., Devenney, P.S., Hill, R.E., and Lettice, L.A. (2014). Mapping the Shh long-range regulatory domain. Development *141*, 3934–3943.

Balaskas, N., Ribeiro, A., Panovska, J., Dessaud, E., Sasai, N., Page, K.M., Briscoe, J., and Ribes, V. (2012). Gene regulatory logic for reading the Sonic Hedgehog signaling gradient in the vertebrate neural tube. Cell *148*, 273–284.

Buenrostro, J.D., Wu, B., Chang, H.Y., and Greenleaf, W.J. (2015). ATAC-seq: A Method for Assaying Chromatin Accessibility Genome-Wide. Curr. Protoc. Mol. Biol. *109*, 21.29.1-9.

Burbridge, S., Stewart, I., and Placzek, M. (2016). Development of the Neuroendocrine Hypothalamus. In Comprehensive Physiology, (Hoboken, NJ, USA: John Wiley & Sons, Inc.), pp. 623–643.

Casillas, C., and Roelink, H. (2018). Gain-of-function Shh mutants activate Smo cell-autonomously independent of Ptch1/2 function. Mech. Dev. *153*, 30–41.

Castinetti, F., Brinkmeier, M.L., Mortensen, A.H., Vella, K.R., Gergics, P., Brue, T., Hollenberg, A.N., Gan, L., and Camper, S.A. (2015). ISL1 is necessary for maximal thyrotrope response to hypothyroidism. Mol. Endocrinol. *29*, 1510–1521.

Chiang, C., Ying, L.T.T., Lee, E., Young, K.E., Corden, J.L., Westphal, H., and Beachy, P.A. (1996). Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. Nature *383*, 407–413.

Choi, K.-S., Lee, C., and Harfe, B.D. (2012). Sonic hedgehog in the notochord is sufficient for patterning of the intervertebral discs. Mech. Dev. *129*, 255–262.

Choi, Y.-H., Fujikawa, T., Lee, J., Reuter, A., and Kim, K.W. (2013). Revisiting the Ventral Medial Nucleus of the Hypothalamus: The Roles of SF-1 Neurons in Energy Homeostasis. Front. Neurosci. *7*.

Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W., Marraffini, L.A., et al. (2013). Multiplex genome engineering using CRISPR/Cas systems. Science *339*, 819–823.

Corman, T.S., Bergendahl, S.E., and Epstein, D.J. (2018). Distinct temporal requirements for Sonic hedgehog signaling in development of the tuberal hypothalamus. Development *145*, dev167379.

Croizier, S., Chometton, S., Fellmann, D., and Risold, P.-Y. (2015). Characterization of a mammalian prosencephalic functional plan. Front. Neuroanat. *8*, 161.

Dessaud, E., Yang, L.L., Hill, K., Cox, B., Ulloa, F., Ribeiro, A., Mynett, A., Novitch, B.G., and Briscoe, J. (2007). Interpretation of the sonic hedgehog morphogen gradient by a temporal adaptation

mechanism. Nature *450*, 717–720.

Dessaud, E., McMahon, A.P., and Briscoe, J. (2008). Pattern formation in the vertebrate neural tube:
A sonic hedgehog morphogen-regulated transcriptional network. Development *135*, 2489–2503.

Dessaud, E., Ribes, V., Balaskas, N., Yang, L.L., Pierani, A., Kicheva, A., Novitch, B.G., Briscoe, J.,
and Sasai, N. (2010). Dynamic assignment and maintenance of positional identity in the ventral
neural tube by the morphogen sonic hedgehog. PLoS Biol. *8*.

Dworkin, S., Boglev, Y., Owens, H., Goldie, S., Dworkin, S., Boglev, Y., Owens, H., and Goldie,
S.J. (2016). The Role of Sonic Hedgehog in Craniofacial Patterning, Morphogenesis and Cranial
Neural Crest Survival. J. Dev. Biol. *4*, 24.

Ericson, J., Norlin, S., Jessell, T.M., and Edlund, T. (1998). Integrated FGF and BMP signaling
controls the progression of progenitor cell differentiation and the emergence of pattern in the
embryonic anterior pituitary. Development *125*, 1005–1015.

Ferrand, R. (1972). [Experimental study of the factors in cytological differentiation of the
adenohypophysis in the chick embryo]. Arch. Biol. (Liege). *83*, 297–371.

Fujikawa, T., Castorena, C.M., Pearson, M., Kusminski, C.M., Ahmed, N., Battiprolu, P.K., Kim,
K.W., Lee, S., Hill, J.A., Scherer, P.E., et al. (2016). SF-1 expression in the hypothalamus is required
for beneficial metabolic effects of exercise. Elife *5*.

Guillemot, F. (2007). Spatial and temporal specification of neural fates by transcription factor codes.
Development *134*, 3771–3780.

Harfe, B.D., Scherz, P.J., Nissim, S., Tian, H., McMahon, A.P., and Tabin, C.J. (2004). Evidence for
an Expansion-Based Temporal Shh Gradient in Specifying Vertebrate Digit Identities. Cell *118*, 517–
528.

Hecksher-Sørensen, J., Hill, R.E., and Lettice, L. (1998). Double Labeling for Whole-Mount In Situ
Hybridization in Mouse. Biotechniques *24*, 914–918.

Hissnauer, T.N., Baranowsky, A., Pestka, J.M., Streichert, T., Wiegandt, K., Goepfert, C., Beil, F.T.,
Albers, J., Schulze, J., Ueblacker, P., et al. (2010). Identification of molecular markers for articular
cartilage. Osteoarthr. Cartil. *18*, 1630–1638.

Jeong, J.H., Mao, J.H., Tenzen, T., Kottmann, A.H., and McMahon, A.P. (2004). Hedgehog signaling
in the neural crest cells regulates the patterning and growth of facial primordia. Genes Dev. *18*, 937–
951.

Jeong, Y., Leskow, F.C., El-Jaick, K., Roessler, E., Muenke, M., Yocum, A., Dubourg, C., Li, X.,
Geng, X., Oliver, G., et al. (2008). Regulation of a remote Shh forebrain enhancer by the Six3
homeoprotein. Nat. Genet. *40*, 1348–1353.

Jeong, Y.S., El-Jaick, K., Roessler, E., Muenke, M., and Epstein, D.J. (2006). A functional screen for
sonic hedgehog regulatory elements across a 1 Mb interval identifies long-range ventral forebrain
enhancers. Development *133*, 761–772.

Kawamura, K., and Kikuyama, S. (1995). Induction from posterior hypothalamus is essential for the
development of the pituitary proopiomelacortin (POMC) cells of the toad (Bufo japonicus). Cell
Tissue Res. *279*, 233–239.

Letelier, J., De La Calle-Mustienes, E., Pieretti, J., Naranjo, S., Maeso, I., Nakamura, T., Pascual-
Anaya, J., Shubin, N.H., Schneider, I., Martinez-Morales, J.R., et al. (2018). A conserved Shh cis-
regulatory module highlights a common developmental origin of unpaired and paired fins. Nat.
Genet. *50*, 504–509.

Lettice, L.A., Williamson, I., Devenney, P.S., Kilanowski, F., Dorin, J., and Hill, R.E. (2014).
Development of five digits is controlled by a bipartite long-range cis-regulator. Development *141*,
1715–1725.

Lu, F., Kar, D., Gruenig, N., Zhang, Z.W., Cousins, N., Rodgers, H.M., Swindell, E.C., Jamrich, M.,
Schuurmans, C., Mathers, P.H., et al. (2013). Rax is a selector gene for mediobasal hypothalamic cell
types. J. Neurosci. *33*, 259–272.

Mackenzie, M.A.F., Jordan, S.A., Budd, P.S., and Jackson, I.J. (1997). Activation of the Receptor
Tyrosine Kinase Kit Is Required for the Proliferation of Melanoblasts in the Mouse Embryo. Dev.
Biol. *192*, 99–107.

Maddin, H.C., Piekarski, N., Sefton, E.M., and Hanken, J. (2016). Homology of the cranial vault in
birds: new insights based on embryonic fate-mapping and character analysis. R. Soc. Open Sci. *3*,
160356.

Marcorelles, P., and Laquerriere, A. (2010). Neuropathology of holoprosencephaly. Am. J. Med.
Genet. Part C Semin. Med. Genet. *154C*, 109–119.

Marcucio, R.S., Young, N.M., Hu, D., and Hallgrimsson, B. (2011). Mechanisms that underlie co-
variation of the brain and face. Genesis *49*, 177–189.

Ming, J.E., and Muenke, M. (2002). Multiple Hits during Early Embryonic Development: Digenic
Diseases and Holoprosencephaly. Am. J. Hum. Genet. *71*, 1017–1032.

Nagy, A., Gertsenstein, M., Vintersten, K., and Behringer, R. (2009a). Alcian blue staining of the
mouse fetal cartilaginous skeleton. Cold Spring Harb. Protoc. *2009*, pdb.prot5169.

Nagy, A., Gertsenstein, M., Vintersten, K., and Behringer, R. (2009b). Alizarin Red Staining of Post-
natal Bone in Mouse. Cold Spring Harb. Protoc. *2009*, pdb.prot5171-pdb.prot5171.

Nord, A.S., Blow, M.J., Attanasio, C., Akiyama, J.A., Holt, A., Hosseini, R., Phouanenavong, S.,
Plajzer-Frick, I., Shoukry, M., Afzal, V., et al. (2013). Rapid and pervasive changes in genome-wide
enhancer usage during mammalian development. Cell *155*, 1521–1531.

Ohyama, K., Ellis, P., Kimura, S., and Placzek, M. (2005). Directed differentiation of neural cells to
hypothalamic dopaminergic neurons. Development *132*, 5185–5197.

Osterwalder, M., Barozzi, I., Tissiéres, V., Fukuda-Yuzawa, Y., Mannion, B.J., Afzal, S.Y., Lee,
E.A., Zhu, Y., Plajzer-Frick, I., Pickle, C.S., et al. (2018). Enhancer redundancy provides phenotypic
robustness in mammalian development. Nature *554*, 239–243.

Pai, Y.J., Leung, K.Y., Savery, D., Hutchin, T., Prunty, H., Heales, S., Brosnan, M.E., Brosnan, J.T.,
Copp, A.J., and Greene, N.D.E. (2015). Glycine decarboxylase deficiency causes neural tube defects
and features of non-ketotic hyperglycinemia in mice. Nat. Commun. *6*.

Pontecorvi, M., Goding, C.R., Richardson, W.D., and Kessaris, N. (2008). Expression of Tbx2 and
Tbx3 in the developing hypothalamic–pituitary axis. Gene Expr. Patterns *8*, 411–417.

Sagai, T., Amano, T., Maeno, A., Ajima, R., and Shiroishi, T. (2019). SHH signaling mediated by a
prechordal and brain enhancer controls forebrain organization. Proc. Natl. Acad. Sci. U. S. A. *116*,
23636–23642.

Sander, M., Paydar, S., Ericson, J., Briscoe, J., Berber, E., German, M., Jessell, T.M., and
Rubenstein, J.L. (2000). Ventral neural patterning by Nkx homeobox genes: Nkx6.1 controls somatic
motor neuron and ventral interneuron fates. Genes Dev. *14*, 2134–2139.

Schwind, J.L. (1928). The development of the hypophysis cerebri of the albino rat. Am. J. Anat. *41*,
295–319.

Sharpe, J. (2003). Optical projection tomography as a new tool for studying embryo anatomy. J.
Anat. *202*, 175–181.

Takuma, N., Sheng, H.Z., Furuta, Y., Ward, J.M., Sharma, K., Hogan, B.L., Pfaff, S.L., Westphal,
H., Kimura, S., and Mahon, K.A. (1998). Formation of Rathke's pouch requires dual induction from
the diencephalon. Development *125*, 4835–4840.

Treier, M., Gleiberman, A.S., O'Connell, S.M., Szeto, D.P., McMahon, J.A., McMahon, A.P., and
Rosenfeld, M.G. (1998). Multistep signaling requirements for pituitary organogenesis in vivo. Genes
Dev. *12*, 1691–1704.

Tsukiji, N., Amano, T., and Shiroishi, T. (2014). A novel regulatory element for Shh expression in
the lung and gut of mouse embryos. Mech. Dev. *131*, 127–136.

Urban-Ciecko, J., and Barth, A.L. (2016). Somatostatin-expressing neurons in cortical networks. Nat.
Rev. Neurosci. *17*, 401–409.

Wada, N., Javidan, Y., Nelson, S., Carney, T.J., Kelsh, R.N., and Schilling, T.F. (2005). Hedgehog
signaling is required for cranial neural crest morphogenesis and chondrogenesis at the midline in the
zebrafish skull. Development *132*, 3977–3988.

Xie, Y., and Dorsky, R.I. (2017). Development of the hypothalamus: conservation, modification and
innovation. Development *144*, 1588–1599.

Yee, S.P., and Rigby, P.W.J. (1993). The regulation of myogenin gene expression during the
embryonic development of the mouse. Genes Dev. *7*, 1277–1289.

Zhao, L., Zevallos, S.E., Rizzoti, K., Jeong, Y., Lovell-Badge, R., and Epstein, D.J. (2012).
Disruption of SoxB1-Dependent Sonic hedgehog Expression in the Hypothalamus Causes Septo-
optic Dysplasia. Dev. Cell *22*, 585–596.

Figure legends

Fig. 1. Loss of SBE2 leads to reduced *Shh* activity in the rostral diencephalon. (A-B) Peri and post-
natal survival for *Shh$^{null/\Delta SBE2}$* compound heterozygotes is shown. Statistical significance was assessed
using the Student's T-test. (C-E) *In situ* hybridisation for *Shh* in E10.5 day heads bisected along the
midline is shown for control, *Shh$^{\Delta SBE2/\Delta SBE2}$* homozygotes and *Shh$^{null/\Delta SBE2}$* compound heterozygotes.
The red and green arrows point to the rostral diencephalic region where *Shh* expression is absent. (F-
H) *In situ* hybridisation for *Ptch1* in E10.5 day bisected heads is shown. Wildtype levels of *Ptch1*
expression in rostral diencephalon in control embryos (indicated by black arrow in F), reduced levels
in *Shh$^{\Delta SBE2/\Delta SBE2}$* (indicated by red arrows in G) and undetectable levels in *Shh$^{null/\Delta SBE2}$* (indicated by
green arrow in H) is shown. Scale bars (C-H) appear in the bottom left hand corner. (I-K) *In situ*
hybridisation for *Shh* in E9.5 embryos is depicted for control, *Shh$^{\Delta SBE2/\Delta SBE2}$* homozygotes and
Shh$^{null/\Delta SBE2}$ compound heterozygotes. The red and green arrows point to the rostral diencephalic region
where *Shh* expression is lost. (L-N) *In situ* hybridisation for *Ptch1* in E9.5 embryos is shown. The red
and green arrows point to the rostral diencephalon. Scale bars (C-N) appear in the bottom left hand
corner. (n=3)

Fig. 2. *Shh$^{null/\Delta SBE2}$* embryos have an altered head shape and gross deformities. (A) Morphometric
analysis of overall head size (log Csize- log centroid size) in control and *Shh$^{null/\Delta SBE2}$* heads at E17.5.
(B) Morphometric analysis of overall head shape in control and *Shh$^{null/\Delta SBE2}$* heads at E17.5. (C-E) Ply
surface of a control E17.5 day CT scanned skull, from a dorsal, lateral and ventral angle. (F-H)
Morphed ply of a randomly chosen control E17.5.surface, morphed based on the average of *Shh$^{null/\Delta SBE2}$*
PC1 landmark changes, presented from a dorsal, lateral and ventral angle. (control n=6; mutant n=8).
(I-L) OPT scan stills from E17.5 day control and *Shh$^{null/\Delta SBE2}$* heads, presenting the midline sagittal
plane and the coronal nasal region. Red arrows point to the region of absent hypothalamic tissue seen
in *Shh$^{null/\Delta SBE2}$* embryos. Red lines serve to demonstrate the reduced nasal cavities found in *Shh$^{null/\Delta SBE2}$*
embryos. Scale bars are depicted in the bottom left or right hand corner. (n=3)

Fig. 3. *Shh$^{null/\Delta SBE2}$* embryos present deformities in the bones of the cranial vault. (A-D) Dual
staining highlighting chondrogenic (blue) and skeletal (red) craniofacial elements in control and
Shh$^{null/\Delta SBE2}$ E17.5 heads are shown. Pale green arrows indicate the basisphenoid bone, yellow arrows
indicate the vomer bone and the white arrows indicate the pterygoid bone. (E-F) H&E staining of E17.5
day parasagittal cryosections from control and *Shh$^{null/\Delta SBE2}$* heads. Black arrows point to the region of
the basisphenoid bone which is intercepted by the adenohypophysis in *Shh$^{null/\Delta SBE2}$* embryos. A zoomed
out view of an E17.5 day head parasagittal cryosection is depicted in (G). The yellow box presents the
region of the head depicted in (E-F). AP- adenohypophysis; NP- neurohypophysis; BS- basisphenoid
bone; BP- base plate of the skull; NA- nasal prominence; FB- forebrain; HB- hindbrain; TG- tongue.
Scale bars are shown in the bottom right or left hand corner. (n=3)

Fig. 4. Bone formation in the viscerocranium of *Shh$^{null/\Delta SBE2}$* embryos. (A-D) *In situ* hybridisation
for *Ptch1* in E12.5 day coronal cryosections of control and *Shh$^{null/\Delta SBE2}$* embryos. (A&C) present the
palatine region and (B&D) present the basisphenoid region. Black arrows point to the region of *Ptch1*
expression seen below the hypothalamic tissue which is absent in *Shh$^{null/\Delta SBE2}$* embryos. (E-H)
Immunofluorescent staining for COL2A1 in E12.5 day coronal cryosections of control and *Shh$^{null/\Delta SBE2}$*

embryos. (E&G) present the palatine region and (F&H) present the basisphenoid region. Red arrows
point to the region of COL2A1 expression seen below the hypothalamic tissue which mostly is absent
in $Shh^{null/\Delta SBE2}$ embryos. 3V- third ventricle; HP- hypothalamus; TG- tongue; RP- Rathke's pouch.
(n=3)

Fig. 5. The AP hypothalamic boundary is shifted ventrally in $Shh^{null/\Delta SBE2}$ embryos. (A-C) H&E
staining of E13.5 day sagittal cryosections of control, $Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$ embryos. (D-F) *In*
situ hybridisation for *Fgf10* in E11.5 day sagittal cryosections of control, $Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$
embryos. Red arrows represent the distance between the MGE and the anterior boundary of *Fgf10*
expression. (G-I) Immunofluorescent staining for TCF4 in E11.5 day sagittal cryosections of control,
$Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$ embryos. White arrows represent the distance between the posterior
infundibulum and the posterior boundary of TCF4 expression. (J)- Quantification of the distance
depicted by the red arrows in (D-F) in control, $Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$ embryos. Statistical
significance was assessed using the Student's T-test. Error bars represent median ± s.e.m.
p***<0.001.(K)- Quantification of the distance between depicted by the white arrows in (G-I) in
control, $Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$ embryos. Statistical significance was assessed using the
Student's T-test. Error bars represent median ± s.e.m. p***<0.001. (L & N) Immunofluorescent
staining for NKX2-1 in E17.5 day sagittal cryosections of control and $Shh^{null/\Delta SBE2}$ embryos. (M & O)
Immunofluorescent staining for CASP3 in E17.5 day sagittal cryosections of control and $Shh^{null/\Delta SBE2}$
embryos. Red circles demarcate the region of cell death apparent in mutant embryos. NS- not
significant. BH- basal hypothalamus; IF- infundibulum; RP- Rathke's pouch; MGE- medial ganglionic
eminence. Scale bars are shown in the bottom left hand corner. (n=3)

Fig. 6. DV hypothalamic patterning is affected by loss of SBE2. (A-C) Immunofluorescent staining
for NKX2-2 in E13.5 day ventral hypothalamic (VH) parasagittal cryosections of control,
$Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$ embryos. Red arrows point to the region of NKX2-2 expression in the
basal hypothalamus which is reduced or absent upon removal of SBE2 enhancer function. (D-F)
Immunofluorescent staining for PAX6 in E13.5 day VH parasagittal cryosections of control,
$Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$ embryos. White arrows point to the region of ectopic PAX6 expression
in the BH which is seen upon removal of SBE2 function. (G-I) Immunofluorescent staining for OLIG2
in E13.5 day VH parasagittal cryosections of control, $Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$ embryos. White
arrows point to the ectopic regions of expression of OLIG2 in the BH which are seen upon removal of
SBE2 function. (J-L) Immunofluorescent staining for NKX6-1 in E13.5 day VH parasagittal
cryosections of control, $Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$ embryos. Red arrows point to the region of
NKX6-1 expression in the AH which is absent upon removal of SBE2 function whilst white arrows
point to the region of ectopic expression in the BH seen in $Shh^{\Delta SBE2/\Delta SBE2}$ embryos. AH- alar
hypothalamus; BH- basal hypothalamus; IF- infundibulum; RP- Rathke's pouch. Scale bars are shown
in the bottom right hand corner. (n=3)

Fig. 7. DV hypothalamic neuronal populations are affected by loss of SBE2. (A-C)
Immunofluorescent staining for NKX2-1 in E13.5 day VH parasagittal cryosections of control,
$Shh^{\Delta SBE2/\Delta SBE2}$ and $Shh^{null/\Delta SBE2}$ embryos. Red arrows point to the regions devoid of NKX2-1 expression
which are found upon removal of SBE2 function. (D-F) Immunofluorescent staining for ISL1 in E13.5

day VH parasagittal cryosections of control, *Shh^{ΔSBE2/ΔSBE2}* and *Shh^{null/ΔSBE2}* embryos. Red arrows point
to the regions of the BH where ISL1 is mostly absent in *Shh^{null/ΔSBE2}* embryos. (G-I) Immunofluorescent
staining for SF1in E13.5 day VH parasagittal cryosections of control, *Shh^{ΔSBE2/ΔSBE2}* and *Shh^{null/ΔSBE2}*
embryos. Red arrows point to the region of the BH where SF1 expression is reduced or absent upon
removal of SBE2 function. (J-L) Immunofluorescent staining for somatostatin in E13.5 day VH
parasagittal cryosections of control, *Shh^{ΔSBE2/ΔSBE2}* and *Shh^{null/ΔSBE2}* embryos. Red arrows point to the
region of the BH where somatostatin is lost in *Shh^{null/ΔSBE2}* embryos. A zoomed in view of the region
of somatostatin expression is presented in the green boxes in the upper right hand corner of (J & K).
AH- alar hypothalamus; BH- basal hypothalamus; IF- infundibulum; RP- Rathke's pouch. Scale bars
are shown in the bottom right hand corner. (n=3)

A	Embryonic day 17.5 (E17.5)			
	Total no.	Control	Single het	Compound het
Observed	41	10	19	12
Expected	-	10	20	10

p=0.7985

B	Post-natal day 14			
	Total no.	Control	Single het	Compound het
Observed	24	4	20	0
Expected	-	6	12	6

p=0.0025

Control | $Shh^{\Delta SBE2/\Delta SBE2}$ | $Shh^{null/\Delta SBE2}$

Shh

Ptch1

E10.5

Shh

Ptch1

E9.5

Figure 1

Figure 1

Control

Mild

Moderate

Severe

Shh^{null/ΔSBE2}

Control

Shh^{null/ΔSBE2}

Figure 2

Figure 3

Figure 4

Figure 6

Figure 7