Replacement of microglia by brain-engrafted macrophages provide long term protection

against brain irradiation and concussive injury

Xi Feng[1,2], Elma S. Frias[1,2], Maria S. Paladini[1,2], David Chen[1,2,3], Zoe Boosalis[1,2],McKenna

Becker[1,2], Sonali Gupta[1,2], Sharon Liu[3], Nalin Gupta[3,6,7] and Susanna Rosi[1,2,3,4,5*]

[1] Brain and Spinal Injury Center

[2] Department of Physical Therapy and Rehabilitation Science

[3] Department of Neurological Surgery

[4] Weill Institute for Neuroscience

[5] Kavli Institute of Fundamental Neuroscience

[6] Brain Tumor Research Center

[7] Department of Pediatrics

University of California San Francisco

* Corresponding Author

Susanna Rosi Ph.D.: Susanna.Rosi@ucsf.edu

1001 Potrero Ave, Building 1, Room 101, Zuckerberg San Francisco General Hospital
Department of Physical Therapy and Rehabilitation Science, University of California, San
Francisco, San Francisco, CA 94110

Running Title: monocyte-derived macrophages protects against brain injury

Keywords: microglia, brain engrafted macrophages, cognition, brain irradiation, CSF-1R inhibitor

Abstract

Brain resident microglia have a distinct origin compared to macrophages in other organs.

Under physiological conditions, microglia are maintained by self-renewal from the local pool,

independent of hematopoietic progenitors. Pharmacological depletion of microglia during

therapeutic whole-brain irradiation prevents synaptic loss and long-term recognition memory

deficits but the mechanisms behind these protective effects are unknown. Here we demonstrate

that after a combination of therapeutic whole-brain irradiation and microglia depletion,

macrophages originating from circulating monocytes engraft into the brain and replace the

microglia pool. Comparisons of transcriptomes reveal that brain-engrafted macrophages have an

intermediate phenotype that resembles both monocytes and embryonic microglia. Brain-

engrafted macrophages display reduced phagocytic activity for synaptic compartments compared

to microglia from normal brains in response to a secondary concussive brain injury. In addition to

sparing mice from brain radiotherapy-induced long-term cognitive deficits, replacement of

microglia by brain-engrafted macrophages can prevent concussive injury-induced memory loss.

These results demonstrate the long-term functional role of brain-engrafted macrophages as a

possible therapeutic tool against radiation-induced cognitive deficits.

Introduction

Brain resident microglia are the innate immune cells of the central nervous system (CNS). Arisen from the yolk sac during embryonic development, microglia actively survey the environment to maintain normal brain functions [1, 2]. Under physiological conditions, microglia are maintained solely by self-renewal from the local pool [3]. Following brain injury and other pathological conditions, microglia become activated and play a central role in the clearance of cellular debris, but if not controlled this process can result in aberrant synaptic engulfment [4-7]. Temporary depletion of microglia can be achieved by using pharmacologic inhibitors of the colony-stimulating factor 1 receptor (CSF-1R) [8]. In the normal brain, treatment with CSF-1R inhibitors (CSF-1Ri) can deplete up to 99% of microglia without causing detectable changes to cognitive functions [8, 9]. Full repopulation occurs within 14 days of inhibitor withdrawal and the repopulated microglia are morphologically and functionally identical to the microglia in naïve brains [9]. Microglia depletion and repopulation by local progenitors has been shown to be beneficial for disease- ,injury- , and age-associated neuropathological and behavioral conditions [10-15]. However, the mechanisms for these protective effects are unknown.

Whole-brain radiotherapy (WBRT), delivered in multiple fractions, is routinely used to treat patients with brain tumors. It is estimated that more than 200,000 patients receive WBRT yearly in the US alone [16]. While it is effective in improving intracranial tumor control, WBRT leads to deterioration of patients' cognitive functions and quality of life [17-19]. Currently, there is no treatment available to prevent or mitigate these adverse effects. Previous studies demonstrated that WBRT causes deleterious effects to the CNS microenvironment by a number of mechanisms including apoptosis of neural progenitor cells, disruption of the blood-brain barrier, activation of microglia and accumulation of peripherally derived macrophages [20-25]. We and others have reported that depletion of microglia during or shortly after brain irradiation in animal models can prevent loss of

dendritic spines in hippocampal neurons and cognitive impairments that develop at later time

points [12-14]. These reports suggest that microglial plays a critical role in inducing synaptic

abnormalities and consequently, cognitive deficits after brain irradiation. The underlying molecular

pathways responsible for the protective effects of repopulated microglia against radiotherapy-

induced neuronal alterations remain unknown.

In the current study, 1) we first defined signature of repopulating cells and analyzed the

transcriptional profile of repopulated brain macrophages from irradiated mouse brains after CSF-

1R inhibitor-mediated depletion. 2) We next sought to establish the origin of repopulated cells

coming from the peripheral monocytic lineage. 3) We identified the functionality of repopulated

macrophages by measuring the ability to engulf synaptic compartments compared to brain

resident microglia. Lastly, 4) we determined the protective properties of brain-engrafted

macrophages (BEMs) against a secondary concussive brain injury-induced cognitive deficits.

Together, these results uncover the mechanism by which brain-engrafted macrophages preserve

hippocampal synapses and memory functions after radiation injury and also in response to an

additional brain injury.

Results

Microglia depletion and repopulation prevents long term radiation-induced memory deficits and loss of hippocampal PSD95.

Temporary microglia depletion during or shortly after exposure to brain irradiation prevents cognitive deficits, suggesting microglia's key role in modifying neuronal and cognitive functions [12-14]. Changes in expression levels of pro-inflammatory cytokine/chemokines have been shown to correlate with cognitive performance in mice [12, 15, 20], however, the exact change in the transcriptional profile of repopulated microglia after brain irradiation is unknown and is an important tool to dissect the roles that repopulated microglia play in preventing of radiation-induced memory deficits. We performed RNA sequencing using repopulated microglia, FACS sorted from irradiated and control mouse brains after CSF-1Ri treatment, and compared with transcriptomes of microglia obtained from mice without CSF-1Ri treatment (**Figure 1a,b**). A CSF-1R inhibitor was used to fully deplete microglia in 8-weeks old male mice, for a duration of 21 days. Three fractions of therapeutic whole-brain irradiation were given to each mouse every other day over five days starting from day 7 of CSF-1R inhibitor treatment. Novel Object Recognition (NOR) test was used to measure recognition memory 4 weeks after the last fraction of WBRT. Consistent with our previous report, fractionated WBRT resulted in impairment in recognition memory, which was prevented by CSF-1R inhibitor treatment (**Figure 1a, lower panel**). One day after the NOR test, mice were euthanized and whole brains were used to sort microglia (CD45^{low}CD11b$^+$ population) for RNA extraction. We used samples from the five best performers in non-impaired groups (control diet + sham, CSF-1R inhibitor + sham/WBTR) and the five worst performers in the memory impaired group (control diet + WBRT) for RNA sequencing (**Figure 1a**).

We previously demonstrated that brain irradiation resulted in reduced density of dendritic spines in hippocampal neurons [13]. To accurately determine the effect of WBRT in the intrinsic synaptic protein levels we measured pre- and post-synaptic markers in the hippocampus by flow-

synaptometry [26, 27]. Fractionated hippocampal cell membranes containing synaptosomes were

enriched and particles between 1 – 3 μm were analyzed to measure synaptic protein levels using

mean fluorescent intensities by FACS (**Figure 1e**). We observed no changes in pre-synaptic

Synapsin-1 protein levels in the hippocampi across all groups (**Figure 1f**). However, we measured

a significant reduction in post-synaptic protein PSD-95 after WBRT, which was completely

prevented by CSF-1R inhibitor mediated microglia depletion (**Figure 1g**). These results cement

the role of microglia in the radiation-induced loss of post-synaptic components after WBRT.

Microglia depletion and repopulation eliminates radiation-induced transcriptome signatures

To identify biological pathways involved in radiation-induced memory deficits, we listed genes

differentially expressed in microglia after WBRT with and without microglia depletion and

repopulation for Gene Ontology Biological Process (GOBP) enrichment analysis. 204 genes were

found to be differentially expressed (DE genes) only in microglia isolated from irradiated brains

(**Figure 1b and Supplementary Table 1**). No enriched GOBP terms were found from the 87

WBRT down-regulated genes (Supplementary Table 1). There were 193 enriched GOBP terms

from the 117 WBRT up-regulated genes, the top 20 enriched GOBP terms are listed in **Figure 1c.**

Almost half (96) of these enriched GOBP terms were associated with increased response to cell

cycle regulation, radiation, DNA repair and stress; the rest enriched GOBP terms were associated

with increased metabolism (21), development (12), regulation of protein kinase activity (8), cellular

adhesion (4) and other functions (**Figure 1d, Supplementary Table** 1). Notably, regardless of

WBRT, the expression of these WBRT-induced DE genes did not change in cells isolated from

brains treated with CSF-1Ri. These results demonstrate that the transcriptomic changes in

microglia induced by WBRT can be completely eliminated after microglia depletion and

repopulation.

qPCR validation of the RNAseq results

To validate the RNAseq results, we next performed qPCR analyses using sorted microglia from animals in the same cohort (**Figure 1b,** and **Supplementary Table 1**). The expression of the toll-like receptor 3 (TLR3) family gene *Lgals9* was significantly increased by irradiation (WBRT + Control diet versus Sham + Control diet) and was at levels comparable to the shams (Sham + Control diet) when treated with CSF-1Ri despite of irradiation (**Supplementary Figure S1a**). *TNFα*, another TLR3 family member which also belongs to GOBP "regulation of response to reactive oxygen species (ROS)", was significantly upregulated by irradiation (WBRT + Control diet versus Sham + Control diet); its expression levels are comparable between the Sham + Control diet and the WBRT + CSF1Ri treated groups. However, *TNFα* remained elevated in microglia from mice treated only by CSF-1Ri (Supplementary Figure S1b). Another gene from the GOBP "regulation of response to ROS", *Sesn2*, was also significantly upregulated by WBRT (**Supplementary Figure S1c**). *Sesn2* remained at the control sham levels in CSF-1Ri only group and was significantly down-regulated in the WBRT + CSF-1Ri group. *Mdm2*, a gene that belongs to GOBP "cellular response to ionizing radiation", was increased after WBRT, and significantly downregulated in in CSF-1Ri treated groups (Supplementary Figure S1d). Other WBRT-induced expression of radiation induced genes, *Ddias*, *Rad51, FoxM1* and *Check 1*, were all at the control sham levels in repopulated microglia regardless of the exposure to WBRT (**Supplementary Figure S1 e – h**). In conclusion, the qPCR validation confirmed that the transcriptomic changes seen in our RNAseq dataset were reliable. These results suggest that CSF-1Ri mediated microglia depletion during WBRT followed by repopulation eliminated radiation-induced signatures in the microglia transcriptome.

Repopulated microglia after WBRT originate from peripheral monocytes

The fractalkine receptor CX3CR1 is expressed in both microglia and peripheral monocytes [28], while chemokine receptor CCR2 is mainly expressed in monocytes [29]. In the Cx3cr1$^{GFP/+}$Ccr2$^{RFP/+}$ reporter mice, the different expression patterns of GFP and RFP can be used to distinguish microglia (GFP+RFP-) from monocytes (GFP+RFP+) [29]. To investigate the cell-of-origin of repopulated microglia in our experimental paradigm we generated bone marrow chimeras with head-protected irradiation using fluorescent labeled bone marrow from Cx3cr1$^{GFP/+}$Ccr2$^{RFP/+}$ donor mice (**Figure 2a**). This allowed partial replacement of bone marrow cells without changing the permeability of the blood-brain-barrier [2, 30, 31]. At 6 weeks after bone marrow transplantation about two thirds of peripheral monocytes were replaced by transplanted cells with fluorescent labels (**Figure 2b**). Bone marrow chimera animals were then treated with WBRT and CSF-1R inhibitor following the same experimental timeline used for RNA sequencing (**Figure 2a**). Next, we compared the compositions of myeloid cells in the brain after CSF-1R inhibitor-mediated depletion and repopulation. Flow cytometry analyses performed 33 days after WBRT revealed that microglia depletion and repopulation alone (Sham + CSF-1Ri) only resulted in limited accumulation of transplanted cells in the brain (**Figure 2c, BMT only**). However, in mice that received WBRT and CSF-1Ri, two thirds of the microglia were replaced by Cx3cr1-GFP labeled cells (**Figure 2c, BMT +fWBI**). These results suggest that microglia depletion during WBRT resulted in significant contribution of the CNS microglia pool by peripheral monocyte-derived BEMs.

Brain-engrafted macrophages retain monocyte signatures

We next assessed the transcriptomic signature of the BEMs after microglia depletion and WBRT by comparing our RNAseq dataset with a previous report by Lavin and Winter *et al* [32]. To minimize

false discovery and noise signals, we examined 1201 genes from this published dataset with a

fold change greater than 1.50 or smaller than 0.667 for down-regulated genes (p<0.05, monocyte

compared to naïve microglia), and found that 1066 genes were expressed in our samples

(**Supplementary Table 2**). Strikingly, the hierarchical clustering of 525 monocyte- and 541

microglia-signature genes revealed that the expression profile of monocyte-derived BEMs (WBRT

+ CSF-1Ri) does not cluster with naïve (Sham + Control diet), irradiated (WBRT + Control diet)

or repopulated (Sham + CSF-1Ri) microglia (**Figure 2d**). Similarity matrix analysis using this

microglia/monocyte signature gene list revealed that the expression pattern in BEMs is different

from naïve, irradiated and repopulated microglia (**Figure 2e**). Next, we counted genes in each

group that expressed in the same trends as microglia or monocyte signature genes from the Lavin

data set to determine the similarity scores to these two cell populations. We found that naïve,

irradiated and repopulated microglia had 60%, 57% and 51% (718, 685 and 612) genes

expressed in the same trends as microglia signature genes, respectively, with minimum similarity

(2-3%) to monocyte signature genes; while BEMs expressed both microglia (28%, 331 genes)

and monocyte signature genes (32%, 386 genes) (**Figure 2f**). These results further confirm that

after microglia depletion and WBRT BEMs originate from blood monocytes.

qPCR validation of microglia- and monocyte-specific genes

To validate microglia and monocyte signature genes that were differentially expressed in our

RNAseq results we next performed qPCR analyses (**Supplementary Figure 2 and**

Supplementary Table 2). Microglia signature genes *Sall1*, *P2ry12*, *Tmem119* and *Trem2* were

expressed at comparable levels in naïve, irradiated and repopulated microglia, while at

significantly lower level in BEMs (**Supplementary Figure 2 a – d**). On the other hand, expression

of monocyte signature gene *Runx3*, was significantly higher in BEMs than other groups

(**Supplementary Figure 2 e**). Notably, previously reported brain-engrafted macrophage specific

genes *Lpar6* and *Pmepa1* [33] have significantly higher expression levels in BEMs after CSF-1Ri

and WBRT treatments compared to other groups (**Supplementary Figure 2 f and g**). In addition,

the expression of *Ccr2*, a monocyte signature gene that was not differentially expressed in our

RNAseq dataset, was also not differentially expressed among the four experimental groups by

qPCR (**Supplementary Figure 2 h**). Taken together, these qPCR results validate our RNAseq

results and cement the notion that BEMs after microglia depletion and WBRT originate from

peripheral monocytes.

Monocyte-derived brain-engrafted macrophages resemble embryonic microglia signatures

Because monocyte-derived BEMs were exposed to the brain microenvironment for a short period

of time, we hypothesized that they were functionally immature. To test this hypothesis, we first

examined genes that were highly expressed at different developmental stages in microglia, and

used yolk sac/embryonic and adult-specific genes as references (called embryonic and adult

signature genes hereon) [34]. Hierarchical clustering of 1617 embryonic and 785 adult microglia

signature genes revealed that transcriptomes of BEMs were highly similar to embryonic microglia,

while the transcriptomes of microglia from other groups were similar to adult microglia and did not

resemble the embryonic one (**Supplementary Figure S3a**). In addition, a similarity matrix

analysis using all 2402 overlapped genes between two datasets showed that BEMs had the

lowest similarity with microglia from other groups (**Supplementary Figure S3b**). In addition, 54%

of the listed genes (n=1306) in BEMs expressed in the same trends as *yolk sac*/embryonic

microglia compared to adult microglia (**Supplementary Figure S3c**). In contrast, naïve (Sham),

irradiated (WBRT only) and repopulated microglia (CSF-1Ri only) had much lower embryonic

signature similarity scores (16%, 19% and 17%, n=381, 445 and 405, respectively,

Supplementary Figure S3c). Notably, naïve, irradiated and repopulated microglia transcriptomes

had high adult signature similarity scores (69%, 59% and 63%, n=1649, 1409 and 1507,

respectively), while BEMs had the lowest adult similarity score (32%, n=759). These data suggest

that the monocyte-derived BEMs start to resemble microglia by expressing more embryonic than

adult microglia transcriptomic signature genes.

Radiation-induced aberrant phagocytosis activity is abrogated in brain-engrafted macrophages

Aberrant loss of synapses during neuroinflammatory conditions has been linked with increased

engulfment of synaptic compartments by microglia [35]. To determine if WBRT affects the

phagocytosis potency of microglia, we injected pre-labeled synaptosomes from a naïve donor

mouse into the hippocampi of mice after WBRT and CSF-1R inhibitor treatment and measured

engulfment by microglia using flow cytometry (**Figure 3a**). After WBRT there was a significant

increase in the number of microglia engulfing synaptosomes in the hippocampus compared to

naïve non-irradiated animals (**Figure 3b**). Strikingly, synapse engulfment activity was unchanged

compared to naïve animals in animals treated with CSF-1R inhibitor during WBRT (**Figure 3b**).

Immunofluorescent imaging at the injection sites confirmed that the injected synaptosomes were

indeed engulfed by microglia, and the increased trend of engulfment by irradiated microglia

remained unchanged (**Figures 3 c and d, Supplementary Figure 4a).** Notably, after

hippocampal injection of fluorescent labeled latex beads into the hippocampus, we found that

WBRT resulted in increased engulfment of latex beads was also inhibited by CSF-1R inhibitor

treatment, suggesting that the WBRT-induced increase of engulfment was not specific to

synaptosomes, but rather a general increase of phagocytosis potency (**Supplementary Figure**

4b). These data are the first to demonstrate that WBRT results in an increase in microglial

phagocytosis activity in the hippocampus that can be prevented by transient microglia depletion

and full repopulation.

Irradiation-induced complement and phagocytic receptors expression in microglia are absent in

BEM after WBRT.

Microglial complement receptors play essential roles in physiologic synaptic elimination during

development and aberrant elimination during neuroinflammatory conditions [35, 36]. To understand

the mechanisms of increased microglia phagocytic activity after WBRT, we measured expression

levels of a list of complement receptors, phagocytic markers and lysosome proteins in microglia

by flow cytometry. The expression of complement receptor C5aR was significantly elevated in

microglia at one month after WBRT. However, in animals treated with CSF-1Ri C5aR expression

was unchanged from naïve animals (**Figure 3e**). The same trend was observed in the expression

levels of CD68 and lysosomal-associated membrane protein 1 (LAMP-1) (**Figure 3 f and g**).

These results were consistent with our data demonstrating decreased PSD95 levels (Figure 3b)

and increased microglial phagocytosis activity in the hippocampus after WBRT (**Figure 3 b and**

d). In addition, complement receptor CR3 (CD11b) was significantly elevated in microglia after

WBRT or CSF-1Ri treatments alone, and remained unchanged in BEMs with combined WBRT

and CSF-1Ri treatments (**Supplementary Figure S5a**). No changes in the complement receptor

C3ar1 were measured after WBRT or CSF-1R inhibitor treatment (**Supplementary Figure S5b**).

These results demonstrate that the increased microglia phagocytosis of synaptosomes after

WBRT was associated with increased phagocytic and lysosome proteins, and was likely through

the complement pathways.

Brain-engrafted macrophages after microglia depletion persist in the brain

To determine whether BEMs are present long-term in the brain, we monitored this cellular

population for 6 months after WBRT. To eliminate the limitation of using bone marrow obtained

from the Cx3cr1$^{+/GFP}$Ccr2$^{+/RFP}$ knock-in reporter mouse strain, we used an actin-GFP transgenic

line as bone marrow donors and generated chimeras using the same body-only irradiation

protocol (**Figure 4 a**). Six weeks later, mice were treated with CSF-1R inhibitor and WBRT and

then used to trace BEMs at varies time points (**Figure 4 a**). Whole coronal sections at the level of

the dorsal hippocampus were stained with Iba1 and imaged to quantify total Iba1+ and GFP+

cells (**Supplementary Figure 6 a**). We found that all GFP+ cells in the brain were also Iba1+,

suggesting that BEMs were indeed derived from the periphery. In addition, the morphology of

Iba1+GFP+ BEMs were analyzed and compared to Iba1+GFP- microglia (**Figure 4 b and d**). We

found that round-shaped Iba1+GFP+ BEM cells entered the brain starting from 7 days after WBRT,

and started to obtain more processes that resembled microglia morphology (**Figure 4 b**). However,

Sholl analysis demonstrated that the morphology of BEMs remained stable from 33 days after

WBRT and never reached the structural complexity of microglia (**Figure 4c, Supplementary**

Figure S7). We found that 40 – 90% of Iba1+ cells are also GFP+ at 14 days after WBRT. This

ratio remained at high levels at 1, 3 and 6 months after WBRT (**Figure 4 e**). Interestingly, although

the Iba1+ and Iba1+GFP+ cell numbers are not fully recovered at 14 days after WBRT, the

microglia replacement ratio was similar to the level of later time points (Figure 5 e and

Supplementary Figure S6 b and c), suggesting a non-competitive manner of brain parenchyma

occupancy by microglia and BEMs. These data demonstrate that BEMs enter the brain shortly

after WBRT, adapt to a microglia-like morphology and maintain a stable population.

BEMs provide long-term protection against WBRT-induced memory deficits and hippocampal

dendritic spine loss

To measure the long-term cognitive outcomes, we treated a batch of wildtype animals, and tested

their recognition memory at 3 and 6 months after WBRT (**Figure 5 a**). We found that WBRT

resulted in persistent loss of recognition memory also at 3 and 6 months, while CSF-1Ri treatment

alone did not alter recognition memory performance (**Figure 5 b and c**). Strikingly, mice that

received WBRT along with temporary microglia depletion did not show any memory deficits and

performed undistinguishable from control animals at 3 and 6 months (**Figure 5 b and c**). Our

previous report demonstrated that WBRT-induced dendritic spine loss in hippocampal neurons

was fully prevented by temporary microglia depletion during irradiation (**Figure 5 d,** replotted

using previously published data) [13]. In this study, we sought to understand if the protective effects

persisted up to 6 months after WBRT. Our results clearly show that radiation-induced loss of

dendritic spines in hippocampal neurons persists to this time point, and that the protective effect

of microglia depletion and subsequent replacement by BEMs is long lasting (**Figure 5 e**). Taken

together, brief depletion of microglia during WBRT induces sustainable BEMs in the brain and

provides long-term protection against irradiation-induced deficits in recognition memory.

Replacement of microglia by BEMs protects against concussive injury-induced memory loss

To investigate the function of BEMs after they replaced microglia, a single mild concussive Closed

Head Injury (CHI) was given to mice 30 days after CSF-1R inhibitor treatment and WBRT;

microglia/BEMs morphology and phagocytic activities were measured following recognition

memory test by NOR (**Fig 6 a**). By FACS analysis at 24 days post injury, we found that

phagocytosis activity increased (p=0.0419) after CHI in microglia but not in BEMs (**Figure 6 b**).

Quantification of immunofluorescent staining of Iba-1/PSD-95 co-localization revealed a trend of

increased engulfment towards pre-stained synaptosomes by microglia but not by BEMs (**Fig 6 c**

and d). In addition, the structural complexity of microglia decreased in Sholl analysis, while the

morphology of BEMs remained unchanged after CHI (**Figure 6 e**). Furthermore, at 20 days after

injury, CHI-induced recognition memory deficits were spared in mice whose microglia were

replaced by BEMs (**Figure 6 f**). These results demonstrate that unlike resident microglia which

transition to a less ramified morphology and exhibit increased phagocytic activity towards injected

synaptosomes, BEMs remain unchanged in both morphology and phagocytic activity in response

to CHI. More importantly, our data suggest that replacement of microglia by BEMs can protect

against CHI-induced memory loss.

Discussion

Here we provide evidence for the direct involvement of microglia phagocytic activity

towards synaptic compartments as a mechanistic cause for loss of dendritic spines with

consequent impairments in memory functions after WBRT. Replacement of microglia with

monocyte derived BEMs prevents loss of synapses and consequent memory deficits. Importantly,

BEMs replacing microglia are also protective against a second injury to the brain. Together our

results unravel novel immediate and long lasting therapeutic benefits of microglia depletion and

repopulation during WBRT.

Microglia play pivotal roles in reshaping synaptic networks during neonatal brain

development [37, 38]. They engulf synaptic elements by active synaptic pruning in an activity- and

complement-dependent manner [38]. Microglia-driven aberrant loss of synapses and consequent

impairment of cognitive functions have also been reported in animal models of AD [35], infection [39],

injury [40, 41], and aging [42]. Using RNA sequencing, we compared the transcriptomes of microglia

from irradiated and non-irradiated brains after CSF-1Ri-mediated microglia depletion and

repopulation. WBRT induces increased expression of genes that mainly belong to cell cycle

regulation, DNA damage repair and stress-induced biological processes (**Figure 1d**). As a result,

activated microglia have higher engulfing potential towards both intrinsic and extrinsic synaptic

compartments (**Figures 1 g, Figure 3 b - d**). This view is further supported by the increased

expression of endosome/lysosome proteins CD68 and CD107a with the complement receptors

CR3 and C5ar1 measured in microglia chronically after WBRT (**Figure 3 e – g, and**

Supplementary Figure 5). Notably, both endosome/lysosome proteins and complement receptor

expressions were comparable to naïve microglia (sham + control diet) in BEMs (WBRT + CSF-

1Ri) and repopulated microglia (CSF-1Ri only). These results suggest that the loss of

hippocampal synapses after WBRT may be dependent on the activation of the alternative

complement pathway. Interestingly, although BEMs are morphologically similar to adult microglia,

they retain a transcriptomic signature similar to both circulating monocytes and embryonic

microglia (**Figure 2 and Supplementary Figure 3**). It is plausible that the BEMs are in a

transitional state between peripheral monocytes and CNS microglia at early developmental

stages.

The decrease in post-synaptic protein PSD95 level in hippocampal synaptosomes is also

paralleled with reductions in hippocampal dendritic spines (**Figures 5 d, e and Feng et al**[13]).

However, pre-synaptic Synapsin 1 protein levels are not affected by WBRT or microglia depletion,

suggesting that WBRT mainly induces loss of post-synaptic compartments. Interestingly, although

the phagocytosis potency of repopulated microglia and BEMs are both low (**Figures 3 b and d**),

microglia depletion and repopulation alone does not affect dendritic spine density (**Figures 5 d**

and e). On the other hand, microglia replacement by BEMs results in increased dendritic spine

density compared to those with radiation alone, and microglia depletion alone (**Figure 5 d, data**

re-plotted from Feng et al[13]). Strikingly, the protective effect of microglia depletion during WBRT

results in intact memory functions and extends to 3- and 6-months following irradiation (**Figure 5**

b and c). The dendritic spine density in mice that received WBRT and CSF-1Ri remained higher

than those who only received CSF-1Ri (**Figure 5 e**) suggesting that in an non-reactivate state

(evidenced by no changes in genes involved in cell cycle and radiation response, in microglial

phagocytosis and lysosome proteins, or in phagocytosis activity towards injected synaptosomes

and latex beads) of repopulated microglia and BEMs may have intrinsic differences in maintaining

the homeostasis of dendritic spines, which appears to diminish over time.

In the CNS, microglia maintain a stable population by self-renewal in either a random

manner or through clonal expansion [3, 43]. CSF-1R inhibitor treatment alone results in acute

depletion of up to 99% of CNS resident microglia, with repopulated microglia arising solely from

the residual microglia and their progenitor cells that remain after treatment [8, 44]. The repopulated

microglia have transcriptional and functional profiles similar to naïve microglia [9]. Peripheral

macrophages can engraft into the brain but remain morphologically, transcriptionally and

functionally different from CNS resident microglia [45, 46,]. Under specific circumstances, monocytes

entering the CNS can become microglia-like cells. This is most clearly demonstrated in

experiments where lethal whole-body irradiation was followed by bone marrow transplantation

with labeled monocytes (Ccr2$^+$Ly6C^{high}), resulting in accumulation of these cells in the brain [30]. In

neonatal mouse brains monocytes can enter the brain parenchyma without head irradiation and

become microglia-like cells at a low frequency [47]. In addition, chronic depletion of microglia without

irradiation also results in myeloid cells entering the CNS and becoming BEMs [33]. However, the

roles of BEMs in cognitive functions are largely unknown. Here we report that concurrent microglia

depletion and therapeutic brain irradiation causes peripheral monocytes to enter the brain

parenchyma and become microglia-like BEMs. BEMs enter the brain at 14 days after the

completion of brain irradiation, or 4 days after the CSF-1Ri withdrawal (**Figure 4 e, and**

Supplementary Figure 6). Notably, although the ratio of BEMs was high at this time point the

total number of Iba1 positive cells is not fully recovered (**Supplementary Figure 6 b**). This ratio

remains at high levels in head-irradiated mice throughout the current study (**Figure 5 e**),

suggesting that microglia depletion during WBRT results in sustainable replacement of microglia

by BEMs. Importantly, this observation correlates with long-term protection against WBRT-

induced loss of recognition memory and dendritic spines in hippocampal granule neurons (**Figure**

6 b – e).

In the clinic, cancer patients are unlikely to receive a second round of radiotherapy to the

brain. Therefore, instead of introducing a second round of WBRT, after they occupied the brain

we gave BEM-bearing mice CHI that causes memory deficits[48, 49], and further examined BEMs'

response to a single head trauma. Our data show that microglia had increased phagocytic

potential to injected synaptosomes after CHI, while phagocytic activity of BEMs did not change

and remained at a similar level as naïve microglia (**Figure 6 b and c**). This is further demonstrated

by Sholl analysis of BEMs showing no change in morphology after CHI (**Figure 6 e**). Most

importantly, CHI-induced memory deficit was prevented in BEM-bearing mice (**Figure 7 f**). These

data are the first to demonstrate that BEMs can prevent brain injury-induced cognitive dysfunction.

Further lineage tracing, transcriptomic and functional studies with different microglia

depletion models will help answer the following questions: 1) whether the delayed engraftment of

BEMs after WBRT seen at day 14 is due to the suppression of cell trafficking from the periphery

to the CNS or survival of newly engrafted BEMs that depend on the CSF-1R. 2) The mechanism

of the long-term maintenance of BEMs after WBRT. Whether it is achieved by continuous

engraftment of new BEMs or by colonization of BEMs that are already in the brain parenchyma.

In conclusion we report evidence for the mechanism by which microglia depletion and

repopulation after WRBT prevents memory loss. Our results demonstrate that replacement of

CNS resident microglia by peripheral monocyte-derived BEMs results in a transcriptional and

functional reset of immune cells in the brain to an inactive state, which spares the brain from

WBRT-induced dendritic spine loss in hippocampal neurons and recognition memory deficits.

Most importantly, replacement of microglia by BEMs protects against concussive brain injury-

induced cognitive deficits. These results suggest that replacement of depleted microglia pool by

peripheral monocyte-derived BEMs represents a potent treatment for irradiation-induced memory

deficits.

Materials and Methods

Animals: All experiments were conducted in compliance with protocols approved by the Institutional Animal Care and Use Committee at the University of California, San Francisco (UCSF), following the National Institutes of Health Guidelines for Animal Care. 7 weeks old C57BL/6J male mice were purchased from the Jackson Laboratory and housed at UCSF animal facilities and were provided with food and water ad libitum. All mice were habituated for one week before any treatments or procedures. 8–10 weeks old $Cx3cr1^{GFP/+}Ccr2^{RFP/+}$ mice were breed by crossing the $Cx3cr1^{GFP/GFP}Ccr2^{RFP/RFP}$ line with wildtype C57BL/6J mice, and used as donors for the bone marrow chimeras.

CSF-1Ri treatment: CSF-1Ri (PLX5622 formulated in AIN-76A standard chow at 1200 ppm, Research Diets, Inc) were provided by Plexxikon, Inc (Berkeley, CA). Mice were given free access to either CSF-1Ri chow or control diet (AIN-76A without PLX5622) for 21 days. Approximately 4.8 mg of PLX5622 was ingested by each mouse per day in the treated group (calculation based on 4 g/mouse daily consumption).

Fractionated whole-brain radiotherapy (WBRT): 8 weeks old mice were injected with ketamine (90mg/kg) /Xylazine (10 mg/kg) mix. When fully immobilized mice were placed in irradiator with cesium-137 source at the dose rate of 2.58 Gy/min. The body was shielded with a lead collimator that limited the radiation beam to a width of 1 cm to cover the brain. Three radiation fractions (3.3 Gy) were delivered every other day over 5 days. Sham animals received ketamine/xylazine without irradiation.

Bone marrow chimeras: 8 weeks old C57BL/6J mice were used as bone marrow recipients. 8 weeks old males received two doses of 6 Gy cersium-137 irradiation at the dose rate of 2.58 Gy/min with head protected by lead plates 6 hours apart. Bone marrow cells from 6–10 weeks old donors $Cx3cr1^{+/GFP}Ccr2^{+/RFP}$ or B6-EGFP (The Jackson Laboratory, stock No 003291) were isolated and resuspended in sterile saline at a concentration of 100 million cells/ml. 0.1 ml of bone

marrow cells were injected into recipients via retro-orbital injection immediately after the second

head protected irradiation. Bone marrow chimeras were housed with 1.1 mg/ml neomycin as

drinking water for 4 weeks and allowed an additional 2 weeks to recover before any treatments.

Concussive TBI – Closed head injury: 12 weeks old C57BL/6J mice were randomly assigned

to each TBI or sham surgery group. Animals were anesthetized and maintained at 2-2.5%

isoflurane during CHI or sham surgery. Animals were secured to a stereotaxic frame with

nontraumatic ear bars and the head of the animal was supported with foam. Contusion was

induced using a 5-mm tip attached to an electromagnetic impactor (Leica) at the following

coordinates: anteroposterior, −1.50 mm and mediolateral, 0 mm with respect to bregma. The

contusion was produced with an impact depth of 1 mm from the surface of the skull with a velocity

of 5.0 m/s sustained for 300 ms. Animals that had a fractured skull after injury were excluded from

the study. Sham animals were secured to a stereotaxic frame with nontraumatic ear bars and

received the midline skin incision but no impact. After CHI or sham surgery, the scalp was sutured

and the animal was allowed to recover in an incubation chamber set to 37 °C. All animals

recovered from the surgical procedures as exhibited by normal behavior and weight maintenance

monitored throughout the duration of the experiments.

Synaptosome isolation staining and injection: Fresh hippocampi from a naïve mouse was

homogenized and spun down in 0.32M sucrose solution (dissolved in 50 mM HEPES buffer).

Supernatant was centrifuged in 0.65M sucrose solution at 12,000 rpm for 30 minutes at 4°C. The

synaptosome containing pellet was resuspended in 1 x ice-cold PBS, diluted to 100 µg/ml, and

stained with PSD-95 antibody (Millipore) on ice for 30 minutes followed by a secondary antibody

staining (Invitrogen, goat anti-mouse AF488). Stained synaptosomes were washed and diluted

20 times in PBS and stored at -80°C. 2 µl of pre-stained synaptosomes were injected into the

right hippocampus of each recipient mouse at the coordinate relative to the bregma: AP + 1.6 mm,

ML + 1.6 mm and DV -2.0 mm. Mice were euthanized 3 days later. The left hemispheres

(uninjected) were used for phagocytosis markers staining and the right hemispheres (injected)

were used to assess synaptosome phagocytosis levels by flow cytometry or immunofluorescent

staining.

Immunofluorescent Staining: hemi-brains with synaptosome injection were fixed in 4% PFA

overnight, cryo-protected in 30% sucrose solution in 1 x PBS and sliced in 20 μm sections. Sliced

tissues were stained with Iba1 (Fujifilm Wako Pure Chemical Corporation, 019-19741) followed

by a secondary antibody staining (goat anti-rabbit AF568, Invitrogen, A-11011). DAPI was used

for nuclear staining. Images close to the injection site (Supplementary Figure 1a) were taken using

a Zeiss Imager Z1 microscope under a 20x objective lens. Tissues from bone marrow chimeras

were processed and stained as described above. Images were taken using a CSU-W1 Nikon

Spinning Disk Confocal microscope under 10x air, 20x air or 100x immerse oil lenses. All images

were analyzed using the Fiji/ImageJ software by experimenters blinded to sample information.

Behavior test: Novel Object Recognition (NOR) task was used to test hippocampal dependent

recognition memory at one, three and six months after the last dose of irradiation. All tests took

place during the dark cycle in a room with dim red light as previously described [13, 14]. Briefly, mice

were habituated in an open arena (30 cm x 30 cm x 30 cm, L x W x H) for 10 minutes on day one

and day two. On day three, two identical objects were put into the arena at a distance of 21 cm

and mice were allowed to explore for 5 minutes. On day four, one object was replaced by a novel

object and mice were allowed to explore for 5 minutes. All trials were recorded by an overhead

camera and analyzed using Ethovision software. Data are presented as discrimination Index,

calculated using fomular $DI = (T_{Novel} - T_{Familiar})/(T_{Novel} + T_{Familiar})$.

Flow cytometry: mice were perfused with cold PBS after euthanasia. Brains were immediately

removed and dissociated using a Neural Tissue Dissociation kit (P) (Miltenyi Biotec). Brain cells

were resuspended in 30% Percoll solution diluted in RPMI medium, and centrifuged at 800 g for

30 minutes at 4°C. Cell pellets were washed with FACS buffer (1 x DPBS with 0.5% BSA fraction

V and 2% FBS), blocked with mouse CD16/32 Fc block (BD Biosciences #553141) and stained

with fluorophore conjugated antibodies (CD11b-AF700, CD45-FITC, BD Pharmingen 557690 and

553080, C5aR-PE, CD68-PE and CD107a-PE, Miltenyi Biotec 130-106-174, 130-102-923 and

130-102-219), washed with FACS buffer and used for sort or analyses of bone marrow chimera

efficiency. Data were collected on an Aria III sorter using the FACSDIVA software (BD

Biosciences, V8.0.1), and analyzed with Flowjo software (FlowJo, LLC, V10.4.2).

Flow synaptometry: after isolation (described above) synaptosomes were stained with PSD-95

(Abcam ab13552) or Synapsin-1 (Millipore #1543) antibodies on ice for 30 minutes, washed and

followed by a secondary antibody staining (Invitrogen, goat anti-mouse AF488, A-11001). Stained

synaptosomes were used immediately for analysis of mean fluorescent intensity measurement.

Fluorescent latex beads of 1 μm, 2 μm, 3 μm and 6 μm were used as references of particle sizes

in the FSC-A vs SSC-A dot plot. Events between 1 μm and 3 μm were used to measure mean

fluorescent intensities of isolated synaptosomes under the FITC channel. Data were collected on

an Aria III sorter using the FACSDIVA software, and analyzed with Flowjo software. At least

100,000 events were collected from each sample for the analyses.

RNA sequencing: mRNA was isolated from 100,000 to 400,000 sorted microglia or BEMs using

the Dynabeads mRNA DIRECT Purification Kit (Invitrogen #61011) following the manufacturer's

instructions. RNA sequencing libraries were generated using the Ovation RNA-seq system V2

and Ultralow Library Construction System sample prep kits (NuGEN). Libraries were sequenced

on the HiSeq 2500 to generate single end 50bp reads according to the manufacturer's instructions.

Normalized per-gene read counts were used to compare relative gene expression levels across

samples. Only genes with average read counts greater than 10 were included for analyses.

Heatmaps were drawn using the online analysis software Morpheus (Broad Institute,

https://software.broadinstitute.org/morpheus), followed by hierarchical clustering using the One

minus pearson correlation method. Gene Ontology analysis was performed using the Statistical

overrepresentation test (GO biological process complete, PANTHER version 14) [50]. Bar graphs

to visualize fold enrichment and p values of enriched GO biological pathways were drawn using

the GraphPad Prism software (V 7.01, GraphPad Software, Inc). For analysis of

monocyte/microglia signature genes, dataset from Lavin and Winter et al was used as reference

(GSE63340) [32]. Genes significantly up or down regulated (p<0.05, fold-change > 1.5 or <0.667)

in monocytes vs microglia comparisons are defined as monocyte or microglia signature genes,

respectively. Heatmaps were drawn as described above, and similarity matrix were drawn using

the Morpheus online tool with Pearson correlation. Monocyte/microglia similarity scores were

calculated based on the numbers of genes in each treatment group from this study that expressed

in the same trend as monocyte/microglia signature genes (genes with fold-change between

0.6667 and 1.500 or with p>0.05 were defined as unspecified). For juvenile/embryonic signature

analysis, dataset from Matcovitch-Natan and Winter et al was used as reference (GSE79819) [34].

Gene listed to be highly expressed in Yolk Sac and embryonic day 10.5–12.5 were defined as

embryonic/juvenile   microglia   signatures,   genes   highly   expressed   in   adult

cortex/hippocampus/spinal cord were defined as adult microglia signatures. Heatmaps, similarity

matrix and similarity scores were drawn or calculated as described above.

qPCR: mRNAs were extracted from sorted microglia using the Dynabeads mRNA DIRECT

Purification Kit (Invitrogen #61011), and reverse transcribed into cDNAs using reverse

transcription kit (info) . qPCR reactions were set up in duplicate reactions using the PowerUp

SYBR Green Master Mix kit (Applied Biosystems #A25777) using an Mx3000P qPCR System

(Agilent, Santa Clara, CA) following the manufacturer's instructions. Data were analyzed using

the standard curve method. Standard cDNAs were generated with total RNAs from mixed naïve

and irradiated mouse brains. qPCR primers sequences are listed in Supplementary Table 4.

Sholl analysis: Images of GFP+ (BEMs from bone marrow chimeras) or Iba1+ (AF555, all

microglia cells, and BEMs from non-bone marrow chimeras) cells were acquired from stained

frozen sections (20um) using a confocal microscope under 100x objectives (CSU-W1 Spinning

Disk/High Speed Widefield, Nikon). Max Z-projections from Z-stack images (0.26um step size)

were used for Sholl analysis[51] in Fiji[52] software using the following settings: manually defined cell

center at the cell body, the numbers of intersections between cellular processes and circles with

incremental radius (2um step size, up to 60um) were recorded, plotted and compared across

samples.

Statistical analyses: Two-way ANOVA was used to determine radiation and CSF-1Ri treatment

effects for NOR, qPCR, flowsynaptometry, flow cytometry, immunofluorescent staining counts

and dendritic spine count results, with Tukey's post hoc multiple comparisons. One-way ANOVA

with Sidak's post hoc multiple comparisons was used to determine effect of developmental stages

for dataset published by Matcovitch-Natan and Winter et al. Unpaired t-test was used to determine

differentially expressed microglia/monocyte signature genes from dataset published by Lavin and

Winter et al. Unpaired t-test was used calculate the p value of the comparison of BEMs

contributions between the BMT and BMT + WBRT groups. Exact p values and numbers of animals

used in each experiment were listed in each related figure legend. All error bars represent mean

± SEM.

Reference

1.   Alliot, F., Godin, I. & Pessac, B. Microglia derive from progenitors, originating from the yolk sac,
and which proliferate in the brain. *Developmental Brain Research* **117**, 145-152 (1999).
2.   Ginhoux, F. *et al.* Fate Mapping Analysis Reveals That Adult Microglia Derive from Primitive
Macrophages. *Science* **330**, 841 (2010).
3.   Ajami, B., Bennett, J.L., Krieger, C., Tetzlaff, W. & Rossi, F.M.V. Local self-renewal can sustain
CNS microglia maintenance and function throughout adult life. *Nature Neuroscience* **10**, 1538
(2007).
4.   Prinz, M., Priller, J., Sisodia, S.S. & Ransohoff, R.M. Heterogeneity of CNS myeloid cells and their
roles in neurodegeneration. *Nat Neurosci* **14**, 1227-1235 (2011).
5.   Saijo, K. & Glass, C.K. Microglial cell origin and phenotypes in health and disease. *Nat Rev
Immunol* **11**, 775-787 (2011).

6.   Hong, S., Dissing-Olesen, L. & Stevens, B. New insights on the role of microglia in synaptic
583   pruning in health and disease. *Current opinion in neurobiology* **36**, 128-134 (2015).
584   7.   Hong, S. & Stevens, B. Microglia: Phagocytosing to Clear, Sculpt, and Eliminate. *Developmental*
585   *Cell* **38**, 126-128 (2016).
586   8.   Elmore, M.R. *et al.* Colony-stimulating factor 1 receptor signaling is necessary for microglia
587   viability, unmasking a microglia progenitor cell in the adult brain. *Neuron* **82**, 380-397 (2014).
588   9.   Elmore, M.R., Lee, R.J., West, B.L. & Green, K.N. Characterizing newly repopulated microglia in
589   the adult mouse: impacts on animal behavior, cell morphology, and neuroinflammation. *PLoS*
590   *One* **10**, e0122912 (2015).
591   10.   Dagher, N.N. *et al.* Colony-stimulating factor 1 receptor inhibition prevents microglial plaque
592   association and improves cognition in 3xTg-AD mice. *Journal of neuroinflammation* **12**, 139
593   (2015).
594   11.   Spangenberg, E.E. *et al.* Eliminating microglia in Alzheimer's mice prevents neuronal loss without
595   modulating amyloid-β pathology. *Brain* (2016).
596   12.   Acharya, M.M. *et al.* Elimination of microglia improves cognitive function following cranial
597   irradiation. *Scientific reports* **6**, 31545 (2016).
598   13.   Feng, X. *et al.* Colony-stimulating factor 1 receptor blockade prevents fractionated whole-brain
599   irradiation-induced memory deficits. *Journal of neuroinflammation* **13**, 215 (2016).
600   14.   Feng, X., Liu, S., Chen, D., Rosi, S. & Gupta, N. Rescue of cognitive function following fractionated
601   brain irradiation in a novel preclinical glioma model. *eLife* **7**, e38865 (2018).
602   15.   Krukowski, K. *et al.* Temporary microglia-depletion after cosmic radiation modifies phagocytic
603   activity and prevents cognitive deficits. *Scientific reports* **8**, 7857 (2018).
604   16.   Rapp, S.R. *et al.* Donepezil for Irradiated Brain Tumor Survivors: A Phase III Randomized Placebo-
605   Controlled Clinical Trial. *Journal of Clinical Oncology* **33**, 1653-1659 (2015).
606   17.   Stupp, R. *et al.* Radiotherapy plus Concomitant and Adjuvant Temozolomide for Glioblastoma.
607   *New England Journal of Medicine* **352**, 987-996 (2005).
608   18.   Greene-Schloesser, D. & Robbins, M.E. Radiation-induced cognitive impairment--from bench to
609   bedside. *Neuro Oncol* **14 Suppl 4**, iv37-44 (2012).
610   19.   Shinohara, C., Gobbel, G.T., Lamborn, K.R., Tada, E. & Fike, J.R. Apoptosis in the Subependyma of
611   Young Adult Rats after Single and Fractionated Doses of X-Rays. *Cancer Research* **57**, 2694-2702
612   (1997).
613   20.   Morganti, J.M., Jopson, T.D., Liu, S., Gupta, N. & Rosi, S. Cranial Irradiation Alters the Brain's
614   Microenvironment and Permits CCR2[+] Macrophage Infiltration. *PLoS ONE* **9**, e93650 (2014).
615   21.   Warrington, J.P. *et al.* Whole Brain Radiation-Induced Vascular Cognitive Impairment:
616   Mechanisms and Implications. *Journal of Vascular Research* **50**, 445-457 (2013).
617   22.   Hua, K., Schindler, M.K., McQuail, J.A., Forbes, M.E. & Riddle, D.R. Regionally distinct responses
618   of microglia and glial progenitor cells to whole brain irradiation in adult and aging rats. *PLoS One*
619   **7**, e52728 (2012).
620   23.   Chiang, C.S., McBride, W.H. & Withers, H.R. Radiation-induced astrocytic and microglial
621   responses in mouse brain. *Radiotherapy and Oncology* **29**, 60-68 (1993).
622   24.   Mizumatsu, S. *et al.* Extreme Sensitivity of Adult Neurogenesis to Low Doses of X-Irradiation.
623   *Cancer Research* **63**, 4021-4027 (2003).
624   25.   Monje, M.L. *et al.* Impaired human hippocampal neurogenesis after treatment for central
625   nervous system malignancies. *Annals of Neurology* **62**, 515-520 (2007).
626   26.   Fein, J.A. *et al.* Co-localization of amyloid beta and tau pathology in Alzheimer's disease
627   synaptosomes. *The American journal of pathology* **172**, 1683-1692 (2008).

27.   Prieto, G.A. *et al.* Synapse-specific IL-1 receptor subunit reconfiguration augments vulnerability
to IL-1β in the aged hippocampus. *Proceedings of the National Academy of Sciences* **112**, E5078
(2015).
28.   Jung, S. *et al.* Analysis of Fractalkine Receptor CX₃CR1 Function by
Targeted Deletion and Green Fluorescent Protein Reporter Gene Insertion. *Molecular and*
cellular biology **20**, 4106 (2000).
29.   Saederup, N. *et al.* Selective Chemokine Receptor Usage by Central Nervous System Myeloid
Cells in CCR2-Red Fluorescent Protein Knock-In Mice. *PLoS ONE* **5**, e13693 (2010).
30.   Mildner, A. *et al.* Microglia in the adult brain arise from Ly-6ChiCCR2+ monocytes only under
defined host conditions. *Nat Neurosci* **10**, 1544-1553 (2007).
31.   Priller, J. *et al.* Targeting gene-modified hematopoietic cells to the central nervous system: Use
of green fluorescent protein uncovers microglial engraftment. *Nature Medicine* **7**, 1356-1361
(2001).
32.   Lavin, Y. *et al.* Tissue-resident macrophage enhancer landscapes are shaped by the local
microenvironment. *Cell* **159**, 1312-1326 (2014).
33.   Cronk, J.C. *et al.* Peripherally derived macrophages can engraft the brain independent of
irradiation and maintain an identity distinct from microglia. *The Journal of experimental*
medicine **215**, 1627-1647 (2018).
34.   Matcovitch-Natan, O. *et al.* Microglia development follows a stepwise program to regulate brain
homeostasis. *Science* **353**, aad8670 (2016).
35.   Hong, S. *et al.* Complement and microglia mediate early synapse loss in Alzheimer mouse
models. *Science* (2016).
36.   Stevens, B. *et al.* The classical complement cascade mediates CNS synapse elimination. *Cell* **131**,
1164-1178 (2007).
37.   Vainchtein, I.D. *et al.* Astrocyte-derived interleukin-33 promotes microglial synapse engulfment
and neural circuit development. *Science* **359**, 1269 (2018).
38.   Schafer, Dorothy P. *et al.* Microglia Sculpt Postnatal Neural Circuits in an Activity and
Complement-Dependent Manner. *Neuron* **74**, 691-705 (2012).
39.   Vasek, M.J. *et al.* A complement-microglial axis drives synapse loss during virus-induced memory
impairment. *Nature* **534**, 538-543 (2016).
40.   Norris, G.T. *et al.* Neuronal integrity and complement control synaptic material clearance by
microglia after CNS injury. *The Journal of experimental medicine* (2018).
41.   Krukowski, K. *et al.* Traumatic Brain Injury in Aged Mice Induces Chronic Microglia Activation,
Synapse Loss, and Complement-Dependent Memory Deficits. *International Journal of Molecular*
Sciences **19** (2018).
42.   Shi, Q. *et al.* Complement C3-Deficient Mice Fail to Display Age-Related Hippocampal Decline.
The Journal of neuroscience : the official journal of the Society for Neuroscience **35**, 13029-13042
(2015).
43.   Tay, T.L. *et al.* A new fate mapping system reveals context-dependent random or clonal
expansion of microglia. *Nature Neuroscience* **20**, 793 (2017).
44.   Huang, Y. *et al.* Repopulated microglia are solely derived from the proliferation of residual
microglia after acute depletion. *Nat Neurosci* **21**, 530-540 (2018).
45.   Ajami, B., Bennett, J.L., Krieger, C., McNagny, K.M. & Rossi, F.M.V. Infiltrating monocytes trigger
EAE progression, but do not contribute to the resident microglia pool. *Nat Neurosci* **14**, 1142-
1149 (2011).
46.   Yamasaki, R. *et al.* Differential roles of microglia and monocytes in the inflamed central nervous
system. *The Journal of experimental medicine* **211**, 1533-1549 (2014).

47.   Chen, H.-R. *et al.* Fate mapping via CCR2-CreER mice reveals monocyte-to-microglia transition in
676   development and neonatal stroke. *Science Advances* **6**, eabb2119 (2020).

677   48.   Chou, A. *et al.* Inhibition of the integrated stress response reverses cognitive deficits after
678   traumatic brain injury. *Proceedings of the National Academy of Sciences* **114**, E6420 (2017).

679   49.   Krukowski, K. *et al.* Integrated Stress Response Inhibitor Reverses Sex-Dependent Behavioral and
680   Cell-Specific Deficits after Mild Repetitive Head Trauma. *Journal of Neurotrauma* **37**, 1370-1380
681   (2019).

682   50.   Mi, H., Muruganujan, A., Ebert, D., Huang, X. & Thomas, P.D. PANTHER version 14: more
683   genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. *Nucleic*
684   *acids research* **47**, D419-D426 (2019).

685   51.   Ferreira, T.A. & Blackman, A.V. Neuronal morphometry directly from bitmap images. **11**, 982-
686   984 (2014).

687   52.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature methods* **9**,
688   676-682 (2012).

689   53.   Feng, X. *et al.* Loss of CX3CR1 increases accumulation of inflammatory monocytes and promotes
690   gliomagenesis. *Oncotarget; Vol 6, No 17* (2015).

Acknowledgements

We thank the Nikon Imaging Center at UCSF for assistance with confocal microscope. We thank the Laboratory for Cell Analysis at UCSF for assistance with FACS sort and analysis.

Funding: This work was funded by grants from the National Institutes of Health no. R01CA133216, R01CA213441 and R01AG056770 (S.R.).

Author contributions

X.F. conceptualized the study, designed and performed the experiments, analyzed the data and wrote the manuscript. E.F. performed experiments, analyzed the data and wrote the manuscript. M.P performed experiments, analyzed the data and revised the manuscript. D.C. assisted in the in vivo phagocytosis assay and data analysis. Z.B. helped with the in vivo phagocytosis assay imaging and data analysis. M.B. assisted in the Sholl analysis of microglia and BEMs. S.G analyzed the long-term dendritic spine counts data. S.L. provided assistance in experiments related to WBRT and BM chimeras. N.G. provided critical inputs to the study and revised the manuscript. S.R. conceptualized and supervised the study and revised the manuscript. All authors approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Figures

Figure 1: Microglia depletion and repopulation prevents long term radiation-induced memory deficits and loss of hippocampal PSD95. a experimental design and Novel Object Recognition (NOR) test result. CSF-1R inhibitor was used to deplete microglia during 3 doses of 3.3 Gy of whole-brain radiotherapy (WBRT). A 4-day NOR protocol was used to measure recognition memory, which ended on day 32 post WBRT. Microglia were isolated using fluorescent activated cell sorting (FACS) on day 33. and dot plots showing NOR results. Statistical analysis was performed using two-way ANOVA with Dunnett's multiple comparisons test. There is no CSF-1Ri treatment effect ($F_{(1,38)}$=1.787, p=0.1893), but significant WBRT effect ($F_{(1,38)}$=13.23, p=0.0008) and interaction between CSF-1Ri treatment and WBRT ($F_{(1,38)}$=6.07, p=0.0184), N = 9-12, animals with insufficient exploration time on NOR test day were excluded. Hollow squares represent animals used in RNA sequencing. **b** hierarchically clustered heatmap showing significantly altered microglial genes by WBRT, but not changed with CSF-1Ri treatment. **c** bar graphs summarizing fold enrichment and p values of the top 20 enriched Biological Processes by Gene Ontology analysis from up-regulated microglial genes after WBRT (full list in

 Supplementary Table1). No significantly enriched terms were identified by GO analysis from

 down-regulated genes by WBRT. **d** a pie chart summarizing all enriched GOBP terms. ns= not

 significant, ***p<0.0001. **e** scatter plots showing gating strategy in flowsynaptometry analyses.

 Fluorescent beads at various sizes were used as standard to gate isolated hippocampal cell

 membrane fractions. Particles between 1 μm and 3 μm were considered synaptosomes and used

 to determine Synapsin1 and PSD95 protein levels by mean fluorescent intensities (MFIs). **f** dot

 plots to compare Synapsin1 and PSD95 MFI levels in hippocampal cell fractions. Statistical

 analyses were performed using two-way ANOVA with Tukey's multiple comparisons test. ns = not

 significant, *p<0.05, ***p<0.001. N=6.

Figure 2: Repopulated microglia-like cells after depletion and WBRT originate from peripheral monocytes and retain monocytic signatures. a experimental design of head-protected bone marrow transplantation (BMT) followed by CSF-1Ri-mediated microglia depletion and WBRT. Lower panel shows fur colors before euthanasian for brain analysis. **b** representative FACS analysis gating strategy to analyze bone marrow chimera efficiency 6 weeks after BMT, about two thirds of the CD11b$^+$Ly6C^{high} monocytes are replaced by GFP$^+$RFP$^+$ cells derived from donor bone marrow cells. **c** representative FACS analysis gating strategy and brain myeloid composition results. Upper panel shows FACS gating using CD45 and CD11b staining; microglia and microglia-like cells are defined by positive CD11b staining and low or intermediate CD45 levels. Lower panel shows scatter plots of GFP/RFP fluorescent levels of the CD11b$^+$CD45$^{low/intermediate}$ population in the brain, and a dot plot comparing percentages of peripheral myeloid cell derived microglia-like cells. Statistical analysis was performed using unpaired t-test, ***p<0.001. **d** hierarchically clustered heatmaps to compare microglia and monocyte signatures. A signature gene list was defined using a dataset published by Lavin and Winter et al, GSE63340. Defined list and expression details are in Supplementary Table 2). **e** Similarity matrix comparisons using defined monocyte and microglia signature genes. **f** bar graph showing similarity scores to compare relative numbers of genes (in percentage of the defined list) that express in the same trends as monocytes or microglia based on the Lavin and Winter *et al* dataset.

Figure 3: Repopulated microglia and brain-engrafted macrophages are not activated and phagocyte less synaptic compartments. a experimental design for in vivo synaptosome phagocytosis assays. Injection of pre-stained synaptosomes was timed to be the same as previous experiments. Three days later, on day 36 after WBRT, ipsilateral hemispheres were harvested and used for engulfment measurement using FACS or Immunofluorescent staining. **b** FACS analysis result showing levels of microglia that engulfed pre-stained PSD-95 signals. **c** representative images showing engulfment of pre-stained synaptosomes by microglia near injection site. White arrows point at microglia that have engulfed pre-stained synaptosomes. scale bar = 20 μm. **d** dot plot to show quantification result of synaptosome engulfment by

immunofluorescent staining. **e – g** dot plots showing cell surface C5aR, and intracellular CD68

and CD107a protein levels in microglia and BEMs. Statistical analyses were performed using two-

way ANOVA with Tukey's multiple comparisons test. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

N = 5 – 6.

Figure 4: BEMs gradually adapt to microglia-like morphology and persist in the brain. a,

schematic of experimental design for long-term assessment of BEMs. **b,** representative images

of microglia/BEMs counting, scale bar = 20 μm. **c,** Sholl analysis results showing numbers of

intersections at different distances to cell center, BEMs at 7, 14, 33, 90 and 180 days after WBRT

were compared to naïve microglia age-matched to 90 days after WBRT., representative images

showing differential Iba1 and GFP expressing bprofiles of microglia (Iba1+ GFP-) and BEMs

(Iba1+ GFP+) in a BEM bearing brain at 33 days after WBRT. **e**, dot plot to show percentage of

replacement of microglia by BEMs, each dot represent an individual mouse. n = 2 - 3. Statistical

analyses were performed using unpaired t-test at each distance point (c) or time point (e). See

Supplementary Figure 7 for detailed comparisons between microglia and BEMs at each time point.

Figure 5: BEMs provide long-term protection against WBRT-induced dendritic spine and

memory loss. a schematic of experimental design for long-term memory and dendritic spine

density analyses. **b** and **c** dot plots to show NOR test results at 3 and 6 months after WBRT,

respectively. N = 6–12. **d** dendritic spine counts of hippocampal granule neurons at 1 month after

WBRT (figure reproduced using our previously published data Feng et al[53]). **e** dendritic spine

counts of hippocampal granule neurons at 6 months after WBRT, N = 5 – 6.. Statistical analyses

were performed using unpaired t-test for each time point (**c**) or two-way ANOVA with Tukey's

multiple comparisons test (**e - h**). *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 6: BEMs protects against concussive injury-induced memory deficits. a, schematic

of experimental design for concussive injury, cognitive test and following analyses. b, dot plot to

show the result of in vivo phagocytosis assay by FACS after injection of pre-stained

synaptosomes, each dot represents value from an individual mouse, n = 4 -5. c, dot plot showing

result of in vivo phagocytosis assay by IF imaging and quantification, each dot represents mean

counts from an individual mouse, n = 3. d, representative images showing microglia and BEMs

(arrows) phagocyting injected synaptosomes (green dots). e, Sholl analysis result showing

numbers of intersections at different distances to the cell center, n = 5 - 6. f, dot plot showing NOR

test result, each dot represent the performance of an individual mouse, n = 12. Statistical analyses

801   were performed using two-way ANOVA with Tukey's multiple comparisons test (b and c) for each

802   distance point (e) or unpaired t-test (f). *$p<0.05$, **$p<0.01$, ***$p<0.001$.