Disentangling Sources of Gene Tree Discordance in Phylogenomic Datasets: Testing Ancient Hybridizations in Amaranthaceae s.l.

Diego F. Morales-Briones[1*], Gudrun Kadereit[2], Delphine T. Tefarikis[2], Michael J. Moore[3], Stephen A. Smith[4], Samuel F. Brockington[5], Alfonso Timoneda[5], Won C. Yim[6], John C. Cushman[6], Ya Yang[1*]

[1] Department of Plant and Microbial Biology, University of Minnesota-Twin Cities, 1445 Gortner Avenue, St. Paul, MN 55108, USA

[2] Institut für Molekulare Physiologie, Johannes Gutenberg-Universität Mainz, D-55099, Mainz, Germany

[3] Department of Biology, Oberlin College, Science Center K111, 119 Woodland Street, Oberlin, OH 44074-1097, USA

[4] Department of Ecology & Evolutionary Biology, University of Michigan, 830 North University Avenue, Ann Arbor, MI 48109-1048, USA

[5] Department of Plant Sciences, University of Cambridge, Tennis Court Road, Cambridge, CB2 3EA, United Kingdom

[6] Department of Biochemistry and Molecular Biology, University of Nevada, Reno, NV, 89577, USA

[*] Correspondence to be sent to: Diego F. Morales-Briones and Ya Yang. Department of Plant and Microbial Biology, University of Minnesota, 1445 Gortner Avenue, St. Paul, MN 55108, USA, Telephone: +1 612-625-6292 (YY) Email: dfmoralesb@gmail.com; yangya@umn.edu

Abstract. — Gene tree discordance in large genomic datasets can be caused by evolutionary

processes such as incomplete lineage sorting and hybridization, as well as model violation, and

errors in data processing, orthology inference, and gene tree estimation. Species tree methods

that identify and accommodate all sources of conflict are not available, but a combination of

multiple approaches can help tease apart alternative sources of conflict. Here, using a

phylotranscriptomic analysis in combination with reference genomes, we test a hypothesis of

ancient hybridization within the plant family Amaranthaceae s.l. that was previously supported

by morphological, ecological, and Sanger-based molecular data. The dataset included seven

genomes and 88 transcriptomes, 17 generated for this study. We examined gene-tree discordance

using coalescent-based species trees and network inference, gene tree discordance analyses, site

pattern tests of introgression, topology tests, synteny analyses, and simulations. We found that a

combination of processes might have acted simultaneously and/or cumulatively to generate the

high levels of gene tree discordance in the backbone of Amaranthaceae s.l. Furthermore,

uninformative genes and model misspecification also contributed to discordance. No single

source or evolutionary process can account for the strong signal of gene tree discordance,

suggesting that the backbone of Amaranthaceae s.l. might be a product of an ancient and rapid

lineage diversification, and remains —and probably will remain— unresolved. This work

highlights the potential problems of identifiability associated with the sources of gene tree

discordance including, in particular, phylogenetic network methods. Our results also demonstrate

the importance of thoroughly testing for multiple sources of conflict in phylogenomic analyses,

especially in the context of ancient, rapid radiations. We provide several recommendations for

exploring conflicting signals in such situations.

46   **Keywords:** Amaranthaceae; gene tree discordance; hybridization; incomplete lineage sorting;

47   phylogenomics; transcriptomics; species tree; species network.

The exploration of gene tree discordance has become common in the phylogenetic era (Salichos

et al. 2014; Smith et al. 2015; Huang et al. 2016; Pease et al. 2018) and is essential for

understanding the underlying processes that shape the Tree of Life. Discordance among gene

trees can be the product of multiple sources. These include errors and noise in data assembly and

filtering, hidden paralogy, incomplete lineage sorting (ILS), gene duplication/loss (Pamilo and

Nei 1988; Doyle 1992; Maddison 1997; Galtier and Daubin 2008), random noise from

uninformative genes, as well as misspecified model parameters of molecular evolution such as

substitutional saturation, codon usage bias, or compositional heterogeneity (Foster 2004; Cooper

2014; Cox et al. 2014; Liu et al. 2014). Among these potential sources of gene tree discordance,

ILS is the most studied in the systematics literature (Edwards 2009), and several phylogenetic

inference methods have been developed to accommodate ILS as the source of discordance

(reviewed in Edwards et al. 2016; Mirarab et al. 2016; Xu and Yang 2016). More recently,

methods that account for additional processes such as hybridization or introgression have gained

attention. These include methods that estimate phylogenetic networks while accounting for ILS

and hybridization simultaneously (e.g., Solís-Lemus and Ané 2016; Wen et al. 2018), and

methods that detect introgression based on site patterns or phylogenetic invariants (e.g., Green et

al. 2010; Durand et al. 2011; Kubatko and Chifman 2019). Frequently, multiple of these sources

of gene tree discordance can contribute to gene tree heterogeneity (Holder et al. 2001; Buckley et

al. 2006; Meyer et al. 2017; Knowles et al. 2018). However, at present, no method can estimate

species trees from phylogenomic data while modeling multiple sources of conflict and molecular

substitution simultaneously. To overcome these limitations, the use of multiple phylogenetic

tools and data partitioning schemes in phylogenomic datasets is essential to disentangle sources

of gene tree heterogeneity and resolve recalcitrant relationships at deep and shallow nodes of the

Tree of Life (e.g., Alda et al. 2019; Roycroft et al. 2019; Widhelm et al. 2019; Prasanna et al.

2020).

In this study, we explore multiple sources of gene tree conflict to test controversial

hypotheses of ancient hybridization among subfamilies in the plant family Amaranthaceae s.l.

Amaranthaceae s.l. includes the previously segregated family Chenopodiaceae (Hernández-

Ledesma et al. 2015; The Angiosperm Phylogeny Group 2016). With ca. 2050 to 2500 species in

181 genera and a worldwide distribution (Hernández-Ledesma et al. 2015), Amaranthaceae s.l. is

iconic for the repeated evolution of complex traits representing adaptations to extreme

environments such as C_4 photosynthesis in hot and often dry environments (e.g., Kadereit et al.

2012; Bena et al. 2017), various modes of extreme salt tolerance (e.g., Flowers and Colmer 2015;

Piirainen et al. 2017) that in several species are coupled with heavy metal tolerance (Moray et al.

2016), and very fast seed germination and production of multiple diaspore types on one

individual (Kadereit et al. 2017). Several important crops are members of Amaranthaceae s.l.,

such as the pseudocereals quinoa and amaranth, sugar beet, spinach, glassworts, and *Salsola*

soda. Many species of the family are also important fodder plants in arid regions and several are

currently being investigated for their soil ameliorating and desalinating effects. Due to their

economic importance, reference genomes are available for *Beta vulgaris* (sugar beet, subfamily

Betoideae; Dohm et al. 2014), *Chenopodium quinoa* (quinoa, Chenopodioideae; Jarvis et al.

2017), *Spinacia oleracea* (spinach; Chenopodioideae; Xu et al. 2017) and *Amaranthus*

hypochondriacus (amaranth; Amaranthoideae; Lightfoot et al. 2017), representing three of the 13

currently recognized subfamilies of Amaranthaceae s.l. (sensu Kadereit et al. 2003; Kadereit et

al. 2017).

Within the core Caryophyllales the previously segregated families Amaranthaceae s.s.

and Chenopodiaceae have always been regarded as closely related, and their separate family

status has long been the subject of phylogenetic and taxonomic debate (Kadereit et al. 2003;

Masson and Kadereit 2013; Hernández-Ledesma et al. 2015; Walker et al. 2018; Fig. 1). Their

close affinity is supported by a number of shared morphological, anatomical and phytochemical

synapomorphies, and has been substantiated by molecular phylogenetic studies (discussed in

Kadereit et al. 2003). Amaranthaceae s.s. has a predominantly tropical and subtropical

distribution with the highest diversity found in the Neotropics, eastern and southern Africa and

Australia (Müller and Borsch 2005), while Chenopodiaceae predominantly occurs in temperate

regions and semi-arid or arid environments of subtropical regions (Kadereit et al. 2003). The key

problem has always been the species-poor and heterogeneous subfamilies Polycnemoideae and

Betoideae, neither of which fit easily within Chenopodiaceae or Amaranthaceae s.s. (cf. Table 5

in Kadereit et al. 2003). Polycnemoideae is similar in ecology and distribution to

Chenopodiaceae but shares important floral traits such as petaloid tepals, filament tubes and 2-

locular anthers with Amaranthaceae s.s. Morphologically, Betoideae fits into either

Chenopodiaceae or Amaranthaceae s.s. but has a unique fruit type—a capsule that opens with a

circumscissile lid (Kadereit et al. 2006). Both Betoideae and Polycnemoideae possess only a few

species each and each has a strongly disjunct distribution patterns across three continents.

Furthermore, the genera of both subfamilies display a number of morphologically dissociating

features. Both intercontinental disjunctions of species-poor genera and unique morphological

traits led to the hypothesis that Betoideae and Polycnemoideae might have originated from

hybridization events among early-branching lineages in Amaranthaceae s.l. (Hohmann et al.

2006; Masson and Kadereit 2013). To test this hypothesis, a phylotranscriptomic approach is

particularly compelling as it not only provides thousands of low-copy nuclear genes for

dissecting sources of phylogenetic discordance, but also enables future studies associating gene

tree topology with gene function and habitat adaptation.

Previous molecular phylogenetic analyses struggled to resolve the relationships among

Betoideae, Polycnemoideae and the rest of the Amaranthaceae s.l. (Fig. 1). The first

phylogenomic study of Amaranthaceae s.l. using nuclear loci (Walker et al. 2018; Fig. 1e)

revealed that gene tree discordance mainly occurred at deep nodes of the phylogeny involving

Betoideae. Polycnemoideae was sister to Chenopodiaceae, albeit supported by only 17% of gene

trees, which contradicted previous analyses based on plastid data (Fig. 1, a–d). However, only a

single species of Betoideae (the cultivated beet and its wild relative) was sampled in Walker et

al. (2018) and sources of conflicting signals among gene trees remained unexplored.

In this study, we used a large genomic dataset to examine sources of gene tree

discordance in Amaranthaceae s.l. Specifically, we tested whether Polycnemoideae and

Betoideae result from independent hybridizations between Amaranthaceae s.s. and

Chenopodioideae by distinguishing the signal of hybridization from gene tree discordance

produced by ILS, uninformative gene trees, hidden paralogy, misspecifications of model of

molecular evolution, and hard polytomy. We show that phylogenetic networks may not be

identifiable, even with large and well-sampled genomic datasets.

FIGURE 1. Phylogenetic hypothesis of Amaranthaceae s.l. from previous studies. a) Kadereit et al. (2003) using the plastid (cpDNA) *rbcL* coding region. b) Müller and Borsch (2005); using the cpDNA *matK* coding region and partial *trnL* intron. c) Hohmann et al. (2006) using the cpDNA *ndhF* coding region. d) Kadereit et al. (2017) using the cpDNA *atpB-rbcL* spacer, *matK* with *trnL* intron, *ndhF-rpL32* spacer, and *rps16* intron e) Walker et al. (2018) using 455 nuclear genes from transcriptome data. Major clades of Amaranthaceae s.l. named following the results of this study. Image credits: *Amaranthus hypochondriacus* by Picture Partners, *Beta vulgaris* by Olha Huchek, *Chenopodium quinoa* by Diana Mower, *Nitrophila mohavensis* by James M. André, and *Salsola soda* by Homeydesign.

MATERIALS AND METHODS

An overview of all dataset and phylogenetic analyses can be found in Figure S1.

164   *Taxon sampling, transcriptome sequencing*

We sampled 92 ingroup species (88 transcriptomes and four genomes) representing 53 genera

(out of ca. 181) of all 13 currently recognized subfamilies and 16 out of 17 tribes of

Amaranthaceae s.l. (sensu [Kadereit et al. 2003; Kadereit et al. 2017]). In addition, 13 outgroups

across the Caryophyllales were included (ten transcriptomes and three genomes; Table S1). We

generated 17 new transcriptomes for this study on an Illumina HiSeq2500 platform (Table S2).

Library preparation was carried out using either poly-A enrichment or ribosomal RNA depletion.

See Supplemental Methods for details on tissue collection, RNA isolation, library preparation,

and quality control.

174   *Transcriptome data processing, assembly, homology and orthology inference*

Read processing, assembly, translation, and homology and orthology inference followed the

'phylogenomic dataset construction' pipeline (Yang and Smith 2014) with multiple updates. We

briefly describe our procedure below, with details in the Supplemental Methods and updated

scripts in https://bitbucket.org/yanglab/phylogenomic_dataset_construction/

We processed raw reads for all 88 transcriptome datasets (except *Bienertia sinuspersici*)

used in this study (Table S1). Reads were corrected for errors, trimmed for sequencing adapters

and low-quality bases, and filtered for organellar reads. *De novo* assembly of processed nuclear

reads was carried out with Trinity v 2.5.1 (Grabherr et al. 2011) with default settings, but without

in silico normalization. Low quality and chimeric transcripts were removed. Filtered transcripts

were clustered into putative genes with Corset v 1.07 (Davidson and Oshlack 2014) and only the

longest transcript of each putative gene was retained (Chen et al. 2019). Lastly, transcripts were

translated, and identical coding sequences (CDS) were removed. Homology inference was

carried out on CDS using reciprocal BLASTN, followed by orthology inference using the

'monophyletic outgroup' approach (Yang and Smith 2014), keeping only ortholog groups with at

least 25 ingroup taxa.

192   *Assessment of recombination*

Coalescent species tree methods assume that there is free recombination between loci and no

recombination within loci. To determine the presence of recombination in our dataset, we used

the pairwise homoplasy index test Φ for recombination, as implemented in PhiPack (Bruen et al.

2006). We tested recombination on the final set of ortholog alignments (with a minimum of 25

taxa) with the default sliding window size of 100 bp. Alignments that showed a strong signal of

recombination with $p \leq 0.05$ were removed from all subsequent phylogenetic analyses.

200   *Nuclear phylogenetic analysis*

We used both concatenation and coalescent-based methods to reconstruct the phylogeny of

Amaranthaceae s.l. Sequences from final orthologs were aligned using MAFFT v 7.307 (Katoh

and Standley 2013) with settings '—genafpair --maxiterate 1000'. Columns with more than 70%

missing data were trimmed with Phyx (Brown et al. 2017), and alignments with at least 1,000

characters and 99 out of 105 taxa were retained. We first estimated a maximum likelihood (ML)

tree of the concatenated matrix with RAxML v 8.2.11 (Stamatakis 2014) using a partition-by-

gene scheme with GTRCAT model for each partition and clade support assessed with 200 rapid

bootstrap (BS) replicates. To estimate a coalescent-based species tree, first we inferred individual

ML gene trees using RAxML with a GTRCAT model and 200 BS replicates to assess clade

support. Gene trees were then used to infer a species tree with ASTRAL-III v5.6.3 (Zhang et al.

2018) using local posterior probabilities (LPP; Sayyari and Mirarab 2016) to assess clade

support.

214   *Detecting and visualizing nuclear gene tree discordance*

To explore discordance among gene trees, we first calculated the internode certainty all (ICA)

value to quantify the degree of conflict on each node of a target tree (i.e., species tree) given

individual gene trees (Salichos et al. 2014). In addition, we calculated the number of conflicting

and concordant bipartitions on each node of the species trees. Both the ICA scores and

conflicting/concordant bipartitions were calculated with Phyparts (Smith et al. 2015), mapping

against the inferred ASTRAL species trees, using individual gene trees with BS support of at

least 50% for the corresponding node. Additionally, in order to distinguish strong conflict from

weakly supported branches, we carried out Quartet Sampling (QS; Pease et al. 2018) with 100

replicates. Quartet Sampling subsamples quartets from the input tree and alignment and assesses

the confidence, consistency, and informativeness of each internal branch by the relative

frequency of the three possible quartet topologies (Pease et al. 2018).

226   To further visualize conflict, we built a cloudogram using DensiTree v2.2.6 (Bouckaert

and Heled 2014). As DensiTree cannot accommodate missing taxa among gene trees, we

reduced the final ortholog alignments to include 41 species (38 ingroup and 3 outgroups) in order

to include as many orthologs as possible while representing all main clades of Amaranthaceae

s.l. (see results). Individual gene trees were inferred as previously described. Trees were time-

calibrated with TreePL v1.0 (Smith and O'Meara 2012) by fixing the crown age of

Amaranthaceae s.l. to 66–72.1 based on a pollen record of *Polyporina cribraria* from the late

Cretaceous (Maastrichtian; Srivastava 1969), and the root for the reduced 41-species dataset

(most common recent ancestor of Achatocarpaceae and Aizoaceae) was set to 95 Ma based on

the time-calibrated plastid phylogeny of Caryophyllales from Yao et al. (2019).

237   *Plastid assembly and phylogenetic analysis*

Although DNase treatment was carried out to remove genomic DNA, due to their high copy

number, plastid sequences are often carried over in RNA-Seq libraries. In addition, as young leaf

tissue was used for RNA-Seq, the presence of RNA from plastid genes is expected to be

represented. To investigate phylogenetic signal from plastid sequences, *de novo* assemblies were

carried out with the Fast-Plast v.1.2.6 pipeline (https://github.com/mrmckain/Fast-Plast) using

the filtered organelle reads. Contigs produced by Spades v 3.9.0 (Bankevich et al. 2012) were

mapped to the closest available reference plastomes (Table S3), one copy of the Inverted Repeat

was removed, and the remaining contigs manually edited in Geneious v.11.1.5 (Kearse et al.

2012) to produce the final oriented contigs.

247   Contigs were aligned with MAFFT with the setting '--auto'. Two samples (*Dysphania*

schraderiana and *Spinacia turkestanica*) were removed due to low sequence occupancy. Using

the annotations of the reference genomes (Table S3), the coding regions of 78 genes were

extracted and each gene alignment was visually inspected in Geneious to check for potential

misassemblies. From each gene alignment, taxa with short sequences (i.e., < 50% of the aligned

length) were removed and the remaining sequences realigned with MAFFT. The genes *rpl32* and

ycf2 were excluded from downstream analyses due to low taxon occupancy (Table S4). For each

individual gene we performed extended model selection (Kalyaanamoorthy et al. 2017) followed

by ML gene tree inference and 1,000 ultrafast bootstrap replicates for branch support (Hoang and

Chernomor 2018) in IQ-TREE v.1.6.1 (Nguyen et al. 2015). For the concatenated matrix we

searched for the best partition scheme (Lanfear et al. 2012) followed by ML gene tree inference

and 1,000 ultrafast bootstrap replicates for branch support in IQ-Tree. Additionally, we evaluated

branch support with QS using 1,000 replicates and gene tree discordance with PhyParts. Lastly,

to identify the origin of the plastid reads (i.e., genomic or RNA), we predicted RNA editing from

CDS alignments using PREP (Mower 2009) with the alignment mode (PREP-aln), and a cutoff

value of 0.8.

264   *Species network analysis using a reduced 11-taxon dataset*

We inferred species networks that model ILS and gene flow using a maximum pseudo-likelihood

approach (Yu and Nakhleh 2015). Species network searches were carried out with PhyloNet

v.3.6.9 (Than et al. 2008) with the command 'InferNetwork_MPL' and using the individual gene

trees as input. Due to computational restrictions, and given our main focus to identify potential

reticulating events among major clades of Amaranthaceae s.l., we reduced our taxon sampling to

one outgroup and ten ingroup taxa to include two representative species from each of the five

well-supported major lineages in Amaranthaceae s.l. (see results). We filtered the final 105-taxon

ortholog set to include genes that have all 11 taxa [referred herein as 11-taxon(net) dataset].

After alignment and trimming we kept genes with a minimum of 1,000 aligned base pairs and

individual ML gene trees were inferred using RAxML with a GTRGAMMA model and 200

bootstrap replicates. We carried out five network searches by allowing one to five reticulation

events and ten runs for each search. To estimate the optimum number of reticulations, we

optimized the branch lengths and inheritance probabilities and computed the likelihood of the

best scored network from each of the five maximum reticulation events searches. Network

likelihoods were estimated given the individual gene trees using the command 'CalGTProb' in

PhyloNet (Yu et al. 2012). Then, we performed model selection using the bias-corrected Akaike

information criterion (AICc; Sugiura 1978), and the Bayesian information criterion (BIC;

Schwarz 1978). The number of parameters was set to the number of branch lengths being

estimated plus the number of hybridization probabilities being estimated. The number of gene

trees used to estimate the likelihood was used to correct for finite sample size. To compare

network models to bifurcating trees, we also estimated bifurcating concatenated ML and

coalescent-based species trees and a plastid tree as previously described with the reduced 11-

species taxon sampling.

289   *Hypothesis testing and detecting introgression using four-taxon datasets*

Given the signal of multiple clades potentially involved in hybridization events detected by

PhyloNet (see results), we next conducted quartet analyses to explore a single event at a time.

First, we further reduced the 11-taxon(net) dataset to six taxa that included one outgroup genome

(*Mesembryanthemum crystallinum*) and one ingroup from each of the five major ingroup clades:

Amaranthus hypochondriacus (genome)*, Beta vulgaris* (genome)*, Chenopodium quinoa*

(genome)*, Caroxylon vermiculatum* (transcriptome)*,* and *Polycnemum majus* (transcriptome) to

represent Amaranthaceae s.s., Betoideae, 'Chenopods I', 'Chenopods II' and Polycnemoideae,

respectively. We carried out a total of ten quartet analyses using all ten four-taxon combinations

that included three out of five ingroup species and one outgroup. We filtered the final set of 105-

taxon orthologs for genes with all four taxa for each combination and inferred individual gene

trees as described before. For each quartet we carried out the following analyses. We first

estimated a species tree with ASTRAL and explored gene tree conflict with PhyParts. We then

explored individual gene tree resolution by calculating the Tree Certainty (TC) score (Salichos et

al. 2014) in RAxML using the majority rule consensus tree across the 200 bootstrap replicates.

Next, we explored potential correlation between TC score and alignment length, GC content and

alignment gap proportion using a linear regression model in R v.3.6.1 (R Core Team 2019).

Lastly, we tested for the fit of gene trees to the three possible rooted quartet topologies for each

gene using the approximately unbiased (AU) tests (Shimodaira 2002). We carried out ten

constraint searches for each of three topologies in RAxML with the GTRGAMMA model, then

calculated site-wise log-likelihood scores for the three constraint topologies in RAxML using

GTRGAMMA and carried out the AU test using Consel v.1.20 (Shimodaira and Hasegawa

2001). In order to detect possible introgression among species of each quartet, first we estimated

a species network with PhyloNet using a full maximum likelihood approach (Yu et al. 2014)

with 100 runs per search while optimizing the likelihood of the branch lengths and inheritance

probabilities for every proposed species network. Furthermore, we also carried out the

ABBA/BABA test to detect introgression (Green et al. 2010; Durand et al. 2011) in each of four-

taxon species trees. We calculated the D-statistic and associated z score for the null hypothesis of

no introgression ($D = 0$) following each quartet ASTRAL species tree for taxon order assignment

using 100 jackknife replicates and a block size of 10,000 bp with evobiR v1.2 (Blackmon and

Adams) in R.

Additionally, to detect any non-random genomic block of particular quartet topology

(Fontaine et al. 2015), we mapped the physical location of genes supporting each alternative

quartet topology onto the *Beta vulgaris* reference genome (See Supplemental Information for

details).

Assessment of substitutional saturation, codon usage bias, compositional heterogeneity, and

325   *model of sequence evolution misspecification*

Analyses were carried out in a 11-taxon dataset [referred herein as 11-taxon(tree)] that included

the same taxa used for species network analyses, but was processed differently to account for

codon structure (see Supplemental Methods for details). Saturation was evaluated by plotting the

uncorrected genetic distances of the concatenated alignment against the inferred distances (see

Supplemental Methods for details). To determine the effect of saturation in the phylogenetic

inferences we estimated individual ML gene trees using an unpartitioned alignment, a partition

by first and second codon positions, and the third codon positions, and by removing all third

codon positions. All tree searches were carried out in RAxML with a GTRGAMMA model and

200 bootstrap replicates. We then estimated a coalescent-based species trees and explored gene

tree discordance with PhyParts.

336   Codon usage bias was evaluated using a correspondence analysis of the Relative

337   Synonymous Codon Usage (RSCU; see Supplemental Methods for details). To determine the

338   effect of codon usage bias in the phylogenetic inferences we estimated individual gene trees

339   using codon-degenerated alignments (see Supplemental Methods for details). Gene tree inference

340   and discordance analyses were carried out on the same three data schemes as previously

341   described.

342   Among-lineage compositional heterogeneity was evaluated on individual genes using a

compositional homogeneity test (Supplemental Methods for details). To assess if compositional

heterogeneity had an effect in species tree inference and gene tree discordance, gene trees that

showed the signal of compositional heterogeneity were removed from saturation and codon

usage analyses and the species tree and discordance analyses were rerun.

To explore the effect of sequence evolution model misspecification, we reanalyzed the

datasets from the saturation and codon usage analyses using inferred gene trees that accounted

for model selection. Additionally, we also explored saturation and model misspecification in

phylogenetic trees from amino acid alignments (see Supplemental Methods for details).

352   *Polytomy test*

To test if the gene tree discordance among the main clades of Amaranthaceae s.l. could be

explained by polytomies instead of bifurcating nodes, we carried out the quartet-based polytomy

test by Sayyari and Mirarab (2018) as implemented in ASTRAL. We performed the polytomy

test using the gene trees inferred from the saturation and codon usage analyses [11-taxon(tree)

dataset]. Because this test can be sensitive to gene tree error (Syyari and Mirarab 2018), we

performed a second test using gene trees where branches with less than 75% of bootstrap support

were collapsed.

361   *Coalescent simulations*

To investigate if gene tree discordance can be explained by ILS alone, we carried out coalescent

simulations similar to Cloutier et al. (2019). An ultrametric species tree with branch lengths in

mutational units (μT) was estimated by constraining an ML tree search of the 11-taxon(net)

concatenated alignment to the ASTRAL species tree topology with a GTR+GAMMA model

while enforcing a strict molecular clock in PAUP v4.0a (build 165; Swofford 2002). The

mutational branch lengths from the constrained tree and branch lengths in coalescent units ($\tau=$

$T/4N_e$) from the ASTRAL species trees were used to estimate the population size parameter theta

($\Theta= \mu T/\tau$; Degnan and Rosenberg 2009) for internal branches. Terminal branches were set with a

population size parameter theta of one. We used the R package Phybase v. 1.4 (Liu and Yu 2010)

that uses the formula from Rannala and Yang (2003) to simulate 10,000 gene trees using the

constraint tree and the estimated theta values. Then we calculated the distribution of Robinson

and Foulds (1981) tree-to-tree distances between the species tree and each gene tree using the R

package Phangorn v2.5.3 (Schliep 2011), and compared this with the distribution of tree-to-tree

distances between the species tree and the simulated gene tree. We ran simulations using the

species tree and associated gene tree distribution from the original no partition 11-taxon(net).

378   *Test of the anomaly zone*

The anomaly zone occurs where a set of short internal branches in the species tree produces gene

trees that differ from the species tree more frequently than those that are concordant [a(x); as

defined in equation 4 of Degnan and Rosenberg (2006)]. To explore if gene tree discordance

observed in Amaranthaceae s.l. is a product of the anomaly zone, we estimated the boundaries of

the anomaly zone [a(x); as defined in equation 4 of Degnan and Rosenberg (2006)] for the

internal nodes of the species tree. Here, x is the branch length in coalescent units in the species

tree that has a descendant internal branch. If the length of the descendant internal branch (y) is

smaller than a(x), then the internode pair is in the anomaly zone and is likely to produce anomaly

gene trees (AGTs). We carried out the calculation of a(x) following Linkem et al. (2016) in the

same 11-taxon(tree) ASTRAL species tree used for coalescent simulations. Additionally, to

establish the frequency of gene trees that were concordant with the estimated species trees, we

quantified the frequency of all 105 possible rooted gene trees with Amaranthaceae s.l. being

monophyletic.

393   **RESULTS**

394   *Transcriptome sequencing, assembly, translation, and quality control*

Raw reads for the 17 newly generated transcriptomes are available from the NCBI Sequence

Read Archive (BioProject: XXXX; Table S2). The number of raw read pairs ranged from 17 to

27 million. For the 16 samples processed using RiboZero, organelle reads accounted for 15% to

52% of read pairs (Table S2). For *Tidestromia oblongifolia* that poly-A enrichment was carried

out in library prep with ~5% of raw reads were from organelle (Table S2). The final number of

orthologs was 13,024 with a mean of 9,813 orthologs per species (Table S1). Of those, 82

orthologs had a strong signal of recombination ($P \leq 0.05$) and were removed from downstream

analyses.

404   *Analysis of the nuclear dataset of Amaranthaceae s.l.*

The final set of nuclear orthologous genes included 936 genes with at least 99 out of 105 taxa

and 1,000 bp in aligned length after removal of low occupancy columns (the 105-taxon dataset).

The concatenated matrix consisted of 1,712,054 columns with a gene and character occupancy of

96% and 82%, respectively. The species tree from ASTRAL and the concatenated ML tree from

RAxML recovered the exact same topology with most clades having maximal support [i.e.,

bootstrap percentage (BS) = 100, local posterior probabilities (LPP) = 1; Fig. 2; Figs S2–S3].

Both analyses recovered Chenopodiaceae as monophyletic with the relationships among major

clades concordant with the cpDNA analysis from Kadereit et al. (2017; Fig. 1d). Betoideae was

placed as sister of Chenopodiaceae, while Polycnemoideae was strongly supported as sister (BS

= 97, LPP = 0.98) to the clade composed of Chenopodiaceae and Betoideae. Amaranthaceae s.s.

had an overall topology concordant to Kadereit et al. (2017), with the exception of *Iresine,* which

was recovered among the Aervoids (Fig. 2; Figs S2–S3).

FIGURE 2. Maximum likelihood phylogeny of Amaranthaceae s.l. inferred from RAxML

analysis of the concatenated 936-nuclear gene supermatrix, which had the same topology as

recovered from ASTRAL. All nodes have maximal support (bootstrap = 100/ASTRAL local

posterior probability = 1) unless noted. Pie charts present the proportion of gene trees that

support that clade (blue), support the main alternative bifurcation (green), support the remaining

alternatives (red), and the proportion (conflict or support) that have < 50% bootstrap support

(gray). Only pie charts for major clades are shown (see Fig. S2 for all node pie charts). Branch

lengths are in number of substitutions per site. The inset (top left) shows the Densitree

cloudogram inferred from 1,242 nuclear genes for the reduced 41-taxon dataset.

The conflict analyses confirmed the monophyly of Amaranthaceae s.l. with 922 out of

930 informative gene trees being concordant (ICA= 0.94) and having full QS support (1/–/1; i.e.,

all sampled quartets supported that branch). Similarly, the monophyly of Amaranthaceae s.s. was

highly supported by 755 of 809 informative gene trees (ICA =0.85) and the QS scores (0.92/0/1).

The backbone of the family, however, was characterized by high levels of gene tree discordance

(Fig. 2; Figs S2–S3). The monophyly of Chenopodiaceae was supported only by 231 out of 632

informative gene trees (ICA = 0.42) and the QS score (0.25/0.19/0.99) suggested weak quartet

support with a skewed frequency for an alternative placement of two well-defined clades within

Chenopodiaceae, herein referred to as 'Chenopods I' and 'Chenopods II' (Fig. 2; Figs S2–S3).

'Chenopods I' and 'Chenopods II' were each supported by the majority of gene trees, 870 (ICA

= 0.89) and 916 (ICA = 0.91), respectively and full QS support. Similarly, high levels of conflict

among informative gene trees were detected in the placement of Betoideae (126 out of 579

informative genes being concordant, ICA = 0.28; QS score 0.31/0.57/1) and Polycnemoideae

(116/511; ICA = 0.29;0.3/0.81/0.99). The Densitree cloudogram also showed significant conflict

along the backbone of Amaranthaceae s.l. (Fig. 2).

Together, analysis of nuclear genes recovered five well-supported clades in

Amaranthaceae s.l.: Amaranthaceae s.s., Betoideae, 'Chenopods I', 'Chenopods II', and

Polycnemoideae. However, relationships among these five clades showed a high level of conflict

among genes [ICA scores and gene counts (pie charts)] and among subsampled quartets (QS

scores), despite having high support from both BS and LPP scores.

450   *Plastid phylogenetic analysis of Amaranthaceae s.l.*

The final alignment from 76 genes included 103 taxa and 55,517 bp in aligned length. The ML

tree recovered the same five main clades within Amaranthaceae s.l. with maximal support (BS =

100; Figs S4–S6). Within each main clade, relationships were fully congruent with Kadereit et

al. (2017) and mostly congruent with our nuclear analyses. However, the relationship among the

five main clades differed from the nuclear tree. Here, the sister relationships between Betoideae

and 'Chenopods I', and between Amaranthaceae s.s. and Polycnemoideae were both supported by

BS =100. The sister relationship between these two larger clades was moderately supported (BS

= 73), leaving 'Chenopods II' as sister to the rest of Amaranthaceae s.l.

459   Conflict analysis confirmed the monophyly of Amaranthaceae s.l. with 51 out of 69

informative gene trees supporting this clade (ICA = 0.29) and full QS support (1/–/1). On the

other hand, and similar to the nuclear phylogeny, significant gene tree discordance was detected

among plastid genes regarding placement of the five major clades (Figs S4–S6). The sister

relationship of Betoideae and 'Chenopods I' was supported by only 20 gene trees (ICA = 0.06),

but it had a strong support from QS (0.84/0.88/0/94). The relationship between Amaranthaceae

s.s. and Polycnemoideae was supported by only 15 gene trees (ICA = 0.07), while QS showed

weak support (0.41/0.21.0.78) with signals of a supported secondary evolutionary history. The

clade uniting Betoideae, 'Chenopods I', Amaranthaceae s.s., and Polycnemoideae was supported

by only four-gene trees, with counter-support from both QS (-0.29/0.42/0.75) and ICA (-0.03),

suggesting that most gene trees and sampled quartets supported alternative topologies.

RNA editing prediction analysis revealed editing sites only on CDS sequences of

reference plastomes (Table S3), suggesting that cpDNA reads in RNA-seq libraries come from

RNA rather than DNA leftover from incomplete DNase digestion during sample processing (See

Discussion for details in plastid assembly from RNA-seq data).

475   *Species network analysis of Amaranthaceae s.l.*

The reduced 11-taxon(net) dataset included 4,138 orthologous gene alignments with no missing

taxon and a minimum of 1,000 bp (aligned length after removal of low occupancy columns). The

11-taxon(net) ASTRAL species tree was congruent with the 105-taxon tree, while both the

nuclear and plastid ML trees from concatenated supermatrices had different topologies than their

corresponding 105-taxon trees (Fig. 3). PhyloNet identified up to five hybridization events

among the clades of Amaranthaceae s.l. (Fig. 3), with the best model having five hybridization

events involving all five clades (Table S5). 'Chenopods II' was involved in hybridization events

in all networks with one to five hybridization events. Model selection indicated that any species

network was a better model than the bifurcating nuclear or plastid trees (Table S5).

FIGURE 3. Species trees and species networks of the reduced 11-taxon(net) dataset of

Amaranthaceae s.l. Nuclear concatenated phylogeny inferred from 4,138-nuclear gene

supermatrix with RAxML. ASTRAL species tree inferred using 4,138 nuclear genes. cpDNA

concatenated tree inferred from 76-plastid gene supermatrix with IQ-tree. Best species network

inferred from PhyloNet pseudolikelihood analyses with 1 to 5 maximum number of reticulations.

Red and blue indicate the minor and major edges, respectively, of hybrid nodes. Number next to

the branches indicates inheritance probabilities for each hybrid node.

495   *Four-taxon analyses*

To test for hybridization events one at a time, we further reduced the 11-taxon(net) dataset to 10

four-taxon combinations that each included one outgroup and one representative each from three

out of the five major ingroup clades. Between 7,756 and 8,793 genes were used for each quartet

analysis (Table S6) and each quartet topology can be found in Figure 4. Only five out of the ten

bifurcating quartet species trees (H0 and more frequent gene tree) were compatible with the

nuclear species tree inferred from the complete 105-taxon dataset. The remaining quartets

corresponded to the second most frequent gene tree topology in the 105-taxon nuclear tree,

except for the quartet of Betoideae, 'Chenopods II' and Polycnemoideae (PBC2, which

correspond to the least frequent gene tree).

FIGURE 4. Gene counts from Approximate-Unbiased (AU) topology test of the 10 quartets from

the five main clades of Amaranthaceae s.l. AU tests were carried out between the three possible

topologies of each quartet. H0 represents the ASTRAL species tree of each quartet. "Equivocal"

indicates gene trees that fail to reject all three alternative topologies for a quartet with $p \leq 0.05$.

Gene counts for each of the three alternative topologies represent gene trees supporting

unequivocally one topology by rejecting the other two alternatives with $p \leq 0.05$. Insets represent

gene counts only for unequivocal topology support. Double arrowed lines in each H0 quartet

represent the direction of introgression from the ABBA/BABA test. Each quartet is named

following the species tree topology, where the first two species are sister to each other. A =

Amaranthaceae s.s. (represented by *Amaranthus hypochondriacus*), B = Betoideae (*Beta*

vulgaris), C1 = Chenopods I (*Chenopodium quinoa*), C2 = Chenopods II (*Caroxylum*

vermiculatum), P = Polycnemoideae (*Polycnemum majus*). All quartets are rooted with

Mesembryanthemum crystallinum.

In each of the ten quartets, the ASTRAL species tree topology (H0) was the most

frequent among individual gene trees (raw counts) but only accounted for 35%–41% of gene

trees, with the other two alternative topologies having balanced to slightly skewed frequencies

(Fig. S7a; Table S7). Gene counts based on the raw likelihood scores from the constraint

analyses showed similar patterns (Fig. S7b; Table S7). When filtered by significant likelihood

support (i.e., $\Delta AICc \geq 2$), the number of trees supporting each of the three possible topologies

dropped between 34% and 45%, but the species tree remained the most frequent topology for all

quartets (Fig. S7b; Table S7). The AU topology tests failed to reject ($P \leq 0.05$) approximately

85% of the gene trees for any of the three possible quartet topologies and rejected all but a single

topology in only 3%–4.5% of cases. Among the unequivocally selected gene trees, the

frequencies among the three alternative topologies were similar to ones based on raw likelihood

scores (Fig S7; Table S7). Therefore, topology tests showed that most genes were uninformative

for resolving the relationships among the major groups of Amaranthaceae s.l.

Across all ten quartets we found that most genes had very low TC scores (for any single

node the maximum TC value is 1; Supplemental Fig. S8), showing that individual gene trees also

had high levels of conflict among bootstrap replicates, which also indicated uninformative genes

and was concordant with the AU topology test results. We were unable to detect any significant

correlation between TC scores and alignment length, GC content or alignment gap fraction

(Table S8), suggesting that filtering genes by any of these criteria is unlikely to increase the

information content of the dataset.

Species network analyses followed by model selection using each of the four-taxon

datasets showed that in seven out of the ten total quartets, the network with one hybridization

event was a better model than any bifurcating tree topology. However, each of the best three

networks from PhyloNet had very close likelihood scores and no significant ΔAICc among them

(Table S6; Fig S9). For the remaining three quartets, the species trees (H0) was the best model.

The ABBA/BABA test results showed a significant signal of introgression within each of

the ten quartets (Table S9; Fig 4). The possible introgression was detected between six out of the

ten possible pairs of taxa. Potential introgression between Betoideae and Amaranthaceae s.s.,

'Chenopods I' or 'Chenopods II', and between 'Chenopods I' and Polycnemoideae was not

detected.

To further evaluate whether alternative quartets were randomly distributed across the

genome, we mapped quartet topologies of the BC1A quartet onto the reference genome of *Beta*

vulgaris. We used the BC1A quartet as an example as all four species in this quartet have

reference genomes. Synteny analysis between the diploid ingroup reference genome *Beta*

vulgaris and the diploid outgroup reference genome *Mesembryanthemum crystallinum* recovered

22,179 collinear genes in 516 syntenic blocks. With the collinear ortholog pair information, we

found that of the 8,258 orthologs of the BC1A quartet, 6,941 contained syntenic orthologous

genes within 383 syntenic blocks. The distribution of the BC1A quartet topologies along the

chromosomes of *Beta vulgaris* did not reveal any spatial clustering of any particular topology

along the chromosomes (Fig. S10).

Assessment of substitutional saturation, codon usage bias, compositional heterogeneity,

563   *and sequence evolution model misspecification*

We assembled a second 11-taxon(tree) dataset that included 5,936 genes and a minimum of 300

bp (aligned length after removal of low occupancy columns) and no missing taxon. The

saturation plots of uncorrected and predicted genetic distances showed that the first and second

codon positions were unsaturated (y = 0.884x), while the slope of the third codon positions (y =

0.571x) showed a signal of saturation (Fig. S11). The correspondence analyses of RSCU show

that some codons are more frequently used in different species, but overall the codon usage was

randomly dispersed among all species and not clustered by clade (Fig. S12). This suggests that

the phylogenetic signal is unlikely to be driven by differences in codon usage bias among clades.

Furthermore, only 549 (~9%) genes showed a signal of compositional heterogeneity (p < 0.05).

The topology and support (LPP = 1.0) for all branches was the same for the ASTRAL species

trees obtained from the different data schemes while accounting for saturation, codon usage,

compositional heterogeneity, and model of sequence evolution, and was also congruent with the

ASTRAL species tree and concatenated ML from the 105-taxon analyses (Fig. S13). In general,

the proportion of gene trees supporting each bipartition remained the same in every analysis and

showed high levels of conflict among the five major clades of Amaranthaceae s.l. (Fig S13).

580   *Polytomy test*

581   The ASTRAL polytomy test resulted in the same bifurcating species tree for the 11-taxon(tree)

582   dataset and rejected the null hypothesis that any branch is a polytomy (p < 0.01 in all cases).

583   These results were identical when using gene trees with collapsed branches.

585   *Coalescent simulations and tests of the anomaly zone*

The distribution of tree-to-tree distances of the empirical and simulated gene trees to the species

tree from the 11-taxon(tree) dataset largely overlapped (Fig 5a), suggesting that ILS alone is able

to explain most of the observed gene tree heterogeneity (Maureira-Butler et al. 2008). The

anomaly zone limit calculations using species trees from the 11-taxon(tree) dataset detected two

pairs of internodes among the five major groups in Amaranthaceae s.l. that fell into the anomaly

zone (the red pair and the green pair, Fig. 5b; Table S10). The gene tree counts showed that the

species tree was not the most common gene tree topology, as expected for the anomaly zone (Fig

5c). The species tree was the fourth most common gene tree topology (119 out of 4,425 gene

trees), while the three most common gene tree topologies occurred 170, 136, and 127 times (Fig.

5c).

FIGURE 5. Coalescent simulations and tests of the anomaly zone from the 11-taxon(tree) dataset

estimated from individual gene trees. a) Distribution of tree-to-tree distances between empirical

gene trees and the ASTRAL species tree, compared to those from the coalescent simulation. b)

ASTRAL species tree showing branch length in coalescent units. Green and red branches

represent the two pairs of internodes that fall in the anomaly zone (see Table S10 for anomaly

zone limits). c) Gene tree counts (top) of the four most common topologies (bottom). Gene trees

that do not support the monophyly of any of the five major clades were ignored.

DISCUSSION

Using a phylotranscriptomic dataset in combination with reference genomes representing major

clades, we have shown that gene tree discordance prevails along the backbone phylogeny of

Amaranthaceae s.l. Interestingly, we found that this discordance is also present within the plastid

dataset. Despite the strong signal of gene tree discordance, we were able to identify five well-

supported major clades within Amaranthaceae s.l. that are congruent with morphology and

previous taxonomic treatments of the group. Using multiple phylogenetic tools and simulations

we comprehensively tested for processes that might have contributed to the gene tree

discordance in Amaranthaceae s.l. Phylogenetic network analyses and ABBA-BABA tests both

supported multiple reticulation events among the five major clades in Amaranthaceae s.l. At the

same time, the patterns of gene tree discordance among these clades can also largely be

explained by uninformative gene trees and ILS. We found evidence that three consecutive short

internal branches produce anomalous trees contributing to the discordance. Molecular evolution

model misspecification (i.e., substitutional saturation, codon usage bias, or compositional

heterogeneity) was less likely to account for the gene tree discordance. Furthermore, synteny

analysis of reference genomes did not support clustering of any particular topology along the

chromosomes. Taken together, no single source can confidently be pointed out to account for the

strong signal of gene tree discordance, suggesting that the discordance results primarily from

ancient and rapid lineage diversification. Furthermore, the backbone of Amaranthaceae s.l.

remains —and likely will remain— unresolved even with genome-scale data. Our work

highlights problems with identifiability in phylogenetic network analyses, and the need to

comprehensively test for multiple sources of conflict in phylogenomic analyses. Below we

discuss our findings in detail and ultimately provide a set of recommendations for dissecting

phylogenetic signal in ancient and rapid diversifications.

631   *Broad taxonomic circumscription of Amaranthaceae*

Both our nuclear and plastid datasets strongly supported five major clades within Amaranthaceae

s.l.: Amaranthaceae s.s, 'Chenopods I', 'Chenopods II', Betoideae, and Polycnemoideae (Figs. 2

& S4). The monophyly of Amaranthaceae s.s., Betoideae, and Polycnemoideae is consistent with

morphology and recent molecular phylogenetic analyses of these lineages using plastid DNA

(Hohmann et al. 2006; Masson and Kadereit 2013; Di Vincenzo et al. 2018). However, our

nuclear analyses (Fig. 2) supported the monophyly of Chenopodiaceae while the plastid analysis

did not. Both nuclear and plastid datasets showed high levels of conflict among two well-defined

clades, 'Chenopods I' and 'Chenopods II' (Figs 2 & S4). Weak support and/or conflicting

topologies along the backbone on the Amaranthaceae s.l. characterize all previous molecular

studies of the lineage (Fig. 1), even with hundreds of loci (Walker et al. 2018). On the other

hand, all studies support the five major clades found in our analysis.

For the sake of taxonomic stability, we therefore suggest retaining Amaranthaceae s.l.

sensu APG IV (The Angiosperm Phylogeny Group 2016), which includes the previously

recognized Chenopodiaceae. Amaranthaceae s.l. is characterized by a long list of anatomical,

morphological and phytochemical characters such as minute sessile flowers with five tepals, a

single whorl of epitepalous stamens and one basal ovule (Kadereit et al. 2003). Here we

recognize five subfamilies within Amaranthaceae s.l.: Amaranthoideae representing

Amaranthaceae s.s. (incl. Gomphrenoideae Schinz), Betoideae Ulbr., Chenopodioideae

represented as 'Chenopods I' here (incl. Corispermoideae Ulbr.), Polycnemoideae Ulbr., and

Salicornioideae Ulbr. represented by 'Chenopods II' (incl. Salsoloideae Ulbr., Suaedoideae Ulbr.

and Camphorosmoideae A.J. Scott). The stem ages of these five subfamilies date back to the

early Tertiary (Paleocene, Fig. 2) which agrees with dates based on plastid markers (Kadereit et

al. 2012; Di Vincenzo et al. 2018; Yao et al. 2019). Due to the gene tree discordance along the

backbone, the geographic origin of Amaranthaceae s.l. remains ambiguous.

657   *Gene tree discordance detected among plastid genes*

Although both our concatenation-based plastid and nuclear phylogenies supported the same five

major clades of Amaranthaceae s.l., the relationships among these clades are incongruent (Figs. 2

& S4). Cytonuclear discordance is well-known in plants and it has been traditionally attributed to

reticulate evolution (Rieseberg and Soltis 1991; Sang et al. 1995; Soltis and Kuzoff 1995). Such

discordance continues to be treated as evidence in support of hybridization in more recent

phylogenomic studies that assume the plastome to be a single, linked locus (e.g., Folk et al.

2017; Vargas et al. 2017; Morales-Briones et al. 2018b; Lee-Yaw et al. 2019). However, recent

work shows that plastid protein-coding genes might not necessarily act as a single locus and high

levels of tree conflict have been detected (Gonçalves et al. 2019; Walker et al. 2019).

667   In Amaranthaceae s.l., previous studies based on plastid protein-coding genes or introns

(Fig. 1; Kadereit et al. 2003; Müller and Borsch 2005; Hohmann et al. 2006; Kadereit et al.

2017) resulted in different relationships among the five main clades and none in agreement with

our 76-gene plastid phylogeny. Our conflict and QS analyses of the plastid dataset (Figs S5–S6)

revealed strong signals of gene tree discordance among the five major clades of Amaranthaceae

s.l., likely due to heteroplasmy, although the exact sources of conflict are yet to be clarified

(Gonçalves et al. 2019). Unlike the results found by Walker et al. (2019), our individual plastid

gene trees had highly supported nodes (i.e., BS $\geq$ 70, Fig S5), suggesting that low phylogenetic

information content is not the main source of conflict in our plastid dataset.

676   Our results support previous studies showing RNA-seq data can be a reliable source for

plastome assembly (Smith 2013; Osuna-Mascaró et al. 2018; Gitzendanner et al. 2018) RNA-seq

libraries can contain some genomic DNA due to incomplete digestion during RNA purification

(Smith 2013) and given the AT-rich nature of plastomes, this allows plastid DNA to survive the

poly-A selection during mRNA enrichment (Schliesky et al. 2012). However, RNA editing

prediction results showed that our Amaranthaceae s.l. cpDNA assemblies came from RNA rather

than DNA contamination regardless of library preparation by poly-A enrichment (71

transcriptomes) or RiboZero (16 transcriptomes). Similarly, Osuna-Mascaró et al. (2018) also

found highly similar plastome assemblies (i.e., general genome structure, and gene number and

composition) from RNA-seq and genomic libraries, supporting the idea that plastomes are fully

transcribed in photosynthetic eukaryotes (Shi et al. 2016). Furthermore, the backbone topology

of our plastid tree built mainly from RNA-seq data (97 out of 105 samples) was consistent with a

recent complete plastome phylogeny of Caryophyllales mainly from genomic DNA (Yao et al.

2019), showing the utility of recovering plastid gene sequences from RNA-seq data.

Nonetheless, RNA editing might be problematic when combining samples from RNA and DNA,

especially when resolving phylogenetic relationships among closely related species.

693   *Identifiability in methods for detecting reticulation events*

All methods that we used to detect ancient hybridization inferred the presence of reticulation

events, but our results suggest that such methods are likely to struggle with ancient, rapid

radiations. Rapid advances have been made in recent years in developing methods to infer

species networks in the presence of ILS (reviewed in Elworth et al. 2019). These methods have

been increasingly used in phylogenetic studies (e.g., Wen et al. 2016; Copetti et al. 2017;

Morales-Briones et al. 2018a; Crowl et al. 2019). To date, however, species network inference is

still computationally intensive and limited to a small number of species and a few hybridization

events (Hejase and Liu 2016; but see Hejase et al. 2018 and Zhu et al. 2019). Furthermore,

studies evaluating the performance of different phylogenetic network inference approaches are

scarce and restricted to simple hybridization scenarios. Kamneva and Rosenberg (2017) showed

that likelihood methods like Yu et al. (2014) are often robust to ILS and gene tree error when

symmetric hybridization (equal genetic contribution of both parents) events are considered.

While this approach usually does not overestimate hybridization events, it fails to detect skewed

hybridization (unequal genetic contribution of both parents) events in the presence of significant

ILS. Methods developed to scale to larger numbers of species and hybridizations like the ones

using pseudo-likelihood approximations (i.e., Solís-Lemus and Ané 2016; Yu and Nakhleh 2015)

are yet to be evaluated independently, but in the case of the Yu and Nakhleh (2015) method

based on rooted triples, cannot distinguish the correct network when other networks can produce

the same set of triples (Yu and Nakhleh 2015). In contrast, the method of Solís-Lemus and Ané

(2016), based on unrooted quartets, is better at avoiding indistinguishable networks, but it is

limited to only level-1 network scenarios.

715   Applying the above methods to our data set yielded inferences of multiple reticulation

events. The result of our 11-taxon(net) phylogenetic analysis using a pseudo-likelihood approach

detected up to five hybridization events involving all five major clades of Amaranthaceae s.l.

(Fig. 3). Model selection, after calculating the full likelihood of the obtained networks, also

chose the 5-reticulation species as the best model. Likewise, we found that any species network

had a better ML score than a bifurcating tree (Table S5). However, further analyses demonstrated

that full likelihood network searches with up to one hybridization event are indistinguishable

from each other (Table S6), resembling a random gene tree distribution. This pattern can

probably be explained by the high levels of gene tree discordance and lack of phylogenetic signal

in the inferred quartet gene trees (Fig. 4), suggesting that the 11-taxon(net) network searches can

potentially overestimate reticulation events due to high levels of gene tree error or ILS.

726   Using the D-Statistic (Green et al. 2010; Durand et al. 2011) we also detected signals of

introgression in seven possible directions among the five main groups of Amaranthaceae s.l.

(Table S9). The inferred introgression events agreed with at least one of the reticulation

scenarios from the phylogenetic network analysis. However, the D-Statistic did not detect any

introgression that involves Betoideae, which was detected in the phylogenetic network analysis

with either four or five reticulations events. The D-Statistic has been shown to be robust to a

wide range of divergence times, but it is sensitive to relative population size (Zheng and Janke

2018), which agrees with the notion that large effective population sizes and short branches

increase the chances of ILS (Pamilo and Nei 1988) and in turn can dilute the signal for the D-

Statistic (Zheng and Janke 2018). Recently, Elworth et al. (2018) found that multiple or 'hidden'

reticulations can cause the signal of the D-statistic to be lost or distorted. Furthermore, when

multiple reticulations are present, the traditional approach of subsetting datasets into quartets can

be problematic as it largely underestimates D values (Elworth et al. 2018). Given short internal

branches in the backbone of Amaranthaceae s.l. and the phylogenetic network results showing

multiple hybridizations, it is plausible that our D-statistic may be affected by these issues. Our

analysis highlights problems with identifiability in relying on D-statistic or phylogenetic network

analysis alone for detecting reticulation events, especially in cases of ancient and rapid

diversification.

745   *ILS and the Anomaly Zone*

ILS is ubiquitous in multi-locus phylogenetic datasets. In its most severe cases ILS produces the

'anomaly zone', defined as a set of short internal branches in the species tree that produce

anomalous gene trees (AGTs) that are more likely than the gene tree that matches the species tree

(Degnan and Rosenberg 2006). Rosenberg (2013) expanded the definition of the anomaly zone

to require that a species tree contain two consecutive internal branches in an ancestor–descendant

relationship in order to produce AGTs. To date, only a few empirical examples of the anomaly

zone have been reported (Linkem et al. 2016; Cloutier et al. 2019). Our results show that the

species tree of Amaranthaceae s.l. has three consecutive short internal branches that lay within

the limits of the anomaly zone (i.e., $y < a[x]$; Fig. 5; Table S10) and that the species tree is not

the most frequent gene tree (Fig. 4). While both lines of evidence support the presence of AGTs,

it is important to point out that our quartet analysis showed that most quartet gene trees were

equivocal (94–96%; Fig. 4) and were therefore uninformative. Huang and Knowles (2009)

pointed out that the gene tree discordance produced from the anomaly zone can alternatively be

produced by uninformative gene trees and that for species trees with short branches the most

probable gene tree topology is a polytomy rather than an AGT. Our ASTRAL polytomy test,

however, rejected a polytomy along the backbone of Amaranthaceae s.l. in any of the gene tree

sets used. While we did not test for polytomies in individual gene trees, our ASTRAL polytomy

test using gene trees with branches of <75% bootstrap support collapsed also rejected the

presence of a polytomy. Therefore the distribution of gene tree frequency in combination with

short internal branches in the species tree supports the presence of an anomaly zone in

Amaranthaceae s.l.

768   *Considerations in distinguishing sources of gene tree discordance*

The exploration of gene tree discordance has become a fundamental step to understand

recalcitrant relationships across the Tree of Life. Recently, new tools have been developed to

identify and visualize gene tree discordance (e.g., Salichos et al. 2014; Smith et al. 2015; Huang

et al. 2016; Pease et al. 2018). However, downstream methods that evaluate processes generating

observed patterns of gene tree discordance are still in their infancy. In this study, by combining

transcriptomes and genomes, we were able to create a rich and dense dataset to start to tease

apart alternative hypotheses concerning the sources of conflict along the backbone phylogeny of

Amaranthaceae s.l. We found that gene tree heterogeneity observed in Amaranthaceae s.l. can be

explained by a combination of processes, including ILS, hybridization, uninformative genes, and

molecular evolution model misspecification, that might have acted simultaneously and/or

cumulatively.

780   Our results highlight the need to test for multiple sources of conflict in phylogenomic

analyses, especially when trying to resolve phylogenetic relationships with extensive

phylogenetic conflict. Furthermore, we need to be aware of the strengths and limitations of

different phylogenetic methods and be cautious about relying on any single analysis, for example

in the usage of phylogenetics species networks over coalescent-based species trees (Blair and

Ané 2019). We make the following recommendation on five essential steps towards exploring

heterogeneous phylogenetic signals in phylogenomic datasets in general. 1) Study design:

consider whether the taxon sampling and marker choice enable testing alternative sources of

conflicting phylogenetic signal. For example, will there be sufficient phylogenetic signal and

sufficient taxon coverage in individual gene trees for methods such as phylogenetic network

analyses? 2) Data processing: care should be taken in data cleaning, partitioning (e.g., nuclear vs.

plastid), and using orthology inference methods that explicitly address paralogy issues (e.g., tree-

based orthology inference and synteny information). 3) Species tree inference using tools that

accommodate the dataset size and data type (e.g., ASTRAL for gene tree-based inferences or

SVDquartet [Chifman and Kubatko, 2014] for SNP-based inferences), followed by visualization

of phylogenetic conflict using tools such as the pie charts (e.g., PhyParts) and quartet-based tools

(e.g., Quartet Sampling) 4) Phylogenetic network analysis using reduced taxon sampling given

results in step 3; 5) Hypothesis testing given the results of steps 3 and 4. Depending on the

scenario, these can include testing for model misspecification, anomaly zone, uninformative gene

tree, and if hybridization is hypothesized, testing putative reticulation events one at a time, as

illustrated in this study. The backbone phylogeny of Amaranthaceae s.l. remains difficult to

resolve despite employing genome-scale data, a situation that may not be atypical across the Tree

of Life. As we leverage more genomic data and explore gene tree discordance in more detail,

these steps will be informative in other clades, especially in those that are products of ancient

and rapid lineage diversification (e.g., Widhelm et al. 2019). Ultimately, such endeavors will be

instrumental in gaining a full understanding of the complexity of the Tree of Life.

807   SUPPLEMENTARY MATERIAL

808 Data available from the Dryad Digital Repository: http://dx.doi.org/10.5061/.[NNNN]

811   ACKNOWLEDGMENTS

The authors thank H. Freitag, J.M. Bena and the Millennium Seed Bank for providing seeds; U.

Martiné for assisting with RNA extraction; N. Wang for sample sequencing. A. Crum, R. Ree, B.

Carstens, and three anonymous reviewers for providing helpful comments; the Minnesota

Supercomputing Institute (MSI) at the University of Minnesota for providing access to

computational resources. This work was supported by the University of Minnesota, the

University of Michigan, the US National Science Foundation (DEB 1354048), and the

Department of Energy, Office of Science, Genomic Science Program (Contract Number DE-

SC0008834).

821   REFERENCES

822   Alda F., Tagliacollo V.A., Bernt M.J., Waltz B.T., Ludt W.B., Faircloth B.C., Alfaro M.E.,

823   Albert J.S., Chakrabarty P. 2019. Resolving Deep Nodes in an Ancient Radiation of

824   Neotropical Fishes in the Presence of Conflicting Signals from Incomplete Lineage

825   Sorting. Syst. Biol. 68:573–593.

826   Bankevich A., Nurk S., Antipov D., Gurevich A.A., Dvorkin M., Kulikov A.S., Lesin V.M.,

827   Nikolenko S.I., Pham S., Prjibelski A.D., Pyshkin A.V., Sirotkin A.V., Vyahhi N., Tesler

828   G., Alekseyev M.A., Pevzner P.A. 2012. SPAdes: A New Genome Assembly Algorithm

829   and Its Applications to Single-Cell Sequencing. J. Comput. Biol. 19:455–477.

830   Bena M.J., Acosta J.M., Aagesen L. 2017. Macroclimatic niche limits and the evolution of C4

831   photosynthesis in Gomphrenoideae (Amaranthaceae). Bot. J. Linn. Soc. 184:283–297.

832   Blackmon H., Adams R.A. 2015 EvobiR: Tools for comparative analyses and teaching

833   evolutionary biology. doi:10.5281/zenodo.30938

Blair C., Ané C. 2019. Phylogenetic Trees and Networks Can Serve as Powerful and

Complementary Approaches for Analysis of Genomic Data. Syst. Biol. syz056

Bouckaert R., Heled J. 2014. DensiTree 2: Seeing Trees Through the Forest. BioRxiv. 012401.

Brown J.W., Walker J.F., Smith S.A. 2017. Phyx - phylogenetic tools for unix. Bioinformatics.

33:1886–1888.

Bruen T.C., Philippe H., Bryant D. 2006. A Simple and Robust Statistical Test for Detecting the

Presence of Recombination. Genetics. 172:2665–2681.

Buckley T.R., Cordeiro M., Marshall D.C., Simon C. 2006. Differentiating between Hypotheses

of Lineage Sorting and Introgression in New Zealand Alpine Cicadas (Maoricicada

Dugdale). Syst. Biol. 55:411–425.

Chen L.-Y., Morales-Briones D.F., Passow C.N., Yang Y. 2019. Performance of gene expression

analyses using de novo assembled transcripts in polyploid species. Bioinformatics.

35:4314–4320.

Chifman J., Kubatko L. 2014. Quartet Inference from SNP Data Under the Coalescent Model.

Bioinformatics. 30:3317–3324.

Cloutier A., Sackton T.B., Grayson P., Clamp M., Baker A.J., Edwards S.V. 2019. Whole-

Genome Analyses Resolve the Phylogeny of Flightless Birds (Palaeognathae) in the

Presence of an Empirical Anomaly Zone. Syst. Biol. 68:937–955

Cooper E.D. 2014. Overly simplistic substitution models obscure green plant phylogeny. Trends

Plant Sci. 19:576–582.

Copetti D., Búrquez A., Bustamante E., Charboneau J.L.M., Childs K.L., Eguiarte L.E., Lee S.,

Liu T.L., McMahon M.M., Whiteman N.K., Wing R.A., Wojciechowski M.F., Sanderson

M.J. 2017. Extensive gene tree discordance and hemiplasy shaped the genomes of North

American columnar cacti. Proc. Natl. Acad. Sci. 114:12003–12008.

Cox C.J., Li B., Foster P.G., Embley T.M., Civáň P. 2014. Conflicting Phylogenies for Early

Land Plants are Caused by Composition Biases among Synonymous Substitutions. Syst.

Biol. 63:272–279.

Crowl A.A., Manos P.S., McVay J.D., Lemmon A.R., Lemmon E.M., Hipp A.L. 2019.

Uncovering the genomic signature of ancient introgression between white oak lineages

(*Quercus*). New Phytol. nph.15842.

Davidson N.M., Oshlack A. 2014. Corset: enabling differential gene expression analysis for de

novo assembled transcriptomes. Genome Biol. 15:57.

Degnan J.H., Rosenberg N.A. 2006. Discordance of Species Trees with Their Most Likely Gene

Trees. PLoS Genet. 2:e68.

Degnan J.H., Rosenberg N.A. 2009. Gene tree discordance, phylogenetic inference and the

multispecies coalescent. Trends Ecol. Evol. 24:332–340.

Di Vincenzo V., Gruenstaeudl M., Nauheimer L., Wondafrash M., Kamau P., Demissew S.,

Borsch T. 2018. Evolutionary diversification of the African achyranthoid clade

(Amaranthaceae) in the context of sterile flower evolution and epizoochory. Ann. Bot.

122:69–85.

Dohm J.C., Minoche A.E., Holtgräwe D., Capella-Gutiérrez S., Zakrzewski F., Tafer H., Rupp

O., Sörensen T.R., Stracke R., Reinhardt R., Goesmann A., Kraft T., Schulz B., Stadler

P.F., Schmidt T., Gabaldón T., Lehrach H., Weisshaar B., Himmelbauer H. 2014. The

genome of the recently domesticated crop plant sugar beet (Beta vulgaris). Nature.

505:546–549.

Doyle J.J. 1992. Gene Trees and Species Trees: Molecular Systematics as One-Character

880   Taxonomy. Syst. Bot. 17:144.

881   Durand E.Y., Patterson N., Reich D., Slatkin M. 2011. Testing for Ancient Admixture between

882   Closely Related Populations. Mol. Biol. Evol. 28:2239–2252.

883   Edwards S.V. 2009. Is A New and General Theory of Molecular Systematics Emerging?

884   Evolution. 63:1–19.

885   Edwards S.V., Xi Z., Janke A., Faircloth B.C., McCormack J.E., Glenn T.C., Zhong B., Wu S.,

886   Lemmon E.M., Lemmon A.R., Leaché A.D., Liu L., Davis C.C. 2016. Implementing and

887   testing the multispecies coalescent model: A valuable paradigm for phylogenomics. Mol.

888   Phylogenet. Evol. 94:447–462.

889   Elworth R.A.L., Allen C., Benedict T., Dulworth P., Nakhleh L.K. 2018. DGEN: A Test Statistic

890   for Detection of General Introgression Scenarios. WABI.

891   Elworth R.A.L., Ogilvie H.A., Zhu J., Nakhleh L. 2019. Advances in Computational Methods for

892   Phylogenetic Networks in the Presence of Hybridization. In: Warnow T., editor.

893   Bioinformatics and Phylogenetics: Seminal Contributions of Bernard Moret. Cham:

894   Springer International Publishing. p. 317–360.

895   Erfan Sayyari, Siavash Mirarab. 2018. Testing for Polytomies in Phylogenetic Species Trees

896   Using Quartet Frequencies. Genes. 9:132.

897   Flowers T.J., Colmer T.D. 2015. Plant salt tolerance: adaptations in halophytes. Ann. Bot.

898   115:327–331.

899   Folk R.A., Mandel J.R., Freudenstein J.V. 2017. Ancestral Gene Flow and Parallel Organellar

900   Genome Capture Result in Extreme Phylogenomic Discord in a Lineage of Angiosperms.

901   Syst. Biol. 66:320-337.

Fontaine M.C., Pease J.B., Steele A., Waterhouse R.M., Neafsey D.E., Sharakhov I.V., Jiang X.,

Hall A.B., Catteruccia F., Kakani E., Mitchell S.N., Wu Y.-C., Smith H.A., Love R.R.,

Lawniczak M.K., Slotman M.A., Emrich S.J., Hahn M.W., Besansky N.J. 2015.

Extensive introgression in a malaria vector species complex revealed by phylogenomics.

Science. 347:1258524.

Foster P.G. 2004. Modeling Compositional Heterogeneity. Syst. Biol. 53:485–495.

Galtier N., Daubin V. 2008. Dealing with incongruence in phylogenomic analyses. Philos. Trans.

R. Soc. B Biol. Sci. 363:4023–4029.

Gitzendanner M.A., Soltis P.S., Yi T.-S., Li D.-Z., Soltis D.E. 2018. Plastome Phylogenetics: 30

Years of Inferences Into Plant Evolution. Plastid Genome Evolution. Elsevier. p. 293–

313.

Gonçalves D.J.P., Simpson B.B., Ortiz E.M., Shimizu G.H., Jansen R.K. 2019. Incongruence

between gene trees and species trees and phylogenetic signal variation in plastid genes.

Mol. Phylogenet. Evol. 138:219–232.

Grabherr M.G., Haas B.J., Yassour M., Levin J.Z., Thompson D.A., Amit I., Adiconis X., Fan

L., Raychowdhury R., Zeng Q., Chen Z., Mauceli E., Hacohen N., Gnirke A., Rhind N.,

di Palma F., Birren B.W., Nusbaum C., Lindblad-Toh K., Friedman N., Regev A. 2011.

Full-length transcriptome assembly from RNA-Seq data without a reference genome.

Nat. Biotechnol. 29:644–652.

Green R.E., Krause J., Briggs A.W., Maricic T., Stenzel U., Kircher M., Patterson N., Li H., Zhai

W., Fritz M.H.Y., Hansen N.F., Durand E.Y., Malaspinas A.S., Jensen J.D., Marques-

Bonet T., Alkan C., Prufer K., Meyer M., Burbano H.A., Good J.M., Schultz R., Aximu-

Petri A., Butthof A., Hober B., Hoffner B., Siegemund M., Weihmann A., Nusbaum C.,

Lander E.S., Russ C., Novod N., Affourtit J., Egholm M., Verna C., Rudan P., Brajkovic

D., Kucan Z., Gusic I., Doronichev V.B., Golovanova L.V., Lalueza-Fox C., de la Rasilla

M., Fortea J., Rosas A., Schmitz R.W., Johnson P.L.F., Eichler E.E., Falush D., Birney

E., Mullikin J.C., Slatkin M., Nielsen R., Kelso J., Lachmann M., Reich D., Paabo S.

2010. A Draft Sequence of the Neandertal Genome. Science. 328:710–722.

Hejase H.A., Liu K.J. 2016. A scalability study of phylogenetic network inference methods using

empirical datasets and simulations involving a single reticulation. BMC Bioinformatics.

17:422.

Hejase H.A., VandePol N., Bonito G.M., Liu K.J. 2018. FastNet: Fast and Accurate Statistical

Inference of Phylogenetic Networks Using Large-Scale Genomic Sequence Data. Comp.

Genomics.:242–259.

Hernández-Ledesma P., Berendsohn W.G., Borsch T., Mering S.V., Akhani H., Arias S.,

Castañeda-Noa I., Eggli U., Eriksson R., Flores-Olvera H., Fuentes-Bazán S., Kadereit

G., Klak C., Korotkova N., Nyffeler R., Ocampo G., Ochoterena H., Oxelman B.,

Rabeler R.K., Sanchez A., Schlumpberger B.O., Uotila P. 2015. A taxonomic backbone

for the global synthesis of species diversity in the angiosperm order Caryophyllales.

Willdenowia. 45:281.

Hoang D.T., Chernomor O. 2018. UFBoot2: Improving the Ultrafast Bootstrap Approximation.

Mol. Biol. Evol. 35:518–522.

Hohmann S., Kadereit J.W., Kadereit G. 2006. Understanding Mediterranean-Californian

disjunctions: molecular evidence from Chenopodiaceae-Betoideae. TAXON. 55:67–78.

Holder M.T., Anderson J.A., Holloway A.K. 2001. Difficulties in Detecting Hybridization. Syst.

Biol. 50:978–982.

Huang H., Knowles L.L. 2009. What Is the Danger of the Anomaly Zone for Empirical

Phylogenetics? Syst. Biol. 58:527–536.

Huang W., Zhou G., Marchand M., Ash J.R., Morris D., Van Dooren P., Brown J.M., Gallivan

K.A., Wilgenbusch J.C. 2016. TreeScaper: Visualizing and Extracting Phylogenetic

Signal from Sets of Trees. Mol. Biol. Evol. 33:3314–3316.

Jarvis D.E., Ho Y.S., Lightfoot D.J., Schmöckel S.M., Li B., Borm T.J.A., Ohyanagi H., Mineta

K., Michell C.T., Saber N., Kharbatia N.M., Rupper R.R., Sharp A.R., Dally N.,

Boughton B.A., Woo Y.H., Gao G., Schijlen E.G.W.M., Guo X., Momin A.A., Negrão

S., Al-Babili S., Gehring C., Roessner U., Jung C., Murphy K., Arold S.T., Gojobori T.,

Linden C.G.V.D., van Loo E.N., Jellen E.N., Maughan P.J., Tester M. 2017. The genome

of *Chenopodium quinoa*. Nature. 542:307–312.

Kadereit G., Ackerly D., Pirie M.D. 2012. A broader model for C4 photosynthesis evolution in

plants inferred from the goosefoot family (Chenopodiaceae s.s.). Proc. R. Soc. B Biol.

Sci. 279:3304–3311.

Kadereit G., Borsch T., Weising K., Freitag H. 2003. Phylogeny of Amaranthaceae and

Chenopodiaceae and the Evolution of C4 Photosynthesis. Int. J. Plant Sci. 164:959–986.

Kadereit G., Hohmann S., Kadereit J.W. 2006. A synopsis of Chenopodiaceae subfam. Betoideae

and notes on the taxonomy of *Beta*. Willdenowia. 36:9–19.

Kadereit G., Newton R.J., Vandelook F. 2017. Evolutionary ecology of fast seed germination—

A case study in Amaranthaceae/Chenopodiaceae. Perspect. Plant Ecol. Evol. Syst. 29:1–

11.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.

970   ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods.

971   14:587–589.

972   Kamneva O.K., Rosenberg N.A. 2017. Simulation-Based Evaluation of Hybridization Network

973   Reconstruction Methods in the Presence of Incomplete Lineage Sorting. Evol.

974   Bioinforma. 13:117693431769193.

975   Katoh K., Standley D.M. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

976   Improvements in Performance and Usability. Mol. Biol. Evol. 30:772–780.

977   Kearse M., Moir R., Wilson A., Stones-Havas S., Cheung M., Sturrock S., Buxton S., Cooper A.,

978   Markowitz S., Duran C., Thierer T., Ashton B., Meintjes P., Drummond A. 2012.

979   Geneious Basic: An integrated and extendable desktop software platform for the

980   organization and analysis of sequence data. Bioinformatics. 28:1647–1649.

981   Knowles L.L., Huang H., Sukumaran J., Smith S.A. 2018. A matter of phylogenetic scale:

982   Distinguishing incomplete lineage sorting from lateral gene transfer as the cause of gene

983   tree discord in recent versus deep diversification histories. Am. J. Bot. 105:376–384.

984   Kubatko L.S., Chifman J. 2019. An invariants-based method for efficient identification of hybrid

985   species from large-scale genomic data. BMC Evol. Biol. 19:112.

986   Lanfear R., Calcott B., Ho S.Y.W., Guindon S. 2012. PartitionFinder: Combined Selection of

987   Partitioning Schemes and Substitution Models for Phylogenetic Analyses. Mol. Biol.

988   Evol. 29:1695–1701.

989   Lee-Yaw J.A., Grassa C.J., Joly S., Andrew R.L., Rieseberg L.H. 2019. An evaluation of

990   alternative explanations for widespread cytonuclear discordance in annual sunflowers

991   (*Helianthus*). New Phytol. 221:515–526.

Lightfoot D.J., Jarvis D.E., Ramaraj T., Lee R., Jellen E.N., Maughan P.J. 2017. Single-molecule

sequencing and Hi-C-based proximity-guided assembly of amaranth (*Amaranthus*

hypochondriacus) chromosomes provide insights into genome evolution. BMC Biol.

15:74.

Linkem C.W., Minin V.N., Leaché A.D. 2016. Detecting the Anomaly Zone in Species Trees

and Evidence for a Misleading Signal in Higher-Level Skink Phylogeny (Squamata:

Scincidae). Syst. Biol. 65:465–477.

Liu L., Yu L. 2010. Phybase: an R package for species tree analysis. Bioinformatics. 26:962–

963.

Liu Y., Cox C.J., Wang W., Goffinet B. 2014. Mitochondrial Phylogenomics of Early Land

Plants: Mitigating the Effects of Saturation, Compositional Heterogeneity, and Codon-

Usage Bias. Syst. Biol. 63:862–878.

Maddison W.P. 1997. Gene Trees in Species Trees. Syst. Biol. 46:532–536.

Masson R., Kadereit G. 2013. Phylogeny of Polycnemoideae (Amaranthaceae): Implications for

biogeography, character evolution and taxonomy. TAXON. 62:100–111.

Maureira-Butler I.J., Pfeil B.E., Muangprom A., Osborn T.C., Doyle J.J. 2008. The Reticulate

History of *Medicago* (Fabaceae). Syst. Biol. 57:466–482.

Mclean B.S., Bell K.C., Allen J.M., Helgen K.M., Cook J.A. 2019. Impacts of Inference Method

and Data set Filtering on Phylogenomic Resolution in a Rapid Radiation of Ground

Squirrels (Xerinae: Marmotini). Syst. Biol. 68:298–316.

Meyer B.S., Matschiner M., Salzburger W. 2017. Disentangling Incomplete Lineage Sorting and

Introgression to Refine Species-Tree Estimates for Lake Tanganyika Cichlid Fishes. Syst.

Biol. 66:531–550.

Mirarab S., Bayzid M.S., Warnow T. 2016. Evaluating Summary Methods for Multilocus

Species Tree Estimation in the Presence of Incomplete Lineage Sorting. Syst. Biol.

65:366–380.

Morales-Briones D.F., Liston A., Tank D.C. 2018a. Phylogenomic analyses reveal a deep history

of hybridization and polyploidy in the Neotropical genus *Lachemilla* (Rosaceae). New

Phytol. 218:1668–1684.

Morales-Briones D.F., Romoleroux K., Kolář F., Tank D.C. 2018b. Phylogeny and Evolution of

the Neotropical Radiation of *Lachemilla* (Rosaceae): Uncovering a History of Reticulate

Evolution and Implications for Infrageneric Classification. Syst. Bot. 43:17–34.

Moray C., Goolsby E.W., Bromham L. 2016. The Phylogenetic Association Between Salt

Tolerance and Heavy Metal Hyperaccumulation in Angiosperms. Evol. Biol. 43:119–

130.

Mower J.P. 2009. The PREP suite: predictive RNA editors for plant mitochondrial genes,

chloroplast genes and user-defined alignments. Nucleic Acids Res. 37:W253–W259.

Müller K., Borsch T. 2005. Phylogenetics of Amaranthaceae Based on *matK/trnK* Sequence

Data: Evidence from Parsimony, Likelihood, and Bayesian Analyses. Ann. Mo. Bot.

Gard. 92:66–102.

Nguyen L.-T., Schmidt H.A., von Haeseler A., Minh B.Q. 2015. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol. Biol. Evol.

32:268–274.

Osuna-Mascaró C., Rubio de Casas R., Perfectti F. 2018. Comparative assessment shows the

reliability of chloroplast genome assembly using RNA-seq. Sci. Rep. 8:17404.

Pamilo P., Nei M. 1988. Relationships between Gene Trees and Species Trees. Mol. Biol. Evol.

5:568–583.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2018. Quartet Sampling

distinguishes lack of support from conflicting support in the green plant tree of life. Am.

J. Bot. 105:385–403.

Piirainen M., Liebisch O., Kadereit G. 2017. Phylogeny, biogeography, systematics and

taxonomy of Salicornioideae (Amaranthaceae/Chenopodiaceae) – A cosmopolitan, highly

specialized hygrohalophyte lineage dating back to the Oligocene. Taxon. 66:109–132.

Prasanna A.N., Gerber D., Kijpornyongpan T., Aime M.C., Doyle V.P., Nagy L.G. 2020. Model

Choice, Missing Data, and Taxon Sampling Impact Phylogenomic Inference of Deep

Basidiomycota Relationships. 69:17–37

R Core Team. 2019. R: A Language and Environment for Statistical Computing. Vienna,

Austria: R Foundation for Statistical Computing.

Rannala B., Yang Z. 2003. Bayes Estimation of Species Divergence Times and Ancestral

Population Sizes Using DNA Sequences From Multiple Loci. Genetics. 166:1645–1656.

Rieseberg L.H., Soltis D.E. 1991. Phylogenetic consequences of cytoplasmic gene flow in plants.

Evol. Trends Plants. 5:65–84.

Robinson D.F., Foulds L.R. 1981. Comparison of phylogenetic trees. Math. Biosci. 53:131–147.

Rosenberg N.A. 2013. Discordance of Species Trees with Their Most Likely Gene Trees: A

Unifying Principle. Mol. Biol. Evol. 30:2709–2713.

Roycroft E.J., Moussalli A., Rowe K.C. 2019. Phylogenomics Uncovers Confidence and

Conflict in the Rapid Radiation of Australo-Papuan Rodents. Syst. Biol. syz044.

Salichos L., Stamatakis A., Rokas A. 2014. Novel Information Theory-Based Measures for

Quantifying Incongruence among Phylogenetic Trees. Mol. Biol. Evol. 31:1261–1271.

Sang T., Crawford D.J., Stuessy T.F. 1995. Documentation of reticulate evolution in peonies

(Paeonia) using internal transcribed spacer sequences of nuclear ribosomal DNA:

implications for biogeography and concerted evolution. Proc. Natl. Acad. Sci. 92:6813–

6817.

Sayyari E., Mirarab S. 2016. Fast Coalescent-Based Computation of Local Branch Support from

Quartet Frequencies. Mol. Biol. Evol. 33:1654–1668.

Schliep K.P. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27:592–593.

Schliesky S., Gowik U., Weber A.P.M., Bräutigam A. 2012. RNA-Seq Assembly – Are We

There Yet? Front. Plant Sci. 3.

Schwarz G. 1978. Estimating the Dimension of a Model. Ann. Stat. 6:461–464.

Sharp P.M., Li W.-H. 1986. An evolutionary perspective on synonymous codon usage in

unicellular organisms. J. Mol. Evol. 24:28–38.

Shi C., Wang S., Xia E.-H., Jiang J.-J., Zeng F.-C., Gao L.-Z. 2016. Full transcription of the

chloroplast genome in photosynthetic eukaryotes. Sci. Rep. 6:30135.

Shimodaira H. 2002. An Approximately Unbiased Test of Phylogenetic Tree Selection. Syst.

Biol. 51:492–508.

Shimodaira H., Hasegawa M. 2001. CONSEL: for assessing the confidence of phylogenetic tree

selection. Bioinformatics. 17:1246–1247.

Smith D.R. 2013. RNA-Seq data: a goldmine for organelle research. Brief. Funct. Genomics.

12:454–456.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals

conflict, concordance, and gene duplications with examples from animals and plants.

BMC Evol. Biol. 15:745.

Smith S.A., O'Meara B.C. 2012. treePL: divergence time estimation using penalized likelihood

for large phylogenies. Bioinformatics. 28:2689–2690.

Solís-Lemus C., Ané C. 2016a. Inferring Phylogenetic Networks with Maximum

Pseudolikelihood under Incomplete Lineage Sorting. PLOS Genet. 12:e1005896.

Soltis D.E., Kuzoff R.K. 1995. Discordance between nuclear and chloroplast phylogenies in the

Heuchera group (Saxifragaceae). Evolution. 49:727–742.

Srivastava S.K. 1969. Assorted angiosperm pollen from the Edmonton Formation

(Maestrichtian), Alberta, Canada. Can. J. Bot. 47:975–989.

Stamatakis A. 2014. RAxML version 8 - a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 30:1312–1313.

Sugiura N. 1978. Further analysts of the data by akaike' s information criterion and the finite

corrections. Commun. Stat. - Theory Methods. 7:13–26.

Swofford D. 2002. PAUP*. Phylogenetic analysis using parsimony (*and other methods) version

4. Sunderland MA Sinauer Assoc.

Than C., Ruths D., Nakhleh L. 2008. PhyloNet: a software package for analyzing and

reconstructing reticulate evolutionary relationships. BMC Bioinformatics. 9:322–16.

The Angiosperm Phylogeny Group, Chase M.W., Christenhusz M.J.M., Fay M.F., Byng J.W.,

Judd W.S., Soltis D.E., Mabberley D.J., Sennikov A.N., Soltis P.S., Stevens P.F. 2016.

An update of the Angiosperm Phylogeny Group classification for the orders and families

of flowering plants: APG IV. Bot. J. Linn. Soc. 181:1–20.

Vargas O.M., Ortiz E.M., Simpson B.B. 2017. Conflicting phylogenomic signals reveal a pattern

of reticulate evolution in a recent high-Andean diversification (Asteraceae: Astereae:

Diplostephium). New Phytol. 214:1736–1750.

Walker J.F., Walker-Hale N., Vargas O.M., Larson D.A., Stull G.W. 2019. Characterizing gene

tree conflict in plastome-inferred phylogenies. PeerJ. 7:e7747.

Walker J.F., Yang Y., Feng T., Timoneda A., Mikenas J., Hutchison V., Edwards C., Wang N.,

Ahluwalia S., Olivieri J., Walker-Hale N., Majure L.C., Puente R., Kadereit G.,

Lauterbach M., Eggli U., Flores-Olvera H., Ochoterena H., Brockington S.F., Moore

M.J., Smith S.A. 2018. From cacti to carnivores: Improved phylotranscriptomic sampling

and hierarchical homology inference provide further insight into the evolution of

Caryophyllales. Am. J. Bot. 105:446–462.

Wen D., Yu Y., Hahn M.W., Nakhleh L. 2016. Reticulate evolutionary history and extensive

introgression in mosquito species revealed by phylogenetic network analysis. Mol. Ecol.

25:2361–2372.

Wen D., Yu Y., Zhu J., Nakhleh L. 2018. Inferring Phylogenetic Networks Using PhyloNet.

Syst. Biol. 67:735–740.

Widhelm T.J., Grewe F., Huang J.-P., Mercado-Díaz J.A., Goffinet B., Lücking R., Moncada B.,

Mason-Gamer R., Lumbsch H.T. 2019. Multiple historical processes obscure

phylogenetic relationships in a taxonomically difficult group (Lobariaceae, Ascomycota).

Sci. Rep. 9:8968.

Xu B., Yang Z. 2016. Challenges in Species Tree Estimation Under the Multispecies Coalescent

Model. Genetics. 204:1353–1368.

Xu C., Jiao C., Sun H., Cai X., Wang X., Ge C., Zheng Y., Liu W., Sun X., Xu Y., Deng J.,

Zhang Z., Huang S., Dai S., Mou B., Wang Q., Fei Z., Wang Q. 2017. Draft genome of

spinach and transcriptome diversity of 120 Spinacia accessions. Nat. Commun. 8:15275.

Yang Y., Moore M.J., Brockington S.F., Timoneda A., Feng T., Marx H.E., Walker J.F., Smith

S.A. 2017. An Efficient Field and Laboratory Workflow for Plant Phylotranscriptomic

Projects. Appl. Plant Sci. 5:1600128.

Yang Y., Smith S.A. 2014. Orthology Inference in Nonmodel Organisms Using Transcriptomes

and Low-Coverage Genomes: Improving Accuracy and Matrix Occupancy for

Phylogenomics. Mol. Biol. Evol. 31:3081–3092.

Yao G., Jin J.-J., Li H.-T., Yang J.-B., Mandala V.S., Croley M., Mostow R., Douglas N.A.,

Chase M.W., Christenhusz M.J.M., Soltis D.E., Soltis P.S., Smith S.A., Brockington S.F.,

Moore M.J., Yi T.-S., Li D.-Z. 2019. Plastid phylogenomic insights into the evolution of

Caryophyllales. Mol. Phylogenet. Evol. 134:74–86.

Yu Y., Degnan J.H., Nakhleh L. 2012. The Probability of a Gene Tree Topology within a

Phylogenetic Network with Applications to Hybridization Detection. PLoS Genet.

8:e1002660–10.

Yu Y., Dong J., Liu K.J., Nakhleh L. 2014. Maximum likelihood inference of reticulate

evolutionary histories. Proc. Natl. Acad. Sci. 111:16448–16453.

Yu Y., Nakhleh L. 2015. A maximum pseudo-likelihood approach for phylogenetic networks.

BMC Genomics. 16:S10.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:523.

Zhao T., Schranz M.E. 2019. Network-based microsynteny analysis identifies major differences

and genomic outliers in mammalian and angiosperm genomes. Proc. Natl. Acad. Sci.

116:2165–2174.

Zheng Y., Janke A. 2018. Gene flow analysis method, the D-statistic, is robust in a wide

parameter space. BMC Bioinformatics. 19:10.

Zhu J., Liu X., Ogilvie H.A., Nakhleh L.K. 2019. A divide-and-conquer method for scalable

phylogenetic network inference from multilocus data. Bioinformatics. 35:i370–i378.