Disentangling Sources of Gene Tree Discordance in Phylotranscriptomic Datasets: A Case

Study from Amaranthaceae s.l.

Diego F. Morales-Briones[1]*, Gudrun Kadereit[2], Delphine T. Tefarikis[2], Michael J. Moore[3],

Stephen A. Smith[4], Samuel F. Brockington[5], Alfonso Timoneda[5], Won C. Yim[6], John C.

Cushman[6], Ya Yang[1]*

[1] Department of Plant and Microbial Biology, University of Minnesota-Twin Cities, 1445

Gortner Avenue, St. Paul, MN 55108, USA

[2] Institut für Molekulare Physiologie, Johannes Gutenberg-Universität Mainz, D-55099, Mainz,

Germany

[3] Department of Biology, Oberlin College, Science Center K111, 119 Woodland Street, Oberlin,

OH 44074-1097, USA

[4] Department of Ecology & Evolutionary Biology, University of Michigan, 830 North University

Avenue, Ann Arbor, MI 48109-1048, USA

[5] Department of Plant Sciences, University of Cambridge, Tennis Court Road, Cambridge, CB2

3EA, United Kingdom

[6] Department of Biochemistry and Molecular Biology, University of Nevada, Reno, NV, 89577,

USA

* Correspondence to be sent to: Diego F. Morales-Briones and Ya Yang. Department of Plant and

Microbial Biology, University of Minnesota, 1445 Gortner Avenue, St. Paul, MN 55108, USA,

Telephone: +1 612-625-6292 (YY) Email: dfmoralesb@gmail.com; yangya@umn.edu

Abstract.— Phylogenomic datasets have become common and fundamental to understanding the

phylogenetic relationships of recalcitrant groups across the Tree of Life. At the same time,

working with large genomic or transcriptomic datasets requires special attention to the processes

that generate gene tree discordance, such as data processing and orthology inference, incomplete

lineage sorting, hybridization, model violation, and uninformative gene trees. Methods to

estimate species trees from phylogenomic datasets while accounting for all sources of conflict

are not available, but a combination of multiple approaches can be a powerful tool to tease apart

alternative sources of conflict. Here using a phylotranscriptomic analysis in combination with

reference genomes, we explore sources of gene tree discordance in the backbone phylogeny of

the plant family Amaranthaceae s.l. The dataset was analyzed using multiple phylogenetic

approaches, including coalescent-based species trees and network inference, gene tree

discordance analyses, site pattern test of introgression, topology test, synteny analyses, and

simulations. We found that a combination of processes might have acted, simultaneously and/or

cumulatively, to generate the high levels of gene tree discordance in the backbone of

Amaranthaceae s.l. Furthermore, other analytical shortcomings like uninformative genes as well

as misspecification of the model of molecular evolution seem to contribute to tree discordance

signal in this family. Despite the comprehensive phylogenomic dataset and detailed analyses

presented here, no single source can confidently be pointed out to account for the strong signal of

gene tree discordance, suggesting that the backbone of Amaranthaceae s.l. might be a product of

an ancient and rapid lineage diversification, and remains —and probably will remain—

unresolved even with genome-scale data. Our work highlights the need to test for multiple

sources of conflict in phylogenomic analyses and provide a set of recommendations moving

forward in disentangling ancient and rapid diversification.

Keywords: Amaranthaceae; gene tree discordance; hybridization; incomplete lineage sorting;

phylogenomics; transcriptomics; species tree; species network.

The detection of gene tree discordance is ubiquitous in the phylogenomic era. As large

phylogenomic datasets are becoming more common (e.g. Jarvis et al. 2014; Misof et al. 2014;

Wickett et al. 2014; Hughes et al. 2018; Walker et al. 2018; Laumer et al. 2019; Varga et al.

2019), exploring gene tree heterogeneity in such datasets (e.g. Salichos et al. 2014; Smith et al.

2015; Huang et al. 2016; Arcila et al. 2017; Pease et al. 2018, is essential for inferring

phylogenetic relationships while accommodating and understanding the underlying processes

that produce gene tree conflict.

Discordance among gene trees can be the product of multiple sources. These include

errors and noise in data assembly and filtering, hidden paralogy, incomplete lineage sorting

(ILS), gene duplication/loss (Pamilo and Nei 1988; Doyle 1992; Maddison 1997; Galtier and

Daubin 2008), random noise from uninformative genes, as well as misspecified model

parameters of molecular evolution such as substitutional saturation, codon usage bias, or

compositional heterogeneity (Foster 2004; Cooper 2014; Cox et al. 2014; Li et al. 2014; Liu et

al. 2014). Among these potential sources of gene tree discordance, ILS is the most studied in the

systematics literature (Edwards 2009), and a number of phylogenetic inference methods have

been developed that accommodate ILS as the source of discordance (reviewed in Edwards et al.

2016; Mirarab et al. 2016; Xu and Yang 2016). More recently, methods that account for

additional processes such as hybridization or introgression have gained attention. These include

methods that estimate phylogenetic networks while accounting for ILS and hybridization

simultaneously (Yu et al. 2014; Yu and Nakhleh 2015; Solís-Lemus and Ané 2016; Wen et al.

2016b; Wen and Nakhleh 2018; Zhang et al. 2018a; Zhu et al. 2018; Zhu et al. 2019), and

methods that detect introgression based on site patterns or phylogenetic invariants (Green et al.

2010; Durand et al. 2011; Patterson et al. 2012; Eaton and Ree 2013; Pease and Hahn 2015;

Elworth et al. 2018; Glémin et al. 2019; Kubatko and Chifman 2019).

The above sources of gene tree discordance can act alone, but most often multiple

sources may contribute to gene tree heterogeneity (Holder et al. 2001; Buckley et al. 2006;

Maureira-Butler et al. 2008; Joly et al. 2009; Meyer et al. 2017; Knowles et al. 2018; Glémin et

al. 2019). However, at present no method can estimate species trees from phylogenomic data

while modeling multiple sources of conflict and molecular substitution simultaneously. To

overcome these limitations, the use of multiple phylogenetic tools and data partitioning schemes

in phylogenomic datasets have become a common practice in order to disentangle sources of

gene tree heterogeneity and resolve recalcitrant relationships at deep and shallow nodes of the

Tree of Life (e.g. Duchêne et al. 2018; Prasanna et al. 2019; Alda et al. 2019; Roycroft et al.

2019; Widhelm et al. 2019).

Here we explore these issues in the plant family Amaranthaceae s.l., including the

previously segregated family Chenopodiaceae (Hernández-Ledesma et al. 2015; The

Angiosperm Phylogeny Group et al. 2016). With c. 2050 to 2500 species in 181 genera and a

worldwide distribution (Hernández-Ledesma et al. 2015), Amaranthaceae s.l. are iconic for the

repeated evolution of complex traits representing adaptations to extreme environments such as

C_4 photosynthesis in hot and often dry environments (e.g. Kadereit et al. 2012; Bena et al. 2017),

various modes of extreme salt tolerance (e.g. Flowers and Colmer 2015; Piirainen et al. 2017)

that in several species are coupled with heavy metal tolerance (Moray et al. 2016), and very fast

seed germination and production of multiple diaspore types on one individual (Kadereit et al.

2017). Amaranthaceae s.l. contains a number of crops, some of them with a long cultivation

history, such as the pseudocereals quinoa and amaranth (Jarvis et al. 2017), and some that have

been taken under cultivation more recently, such as sugar beet (Dohm et al. 2014), spinach,

glassworts, and *Salsola soda*. Many species of the family are important fodder plants in arid

regions and several are currently being investigated for their soil ameliorating and desalinating

effects. Reference genomes are available for *Beta vulgaris* (sugar beet, subfamily Betoideae;

Dohm et al. 2014), *Chenopodium quinoa* (quinoa, Chenopodioideae; Jarvis et al. 2017), *Spinacia*

oleracea (spinach; Chenopodioideae; Xu et al. 2017) and *Amaranthus hypochondriacus*

(amaranth; Amaranthoideae; Lightfoot et al. 2017), representing three of the 13 currently

recognized subfamilies (sensu Kadereit et al. 2003; Kadereit et al. 2017).

123   Within the core Caryophyllales the previously recognized families Amaranthaceae s.s.

and Chenopodiaceae have always been regarded as closely related and their separate family

status has been subjected to phylogenetic and taxonomic debate repeatedly (see Kadereit et al.

2003; Masson and Kadereit 2013; Hernández-Ledesma et al. 2015; Walker et al. 2018; Fig. 1).

Their common ancestry was first concluded from a number of shared morphological, anatomical

and phytochemical synapomorphies and later substantiated by molecular phylogenetic studies

with the Achatocarpaceae as sister group (see Kadereit et al. 2003 and references therein).

Amaranthaceae s.s. has a predominant tropical and subtropical distribution with the highest

diversity found in the Neotropics, eastern and southern Africa and Australia (Müller and Borsch

2005), while the previously segregated family Chenopodiaceae predominantly occurs in

temperate regions and semi-arid or arid environments of subtropical regions (Kadereit et al.

2003). The key problem has always been the species-poor and heterogeneous subfamilies

Polycnemoideae and Betoideae, which do not fit comfortably morphologically in either the

Chenopodiaceae or Amaranthaceae s.s. (cf. Table 5 in Kadereit et al. 2003). Polycnemoideae are

similar in ecology and distribution to Chenopodiaceae but share important floral traits such as

FIGURE 1. Phylogenetic hypothesis of Amaranthaceae s.l. from previous studies. a) Kadereit et al. (2003) using the chloroplast (cpDNA) *rbcL* coding region. b) Müller and Borsch (2005); using the chloroplast *matK* coding region and partial *trnL* intron. c) Hohmann et al. (2006) using the chloroplast *ndhF* coding region. d) Kadereit et al. (2017) using the chloroplast *atpB-rbcL* spacer, *matK* with *trnL* intron, *ndhF-rpL32* spacer, and *rps16* intron e) Walker et al. (2018) using 455 nuclear genes from transcriptome data. Major clades of Amaranthaceae s.l. named following the results of this study. Image credits: *Amaranthus hypochondriacus* by Picture Partners, *Beta vulgaris* by Olha Huchek, *Chenopodium quinoa* by Diana Mower, *Nitrophila mohavensis* by James M. André, and *Salsola soda* by Homeydesign.

petaloid tepals, filament tubes and 2-locular anthers with Amaranthaceae s.s. Morphologically, Betoideae fit into either of the two traditionally circumscribed families but have a unique fruit type—a capsule that opens with a circumscissile lid (Kadereit et al. 2006). Both Betoideae and

Polycnemoideae show strongly disjunct distribution patterns, occurring each with only a few

species on three different continents. Furthermore, the genera of both subfamilies display a

number of morphologically dissociating features. Both intercontinental disjunctions of species-

poor genera and unique morphological traits led to the hypothesis that Betoideae and

Polycnemoideae might be relicts of, or from hybridization among early-branching lineages in

Amaranthaceae s.l. (Hohmann et al. 2006; Masson and Kadereit 2013).

Previous molecular phylogenetic analyses struggled to resolve the relationships among

Betoideae, Polycnemoideae and the rest of the Amaranthaceae s.l. (Kadereit et al. 2003; Müller

and Borsch 2005; Kadereit et al. 2012; Masson and Kadereit 2013; Walker et al. 2018). The first

phylogenomic study of Amaranthaceae s.l. by Walker et al. (2018) revealed that gene tree

discordance mainly occurred at deeper nodes of the phylogeny involving Betoideae.

Polycnemoideae was resolved as sister to Chenopodiaceae in Walker et al. (2018), albeit with

low (17%) gene tree concordance, which contradicted previous analyses based on chloroplast

data (Masson and Kadereit 2013). However, only a single species of Betoideae (the cultivated

beet and its wild relative) was sampled in Walker et al. (2018). In addition, sources of conflicting

signals among species trees remained unexplored.

In this study, we leverage 71 publicly available transcriptomes, 17 newly sequenced

transcriptomes, and 4 reference genomes that span all 13 subfamilies of Amaranthaceae s.l. and

include increased taxon sampling in Betoideae. Consistent with previous analyses, we identified

high levels of gene tree discordance in the backbone phylogeny of Amaranthaceae s.l. Using a

combination of phylogenetic approaches, we explored multiple sources that can explain such

conflict. We tested for 1) ancient hybridization, focusing on the hypothesis of the hybrid origin

of Polycnemoideae and Betoideae, between Amaranthaceae s.s. and Chenopodioideae, 2)

discordance produced by misspecifications of model of molecular evolution, and 3) discordance

due to ILS as a result of short internal branches in the backbone phylogeny of Amaranthaceae s.l.

In addition, we comprehensively updated the phylotranscriptomic pipeline of Yang and Smith

(2014) with additional features of filtering isoforms and spurious tips. Our results showed that

both species network and site pattern methods that model gene flow while accounting for ILS

detected signals of multiple hybridization events in Amaranthaceae s.l. However, when these

hybridization events were analyzed individually, most of the gene tree discordance could be

explained by uninformative gene trees. In addition, the high level of gene tree discordance in

Amaranthaceae s.l. could also be explained by three consecutive short branches that produce

anomalous gene trees. Combined, our results showed that multiple processes might have

contributed to the gene tree discordance in Amaranthaceae s.l., and that we might not be able to

distinguish among these processes even with genomic-scale sampling and synteny information.

Finally, we make recommendations on strategies for disentangling multiple sources of gene tree

discordance in phylogenomic datasets.

191   **MATERIALS AND METHODS**

193   An overview of all dataset and phylogenetic analyses can be found in Figure S1. Scripts for raw

194   data processing, assembly, translation, and homology and orthology search can be found at

195   https://bitbucket.org/yanglab/phylogenomic_dataset_construction/ as part of an updated

196   'phylogenomic dataset construction' pipeline (Yang and Smith 2014).

199   *Taxon sampling, transcriptome sequencing*

We sampled 92 species (88 transcriptomes and four genomes) representing all 13 currently

recognized subfamilies and 16 out of 17 tribes of Amaranthaceae s.l. (sensu [Kadereit et al.

2003; Kadereit et al. 2017]). In addition, 13 outgroups across the Caryophyllales were included

(ten transcriptomes and three genomes; Table S1). We generated 17 new transcriptomes for this

study (Table S2). For *Tidestromia oblongifolia*, tissue collection, RNA isolation, library

preparation was carried out using the KAPA Stranded mRNA-Seq Kits (KAPA Biosystems,

Wilmington, Massachusetts, USA). The library was multiplexed with 10 other samples from a

different project on an Illumina HiSeq2500 platform with V4 chemistry at the University of

Michigan Sequencing Core (Yang et al. 2017). For the remaining 16 samples total RNA was

isolated from c. 70-125 mg leaf tissue collected in liquid nitrogen using the RNeasy Plant Mini

Kit (Qiagen) following the manufacturer's protocol (June 2012). A DNase digestion step was

included with the RNase-Free DNase Set (Qiagen). Quality and quantity of RNA were checked

on the NanoDrop (Thermo Fisher Scientific) and the 2100 Bioanalyzer (Agilent Technologies).

Library preparation was carried out using the TruSeq® Stranded Total RNA Library Prep Plant

with RiboZero probes (96 Samples. Illumina, #20020611). Indexed libraries were normalized,

pooled and size selected to 320bp +/- 5% using the Pippin Prep HT instrument to generate

libraries with mean inserts of 200 bp, and sequenced on the Illumina HiSeq2500 platform with

V4 chemistry at the University of Minnesota Genomics Center. Reads from all 17 libraries were

paired-end 125 bp.

222   *Transcriptome data processing and assembly*

We processed raw reads for all 98 transcriptome datasets (except *Bienertia sinuspersici*) used in

this study (88 ingroups + 10 outgroups; Table S1). Sequencing errors in raw reads were corrected

with Rcorrector (Song and Florea 2015) and reads flagged as uncorrectable were removed.

Sequencing adapters and low-quality bases were removed with Trimmomatic v0.36

(SLIDINGWINDOW:4:5 LEADING:5 TRAILING:5 MINLEN:25; Bolger et al. 2014).

Additionally, reads were filtered for chloroplast and mitochondrial reads with Bowtie2 v 2.3.2

(Langmead and Salzberg 2012) using publicly available Caryophyllales organelle genomes from

the Organelle Genome Resources database (RefSeq; [Pruitt et al. 2007]; last accessed on October

17, 2018) as references. Read quality was assessed with FastQC v 0.11.7

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Finally, overrepresented

sequences detected with FastQC were discarded. *De novo* assembly was carried out with Trinity

v 2.5.1 (Haas et al. 2013) with default settings, but without in silico normalization. Assembly

quality was assessed with Transrate v 1.0.3 (Smith-Unna et al. 2016). Low quality and poorly

supported transcripts were removed using individual cut-off values for three contig score

components of Transrate: 1) proportion of nucleotides in a contig that agrees in identity with the

aligned read, $s(Cnuc) \leq 0.25$; 2) proportion of nucleotides in a contig that have one or more

mapped reads, $s(Ccov) \leq 0.25$; and 3) proportion of reads that map to the contig in correct

orientation, $s(Cord) \leq 0.5$. Furthermore, chimeric transcripts (*trans*-self and *trans*-multi-gene)

were removed following the approach described in Yang and Smith (2013) using *Beta vulgaris*

as the reference proteome, and percentage similarity and length cutoffs of 30 and 100,

respectively. In order to remove isoforms and assembly artifacts, filtered reads were remapped to

filtered transcripts with Salmon v 0.9.1 (Patro et al. 2017) and putative genes were clustered with

Corset v 1.07 (Davidson and Oshlack 2014) using default settings, except that we used a minimal

of five reads as threshold to remove transcripts with low coverage (-m 5). Only the longest

transcript of each putative gene inferred by Corset was retained. Our previous benchmark study

have shown that Corset followed by selecting the longest transcript for each putative gene

performed well in reducing isoforms and assembly artifacts, especially in polyploid species

(Chen et al. 2019). Filtered transcripts were translated with TransDecoder v 5.0.2 (Haas et al.

2013) with default settings and the proteome of *Beta vulgaris* and *Arabidopsis thaliana* to

identify open reading frames. Finally, translated amino acid sequences were further reduced with

CD-HIT v 4.7 (-c 0.99; [Fu et al. 2012]) to remove near-identical amino acid sequences.

255   *Homology and orthology inference*

Initial homology inference was carried out following Yang and Smith (2014) with some

modification. First, an all-by-all BLASTN search was performed on coding sequences (CDS)

using an *E* value cutoff of 10 and max_target_seqs set to 100. Raw BLAST output was filtered

with a hit fraction of 0.4. Then putative homologs groups were clustered using MCL v 14-137

(van Dongen 2000) with a minimal minus log-transformed *E* value cutoff of 5 and an inflation

value of 1.4. Finally, only clusters with a minimum of 25 taxa were retained. Individual clusters

were aligned using MAFFT v 7.307 (Katoh and Standley 2013) with settings '–genafpair –

maxiterate 1000'. Aligned columns with more than 90% missing data were removed using Phyx

(Brown et al. 2017). Homolog trees were built using RAxML v 8.2.11 (Stamatakis 2014) with a

GTR-CAT model and clade support assessed with 200 rapid bootstrap (BS) replicates. Spurious

or outlier long tips were detected and removed with TreeShrink v 1.0.0 (Mai and Mirarab 2018).

Monophyletic and paraphyletic tips that belonged to the same taxon were removed keeping the

tip with the highest number of characters in the trimmed alignment. After visual inspection of ca.

50 homolog trees, internal branches longer than 0.25 were likely representing deep paralogs.

These branches were cut apart, keeping resulting subclades with a minimum of 25 taxa.

Homolog tree inference, tip masking, outlier removal, and deep paralog cutting was carried out

for a second time using the same settings to obtain final homologs. Orthology inference was

carried out following the 'monophyletic outgroup' approach from Yang and Smith (2014),

keeping only ortholog groups with at least 25 ingroup taxa. The 'monophyletic outgroup'

approach filters for clusters that have outgroup taxa being monophyletic and single-copy, and

therefore filters for single- and low-copy genes. It then roots the gene tree by the outgroups,

traverses the rooted tree from root to tip, and removes the side with less taxa when gene

duplication is detected at any given node.

280   *Chloroplast assembly*

Although DNase treatment is carried out to remove genomic DNA, due to its high copy number,

chloroplast sequences are often carried over in RNA-seq libraries. In addition, as young leaf

tissue was used for RNA-seq, RNA from chloroplast genes are expected to be represented,

especially in libraries prepared using a RiboZero approach. To investigate phylogenetic signal

from plastome sequences, *de novo* assemblies were carried out with the Fast-Plast v.1.2.6

pipeline (https://github.com/mrmckain/Fast-Plast) using the organelle reads from the filtering

step.lNo complete or single-contig plastomes were obtained. Filtered contigs produced by

Spades v 3.9.0 (Bankevich et al. 2012) were mapped to the closest available reference plastome

(with an Inverted Repeat removed; Table S3) and manually edited in Geneious v.11.1.5 (Kearse

et al. 2012) to produce final oriented contigs.

292   *Assessment of recombination*

293   Coalescent species tree methods assume that there is no recombination within loci and free

294   recombination between loci. To determine the presence of recombination in our dataset, we used

295   the Φ (pairwise homoplasy index) test for recombination, as implemented in PhiPack (Bruen et

296   al. 2006). We tested recombination on the final set of ortholog alignments (with a minimum of

297   25 taxa) with the default sliding window size of 100 bp.

299   *Nuclear phylogenetic analysis*

We used concatenation and coalescent-based methods to reconstruct the phylogeny of

Amaranthaceae s.l. Sequences from final orthologs were aligned with MAFFT, columns were

trimmed with Phyx requiring a minimal occupancy of 30%, and alignments with at least 1,000

characters and 99 out of 105 taxa were retained. We first estimated a maximum likelihood (ML)

tree of the concatenated matrix with RAxML using a partition-by-gene scheme with GTR-CAT

model for each partition and clade support assessed with 200 rapid bootstrap (BS) replicates. To

estimate a coalescent-based species tree, first we inferred individual ML gene trees using

RAxML with a GTR-CAT model and 200 BS replicates to assess clade support. Individual gene

trees were then used to estimate a species tree with ASTRAL-III v5.6.3 (Zhang et al. 2018b)

using local posterior probabilities (LPP; Sayyari and Mirarab 2016) to assess clade support.

311   *Detecting and visualizing nuclear gene tree discordance*

To explore discordance among gene trees, we first calculated the internode certainty all (ICA)

value to quantify the degree of conflict on each node of a target tree (i.e. species tree) given

individual gene trees (Salichos et al. 2014). In addition, we calculated the number of conflicting

and concordant bipartitions on each node of the species trees. We calculated both the ICA scores

and the number of conflicting/concordant bipartitions with Phyparts (Smith et al. 2015), mapping

against the estimated ASTRAL species trees, using individual gene trees with BS support of at

least 50% for the corresponding node. Additionally, in order to distinguish strong conflict from

weakly supported branches, we evaluated tree conflict and branch support with Quartet Sampling

(QS; Pease et al. 2018) using 100 replicates. Quartet Sampling subsamples quartets from the

input tree and alignment and assess the confidence, consistency, and informativeness of each

internal branch by the relative frequency of the three possible quartet topologies (Pease et al.

2018)

Furthermore, in order to visualize conflict, we built a cloudogram using DensiTree v2.2.6

(Bouckaert and Heled 2014). We filtered the final ortholog alignments to include only 41 species

(38 ingroup and 3 outgroups) in order to include as many orthologs as possible while

representing all main clades of Amaranthaceae s.l. (see results). Individual gene trees were

inferred as previously described. Trees were time-calibrated with TreePL v1.0 (Smith and

O'Meara 2012) by fixing the crown age of Amaranthaceae s.l. to 66–72.1 based on a pollen

record of *Polyporina cribraria* from the late Cretaceous (Maastrichtian; Srivastava 1969), and

the root for the reduced 41-species dataset (most common recent ancestor of Achatocarpaceae

and Aizoaceae) was set to 95 Ma based on the time-calibrated plastome phylogeny of

Caryophyllales from Yao et al. (2019).

337   *Chloroplast phylogenetic analysis*

Assembled contigs (excluding one inverted repeat region) were aligned with MAFFT with the

setting '--auto'. Two samples (*Dysphania schraderiana* and *Spinacia turkestanica*) were

removed due to low sequence occupancy. Using the annotations of the reference genomes (Table

S3), the coding regions of 78 genes were extracted and each gene alignment was visually

inspected in Geneious to check for potential misassemblies. From each gene alignment taxa with

short sequences (i.e. < 50% of the aligned length) were removed and realigned with MAFFT.

The genes *rpl32* and *ycf2* were excluded from downstream analyses due to low taxon occupancy

(Table S4). For each individual gene we performed extended model selection (Kalyaanamoorthy

et al. 2017) followed by ML gene tree inference and 1,000 ultrafast bootstrap replicates for

branch support (Hoang and Chernomor 2018) in IQ-Tree v.1.6.1 (Nguyen et al. 2015). For the

concatenated matrix we searched for the best partition scheme (Lanfear et al. 2012) followed by

ML gene tree inference and 1,000 ultrafast bootstrap replicates for branch support in IQ-Tree.

Additionally, we evaluated branch support with QS using 1,000 replicates and gene tree

discordance with PhyParts in the ML and species tree. Finally, to identify the origin of the

chloroplast reads (i.e. genomic or RNA), we predicted RNA editing from CDS alignments using

PREP (Mower 2009) with the alignment mode (PREP-aln), and a cutoff value of 0.8.

355   *Species network analysis using a reduced 11-taxon dataset*

We inferred species networks that model ILS and gene flow using a maximum pseudo-likelihood

approach (Yu and Nakhleh 2015). Species network searches were carried out with PhyloNet

v.3.6.9 (Than et al. 2008) with the command 'InferNetwork_MPL' and using the individual gene

trees as input. Due to computational restrictions, and given our main focus was to search for

potential reticulating events among major clades of Amaranthaceae s.l., we reduced our taxon

sampling to one outgroup and ten ingroup taxa including two representative species from each of

the five well-supported major lineages in Amaranthaceae s.l. (see results). We filtered the final

105-taxon ortholog alignments to include genes that have all 11 taxa [referred herein as 11-

taxon(net) dataset]. After realignment and trimming we kept genes with a minimum of 1,000

aligned base pairs and individual ML gene trees were inferred with RAxML with a GTR-

GAMMA model and 200 bootstrap replicates. We carried out 10 independent network searches

allowing for up to five hybridization events for each search. To estimate the optimum number of

hybridizations, first we optimized the branch lengths and inheritance probabilities and computed

the likelihood of the best scored network from each of the five maximum hybridization events

searches. Network likelihoods were estimated given the individual gene trees, as implemented in

Yu et al. (2012), using the command 'CalGTProb' in PhyloNet. Then, we performed model

selection using the bias-corrected Akaike information criterion (AICc; Sugiura 1978), and the

Bayesian information criterion (BIC; Schwarz 1978). The number of parameters was set to the

number of branch lengths being estimated plus the number of hybridization probabilities being

estimated. The number of gene trees used to estimate the likelihood was used to correct for finite

sample size. To compare network models to bifurcating trees, we also estimated ML and

coalescent-based species trees as well as a chloroplast tree with the same taxon sampling used in

the network searches. Tree inferences were carried out as previously described for the ML,

coalescent-based, and chloroplast trees, respectively.

Hypothesis testing and detecting introgression using four-taxon datasets

Given the signal of multiple clades potentially involved in hybridization events detected by

PhyloNet (see results), we next conducted quartet analyses to explore a single event at a time.

First, we further reduced the 11-taxon(net) dataset to six taxa that included one outgroup genome

(*Mesembryanthemum crystallinum*) and one ingroup from each of the five major ingroup clades:

Amaranthus hypochondriacus (genome)*, Beta vulgaris* (genome)*, Chenopodium quinoa*

(genome)*, Caroxylon vermiculatum* (transcriptome)*,* and *Polycnemum majus* (transcriptome) to

represent Amaranthaceae s.s., Betoideae, 'Chenopods I', 'Chenopods II' and Polycnemoideae,

respectively. We carried out a total of ten quartet analyses using all ten four-taxon combinations

that included three out of five ingroup species and one outgroup. We filtered the final set of 105-

taxon ortholog alignments for genes with all four taxa for each combination and inferred

individual gene trees as described before. For each quartet we carried out the following analyses.

We first estimated a species tree with ASTRAL and explored gene tree conflict with PhyParts.

We then explored individual gene tree resolution by calculating the Tree Certainty (TC) score

(Salichos et al. 2014) in RAxML using the majority rule consensus tree across the 200 bootstrap

replicates. Next, we explored potential correlation between TC score and alignment length, GC

content and alignment gap proportion using a linear regression model in R v.3.6.1 (R Core Team

2019). Finally, we tested for the fit of gene trees to the three possible rooted quartet topologies

for each gene using the approximately unbiased (AU) tests (Shimodaira 2002). We carried out

ten constraint searches for each of three topologies in RAxML with the GTR-GAMMA model,

then calculated site-wise log-likelihood scores for the three constraint topologies in RAxML

using the GTR-GAMMA and carried out the AU test using Consel v.1.20 (Shimodaira and

Hasegawa 2001). In order to detect possible introgression among species of each quartet, first we

estimated a species network with PhyloNet using a full maximum likelihood approach (Yu et al.

2014) with 100 independent searches while optimizing the likelihood of the branch lengths and

inheritance probabilities for every proposed species network. Furthermore, we also carried out

the ABBA/BABA test to detect introgression (Green et al. 2010); Durand et al. 2011; Patterson

et al. 2012) in each quartet. We calculated the D-statistic and associated z score for the null

hypothesis of no introgression ($D = 0$) following each quartet ASTRAL species tree for taxon

order assignment using 100 jackknife replicates and a block size of 10,000 bp with evobiR v1.2

(Blackmon and Adams) in R.

414   Additionally, to visualize any genomic patterns of the phylogenetic history of *Beta*

vulgaris regarding its relationship with Amaranthaceae s.s. and Chenopodiaceae, we first

identified syntenic regions between the genomes of *Beta vulgaris* and the outgroup

Mesembryanthemum crystallinum using the SynNet pipeline

(https://github.com/zhaotao1987/SynNet-Pipeline; Zhao and Schranz 2019). We used

DIAMOND v.0.9.24.125 (Buchfink et al. 2015) to perform all-by-all inter- and intra-pairwise

protein searches with default parameters, and MCScanX (Wang et al. 2012) for pairwise synteny

block detection with default parameters, except match score (-k) that was set to five. Then, we

plot the nine chromosomes of *Beta vulgaris* by assigning each of the 8,258 orthologs of the

quartet composed of *Mesembryanthemum crystallinum* (outgroup), *Amaranthus*

hypochondriacus, Beta vulgaris, and *Chenopodium quinoa* (BC1A) to synteny blocks and to one

of the three possible quartet topologies based on best likelihood score.

Assessment of substitutional saturation, codon usage bias, compositional heterogeneity, and

model of sequence evolution misspecification

We refiltered the final 105-taxon ortholog alignments to again include genes that have the same

11 taxa (referred herein as 11-taxon(tree) dataset used for the species network analyses. We

realigned individual genes using MACSE v.2.03 (Ranwez et al. 2018) to account for codon

structure and frameshifts. Codons with frameshifts were replaced with gaps, and ambiguous

alignment sites were removed using GBLOCKS v0.9b (Castresana 2000) while accounting for

codon alignment (-t=c -b1=6 -b2=6 -b3=2 -b4=2 -b5=h). After realignment and removal of

ambiguous sites, we kept genes with a minimum of 300 aligned base pairs. To detect potential

saturation, we plotted the uncorrected genetic distances against the inferred distances as

described in Philippe and Forterre (1999). The level of saturation was determined by the slope of

the linear regression between the two distances where a shallow slope (i.e < 1) indicates

saturation. We estimated the level of saturation by concatenating all genes and dividing the first

and second codon positions from the third codon positions. We calculated uncorrected, and

inferred distances with the TN93 substitution model using APE v5.3 (Paradis and Schliep 2019)

in R. To determine the effect of saturation in the phylogenetic inferences we estimated individual

gene trees using three partition schemes. We inferred ML trees with an unpartitioned alignment,

a partition by first and second codon positions, and the third codon positions, and by removing

all third codon positions. All tree searches were carried out in RAxML with a GTR+GAMMA

model and 200 bootstrap replicates. A species tree for each of the three data schemes was

estimated with ASTRAL and gene tree discordance was examined with PhyParts.

Codon usage bias was evaluated using a correspondence analysis of the Relative

Synonymous Codon Usage (RSCU), which is defined as the number of times a particular codon

is observed relative to the number of times that the codon would be observed in the absence of

any codon usage bias (Sharp and Li 1986). RSCU for each codon in the 11-taxon concatenated

alignment was estimated with CodonW v.1.4.4 (Peden 1999). Correspondence analysis was

carried out using FactoMineR v1.4.1(Lê et al. 2008) in R. To determine the effect of codon usage

bias in the phylogenetic inferences we estimated individual gene trees using codon-degenerated

alignments. Alignments were recoded to eliminate signals associated with synonymous

substitutions by degenerating the first and third codon positions using ambiguity coding using

DEGEN v1.4 (Regier et al. 2010; Zwick et al. 2012). Gene tree inference and discordance

analyses were carried out on the same three data schemes as previously described.

460   To examine the presence of among-lineage compositional heterogeneity, individual genes

were evaluated using the compositional homogeneity test that uses a null distribution from

simulations as proposed by Foster (2004). We performed the compositional homogeneity test by

optimizing individual gene trees with a GTR-GAMMA model and 1,000 simulations in P4

(Foster 2004). To assess if compositional heterogeneity had an effect in species tree inference

and gene tree discordance, gene trees that showed the signal of compositional heterogeneity were

removed from saturation and codon usage analyses and the species tree and discordance analyses

were rerun.

468   To explore the effect of sequence evolution model misspecification, we reanalyzed the

datasets from the saturation and codon usage analyses using inferred gene trees that accounted

for model selection. We performed extended model selection followed by ML gene tree

inference and 1,000 ultrafast bootstrap replicates for branch support in IQ-Tree. Species tree

inference, conflict analysis and removal of genes with compositional heterogeneity were carried

out as previously described.

Finally, we also used amino acid alignments from MACSE to account for substitutional

saturation. Amino acid positions with frameshifts were replaced with gaps, and ambiguous

alignment sites were removed with Phyx requiring a minimal occupancy of 30%. We inferred

individual gene trees with IQ-tree to account for a model of sequence evolution and carried out

species tree inference, conflict analysis, and removal of genes with compositional heterogeneity

as described for the nucleotide alignments.

481   *Polytomy test*

To explore if the gene tree discordance among the main clades of Amaranthaceae s.l. could be

explained by polytomies instead of bifurcating nodes, we carried out the polytomy test by

Sayyari and Mirarab (2018) as implemented in ASTRAL. This test uses quartet frequencies to

assess whether a branch should be replaced with a polytomy while accounting for ILS. We

performed the polytomy test using the gene trees inferred from the saturation and codon usage

analyses [11-taxon(tree) dataset]. Because this test can be sensitive to gene tree error (Syyari and

Mirarab 2018), we ran the analyses using the original gene trees and also using gene trees where

branches with less than 75% of bootstrap support were collapsed.

491   *Coalescent simulations*

To investigate if gene tree discordance can be explained by ILS alone, we carried out coalescent

simulations similar to Cloutier et al. (2019) An ultrametric species tree with branch lengths in

mutational units (μT) was estimated by constraining an ML tree search of the 11-taxon(net)

concatenated alignment (from individual MAFFT gene alignment) to the ASTRAL species tree

topology with a GTR+GAMMA model while enforcing a strict molecular clock in PAUP v4.0a

(build 165; Swofford 2002). The mutational branch lengths from the constrained tree and branch

lengths in coalescent units ($\tau = T/4N_e$) from the ASTRAL species trees were used to estimate the

population size parameter theta ($\Theta = \mu T/\tau$; Degnan and Rosenberg 2009) for internal branches.

Terminal branches were set with a population size parameter theta of one. We used the R

package Phybase v. 1.4 (Liu and Yu 2010) which uses the formula from Rannala and Yang

(2003) to simulate 10,000 gene trees using the constraint tree and the estimated theta values.

Then the tree-to-tree distances using the Robinson and Foulds (1981) metric was calculated

between the species tree and each gene tree and compared with the distribution of tree-to-tree

distances between the species tree and the simulated gene tree. Tree-to-tree distances were

calculated using the R package Phangorn v2.5.3 (Schliep 2011). We ran simulations in seven

species trees and associated gene tree distribution to represent the trees and gene tree

distributions from the saturation, codon usage and model selection analyses that accounted for

branch length variation in the species trees and individual gene tree inference. Following

Maureira-Butler et al. (2008), if the tree-to-tree distances between the species trees and gene

trees were larger than 95% of the distribution of tree-to-tree distances of the species trees and the

simulated gene trees then ILS alone is considered unlikely to explain the gene tree heterogeneity.

514   *Test of anomaly zone*

Anomaly zone occurs where a set of short internal branches in the species tree produces gene

trees that differ from the species tree more frequently than those that are concordant [a(x); as

defined in equation 4 of Degnan and Rosenberg (2006)]. To explore if gene tree discordance

observed in Amaranthaceae s.l. is a product of the anomaly zone, we estimated the boundaries of

the anomaly zone [a(x); as defined in equation 4 of Degnan and Rosenberg (2006)] for the

internal nodes of the species tree. Here, x is the branch length (coalescent units) in the species

tree that has a descendant internal branch. If the length of the descendant internal branch (y) is

smaller than a(x), then the internode pair is in the anomaly zone and is likely to produce anomaly

gene trees (AGTs). We carried out the calculation of a(x) following Linkem et al. (2016) in the

same 11-taxon(tree) ASTRAL species trees used for coalescent simulations to account for branch

length variation. Additionally, to establish the frequency of gene trees that were concordant with

the estimated species trees, we quantified the frequency of all 105 possible rooted gene trees

(when clades of Amaranthaceae sl. are monophyletic). We calculated tree-to-tree distances

between the 105 possible topologies and all 5,936 gene trees and counted how many times a

topology had a distance of zero among the set of gene trees.

<center>**RESULTS**</center>

<center>*Transcriptome sequencing, assembly, translation, and quality control*</center>

We generated 17 new transcriptomes of Amaranthaceae s.l. for this study. Raw reads are

available from the NCBI Sequence Read Archive (BioProject: XXXX; Table S2). The number of

raw read pairs ranged from 17 to 27 million. For the 16 samples processed using RiboZero

organelle reads accounted for 15% to 52% of read pairs (Table S2). For *Tidestromia oblongifolia*

that poly-A enrichment was carried out in library prep with ~5% of raw reads were from

organelle (Table S2). Number of final CDS (after quality control and redundancy reduction) used

for all-by-all homology search can be found in Table S5. The final number of orthologs from the

'monophyletic outgroup' approach was 13,024 with a mean of 9,813 orthologs per species

(Table S6).

Assessment of recombination

545   The test for recombination, Φ, identified 82 out of the 13,024 genes from the final set of

546   orthologs (with a minimum of 25 taxa) with a strong signal of recombination (p ≤ 0.05; Table

547   S7). Alignments that showed signal of recombination were removed from all subsequent

548   phylogenetic analyses.

Analysis of the nuclear dataset of Amaranthaceae s.l.

The final set of nuclear orthologous genes included 936 genes with at least 99 out of 105 taxa

and 1,000 bp in aligned length after removal of low occupancy columns (the 105-taxon dataset,

Fig. S1). The concatenated matrix consisted of 1,712,054 columns with a gene and character

occupancy of 96% and 82%, respectively. The species tree from ASTRAL and the concatenated

ML tree from RAxML recovered the exact same topology with most clades with the highest

support [i.e. bootstrap percentage (BS) = 100, local posterior probabilities (LPP) = 1; Fig. 2; Figs

S2–S3]. Our phylogenetic analyses recovered Chenopodiaceae as monophyletic with the

subfamilies and relationships among them similar to Kadereit et al. (2017). Betoideae was placed

as sister of Chenopodiaceae, while Polycnemoideae was placed as sister (BS = 97, LPP = 0.98)

to the clade composed of Chenopodiaceae and Betoideae. Finally, we recovered Amaranthaceae

s.s. with an overall topology concordant to Kadereit et al. (2017), with the exception of *Iresine*

that is placed among the Aervoids (Fig. 2; Figs S2–S3).

FIGURE 2. Maximum likelihood phylogeny of Amaranthaceae s.l. inferred from RAxML

analysis of the concatenated 936-nuclear gene supermatrix. All nodes have full support

(Bootstrap = 100/Local posterior probability = 100) unless noted next to nodes. Boxes contain

gene tree conflict and Quartet Sampling (QS) scores for major clades (see Figs S2–S3 for all

node scores). In each box, numbers on the upper left indicate the number of gene trees

concordant/conflicting with that node in the species tree, and the number on the lower left

indicate the Internode Certainty All (ICA) score. Pie charts present the proportion of gene trees

that support that clade (blue), the proportion that support the main alternative bifurcation (green),

the proportion that support the remaining alternatives (red), and the proportion (conflict or

support) that have < 50% bootstrap support (gray). Number on the right of the pie chart indicates

QS scores: Quartet concordance/Quartet differential/Quartet informativeness. QS scores in blue

indicate support for individual major clades of Amaranthaceae s.l., while red scores indicate

strong support for alternative relationships among them. Branch lengths are in number of

substitutions per site (scale bar on the bottom).

The conflict analyses confirmed the monophyly of Amaranthaceae s.l. with most gene

trees being concordant (922; ICA= 0.94) and full QS support (1/–/1; i.e. all sampled quartets

supported that branch), but also recovered significant discordance in the backbone of the family

(Fig. 2; Figs S2–S3). The monophyly of Chenopodiaceae s.s. was supported only by 231 out of

632 informative gene trees (ICA = 0.42) and the QS score (0.25/0.19/0.99) suggested weak

quartet support with a skewed frequency for an alternative placement of two well-defined clades

within Chenopodiaceae s.s., herein referred to as 'Chenopods I' and 'Chenopods II' (Fig. 2; Figs

S2–S3). 'Chenopods I' and 'Chenopods II' were each supported by the majority of gene trees,

870 (ICA = 0.89) and 916 (ICA = 0.91), respectively and full QS support. The placement of

Betoideae and Polycnemoideae as successive sisters of Chenopodiaceae also showed significant

conflict (Fig. 2; Figs S2–S3). The placement of Betoideae was supported only by 126 out of 579

informative gene trees (ICA = 0.28) and the QS score (0.31/0.57/1) also showed low support

with the presence of supported alternative placements close to the same frequency. Similarly, the

placement of Polycnemoideae was supported by only 116 out of 511 informative gene trees (ICA

= 0.29) and low QS support (0.3/0.81/0.99) with alternative topologies close to equal

frequencies. The monophyly of Amaranthaceae s.s. was highly supported by 755 gene trees (ICA

=0.85) and the QS score (0.92/0/1) also indicated high quartet support and no support for a single

alternative topology.

Congruent with the overall low support in the backbone of Amaranthaceae s.l. from BS,

LPP, ICA, QS, and PhyParts, the cloudogram of 41 species using 1,242 gene trees also showed

significant conflict in the backbone of Amaranthaceae s.l. where no clear pattern can be

identified regarding the relationships of the five main clades of Amaranthaceae s.l. (Fig. 3). In

summary, analysis of nuclear genes recovered five well-supported clades in Amaranthaceae s.l.:

Amaranthaceae s.s., Betoideae, 'Chenopods I', 'Chenopods II', and Polycnemoideae. However,

relationships among these five clades showed a high level of conflict among genes (ICA scores

and gene counts [pie charts]) and among subsampled quartets (QS scores), despite having high

support from both BS and LPP scores.

FIGURE 3. ASTRAL species tree (left) and cloudogram (right) inferred from 1,242 nuclear genes

for the 41-taxon dataset of Amaranthaceae s.l. Pie charts on nodes present the proportion of gene

trees that support that clade (blue), the proportion that support the main alternative bifurcation

(green), the proportion that support the remaining alternatives (red), and the proportion (conflict

or support) that have < 50% bootstrap support (gray).

617   *Chloroplast phylogenetic analysis of Amaranthaceae s.l.*

The final alignment from 76 genes included 103 taxa and 55,517 bp in aligned length. The ML

tree recovered the same five main clades within Amaranthaceae s.l. with the highest support (BS

= 100; Fig. 4; Figs S4–S5). Within each main clade, relationships were fully congruent with

(Kadereit et al. 2017) and mostly congruent with our nuclear analyses. However, the relationship

among the five main clades differed from the nuclear tree. Here, Betoideae was retrieved as

sister (BS = 100) of 'Chenopods I', while Amaranthaceae s.s. and Polycnemoideae were also

recovered as sister clades (BS =100). Furthermore, the clade formed by Betoideae and

'Chenopods I', and Amaranthaceae s.s. and Polycnemoideae were recovered as sister groups (BS

= 73), leaving 'Chenopods II' as sister to the former two. Conflict analysis confirmed the

monophyly of Amaranthaceae s.l. with 51 out of 76 gene trees supporting this clade (ICA = 0.29)

and full QS support (1/–/1). On the other hand, and similar to the nuclear phylogeny, conflict and

QS analyses showed significant discordance in the backbone of the family (Fig. 4; Figs S4–S5).

The sister relationship of Betoideae and 'Chenopods I' was supported by only 20 gene trees (ICA

= 0.06), but it had a strong support from QS (0.84/0.88/0/94). The relationship between

Amaranthaceae s.s. and Polycnemoideae was supported only by 15 gene trees (ICA = 0.07),

while QS showed weak support (0.41/0.21.0.78) with signals of a supported secondary

evolutionary history. The clade uniting Betoideae, 'Chenopods I', Amaranthaceae s.s., and

Polycnemoideae was supported by only four-gene trees, with counter-support from both QS (-

0.29/0.42/0.75) and ICA (-0.03), suggesting that most gene trees and sampled quartets supported

alternative topologies. RNA editing prediction analysis revealed editing sites only on CDS

sequences of reference plastome genomes (Table S3), suggesting that cpDNA reads in RNA-seq

libraries come from RNA rather than DNA contamination from incomplete DNase digestion

during sample processing.

FIGURE 4. Maximum likelihood phylogeny of Amaranthaceae s.l. inferred from IQ-tree analysis

of concatenated 76-chloroplast gene supermatrix. All nodes have full support (Bootstrap =

100/Local posterior probability = 100) unless noted next to nodes. Boxes contain gene tree

conflict and Quartet Sampling (QS) scores for major clades (see Figs S2–S3 for all node scores).

In each box, numbers on the upper left indicate the number of gene trees concordant/conflicting

with that node in the species tree, and the number on the lower left indicate the Internode

Certainty All (ICA) score. Pie charts present the proportion of gene trees that support that clade

(blue), the proportion that support the main alternative bifurcation (green), the proportion that

support the remaining alternatives (red), and the proportion (conflict or support) that have < 50%

bootstrap support (gray). Numbers on the right of the pie chart indicate QS scores: Quartet

concordance/Quartet differential/Quartet informativeness. QS scores in blue indicate support for

individual major clades of Amaranthaceae s.l., while red scores indicate strong support for

alternative relationships among them. Branch lengths are in number of substitutions per site

(scale bar on the bottom).

658   *Species network analysis of Amaranthaceae s.l.*

Due to the computational limit of species network analyses, we reduced our full 105-taxon

dataset to ten ingroup taxa plus one outgroup taxon. In this reduced dataset two taxa were used to

represent the diversity for each of the five well-supported ingroup clades within Amaranthaceae

s.l. The reduced 11-taxon(net) dataset included 4,138 orthologous gene alignments with no

missing taxon and a minimum of 1,000 bp (aligned length after removal of low occupancy

columns). The 11-taxon(net) ASTRAL species tree was congruent with the 105-taxon tree, while

both the nuclear and chloroplast ML trees from concatenated supermatrices both had different

topologies than their corresponding 105-taxon trees (Fig. 5). PhyloNet identified up to five

hybridization events among the clades of Amaranthaceae s.l. (Fig. 5), with the best model having

five hybridization events involving all five clades (Table 1). 'Chenopods II' was involved in

hybridization events in all networks with one to five hybridization events. Model selection

indicated that any species network was a better model than the bifurcating nuclear or chloroplast

trees (Table 1).

FIGURE 5. Species trees and species networks of the reduced 11-taxon(net) dataset of

Amaranthaceae s.l. a) Maximum likelihood phylogeny inferred from RAxML analysis of the

concatenated 4,138-nuclear gene supermatrix. b) Species tree inferred with ASTRAL using

4,138 nuclear genes. c) Maximum likelihood tree inferred from IQ-tree analysis of the

concatenated 76-chloroplast gene supermatrix. d–h). Best species network inferred from

PhyloNet pseudolikelihood analyses with 1 to 5 maximum number of hybridizations. Red and

blue indicates the minor and major edges, respectively, of hybrid nodes. Number next to the

branches indicates inheritance probabilities for each hybrid node.

690 **TABLE 1.** Model selection between maximum number of hybridizations in species networks searches.

Topology	Maximum number of hybridizations allowed	Number of inferred hybridizations	ln(*L*)	Parameters	Number of loci	AICc	ΔAICc	BIC	ΔBIC
RAxML ML tree	NA	NA	-24486.33124	19	4138	49048.84703	20589.66354	49130.89387	20546.62287
ASTRAL species tree	NA	NA	-23448.39741	19	4138	46972.97939	18513.79589	47055.02622	18470.75522
Chloroplast ML tree	NA	NA	-24568.33287	19	4138	49212.8503	20753.66681	49294.89713	20710.62614
Network 1	1	1	-21177.79113	21	4138	42439.80675	13980.62326	42530.46958	13946.19859
Network 2	2	2	-17275.62523	23	4138	34643.51881	6184.335324	34742.79372	6158.522728
Network 3	3	2	-16741.99114	23	4138	33576.25064	5117.067147	33675.52555	5091.254551
Network 4	4	3	-15415.80012	25	4138	30931.91638	2472.73289	31039.79943	2455.528435
Network 5	**5**	**5**	**-14171.37996**	**29**	**4138**	**28459.18349**	**0**	**28584.27099**	**0**

699   *Four-taxon analyses*

To test for hybridization events one at a time, we further reduced the 11-taxon(net) dataset to 10

four-taxon combinations that each included one outgroup and one representative each from three

out of the five major ingroup clades. Between 7,756 and 8,793 genes were used for each quartet

analysis (Table 2) and each quartet topology can be found in Figure 6. Only five out of the ten

bifurcating quartet species trees (H0 and more frequent gene tree) were compatible with the

nuclear species tree inferred from the complete 105-taxon dataset. The other five quartets

compatible with the complete-taxon species tree corresponded to the second most frequent

quartet gene trees, except for the quartet of Betoideae, 'Chenopods II' and Polycnemoideae

(PBC2, which correspond to the least frequent gene tree).

FIGURE 6. Gene counts from Approximate-Unbiased (AU) topology test of the 10 quartets from

the five main clades of Amaranthaceae s.l. AU tests were carried out between the three possible

topologies of each quartet. H0 represents the ASTRAL species tree of each quartet. Equivocal

indicates gene trees that fail to reject all three alternative topologies for a quartet with $p \leq 0.05$.

Gene counts for each of the three alternative topologies represent gene trees supporting

unequivocally one topology by rejecting the other two alternatives with $p \leq 0.05$. Insets represent

gene count only for unequivocally topology support. Each quartet is named following the species

tree topology, where the first two species are sister to each other (all topologies can be found in

Figure S1). A = Amaranthaceae s.s. (represented by *Amarantus hypocondriacus*), B = Betoideae

(*Beta vulgaris*), C1 = Chenopods I (*Chenopodium quinoa*), C2 = Chenopods II (*Caroxylum*

vermiculatum), P = Polycnemoideae (*Polycnemum majus*). All quartets are rooted with

Mesembryanthemum crystallinum.

Similar to the 105-taxon and the 11-taxon(net) datasets, the conflict analyses recovered

significant conflict among all three possible rooted quartet topologies in all ten quartets. In each

of the ten quartets, the ASTRAL species tree topology (H0) was the most frequent among

individual gene trees (raw counts) but only with 35%–41% of occurrences while the other two

topologies varied between similar or slightly skewed frequencies (Fig. S6a; Table S8). Gene

counts based on the raw likelihood scores from the constraint analyses showed similar patterns

(Fig. S6b; Table S8). Furthermore, when gene counts were filtered by significant likelihood

support (i.e. **Δ**AICc $\geq$ 2), the number of trees supporting each of the three possible topologies

dropped between 34% and 45%, but the species tree remained to be the most frequent topology

for all quartets (Fig. S6b; Table S8). The AU topology tests failed to reject ($p \leq 0.05$)

approximately 85% of the gene trees for any of the three possible quartet topologies and rejected

all but a single topology in only 3%–4.5% of cases. Among the unequivocally selected gene

trees, the frequencies among the three alternative topologies were similar to ones based on raw

likelihood scores and overall the species tree was the most common topology for each quartet

(Fig 6; Table S8). Furthermore, the topology test clearly showed that most genes were

uninformative for resolving the relationships among the major groups of Amaranthaceae s.l.

Across all ten quartets we found that most genes had very low TC scores (for any single

node the maximum TC value is 1; Supplemental Fig. S7), showing that individual gene trees had

also large conflict among bootstrap replicates, which is also a signal of uninformative genes and

is concordant with the AU topology test results. Additionally, the linear models did not show any

significant correlation between TC scores and alignment length, GC content or alignment gapless

(Table S9), suggesting that filtering genes by any of these criteria are unlikely to increase the

information content of the dataset.

Species network analyses followed by model selection using each of the four-taxon

datasets showed that in seven out of the ten total quartets, the network with one hybridization

event was a better model than any bifurcating tree topology. However, each of the best three

networks from PhyloNet had very close likelihood scores and no significant **Δ**AICc among them.

For the remaining three quartets the most common bifurcating tree (H0; C1PA, C1BP, PBC2)

was the best model (Table 2; Figs 6, S6, S8).

761 **TABLE 2.** Model selection between quartet tree topologies and species networks. Trees correspond to each of the three possible quartet

762 topologies where H0 is the ASTRAL quartet species tree. Networks correspond to the best three networks for searches with one

763 hybridization event allowed.

Quartet[a]	Topology[b]	ln(L)	Parameters	Number of loci	AICc	ΔAICc	BIC	ΔBIC
BC1A								
	H0	-9014.809786	5	8258	18049.62684	24.73436754	18074.71426	14.70279692
	H1	-9072.456373	5	8258	18164.92002	140.0275408	18190.00743	129.9959702
	H2	-9073.888783	5	8258	18167.78484	142.8923611	18192.87225	132.8607905
	Net 1	**-8998.43945**	**7**	**8258**	**18024.89248**	**0**	**18060.01146**	**0**
	Net 2	-8998.439526	7	8258	18024.89263	0.000151947	18060.01162	0.000151947
	Net 3	-8998.441478	7	8258	18024.89653	0.004056302	18060.01552	0.004056302
ABC2								
	H0	-8516.854413	5	7811	17053.71651	12.87079823	17078.52527	2.950887757
	H1	-8581.563051	5	7811	17183.13379	142.2880731	17207.94254	132.3681626
	H2	-8582.670875	5	7811	17185.34944	144.5037223	17210.15819	134.5838118
	Net 1	**-8506.415681**	**7**	**7811**	**17040.84572**	**0**	**17075.57438**	**0**
	Net 2	-8506.415769	7	7811	17040.84589	0.000176519	17075.57456	0.000176519
	Net 3	-8506.42071	7	7811	17040.85577	0.010057548	17075.58444	0.010057548
BC1C2								
	H0	-9140.191425	5	8385	18300.39001	156.347016	18325.55385	146.2848258
	H1	-9201.981045	5	8385	18423.96925	279.9262567	18449.13309	269.8640665
	H2	-9214.405292	5	8385	18448.81775	304.7747517	18473.98158	294.7125615

Net 1	**-9058.014812**	**7**	**8385**	**18144.04299**	**0**	**18179.26902**	**0**
Net 2	-9058.019338	7	8385	18144.05205	0.009052497	18179.27807	0.009052497
Net 3	-9058.024046	7	8385	18144.06146	0.018468011	18179.28749	0.018468011
C1PA							
H0	**-8932.927759**	**5**	**8134**	**17885.8629**	**0**	**17910.87456**	**0**
H1	-8936.145955	5	8134	17892.29929	6.436391285	17917.31095	6.436391285
H2	-8936.481125	5	8134	17892.96963	7.106730999	17917.98129	7.106730999
Net 1	-8932.077808	7	8134	17892.1694	6.306498884	17927.18227	16.30771403
Net 2	-8932.078011	7	8134	17892.16981	6.306905172	17927.18268	16.30812032
Net 3	-8932.078714	7	8134	17892.17121	6.308310587	17927.18408	16.30952573
PAC2							
H0	-8530.661274	5	7784	17081.33026	40.10000797	17106.12168	30.18704595
H1	-8552.9448	5	7784	17125.89731	84.66706025	17150.68873	74.75409823
H2	-8548.291438	5	7784	17116.59059	75.36033576	17141.382	65.44737374
Net 1	**-8506.607925**	**7**	**7784**	**17041.23025**	**0**	**17075.93463**	**0**
Net 2	-8506.609795	7	7784	17041.23399	0.00373969	17075.93837	0.00373969
Net 3	-8506.618966	7	7784	17041.25233	0.02208072	17075.95671	0.02208072
C1C2P							
H0	-9119.250871	5	8341	18258.50894	12.50997925	18283.64643	2.458344441
H1	-9163.685997	5	8341	18347.37919	101.38023	18372.51669	91.32859519
H2	-9164.83263	5	8341	18349.67246	103.6734974	18374.80995	93.62186263
Net 1	**-9108.992761**	**7**	**8341**	**18245.99896**	**0**	**18281.18809**	**0**
Net 2	-9108.994383	7	8341	18246.00221	0.003244509	18281.19133	0.003244509

	Net 3	-9108.994843	7	8341	18246.00313	0.0041636	18281.19225	0.0041636
C1C2A								
	H0	-8447.623029	5	7756	16915.2538	63.6063012	16940.02717	53.70057058
	H1	-8520.509174	5	7756	17061.02609	209.378593	17085.79946	199.4728624
	H2	-8522.764578	5	7756	17065.5369	213.889401	17090.31027	203.9836704
	Net 1	**-8411.816521**	**7**	**7756**	**16851.6475**	**0**	**16886.3266**	**0**
	Net 2	-8411.819912	7	7756	16851.65428	0.006781956	16886.33338	0.006781956
	Net 3	-8411.820308	7	7756	16851.65507	0.007573446	16886.33417	0.007573446
ABP								
	H0	-9008.115816	5	8206	18036.23895	3.307596079	18061.29474	-6.711300872
	H1	-9015.941176	5	8206	18051.88967	18.95831519	18076.94546	8.939418238
	H2	-9014.738462	5	8206	18049.48424	16.55288764	18074.54003	6.533990688
	Net 1	**-9002.458846**	**7**	**8206**	**18032.93135**	**0**	**18068.00604**	**0**
	Net 2	-9002.460142	7	8206	18032.93395	0.002592568	18068.00863	0.002592568
	Net 3	-9002.464397	7	8206	18032.94246	0.011102577	18068.01714	0.011102577
C1BP								
	H0	**-9557.910518**	**5**	**8793**	**19135.82787**	**0**	**19161.22959**	**0**
	H1	-9661.475396	5	8793	19342.95762	207.1297559	19368.35935	207.1297559
	H2	-9661.009687	5	8793	19342.0262	206.1983365	19367.42793	206.1983365
	Net 1	-9556.24034	7	8793	19140.49343	4.665563813	19176.05266	14.82306554
	Net 2	-9556.243036	7	8793	19140.49882	4.670955519	19176.05805	14.82845724
	Net 3	-9556.246261	7	8793	19140.50527	4.677405326	19176.0645	14.83490705
PBC2								
	H0	**-9158.309463**	**5**	**8379**	**18336.62609**	**0**	**18361.78635**	**0**

H1	-9206.127177	5	8379	18432.26152	95.63542753	18457.42177	95.63542753
H2	-9205.933131	5	8379	18431.87343	95.24733612	18457.03368	95.24733612
Net 1	-9158.016519	7	8379	18344.04642	7.42032489	18379.26742	17.48107897
Net 2	-9158.017286	7	8379	18344.04795	7.421858749	18379.26896	17.48261282
Net 3	-9158.017377	7	8379	18344.04813	7.422042036	18379.26914	17.48279611

764 [a]Each quartet is named following the species tree topology, where the first two are sister. A = Amaranthaceae. s.s. (*Amaranthus*

765 *hypochondriacus*), B = Betoideae (*Beta vulgaris*), C1 = Chenopods I (*Chenopodium quinoa*), C2 = Chenopods II (*Caroxylum*

766 *vermiculatum*), P = Polycnemoideae (*Polycnemum majus*).

767 [b]All quartet tree topologies can be found in Figure 6 and quartet network topologies in Figure S8.

The ABBA/BABA test results showed a significant signal of introgression within each of

the ten quartets (Table 3). The possible introgression was detected between six out of the ten

possible pairs of taxa. Potential introgression between Betoideae and Amaranthaceae s.s.,

'Chenopods I' or 'Chenopods II', and between 'Chenopods I' and Polycnemoideae was not

detected.

TABLE 3. ABBA/BABA test results of Amaranthaceae s.l. five main groups quartets.

Quartet (H0)[a]	Number of loci	Sites in alignment	ABBA	BABA	Raw D-statistic	Z-score	P-value	Introgression direction
BC1A[b]	8258	12778649	287226	254617	0.06018164	41.1085	≤ 0.001	A⇔C1
ABC2	7811	12105324	252772	376755	-0.1969463	124.4161	≤ 0.001	A⇔C2
BC1C2	8385	13192317	306570	258349	0.08535914	54.59751	≤ 0.001	C1⇔C2
C1PA[b]	8134	12635201	342350	286813	0.08827124	64.62297	≤ 0.001	A⇔P
PAC2	7784	12049734	344726	405627	-0.08116313	42.88069	≤ 0.001	C2⇔P
C1C2P[b]	8341	13127397	445384	276652	0.2336892	136.0151	≤ 0.001	C2⇔P
C1C2A[b]	7756	12114778	396219	292561	0.1504951	101.3243	≤ 0.001	A⇔C2
ABP	8206	12622625	276319	312060	-0.06074486	36.64264	≤ 0.001	A⇔P
C1BP[b]	8793	13712853	273286	261620	0.02180944	18.08364	≤ 0.001	B⇔P
PBC2	8379	13074019	217549	415616	-0.3128205	196.8972	≤ 0.001	C2⇔P

[a]Each quartet is named following the species tree topology, where the first two are sister. A =

Amaranthaceae. s.s. (*Amaranthus hypochondriacus*), B = Betoideae (*Beta vulgaris*), C1 =

Chenopods I (*Chenopodium quinoa*), C2 = Chenopods II (*Caroxylum vermiculatum*), P =

Polycnemoideae (*Polycnemum majus*). H0 topologies can be found in Figure 6

[b]Quartet compatible with the complete 105-taxon species trees

The synteny analysis between the diploid ingroup reference genome *Beta vulgaris* and

the diploid outgroup reference genome *Mesembryanthemum crystallinum* recovered 22,179 (out

of 52,357) collinear genes in 516 syntenic blocks. With the collinear ortholog pair information,

we found that of the 8,258 orthologs of the BC1A quartet 6,941 contained orthologous genes

within 383 syntenic blocks. The distribution of the BC1A quartet topologies along the

chromosomes of *Beta vulgaris* did not reveal any spatial clustering along the chromosomes (Fig.

S9).

799   *Assessment of substitutional saturation, codon usage bias, compositional heterogeneity,*

800   *and sequence evolution model misspecification*

We assembled a second 11-taxon(tree) dataset that included 5,936 genes and a minimum of 300

bp (aligned length after removal of low occupancy columns) and no missing taxon. The

saturation plots of uncorrected and predicted genetic distances showed that the first and second

codon position are unsaturated (y = 0.8841002x), while the slope of the third codon positions (y

= 0.5710071x) showed a clear signal of saturation (Fig. S10). The correspondence analyses of

RSCU show that some codons are more frequently used in different species, but overall the

codon usage seems to be randomly dispersed among all species and not clustered by clade (Fig.

S11). This suggests that the phylogenetic signal is unlikely to be driven by differences in codon

usage bias among clades. Furthermore, 549 (~9%) genes showed signal of compositional

heterogeneity ($p < 0.05$) (Table S10). The topology and support (LPP = 1.0) for all branches was

the same for the ASTRAL species trees obtained from the different data schemes while

accounting for saturation, codon usage, compositional heterogeneity, and model of sequence

evolution, and was also congruent with the ASTRAL species tree and concatenated ML from the

full-taxon analyses (Fig. 7). In general, the proportion of gene trees supporting each bipartition

remained the same in every analysis and showed high levels of conflict among the main clades of

Amaranthaceae s.l. (Fig 7). Gene trees inferred accounting for selection of model of sequence

evolution had higher bootstrap support resulting in higher proportion of both concordant and

discordant trees (Fig 7b, 7d, 7e), but the proportion among them is the same as in the gene trees

that used a single model of sequence evolution (Fig 7a–7c).

FIGURE 7. ASTRAL species trees from the 11-taxon(net) dataset estimated from gene trees

inferred using multiple data schemes. a) Gene trees inferred with RAxML with a GTR-GAMMA

model. b) Gene trees inferred with IQ-tree allowing for automatic model selection of sequence

evolution. c) Gene trees inferred with RAxML with a GTR-GAMMA model and removal of

genes that had signal of compositional heterogeneity. d) Gene trees inferred with IQ-tree

allowing for automatic model selection of sequence evolution and removal of genes that had

signal of compositional heterogeneity. a–d) Gene trees were inferred with no partition, codon

partition (first and second codon, and third codon) and, only first and second codon positions

(third codon position removed and no partition). Gene trees were inferred using codon

alignments with standard nucleotide coding, and alignments with degenerated coding of the first

and third codon positions. e) All gene trees and gene trees after removal of genes that had signal

of compositional heterogeneity, inferred with IQ-tree using amino acid sequences allowing for

automatic model selection of sequence evolution. Pie charts on nodes present the proportion of

gene trees that support that clade (blue), the proportion that support the main alternative

bifurcation (green), the proportion that support the remaining alternatives (red), and the

proportion (conflict or support) that have < 50% bootstrap support (gray).

839   *Polytomy test*

840   The ASTRAL polytomy test resulted in the same bifurcating species tree for the 11-taxon(tree)

841   dataset and rejected the null hypothesis that any branch is a polytomy ($p < 0.01$ in all cases).

842   These results were identical when using gene trees with collapsed branches.

844   *Coalescent simulations*

The distribution of tree-to-tree distances of the empirical and simulated gene trees to the species

tree largely overlapped in all seven partition schemes tested (Fig. 8), suggesting that ILS alone

was able to largely account for the gene tree heterogeneity seen in the 11-taxon(tree) dataset. The

least overlap between empirical vs. simulated gene trees was observed in the dataset that only

included the first and second codon positions in CDS, and in the amino acid dataset. This can be

attributed to higher gene tree inference error due to removal of informative sites from the third

codon position.

FIGURE 8. Distribution of tree-to-tree distances from empirical gene trees and species tree versus

coalescent simulation. Simulations were carried out using the ASTRAL species trees from the

11-taxon(tree) dataset estimated from gene trees inferred using seven data schemes. Species trees

used for the coalescent simulation can be seen in Figure 9.

861   *Test of anomaly zone*

The anomaly zone limit calculations using species trees from the 11-taxon(tree) dataset revealed

that two pairs of internodes in the Amaranthaceae s.l. species tree fell into the anomaly zone.

These internodes are characterized by having very short branches relative to the rest of the tree.

The branch lengths among species trees from the seven different data schemes varied among the

trees, but the same internodes were identified under the anomaly zone in all cases. The first pair

of internodes is located between the clade comprised of all Amaranthaceae s.l. and the clade that

includes Chenopods I, Chenopods II, and Betoideae. The second pair of internodes is located

between the clade that includes Chenopods I, Chenopods II, and Betoideae and the clade

composed of Chenopods I and Chenopods II (Table 4, Fig. 9).

TABLE 4. Anomaly zone limit calculations in 11-taxon species trees. Bold rows show pair of

internodes in the anomaly zone when $y < a(x)$.

Species tree[a]	Clade (x)[b]	Clade (y)[b]	x	y	a(x)
No partition	(C1, C2)	(C1)	0.1467	2.722	0.1799
	(C1, C2)	(C2)	0.1467	2.1102	0.1799
	((C1, C2), B)	**(C1, C2)**	**0.1045**	**0.1467**	**0.3084**
	((C1, C2), B)	(B)	0.1045	2.6081	0.3084
	(((C1, C2), B), P)	**((C1, C2), B)**	**0.0846**	**0.1045**	**0.4003**
	(((C1, C2), B), P)	(P)	0.0846	3.5424	0.4003
Codon partition	(C1, C2)	(C1)	0.1433	2.6766	0.1882
	(C1, C2)	(C2)	0.1433	2.0655	0.1882
	((C1, C2), B)	**(C1, C2)**	**0.0931**	**0.1433**	**0.3572**
	((C1, C2), B)	(B)	0.0931	2.5862	0.3572
	(((C1, C2), B), P)	**((C1, C2), B)**	**0.0734**	**0.0931**	**0.4669**
	(((C1, C2), B), P)	(P)	0.0734	3.5366	0.4669
Codon 1&2	(C1, C2)	(C1)	0.118	1.3092	0.26
	(C1, C2)	(C2)	0.118	1.7645	0.26
	((C1, C2), B)	**(C1, C2)**	**0.1009**	**0.118**	**0.3231**
	((C1, C2), B)	(B)	0.1009	1.6229	0.3231
	(((C1, C2), B), P)	**((C1, C2), B)**	**0.0673**	**0.1009**	**0.5102**

	(((C1, C2), B), P)	(P)	0.0673	2.3625	0.5102
	(C1, C2)	(C1)	0.1439	2.1199	0.1865
	(C1, C2)	(C2)	0.1439	2.7013	0.1865
No partition - Model selection	**((C1, C2), B)**	**(C1, C2)**	**0.1079**	**0.1439**	**0.2951**
	((C1, C2), B)	(B)	0.1079	2.5846	0.2951
	(((C1, C2), B), P)	**((C1, C2), B)**	**0.0803**	**0.1079**	**0.4242**
	(((C1, C2), B), P)	(P)	0.0803	3.58	0.4242
	(C1, C2)	(C1)	0.1427	2.6398	0.1896
	(C1, C2)	(C2)	0.1427	2.0911	0.1896
Codon partition - Model selection	**((C1, C2), B)**	**(C1, C2)**	**0.0945**	**0.1427**	**0.351**
	((C1, C2), B)	(B)	0.0945	2.6252	0.351
	(((C1, C2), B), P)	**((C1, C2), B)**	**0.0721**	**0.0945**	**0.4758**
	(((C1, C2), B), P)	(P)	0.0721	3.5978	0.4758
	(C1, C2)	(C1)	0.1232	1.9043	0.2432
	(C1, C2)	(C2)	0.1232	1.415	0.2432
Codon 1&2 - Model selection	**((C1, C2), B)**	**(C1, C2)**	**0.1024**	**0.1232**	**0.317**
	((C1, C2), B)	(B)	0.1024	1.7399	0.317
	(((C1, C2), B), P)	**((C1, C2), B)**	**0.07**	**0.1024**	**0.4906**
	(((C1, C2), B), P)	(P)	0.07	2.5666	0.4906
	(C1, C2)	(C1)	0.115	1.887	0.269
	(C1, C2)	(C2)	0.115	1.317	0.269
Amino Acid - Model selection	**((C1, C2), B)**	**(C1, C2)**	**0.122**	**0.115**	**0.247**
	((C1, C2), B)	(B)	0.122	1.744	0.247
	(((C1, C2), B), P)	**((C1, C2), B)**	**0.077**	**0.122**	**0.444**
	(((C1, C2), B), P)	(P)	0.077	2.471	0.444

874 [a]Species tree topologies can be found in Figure 7.

875 [b] B = Betoideae (*Beta vulgaris*), C1 = Chenopods I (*Chenopodium quinoa*), C2 = Chenopods II

876 (*Caroxylum vermiculatum*), P = Polycnemoideae (*Polycnemum majus*).

FIGURE 9. ASTRAL species trees from the 11-taxon(tree) dataset estimated from individual gene trees inferred with seven data schemes. Number next or above branches represent branch length in coalescent units. Colored branches represent pairs of internodes that fall in the anomaly zone (see Table 4 for anomaly zone limits).

The gene tree counts showed that the species tree was not the most common gene tree topology in four of the seven data schemes analyzed, as expected for the anomaly zone (Fig. S12). When gene trees were inferred with no partition or partitioned by codon, the species tree was the fourth most common gene tree topology (119 out of 5,936 gene trees), while the most common gene tree topologies occurred between 170 and 149 times (Fig. 10). Similar patterns were identified for gene trees inferred while accounting from model of sequence evolution selection (Fig. 10). Interestingly, for the gene tree sets inferred using only the first and second codons, and amino acids, the species tree was the most common topology.

FIGURE 10. Gene tree counts (left) of the four most common topologies (right) of 11-taxon(tree)

dataset inferred with seven data schemes. Gene trees that do not support the monophyly of any of

the five major clades were ignored.

902   **DISCUSSION**

Using a phylotranscriptomic dataset in combination with reference genomes representing major

clades, we have shown the prevalence of gene tree discordance in the backbone phylogeny of

Amaranthaceae s.l. Interestingly, we found that this discordance is also present within the

chloroplast dataset. Despite the strong signal of gene tree discordance, we were able to identify

five well-supported major clades within Amaranthaceae s.l. that are congruent with morphology

and previous taxonomic treatments of the group. Using multiple phylogenetic tools and

simulations we comprehensively tested for processes that might have contributed to the gene tree

discordance in Amaranthaceae s.l. Phylogenetic network analyses and ABBA-BABA tests both

supported multiple reticulation events among the five major clades in Amaranthaceae s.l. At the

same time, the patterns of gene tree discordance among these clades can also largely be

explained by uninformative gene trees and ILS. We found evidence that three consecutive short

internal branches produce anomalous trees contributing to the discordance. Molecular evolution

model misspecification (i.e. substitutional saturation, codon usage bias, or compositional

heterogeneity) was less likely to account for the gene tree discordance. Taken together, no single

source can confidently be pointed out to account for the strong signal of gene tree discordance,

suggesting that the discordance results primarily from ancient and rapid lineage diversification.

Furthermore, the backbone of Amaranthaceae s.l. and remains —and probably will remain—

unresolved even with genome-scale data. Our work highlights the need to test for multiple

sources of conflict in phylogenomic analyses and provide a set of recommendations moving

forward in resolving ancient and rapid diversification.

Five well-supported major clades in Amaranthaceae s.l.

Both our nuclear and chloroplast datasets strongly supported five major clades within

Amaranthaceae s.l.: Amaranthaceae s.s, 'Chenopods I', 'Chenopods II', Betoideae, and

Polycnemoideae (Figs. 2 & 4). We recovered Amaranthaceae s.s., Betoideae, and

Polycnemoideae as monophyletic, which is consistent with morphology and the most recent

molecular analyses of these lineages (Hohmann et al. 2006; Masson and Kadereit 2013; Di

Vincenzo et al. 2018). In the case of Chenopodiaceae s.s., the nuclear analyses (Fig. 2) suggested

the monophyly of this previously segregated family, but gene tree discordance analyses revealed

high levels of conflict among two well-defined clades (Fig. 2), 'Chenopods I' and 'Chenopods

II'. Moreover, the chloroplast analyses did not support the monophyly of Chenopodiaceae s.s.

While we also find evidence of gene tree discordance in the backbone cpDNA phylogeny (see

below), a sister relationship between 'Chenopods I' and Betoideae had strong QS support

(0.84/0.88/0.94; Fig. 4). Weak support and/or conflicting topologies along the backbone on the

Amaranthaceae s.l. characterize all previous molecular studies of the lineage (Fig. 1), even with

hundreds of loci (Walker et al. 2018). On the other hand, all studies support the five major clades

found in our analysis.

For the sake of taxonomic stability, we therefore suggest retaining Amaranthaceae s.l.

sensu APG IV (The Angiosperm Phylogeny Group et al. 2016), which includes the previously

recognized Chenopodiaceae. Here we recognize five subfamilies within Amaranthaceae s.l.:

Amaranthoideae representing Amaranthaceae s.s. (incl. Gomphrenoideae Schinz), Betoideae

Ulbr., Chenopodioideae represented as 'Chenopods I' here (incl. Corispermoideae Ulbr.),

Polycnemoideae Ulbr., and Salicornioideae Ulbr. represented by 'Chenopods II' (incl.

Salsoloideae Ulbr., Suaedoideae Ulbr. and Camphorosmoideae A.J. Scott). The stem ages of

these five subfamilies date back to the early Tertiary (Paleocene, Fig. 3) which agrees with dates

based on chloroplast markers (Kadereit et al. 2012; Di Vincenzo et al. 2018; Yao et al. 2019).

Due to the gene tree discordance along the backbone, the geographic origin of Amaranthaceae

s.l. remains ambiguous.

953   *Gene tree discordance detected among chloroplast genes*

Our concatenation-based chloroplast phylogeny (Fig. 4) retrieved the same five major clades of

Amaranthaceae s.l. as in the nuclear phylogeny, but the relationships among the major clades are

incongruent with the nuclear phylogeny (Fig. 2). Cytonuclear discordance is a well-known

process in plants and it has been traditionally attributed to reticulate evolution (Rieseberg and

Soltis 1991; Sang et al. 1995; Soltis and Kuzoff 1995). Such discordance continues to be treated

as evidence in support of hybridization in more recent phylogenomic studies that assume the

chloroplast to be a single, linked locus (e.g. Folk et al. 2017; Vargas et al. 2017; Morales-Briones

et al. 2018b; Lee-Yaw et al. 2019). However, cytonuclear discordance can also be attributed to

other processes like ILS (Doyle 1992; Ballard and Whitlock 2004). Recent work shows that

chloroplast protein-coding genes may not necessarily act as a single locus, and high levels of tree

conflict has been detected (Gonçalves et al. 2019; Walker et al. 2019).

In Amaranthaceae s.l., previous studies based on chloroplast protein-coding genes or

introns (Kadereit et al. 2003; Müller and Borsch 2005; Hohmann et al. 2006; Kadereit et al.

2017) resulted in different relationships among the five main clades and none in agreement with

our 76-gene phylogeny. Our conflict and QS analyses of the chloroplast dataset (Fig. 4; Figs S4–

S5) revealed strong signals of gene tree discordance among the five major clades of

Amaranthaceae s.l. The strong conflicting signal in the chloroplast genome may be attributed to

heteroplasmy and difference in individual gene phylogenetic information (Walker et al. 2019),

although the exact sources of conflict are yet to be clarified (Gonçalves et al. 2019). Unlike the

results found by Walker et al. (2019), nodes showing conflicting signals in individual gene trees

in our dataset were mostly highly supported (i.e. BS $\geq$ 70, Fig S4), suggesting that low

phylogenetic information is not the source of conflict in our chloroplast dataset.

Our results support previous studies showing RNA-seq data can be a reliable source for

plastome assembly (Smith 2013; Osuna-Mascaró et al. 2018). While the approach has been used

for deep-scale phylogenomic reconstruction in green plants (Gitzendanner et al. 2018), at present

extracting plastome data is not part of routine phylotranscriptomic pipelines. RNA-seq libraries

can contain some genomic DNA due to incomplete digestion during RNA purification (Smith

2013) and given the AT-rich nature of plastomes, this allows plastome DNA to survive the poly-

A selection during mRNA enrichment (Schliesky et al. 2012). However, our results showed that

Amaranthaceae s.l. cpDNA assemblies came from RNA rather than DNA contamination

regardless of library preparation strategies. Similarly, Osuna-Mascaró et al. (2018) also found

highly similar plastome assemblies (i.e. general genome structure, and gene number and

composition) from RNA-seq and genomic libraries, supports the idea that plastome genomes are

fully transcribed in photosynthetic eukaryotes (Shi et al. 2016). Here we implemented additional

steps to the Yang and Smith (2014) pipeline to filter chloroplast and mitochondrial reads prior to

de novo transcriptome assembly, which allowed us to assemble plastome sequences from RNA-

seq libraries, build a plastome phylogeny, and compare it to gene trees constructed from nuclear

genes. Furthermore, the backbone topology of our cpDNA tree built mainly from RNA-seq data

(97 out of 105 samples) was consistent with a recent complete plastome phylogeny of

Caryophyllales (Yao et al. 2019), showing the potential value of using cpDNA from RNA-seq

data. Nonetheless, RNA editing might be problematic when combining samples from RNA and

DNA, especially when trying to resolve phylogenetic relationships among closely related

species.

998   *Hybridization*

Rapid advances have been made in recent years in developing methods to infer species networks

in the presence of ILS (reviewed in Elworth et al. 2019). These methods have been increasingly

used in phylogenetic studies (e.g. Marcussen et al. 2014; Wen et al. 2016a; Copetti et al. 2017);

Morales-Briones et al. 2018a; Crowl et al. 2019). To date, however, species network inference is

still computationally intensive and limited to a small number of species and a few hybridization

events (Hejase and Liu 2016; but see Hejase et al. 2018 and Zhu et al. 2019). Furthermore,

studies evaluating the performance of different phylogenetic network inference approaches are

scarce and restricted to simple hybridization scenarios. (Kamneva and Rosenberg 2017) showed

that likelihood methods like Yu et al. (2014) are often robust to ILS and gene tree error when

symmetric hybridization (equal genetic contribution of both parents) events are considered, and

while it usually does not overestimate hybridization events, it fails to detect skewed

hybridization (unequal genetic contribution of both parents) events in the presence of significant

ILS. Methods developed to scale to larger numbers of species and hybridizations like the ones

using pseudo-likelihood approximations (i.e. Solís-Lemus and Ané 2016; Yu and Nakhleh 2015)

are yet to be evaluated independently, but in the case of Yu and Nakhleh (2015), a method based

on rooted triples, it has been shown that this method cannot distinguish the correct network when

other networks can produce the same set of triples (Yu and Nakhleh 2015). The result of our 11-

taxon(net) phylogenetic analysis using a pseudo-likelihood approach detected up to five

hybridization events involving all five major clades of Amaranthaceae s.l. (Fig. 5). Model

selection, after calculating the full likelihood of the obtained networks, chose the 5-reticulation

species as the best model. Also, any species network had a better score than a bifurcating tree

(Table 1). However, a further look of these hybridization events by breaking the 11-taxon dataset

into ten quartets showed that full likelihood networks searches with up to one hybridization event

are indistinguishable from each other (Table 2), resembling a random gene tree distribution. This

pattern can probably be explained by the high levels of gene tree discordance and lack of

phylogenetic signal in the inferred quartet gene trees (Fig. 6), suggesting that the 11-taxon(net)

network searches can potentially overestimate reticulation events due to high levels of gene tree

error or ILS.

Using the D-Statistic (Green et al. 2010; Durand et al. 2011) we also found signals of

introgression in seven possible directions among the five main groups of Amaranthaceae s.l.

(Table 3). The inferred introgression events agreed with at least one of the reticulation scenarios

from the phylogenetic network analysis. However, the D-Statistic did not detect any

introgression that involves Betoideae, which was detected in the phylogenetic network analysis

with either four or five reticulations events. The D-Statistic has been shown to be robust to a

wide range of divergence times, but it is sensitive to relative population size (Zheng and Janke

2018), which agrees with the notion that large effective population sizes and short branches

increase the chances of ILS (Pamilo and Nei 1988) and in turn can dilute the signal for the D-

Statistic (Zheng and Janke 2018). Recently, Elworth et al. (2018) found that multiple or 'hidden'

reticulations can cause the signal of the D-statistic to be lost or distorted. Furthermore, when

multiple reticulations are present, the traditional approach of subsetting datasets into quartets can

be problematic as it largely underestimates D values (Elworth et al. 2018). Given short internal

branches in the backbone of Amaranthaceae s.l. and the phylogenetic network results showing

multiple hybridizations, it is plausible that our *D*-statistic may be affected by these issues. Our

analysis highlights the uncertainty of relying *D*-statistic as the only test for detecting reticulation

events, especially in cases of ancient and rapid diversification.

1044

1045   *ILS and the Anomaly Zone*

Incomplete Lineage Sorting, or ILS, is ubiquitous in multi-locus phylogenetic datasets. In its

most severe cases ILS produces the 'anomaly zone', defined as a set of short internal branches in

the species tree that produce anomalous gene trees (AGTs) that are more likely than the gene tree

that matches the species tree (Degnan and Rosenberg 2006). Rosenberg (2013) expanded the

definition of the anomaly zone to require that a species tree contain two consecutive internal

branches in an ancestor–descendant relationship in order to produce AGTs. To date, only a few

examples of an empirical anomaly zone have been reported (Linkem et al. 2016; Cloutier et al.

2019). Furthermore, Huang and Knowles (2009) have pointed out that the gene tree discordance

produced from the anomaly zone can also be produced by uninformative gene trees and that for

species trees with short branches the most probable gene tree topology is a polytomy rather than

an AGT. Our results show that the species tree of Amaranthaceae s.l. have three consecutive

short internal branches that lay within the limits of the anomaly zone (i.e. $y < a(x)$; Fig. 9; Table

4). While this is clear evidence that gene tree discordance in Amaranthaceae s.l may be product

of AGTs, it is important to point out that our quartet analysis showed that most quartet gene trees

were equivocal (94–96%; Fig. 6), and were therefore uninformative gene trees. Nonetheless, the

ASTRAL polytomy test rejected a polytomy along the backbone of Amaranthaceae s.l. in any of

the gene tree sets used. While we did not test for polytomies in individual gene trees, our

ASTRAL polytomy test was also carried using gene trees with branches collapsed if they had

<75% bootstrap support, and obtained the same species tree with polytomy being rejected.

Furthermore, we found that for most of the partition schemes tested, the species tree is not the

most frequent gene tree (Fig. 10). The distribution of gene tree frequency in combination with

short internal branches in the species tree supports the presence of an anomaly zone in

Amaranthaceae s.l.

1069

1070   *Considerations in distinguishing sources of gene tree discordance*

With the frequent generation of phylogenomic datasets, the need to explore and disentangle gene

tree discordance has become a fundamental step to understand the phylogenetic relationships of

recalcitrant groups across the Tree of Life. Recently, development of tools to identify and

visualize gene tree discordance has received great attention (e.g. Salichos et al. 2014; Smith et al.

2015; Huang et al. 2016; Pease et al. 2018). New tools have facilitated the detection of conflict,

which has led to the development of downstream phylogenetic analyses to attempt to

characterize it. Although exploring sources of conflicting signal in phylogenomic data is now

common, this is typically focused on data filtering approaches and its effect on concatenation-

based vs. coalescent-based tree inference methods (e.g. Alda et al. 2019; Mclean et al. 2019;

Roycroft et al. 2019). Methods to estimate species trees from phylogenomic dataset while

accounting for multiple sources of conflict and molecular substitution simultaneously are not

available, but by combining transcriptomes and genomes, we were able to create a rich and dense

dataset to start to tease apart alternative hypotheses concerning the sources of conflict in the

backbone phylogeny of Amaranthaceae s.l. Nonetheless, we could not attribute the strong gene

tree discordance signal to a single main source. Instead, we found that gene tree heterogeneity

observed in Amaranthaceae s.l. is likely to be explained by a combination of processes, including

ILS, hybridization, uninformative genes, and molecular evolution model misspecification, that

might have acted simultaneously and/or cumulatively.

Our results highlight the need to test for multiple sources of conflict in phylogenomic

analyses, especially when trying to resolve phylogenetic relationships in groups with a long

history of phylogenetic conflict. We consider that special attention should be put in data

processing, orthology inference, as well as the informativeness of individual gene trees.

Furthermore, we need to be aware of the strengths and limitations of different phylogenetic

methods and be cautious against relying on a single analysis, for example in the usage of

phylogenetics species networks over coalescent-based species trees (also see Blair and Ané

2019). While the backbone phylogeny of Amaranthaceae s.l. remains difficult to resolve despite

employing genome-scale data, a question emerges whether this is an atypical case, or as we

leverage more phylogenomic datasets and explore gene tree discordance in more detail, we could

find similar patterns in other groups, especially in those that are products of ancient and rapid

lineage diversification (Widhelm et al. 2019). Ultimately, such endeavor will be instrumental in

our fundamental understanding of the biology of the organisms.

1102

1103   SUPPLEMENTARY MATERIAL

1104 Data available from the Dryad Digital Repository: http://dx.doi.org/10.5061/.[NNNN]

1105

1106   ACKNOWLEDGMENTS

The authors thank H. Freitag, J.M. Bena and the Millennium Seed Bank for providing seeds;

Ursula Martiné for assisting with RNA extraction; Alexandra Crum for help revising the

manuscript; X anonymous reviewers for providing helpful comments; the Minnesota

Supercomputing Institute (MSI) at the University of Minnesota for providing access to

computational resources. This work was supported by the University of Minnesota, University of

Michigan, and the US National Science Foundation (DEB 1354048).

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

REFERENCES

1132

1133

Alda F., Tagliacollo V.A., Bernt M.J., Waltz B.T., Ludt W.B., Faircloth B.C., Alfaro M.E.,

Albert J.S., Chakrabarty P. 2019. Resolving Deep Nodes in an Ancient Radiation of

Neotropical Fishes in the Presence of Conflicting Signals from Incomplete Lineage

Sorting. Syst. Biol. 68:573–593.

Arcila D., Ortí G., Vari R., Armbruster J.W., Stiassny M.L.J., Ko K.D., Sabaj M.H., Lundberg J.,

Revell L.J., Betancur-R. R. 2017. Genome-wide interrogation advances resolution of

recalcitrant groups in the tree of life. Nat. Ecol. Evol. 1:0020.

Ballard J.W.O., Whitlock M.C. 2004. The incomplete natural history of mitochondria. Mol. Ecol.

13:729–744.

Bankevich A., Nurk S., Antipov D., Gurevich A.A., Dvorkin M., Kulikov A.S., Lesin V.M.,

Nikolenko S.I., Pham S., Prjibelski A.D., Pyshkin A.V., Sirotkin A.V., Vyahhi N., Tesler

G., Alekseyev M.A., Pevzner P.A. 2012. SPAdes: A New Genome Assembly Algorithm

and Its Applications to Single-Cell Sequencing. J. Comput. Biol. 19:455–477.

Bena M.J., Acosta J.M., Aagesen L. 2017. Macroclimatic niche limits and the evolution of C4

photosynthesis in Gomphrenoideae (Amaranthaceae). Bot. J. Linn. Soc. 184:283–297.

Blackmon H., Adams R.A. 2015 EvobiR: Tools for comparative analyses and teaching

evolutionary biology. doi:10.5281/zenodo.30938

Blair C., Ané C. 2019. Phylogenetic Trees and Networks Can Serve as Powerful and

Complementary Approaches for Analysis of Genomic Data. Syst. Biol. syz056,

https://doi.org/10.1093/sysbio/syz056

Bolger A.M., Lohse M., Usadel B. 2014. Trimmomatic - a flexible trimmer for Illumina

sequence data. Bioinformatics. 30:2112–2120.

Bouckaert R., Heled J. 2014. DensiTree 2: Seeing Trees Through the Forest. BioRxiv. 012401.

Brown J.W., Walker J.F., Smith S.A. 2017. Phyx - phylogenetic tools for unix. Bioinformatics.

33:1886–1888.

Bruen T.C., Philippe H., Bryant D. 2006. A Simple and Robust Statistical Test for Detecting the

Presence of Recombination. Genetics. 172:2665–2681.

Buchfink B., Xie C., Huson D.H. 2015. Fast and sensitive protein alignment using DIAMOND.

Nat. Methods. 12:59–60.

Buckley T.R., Cordeiro M., Marshall D.C., Simon C. 2006. Differentiating between Hypotheses

of Lineage Sorting and Introgression in New Zealand Alpine Cicadas (Maoricicada

Dugdale). Syst. Biol. 55:411–425.

Castresana J. 2000. Selection of Conserved Blocks from Multiple Alignments for Their Use in

Phylogenetic Analysis. Mol. Biol. Evol. 17:540–552.

Chen L.-Y., Morales-Briones D.F., Passow C.N., Yang Y. 2019. Performance of gene expression

analyses using de novo assembled transcripts in polyploid species. Bioinformatics.

btz620, https://doi.org/10.1093/bioinformatics/btz620

Cloutier A., Sackton T.B., Grayson P., Clamp M., Baker A.J., Edwards S.V. 2019. Whole-

Genome Analyses Resolve the Phylogeny of Flightless Birds (Palaeognathae) in the

Presence of an Empirical Anomaly Zone. Syst. Biol. syz019,

https://doi.org/10.1093/sysbio/syz019

Cooper E.D. 2014. Overly simplistic substitution models obscure green plant phylogeny. Trends

Plant Sci. 19:576–582.

Copetti D., Búrquez A., Bustamante E., Charboneau J.L.M., Childs K.L., Eguiarte L.E., Lee S.,

Liu T.L., McMahon M.M., Whiteman N.K., Wing R.A., Wojciechowski M.F., Sanderson

M.J. 2017. Extensive gene tree discordance and hemiplasy shaped the genomes of North

American columnar cacti. Proc. Natl. Acad. Sci. 114:12003–12008.

Cox C.J., Li B., Foster P.G., Embley T.M., Civáň P. 2014. Conflicting Phylogenies for Early

Land Plants are Caused by Composition Biases among Synonymous Substitutions. Syst.

Biol. 63:272–279.

Crowl A.A., Manos P.S., McVay J.D., Lemmon A.R., Lemmon E.M., Hipp A.L. 2019.

Uncovering the genomic signature of ancient introgression between white oak lineages

(*Quercus*). New Phytol. nph.15842, https://doi.org/10.1111/nph.15842

Davidson N.M., Oshlack A. 2014. Corset: enabling differential gene expression analysis for de

novo assembled transcriptomes. Genome Biol. 15:57.

Degnan J.H., Rosenberg N.A. 2006. Discordance of Species Trees with Their Most Likely Gene

Trees. PLoS Genet. 2:e68.

Degnan J.H., Rosenberg N.A. 2009. Gene tree discordance, phylogenetic inference and the

1192   multispecies coalescent. Trends Ecol. Evol. 24:332–340.

1193   Di Vincenzo V., Gruenstaeudl M., Nauheimer L., Wondafrash M., Kamau P., Demissew S.,

1194   Borsch T. 2018. Evolutionary diversification of the African achyranthoid clade

1195   (Amaranthaceae) in the context of sterile flower evolution and epizoochory. Ann. Bot.

1196   122:69–85.

1197   Dohm J.C., Minoche A.E., Holtgräwe D., Capella-Gutiérrez S., Zakrzewski F., Tafer H., Rupp

1198   O., Sörensen T.R., Stracke R., Reinhardt R., Goesmann A., Kraft T., Schulz B., Stadler

1199   P.F., Schmidt T., Gabaldón T., Lehrach H., Weisshaar B., Himmelbauer H. 2014. The

1200   genome of the recently domesticated crop plant sugar beet (Beta vulgaris). Nature.

1201   505:546–549.

1202   van Dongen S.M. 2000. Graph Clustering by Flow Simulation. PhD diss., Ultrecht University.

1203   Doyle J.J. 1992. Gene Trees and Species Trees: Molecular Systematics as One-Character

1204   Taxonomy. Syst. Bot. 17:144.

1205   Duchêne D.A., Bragg J.G., Duchêne S., Neaves L.E., Potter S., Moritz C., Johnson R.N., Ho

1206   S.Y.W., Eldridge M.D.B. 2018. Analysis of Phylogenomic Tree Space Resolves

1207   Relationships Among Marsupial Families. Syst. Biol. 67:400–412.

1208   Durand E.Y., Patterson N., Reich D., Slatkin M. 2011. Testing for Ancient Admixture between

1209   Closely Related Populations. Mol. Biol. Evol. 28:2239–2252.

Eaton D.A.R., Ree R.H. 2013. Inferring Phylogeny and Introgression using RADseq Data: An

Example from Flowering Plants (Pedicularis: Orobanchaceae). Syst. Biol. 62:689–706.

Edwards S.V. 2009. Is A New and General Theory of Molecular Systematics Emerging?

Evolution. 63:1–19.

Edwards S.V., Xi Z., Janke A., Faircloth B.C., McCormack J.E., Glenn T.C., Zhong B., Wu S.,

Lemmon E.M., Lemmon A.R., Leaché A.D., Liu L., Davis C.C. 2016. Implementing and

testing the multispecies coalescent model: A valuable paradigm for phylogenomics. Mol.

Phylogenet. Evol. 94:447–462.

Elworth R.A.L., Allen C., Benedict T., Dulworth P., Nakhleh L.K. 2018. DGEN: A Test Statistic

for Detection of General Introgression Scenarios. WABI.

Elworth R.A.L., Ogilvie H.A., Zhu J., Nakhleh L. 2019. Advances in Computational Methods for

Phylogenetic Networks in the Presence of Hybridization. In: Warnow T., editor.

Bioinformatics and Phylogenetics: Seminal Contributions of Bernard Moret. Cham:

Springer International Publishing. p. 317–360.

Erfan Sayyari, Siavash Mirarab. 2018. Testing for Polytomies in Phylogenetic Species Trees

Using Quartet Frequencies. Genes. 9:132.

Flowers T.J., Colmer T.D. 2015. Plant salt tolerance: adaptations in halophytes. Ann. Bot.

115:327–331.

Folk R.A., Mandel J.R., Freudenstein J.V. 2017. Ancestral Gene Flow and Parallel Organellar

Genome Capture Result in Extreme Phylogenomic Discord in a Lineage of Angiosperms.

Syst. Biol. 66:320-337.

Foster P.G. 2004. Modeling Compositional Heterogeneity. Syst. Biol. 53:485–495.

Fu L., Niu B., Zhu Z., Wu S., Li W. 2012. CD-HIT: accelerated for clustering the next-

generation sequencing data. Bioinformatics. 28:3150–3152.

Galtier N., Daubin V. 2008. Dealing with incongruence in phylogenomic analyses. Philos. Trans.

R. Soc. B Biol. Sci. 363:4023–4029.

Gitzendanner M.A., Soltis P.S., Yi T.-S., Li D.-Z., Soltis D.E. 2018. Plastome Phylogenetics: 30

Years of Inferences Into Plant Evolution. Plastid Genome Evolution. Elsevier. p. 293–

313.

Glémin S., Scornavacca C., Dainat J., Burgarella C., Viader V., Ardisson M., Sarah G., Santoni

S., David J., Ranwez V. 2019. Pervasive hybridizations in the history of wheat relatives.

Sci. Adv. 5:eaav9188.

Gonçalves D.J.P., Simpson B.B., Ortiz E.M., Shimizu G.H., Jansen R.K. 2019. Incongruence

between gene trees and species trees and phylogenetic signal variation in plastid genes.

Mol. Phylogenet. Evol. 138:219–232.

Green R.E., Krause J., Briggs A.W., Maricic T., Stenzel U., Kircher M., Patterson N., Li H., Zhai

W., Fritz M.H.Y., Hansen N.F., Durand E.Y., Malaspinas A.S., Jensen J.D., Marques-

Bonet T., Alkan C., Prufer K., Meyer M., Burbano H.A., Good J.M., Schultz R., Aximu-

Petri A., Butthof A., Hober B., Hoffner B., Siegemund M., Weihmann A., Nusbaum C.,

Lander E.S., Russ C., Novod N., Affourtit J., Egholm M., Verna C., Rudan P., Brajkovic

D., Kucan Z., Gusic I., Doronichev V.B., Golovanova L.V., Lalueza-Fox C., de la Rasilla

M., Fortea J., Rosas A., Schmitz R.W., Johnson P.L.F., Eichler E.E., Falush D., Birney

E., Mullikin J.C., Slatkin M., Nielsen R., Kelso J., Lachmann M., Reich D., Paabo S.

2010. A Draft Sequence of the Neandertal Genome. Science. 328:710–722.

Haas B.J., Papanicolaou A., Yassour M., Grabherr M., Blood P.D., Bowden J., Couger M.B.,

Eccles D., Li B., Lieber M., MacManes M.D., Ott M., Orvis J., Pochet N., Strozzi F.,

Weeks N., Westerman R., William T., Dewey C.N., Henschel R., LeDuc R.D., Friedman

N., Regev A. 2013. De novo transcript sequence reconstruction from RNA-seq using the

Trinity platform for reference generation and analysis. Nat. Protoc. 8:1494–1512.

Hejase H.A., Liu K.J. 2016. A scalability study of phylogenetic network inference methods using

empirical datasets and simulations involving a single reticulation. BMC Bioinformatics.

17:422.

Hejase H.A., VandePol N., Bonito G.M., Liu K.J. 2018. FastNet: Fast and Accurate Statistical

Inference of Phylogenetic Networks Using Large-Scale Genomic Sequence Data. Comp.

Genomics.:242–259.

Hernández-Ledesma P., Berendsohn W.G., Borsch T., Mering S.V., Akhani H., Arias S.,

Castañeda-Noa I., Eggli U., Eriksson R., Flores-Olvera H., Fuentes-Bazán S., Kadereit

G., Klak C., Korotkova N., Nyffeler R., Ocampo G., Ochoterena H., Oxelman B.,

Rabeler R.K., Sanchez A., Schlumpberger B.O., Uotila P. 2015. A taxonomic backbone

for the global synthesis of species diversity in the angiosperm order Caryophyllales.

Willdenowia. 45:281.

Hoang D.T., Chernomor O. 2018. UFBoot2: Improving the Ultrafast Bootstrap Approximation.

Mol. Biol. Evol. 35:518–522.

Hohmann S., Kadereit J.W., Kadereit G. 2006. Understanding Mediterranean-Californian

disjunctions: molecular evidence from Chenopodiaceae-Betoideae. TAXON. 55:67–78.

Holder M.T., Anderson J.A., Holloway A.K. 2001. Difficulties in Detecting Hybridization. Syst.

Biol. 50:978–982.

Huang H., Knowles L.L. 2009. What Is the Danger of the Anomaly Zone for Empirical

Phylogenetics? Syst. Biol. 58:527–536.

Huang W., Zhou G., Marchand M., Ash J.R., Morris D., Van Dooren P., Brown J.M., Gallivan

K.A., Wilgenbusch J.C. 2016. TreeScaper: Visualizing and Extracting Phylogenetic

Signal from Sets of Trees. Mol. Biol. Evol. 33:3314–3316.

Hughes L.C., Ortí G., Huang Y., Sun Y., Baldwin C.C., Thompson A.W., Arcila D., Betancur-R.

R., Li C., Becker L., Bellora N., Zhao X., Li X., Wang M., Fang C., Xie B., Zhou Z.,

Huang H., Chen S., Venkatesh B., Shi Q. 2018. Comprehensive phylogeny of ray-finned

fishes (Actinopterygii) based on transcriptomic and genomic data. Proc. Natl. Acad. Sci.

115:6249–6254.

Jarvis D.E., Ho Y.S., Lightfoot D.J., Schmöckel S.M., Li B., Borm T.J.A., Ohyanagi H., Mineta

K., Michell C.T., Saber N., Kharbatia N.M., Rupper R.R., Sharp A.R., Dally N.,

Boughton B.A., Woo Y.H., Gao G., Schijlen E.G.W.M., Guo X., Momin A.A., Negrão

S., Al-Babili S., Gehring C., Roessner U., Jung C., Murphy K., Arold S.T., Gojobori T.,

Linden C.G.V.D., van Loo E.N., Jellen E.N., Maughan P.J., Tester M. 2017. The genome

of *Chenopodium quinoa*. Nature. 542:307–312.

Jarvis E.D., Mirarab S., Aberer A.J., Li B., Houde P., Li C., Ho S.Y.W., Faircloth B.C., Nabholz

B., Howard J.T., Suh A., Weber C.C., da Fonseca R.R., Li J., Zhang F., Li H., Zhou L.,

Narula N., Liu L., Ganapathy G., Boussau B., Bayzid Md.S., Zavidovych V.,

Subramanian S., Gabaldón T., Capella-Gutiérrez S., Huerta-Cepas J., Rekepalli B.,

Munch K., Schierup M., Lindow B., Warren W.C., Ray D., Green R.E., Bruford M.W.,

Zhan X., Dixon A., Li S., Li N., Huang Y., Derryberry E.P., Bertelsen M.F., Sheldon

F.H., Brumfield R.T., Mello C.V., Lovell P.V., Wirthlin M., Schneider M.P.C.,

Prosdocimi F., Samaniego J.A., Velazquez A.M.V., Alfaro-Núñez A., Campos P.F.,

Petersen B., Sicheritz-Ponten T., Pas A., Bailey T., Scofield P., Bunce M., Lambert D.M.,

Zhou Q., Perelman P., Driskell A.C., Shapiro B., Xiong Z., Zeng Y., Liu S., Li Z., Liu B.,

Wu K., Xiao J., Yinqi X., Zheng Q., Zhang Y., Yang H., Wang J., Smeds L., Rheindt

F.E., Braun M., Fjeldsa J., Orlando L., Barker F.K., Jønsson K.A., Johnson W., Koepfli

K.-P., O'Brien S., Haussler D., Ryder O.A., Rahbek C., Willerslev E., Graves G.R.,

Glenn T.C., McCormack J., Burt D., Ellegren H., Alström P., Edwards S.V., Stamatakis

A., Mindell D.P., Cracraft J., Braun E.L., Warnow T., Jun W., Gilbert M.T.P., Zhang G.

2014. Whole-genome analyses resolve early branches in the tree of life of modern birds.

Science. 346:1320.

Joly S., McLenachan P.A., Lockhart P.J. 2009. A Statistical Approach for Distinguishing

Hybridization and Incomplete Lineage Sorting. Am. Nat. 174:E54–E70.

Kadereit G., Ackerly D., Pirie M.D. 2012. A broader model for C4 photosynthesis evolution in

plants inferred from the goosefoot family (Chenopodiaceae s.s.). Proc. R. Soc. B Biol.

Sci. 279:3304–3311.

Kadereit G., Borsch T., Weising K., Freitag H. 2003. Phylogeny of Amaranthaceae and

Chenopodiaceae and the Evolution of C4 Photosynthesis. Int. J. Plant Sci. 164:959–986.

Kadereit G., Hohmann S., Kadereit J.W. 2006. A synopsis of Chenopodiaceae subfam. Betoideae

and notes on the taxonomy of *Beta*. Willdenowia. 36:9–19.

Kadereit G., Newton R.J., Vandelook F. 2017. Evolutionary ecology of fast seed germination—

A case study in Amaranthaceae/Chenopodiaceae. Perspect. Plant Ecol. Evol. Syst. 29:1–

11.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods.

14:587–589.

Kamneva O.K., Rosenberg N.A. 2017. Simulation-Based Evaluation of Hybridization Network

Reconstruction Methods in the Presence of Incomplete Lineage Sorting. Evol.

Bioinforma. 13:117693431769193.

Katoh K., Standley D.M. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. Mol. Biol. Evol. 30:772–780.

Kearse M., Moir R., Wilson A., Stones-Havas S., Cheung M., Sturrock S., Buxton S., Cooper A.,

Markowitz S., Duran C., Thierer T., Ashton B., Meintjes P., Drummond A. 2012.

Geneious Basic: An integrated and extendable desktop software platform for the

organization and analysis of sequence data. Bioinformatics. 28:1647–1649.

Knowles L.L., Huang H., Sukumaran J., Smith S.A. 2018. A matter of phylogenetic scale:

Distinguishing incomplete lineage sorting from lateral gene transfer as the cause of gene

tree discord in recent versus deep diversification histories. Am. J. Bot. 105:376–384.

Kubatko L.S., Chifman J. 2019. An invariants-based method for efficient identification of hybrid

species from large-scale genomic data. BMC Evol. Biol. 19:112.

Lanfear R., Calcott B., Ho S.Y.W., Guindon S. 2012. PartitionFinder: Combined Selection of

Partitioning Schemes and Substitution Models for Phylogenetic Analyses. Mol. Biol.

Evol. 29:1695–1701.

Langmead B., Salzberg S.L. 2012. Fast gapped-read alignment with Bowtie 2. Nat. Methods.

9:357–359.

Laumer C.E., Fernández R., Lemer S., Combosch D., Kocot K.M., Riesgo A., Andrade S.C.S.,

Sterrer W., Sørensen M.V., Giribet G. 2019. Revisiting metazoan phylogeny with

genomic sampling of all phyla. Proc. R. Soc. B Biol. Sci. 286:20190831.

Lê S., Josse J., Husson F. 2008. FactoMineR : An R Package for Multivariate Analysis. J. Stat.

Softw. 25: 1–18.

Lee-Yaw J.A., Grassa C.J., Joly S., Andrew R.L., Rieseberg L.H. 2019. An evaluation of

alternative explanations for widespread cytonuclear discordance in annual sunflowers

(*Helianthus*). New Phytol. 221:515–526.

Li B., Lopes J.S., Foster P.G., Embley T.M., Cox C.J. 2014. Compositional Biases among

Synonymous Substitutions Cause Conflict between Gene and Protein Trees for Plastid

Origins. Mol. Biol. Evol. 31:1697–1709.

Lightfoot D.J., Jarvis D.E., Ramaraj T., Lee R., Jellen E.N., Maughan P.J. 2017. Single-molecule

sequencing and Hi-C-based proximity-guided assembly of amaranth (*Amaranthus*

hypochondriacus) chromosomes provide insights into genome evolution. BMC Biol.

15:74.

Linkem C.W., Minin V.N., Leaché A.D. 2016. Detecting the Anomaly Zone in Species Trees

and Evidence for a Misleading Signal in Higher-Level Skink Phylogeny (Squamata:

Scincidae). Syst. Biol. 65:465–477.

Liu L., Yu L. 2010. Phybase: an R package for species tree analysis. Bioinformatics. 26:962–

963.

Liu Y., Cox C.J., Wang W., Goffinet B. 2014. Mitochondrial Phylogenomics of Early Land

Plants: Mitigating the Effects of Saturation, Compositional Heterogeneity, and Codon-

Usage Bias. Syst. Biol. 63:862–878.

Maddison W.P. 1997. Gene Trees in Species Trees. Syst. Biol. 46:532–536.

Mai U., Mirarab S. 2018. TreeShrink: fast and accurate detection of outlier long branches in

collections of phylogenetic trees. BMC Genomics. 19:4046.

Marcussen T., Sandve S.R., Heier L., Spannagl M., Pfeifer M., Jakobsen K.S., Wulff B.B.H.,

1371   Steuernagel B., Mayer K.F.X., Olsen O.-A. 2014. Ancient hybridizations among the

1372   ancestral genomes of bread wheat. Science. 345:1250092.

1373   Masson R., Kadereit G. 2013. Phylogeny of Polycnemoideae (Amaranthaceae): Implications for

1374   biogeography, character evolution and taxonomy. TAXON. 62:100–111.

1375   Maureira-Butler I.J., Pfeil B.E., Muangprom A., Osborn T.C., Doyle J.J. 2008. The Reticulate

1376   History of *Medicago* (Fabaceae). Syst. Biol. 57:466–482.

1377   Mclean B.S., Bell K.C., Allen J.M., Helgen K.M., Cook J.A. 2019. Impacts of Inference Method

1378   and Data set Filtering on Phylogenomic Resolution in a Rapid Radiation of Ground

1379   Squirrels (Xerinae: Marmotini). Syst. Biol. 68:298–316.

1380   Meyer B.S., Matschiner M., Salzburger W. 2017. Disentangling Incomplete Lineage Sorting and

1381   Introgression to Refine Species-Tree Estimates for Lake Tanganyika Cichlid Fishes. Syst.

1382   Biol. 66:531–550.

1383   Mirarab S., Bayzid M.S., Warnow T. 2016. Evaluating Summary Methods for Multilocus

1384   Species Tree Estimation in the Presence of Incomplete Lineage Sorting. Syst. Biol.

1385   65:366–380.

1386   Misof B., Liu S., Meusemann K., Peters R.S., Donath A., Mayer C., Frandsen P.B., Ware J.,

1387   Flouri T., Beutel R.G., Niehuis O., Petersen M., Izquierdo-Carrasco F., Wappler T., Rust

1388   J., Aberer A.J., Aspock U., Aspock H., Bartel D., Blanke A., Berger S., Bohm A.,

1389   Buckley T.R., Calcott B., Chen J., Friedrich F., Fukui M., Fujita M., Greve C., Grobe P.,

1390   Gu S., Huang Y., Jermiin L.S., Kawahara A.Y., Krogmann L., Kubiak M., Lanfear R.,

1391   Letsch H., Li Y., Li Z., Li J., Lu H., Machida R., Mashimo Y., Kapli P., McKenna D.D.,

1392   Meng G., Nakagaki Y., Navarrete-Heredia J.L., Ott M., Ou Y., Pass G., Podsiadlowski

1393   L., Pohl H., von Reumont B.M., Schutte K., Sekiya K., Shimizu S., Slipinski A.,

1394   Stamatakis A., Song W., Su X., Szucsich N.U., Tan M., Tan X., Tang M., Tang J.,

1395   Timelthaler G., Tomizuka S., Trautwein M., Tong X., Uchifune T., Walzl M.G.,

1396   Wiegmann B.M., Wilbrandt J., Wipfler B., Wong T.K.F., Wu Q., Wu G., Xie Y., Yang

1397   S., Yang Q., Yeates D.K., Yoshizawa K., Zhang Q., Zhang R., Zhang W., Zhang Y.,

1398   Zhao J., Zhou C., Zhou L., Ziesmann T., Zou S., Li Y., Xu X., Zhang Y., Yang H., Wang

1399   J., Wang J., Kjer K.M., Zhou X. 2014. Phylogenomics resolves the timing and pattern of

1400   insect evolution. Science. 346:763–767.

1401 Morales-Briones D.F., Liston A., Tank D.C. 2018a. Phylogenomic analyses reveal a deep history

1402   of hybridization and polyploidy in the Neotropical genus *Lachemilla* (Rosaceae). New

1403   Phytol. 218:1668–1684.

1404 Morales-Briones D.F., Romoleroux K., Kolář F., Tank D.C. 2018b. Phylogeny and Evolution of

1405   the Neotropical Radiation of *Lachemilla* (Rosaceae): Uncovering a History of Reticulate

1406   Evolution and Implications for Infrageneric Classification. Syst. Bot. 43:17–34.

1407 Moray C., Goolsby E.W., Bromham L. 2016. The Phylogenetic Association Between Salt

1408   Tolerance and Heavy Metal Hyperaccumulation in Angiosperms. Evol. Biol. 43:119–

1409   130.

1410 Mower J.P. 2009. The PREP suite: predictive RNA editors for plant mitochondrial genes,

1411   chloroplast genes and user-defined alignments. Nucleic Acids Res. 37:W253–W259.

Müller K., Borsch T. 2005. Phylogenetics of Amaranthaceae Based on *matK/trnK* Sequence

Data: Evidence from Parsimony, Likelihood, and Bayesian Analyses. Ann. Mo. Bot.

Gard. 92:66–102.

Nguyen L.-T., Schmidt H.A., von Haeseler A., Minh B.Q. 2015. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol. Biol. Evol.

32:268–274.

Osuna-Mascaró C., Rubio de Casas R., Perfectti F. 2018. Comparative assessment shows the

reliability of chloroplast genome assembly using RNA-seq. Sci. Rep. 8:17404.

Pamilo P., Nei M. 1988. Relationships between Gene Trees and Species Trees. Mol. Biol. Evol.

5:568–583.

Paradis E., Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary

analyses in R. Bioinformatics. 35:526–528.

Patro R., Duggal G., Love M.I., Irizarry R.A., Kingsford C. 2017. Salmon provides fast and bias-

aware quantification of transcript expression. Nat. Methods. 14:417–419.

Patterson N., Moorjani P., Luo Y., Mallick S., Rohland N., Zhan Y., Genschoreck T., Webster

T., Reich D. 2012. Ancient Admixture in Human History. Genetics. 192:1065–1093.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2018. Quartet Sampling

distinguishes lack of support from conflicting support in the green plant tree of life. Am.

J. Bot. 105:385–403.

Pease J.B., Hahn M.W. 2015. Detection and Polarization of Introgression in a Five-Taxon

Phylogeny. Syst. Biol. 64:651–662.

Peden J. 1999. Analysis of Codon Usage. PhD diss., University of Nottingham.

Philippe H., Forterre P. 1999. The Rooting of the Universal Tree of Life Is Not Reliable. J. Mol.

Evol. 49:509–523.

Piirainen M., Liebisch O., Kadereit G. 2017. Phylogeny, biogeography, systematics and

taxonomy of Salicornioideae (Amaranthaceae/Chenopodiaceae) – A cosmopolitan, highly

specialized hygrohalophyte lineage dating back to the Oligocene. Taxon. 66:109–132.

Prasanna A.N., Gerber D., Kijpornyongpan T., Aime M.C., Doyle V.P., Nagy L.G. 2019. Model

Choice, Missing Data, and Taxon Sampling Impact Phylogenomic Inference of Deep

Basidiomycota Relationships. syz029, https://doi.org/10.1093/sysbio/syz029

Pruitt K.D., Tatusova T., Maglott D.R. 2007. NCBI reference sequences (RefSeq): a curated

non-redundant sequence database of genomes, transcripts and proteins. Nucleic Acids

Res. 35:D61–D65.

R Core Team. 2019. R: A Language and Environment for Statistical Computing. Vienna,

Austria: R Foundation for Statistical Computing.

Rannala B., Yang Z. 2003. Bayes Estimation of Species Divergence Times and Ancestral

Population Sizes Using DNA Sequences From Multiple Loci. Genetics. 166:1645–1656.

Ranwez V., Douzery E.J.P., Cambon C., Chantret N., Delsuc F. 2018. MACSE v2: Toolkit for

the Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. Mol.

Biol. Evol. 35:2582–2584.

Regier J.C., Shultz J.W., Zwick A., Hussey A., Ball B., Wetzer R., Martin J.W., Cunningham

C.W. 2010. Arthropod relationships revealed by phylogenomic analysis of nuclear

protein-coding sequences. Nature. 463:1079–1083.

Rieseberg L.H., Soltis D.E. 1991. Phylogenetic consequences of cytoplasmic gene flow in plants.

Evol. Trends Plants. 5:65–84.

Robinson D.F., Foulds L.R. 1981. Comparison of phylogenetic trees. Math. Biosci. 53:131–147.

Rosenberg N.A. 2013. Discordance of Species Trees with Their Most Likely Gene Trees: A

Unifying Principle. Mol. Biol. Evol. 30:2709–2713.

Roycroft E.J., Moussalli A., Rowe K.C. 2019. Phylogenomics Uncovers Confidence and

Conflict in the Rapid Radiation of Australo-Papuan Rodents. Syst. Biol. syz044,

https://doi.org/10.1093/sysbio/syz044

Salichos L., Stamatakis A., Rokas A. 2014. Novel Information Theory-Based Measures for

Quantifying Incongruence among Phylogenetic Trees. Mol. Biol. Evol. 31:1261–1271.

Sang T., Crawford D.J., Stuessy T.F. 1995. Documentation of reticulate evolution in peonies

(Paeonia) using internal transcribed spacer sequences of nuclear ribosomal DNA:

implications for biogeography and concerted evolution. Proc. Natl. Acad. Sci. 92:6813–

6817.

Sayyari E., Mirarab S. 2016. Fast Coalescent-Based Computation of Local Branch Support from

Quartet Frequencies. Mol. Biol. Evol. 33:1654–1668.

Schliep K.P. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27:592–593.

Schliesky S., Gowik U., Weber A.P.M., Bräutigam A. 2012. RNA-Seq Assembly – Are We

There Yet? Front. Plant Sci. 3.

Schwarz G. 1978. Estimating the Dimension of a Model. Ann. Stat. 6:461–464.

Sharp P.M., Li W.-H. 1986. An evolutionary perspective on synonymous codon usage in

unicellular organisms. J. Mol. Evol. 24:28–38.

Shi C., Wang S., Xia E.-H., Jiang J.-J., Zeng F.-C., Gao L.-Z. 2016. Full transcription of the

chloroplast genome in photosynthetic eukaryotes. Sci. Rep. 6:30135.

Shimodaira H. 2002. An Approximately Unbiased Test of Phylogenetic Tree Selection. Syst.

Biol. 51:492–508.

Shimodaira H., Hasegawa M. 2001. CONSEL: for assessing the confidence of phylogenetic tree

selection. Bioinformatics. 17:1246–1247.

Smith D.R. 2013. RNA-Seq data: a goldmine for organelle research. Brief. Funct. Genomics.

12:454–456.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals

conflict, concordance, and gene duplications with examples from animals and plants.

BMC Evol. Biol. 15:745.

Smith S.A., O'Meara B.C. 2012. treePL: divergence time estimation using penalized likelihood

1489   for large phylogenies. Bioinformatics. 28:2689–2690.

1490   Smith-Unna R., Boursnell C., Patro R., Hibberd J.M., Kelly S. 2016. TransRate: reference-free

1491   quality assessment of de novo transcriptome assemblies. Genome Res. 26:1134–1144.

1492   Solís-Lemus C., Ané C. 2016a. Inferring Phylogenetic Networks with Maximum

1493   Pseudolikelihood under Incomplete Lineage Sorting. PLOS Genet. 12:e1005896.

1494   Soltis D.E., Kuzoff R.K. 1995. Discordance between nuclear and chloroplast phylogenies in the

1495   Heuchera group (Saxifragaceae). Evolution. 49:727–742.

1496   Song L., Florea L. 2015. Rcorrector: efficient and accurate error correction for Illumina RNA-

1497   seq reads. GigaScience. 4:48.

1498   Srivastava S.K. 1969. Assorted angiosperm pollen from the Edmonton Formation

1499   (Maestrichtian), Alberta, Canada. Can. J. Bot. 47:975–989.

1500   Stamatakis A. 2014. RAxML version 8 - a tool for phylogenetic analysis and post-analysis of

1501   large phylogenies. Bioinformatics. 30:1312–1313.

1502   Sugiura N. 1978. Further analysts of the data by akaike' s information criterion and the finite

1503   corrections. Commun. Stat. - Theory Methods. 7:13–26.

1504   Swofford D. 2002. PAUP*. Phylogenetic analysis using parsimony (*and other methods) version

1505   4. Sunderland MA Sinauer Assoc.

1506   Than C., Ruths D., Nakhleh L. 2008. PhyloNet: a software package for analyzing and

1507   reconstructing reticulate evolutionary relationships. BMC Bioinformatics. 9:322–16.

The Angiosperm Phylogeny Group, Chase M.W., Christenhusz M.J.M., Fay M.F., Byng J.W.,

Judd W.S., Soltis D.E., Mabberley D.J., Sennikov A.N., Soltis P.S., Stevens P.F. 2016.

An update of the Angiosperm Phylogeny Group classification for the orders and families

of flowering plants: APG IV. Bot. J. Linn. Soc. 181:1–20.

Varga T., Krizsán K., Földi C., Dima B., Sánchez-García M., Sánchez-Ramírez S., Szöllősi G.J.,

Szarkándi J.G., Papp V., Albert L., Andreopoulos W., Angelini C., Antonín V., Barry

K.W., Bougher N.L., Buchanan P., Buyck B., Bense V., Catcheside P., Chovatia M.,

Cooper J., Dämon W., Desjardin D., Finy P., Geml J., Haridas S., Hughes K., Justo A.,

Karasiński D., Kautmanova I., Kiss B., Kocsubé S., Kotiranta H., LaButti K.M., Lechner

B.E., Liimatainen K., Lipzen A., Lukács Z., Mihaltcheva S., Morgado L.N., Niskanen T.,

Noordeloos M.E., Ohm R.A., Ortiz-Santana B., Ovrebo C., Rácz N., Riley R., Savchenko

A., Shiryaev A., Soop K., Spirin V., Szebenyi C., Tomšovský M., Tulloss R.E., Uehling

J., Grigoriev I.V., Vágvölgyi C., Papp T., Martin F.M., Miettinen O., Hibbett D.S., Nagy

L.G. 2019. Megaphylogeny resolves global patterns of mushroom evolution. Nat. Ecol.

Evol. 3:668–678.

Vargas O.M., Ortiz E.M., Simpson B.B. 2017. Conflicting phylogenomic signals reveal a pattern

of reticulate evolution in a recent high-Andean diversification (Asteraceae: Astereae:

Diplostephium). New Phytol. 214:1736–1750.

Walker J.F., Walker-Hale N., Vargas O.M., Larson D.A., Stull G.W. 2019. Characterizing gene

tree conflict in plastome-inferred phylogenies. PeerJ. 7:e7747.

Walker J.F., Yang Y., Feng T., Timoneda A., Mikenas J., Hutchison V., Edwards C., Wang N.,

Ahluwalia S., Olivieri J., Walker-Hale N., Majure L.C., Puente R., Kadereit G.,

Lauterbach M., Eggli U., Flores-Olvera H., Ochoterena H., Brockington S.F., Moore

M.J., Smith S.A. 2018. From cacti to carnivores: Improved phylotranscriptomic sampling

and hierarchical homology inference provide further insight into the evolution of

Caryophyllales. Am. J. Bot. 105:446–462.

Wang Y., Tang H., DeBarry J.D., Tan X., Li J., Wang X., Lee T. -h., Jin H., Marler B., Guo H.,

Kissinger J.C., Paterson A.H. 2012. MCScanX: a toolkit for detection and evolutionary

analysis of gene synteny and collinearity. Nucleic Acids Res. 40:e49–e49.

Wen D., Nakhleh L. 2018. Coestimating Reticulate Phylogenies and Gene Trees from Multilocus

Sequence Data. Syst. Biol. 67:439–457.

Wen D., Yu Y., Hahn M.W., Nakhleh L. 2016a. Reticulate evolutionary history and extensive

introgression in mosquito species revealed by phylogenetic network analysis. Mol. Ecol.

25:2361–2372.

Wen D., Yu Y., Nakhleh L. 2016b. Bayesian Inference of Reticulate Phylogenies under the

Multispecies Network Coalescent. PLOS Genet. 12:e1006006.

Wickett N.J., Mirarab S., Nguyen N., Warnow T., Carpenter E., Matasci N., Ayyampalayam S.,

Barker M.S., Burleigh J.G., Gitzendanner M.A., Ruhfel B.R., Wafula E., Der J.P.,

Graham S.W., Mathews S., Melkonian M., Soltis D.E., Soltis P.S., Miles N.W., Rothfels

C.J., Pokorny L., Shaw A.J., DeGironimo L., Stevenson D.W., Surek B., Villarreal J.C.,

Roure B., Philippe H., dePamphilis C.W., Chen T., Deyholos M.K., Baucom R.S.,

Kutchan T.M., Augustin M.M., Wang J., Zhang Y., Tian Z., Yan Z., Wu X., Sun X.,

Wong G.K.-S., Leebens-Mack J. 2014. Phylotranscriptomic analysis of the origin and

early diversification of land plants. Proc. Natl. Acad. Sci. 111:E4859–E4868.

Widhelm T.J., Grewe F., Huang J.-P., Mercado-Díaz J.A., Goffinet B., Lücking R., Moncada B.,

Mason-Gamer R., Lumbsch H.T. 2019. Multiple historical processes obscure

phylogenetic relationships in a taxonomically difficult group (Lobariaceae, Ascomycota).

Sci. Rep. 9:8968.

Xu B., Yang Z. 2016. Challenges in Species Tree Estimation Under the Multispecies Coalescent

Model. Genetics. 204:1353–1368.

Xu C., Jiao C., Sun H., Cai X., Wang X., Ge C., Zheng Y., Liu W., Sun X., Xu Y., Deng J.,

Zhang Z., Huang S., Dai S., Mou B., Wang Q., Fei Z., Wang Q. 2017. Draft genome of

spinach and transcriptome diversity of 120 Spinacia accessions. Nat. Commun. 8:15275.

Yang Y., Moore M.J., Brockington S.F., Timoneda A., Feng T., Marx H.E., Walker J.F., Smith

S.A. 2017. An Efficient Field and Laboratory Workflow for Plant Phylotranscriptomic

Projects. Appl. Plant Sci. 5:1600128.

Yang Y., Smith S.A. 2013. Optimizing de novo assembly of short-read RNA-seq data for

phylogenomics. BMC Genomics. 14:328.

Yang Y., Smith S.A. 2014. Orthology Inference in Nonmodel Organisms Using Transcriptomes

and Low-Coverage Genomes: Improving Accuracy and Matrix Occupancy for

Phylogenomics. Mol. Biol. Evol. 31:3081–3092.

Yao G., Jin J.-J., Li H.-T., Yang J.-B., Mandala V.S., Croley M., Mostow R., Douglas N.A.,

Chase M.W., Christenhusz M.J.M., Soltis D.E., Soltis P.S., Smith S.A., Brockington S.F.,

Moore M.J., Yi T.-S., Li D.-Z. 2019. Plastid phylogenomic insights into the evolution of

Caryophyllales. Mol. Phylogenet. Evol. 134:74–86.

Yu Y., Degnan J.H., Nakhleh L. 2012. The Probability of a Gene Tree Topology within a

Phylogenetic Network with Applications to Hybridization Detection. PLoS Genet.

8:e1002660–10.

Yu Y., Dong J., Liu K.J., Nakhleh L. 2014. Maximum likelihood inference of reticulate

evolutionary histories. Proc. Natl. Acad. Sci. 111:16448–16453.

Yu Y., Nakhleh L. 2015. A maximum pseudo-likelihood approach for phylogenetic networks.

BMC Genomics. 16:S10.

Zhang C., Ogilvie H.A., Drummond A.J., Stadler T. 2018a. Bayesian Inference of Species

Networks from Multilocus Sequence Data. Mol. Biol. Evol. 35:504–517.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018b. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:523.

Zhao T., Schranz M.E. 2019. Network-based microsynteny analysis identifies major differences

and genomic outliers in mammalian and angiosperm genomes. Proc. Natl. Acad. Sci.

116:2165–2174.

Zheng Y., Janke A. 2018. Gene flow analysis method, the D-statistic, is robust in a wide

parameter space. BMC Bioinformatics. 19:10.

Zhu J., Liu X., Ogilvie H.A., Nakhleh L.K. 2019. A divide-and-conquer method for scalable

phylogenetic network inference from multilocus data. Bioinformatics. 35:i370–i378.

Zhu J., Wen D., Yu Y., Meudt H.M., Nakhleh L. 2018. Bayesian inference of phylogenetic

networks from bi-allelic genetic markers. PLOS Comput. Biol. 14:e1005932.

Zwick A., Regier J.C., Zwickl D.J. 2012. Resolving Discrepancy between Nucleotides and

Amino Acids in Deep-Level Arthropod Phylogenomics: Differentiating Serine Codons in

21-Amino-Acid Models. PLoS ONE. 7:e47450.