Title: Thetaburst TMS to the posterior superior temporal sulcus decreases resting-state fMRI connectivity across the face processing network

Short Title: TMS to the STS disrupts the face network

Authors: Daniel A Handwerker[1], Geena Ianni [2], Benjamin Gutierrez [1], Vinai Roopchansingh [3], Javier Gonzalez-Castillo [1], Gang Chen [4], Peter A Bandettini[1, 3], Leslie G Ungerleider [2], & David Pitcher [5]

1. Section on Functional Imaging Methods, Laboratory of Brain and Cognition, National Institute of Mental Health, Bethesda, MD, 20892, U.S.A.

2. Section on Neurocircuitry, Laboratory of Brain and Cognition, National Institute of Mental Health, Bethesda, MD, 20892, U.S.A.

3. Functional MRI Facility, National Institute of Mental Health, Bethesda, MD, 20892, U.S.A.

4. Scientific and Statistical Computing Core, National Institute of Mental Health, Bethesda, MD, 20892, U.S.A.

5. Department of Psychology, University of York, Heslington, York, YO10 5DD, U.K.

Corresponding authors: Daniel A Handwerker - Email: handwerkerd@mail.nih.gov

David Pitcher – Email: david.pitcher@york.ac.uk

Keywords: Resting-state fMRI, thetaburst TMS (TBS), face network, superior temporal sulcus (STS), multi-echo fMRI

Classifications: 30.010: Mapping of functional networks, 30.015: Resting state networks, 50.015: Network manipulation and control

Author Contributions: D.A.H, P.A.B, L.G.U & D.P designed research; D.A.H, G.I, B.G, V.R, J.G-C & D.P performed research; D.A.H, G.C. & D.P analyzed data; D.A.H, P.A.B, L.G.U & D.P wrote the paper

Technical Terms:

TBS vs TMS: Transcranial magnetic stimulation (TMS) is a method that induces current in neural tissue by using a rapidly changing magnetic field. The pattern of magnetic field changes can vary. Thetaburst TMS (TBS) is a type of TMS where the same stimulation pattern fluctuates at around a 5Hz cycle.

Multi-echo fMRI: During typical fMRI, protons are excited and there is a delay, the echo time, before data are collected. That delay is typically designed to result in a high contrast for blood oxygenation differences. In multi-echo fMRI, data are collected at several echo times each time protons are excited. This results in data that have different levels of contrast for blood oxygenation differences. This added information can be used to empirically decrease noise.

Face network: A group of brain regions that show significant activity changes in response to visual face stimuli. While these regions have been defined using univariate analyses with task-based fMRI, they often significantly correlate with each other at rest. In this manuscript, the following regions were a priori defined as part of the face network: posterior superior temporal sulcus (pSTS), amygdala, fusiform face area (FFA), and occipital face area (OFA).

Matrix based analysis (MBA): A recent approach that uses a Bayesian multilevel modeling framework to identify pairs of ROIs where a decrease in correlation magnitude was larger than expected along with a measure of statistical evidence. With this approach, correlations between all pairs of ROIs are assessed as part of a single model rather than many independent statistical tests.

1 **Abstract**
2
Humans process faces using a network of face-selective regions distributed across the brain.

Neuropsychological patient studies demonstrate that focal damage to nodes in this network

can impair face recognition, but such patients are rare. We approximated the effects of

damage to the face network in neurologically normal human participants using thetaburst

transcranial magnetic stimulation (TBS). Multi-echo functional magnetic resonance imaging

(fMRI) resting-state data were collected pre- and post-TBS delivery over the face-selective

right superior temporal sulcus (rpSTS), or a control site in the right motor cortex. Results

showed that TBS delivered over the rpSTS reduced resting-state connectivity across the

extended face-processing network. This connectivity reduction was observed not only

between the rpSTS and other face-selective areas, but also between non-stimulated face-

selective areas across the ventral, medial and lateral brain surfaces (e.g. between the right

amygdala and bilateral fusiform face areas and occipital face areas). TBS delivered over the

motor cortex did not produce significant changes in resting-state connectivity across the face-

processing network. These results demonstrate that, even without task-induced fMRI signal

changes, disrupting a single node in a brain network can decrease the functional connectivity

between nodes in that network that have not been directly stimulated.

19
Author Summary

Human behavior is dependent on brain networks that perform different cognitive functions.

We combined thetaburst transcranial magnetic stimulation (TBS) with resting-state fMRI to

study the face processing network. Disruption of the face-selective right posterior superior

temporal sulcus (rpSTS) reduced fMRI connectivity across the face network. This

impairment in connectivity was observed not only between the rpSTS and other face-

selective areas, but also between non-stimulated face-selective areas on the ventral and

medial brain surfaces (e.g. between the right amygdala and bilateral fusiform face areas and

occipital face areas). Thus, combined TBS/fMRI can be used to approximate and measure the

effects of focal brain damage on brain networks, and suggests such an approach may be

useful for mapping intrinsic network organization.

32
33   **<u>Introduction</u>**
34
The ubiquitous presence of faces in our environment makes them a salient stimulus for

studying the cognitive functions of the human brain. Functional magnetic resonance imaging

studies have identified regions across the brain that exhibit a stronger neural response to faces

than to objects (Gauthier et al., 2000; Kanwisher, McDermott, & Chun, 1997; McCarthy,

Puce, Gore, & Allison, 1997; Phillips et al., 1997; Puce, Allison, Bentin, Gore, & McCarthy,

1998). These face-selective regions are linked to form distributed nodes of a face-processing

network (Calder & Young, 2005; Haxby, Hoffman, & Gobbini, 2000). Neuropsychological

patients with damage to these face-selective areas exhibit face-selective recognition

impairments (Barton, 2008; Bouvier & Engel, 2006; Landis, Cummings, Christen, Bogen, &

Imhof, 1986; Rezlescu, Barton, Pitcher, & Duchaine, 2014; Rossion et al., 2003), providing

strong evidence that faces are processed in a specialized network. Despite the importance

they have afforded the study of brain function, such patients are rare. In addition, the

interpretation of the data they produce is limited by individual differences in pre-morbid

ability (Farah, 2004), and compensatory effects of plasticity that may have occurred after the

incident (Robertson & Murre, 1999).

In the present study we used thetaburst transcranial magnetic stimulation (TBS) in

combination with multi-echo fMRI acquisition (Posse et al., 1999). Resting-state fMRI was

used as a proxy for the impact of disruption on one node in the face network of

neurologically normal experimental participants. Over two sessions, these participants were

scanned using multi-echo fMRI pre- and post-TBS delivery over the face-selective right

superior temporal sulcus (rpSTS), the stimulation site of interest, or the hand area of the right

motor cortex, the control site. Multi-echo fMRI denoising is a relatively new method that

quantitatively identifies and removes non-blood oxygen-weighted noise from fMRI data

(Kundu et al., 2013). Prior to TBS, participants viewed 3-second videos of moving faces and

objects (Pitcher, Dilks, Saxe, Triantafyllou, & Kanwisher, 2011) during fMRI scanning to

functionally localize face-selective regions of interest (ROIs).

The face-selective areas in the rpSTS, the fusiform face area (FFA) (Kanwisher et al., 1997;

McCarthy et al., 1997), and the occipital face area (OFA) (Gauthier et al., 2000) comprise the

core nodes of the face processing network (Calder & Young, 2005; Haxby et al., 2000). In

addition to these core nodes, fMRI studies have also identified face-selective voxels in the

amygdala (Phillips et al., 1997). Face processing models propose these areas perform

different cognitive functions (e.g. recognizing identity, facial expression or eye gaze

direction), but share task-relevant information. The existence of functional connections

between face areas is supported by evidence showing that disruption to one of these areas can

impair a range of face recognition tasks. For example, TMS delivered over the rpSTS

disrupts the McGurk effect (Beauchamp, Nath, & Pasalar, 2010) as well as performance on

face-processing tasks involving eye-gaze (Pourtois et al., 2004), trustworthiness judgments

(Dzhelyova, Ellison, & Atkinson, 2011) and discriminating facial expressions (Pitcher, 2014;

Sliwinska & Pitcher, 2018). Neuropsychological patients with lesions to the rpSTS also show

face processing impairments. A patient with a lesion to the rpSTS and angular gyrus was

impaired on an unfamiliar face matching task (Sakurai, Hamada, Tsugawa, & Sugimoto,

2016), while another patient with a lesion to right superior temporal gyrus was impaired at

discriminating gaze detection (Akiyama et al., 2006).

Previous studies have shown TMS-induced face network activation changes during tasks

(Pitcher, Duchaine, & Walsh, 2014; Pitcher, Japee, Rauth, & Ungerleider, 2017). The nodes

of the face network also show functional integration during resting state fMRI (Li, Song, &

Liu, 2019; X. Wang et al., 2016; Zhang, Tian, Liu, Li, & Lee, 2009). In this study, we tested

whether the same face network areas defined in those studies show correlation decreases in

response to TMS to the rpSTS, even in the absence of a face stimulus. If so, then this would

build evidence that behaviorally defined face processing regions are part of a network whose

nodes regularly communicate and interact with each other without this communication being

induced by a specific set of stimuli. Results demonstrated that TBS delivered over the rpSTS

caused a reduction in correlations between the stimulated node and unstimulated nodes of the

face processing network.

92
93 **<u>Results</u>**
94
For each volunteer, face-selective regions of interest (ROIs) were localized using voxels with

larger responses to face videos than to object videos for the left posterior superior temporal

sulcus (lpSTS), bilateral fusiform face areas (FFA), bilateral occipital face areas (OFA) and

bilateral amygdala. The right posterior superior temporal sulcus (rpSTS) ROIs were face-

selective voxels in gray matter that were also within an 18mm diameter sphere centered on

the stimulation site. This added restriction means the voxels in the rpSTS ROI were likely to

have been directly affected by TBS stimulation. The stimulation site was defined using the

same functional localizer scan, but collected on a preceding session. The right hand motor

stimulation site was anatomically defined using an anatomical scan from a preceding session.

ROIs for the bilateral hand areas were manually drawn following the gray matter anatomy of

the hand knob in the precentral sulcus. The locations for the regions of interest are shown in

Figure S1.

For ROI-based analyses, mean time series of the ten-minute resting state fMRI data were

calculated using the voxels within each ROI. The correlations between these ROIs were

calculated for the resting runs pre- and post-TBS stimulation. The correlation coefficients

were Fisher Z transformed. Since the ROIs are *apriori* selected from a network where we

expect to see disruption (Pitcher et al., 2014; Pitcher et al., 2017) – with the hand motor

regions as a control, the statistical changes of interest will focus on the ROI analyses. A

matrix based analysis (MBA) through Bayesian multilevel modeling was used to identify

pairs of ROIs where a decrease in correlation magnitude was larger than expected along with

a measure of statistical evidence (Chen et al., 2019). The advantage of this approach is that,

instead of adopting a univariate GLM with the assumption that each ROI pair is an

independent entity that shares no commonality or similarity with its peers, the magnitude

estimates and uncertainties of all ROI pairs are assessed as part of a single integrative model.

Figure 1 shows the effects of stimulation on the correlations pre – post TBS for all pairs of

ROIs (bilateral STS, FFA, OFA, amygdala and hand motor cortices) using Bayesian

multilevel modeling. For rpSTS stimulation, multiple ROI pairs within the face processing

network showed decreased connectivity that would be likely to be greater than zero with a

posterior probability of at least 0.95 (Fig 1A). For pre – post TBS to the motor cortex (Fig

1B) and interaction effects, there was no strong evidence that any ROI pairs were likely to be

greater than zero. All posterior probabilities were less than 0.85 and all effect sizes were less

than 0.07 (Fig. S2).

Fisher Z transformed correlation magnitude changes from pre - post TMS to the (A) rpSTS and (B) right motor cortex. Magnitudes are the Matrix Based Analysis (MBA) model fits across the population. Changes that are likely to be greater than 0 with a posterior probability of at least 95% are full squares and include an *. The squares' edge lengths decrease linearly to their minimum size for a 90% likelihood or less. The green line marks the ROI pairs that are within the pre-defined face-selective network. Fig. S2 contains an unthresholded version of this figure and the interaction effect.

129
To help visually compare the effect sizes for rpSTS vs motor stimulation, Figure 2 shows

boxplots for the correlations between the rpSTS and the other ROIs for the runs pre and post

TBS. The colored section of the pie charts in this figure show the number of volunteers that

had a pre – post TBS correlation decrease. These pie charts show that 11-13 of the 16

volunteers showed a correlation decrease post rpSTS stimulation. The supplementary

material includes boxplots for correlation magnitudes of all ROI pairs (Figs S3 and S4).

Whether the correlation is between a pair of ROIs that includes the directly stimulated rpSTS

or another pair of face network ROIs, the correlations to other ROIs in the face network

decreased across the majority of volunteers when the rpSTS was stimulated.

Correlations between the rpSTS ROI and ROIs in a pre-defined face network as well as the bilateral primary motor hand regions. These data correspond to the first row of the matrix in figure 1. The full matrix of bar plots in in Fig. S2. Magnitudes are the MBA model fits across the population for each condition. Boxplots show 25-75% of the distribution. The white line is the median. Whiskers are the maximum and minimum values excluding outliers. MBA used to calculate posterior probabilities that a difference is greater than 0. The correlations between the rpSTS and other face regions consistently decrease post rpSTS stimulation, but not post rMotor stimulation.

There is a possibility that TBS to the rpSTS would cause correlation decreases between the

rpSTS and the entire brain. To test for this possibility, all subjects' data were aligned to each

other and a group map was calculated with an ANOVA using multivariate modeling (Figure

3). While there are decreases in activity that do not all cross meaningful statistical thresholds

both within and outside the face network, the correlation decreases post rpSTS stimulation

are clearly not a global effect.

Figure 3

Whole brain correlation changes (Fisher Z transformed) pre - post TBS stimulation. Voxel coloring shows the magnitude of the correlation difference. Voxels are opaque for p>=0.01 to transparent for p=0.The green outlines are voxels in the ROIs of the pre-defined face network in at least 3 volunteers. For TMS to the rpSTS, many of the larger signal drops are within the face network ROIs. TMS to right motor shows decreases across more of the brain that isn't localized to the face network.

146
147
148   **<u>Discussion</u>**
149
Models of the human face processing network propose that face-selective areas perform

different functional roles but share task-relevant information when processing faces (Calder

& Young, 2005; Haxby et al., 2000). In the current study, we approximated the impact of

disruption to the face network by delivering TBS over the face-selective area in the right

posterior superior temporal sulcus (rpSTS). The impact of the TBS stimulation was measured

with resting-state fMRI, with the goal of examining whether the relationship between face

network regions is altered even in the absence of a face-selective cognitive task. Our results

demonstrated that disruption of the rpSTS reduced resting-state connectivity, measured by

correlated activity, across the nodes of the face processing network. Figure 1 shows a

reduction in connectivity between the rpSTS and bilateral FFA, OFA, and amygdala as well

as between 14 of 21 pairs of ROIs in the face network that were not directly stimulated. This

result demonstrates that disrupting a single face-selective area causes widespread

connectivity decreases across the extended face processing network.

In addition, functional connectivity between different nodes of the face network has been

shown to correlate with behavioural measures of face recognition. Zhu and colleagues

demonstrated that connectivity between the OFA and FFA correlated with performance on

face, but not object, matching discrimination tasks (Zhu, Zhang, Luo, Dilks, & Liu, 2011).

O'Neill and colleagues demonstrated that connectivity between the FFA and perirhinal cortex

correlated with an upright, but not inverted, face matching task (O'Neil, Hutchison, McLean,

& Kohler, 2014). Most recently, Ramot and colleagues demonstrated that face recognition

memory was correlated with connectivity between the face network and regions of the medial

temporal lobe (including the hippocampus) (Ramot, Walsh, & Martin, 2019). Taken together,

these findings suggest that synchronized spontaneous neural activity between face regions

might be responsible for the different behavioural aspects of face recognition. The current

study builds upon and extends these prior findings to show that transient disruption of a

single node in the face network can disrupt functional connectivity across the wider network.

Figures 2, S3, and S4 visualize the effect of TBS across subjects to show that correlations

between face network regions have correlation decreases for the majority of volunteers post

rpSTS stimulation. This widespread pattern of distributed correlation decreases suggests that

the nodes of the face network are tightly interconnected, and that distruption of one node can

affect other nodes in the network. Importantly, TBS delivered over the right motor cortex

(which acted as a control site) did not produce above chance reductions in functional

connectivity across the face network. The observed network-wide drop in resting-state

connectivity is consistent with models proposing that the rpSTS is a core component of the

face-processing network (Calder & Young, 2005; Haxby et al., 2000). While we do show an

effect of rpSTS stimulation and not an effect for motor stimulation, we do not observe a

significant effect of motor cortex stimulation between motor ROIs, nor an interaction effect.

Our results do not completely match our predictions, possibly due to the combined effect of

two inherently noisy techniques, namely, resting-state fMRI and TBS. Resting-state fMRI is

known to be more strongly affected than task fMRI by many noise sources (e.g. inter-subject

cognitive variation, physiological noise, head motion, scanner instabilities). Also, there is still

not a full understanding of all factors that modulate TBS effect size including how much

spatial precision matters, whether different brain regions are more sensitive to TBS than

others, and the causes of inter-subject variability in TBS sensitivity. Specifically in this study,

we note our post-stimulation observation of spatial variation between the hand motor ROI

and the hand motor stimulation site (Fig S1). This variation in stimulation site may have led

to an observed increase in variance of correlation changes post motor stimulation vs rpSTS

stimulation (Fig S3 and S4). That added variance may have reduced the posterior

probabilities involving motor stimulation data. Although we did our best to account for

known sources of noise, residual artifacts can be expected to exist, and those may have a

multiplicative effect when combining the techniques, further hindering our statistical power.

Additionally, figure 3 shows large, but not statistically significant, correlation decreases in

response to TBS to the rpSTS. These occur both within and outside the a priori defined face

network regions. While we will not attempt to interpret these below-threshold correlation

decreases, it is important to note that rpSTS function is not limited to face processing.

Additionally, the stimulation of an approximately $1cm^3$ volume of cortex may affect brain

regions outside of a face network.

In this study, we could not directly test whether TBS to the rpSTS caused alterations in face

processing because we cannot both collect resting state fMRI and also present face-selective

stimuli during the post TBS time window when the TBS effect is observable. However, in

previous studies using face-selective stimulus evoked BOLD mmeasures and the same TBS

stimulation protocol as used here, changes were reported in responses to face stimuli (Pitcher

et al., 2014; Pitcher et al., 2017). The reduction in network connectivity we observed between

non-stimulated face-selective regions on the medial and ventral brain surfaces extends the

findings of this prior work of Pitcher et al. Our previous combined TBS/fMRI studies

examined the impact of disruption in face-selective regions while participants viewed face

videos. These studies showed that TBS delivered over the rpSTS reduced the neural response

to faces in the rpSTS itself, as well as in other face-selective areas including the right

fusiform face area (rFFA) (Pitcher, 2014) and face-selective voxels in the right amygdala

(Pitcher et al., 2017). However, while these studies reported a reduction in the neural

response to faces in remote face-selective areas, neither study was capable of detecting

changes in functional connectivity between face-selective areas. While prior studies

demonstrated functional connectivity between the FFA and the amygdala (Fairhall & Ishai,

2007; Herrington, Taylor, Grupe, Curby, & Schultz, 2011; Vuilleumier, Armony, Driver, &

Dolan, 2001, 2003), our study shows that correlations between these areas are reduced post-

TBS delivery over a remote node in the network, in this case the rpSTS.

Non-human primate neuroanatomical studies in macaques report a cortical pathway

projecting from the STS to dorsal sub-regions of the amygdala (Aggleton, Burton, &

Passingham, 1980; Stefanacci & Amaral, 2000, 2002). In addition, neuroimaging and lesion

studies of humans and macaques demonstrate that the amygdala is engaged in facial

expression recognition (Adolphs, Tranel, Damasio, & Damasio, 1994; Adolphs et al., 1999;

Calder et al., 1996; Hadj-Bouziane et al., 2012; Hoffman, Gothard, Schmid, & Logothetis,

2007; Morris et al., 1996). We previously proposed that humans have a cortical pathway

projecting along the STS into the amygdala that processes changeable facial aspects (e.g

expression), which was supported using combined TBS/fMRI during face, body, and object

viewing (Pitcher et al., 2017). By replicating this effect using functional connectivity and

without a viewing task, we provide further support for a pathway between these regions.

Neuropsychological patients exhibiting face-selective recognition impairments were essential

to the development of cognitive and brain models of face processing (Bruce & Young, 1986;

Haxby et al., 2000). Studies of patients with right lateral lesions (in the area of the pSTS)

showing impairments with eye gaze direction detection (Akiyama et al., 2006; Campbell,

Heywood, Cowey, Regard, & Landis, 1990) and unfamiliar face identity matching (Sakurai

et al., 2016) have been reported. However, prosopagnosic patients with lateral lesions are less

common than those with ventral lesions. Our study further demonstrates that transiently

disrupting the brains of neurologically normal experimental participants with TBS offers an

alternative and safe proxy for modeling the effects of focal cortical disruption on cognitive

networks. TBS can be combined with functional magnetic resonance imaging (fMRI) to

study the networks that process scenes (Mullin & Steeves, 2013), faces (Rafique, Solomon-

Harris, & Steeves, 2015; Solomon-Harris, Rafique, & Steeves, 2016), decision making

(Rahnev, Nee, Riddle, Larson, & D'Esposito, 2016) and memory (J. X. Wang et al., 2014).

Crucially, while we do see some decreases between a few face network ROIs and the motor

ROIs (Fig 1), we examined the voxel-wise correlations across the brain to show that the

largest distruptions of TBS to the rpSTS were in the face selective network. Still, the

decreases in some connections to motor cortex ROIs hint at how altering a single node can

potentially distrupt multiple distinct networks. Each brain region does not contribute to only

one self-contained network of brain regions.

Neurologists have studied the impact of focal brain lesions in patients for over two hundred

years. The study of these patients has been highly influential and has produced insights into

human cognition. However, patients with focal lesions in brain regions of interest are

extremely rare. In addition, the interpretation of the data they produce is tempered by factors

like individual differences in pre-morbid ability (Robertson & Murre, 1999) and the unknown

effects of neural plasticity that may have occurred after their incident (Farah, 2004). TBS

combined with resting state fMRI has been used to study the default mode network

(Abellaneda-Perez et al., 2019; Shang et al., 2019), attention (Anderkova et al., 2018),

visualspatial neglect (Fu et al., 2017), cerebellar connectivity (Rastogi et al., 2017), mental

illness, (Baeken, Duprat, Wu, De Raedt, & van Heeringen, 2017) and memory in humans

(Mancini et al., 2017). These studies show that measuring transient TBS induced disruption

with resting state fMRI offers a systematic experimental methodology for extending more

than two hundred years of neuropsychological research.

In conclusion, models propose that face-selective brain areas can be linked together to form

the distributed components of a face processing network (Calder & Young, 2005; Haxby et

al., 2000). The current findings support these models by showing that transient disruption to

one of these components, the rpSTS, results in widespread correlation decreases across the

distributed nodes of this network that persists for ten minutes.

<u>Materials and Methods</u>

Participants

17 right-handed participants (11 female, age mean±stdev=27±5), with no major medical

illness, no neurological or psychiatric illness, no serious head injury, no learning disability,

no history of drug or alcohol abuse in the past 3 months, no prescription drugs or

supplements affecting brain function, no serious vision or hearing problems and with normal,

or corrected-to-normal, vision gave informed consent as directed by the National Institutes of

Health (NIH) Institutional Review Board (ClinicalTrials.gov identifier: NCT01617408). One

participant was excluded due to high motion (207 of 1200 volumes in the 4 resting runs

censored due to motion or outliers vs 0-40 for all other volunteers).

Experimental Design

<u>Procedure</u>

Participants completed three separate fMRI sessions, each performed on a different day. The

first session was an fMRI experiment designed to individually localize the TBS stimulation

sites in each participant. Participants viewed face and object videos and a high resolution

structural scan was also taken. The data collected in this initial session were used for TBS

target site identification only. During the two subsequent fMRI sessions, participants were

scanned pre and post receiving TBS stimulation of either the right posterior superior temporal

sulcus (rpSTS) or the right primary motor cortex hand knob (rMotor). Stimulation site order

was balanced across participants. The two TBS sessions were 7 – 182 days apart

(median=23). All the data presented were collected during these two TBS sessions.

<u>Stimulation Site Localizer Session and Selection</u>

Stimulation sites were localized using individual structural and functional images collected

during an fMRI localizer task that each participant completed prior to the combined

TBS/fMRI sessions. For the independent localizer runs used to identify face-selective regions

of interest (ROIs), participants viewed 3-second video clips of faces, objects, and scrambled

objects. There were sixty movie clips for each category in which distinct exemplars appeared

multiple times. Movies of faces were filmed on a black background, and framed close-up to

reveal only the faces of 7 children as they danced or played with toys or adults (who were out

of frame). Fifteen different moving objects were selected that minimized any suggestion of

animacy of the object itself or of a hidden actor pushing the object (these included mobiles,

windup toys, toy planes and tractors, balls rolling down sloped inclines). Within each block,

stimuli were randomly selected from within the entire set for that stimulus category. This

meant that the same actor or object could appear within the same block. These stimuli were

used in a previous fMRI study of face perception (Pitcher et al., 2011) and previous

combined TBS/fMRI studies (Pitcher et al., 2014; Pitcher et al., 2017).

The rpSTS was identified using a contrast of faces greater than objects. The rpSTS

stimulation target was the peak voxel in the significant cluster in the rpSTS. A T1 weighted

anatomical scan was also collected during this session and the right motor cortex was the

most superior location on the motor cortex hand knob. A posthoc re-examination identified

two vounteers where the motor stimulation zone may not have directly stimulated the hand

motor region (Fig S1). This may have contributed to a greater variance of responses to hand

motor stimulation across the population and thus the lower posterior probabilities for

correlation decreases post motor stimulation and interaction effects between the stimulation

sites.

<u>Combined fMRI/TBS sessions</u>

Data acquisition: Participants were scanned using a GE 3-Tesla MR 750 scanner at the

National Institutes of Health (NIH). fMRI images were acquired using a 32-channel head coil

(36 slices, $3 \times 3 \times 3$ mm, FOV=21.6 cm, grid = 72x72, flip angle = 77°, ASSET=3, TR = 2 s,

TEs = 14.8, 27.1, & 39.5 ms). In addition, high-resolution MPRAGE anatomical scans (T1-

weighted, 1 x 1 x 1 mm resolution) were acquired to anatomically localize functional

activations and register functional data between sessions.

335   Before each MRI session, the stimulation site (either rpSTS or rMotor) was

located using the Brainsight TMS-MRI coregistration system (Rogue Research), the

information was extracted from the subject-specific localizer scans (see the previous

"Stimulation Site Localizer Session and Selection" section) and the proper coil locations

were marked on each participant's scalp using a marker pen. During the MRI session, two

10-minute resting-state scans (310 volumes) were acquired followed by functional localizer

scans similar to those used for the stimulation site localizer session and then an MPRAGE

anatomical scan. The volunteers were then removed from the scanner.

343   A Magstim Super Rapid Stimulator (Magstim; Whitland, UK) was used to deliver the

TBS via a figure-eight coil with a wing diameter of 70 mm. TBS was delivered at an intensity

of 80% of active motor threshold or 30% of machine output (whichever was higher) over

each participant's functionally localized rpSTS or the right motor hand knob. We used a

continuous TBS paradigm (Huang, Edwards, Rounis, Bhatia, & Rothwell, 2005) of 3 pulses

at 50 Hz repeated at 200-ms intervals for a 60-second uninterrupted train of 900 pulses. This

is the same protocol that was used in our previous combined TBS/fMRI studies of the rpSTS

(Pitcher et al., 2014; Pitcher et al., 2017). We matched the protocol to aid with comparisons

across studies. The Stimulator coil handle was held pointing upwards and parallel to the

midline.

As soon as stimulation ended, volunteers were returned to the MRI scanner where the

following data were acquired: a brief anatomical reference scan to prescribe a slice placement

that was visually similar to the pre-TBS scan, an ASSET calibration scan, and two 10-minute

resting-state scans. The resting scans began 2.5-5 minutes after the last TBS pulse was

delivered depending on how quickly the participant was able to return to the scanner. After

the resting scan, another MPRAGE anatomical scan was acquired. Three of the 34 MRI

sessions did not include an anatomical scan during the post-TBS part of the session due to

time constraints.

The second 10-minute resting scans both pre and post TBS had more head motion in

many subjects. For 14 of the 64 pre or post TBS run pairs, there was at least 5 more motion

censored volumes in the second run, versus 1 for first run – second run. Similarly the median

delta motion (frame-to-frame difference) was larger for 49 of the second runs vs first runs.

The average across runs of median delta motion difference between run pairs was 0.01 larger

in the second runs. The average median delta motion was 0.05 so this was a 19% increase in

median motion. There was only a 7% difference in median delta motion for the first and third

runs, which were used in these analyses. While these are subtle differences, the systematic

nature of this difference added a confound that made these data difficult to interpret, so the

second runs were excluded from all analyses.

Data processing

Data were processed using AFNI (Cox, 1996). For each fMRI run: The first four pre-steady

state volumes were removed; Data were despiked to remove non-biological large signal

fluctuations and slice time corrected. Spatial alignment transforms for motion correction

within runs and registration across runs were calculated using the middle echo (TE=27.1ms)

time series. Non-uniformity of image intensity for the first volume of this time series was

corrected using the bias correction code in SPM8. This reference fMRI volume was aligned

to the MPRAGE anatomical from the same pre or post TBS session, if available. Otherwise,

it was aligned to the anatomical scan from the same day. The anatomical scans were aligned

to each other and averaged to make a single, higher quality anatomical volume per volunteer.

Registration quality for each pair of anatomical scans and for each fMRI to anatomical

alignment were visually inspected for accuracy. The motion correction and alignment

transforms were combined into a single affine transform matrix and applied to fMRI time

series for all echoes so that fMRI data from each subject was in a single subject-specific

space across sessions.

The multi-echo data were then denoised using the MEICA algorithm. MEICA uses the

expected properties of signal changes across multiple echoes times to identify and remove

signal fluctuations that are unlikely to reflect the blood oxygenation differences that are

central to fMRI. By doing this, MEICA has been shown to increase the signal-to-noise ratio

and remove data artifacts (Kundu et al., 2013; Kundu, Inati, Evans, Luh, & Bandettini, 2012).

The version of the denoising code that we used is available at:

https://github.com/handwerkerd/MEICA_FaceNetworkTMS. We will refer to the output of

the full MEICA algorithm as "denoised" data. The denoised data will be used for all analyses

in this manuscript.

Univariate statistical maps and correlations were calculated within subjects. For functional

localizer runs, time series were intensity normalized by dividing by the mean value in each

run. For resting-state runs, volumes when the first derivative Euclidian norm of the 6 motion

parameters was greater than 0.2 were censored. Volumes were also censored if more than

10% of the voxels in a volume contained large signal fluctuations that were considered

outliers. Subject had 0-40 censored volumes, across all 4 pre and post TBS resting runs (1200

volumes total). Censored volumes per 300 volume run were: mean=2, median=0,

maximum=22. The motion parameters and their first derivatives were regressed from the time

series. In the same step, signals from a Cerebral Spinal Fluid (CSF) ROI and a white matter

ROI that excluded voxels within 8 voxels of either stimulation site were averaged and

regressed from the data. The resting-state data were bandpass filtered (0.01-0.1Hz) with

censoring and the data were spatially smoothed with a 5-mm full-width-half-maximum

Gaussian kernel.

<u>ROI specifications and cross-subject alignment</u>

The statistical tests visualized in Figures 1 and 2 used regions of interest that were defined for

each volunteer. The average of each volunteer's anatomical scans, as described in the

previous section, were processed in Freesurfer (Fischl et al., 2004) to estimate volunteer-

specific anatomical boundaries for the fusiform gyri, lateral occipital gyri, and amygdala.

Any fMRI voxel that included at least 50% the Freesurfer estimated region was included in

the region. These anatomical regions were intersected with faces>objects functional

activation maps from the localizer scans collected during the TBS sessions (false-discovery-

rate threshold q<0.05). To make sure ROIs for each brain region were identified in each

volunteer, if 10% of the voxels in the anatomical ROI, or 10 voxels for anatomical ROIs with

less than 100 voxels did not cross this threshold, then the threshold was increased until this

threshold of faces>objects voxels was crossed. In 5 subjects, this liberal threshold was not

crossed in the amygdala ROIs, which has a mean of 51 voxels in the anatomical ROIs. In 4 of

these subjects, the same volunteers participated in a separate study that included the identical

functional localizer task with more significant faces>objects voxels in the regions of interest

(Pitcher et al., 2017). Those localizer scans were used to define the functional ROIs in those 4

volunteers. The voxels in the fusiform gyrus with the face>objects contrast were designated

the Fusiform Face Area (FFA) ROI and the voxels in the lateral occipital gyrus were

designated the Occipital Face Area (OFA) ROI. All volunteers had functionally localized

rpSTS and rFFA ROIs with q<0.05. The threshold was q>0.05 for 1 volunteer in lFFA and

rOFA, 2 volunteers in lpSTS, 3 volunteers in lOFA, 4 volunteers in rAmyg, and 8 volunteers

in lAmyg. The left and right hand knob motor ROIs were hand drawn on each volunteer's

structural scan by a single person and checked for accuracy by a second person.

For whole-brain group maps, the within-subject aligned anatomical volumes from

each subject were registered to a standard MNI space and then nonlinearly, iteratively warped

to each other to make a within-study anatomical template (Figures 3 and S1).

<u>Acknowledgements</u>

The research reported here was supported by the Intramural Research Program of the NIMH

(ZIAMH002783 and ZIAMH002918) and by BBSRC grant BB/P006981/1 awarded to D.P.

Clinical Protocol #: NCT01617408. This work utilized the computational resources of the

NIH HPC Biowulf cluster (http://hpc.nih.gov). We thank Nancy Kanwisher for providing

face localizer stimuli. We thank Shruti Japee for reviewing functional localizer results and for

identifying functional localizer scans from these volunteers that were collected as part of

other data sets.

<u>References</u>

Abellaneda-Perez, K., Vaque-Alcazar, L., Vidal-Pineiro, D., Jannati, A., Solana, E., Bargallo,

N., . . . Bartres-Faz, D. (2019). Age-related differences in default-mode network

connectivity in response to intermittent theta-burst stimulation and its relationships

with maintained cognition and brain integrity in healthy aging. *NeuroImage, 188*,

794-806. doi:10.1016/j.neuroimage.2018.11.036

Adolphs, R., Tranel, D., Damasio, H., & Damasio, A. (1994). Impaired recognition of

emotion in facial expressions following bilateral damage to the human amygdala.

Nature, 372(6507), 669-672. doi:10.1038/372669a0

Adolphs, R., Tranel, D., Hamann, S., Young, A. W., Calder, A. J., Phelps, E. A., . . .

Damasio, A. R. (1999). Recognition of facial emotion in nine individuals with

bilateral amygdala damage. *Neuropsychologia, 37*(10), 1111-1117.

doi:10.1016/s0028-3932(99)00039-1

Aggleton, J. P., Burton, M. J., & Passingham, R. E. (1980). Cortical and subcortical afferents

to the amygdala of the rhesus monkey (Macaca mulatta). *Brain Res, 190*(2), 347-368.

doi:10.1016/0006-8993(80)90279-6

Akiyama, T., Kato, M., Muramatsu, T., Saito, F., Umeda, S., & Kashima, H. (2006). Gaze

but not arrows: a dissociative impairment after right superior temporal gyrus damage.

Neuropsychologia, 44(10), 1804-1810. doi:10.1016/j.neuropsychologia.2006.03.007

Anderkova, L., Pizem, D., Klobusiakova, P., Gajdos, M., Koritakova, E., & Rektorova, I.

(2018). Theta Burst Stimulation Enhances Connectivity of the Dorsal Attention

Network in Young Healthy Subjects: An Exploratory Study. *Neural Plast, 2018*,

3106918. doi:10.1155/2018/3106918

Baeken, C., Duprat, R., Wu, G. R., De Raedt, R., & van Heeringen, K. (2017). Subgenual

Anterior Cingulate-Medial Orbitofrontal Functional Connectivity in Medication-

Resistant Major Depression: A Neurobiological Marker for Accelerated Intermittent

Theta Burst Stimulation Treatment? *Biol Psychiatry Cogn Neurosci Neuroimaging,*

2(7), 556-565. doi:10.1016/j.bpsc.2017.01.001

Barton, J. J. (2008). Prosopagnosia associated with a left occipitotemporal lesion.

Neuropsychologia, 46(8), 2214-2224. doi:10.1016/j.neuropsychologia.2008.02.014

Beauchamp, M. S., Nath, A. R., & Pasalar, S. (2010). fMRI-Guided transcranial magnetic

stimulation reveals that the superior temporal sulcus is a cortical locus of the McGurk

effect. *J Neurosci, 30*(7), 2414-2417. doi:10.1523/JNEUROSCI.4865-09.2010

Bouvier, S. E., & Engel, S. A. (2006). Behavioral deficits and cortical damage loci in cerebral

achromatopsia. *Cereb Cortex, 16*(2), 183-191. doi:10.1093/cercor/bhi096

Bruce, V., & Young, A. (1986). Understanding face recognition. *British Journal of*

Psychology, 77(3), 305-327. doi:10.1111/j.2044-8295.1986.tb02199.x

Calder, A. J., & Young, A. W. (2005). Understanding the recognition of facial identity and

facial expression. *Nat Rev Neurosci, 6*(8), 641-651. doi:10.1038/nrn1724

Calder, A. J., Young, A. W., Rowland, D., Perrett, D. I., Hodges, J. R., & Etcoff, N. L.

(1996). Facial emotion recognition after bilateral amygdala damage: Differentially

severe impairment of fear. *Cognitive Neuropsychology, 13*(5), 699-745.

doi:10.1080/026432996381890

Campbell, R., Heywood, C. A., Cowey, A., Regard, M., & Landis, T. (1990). Sensitivity to

496   eye gaze in prosopagnosic patients and monkeys with superior temporal sulcus

497   ablation. *Neuropsychologia, 28*(11), 1123-1142. doi:10.1016/0028-3932(90)90050-x

498   Chen, G., Burkner, P. C., Taylor, P. A., Li, Z., Yin, L., Glen, D. R., . . . Pessoa, L. (2019). An

499   integrative Bayesian approach to matrix-based analysis in neuroimaging. *Hum Brain*

500   *Mapp, 40*(14), 4072-4090. doi:10.1002/hbm.24686

501   Cox, R. W. (1996). AFNI: software for analysis and visualization of functional magnetic

502   resonance neuroimages. *Comput Biomed Res, 29*(3), 162-173.

503   doi:10.1006/cbmr.1996.0014

504   Dzhelyova, M. P., Ellison, A., & Atkinson, A. P. (2011). Event-related repetitive TMS

505   reveals distinct, critical roles for right OFA and bilateral posterior STS in judging the

506   sex and trustworthiness of faces. *J Cogn Neurosci, 23*(10), 2782-2796.

507   doi:10.1162/jocn.2011.21604

508   Fairhall, S. L., & Ishai, A. (2007). Effective connectivity within the distributed cortical

509   network for face perception. *Cereb Cortex, 17*(10), 2400-2406.

510   doi:10.1093/cercor/bhl148

511   Farah, M. J. (2004). *Visual agnosia* (2nd ed.). Cambridge, Mass.: MIT Press.

512   Fischl, B., van der Kouwe, A., Destrieux, C., Halgren, E., Segonne, F., Salat, D. H., . . . Dale,

513   A. M. (2004). Automatically parcellating the human cerebral cortex. *Cereb Cortex,*

514   *14*(1), 11-22. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/14654453

515   Fu, W., Cao, L., Zhang, Y., Huo, S., Du, J., Zhu, L., & Song, W. (2017). Continuous theta-

516   burst stimulation may improve visuospatial neglect via modulating the attention

network: a randomized controlled study. *Top Stroke Rehabil, 24*(4), 236-241.

doi:10.1080/10749357.2016.1253139

Gauthier, I., Tarr, M. J., Moylan, J., Skudlarski, P., Gore, J. C., & Anderson, A. W. (2000).

The fusiform "face area" is part of a network that processes faces at the individual

level. *J Cogn Neurosci, 12*(3), 495-504. doi:10.1162/089892900562165

Hadj-Bouziane, F., Liu, N., Bell, A. H., Gothard, K. M., Luh, W. M., Tootell, R. B., . . .

Ungerleider, L. G. (2012). Amygdala lesions disrupt modulation of functional MRI

activity evoked by facial expression in the monkey inferior temporal cortex. *Proc Natl*

Acad Sci U S A, 109(52), E3640-3648. doi:10.1073/pnas.1218406109

Haxby, J. V., Hoffman, E. A., & Gobbini, M. I. (2000). The distributed human neural system

for face perception. *Trends Cogn Sci, 4*(6), 223-233. doi:10.1016/S1364-

6613(00)01482-0

Herrington, J. D., Taylor, J. M., Grupe, D. W., Curby, K. M., & Schultz, R. T. (2011).

Bidirectional communication between amygdala and fusiform gyrus during facial

recognition. *NeuroImage, 56*(4), 2348-2355. doi:10.1016/j.neuroimage.2011.03.072

Hoffman, K. L., Gothard, K. M., Schmid, M. C., & Logothetis, N. K. (2007). Facial-

expression and gaze-selective responses in the monkey amygdala. *Curr Biol, 17*(9),

766-772. doi:10.1016/j.cub.2007.03.040

Huang, Y. Z., Edwards, M. J., Rounis, E., Bhatia, K. P., & Rothwell, J. C. (2005). Theta burst

stimulation of the human motor cortex. *Neuron, 45*(2), 201-206.

doi:10.1016/j.neuron.2004.12.033

Kanwisher, N., McDermott, J., & Chun, M. M. (1997). The fusiform face area: a module in

539   human extrastriate cortex specialized for face perception. *J Neurosci, 17*(11), 4302-

540   4311. doi:10.1523/JNEUROSCI.17-11-04302.1997

541   Kundu, P., Brenowitz, N. D., Voon, V., Worbe, Y., Vertes, P. E., Inati, S. J., . . . Bullmore, E.

542   T. (2013). Integrated strategy for improving functional connectivity mapping using

543   multiecho fMRI. *Proc Natl Acad Sci U S A, 110*(40), 16187-16192.

544   doi:10.1073/pnas.1301725110

545   Kundu, P., Inati, S. J., Evans, J. W., Luh, W. M., & Bandettini, P. A. (2012). Differentiating

546   BOLD and non-BOLD signals in fMRI time series using multi-echo EPI.

547   *NeuroImage, 60*(3), 1759-1770. doi:10.1016/j.neuroimage.2011.12.028

548   Landis, T., Cummings, J. L., Christen, L., Bogen, J. E., & Imhof, H. G. (1986). Are unilateral

549   right posterior cerebral lesions sufficient to cause prosopagnosia? Clinical and

550   radiological findings in six additional patients. *Cortex, 22*(2), 243-252.

551   doi:10.1016/s0010-9452(86)80048-x

552   Li, J., Song, Y., & Liu, J. (2019). Functional connectivity pattern in the core face network

553   reflects different mechanisms of holistic face processing measured by the whole-part

554   effect and composite-face effect. *Neuroscience, 408*, 248-258.

555   doi:10.1016/j.neuroscience.2019.04.017

556   Mancini, M., Mastropasqua, C., Bonni, S., Ponzo, V., Cercignani, M., Conforto, S., . . .

557   Bozzali, M. (2017). Theta Burst Stimulation of the Precuneus Modulates Resting

558   State Connectivity in the Left Temporal Pole. *Brain Topogr, 30*(3), 312-319.

559   doi:10.1007/s10548-017-0559-x

McCarthy, G., Puce, A., Gore, J. C., & Allison, T. (1997). Face-specific processing in the

human fusiform gyrus. *J Cogn Neurosci, 9*(5), 605-610.

doi:10.1162/jocn.1997.9.5.605

Morris, J. S., Frith, C. D., Perrett, D. I., Rowland, D., Young, A. W., Calder, A. J., & Dolan,

R. J. (1996). A differential neural response in the human amygdala to fearful and

happy facial expressions. *Nature, 383*(6603), 812-815. doi:10.1038/383812a0

Mullin, C. R., & Steeves, J. K. (2013). Consecutive TMS-fMRI reveals an inverse

relationship in BOLD signal between object and scene processing. *J Neurosci, 33*(49),

19243-19249. doi:10.1523/JNEUROSCI.2537-13.2013

O'Neil, E. B., Hutchison, R. M., McLean, D. A., & Kohler, S. (2014). Resting-state fMRI

reveals functional connectivity between face-selective perirhinal cortex and the

fusiform face area related to face inversion. *NeuroImage, 92*, 349-355.

doi:10.1016/j.neuroimage.2014.02.005

Phillips, M. L., Young, A. W., Senior, C., Brammer, M., Andrew, C., Calder, A. J., . . .

David, A. S. (1997). A specific neural substrate for perceiving facial expressions of

disgust. *Nature, 389*(6650), 495-498. doi:10.1038/39051

Pitcher, D. (2014). Facial expression recognition takes longer in the posterior superior

temporal sulcus than in the occipital face area. *J Neurosci, 34*(27), 9173-9177.

doi:10.1523/JNEUROSCI.5038-13.2014

Pitcher, D., Dilks, D. D., Saxe, R. R., Triantafyllou, C., & Kanwisher, N. (2011). Differential

selectivity for dynamic versus static information in face-selective cortical regions.

NeuroImage, 56(4), 2356-2363. doi:10.1016/j.neuroimage.2011.03.067

Pitcher, D., Duchaine, B., & Walsh, V. (2014). Combined TMS and FMRI reveal dissociable

cortical pathways for dynamic and static face perception. *Curr Biol, 24*(17), 2066-

2070. doi:10.1016/j.cub.2014.07.060

Pitcher, D., Japee, S., Rauth, L., & Ungerleider, L. G. (2017). The Superior Temporal Sulcus

Is Causally Connected to the Amygdala: A Combined TBS-fMRI Study. *J Neurosci,*

37(5), 1156-1161. doi:10.1523/JNEUROSCI.0114-16.2016

Posse, S., Wiese, S., Gembris, D., Mathiak, K., Kessler, C., Grosse-Ruyken, M. L., . . .

Kiselev, V. G. (1999). Enhancement of BOLD-contrast sensitivity by single-shot

multi-echo functional MR imaging. *Magn Reson Med, 42*(1), 87-97.

doi:10.1002/(sici)1522-2594(199907)42:1<87::aid-mrm13>3.0.co;2-o

Pourtois, G., Sander, D., Andres, M., Grandjean, D., Reveret, L., Olivier, E., & Vuilleumier,

P. (2004). Dissociable roles of the human somatosensory and superior temporal

cortices for processing social face signals. *European Journal of Neuroscience, 20*(12),

3507-3515. doi:10.1111/j.1460-9568.2004.03794.x

Puce, A., Allison, T., Bentin, S., Gore, J. C., & McCarthy, G. (1998). Temporal cortex

activation in humans viewing eye and mouth movements. *J Neurosci, 18*(6), 2188-

2199. doi:10.1523/JNEUROSCI.18-06-02188.1998

Rafique, S. A., Solomon-Harris, L. M., & Steeves, J. K. (2015). TMS to object cortex affects

both object and scene remote networks while TMS to scene cortex only affects scene

networks. *Neuropsychologia, 79*(Pt A), 86-96.

doi:10.1016/j.neuropsychologia.2015.10.027

Rahnev, D., Nee, D. E., Riddle, J., Larson, A. S., & D'Esposito, M. (2016). Causal evidence

for frontal cortex organization for perceptual decision making. *Proc Natl Acad Sci U*

S A, 113(21), 6059-6064. doi:10.1073/pnas.1522551113

Ramot, M., Walsh, C., & Martin, A. (2019). Multifaceted Integration: Memory for Faces Is

Subserved by Widespread Connections between Visual, Memory, Auditory, and

Social Networks. *J Neurosci, 39*(25), 4976-4985. doi:10.1523/JNEUROSCI.0217-

19.2019

Rastogi, A., Cash, R., Dunlop, K., Vesia, M., Kucyi, A., Ghahremani, A., . . . Chen, R.

(2017). Modulation of cognitive cerebello-cerebral functional connectivity by lateral

cerebellar continuous theta burst stimulation. *NeuroImage, 158*, 48-57.

doi:10.1016/j.neuroimage.2017.06.048

Rezlescu, C., Barton, J. J., Pitcher, D., & Duchaine, B. (2014). Normal acquisition of

expertise with greebles in two cases of acquired prosopagnosia. *Proc Natl Acad Sci U*

S A, 111(14), 5123-5128. doi:10.1073/pnas.1317125111

Robertson, I. H., & Murre, J. M. (1999). Rehabilitation of brain damage: brain plasticity and

principles of guided recovery. *Psychol Bull, 125*(5), 544-575. doi:10.1037/0033-

2909.125.5.544

Rossion, B., Caldara, R., Seghier, M., Schuller, A. M., Lazeyras, F., & Mayer, E. (2003). A

network of occipito-temporal face-sensitive areas besides the right middle fusiform

gyrus is necessary for normal face processing. *Brain, 126*(Pt 11), 2381-2395.

doi:10.1093/brain/awg241

Sakurai, Y., Hamada, K., Tsugawa, N., & Sugimoto, I. (2016). Ventral simultanagnosia and

prosopagnosia for unfamiliar faces due to a right posterior superior temporal sulcus

and angular gyrus lesion. *Neurocase, 22*(1), 122-129.

doi:10.1080/13554794.2015.1066827

Shang, Y., Chang, D., Zhang, J., Peng, W., Song, D., Gao, X., & Wang, Z. (2019). Theta-

burst transcranial magnetic stimulation induced functional connectivity changes

between dorsolateral prefrontal cortex and default-mode-network. *Brain Imaging*

Behav. doi:10.1007/s11682-019-00139-y

Sliwinska, M. W., & Pitcher, D. (2018). TMS demonstrates that both right and left superior

temporal sulci are important for facial expression recognition. *NeuroImage, 183*, 394-

400. doi:10.1016/j.neuroimage.2018.08.025

Solomon-Harris, L. M., Rafique, S. A., & Steeves, J. K. (2016). Consecutive TMS-fMRI

reveals remote effects of neural noise to the "occipital face area". *Brain Res, 1650*,

134-141. doi:10.1016/j.brainres.2016.08.043

Stefanacci, L., & Amaral, D. G. (2000). Topographic organization of cortical inputs to the

lateral nucleus of the macaque monkey amygdala: a retrograde tracing study. *J Comp*

Neurol, 421(1), 52-79. doi:10.1002/(sici)1096-9861(20000522)421:1<52::aid-

cne4>3.0.co;2-o

Stefanacci, L., & Amaral, D. G. (2002). Some observations on cortical inputs to the macaque

monkey amygdala: an anterograde tracing study. *J Comp Neurol, 451*(4), 301-323.

doi:10.1002/cne.10339

Vuilleumier, P., Armony, J. L., Driver, J., & Dolan, R. J. (2001). Effects of attention and

emotion on face processing in the human brain: An event-related fMRI study.

Neuron, 30(3), 829-841. doi:10.1016/S0896-6273(01)00328-2

Vuilleumier, P., Armony, J. L., Driver, J., & Dolan, R. J. (2003). Distinct spatial frequency

sensitivities for processing faces and emotional expressions. *Nature Neuroscience,*

6(6), 624-631. doi:10.1038/nn1057

Wang, J. X., Rogers, L. M., Gross, E. Z., Ryals, A. J., Dokucu, M. E., Brandstatt, K. L., . . .

Voss, J. L. (2014). Targeted enhancement of cortical-hippocampal brain networks and

associative memory. *Science, 345*(6200), 1054-1057. doi:10.1126/science.1252900

Wang, X., Zhen, Z., Song, Y., Huang, L., Kong, X., & Liu, J. (2016). The Hierarchical

Structure of the Face Network Revealed by Its Functional Connectivity Pattern. *J*

Neurosci, 36(3), 890-900. doi:10.1523/JNEUROSCI.2789-15.2016

Zhang, H., Tian, J., Liu, J., Li, J., & Lee, K. (2009). Intrinsically organized network for face

perception during the resting state. *Neurosci Lett, 454*(1), 1-5.

doi:10.1016/j.neulet.2009.02.054

Zhu, Q., Zhang, J., Luo, Y. L., Dilks, D. D., & Liu, J. (2011). Resting-state neural activity

across face-selective cortical regions is behaviorally relevant. *J Neurosci, 31*(28),

10323-10330. doi:10.1523/JNEUROSCI.0873-11.2011

Figure Legends

Figure 1: Fisher Z transformed correlation magnitude changes from pre - post TMS to the (A)

rpSTS and (B) right motor cortex. Magnitudes are the Matrix Based Analysis (MBA) model

fits across the population. Changes that are likely to be greater than 0 with a posterior

probability of at least 95% are full squares and include an *. The squares' edge lengths

decrease linearly to their minimum size for a 90% likelihood or less. The green line marks the

ROI pairs that are within the pre-defined face-selective network. Fig. S2 contains an

unthresholded version of this figure and the interaction effect.

Figure 2: Correlations between the rpSTS ROI and ROIs in a pre-defined face network as

well as the bilateral primary motor hand regions. These data correspond to the first row of the

matrix in figure 1. The full matrix of bar plots in in Fig. S2. Magnitudes are the MBA model

fits across the population for each condition. Boxplots show 25-75% of the distribution. The

white line is the median. Whiskers are the maximum and minimum values excluding outliers.

MBA used to calculate posterior probabilities that a difference is greater than 0. The

correlations between the rpSTS and other face regions consistently decrease post rpSTS

stimulation, but not post rMotor stimulation.

Figure 3: Whole brain correlation changes (Fisher Z transformed) pre - post TBS stimulation.

Voxel coloring shows the magnitude of the correlation difference. Voxels are opaque for

$p>=0.01$ to transparent for $p=0$.The green outlines are voxels in the ROIs of the pre-defined

face network in at least 3 volunteers. For TMS to the rpSTS, many of the larger signal drops

are within the face network ROIs. TMS to right motor shows decreases across more of the

brain that isn't localized to the face network.

Figure 1

Fisher Z transformed correlation magnitude changes from pre - post TMS to the (A) rpSTS and (B) right motor cortex. Magnitudes are the Matrix Based Analysis (MBA) model fits across the population. Changes that are likely to be greater than 0 with a posterior probability of at least 95% are full squares and include an *. The squares' edge lengths decrease linearly to their minimum size for a 90% likelihood or less. The green line marks the ROI pairs that are within the pre-defined face-selective network. Fig. S2 contains an unthresholded version of this figure and the interaction effect.

Figure 2

Correlations between the rpSTS ROI and ROIs in a pre-defined face network as well as the bilateral primary motor hand regions. These data correspond to the first row of the matrix in figure 1. The full matrix of bar plots in in Fig. S2. Magnitudes are the MBA model fits across the population for each condition. Boxplots show 25-75% of the distribution. The white line is the median. Whiskers are the maximum and minimum values excluding outliers. MBA used to calculate posterior probabilities that a difference is greater than 0. The correlations between the rpSTS and other face regions consistently decrease post rpSTS stimulation, but not post rMotor stimulation.

Figure 3

Corr. to the rpSTS pre - post TMS

FisherZ
pre-post
TMS<-0.2

FisherZ
pre-post
TMS>0.2

Whole brain correlation changes (Fisher Z
transformed) pre - post TBS stimulation. Voxel
coloring shows the magnitude of the correlation
difference. Voxels are opaque for p>=0.01 to
transparent for p=0.The green outlines are voxels
in the ROIs of the pre-defined face network in at
least 3 volunteers. For TMS to the rpSTS, many
of the larger signal drops are within the face
network ROIs. TMS to right motor shows
decreases across more of the brain that isn't
localized to the face network.