Lytic infection with murine gammaherpesvirus 68 activates host and viral RNA polymerase III-dependent promoters to enhance non-coding RNA expression

Ashley N. Knox,[a] Alice Mueller,[b] Eva M. Medina,[a] Eric T. Clambey,[c] Linda F. van Dyk[a]#

[a] Department of Immunology and Microbiology, University of Colorado Anschutz Medical Campus, Aurora, Colorado, USA

[b] Department of Molecular, Cellular, and Developmental Biology, University of Colorado, Boulder, Colorado, USA

[c] Department of Anesthesiology, University of Colorado Anschutz Medical Campus, Aurora, Colorado, USA

Running Title: γHV68 infection alters pol III promoter activity

#Address correspondence to Linda van Dyk, linda.vandyk@cuanschutz.edu

Abstract: 218 words

Importance: 141 words

Text: 5,479 words

ABSTRACT

RNA polymerase III (pol III) transcribes multiple non-coding (nc) RNAs that are essential for cellular function. Pol

III-dependent transcription is also engaged during certain viral infections, including the gammaherpesviruses

(γHVs), where pol III-dependent viral ncRNAs promote pathogenesis. Additionally, several host ncRNAs are

upregulated during γHV infection and play integral roles in pathogenesis by facilitating viral establishment and

gene expression. Here we sought to investigate how pol III-dependent transcriptional activity was regulated

during gammaherpesvirus infection, using the murine gammaherpesvirus 68 (γHV68) system. To compare the

transcriptional regulation of host and viral pol III-dependent ncRNAs, we analyzed a series of pol III-dependent

promoters using a newly-generated luciferase reporter optimized to measure pol III activity. We measured

promoter activity from these constructs at the translation level via luciferase activity and at the transcription level

via RT-qPCR. We further measured endogenous ncRNA expression at single cell-resolution by flow cytometry.

These studies demonstrated that lytic infection with γHV68 increased the transcriptional activity of multiple host

and viral pol III-dependent promoters, and further identified the ability of accessory sequences to influence both

baseline and inducible promoter activity after infection. These studies highlight how lytic gammaherpesvirus

infection alters the transcriptional landscape of host cells to increase pol III-derived transcription, a process that

may further modify cellular function and enhance viral gene expression and pathogenesis.

IMPORTANCE

Gammaherpesviruses are a prime example of how viruses can alter the host transcriptional landscape to

establish infection. Despite major insights into how these viruses modify RNA polymerase II-dependent

generation of messenger RNAs, how these viruses influence the activity of host RNA polymerase III remains

much less clear. Small non-coding RNAs produced by RNA polymerase III are increasingly recognized to play

critical regulatory roles in cell biology and virus infection. Studies of RNA polymerase III dependent transcription

are complicated by its use of multiple promoter types and diverse RNAs with variable stability and processing

requirements. Here, we established a reporter system to directly study RNA polymerase III-dependent promoter

responses during gammaherpesvirus infection and utilized single-cell flow cytometry-based methods to reveal

that gammaherpesvirus lytic replication broadly induces pol III activity to enhance host and viral non-coding RNA

expression within the infected cell.

INTRODUCTION

Gammaherpesviruses (γHVs) are large, lymphotropic viruses that establish a life-long infection in their hosts, with long-term latency in lymphocytes (1, 2). The γHVs include the human pathogens, Epstein-Barr virus (EBV) and Kaposi's sarcoma-associated herpesvirus (KSHV or HHV-8), and murine gammaherpesvirus 68 (γHV68 or MHV-68; ICTV nomenclature *Murid herpesvirus 4*, MuHV-4) (3). These viruses establish a primary lytic infection in their host that is followed by a prolonged quiescent infection termed latency. Latency is maintained in healthy individuals by a homeostatic relationship between the virus and the host immune response; if this balance is disrupted (e.g., by immunosuppression), γHVs can reactivate from latency and actively replicate. Disruption between the balance of γHV infection and host immune control is associated with γHV multiple pathologies, including a range of malignancies (4).

The γHVs contain several types of non-coding (nc) RNAs, including nuclear ncRNAs and functional miRNAs; these diverse RNAs including ncRNAs transcribed by RNA polymerase II (e.g. the KSHV PAN RNA and the KSHV and EBV miRNAs) or by RNA pol III (e.g. the EBV-encoded small RNAs (EBERs) and the γHV68 tRNA-miRNA-encoded RNAs (TMERs)) (5-13). Viral ncRNAs are considered to have important host-modulatory functions, interacting with host proteins and regulating host and viral gene expression. For example, the EBV EBERs are highly expressed during latency, and were discovered through their interaction with the host lupus-associated antigen (La) protein, which putatively mediates EBER interaction with TLR3 (14-17).The EBERs have further been shown to interact with several host proteins including ribosomal protein L22, protein kinase R (PKR), and retinoic-acid inducible gene I (RIG-I) (18). These interactions can trigger sustained host innate immune responses that are implicated in the development of EBV-associated malignancies (16, 19-21). γHV68, a highly tractable small animal model of γHV infection, also encodes several pol III-transcribed ncRNAs known as the tRNA-miRNA-encoded RNAs (TMERs) (22, 23). The TMERs are dispensable for lytic replication and establishment of latency; however, these transcripts are required for pathogenesis during acute infection of an immunocompromised host (7, 24-26). The TMERs contain bi-functional elements with a tRNA-like structure on the 5' end and hairpin loops that are processed into biologically-active miRNAs (7), capable of targeting a number of RNAs for post-transcriptional regulation (27). Our lab has previously shown that the tRNA-like structure is sufficient to rescue pathogenesis of a TMER deficient viral recombinant, suggesting that like the EBERs, the TMERs may contribute to pathogenesis through their interactions with host proteins (26). Though TMER-host protein interactions have yet to be fully explored, it is notable that several characteristics of the EBERs, such as a 5'-triphosphate and 3'-polyU, are imparted by RNA polymerase III (pol III) transcription (28).These motifs can be recognized by host RNA-binding proteins, such as RIG-I or La, to trigger an innate immune response (16, 20, 28, 29).

Pol III is often considered to perform "house-keeping" functions, as it transcribes host genes required for cell growth and maintenance (e.g. U6 snRNA, tRNAs, and 5S rRNA) (30). Despite this, it is clear that the γHVs

can usurp pol III-dependent transcription mechanisms for their own purposes. Latent EBV infection has been
shown to upregulate components of pol III and ultimately increase the expression of host pol III transcripts –
particularly vault RNAs – that allow increased establishment of viral infection and gene expression (31-33).
Similarly, γHV68 infection drives upregulation of host pol III-dependent short-interspersed nuclear element
(SINE) RNAs, which in turn, mediate increased viral gene expression (34, 35). Additionally, our lab has reported
that reactivation of a latently-infected γHV68 cell line results in increased expression of TMERs within a rare
subset of the population (36). Notably, dysregulation of pol III is a common feature of many cancer cells,
implicating γHV infection-driven alteration of pol III activity as one potential contributor to γHV-associated
malignancies (37). Therefore, understanding how γHV infection alters pol III activity is integral to elucidating
mechanisms of γHV pathogenesis.

Due to the importance of γHV ncRNAs during infection and the unique transcriptional regulation afforded
by RNA pol III, the overall objective of this study was to characterize the activity of host and viral pol III-dependent
promoters during lytic γHV68 infection. To this end, we used a luciferase-reporter system optimized to report pol
III-dependent transcriptional activity to investigate how virus infection regulated pol III promoter activity *in vitro*.
We found that γHV68 infection upregulates the activity of multiple viral and host pol III promoters, a process
further associated with the induction of pol III-dependent targets. These studies indicate that lytic γHV infection
can broadly enhance RNA pol III promoter activity to modify the ncRNA landscape of infected cells.

RESULTS

RNA polymerase III can transcribe RNA from a variety of gene-internal (type 1 and 2) and gene-

external (type 3) promoters (Fig 1A). These promoters contain distinct motifs that determine which transcription

factors bind to the promoter to recruit pol III (38). To understand how γHV lytic replication influenced RNA pol

III promoter activity without the confounding factors of RNA sequence or stability, we sought to study a series

of viral and host pol III promoters using a luciferase assay system previously used to study pol III promoter

activity (39). Several host and viral pol III promoters encompassing a variety of promoter types and motifs were

selected for analysis (Supplemental Figure 1). Pol III promoters included host (U6, tRNA-Tyr, and vault RNA 1-

1) and viral promoters (γHV68 TMER1, TMER4, TMER5; EBV EBER1 and EBER2; and adenovirus VA1). A

subset of studies further compared promoter variants, encompassing either the minimal promoter elements

("minimal") or including additional accessory sequences ("full"). See Supplemental Figure 1 for promoter

schematics, and Table 1 for a list of all promoters examined.

To analyze the activity of pol III promoters by luciferase assay, we selected the reporter plasmid pNL1.1

(Promega), as the NanoLuc luciferase in this reporter creates a brighter signal and the protein is smaller than

its Renilla and Firefly counterparts; the smaller protein size is more consistent with pol III processivity of small

ncRNAs. Our analysis of the pNL1.1 sequence revealed a pol III termination signal within the luciferase coding

gene (TTTT). Therefore, to examine the activity of pol III promoters without the potential for early termination,

we introduced silent mutations into the NanoLuc reporter construct to remove the termination signal. This new

vector was named "pNLP3" to reflect that it is a _NanoLuc_ reporter optimized for _p_ol _III_ (Fig 1B). The human U6

promoter was cloned into both pNL1.1 and pNLP3 to compare the effects of removing the pol III termination

signal, with promoter activity measured 24 hours post-infection. We found that removal of the termination

signal increased the luciferase output, indicating that there was more read-through of the full NanoLuc gene

from pNLP3 (Fig 1B). Furthermore, RT-PCR analysis of the NanoLuc transcript transcribed from the U6

promoter in either the pNL1.1 or pNLP3 vector revealed more full-length NanoLuc transcript from the pNLP3

vector (Fig 1C). This indicates that the pNLP3 vector allows for optimal pol III transcription of the reporter gene.

We therefore used the pNLP3 vector as the backbone for analysis of all other pol III promoters included in this

study.

With an optimized pol III reporter construct, we assessed how different pol III promoters respond to

γHV68 infection over time. We first compared the activity of the human U6 (type 3) and γHV68 TMER1 (type 2)

promoters. HEK 293 cells were transfected with pNLP3 vectors containing either the U6 or TMER1 promoter,

co-transfected with an SV40 (pol II promoter)-driven Firefly luciferase vector, then infected with γHV68 (Fig

2A). Cell lysates were collected every 4 hours for 24 hours post-infection to quantify promoter activity as

defined by NanoLuc luciferase activity. This analysis revealed that γHV68 infection resulted in a time-

dependent increase in NanoLuc activity for both the U6 and TMER1 promoters (left panel, Fig 2B-2C) relative

to mock-infected samples. γHV68 infection also resulted in a time-dependent increase in expression of the

control, Firefly luciferase reporter (second panel from left, Fig 2B-2C). The results for dual luciferase assays

are typically reported as relative luminescence units (RLUs), where the reporter luminescence units (LUs) are

normalized to the luminescence units of the control luciferase (i.e., NanoLuc LUs / Firefly LUs). However, since

γHV68 infection simultaneously increased luminescence from both the NanoLuc reporter, and from the control

Firefly reporter, this normalization errantly implied decreased relative U6 promoter activity with infection (Fig

2B). Given the limitation of standardizing pol III promoter activity relative to a pol II promoter control (i.e. SV40

promoter), all subsequent analyses report promoter activity as a fold change in NanoLuc luminescence

comparing mock and γHV68 infected samples. We found that the U6 promoter drives high basal luciferase

activity under mock conditions (left panel, Fig 2B), with a further increase in raw and normalized U6-expressed

NanoLuc LUs throughout infection (Fig 2B). In contrast, the TMER1 promoter was characterized by extremely

low basal luciferase activity (left panel, Fig 2C) in mock conditions; however, this promoter was strongly

induced by infection (Fig 2C). These data suggest that γHV68 lytic infection increases the activity of multiple

pol III promoter types, with a more robust induction of the type 2 promoter of TMER1 compared to the U6

promoter.

We further analyzed the activity of several other promoter types to assess how they are regulated

during γHV68 infection. Experiments indicated that γHV68 infection induced activity from multiple pol III

promoters, including the human U6 and tRNA-Tyr promoters, the EBER1 and EBER2 promoters, and TMER1,

4, and 5 promoters (Fig 3). Though the vaultRNA1-1 promoter was cloned into the reporter, there was no

detectable activity from this construct (unpublished data). These promoters have different levels of baseline

NanoLuc luciferase activity under mock conditions, with the U6 and EBER promoters showing high activity,

and the TMER promoter displaying minimal detectable activity under mock conditions (unpublished data). Of

the promoters tested, the TMER promoters consistently showed the greatest induction of luciferase activity

during infection.

We compared the sequences of TMER promoters to identify which features of these promoters could

potentially contribute to the strong induction of activity during infection. The TMER promoters initially analyzed

contained the TMER promoter, as well as extended sequence around the minimal promoter elements (see

Supplementary Figure 1 for diagrams). Considering that the extra sequence included in these "full" promoters

may contribute to infection-induced activity, we created a panel of "minimal" TMER promoters that contain only

the sequence canonically identified as the minimal promoter elements necessary for transcription, i.e. the

sequence beginning from the A box to the end of the B box (Supplemental Figure 1). Additionally, we removed

the upstream sequence from the EBER promoters; these EBER minimal promoters were used to further

analyze how extended sequences can alter promoter induction during infection. As previously described, these

constructs were transfected into HEK 293 cells, then infected with γHV68. The fold change in NanoLuc activity

relative to mock-treated samples was compared after 24 h of infection (Fig 4). When we compared the relative

inducibility of EBER "full" versus "minimal" promoters, "minimal" promoters showed greater virus-inducibility.

This enhanced inducibility of the "minimal" EBER promoters likely reflects the reduced baseline luminescence

from these promoters (unpublished data). Conversely, TMER minimal promoters displayed a weaker induction

during infection than their "full" counterparts, suggesting the sequence surrounded the TMER minimal

promoters drives stronger expression during infection. These results indicate that the sequence surrounding

minimal pol III promoter elements impacts both the baseline activity and inducibility of these promoters during

infection.

Luciferase readouts of pol III promoter activity allowed us to uniformly analyze pol III promoter activity.

This assay does not directly measure the level of RNAs, however, instead relying on an enzymatic readout of

luciferase protein activity. To ensure that γHV68 infection was inducing pol III activity transcriptionally, we used

the same NanoLuc constructs to measure promoter activity at the RNA level by performing RT-qPCR for the

NanoLuc transcript. Following the same protocol as used for the luciferase assays, HEK 293 cells were

transfected with pGL3 and the pNLP3 vector expressed by pol III promoters of interest (as outlined in Fig 2).

Cells were then infected with γHV68 and RNA was purified from cells 16 or 24 h post-infection. Primers

targeting the NanoLuc gene were used for qPCR following reverse transcription of the RNA. Infection

increased the NanoLuc RNA expression from the U6 and TMER1 promoters, with more modest induction from

the EBER promoters (Fig 5A). These results indicate that γHV68 infection stimulates pol III-promoter activity

from multiple host and viral promoters, measured at both the transcriptional and translational level. To extend

these findings, we further measured NanoLuc RNA expression from "minimal" or "full" TMER promoters. These

studies demonstrated the γHV68 infection increased NanoLuc RNA from the "minimal" promoter relative to

mock infected samples, with further RNA induction from the "full" TMER promoter. These results strongly

suggest that the NanoLuc reporter assay serves as a faithful readout for pol III-dependent transcription,

quantified at both the RNA and protein level. These findings also emphasize that sequences outside of the

minimal TMER promoters contribute to increased expression during infection.

γHV lytic replication critically depends on viral DNA replication and late gene transcription, processes

that are inhibited by phosphonoacetic acid (PAA) (40, 41). We therefore tested the impact of PAA on virus-

induced pol III induction. To do this, HEK 293 cells were transfected with pNLP3-TMER1 and infected as

before, with one set of samples receiving PAA treatment (200μg/mL) after 1 h of viral inoculation. RNA was

isolated 16 h post-infection and RT-qPCR was performed to detect the NanoLuc transcript. Notably, treatment

with PAA during infection had no impact on the induction of NanoLuc RNA compared to γHV68 infected cells,

indicating that viral DNA replication and late gene synthesis was not required for pol III induction (Fig 6A).

Though PAA treatment had no effect on the level of NanoLuc RNA levels (i.e. transcription), PAA treatment

was consistently associated with increase luciferase enzymatic activity, with PAA-treated γHV68-infected

cultures characterized by a greater apparent induction of luciferase activity compared to γHV68-infected

cultures alone. This PAA-driven enhancement of luciferase activity was observed for multiple pol III promoters,

including U6, TMER1, 4 and 5 and EBER 1 and 2 (Fig 6B). The increase in luciferase activity following PAA

treatment, with minimal impact on NanoLuc RNA, strongly suggests that PAA treatment enhanced the

translational output from the promoters tested. These data suggest that viral late gene expression plays an

additional role in translation that is not seen at the transcriptional level, a phenomenon independent of pol III

promoter activity.

Given the reported relationship between the NF-κB pathway and the expression of pol III-dependent

transcripts (31), we analyzed the effect of NF-κB activation or inhibition on the activity of the U6 and TMER1

promoters via luciferase activity. First, we measured induction of an NF-κB reporter plasmid following treatment

with either TNFα, a known inducer of the NF-κB pathway, or following γHV68 infection. Whereas TNFα

induced NF-κB reporter activity at 4 and 24 hours post-treatment, γHV68 infection had no measurable impact

on expression from the NF-κB reporter (Supplemental Figure 2A). Next, we analyzed the impact of NF-κB

manipulation on pol III promoter activity. Treating cells with TNFα modestly increased U6 promoter activity,

albeit to a lesser extent than γHV68 infection (Supplemental Figure 2B). TMER1 promoter activity was not

affected by TNFα treatment. Inhibition of NF-κB with the BAY 11-7082 (BAY 11) compound increased U6-

expressed luciferase activity in virus infected conditions, yet had no significant impact on TMER1 promoter

activity after infection (Supplemental Figure 2C). This indicates a potential role of NF-κB in inhibiting pol III

promoter activity during infection; however, this effect is only observed in the case of a gene-external (i.e. type

3) promoter. Ultimately, these data do not support a significant role of the NF-κB pathway in the observed

induction of pol III promoter activity after γHV68 infection.

While our analysis of pol III promoter activity suggested a general induction during infection, a previous

report suggested that only a subset of host pol III-transcribed genes – the SINE RNAs – are increased during

lytic γHV68 infection (34). These studies were largely based on analysis of total RNA of infected cells. To gain

further insight into endogenous host ncRNA expression during lytic γHV68 infection, we made use of the

PrimeFlow Assay kit, which allows for flow cytometry-based quantitation of RNA expression at the single-cell

level. Murine fibroblast cells (3T12) were mock-treated, infected with wild-type (WT) γHV68, or infected with an

EBER-knock in (EBER-KI.γHV68) recombinant γHV68 that lacks the TMERs and instead contains insertion of

the EBV EBERs into the TMER locus. EBER-KI.γHV68 was competent for viral replication (manuscript in

progress). 3T12 cells were infected with an MOI of 1 and harvested at 16 h post-infection, conditions that result

in a mixed population of virally infected and uninfected cells. Cells were then queried with fluorescent probes to

detect viral gene expression as defined by either the γHV68 TMERs or the EBV EBERs, expressed from the

context of the γHV68 genome. Host ncRNA for U6 snRNA or 4.5 rRNA (the murine equivalent of human 5S

rRNA) expression was then assessed, comparing expression between cells with virus expression (i.e. TMER+

or EBER+) and cells lacking virus expression (i.e. TMER- or EBER-). Gating schemes and probe specificity are

shown in Supplemental Figure 3. Notably, probes for the TMERs and EBERs were specific for their intended

targets, with no detectable TMER or EBER expression in mock-infected samples, TMER+ cells only present in

WT γHV68 infected samples, and EBER+ cells only present in EBER-KI.γHV68 infected samples. We next

compared the relative expression of the U6 (Fig 7A-B) and 4.5S ncRNAs (Fig 7C-D) between cells with active

viral RNA expression versus those cells that did not express these viral ncRNAs. Analysis of the geometric

mean fluorescence intensity (gMFI) for the U6 snRNA and 4.5 rRNA probes revealed increased expression of

U6 and 4.5 rRNA ncRNAs in virally infected cells (i.e. TMER+ or EBER+) compared to uninfected cells (TMER-

or EBER-) (Fig 7E-F). These data demonstrate an increase in host pol III-transcribed ncRNAs in γHV68-

infected cells and emphasize the benefit of single cell analysis of endogenous RNA expression to quantify

ncRNA expression during virus infection.

DISCUSSION

While γHV infection is known to alter expression of host genes, and viral ncRNAs are integral for

pathogenesis, the transcriptional regulation of these ncRNAs has remained unclear. Here, we propose that

different pol III promoter types allow for distinct means for transcriptional regulation during infection. This study

focused on the impact of γHV68 lytic replication on pol III promoter activity, identifying that lytic infection drives

a general upregulation of promoter activity across multiple host and viral pol III-dependent transcripts. This

upregulation was measured by a luciferase assay optimized for pol III transcription, with increased pol III-

dependent promoter activity quantified by luciferase activity and RNA expression, with complementary findings

revealed through the use of flow cytometric analysis of ncRNA expression. These findings emphasize the utility

of the modified NanoLuc luciferase system to analyze pol III promoter activity, and provide clear evidence for

pol III promoters with large differences in basal and inducible promoter activity. They further emphasize the

capacity of γHV lytic infection to modify pol III-dependent transcriptional machinery in infected cells, a process

that likely facilitates productive virus replication (34). At this time, it remains unknown whether pol III machinery

or transcription is altered during γHV68 latent infection or reactivation from latency.

Our use of a luciferase reporter to measure the activity of pol III promoters allowed us to directly

compare the functional activity of multiple pol III promoters, while avoiding potential differences that may arise

due to ncRNA sequence, structure, or stability. Though pol III promoters conventionally drive expression of

non-coding RNAs, there is clear precedent that pol III can transcribe translation-competent RNA (42, 43), and

luciferase reporters have been used for high throughput and unbiased analysis of pol III promoter activity (44,

45). Inspired by these studies, we cloned several host and viral ncRNA promoters into a NanoLuc luciferase

reporter to measure their activity during lytic γHV68 infection. We chose the NanoLuc luciferase as our To

further enhance the robustness of this reporter, we identified and removed a pol III termination sequence within

the NanoLuc gene which approximately doubled luciferase reporter activity. Though pol III transcription should

theoretically be terminated in the original NanoLuc reporter, pol III read-through of termination signals has

been reported (46). In total, use of the modified NanoLuc reporter construct afforded a sensitive and robust

readout for assessing pol III promoter activity.

Through use of this pol III reporter assay, we found that γHV68 infection increased promoter activity

across a range of host and viral pol III promoters. This suggests that γHV68 infection generally upregulates pol

III activity. Interestingly, the consequence/magnitude of induction elicited by infection varied between

promoters. For example, the U6 promoter conveyed high basal activity with infection resulting in a modest

induction of U6 promoter activity. Conversely, the TMER promoters exhibited extremely low basal activity in

mock-infected conditions, with dramatic induction after γHV68 infection. The inducibility of the TMERs was

further enhanced by accessory sequences outside of the minimal A and B box elements. One explanation for

this enhanced induction is that these extended sequences may contain additional transcriptional elements that

are integral to the promoter itself. While formally possible, it is notable that accessory sequences across the

TMERs are not conserved (Supplemental Fig 4A). An alternate explanation for the enhanced activity of the full

TMER promoters is that inclusion of the extended sequence includes the full tRNA-like structure of the TMER

genes (Supplemental Fig 4B). It is interesting to speculate that this tRNA-like structure could either lend

greater stability to transcripts, or protect the transcripts from degradation by host exonucleases or the viral

endonuclease, muSOX (47). Although pol III transcription is frequently associated with the transcription of

housekeeping ncRNAs, there is clear precedent that pol III can also participate in inducible gene expression

(e.g. γHV68-induced expression of SINEs) (34, 35). Whether the TMERs have conserved regulatory

mechanisms with host inducible ncRNAs is currently unknown, however, the TMERs share more promoter

similarity to the SINEs (type 2, gene internal) than to the vault RNAs (type 3, gene external).

Our studies demonstrated that infection increased NanoLuc expression at both the RNA and protein

level, indicating that virus infection increased pol III promoter activity and not some secondary measurement.

This induction was further associated with increased expression of the host ncRNAs U6 and 4.5 rRNA. While

these findings emphasize the impact that infection has on pol III-dependent transcription, our studies did reveal

one context in which infection impacted a post-transcriptional measurement. Unexpectedly, inhibition of viral

late gene expression with PAA resulted in increased luciferase activity from nearly all of the promoters

examined – this phenomenon was only seen at the level of NanoLuc protein activity, not at the level of

NanoLuc RNA. The ability of PAA to enhance NanoLuc protein activity, with no commensurate change in RNA

expression, suggests that viral late genes may have a possible role in tempering translation. Manipulation of

host translational machinery by the herpesviruses is a common strategy that is required for optimal virus

replication and the production of virus progeny (48).

Previous reports show that γHV infection can have diverse effects on pol III transcription, ranging from

a general induction of pol III machinery (e.g. in the context of EBV and EBNA1 (33, 49), to the selective

induction of pol III-transcribed RNAs (e.g. induction of specific host vaultRNAs in EBV latently infected cells

(31, 32), and the host SINE RNAs in γHV68-infected cells (34)). One challenge in interpreting these different

findings is that these studies have been done in different states of infection (latent versus lytic), in different cell

types, and in different states of cellular transformation. In many cases, the mechanistic insights gained from

these studies could only be gained through the use of in vitro studies. Future studies will benefit from direct

comparisons of primary virus infection, when possible, coupled with the analysis of endogenous ncRNA

expression. In keeping with this, we anticipate that the γHV68 system will afford unique insights in how the

γHVs regulate pol III-dependent ncRNA expression, allowing the analysis of primary infection coupled with

technologies to measure endogenous ncRNA expression. For example, our single-cell analysis of pol III-

derived transcripts – U6 snRNA and 4.5 rRNA – supported that γHV68-infection not only increases pol III-

dependent promoter activity, but also increases the endogenous expression of these transcripts. How host and

viral ncRNAs are regulated as a function of cell type and virus stage of infection remains an important

unanswered question.

In total, our studies revealed a γHV68-dependent induction in the activity of host and viral pol III

promoters. This induction was seen in the expression of a reporter gene, as well as in the endogenous

expression of pol III-dependent transcripts. Though previous reports have focused on the virus-mediated

upregulation of specific host ncRNAs, these experiments suggest a broader effect of lytic γHV68 infection on

pol III activity. This suggests that γHV68 modulation of the host transcriptional landscape goes beyond mRNA

regulation, and that pol III-dependent transcripts are likely to play a wider role in γHV68 pathogenesis than

previously appreciated.

METHODS

Viruses and tissue culture. All viruses were derived from a BAC-derived WT γHV68 (50 Koszinowski

UH. 2000) For some experiments, the TMER total knock-out (TMER-TKO) virus was used; this virus was

generated as previously described (26). Viruses were propagated and titered as previously described (26).

EBER-KI virus contains EBERs 1 and 2 in the TMER-TKO virus backbone. Its generation and characterization

are described in a manuscript in preparation.

Human endothelial kidney (HEK 293) and murine fibroblast (3T12) cells were cultured in Dulbecco's

modified Eagle medium (DMEM; Life Technologies) supplemented with 5% fetal bovine serum (FBS; Atlanta

Biologicals), 2 mM L-glutamine, 10 U/ml penicillin, and 10 µg/ml streptomycin sulfate (complete DMEM). Cells

were cultured at 37°C with 5% CO_2.

Mutagenesis of pNL1.1 to create the pNLP3 NanoLuc luciferase reporter. The promoterless

NanoLuc luciferase reporter vector pNL1.1[*Nluc*] was obtained from Promega, and primers were designed to

introduce silent mutations to remove the pol III termination signal in the NanoLuc coding sequence; these

primers are listed in Supplementary Table 1. Mutagenesis PCR was performed with the following cycles: (i) 95°

for 30s, (ii) 12 cycles of 95°C for 30s, 55°C for 1 min, 68°C for 3 m. The resulting DNA was digested with DpnI

(New England Biolabs Inc) and transformed into XL1-Blue super-competent cells (Agilent). Bacterial colonies

were sequenced to confirm the correct mutations. The resulting plasmid was named "pNLP3" to indicate that it

is a NanoLuc plasmid optimized for pol III.

Generating a pol III promoter-driven NanoLuc reporter panel. All promoters were generated to

include XhoI and HindIII overhang sequences on the 5' and 3' ends respectively. Several promoters were

constructed using ligated oligonucleotides. PCR-amplified promoters and pNLP3 were digested with XhoI and

HindIII, then promoters were ligated into pNLP3 using T4 DNA ligase (New England BioLabs Inc). See

Supplementary Table 1 for primers used to PCR-amplify promoters. Ligated constructs were transformed into

One Shot electro- or chemically competent TOP10 *E. coli* cells (Thermo Fisher Scientific, #C404052 or

#C404010), which were then plated at several dilutions on LB agar containing ampicillin. Resulting colonies

were expanded in LB broth with ampicillin and plasmid was isolated using the QIAprep Spin Miniprep Kit

(Qiagen). All constructs were confirmed by sequencing.

Transfecting cells. For transfections, HEK 293 cells were cultured in 5% FBS DMEM without penicillin

or streptomycin for approximately 24 hours. Transfection solutions contained Opti-MEM (Thermo Fisher

Scientific), NanoLuc plasmid (pNLP3 with inserted pol III promoters), and the Firefly control plasmid (pGL3-

Control; Promega). After plasmids were added to the Opti-MEM, solutions were incubated with X-tremeGENE

HP DNA Transfection Reagent (Sigma-Aldrich) for at least 15 minutes at room temperature. Transfections

were performed in several plate formats, depending on the downstream use; transfection solutions were added

dropwise to the appropriate wells (10uL of solution for 96-well plate, 100uL of solution in 12-well plates, 200uL

of solution in 6-well plates). For all transfections, the molar ratio of NanoLuc plasmid to Firefly plasmid was

kept at 10:1; the total amount of DNA transfected per well was adjusted depending on the plate size

(approximately 10ng for 96-well, 100ng for 12-well, and 200ng for 6-well). Cells were incubated with

transfection solution for 24 hours prior to downstream applications, unless otherwise stated.

To analyze how promoter activity was affected by γHV68 infection, transfected HEK 293 cells were

infected with WT, TMER-TKO, or EBER-KI γHV68 at a multiplicity of infection (MOI) of 1 plaque forming unit

per cell. Cells were cultured for approximately 24 h prior to infection. Cell counts were determined by treating

with 0.05% Trypsin-EDTA (Life Tech, #25300-054) to remove and cells. These cells were mixed with Trypan

Blue dye (Bio-Rad, #145-0021) to obtain a live cell count using the TC20 Automated Cell Counter (Bio-Rad).

Virus stocks were mixed with 5% complete DMEM, then added to cells and incubated with virus for 1 hour.

Viral inoculum was then removed and replaced with complete 5% DMEM. For samples treated with

phosphonoacetic acid (200 µg/mL, Sigma-Aldrich #284270), inoculum was removed and replaced with 5%

complete DMEM containing the drug.

Dual luciferase assays. Following transfection and infection, HEK 293 cell lysates were collected for

analysis of luciferase activity. All luciferase assays were performed using the Nano-Glo® Dual-Luciferase®

Reporter Assay System (Promega). For experiments performed in the 12-well format, supernatant was removed

and cells were scraped, collected into 1.5mL tubes, then pelleted. Cell pellets were resuspended in 250 µL of

Passive Lysis Buffer, per the manufacturer's protocol. 80 µL of the cell suspensions were used for assays. For

luciferase assays performed in the 96-well format, 80 µL of ONE-Glo™ EX Reagent was added directly to the

cells and supernatant, as recommended by the manufacturer. Samples were incubated at room temperature

while on a shaker for three minutes. This solution was then transferred from a transparent 96-well culture plate

to a white-wall luminescence plate prior to reading Firefly luciferase activity on the Tecan Infinite® 200 PRO

plate reader. Then, 80 uL of the NanoDLR™ Stop & Glo® Reagent was added to the solution and incubated at

room temperature on a shaker for at least 10 minutes. Samples were read on the Infinite® 200 PRO again for

the NanoLuc luminescence measurement.

RT-qPCR of NanoLuc RNA. RNA was isolated from transfected HEK 293 cells with TRIzol® Reagent

(Thermo Fisher Scientific, #15596026) and DNase-treated with TURBO™ DNase (Invitrogen, #AM2238)

following the manufacturer's protocols. RNA amplification and removal of DNA was confirmed by RT-PCR or

PCR amplification of a control host gene, 18S. RT-PCR was performed using the OneStep RT-PCR Kit (Qiagen,

#210212), with the following conditions: (i) $50^{o}C$ for 30 m, (ii) $95^{o}C$ for 15 m, (iii) 40 cycles of $94^{o}C$ for 30 s, $52^{o}C$

for 30 s, and $72^{o}C$ for 30 s, (iv) $72^{o}C$ for 10 m, (v) hold at $4^{o}C$. PCR was performed using Taq DNA polymerase

(Qiagen, #201205) with the following conditions: (i) $95^{o}C$ for 5 min, (ii) 40 cycles of $94^{o}C$ for 30 s, $52^{o}C$ for 30 s,

$72^{o}C$ for 30 s, (iii) $72^{o}C$ for 10 m, (iv) hold at $4^{o}C$. RNA samples that showed no product following PCR

amplification were deemed DNA-free, and converted to cDNA using SuperScript III Reverse Transcriptase

(Invitrogen, #18080093) following the manufacturer's protocol. 100ng of the cDNA was then used for qPCR
analysis of the NanoLuc and 18S genes using the iQ™ SYBER® Green Supermix (Bio-Rad, #1708880) with the
following conditions: i) 95°C for 3 m, ii) 40 cycles of 95°C for 15 s, 60°C for 1 m, iii) 95°C for 15 s, 60°C for 1 m,
95°C for 15 s. Amplification of NanoLuc was normalized to 18S expression to calculate the relative difference of
NanoLuc expression using the Pfaffl method: $\frac{NanoLuc\ Primer\ Efficiency^{Target\ \Delta Ct}}{18S\ Primer\ Efficiency^{18S\ \Delta Ct}}$ (51).

PrimeFlow RNA assay. The PrimeFlow RNA assay kit (Thermo Fisher Scientific, # 88-18005-210) was
used to analyze the endogenous expression of non-coding RNAs in mock and γHV68-infected 3T12 cells. Cells
were infected at an MOI of 1 for 16 h, then processed with the PrimeFlow kit following the manufacturer's
protocol. Cells were treated with probes for TMERs (Type 4/AF488), EBERs (Type 1/AF647), and U6 snRNA or
4.5 rRNA (Type 6/AF750). Samples were collected on an LSR II Flow Cytometer (BD Biosciences), and included
single stain and "full minus one" controls. Flow cytometry data were analyzed using FlowJo software (version
10.6.1), with compensation based on single stained beads and cells. Compensated flow cytometry data were
subsequently analyzed for singlet events (based on doublet discrimination), with positive and negative
populations based on control samples.

Software and statistical analysis. Statistical analysis and graphing were done in GraphPad Prism
(Version 8.0d). Statistical significance was tested by unpaired t test (comparing two conditions) or one-way
ANOVA (comparing three or more conditions) and subjected to multiple corrections tests using recommended
settings in Prism. All flow cytometry data were analyzed in FlowJo (version 10.6.1) with flow cytometry data
shown as pseudo-color dot plots.

ACKNOWLEDGEMENTS

This work was supported by the NIH grants R01AI121300 to LFvD and R21 AI134084 to LFvD and ETC,

T32 AI052066 to ANK and CO RNA Biosciences summer internship support to AM.

We thank the members of the Clambey and van Dyk lab for helpful discussions, members of the Colorado

RNA Bioscience Initiative for their insights, the Colorado ClinImmune core for flow cytometry services and the

Genomics Shared Resource of the University of Colorado Cancer Center which receives direct funding support

from the National Cancer Institute through Cancer Center Support Grant P30CA046934.

REFERENCES

1.   Longnecker R, Kieff E, Cohen JI. 2013. Epstein-Barr Virus, Fields Virology, 6th ed. Wolters Kluwer Health
Adis (ESP).
2.   Ganem D. 2006. KSHV infection and the pathogenesis of Kaposi's sarcoma. Annu Rev Pathol 1:273-96.
3.   Barton E, Mandal P, Speck SH. 2011. Pathogenesis and host control of gammaherpesviruses: lessons
from the mouse. Annu Rev Immunol 29:351-97.
4.   Cesarman E. 2011. Gammaherpesvirus and lymphoproliferative disorders in immunocompromised
patients. Cancer Lett 305:163-74.
5.   Samols MA, Hu J, Skalsky RL, Renne R. 2005. Cloning and identification of a microRNA cluster within
the latency-associated region of Kaposi's sarcoma-associated herpesvirus. J Virol 79:9301-5.
6.   Pfeffer S, Sewer A, Lagos-Quintana M, Sheridan R, Sander C, Grasser FA, van Dyk LF, Ho CK, Shuman
S, Chien M, Russo JJ, Ju J, Randall G, Lindenbach BD, Rice CM, Simon V, Ho DD, Zavolan M, Tuschl
T. 2005. Identification of microRNAs of the herpesvirus family. Nat Methods 2:269-76.
7.   Diebel KW, Smith AL, van Dyk LF. 2010. Mature and functional viral miRNAs transcribed from novel RNA
polymerase III promoters. RNA 16:170-85.
8.   Sun R, Lin SF, Gradoville L, Miller G. 1996. Polyadenylylated nuclear RNA encoded by Kaposi sarcoma-
associated herpesvirus. Proc Natl Acad Sci U S A 93:11883-8.
9.   Cai X, Lu S, Zhang Z, Gonzalez CM, Damania B, Cullen BR. 2005. Kaposi's sarcoma-associated
herpesvirus expresses an array of viral microRNAs in latently infected cells. Proc Natl Acad Sci U S A
102:5570-5.
10.   Zhu JY, Strehle M, Frohn A, Kremmer E, Hofig KP, Meister G, Adler H. 2010. Identification and analysis
of expression of novel microRNAs of murine gammaherpesvirus 68. J Virol 84:10266-75.
11.   Reese TA, Xia J, Johnson LS, Zhou X, Zhang W, Virgin HW. 2010. Identification of novel microRNA-like
molecules generated from herpesvirus and host tRNA transcripts. J Virol 84:10344-53.
12.   Kim DN, Lee SK. 2012. Biogenesis of Epstein-Barr virus microRNAs. Mol Cell Biochem 365:203-10.
13.   Rosa MD, Gottlieb E, Lerner MR, Steitz JA. 1981. Striking similarities are exhibited by two small Epstein-
Barr virus-encoded ribonucleic acids and the adenovirus-associated ribonucleic acids VAI and VAII. Mol
Cell Biol 1:785-96.
14.   Ahmed W, Philip PS, Tariq S, Khan G. 2014. Epstein-Barr virus-encoded small RNAs (EBERs) are
present in fractions related to exosomes released by EBV-transformed cells. PLoS One 9:e99163.
15.   Baglio SR, van Eijndhoven MA, Koppers-Lalic D, Berenguer J, Lougheed SM, Gibbs S, Leveille N, Rinkel
RN, Hopmans ES, Swaminathan S, Verkuijlen SA, Scheffer GL, van Kuppeveld FJ, de Gruijl TD, Bultink
IE, Jordanova ES, Hackenberg M, Piersma SR, Knol JC, Voskuyl AE, Wurdinger T, Jimenez CR,
Middeldorp JM, Pegtel DM. 2016. Sensing of latent EBV infection through exosomal transfer of
5'pppRNA. Proc Natl Acad Sci U S A 113:E587-96.
16.   Iwakiri D, Zhou L, Samanta M, Matsumoto M, Ebihara T, Seya T, Imai S, Fujieda M, Kawa K, Takada K.
2009. Epstein-Barr virus (EBV)-encoded small RNA is released from EBV-infected cells and activates
signaling from Toll-like receptor 3. J Exp Med 206:2091-9.
17.   Lerner MR, Andrews NC, Miller G, Steitz JA. 1981. Two small RNAs encoded by Epstein-Barr virus and
complexed with protein are precipitated by antibodies from patients with systemic lupus erythematosus.
Proc Natl Acad Sci U S A 78:805-9.
18.   Iwakiri D. 2016. Multifunctional non-coding Epstein-Barr virus encoded RNAs (EBERs) contribute to viral
pathogenesis. Virus Res 212:30-8.
19.   Li Z, Duan Y, Cheng S, Chen Y, Hu Y, Zhang L, He J, Liao Q, Yang L, Sun LQ. 2015. EBV-encoded RNA
via TLR3 induces inflammation in nasopharyngeal carcinoma. Oncotarget 6:24291-303.
20.   Samanta M, Iwakiri D, Kanda T, Imaizumi T, Takada K. 2006. EB virus-encoded RNAs are recognized
by RIG-I and activate signaling to induce type I IFN. EMBO J 25:4207-14.
21.   Samanta M, Iwakiri D, Takada K. 2008. Epstein-Barr virus-encoded small RNA induces IL-10 through
RIG-I-mediated IRF-3 signaling. Oncogene 27:4150-60.

22. Diebel KW, Claypool DJ, van Dyk LF. 2014. A conserved RNA polymerase III promoter required for
gammaherpesvirus TMER transcription and microRNA processing. Gene 544:8-18.
23. Bowden RJ, Simas JP, Davis AJ, Efstathiou S. 1997. Murine gammaherpesvirus 68 encodes tRNA-like
sequences which are expressed during latency. J Gen Virol 78 (Pt 7):1675-87.
24. Feldman ER, Kara M, Coleman CB, Grau KR, Oko LM, Krueger BJ, Renne R, van Dyk LF, Tibbetts SA.
2014. Virus-encoded microRNAs facilitate gammaherpesvirus latency and pathogenesis in vivo. MBio
5:e00981-14.
25. Steer B, Strehle M, Sattler C, Bund D, Flach B, Stoeger T, Haas JG, Adler H. 2016. The small noncoding
RNAs (sncRNAs) of murine gammaherpesvirus 68 (MHV-68) are involved in regulating the latent-to-lytic
switch in vivo. Sci Rep 6:32128.
26. Diebel KW, Oko LM, Medina EM, Niemeyer BF, Warren CJ, Claypool DJ, Tibbetts SA, Cool CD, Clambey
ET, van Dyk LF. 2015. Gammaherpesvirus small noncoding RNAs are bifunctional elements that regulate
infection and contribute to virulence in vivo. MBio 6:e01670-14.
27. Bullard WL, Kara M, Gay LA, Sethuraman S, Wang Y, Nirmalan S, Esemenli A, Feswick A, Hoffman BA,
Renne R, Tibbetts SA. 2019. Identification of murine gammaherpesvirus 68 miRNA-mRNA hybrids
reveals miRNA target conservation among gammaherpesviruses including host translation and protein
modification machinery. PLoS Pathog 15:e1007843.
28. Chiu YH, Macmillan JB, Chen ZJ. 2009. RNA polymerase III detects cytosolic DNA and induces type I
interferons through the RIG-I pathway. Cell 138:576-91.
29. Yoneyama M, Fujita T. 2007. Function of RIG-I-like receptors in antiviral innate immunity. J Biol Chem
282:15315-8.
30. White RJ. 2005. RNA polymerases I and III, growth control and cancer. Nat Rev Mol Cell Biol 6:69-78.
31. Amort M, Nachbauer B, Tuzlak S, Kieser A, Schepers A, Villunger A, Polacek N. 2015. Expression of the
vault RNA protects cells from undergoing apoptosis. Nat Commun 6:7030.
32. Nandy C, Mrazek J, Stoiber H, Grasser FA, Huttenhofer A, Polacek N. 2009. Epstein-barr virus-induced
expression of a novel human vault RNA. J Mol Biol 388:776-84.
33. Felton-Edkins ZA, Kondrashov A, Karali D, Fairley JA, Dawson CW, Arrand JR, Young LS, White RJ.
2006. Epstein-Barr virus induces cellular transcription factors to allow active expression of EBER genes
by RNA polymerase III. J Biol Chem 281:33871-80.
34. Karijolich J, Abernathy E, Glaunsinger BA. 2015. Infection-Induced Retrotransposon-Derived Noncoding
RNAs Enhance Herpesviral Gene Expression via the NF-kappaB Pathway. PLoS Pathog 11:e1005260.
35. Karijolich J, Zhao Y, Alla R, Glaunsinger B. 2017. Genome-wide mapping of infection-induced SINE
RNAs reveals a role in selective mRNA export. Nucleic Acids Res 45:6194-6208.
36. Oko LM, Kimball AK, Kaspar RE, Knox AN, Coleman CB, Rochford R, Chang T, Alderete B, van Dyk LF,
Clambey ET. 2019. Multidimensional analysis of Gammaherpesvirus RNA expression reveals
unexpected heterogeneity of gene expression. PLoS Pathog 15:e1007849.
37. White RJ. 2004. RNA polymerase III transcription and cancer. Oncogene 23:3208-16.
38. White RJ. 2011. Transcription by RNA polymerase III: more complex than we thought. Nat Rev Genet
12:459-63.
39. Schramm L, Pendergrast PS, Sun Y, Hernandez N. 2000. Different human TFIIIB activities direct RNA
polymerase III transcription from TATA-containing and TATA-less promoters. Genes Dev 14:2650-63.
40. Summers WC, Klein G. 1976. Inhibition of Epstein-Barr virus DNA synthesis and late gene expression
by phosphonoacetic acid. J Virol 18:151-5.
41. Li D, Fu W, Swaminathan S. 2018. Continuous DNA replication is required for late gene transcription and
maintenance of replication compartments in gammaherpesviruses. PLoS Pathog 14:e1007070.
42. Gunnery S, Mathews MB. 1995. Functional mRNA can be generated by RNA polymerase III. Mol Cell
Biol 15:3597-607.
43. Gunnery S, Maivali U, Mathews MB. 1997. Translation of an uncapped mRNA involves scanning. J Biol
Chem 272:21642-6.
44. Rollins J, Veras I, Cabarcas S, Willis I, Schramm L. 2007. Human Maf1 negatively regulates RNA
polymerase III transcription via the TFIIB family members Brf1 and Brf2. Int J Biol Sci 3:292-302.

45. Rumi M, Ishihara S, Aziz M, Kazumori H, Ishimura N, Yuki T, Kadota C, Kadowaki Y, Kinoshita Y. 2006. RNA polymerase II mediated transcription from the polymerase III promoters in short hairpin RNA expression vector. Biochem Biophys Res Commun 339:540-7.

46. Orioli A, Pascali C, Quartararo J, Diebel KW, Praz V, Romascano D, Percudani R, van Dyk LF, Hernandez N, Teichmann M, Dieci G. 2011. Widespread occurrence of non-canonical transcription termination by human RNA polymerase III. Nucleic Acids Res 39:5499-512.

47. Muller M, Hutin S, Marigold O, Li KH, Burlingame A, Glaunsinger BA. 2015. A ribonucleoprotein complex protects the interleukin-6 mRNA from degradation by distinct herpesviral endonucleases. PLoS Pathog 11:e1004899.

48. Glaunsinger BA. 2015. Modulation of the Translational Landscape During Herpesvirus Infection. Annu Rev Virol 2:311-33.

49. Owen TJ, O'Neil JD, Dawson CW, Hu C, Chen X, Yao Y, Wood VH, Mitchell LE, White RJ, Young LS, Arrand JR. 2010. Epstein-Barr virus-encoded EBNA1 enhances RNA polymerase III-dependent EBER expression through induction of EBER-associated cellular transcription factors. Mol Cancer 9:241.

50. Adler H, Messerle M, Wagner M, Koszinowski UH. 2000. Cloning and mutagenesis of the murine gammaherpesvirus 68 genome as an infectious bacterial artificial chromosome. J Virol 74:6964-74.

51. Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res 29:e45.

52. Dumay-Odelot H, Durrieu-Gaillard S, El Ayoubi L, Parrot C, Teichmann M. 2014. Contributions of in vitro transcription to the understanding of human RNA polymerase III transcription. Transcription 5:e27526.

53. van Zon A, Mossink MH, Schoester M, Scheffer GL, Scheper RJ, Sonneveld P, Wiemer EA. 2001. Multiple human vault RNAs. Expression and association with the vault complex. J Biol Chem 276:37715-21.

54. Vachon VK, Conn GL. 2016. Adenovirus VA RNA: An essential pro-viral non-coding RNA. Virus Res 212:39-52.

55. Howe JG, Shu MD. 1989. Epstein-Barr virus small RNA (EBER) genes: unique transcription units that combine RNA polymerase II and III promoter elements. Cell 57:825-34.

56. Howe JG, Shu MD. 1993. Upstream basal promoter element important for exclusive RNA polymerase III transcription of the EBER 2 gene. Mol Cell Biol 13:2655-65.

57. Niller HH, Salamon D, Ilg K, Koroknai A, Banati F, Bauml G, Rucker O, Schwarzmann F, Wolf H, Minarovits J. 2003. The in vivo binding site for oncoprotein c-Myc in the promoter for Epstein-Barr virus (EBV) encoding RNA (EBER) 1 suggests a specific role for EBV in lymphomagenesis. Med Sci Monit 9:HY1-9.

FIGURE LEGENDS

Figure 1. RNA Polymerase III promoters contain distinct motifs. Pol III transcribes RNA from gene-internal

(type 1 and type 2) or gene-external (type 3) promoters. Each promoter type contains distinct motifs, which are

in turn bound by specific transcription factors (TF) that recruit pol III to the promoter. Viral promoters can have

canonical type 2 promoters, such as adenovirus; however, the γHV68 TMERs contain a triplicated, overlapping

A box motif, and the EBV EBER promoters include upstream elements. Gray boxes indicate the gene, while the

"TTTT" marks the end of the gene where pol III transcription is terminated. Diagrams are not to scale. *A* = A

box, *B* = B box, *C* = C box, *IE* = intermediate element, *"TTTT"* = pol III termination signal, *TF* = transcription

factor, *TATA* = TATA box, *PSE* = proximal sequence element, *DSE* = distal sequence elements, *ETAB* = EBER

TATA-like box, *ATF* = activating transcription factor binding sequence, *Sp1* = Sp1-like binding sequence. B)

Optimization of the NanoLuc reporter vector. Mutations were introduced into the pNL1.1 vector to remove the

pol III termination signal (TTTT) from the Nanoluc coding region, resulting in the pNLP3 vector. The human U6

promoter was cloned into each of these reporters, and dual luciferase assays were performed to compare

luciferase output. Data shown is representative of two independent experiments with biological triplicates.

Additionally, RNA was isolated from cells transfected with these two constructs. Cells were mock or WT γHV68-

infected for 16 h, then cellular RNA was used as a template for primers targeting the entire NanoLuc gene (top

gel, 534 nt), or targeting just the NanoLuc sequence upstream of the termination sequence (bottom gel, 234 nt).

Data shown is from one experiment with biological triplicates. NTC = Non-template control, black triangles =

primers, * = location of termination sequence in pNL1.1.

Figure 2. γHV68 infection increases luciferase activity when expressed from two different pol III promoter

types. A) Experimental design. HEK293 cells are transfected with a control Firefly reporter (pGL3) and a pNLP3

reporter expressed from a pol III promoter of interest. 24 h post-transfection, cells are infected with wild-type

(WT) γHV68 at an MOI of 1, or mock treated. Cell lysates are collected 24 h post-transfection for the dual

luciferase assay. Promoter activity was measured for the (B) U6 and (C) TMER1 promoters using a dual

luciferase assay. Cells were harvested every 4 hours post-infection up to 24 h. Each time-point was repeated for

biological triplicates. Raw values for the Firefly and NanoLuc activity for each promoter are shown as

luminescence units (LUs). Promoter activity was analyzed as average relative luminescence units (RLUs =

NanoLuc LUs / Firefly LUs), and as the fold change in NanoLuc luminescence units compared to the 4 h

uninfected samples (NanoLuc fold change = Infected Sample NanoLuc LU / Mock Sample NanoLuc LU). Error

bars = SEM.

Figure 3. γHV68 infection induces activity of multiple pol III promoter types. Promoter activity was

measured for the U6 (n = 8), tRNA (n = 3), EBER1 (n = 3), EBER2 (n = 4), TMER1 (n = 18), TMER4 (n = 4), and

TMER5 (n = 6) promoters during infection. Luciferase assays were performed as previously described, with cell

lysates collected 24 h post-infection. Each independent experiment (n) contained biological triplicates or

duplicates. All promoter activity changes are analyzed as the fold change in NanoLuc activity normalized to

uninfected samples (Infected NanoLuc LU / Mock NanoLuc LU). Error bars = SEM. Significant differences

analyzed by t-test and indicated as asterisks. P-values are indicated as follows: * = P ≤ 0.05, ** = P ≤ 0.01, *** =

P≤ 0.001, **** = P ≤ 0.0001.

Figure 4. γHV68 infection differentially induces activity of promoters containing external sequences.

Promoter activity was measured for the "minimal" or "full" promoters for EBER1, EBER2, TMER1, TMER4, and

TMER5. Minimal promoters contained only the sequence constituting the internal promoter elements. Full

promoters included sequence outside of the internal promoter elements; the full EBER promoters also contained

upstream promoter elements, while full TMER promoters contained the entire tRNA-like section of the gene (see

supplementary figure 1). Luciferase assays were performed as previously described, with cell lysates collected

24 h post-infection. All promoter activity changes are analyzed as the fold change in NanoLuc activity normalized

to uninfected samples. Each experiment (n) included biological triplicates. Error bars = SEM. Significant

differences analyzed by t-test and indicated as asterisks. P-values are indicated as follows: * = P ≤ 0.05, ** = P

≤ 0.01, *** = P≤ 0.001, **** = P ≤ 0.0001.

Figure 5. Effect of γHV68 infection on NanoLuc transcript levels. Promoter activity was measured for A) the

U6, TMER1, and EBER full promoters, or B) the TMER1, TMER4, and TMER5 minimal vs. full promoters. Cells

were treated as previously described and RNA was purified from cells at 16 h post-infection (U6 n = 2, TMER1,

n = 5) or 24hpi (EBER1 n = 1, EBER2 n = 1, TMER4 n = 2, and TMER5, n = 1). RNA was converted to cDNA,

then LightCycler real-time PCR using Syber Green was performed with primers targeting the NanoLuc gene and

a host control gene (18S). The relative difference of NanoLuc was calculated using the Pfaffl method (2001,

Nucleic Acids Research), where the ratio = $(E_{target})^{\Delta CP\ target\ (control-sample)} / (E_{ref})^{\Delta CP\ ref\ (control-sample)}$. Each experiment (n)

included biological triplicates or duplicates. Error bars = SEM. Significant differences analyzed by t-test and

indicated as asterisks. P-values are indicated as follows: * = P ≤ 0.05, ** = P ≤ 0.01, *** = P≤ 0.001, **** = P ≤

0.0001.

Figure 6. γHV68 infection coupled with PAA treatment further increases luciferase activity, but not

transcript level, from pol III promoters. A) Promoter activity was measured via RT-qPCR of the NanoLuc

transcript expressed from the TMER1 promoter during infection and concurrent phosphonoacetic acid (PAA)

treatment. RT-qPCR was performed as previously described. Data is shown as NanoLuc relative difference. N

= 1 with biological triplpicates.

B) Promoter activity was measured via luciferase activity for the following promoters during infection, including
with concurrent PAA treatment: TMER1 (n = 5), TMER4 (n = 2), TMER5 (n = 2), EBER1 (n = 2), EBER2 (n = 3),
U6 (n = 2), and tRNA (n = 1). Each experiment contained biological duplicates or triplicates. Luciferase assays
were performed as previously described, with cell lysates collected 24 h post-infection. NanoLuc fold change for
infected samples is from previous figures for comparison to PAA-treated samples. All promoter activity changes
are analyzed as the fold change in NanoLuc activity normalized to uninfected samples. Error bars = SEM.
Significant differences analyzed by one-way ANOVA and indicated as asterisks. P-values are indicated as
follows: * = P ≤ 0.05, ** = P ≤ 0.01, *** = P≤ 0.001, **** = P ≤ 0.0001.

Figure 7. Endogenous expression of murine U6 snRNA and 4.5S rRNA during γHV68 infection. Murine
fibroblast cells (3T12) were infected with mock, WT γHV68, or an EBER knock-in (EBER-KI) γHV68 at an MOI=1,
harvested at 16 hpi and subjected to the PrimeFlow RNA Assay and flow cytometric analysis. **(A,B)** Analysis of
U6 snRNA expression comparing background fluorescence ("No probe", left) versus U6 probe hybridization in
Mock (middle), or WT gHV68 or EBER-KI.gHV68 infected cells (right). **(C,D)** Analysis of 4.5S rRNA expression
comparing background fluorescence ("No probe", left) versus 4.5S rRNA probe hybridization in Mock (middle),
or WT gHV68 or EBER-KI.gHV68 infected cells (right). **(E,F)** The geometric mean fluorescence intensity (gMFI)
was calculated for **(E)** U6 snRNA or **(F)** 4.5S rRNA in mock infected, or virus infected samples, comparing cells
that differed in expression of either the TMERs or the EBERs in WT or EBER-KI infections respectively. Data
depict flow cytometric analysis analyzing singlets, defined by sequential discrimination using SSC and FSC
parameters. Data are from 3 biological replicates, with statistical significance assessed using an unpaired t-test.
Statistical significance as follows: **, p<0.01 and ***, p<0.001.

Table 1. Promoters analyzed in this study. Pol III promoters were selected from several organisms and cloned into a luciferase reporter to analyze promoter activity during infection

			Promoter Insert	
Origin	Promoter	Promoter Type	Length (nt)	Reference
Human	5S rRNA	Type 1	42	(52)
Human	tRNA-Tyr	Type 2	53	Gene ID: 100189507
Human	Vault RNA1-1	Type 2	65	(32, 53); GenBank: AF045143.1
Human	U6 snRNA	Type 3	227	GenBank: M14486.1
Adenovirus	VA1	Type 2	58	(54)
γHV68	TMER1 full	Type 2 (3A)	91	(22)
γHV68	TMER1 minimal	Type 2	55	" "
γHV68	TMER4 full	Type 2 (3A)	93	" "
γHV68	TMER4 minimal	Type 2	55	" "
γHV68	TMER5 full	Type 2 (3A)	94	" "
γHV68	TMER5 minimal	Type 2	53	" "
EBV	EBER1	Type 2 (+ upstream)	157	(55-57)
EBV	EBER1 minimal	Type 2 (no upstream)	70	" "
EBV	EBER2	Type 2 (+ upstream)	156	" "
EBV	EBER2 minimal	Type 2 (no upstream)	68	" "

Figure 1
Knox et al

Figure 2
Knox et al

Figure 3
Knox et al

Figure 4
Knox et al

A)

B)

Figure 5
Knox et al

Figure 6
Knox et al

Figure 7
Knox et al