Lytic infection with murine gammaherpesvirus 68 activates host and viral RNA polymerase III promoters and enhances non-coding RNA expression

Ashley N. Knox,[a] Alice Mueller,[b] Eva M. Medina,[a] Eric T. Clambey,[c] Linda F. van Dyk[a]#

[a] Department of Immunology and Microbiology, University of Colorado Anschutz Medical Campus, Aurora, Colorado, USA

[b] Department of Molecular, Cellular, and Developmental Biology, University of Colorado, Boulder, Colorado, USA

[c] Department of Anesthesiology, University of Colorado Anschutz Medical Campus, Aurora, Colorado, USA

Running Title: γHV68 infection activates pol III promoters

#Address correspondence to Linda van Dyk, linda.vandyk@cuanschutz.edu

Abstract: 239 words

Importance: 136 words

Text: 8594 words

ABSTRACT

RNA polymerase III (pol III) transcribes multiple non-coding (nc) RNAs that are essential for cellular function. Pol

III-dependent transcription is also engaged during certain viral infections, including the gammaherpesviruses

(γHVs), where pol III-dependent viral ncRNAs promote pathogenesis. Additionally, several host ncRNAs are

upregulated during γHV infection and play integral roles in pathogenesis by facilitating viral establishment and

gene expression. Here, we sought to investigate how pol III promoters and transcripts are regulated during

gammaherpesvirus infection using the murine gammaherpesvirus 68 (γHV68) system. To compare the

transcription of host and viral pol III-dependent ncRNAs, we analyzed a series of pol III promoters for host and

viral ncRNAs using a luciferase reporter optimized to measure pol III activity. We measured promoter activity

from the reporter gene at the translation level via luciferase activity and at the transcription level via RT-qPCR.

We further measured endogenous ncRNA expression at single cell-resolution by flow cytometry. These studies

demonstrated that lytic infection with γHV68 increased the transcription from multiple host and viral pol III

promoters, and further identified the ability of accessory sequences to influence both baseline and inducible

promoter activity after infection. RNA flow cytometry revealed the induction of endogenous pol III-derived

ncRNAs that tightly correlated with viral gene expression. These studies highlight how lytic gammaherpesvirus

infection alters the transcriptional landscape of host cells to increase pol III-derived RNAs, a process that may

further modify cellular function and enhance viral gene expression and pathogenesis.

IMPORTANCE

Gammaherpesviruses are a prime example of how viruses can alter the host transcriptional landscape to

establish infection. Despite major insights into how these viruses modify RNA polymerase II-dependent

generation of messenger RNAs, how these viruses influence the activity of host RNA polymerase III remains

much less clear. Small non-coding RNAs produced by RNA polymerase III are increasingly recognized to play

critical regulatory roles in cell biology and virus infection. Studies of RNA polymerase III dependent transcription

are complicated by multiple promoter types and diverse RNAs with variable stability and processing

requirements. Here, we characterized a reporter system to directly study RNA polymerase III-dependent

responses during gammaherpesvirus infection and utilized single-cell flow cytometry-based methods to reveal

that gammaherpesvirus lytic replication broadly induces pol III activity to enhance host and viral non-coding RNA

expression within the infected cell.

INTRODUCTION

Gammaherpesviruses (γHVs) are large, dsDNA viruses that establish a life-long infection in their hosts,

with long-term latency in lymphocytes (1, 2). The γHVs include the human pathogens, Epstein-Barr virus (EBV)

and Kaposi's sarcoma-associated herpesvirus (KSHV or HHV-8), and murine gammaherpesvirus 68 (γHV68 or

MHV-68; ICTV nomenclature *Murid herpesvirus 4*, MuHV-4) (3). These viruses establish a primary lytic infection

in their host that is followed by a prolonged quiescent infection termed latency. Latency is maintained in healthy

individuals by a homeostatic relationship between the virus and the host immune response; if this balance is

disrupted (e.g. by immunosuppression), γHVs can reactivate from latency and actively replicate. Disruption

between the balance of γHV infection and host immune control is associated with γHV multiple pathologies,

including a range of malignancies (4).

The γHVs contain several types of non-coding (nc) RNAs, including nuclear ncRNAs and functional

miRNAs; these diverse RNAs include ncRNAs transcribed by RNA polymerase II (e.g. the KSHV PAN RNA and

the KSHV and EBV miRNAs) or by RNA pol III (e.g. the EBV-encoded small RNAs (EBERs) and the γHV68

tRNA-miRNA-encoded RNAs (TMERs)) (5-13). Viral ncRNAs are considered to have important host-modulatory

functions, interacting with host proteins and regulating host and viral gene expression. For example, the EBV

EBERs are expressed during latency, and were discovered through their interaction with the host lupus-

associated antigen (La) protein, which putatively mediates EBER interaction with TLR3 (14-17).The EBERs have

further been shown to interact with several host proteins including ribosomal protein L22, protein kinase R (PKR),

and retinoic-acid inducible gene I (RIG-I) (18). These interactions can trigger sustained host innate immune

responses that are implicated in the development of EBV-associated malignancies (16, 19-21). γHV68, a highly

tractable small animal model of γHV infection, also encodes several pol III-transcribed ncRNAs known as the

tRNA-miRNA-encoded RNAs (TMERs) (22, 23). The TMERs are dispensable for lytic replication and

establishment of latency; however, these transcripts are required for pathogenesis during acute infection of an

immunocompromised host (7, 24, 25). The TMERs contain bi-functional elements with a tRNA-like structure at

the 5' end and hairpins that are processed into biologically-active miRNAs (7), capable of targeting a number of

RNAs for post-transcriptional regulation (26). Our lab has previously shown that the tRNA-like structure is

sufficient to rescue pathogenesis of a TMER deficient viral recombinant, suggesting that like the EBERs, the

TMERs may contribute to pathogenesis through their interactions with host proteins (25). Though TMER-host

protein interactions have yet to be fully explored, it is notable that several characteristics of the EBERs, such as

a 5'-triphosphate and 3'-polyU, are imparted by RNA polymerase III (pol III) transcription (27).These motifs can

be recognized by host RNA-binding proteins, such as RIG-I or La, to trigger an innate immune response (16, 20,

27, 28).

Pol III is often considered to perform "house-keeping" functions, as it transcribes host genes required for

cell growth and maintenance (e.g. U6 snRNA, tRNAs, and 5S rRNA) (29). Despite this, it is clear that the γHVs

can usurp pol III-dependent transcription mechanisms for their own purposes. Latent EBV infection has been
shown to upregulate components of pol III and ultimately increase the expression of host pol III transcripts –
particularly vault RNAs – that allow increased establishment of viral infection and gene expression (30-32).
Similarly, γHV68 infection drives upregulation of host pol III-dependent short-interspersed nuclear element
(SINE) RNAs, which in turn, mediate increased viral gene expression (33-35). Additionally, our lab has reported
that reactivation of a latently-infected γHV68 cell line results in in a rare subset of the population that
demonstrates increased viral transcription and translation, including increased expression of TMERs (36).
Notably, dysregulation of pol III is a common feature of many cancer cells, implicating γHV infection-driven
alteration of pol III activity as one potential contributor to γHV-associated malignancies (37). Therefore,
understanding how γHV infection alters pol III activity is integral to elucidating mechanisms of γHV pathogenesis.

The analysis of pol III activity during γHV infection has been complicated by the nature of pol III-derived
ncRNAs. These transcripts are often short and structured, creating complications in probe specificity to
quantitatively analyze promoter activity/gene expression by conventional means (e.g. northern blot, RT-qPCR).
Probe specificity is also challenging for classes of RNAs with highly conserved promoter features, such as the
γHV68 TMERs or the human tRNAs. However, promoter analysis of the TMERs could reveal rules of
transcription that apply to other conserved and potentially co-regulated ncRNAs, such as the human tRNAs.
Additionally, many pol III-derived ncRNAs may be scarce or abundant so that changes in expression can be
obscured. Therefore, highly sensitive readouts, such as the high dynamic range of luciferase assays or
quantitative assays with single-cell resolution, offer potential improvement in measuring pol III-derived ncRNA
expression. Single cell RNA flow cytometry allows RNA detection with high specificity without the need for unique
primers and probe, and has the additional benefit of measuring RNA levels in individual cells to reveal fine
fluctuations that may be obscured in bulk analyses.

The ideal comparison of promoters allows variation only in the promoter elements coupled to a common
reporter gene and sensitive detection. Traditional analysis of RNA pol II promoter activity has benefitted
significantly from the use of luciferase reporter systems, which provide the advantages of a readout that is high
throughput, has a wide dynamic range for maximal quantitation, and is standardized across varied promoters
and cellular conditions. RNA pol III does not produce coding RNAs in normal biology; however, several studies
have reported the use of luciferase reporters to measure ncRNA derived from pol III or pol I (38, 39). Based on
these studies, we developed a panel of luciferase reporters driven by pol III promoters to determine the efficacy
of a reporter gene approach in analyzing ncRNA promoter activity during viral infection. As with analysis of RNA
pol II reporters, caveats to the enzymatic readout of this system are that it is several steps downstream of RNA
transcription, the efficacy of RNA translation may differ among specific RNAs, and infection may alter translation
in a number of ways yet to be described. However, use of the facile enzymatic readout plus RT-qPCR
quantitation of the reporter RNA allows us to directly compare these measures for highly sensitive quantitative

analysis. As a complementary approach, we quantified ncRNAs expression at the single-cell level in the

presence or absence of virus infection.

Due to the importance of γHV ncRNAs during infection and the unique transcriptional regulation afforded

by RNA pol III, the overall objective of this study was to compare different RNA pol III promoters and their activity

during virus infection using three different methods for sensitive and quantitative analysis. We found that γHV68

infection upregulates the activity of multiple viral and host pol III promoters, a process further associated with the

induction of pol III-dependent targets. These studies indicate that lytic γHV infection can broadly enhance RNA

pol III promoter activity to modify the ncRNA landscape of infected cells.

RESULTS

RNA polymerase III can transcribe RNA from a variety of gene-internal (type 1 and 2) and gene-

external (type 3) promoters (Fig 1A). These promoters contain distinct motifs that determine which transcription

factors bind to the promoter to recruit pol III (40). To understand how γHV lytic replication influenced RNA pol

III promoter activity while limiting the confounding factors of the individual RNA primary transcripts and

modified or processed products, we sought to make use of a luciferase assay previously used to study pol III

promoter activity (41) to study a series of viral and host pol III promoters. We selected the reporter plasmid

pNL1.1 (Promega), because NanoLuc luciferase creates a brighter signal, and the protein is smaller than other

luciferase proteins (NanoLuc 19.1 kDa and 171 nucleotides; Renilla 36.0 kDa and 312 nucleotides; Firefly 60.6

kDa and 550 nucleotides), which is consistent with pol III processivity of small ncRNAs. Our analysis of the

pNL1.1 sequence revealed a pol III termination signal within the luciferase coding gene (TTTT). Therefore, to

examine the activity of pol III promoters without the potential for early termination, we introduced silent

mutations into the NanoLuc reporter construct to remove the termination signal. This altered vector was named

"pNLP3" to reflect that it is a *NanoLuc* reporter optimized for *p*ol *III* (Fig 1B). The human U6 promoter was

cloned into both the parental pNL1.1 and the modified pNLP3 to compare the effects of removing the pol III

termination signal, with promoter activity measured 24 hours post-infection. We found that removal of the

termination signal increased the luciferase output, indicating that there was more read-through of the full

NanoLuc gene from pNLP3 (Fig 1B, left). Furthermore, RT-PCR analysis of the NanoLuc transcript transcribed

from the U6 promoter in either the pNL1.1 or pNLP3 vector revealed more full-length NanoLuc transcript from

the pNLP3 vector (Fig 1B, right). This indicates that the pNLP3 vector allows for optimal pol III transcription of

the reporter gene. We therefore used the pNLP3 vector as the backbone for analysis of all other pol III

promoters included in this study.

With an optimized pol III reporter construct, we assessed how different pol III promoters respond to

γHV68 infection over time. We first compared the activity of the human U6 (type 3) and γHV68 TMER1 (type 2)

promoters. HEK 293 cells were transfected with pNLP3 vectors containing either the U6 or TMER1 promoter,

co-transfected with an SV40 (pol II promoter)-driven Firefly luciferase vector, then infected with γHV68 (Fig

1C). Cell lysates were collected every 4 hours for 24 hours post-infection to quantify promoter activity as

defined by NanoLuc luciferase activity. This analysis revealed that γHV68 infection resulted in a time-

dependent increase in NanoLuc activity for both the U6 and TMER1 promoters (left panel, Fig 1D-1E) relative

to mock-infected samples. γHV68 infection also resulted in a time-dependent increase in expression of the

control, Firefly luciferase reporter (second panel from left, Fig 1D-1E). The results for dual luciferase assays

are typically reported as relative luminescence units (RLUs), where the reporter luminescence units (LUs) are

normalized to the luminescence units of the control luciferase (i.e., NanoLuc LUs / Firefly LUs). However, since

γHV68 infection simultaneously increased luminescence from both the NanoLuc reporter, and from the control

Firefly reporter, this normalization implied decreased relative U6 promoter activity with infection when we

actually observe an increase in NanoLuc activity (Fig 1D). Clearly, the numerous changes incurred in cells

during viral infection limits our ability to standardize pol III promoter activity relative to a pol II promoter control

(i.e. SV40 promoter); therefore, all subsequent analyses report promoter activity as a fold change in NanoLuc

luminescence comparing mock and γHV68 infected samples. This allows us to directly compare the effect of

infection on the reporter in related samples. We found that the U6 promoter drives high basal luciferase activity

under mock conditions (left panel, Fig 1D), with a further increase in raw and normalized U6-expressed

NanoLuc LUs throughout infection (Fig 1D). In contrast, the TMER1 promoter was characterized by extremely

low basal luciferase activity (left panel, Fig 1E) in mock conditions; however, this promoter was strongly

induced by infection (Fig 1E). These data suggest that the luciferase assay can be used for analysis of pol III

promoters and show that γHV68 lytic infection increases the activity of multiple pol III promoter types, with a

more robust induction of the type 2 promoter of TMER1 compared to the U6 promoter.

We further analyzed the activity of several other promoter types to assess how they are impacted

during γHV68 infection. Experiments indicated that γHV68 infection induced activity from multiple pol III

promoters, including the human U6 and tRNA-Tyr promoters, the EBER1 and EBER2 promoters, and TMER1,

4, and 5 promoters (Fig 2). Though the vaultRNA1-1 and adenovirus VA1 promoters were cloned into the

reporter, there was no detectable activity from these constructs (unpublished data). While viral infection

induced luciferase activity from all of the examined promoters, the TMER promoters consistently showed the

greatest induction during infection.

Because the normal role of pol III is in transcription of non-coding RNAs, we wanted to directly measure

the impact of a pol III specific inhibitor (CAS 577784-91-9) on luciferase activity from the TMER1 promoter

compared to pol II promoters (TK-NanoLuc and SV40-Firefly). We treated cells with 40 µM of CAS 577784-91-

9, a drug concentration previously reported in investigation of γHV68 induction of SINE RNAs (33), prior to

transfection and infection. We found that Inhibition of pol III with this drug concentration significantly reduced

the induction of luciferase from the TMER1 promoter without toxicity (unpublished data), and that this drug

concentration had no consistent effect on the pol II promoters SV40 and TK (Fig 3A). In contrast, pol III

inhibition led to significant decreases in luciferase activity from six of the seven tested pol III promoters (Fig

3B), with the exception being the U6 promoter (the only type 1 promoter) which exhibited no decrease in

activity. The apparent resistance of the U6 promoter to inhibition could either be due to the relatively high

activity of the U6 promoter, or to alternate mechanisms of transcription at the U6 promoter (e.g. pol II

recruitment to the promoter (41)) invoked at these inhibitor concentrations. These studies indicate that

inhibiting pol III activity consistently impaired induction of luciferase from multiple host and viral pol III

promoters corresponding to the type 2 pol III promoter subtype.

We also measured the luciferase RNA expressed from the TK (pol II) and TMER1 (pol III) promoters

during infection with pol III inhibition. When mimicking the transfection conditions used for the dual luciferase

assays with 200ng transfected DNA, we found no significant difference in the expression of NanoLuc from

either the TK or TMER1 promoters in any of the examined conditions (Fig 3A, B, in blue). However, given that

there is induction of NanoLuc RNA from the TMER1 promoter during infection (Fig 5), we tested whether

increasing the amount of transfected reporter DNA may allow for greater distinction in NanoLuc RNA levels

under infection and pol III-inhibition conditions. Thus, we repeated these experiments with 2ug of total

transfected DNA. These experiments showed no difference in NanoLuc expressed from the TK promoter

during infection or pol III inhibition as compared to mock. However, there was a significant decrease in

NanoLuc expressed from the TMER1 promoter during infection with pol III inhibition compared to both mock (4-

fold) and to infection (6-fold). These data suggest that the NanoLuc expression from pol III promoters is

dependent on pol III activity.

Our analyses to this point indicated that the TMER promoters expressed the highest induction in

luciferase activity during infection; therefore, we compared the sequences of TMER promoters to identify which

features of these promoters could potentially contribute to this strong induction. The initially analyzed TMER

promoters contained the TMER promoter, as well as extended sequence around the minimal promoter

elements (Fig 4A). Considering that the extra sequence included in these "full" promoters may contribute to

infection-induced activity, we created a panel of "minimal" TMER and EBER promoters that contain only the

minimal RNA pol III promoter elements, i.e. the sequence beginning from the A box to the end of the B box (Fig

4A). We initially compared the activity of these promoters under basal (no infection) conditions to calculate the

average RLUs (NanoLuc LU/Firefly LU) in the absence of virus-induced changes. This analysis showed that

U6 was the most active of all pol III promoters under uninfected conditions, followed by the "full" EBER

promoters (Fig 4B). In contrast, "minimal" EBER promoters showed a significant reduction in baseline

luciferase activity. All TMER promoters (full or minimal) appeared similar to the empty vector, that is, to have

virtually no activity under uninfected conditions. We then compared the induction of NanoLuc luciferase activity

between the EBER and TMER minimal and full promoters during infection to determine the role of extended

sequence on promoter activity. As previously described, these constructs were transfected into HEK 293 cells,

then infected with γHV68. The fold change in NanoLuc activity relative to mock-treated samples was compared

after 24 h of infection (Fig 4C). When we compared the relative inducibility of EBER "full" versus "minimal"

promoters, "minimal" promoters showed greater virus-inducibility. This enhanced inducibility of the "minimal"

EBER promoters likely reflects the reduced basal luminescence from these promoters (Fig 4B). Conversely,

TMER "minimal" promoters displayed a weaker induction during infection than their "full" counterparts,

suggesting the sequence surrounding the TMER minimal promoters drives stronger expression during

infection. These results indicate that the sequence surrounding minimal pol III promoter elements impacts both
the basal activity and inducibility of these promoters during infection.

Luciferase readouts of pol III promoter activity allowed us to uniformly analyze pol III promoter activity.
This assay does not directly measure the level of RNAs, however, instead relying on an enzymatic readout of
luciferase protein activity. To ensure that γHV68 infection was inducing pol III activity transcriptionally, we used
the same NanoLuc constructs to measure promoter activity at the RNA level by performing RT-qPCR for the
NanoLuc transcript. Following the same protocol as used for the luciferase assays, HEK 293 cells were
transfected with pGL3 and the pNLP3 vector expressed by pol III promoters of interest (as outlined in Fig 1).
Cells were then infected with γHV68 and RNA was purified from cells 16 or 24 h post-infection. Primers
targeting the NanoLuc gene were used for qPCR following reverse transcription of the RNA. Infection
increased the NanoLuc RNA expression from the U6 and TMER1 promoters, with more modest induction from
the EBER promoters (Fig 5A). These results indicate that γHV68 infection stimulates pol III-promoter activity
from multiple host and viral promoters, measured at both the transcriptional and translational level. To extend
these findings, we further measured NanoLuc RNA expression from "minimal" or "full" TMER promoters. These
studies demonstrated that γHV68 infection increased NanoLuc RNA from the "minimal" promoter relative to
mock infected samples, with further RNA induction from the "full" TMER promoter (Fig 5B). These results
strongly suggest that the NanoLuc reporter assay serves as a faithful readout for pol III-dependent
transcription, quantified at both the RNA and protein level. These findings also emphasize that sequences
outside of the minimal TMER promoters contribute to increased expression during infection.

γHV lytic replication critically depends on viral DNA replication and late gene transcription, processes
that are inhibited by phosphonoacetic acid (PAA) (42, 43). We therefore tested the impact of PAA on virus-
induced pol III induction. To do this, HEK 293 cells were transfected with the pol III-driven NanoLuc constructs
and infected as before, with one set of samples receiving PAA treatment (200µg/mL) after 1 h of viral
inoculation. PAA treatment was consistently associated with increase luciferase enzymatic activity, with PAA-
treated γHV68-infected cultures characterized by a greater apparent induction of luciferase activity compared
to γHV68-infected cultures alone. This PAA-driven enhancement of luciferase activity was observed for
multiple pol III promoters, including U6, TMER1, 4 and 5, and EBER 1 and 2 (Fig 6A). To determine if this
effect was also observed at the transcriptional level, cells were transfected and infected as before. RNA was
isolated 16 h post-infection and RT-qPCR was performed to detect the NanoLuc transcript. Notably, treatment
with PAA during γHV68 infection had no impact on the induction of NanoLuc RNA compared to untreated
infected cells, indicating that viral DNA replication and late gene synthesis was not required for pol III induction
(Fig 6B). Effectiveness of the PAA treatment was confirmed by measuring expression of a viral late gene, gB
(Fig 6C). The increase in luciferase activity following PAA treatment, with minimal impact on NanoLuc RNA,
strongly suggests that PAA treatment enhanced the translational output from the promoters tested. These data

suggest that viral late gene expression plays an additional role in translation that is not seen at the

transcriptional level, a phenomenon independent of pol III promoter activity.

Given the reported relationship between the NF-κB pathway and the expression of pol III-dependent

transcripts (30), we analyzed the effect of NF-κB activation or inhibition on the activity of the U6 and TMER1

promoters via luciferase activity. First, we measured induction of an NF-κB reporter plasmid following treatment

with either TNFα, a known inducer of the NF-κB pathway, or following γHV68 infection. Whereas TNFα

induced NF-κB reporter activity at 4 and 24 hours post-treatment, γHV68 infection had no measurable impact

on expression from the NF-κB reporter (Fig 6D). Next, we analyzed the impact of NF-κB manipulation on pol III

promoter activity. Treating cells with TNFα modestly increased U6 promoter activity, albeit to a lesser extent

than γHV68 infection, while TMER1 promoter activity was not affected by TNFα treatment (Fig 6D). Inhibition of

NF-κB with the BAY 11-7082 (BAY 11) compound increased U6-expressed luciferase activity in virus infected

conditions, yet had no significant impact on TMER1 promoter activity after infection (Fig 6E). This indicates a

potential role of NF-κB in inhibiting pol III promoter activity during infection; however, this effect is only

observed in the case of a gene-external (i.e. type 3) promoter. Ultimately, these data do not support a

significant role of the NF-κB pathway in the observed induction of pol III promoter activity after γHV68 infection

in these culture conditions.

The NanoLuc-expressing constructs allowed us to measure promoter activity using a reporter system

including a shared readout that minimizes confounding factors of RNA sequence/structure/stability, and our

analysis of pol III promoter activity suggested a general induction during infection. However, a previous report

suggested that only a subset of host pol III-transcribed genes – the SINE RNAs – are increased during lytic

γHV68 infection of NIH3T3 cells and *in vivo* infections of C57BL/6 mice (33). While our reporter studies rely on

cell systems that can be easily transfected with reporter DNA, we wanted to quantify the impact of γHV68

infection on the abundance of endogenous murine and viral pol III RNAs, yet avoid challenges in PCR

amplification, unique primer/probe designs, and bulk analysis. In order to accomplish this, we made use of the

PrimeFlow assay system, a sensitive and robust fluorescent in situ hybridization assay, combined with

multiparameter flow cytometry. This system quantifies steady-state RNA expression at the single-cell level

using direct probe hybridization to endogenous RNAs, with sensitivity provided by amplification based on probe

stacking rather than PCR and primers. For this analysis, murine fibroblast cells (3T12) were mock-treated,

infected with wild-type (WT) γHV68, or infected with an EBER-knock in (EBER-KI.γHV68) recombinant γHV68

that lacks the TMERs and instead contains insertion of the EBV EBERs into the TMER locus. EBER-KI.γHV68

was competent for viral replication (manuscript in progress). 3T12 cells were infected with an MOI of 1 and

harvested at 16 h post-infection, conditions that result in a mixed population of virally infected and uninfected

cells. Cells were then queried with fluorescent probes to detect viral (the γHV68 TMERs or the EBV EBERs)

and host ncRNAs (U6 snRNA or 4.5 rRNA, the murine equivalent of human 5S rRNA). This analysis allowed a

comparison of host ncRNA expression as a function of viral ncRNA expression, comparing cells with or without

virus ncRNA expression. Gating schemes and probe specificity are shown in Supplemental Figure 1. Notably,

probes for the TMERs and EBERs were specific for their intended targets, with no detectable TMER or EBER

signal in mock-infected samples, TMER+ cells only present in WT γHV68 infected samples, and EBER+ cells

only present in EBER-KI.γHV68 infected samples (Fig 7E). We next compared the relative expression of the

U6 (Fig 7A-B) and 4.5S ncRNAs (Fig 7C-D) between cells with active viral RNA expression versus those cells

that did not express these viral ncRNAs. Analysis of the geometric mean fluorescence intensity (gMFI) for the

U6 snRNA and 4.5 rRNA probes revealed increased expression of U6 and 4.5 rRNA ncRNAs in virally infected

cells (i.e. TMER+ or EBER+) compared to uninfected cells (TMER- or EBER-) (Fig 7F-G). These data

demonstrate an increase in host pol III-transcribed ncRNAs in γHV68-infected cells and emphasize the benefit

of single-cell analysis to quantify endogenous ncRNA expression during virus infection.

The above analysis established a correlation between expression of viral and host ncRNAs at the

single-cell level, but raised the potential that detection of endogenous viral ncRNAs could bias our analysis to

cells with higher pol III activity independent of viral effects. To address this concern, we repeated the analysis

with another, pol III-independent read-out of infection, detection of the pol II-dependent γHV68 ORF18 RNA.

Cells were treated as before, with the modification that they were infected at an MOI of 5 rather than 1 so that

differences between infected and non-infected cells were exaggerated. To simplify the RNA probe panel, and

to fortify previous data, we focused on the U6 snRNA as a host measure of pol III activity. Gating schemes and

probe specificity are shown in Supplemental Figure 2. Infected cells were defined as ORF18 RNA positive, as

well as TMER (Fig 8A) or EBER (Fig 8B) positive for WT and EBER-KI γHV68 infection, respectively. This

analysis identified several populations that could be distinguished by different expression levels (negative, mid

or high expression) for either ORF18 or the viral ncRNAs (Fig 8C-D). Within each ORF18-defined population,

we calculated the gMFI for either the TMERs/EBERs (Fig 8E) or U6 snRNA (Fig 8F). ORF18 RNA[high] cells had

the highest expression of the TMERs, in WT infected samples, and the EBERs, in EBER-KI infected samples

compared to ORF18 RNA[mid] or ORF18 negative cells (Fig 8E). Notably, virally-infected ORF18 RNA[high] cells

from either WT or EBER-KI γHV68 infection were also characterized by increased U6 expression relative to

uninfected, ORF18 negative cells in the same cultures (Fig. 8F). Consistent with our previous observations, U6

RNA gMFI was highest in cells with highest expression of either the TMERs or EBERs (Fig 8G). These data

demonstrate that virally infected cultures are characterized by variation in ncRNA expression between

individual cells, and that cells with abundant virus transcription are further characterized by increased host and

viral ncRNA expression.

DISCUSSION

While γHV infection is known to alter expression of host genes, and viral ncRNAs are integral for

pathogenesis, the transcriptional regulation of these ncRNAs has remained unclear. Here, we propose that

different pol III promoter types allow for distinct means for transcriptional regulation during infection. This study

focused on the impact of γHV68 lytic replication on pol III promoter activity. To measure pol III activity, we used

three highly-sensitive methods, two of which were based on a reporter gene and one method based on

measuring authentic RNAs. Though these methods each have different advantages and disadvantages, they

all indicate that lytic infection drives a general upregulation of promoter activity across multiple host and viral

pol III-dependent transcripts, as shown by upregulation of both luciferase activity and luciferase RNA

expression, as well as increased expression of endogenous transcripts as measured by flow cytometric

analysis of pol III RNAs. Our studies further reveal distinct effects of specific promoters and promoter features.

These findings emphasize the utility of the modified NanoLuc luciferase system to analyze pol III promoter

activity, and provide clear evidence for pol III promoters with large differences in basal and inducible promoter

activity. They further emphasize the capacity of γHV lytic infection to modify pol III-dependent transcriptional

machinery in infected cells, a process that likely facilitates productive virus replication (33). At this time, it

remains unknown whether pol III machinery or transcription is altered during γHV68 latent infection or

reactivation from latency.

Our use of a luciferase reporter to measure the activity of pol III promoters allowed us to directly

compare the functional activity of multiple pol III promoters in a high-throughput assay while minimizing

potential differences that may arise due to ncRNA sequence, structure, or stability. While these assays used

NanoLuc RNA and protein as a standard platform for analysis, an important caveat of this analysis is that many

pol III promoters are gene-internal. Based on this, the resulting transcript is comprised of a hybrid of at least

the minimal pol III promoter elements (i.e. the A and B box) directly fused upstream of the NanoLuc RNA,

creating a hybrid RNA that is not 100% identical between constructs. Though pol III promoters conventionally

drive expression of non-coding RNAs, there is clear precedent that pol III can transcribe translation-competent

RNA (44, 45), and luciferase reporters have been used for high throughput and unbiased analysis of pol III

promoter activity (38, 46). Inspired by these studies, we cloned several host and viral ncRNA promoters into a

NanoLuc luciferase reporter to measure their activity during lytic γHV68 infection. We chose the NanoLuc

luciferase as our reporter due to the small size and high activity, which is consistent with pol III transcriptional

capacity and measurement of genes that may be expressed at a low level. To further enhance the robustness

of this reporter, we identified and removed a pol III termination sequence within the NanoLuc gene which

approximately doubled luciferase reporter activity. Though pol III transcription should theoretically be

terminated in the original NanoLuc reporter, pol III read-through of termination signals has been reported (47).

In total, use of the modified NanoLuc reporter construct afforded a sensitive and robust readout for assessing
pol III promoter activity.

Through use of this pol III reporter assay, we found that γHV68 infection increased promoter activity
across a range of host and viral pol III promoters as measured by both luciferase activity and reporter RNA
abundance. This suggests that γHV68 infection generally upregulates pol III activity. Interestingly, the
consequence/magnitude of induction elicited by infection varied between promoters. For example, the U6
promoter conveyed high basal activity, with infection resulting in a modest induction of U6 promoter activity.
Conversely, the TMER promoters exhibited extremely low basal activity in mock-infected conditions, with
dramatic induction after γHV68 infection. The inducibility of the TMERs was further enhanced by accessory
sequences outside of the minimal A and B box elements. One explanation for this enhanced induction is that
these extended sequences may contain additional transcriptional elements that are integral to the promoter
itself. While formally possible, it is notable that accessory sequences across the TMERs are not conserved (6,
22, 23). An alternate explanation for the enhanced activity of the full TMER promoters is that inclusion of the
extended sequence includes the full tRNA-like structure of the TMER genes (Fig 4A). It is interesting to
speculate that this tRNA-like structure could either lend greater stability to transcripts, or protect the transcripts
from degradation by host exonucleases or the viral endonuclease, muSOX (48). Although pol III transcription is
frequently associated with the transcription of housekeeping ncRNAs, there is clear precedent that pol III can
also participate in inducible gene expression (e.g. γHV68-induced expression of SINEs) (33, 34). Whether the
TMERs have conserved regulatory mechanisms with host inducible ncRNAs is currently unknown, however,
the TMERs share promoter similarity to the SINEs (type 2, gene internal).

Our studies demonstrated that infection increased NanoLuc expression at both the RNA and protein
activity level, indicating that virus infection increased pol III promoter activity and not some secondary
measurement. While we saw broad induction of luciferase activity across pol II and pol III promoters, induction
of activity from pol III promoters was inhibited by a pol III-inhibitor, consistent with a model that these
promoters are recruiting pol III to transcribe translation-competent RNA. Notably, induction of the reporter RNA
as measured by RT-qPCR was specific to a pol III promoter (TMER1), and no induction was seen from a pol II
promoter (TK, Fig 3A). This suggests possible caveats in the induction we measure from luciferase activity vs.
luciferase RNA. A difference in reporter protein activity vs. transcript abundance was also seen when we inhibit
viral late gene expression. Unexpectedly, inhibition of viral late gene expression with PAA resulted in increased
luciferase activity from nearly all of the promoters examined – this phenomenon was only seen at the level of
NanoLuc protein activity, not at the level of NanoLuc RNA. The ability of PAA to enhance NanoLuc protein
activity, with no commensurate change in RNA expression, suggests that viral late genes may have a potential
role in tempering translation. Together with the general induction of luciferase activity seen during infection,
these data suggest that viral induction of gene expression may be occurring at both the transcriptional and

translational level. Infection may induce expression of pol III-derived reporter transcripts (RNA) and also
increase general translation (luciferase activity), leading to a compound effect when we measure reporter
protein activity. Manipulation of host translational machinery by the herpesviruses is a common strategy that is
required for optimal virus replication and the production of virus progeny (49).

While experiments with the NanoLuc reporter constructs indicated a general induction in the activity of
the observed pol III promoters, the levels of induction varied by promoter type. Interestingly, promoters with
upstream elements (U6, EBER1, and EBER2) displayed the highest level of basal activity in mock conditions
and the lowest level of virus-mediated induction of luciferase activity, while gene-internal type 2 promoters had
the lowest basal activity and highest induction (Fig 2). This is likely not due to promoters with gene external
elements reaching the limit of detection of the luciferase assay, as demonstrated by the increased induction
seen with PAA treatment (Fig 6). Notably, the luciferase activity expressed from the U6 promoter was not
significantly impacted by pol III inhibition, while all other pol III promoters showed decreased activity (Fig 3).
Among the promoters with external elements, the U6 promoter has the highest induction of NanoLuc
expression when measured at the RNA level and the EBERs had a relatively low induction, with the gene-
internal TMERs moderately induced (Fig 5). The inhibitors used in this project also appear to have promoter-
specific affects, with NF-κB modulation altering the output from the type 1 U6 promoters but not the type 2
TMER promoter (Fig 6). Though all transcripts are transcribed by pol III, these data indicate varying induction
of the different promoters, with varying abilities for the resulting transcripts to be translated into functional
proteins. As evidenced by the U6 promoter, the external elements of type 1 pol III promoters seem to drive
unique responses compared to other pol III promoters and these unique characteristics must be kept in mind
when analyzing different promoter types.

Due to the possibility that viral infection has unique effects on transfected plasmids, we complemented
our reporter studies with a highly sensitive analysis of endogenous and viral gene expression using the
PrimeFlow assay to quantify endogenous RNAs by flow cytometry. Notably, this assay allowed us to identify
gene expression differences within individual cells that may be lost in bulk analysis. Furthermore, the use of a
branching fluorescent probe targeting the RNA of interest allows magnification of low signals without requiring
replication of the gene of interest with potentially inadequate/inefficient primers. These data were consistent
with our previous findings, indicating that viral infection lead to the induction of both viral (TMERs, EBERs) and
host (U6 snRNA, 4.5 rRNA) pol III-derived transcripts. An alternate explanation for these findings is that cells
with inherently higher pol III activity are more susceptible to viral infection, resulting in the observation of both
high viral pol II gene expression (ORF18) coupled with high expression of pol III-derived ncRNAs. As this
assay measures steady-state RNA levels that are influenced by RNA production and decay, changes in
measured RNA levels may be influenced by factors such as transcriptional induction, RNA stability, and/or
degradation.

Previous reports show that γHV infection can have diverse effects on pol III transcription, ranging from
a general induction of pol III machinery (e.g. in the context of EBV and EBNA1 (32, 50)), to the selective
induction of pol III-transcribed RNAs (e.g. induction of specific host vault RNAs in EBV latently infected cells
(30, 31), and the host SINE RNAs in γHV68-infected cells (33)). One challenge in interpreting these different
findings is that these studies have been done in different states of infection (latent versus lytic), in different cell
types, and in different states of cellular transformation. In many cases, the mechanistic insights gained from
these studies could only be gained through the use of *in vitro* studies. In keeping with this, we anticipate that
the γHV68 system will afford unique insights in how the γHVs regulate pol III-dependent ncRNA expression,
allowing the analysis of primary infection coupled with technologies to measure promoter activity and
endogenous ncRNA expression. For example, our single-cell analysis of pol III-derived transcripts – U6 snRNA
and 4.5 rRNA – supported that γHV68-infection not only increases pol III-dependent promoter activity (as
observed in experiments with reporter constructs), but also increases the endogenous expression of these
transcripts. Future strategies to improve high throughput direct comparison of promoters could include the use
of reporter based constructs such as those described in this study along with other reporter-based readouts,
such as recently reported fluorescent RNA aptamers that do not rely on translation (51). How host and viral
ncRNAs are regulated as a function of cell type and virus stage of infection remains an important unanswered
question.

In total, our studies revealed a γHV68-dependent induction in the activity of host and viral pol III
promoters. This induction was seen in the expression of a reporter gene, as well as in the endogenous
expression of pol III-dependent transcripts. Though previous reports have focused on the virus-mediated
upregulation of specific host ncRNAs, these experiments suggest a broader effect of lytic γHV68 infection on
pol III activity. This suggests that γHV68 modulation of the host transcriptional landscape goes beyond mRNA
regulation, and that pol III-dependent transcripts are likely to play a wider role in γHV68 pathogenesis than
previously appreciated.

METHODS

Viruses and tissue culture. All viruses were derived from a BAC-derived WT γHV68 (52 Koszinowski UH. 2000) For some experiments, the TMER total knock-out (TMER-TKO) virus was used; this virus was generated as previously described (25). Viruses were propagated and titered as previously described (25). EBER-KI virus contains EBERs 1 and 2 in the TMER-TKO virus backbone. Its generation and characterization are described in a manuscript in preparation.

Human endothelial kidney (HEK 293) and murine fibroblast (3T12) cells were cultured in Dulbecco's modified Eagle medium (DMEM; Life Technologies) supplemented with 5% fetal bovine serum (FBS; Atlanta Biologicals), 2 mM L-glutamine, 10 U/ml penicillin, and 10 µg/ml streptomycin sulfate (complete DMEM). Cells were cultured at 37°C with 5% CO_2.

Mutagenesis of pNL1.1 to create the pNLP3 NanoLuc luciferase reporter. The promoterless NanoLuc luciferase reporter vector pNL1.1[*Nluc*] was obtained from Promega, and primers were designed to introduce silent mutations to remove the pol III termination signal in the NanoLuc coding sequence; these primers are listed in Supplementary Table 1. Mutagenesis PCR was performed with the following cycles: (i) 95° for 30s, (ii) 12 cycles of 95°C for 30s, 55°C for 1 min, 68°C for 3 m. The resulting DNA was digested with DpnI (New England Biolabs Inc) and transformed into XL1-Blue super-competent cells (Agilent). Bacterial colonies were sequenced to confirm the correct mutations. The resulting plasmid was named "pNLP3" to indicate that it is a <u>N</u>ano<u>L</u>uc plasmid optimized for <u>p</u>ol <u>III</u>.

Generating a pol III promoter-driven NanoLuc reporter panel. All promoters were generated to include XhoI and HindIII overhang sequences on the 5' and 3' ends respectively. Several promoters were constructed using ligated oligonucleotides. Sequences for primers and oligonucleotides used in this work are shown in the table below. PCR-amplified promoters and pNLP3 were digested with XhoI and HindIII, then promoters were ligated into pNLP3 using T4 DNA ligase (New England BioLabs Inc). See Supplementary Table 1 for primers used to PCR-amplify promoters. Ligated constructs were transformed into One Shot electro- or chemically competent TOP10 *E. coli* cells (Thermo Fisher Scientific, #C404052 or #C404010), which were then plated at several dilutions on LB agar containing ampicillin. Resulting colonies were expanded in LB broth with ampicillin and plasmid was isolated using the QIAprep Spin Miniprep Kit (Qiagen). All constructs were confirmed by sequencing.

Name	Sequence	Cloning Template	Purpose
NanoLuc Mut For.	5'– C CAA ATG GGC CAG ATC GAA AAA ATC TTC AAG GTG GTG TAC CC –3'	pNL1.1 (Promega)	Remove termination signal (TTTT) from pNL1.1 to make pNLP3
NanoLuc Mut Rev.	5' – GG GTC CAC CAC CTT GAA GAT TTT TTC GAT CTG GCC CAT TTG G – 3'	pNL1.1 (Promega)	
U6-F-XhoI	5' – GTTATTCTCGAGGAGGGCCTATTTCCCATG – 3'	SHC001 (Functional Genomics Facility)	Clone full U6 promoter into pNLP3 (product length = 227 bp)
U6-R-HindIII	5' – GCCGCCGAAGCTTATATATAAAGCCAAGAAATC – 3'	SHC001 (Functional Genomics Facility)	
TMER-1-F-XhoI	5' – GTTGTTCTCGAGGCCAGAGTAGCTCAATTC – 3	γHV68 Left-End Plasmid	Clone TMER1 promoter into pNLP3 (product = 91 bp)
TMER-1-R-HindIII	5' – GTCGTTAAGCTTAGTTGGACCCACTTCCTC – 3'	γHV68 Left-End Plasmid	
TMER4full_F_XhoI	5' - GTTATTCTCGAGGTCGGGGTAGCTCAATTG - 3'	γHV68 Left-End Plasmid	Clone TMER4 promoter into pNLP3 (product = 93 bp)
TMER4full_R_HindIII	5' - GTTGTGAAGCTTGACGACCCGATCTCAAC - 3'	γHV68 Left-End Plasmid	

Name	Sequence	Template	Purpose
TMER5full_F_XhoI	5' - GTACTACTCGAGGCCAGGGTAGCTCAATTG - 3'	γHV68 Left-End Plasmid	Clone TMER5 promoter into pNLP3 (product = 94 bp)
TMER5full_R_HindIII	5' - GTTGTGAAGCTTTACCGCACCTCCAC - 3'	γHV68 Left-End Plasmid	
EBER1-XhoI-F	5' - GTTGTTCTCGAGCAACTATAGCAAACCCCG - 3'	pSP73-EBER plasmid	Clone full EBER1 promoter, including upstream elements, into pNLP3 (product length = 157 bp)
EBER1-HindIII-R	5' - GTTGTTAAGCTTGGGACTTGTACCCGGG - 3'	pSP73-EBER plasmid	
EBER2-XhoI-F	5' - GTCGTTCTCGAGAGATGCACGCTTAACC - 3'	pSP73-EBER plasmid	Clone full EBER2 promoter, including upstream elements, into pNLP3 (product length = 156 bp)
EBER2-HindIII-R	5' - GGTGTTAAGCTTGGGACTTGACCTCGG - 3'	pSP73-EBER plasmid	
EBER1-Gl-XhoI-F	5' - GTTCTTCTCGAGCGCTGCCCTAGAG - 3'	pSP73-EBER plasmid	Clone EBER1-minimal promoter into pNLP3. Used with EBER1-HindIII-R, product = 70 bp
EBER2-Gl-XhoI-F	5' - GTTGTTCTCGAGCGTTGCCCTAGTGGTTTC - 3'	pSP73-EBER plasmid	Clone EBER2-minimal promoter into pNLP3. Used with EBER2-HindIII-R, product = 68 bp
NanoLuc_Forward	5' - CAC CAT GGT CTT CAC ACT CG - 3'		To perform RT-PCR on pNL1.1 and pNLP3 for the NanoLuc gene
NanoLuc-full_Rev	5' - CTA GAG TCG CGG CCT TAC G - 3'		To perform RT-PCR on pNL1.1 and pNLP3 for the NanoLuc gene
NanoLuc-prestop_Rev	5' - CGA TCT GGC CCA TTT GGT C - 3'		To perform RT-PCR on pNL1.1 and pNLP3 for the NanoLuc gene
Syber-NanoLuc-F	5' - CACTGGTAATCGACGGGGTT - 3'		To perform qPCR for the NanoLuc gene
Syber-NanoLuc-R	5' - TTTTGTTGCCGTTCCACAGG - 3'		To perform qPCR for the NanoLuc gene
pNL1.1_MCR_seq	5' - GTG TGA ATC GAT AGT ACT AAC ATA CGC - 3'		Ideal sequencing of any insert in the MCR of pNL1.1 or pNLP3
5SrRNA-XhoI-F	5' - TCGAGAGCTAAGCAGGGTCGGGCCTGGTTAGTACTTGGATGGGAGAC - 3'		Oligonucleotides annealed then enzyme-treated to create the 5S rRNA promoter, then inserted into pNLP3
5SrRNA-HindIII-R	5' - AGCTTGTCTCCCATCCAAGTACTAACCAGGCCCGACCCTGCTTAGCT - 3'		
Ad2-VAI_Fwd	5' - GTT GTT CTC GAG GTG GTC TGG TGG ATA AAT TCG CAA GGG TAT CAT GGC GGA CGC CCG GGG TTC GAA CCC CAA GCT TGT CGT C - 3'		Oligonucleotides annealed then enzyme-treated to create the adenovirus VA1 promoter, then inserted into pNLP3
Ad2-VAI_Rev	5' - GAC GAC AAG CTT GGG GTT CGA ACC CCG GGC GTC CGC CAT GAT ACC CTT GCG AAT TTA TCC ACC AGA CCA CCT CGA GAA CAA C - 3'		
vaultRNA1-1_Fwd	5' - GTT GTT CTC GAG AGC TCA GCG GTT ACT TCG ACA GTT CTT TAA TTG AAA CAA GCA ACC TGT CTG GGT TGT TCG AGA AAG CTT GTC GTC - 3'		Oligonucleotides annealed then enzyme-treated to create the adenovirus vaultRNA1-1 promoter, then inserted into pNLP3
vaultRNA1-1_Rev	5' - GAG GAC AAG CTT TCT CGA ACA ACC CAG ACA GGT TGC TTG TTT CAA TTA AAG AAC TGT CGA AGT AAC CGC TGA GCT CTC GAG AAC AAC - 3'		
tRNA-Tyr_Fwd	5' - GTT GTT CTC GAG TAG CTC AGT GGT AGA GCA TTT AAC TGT AGA TCA AGA GGT CCC TGG ATC AAC TCA AGC TTG TCG TC - 3'		Oligonucleotides annealed then enzyme-treated to create the human tRNA-Tyrosine(Tyr) promoter, then inserted into pNLP3
tRNA-Tyr_Rev	5' - GAC GAC AAG CTT GAG TTG ATC CAG GGA CCT CTT GAT CTA CAG TTA AAT GCT CTA CCA CTG AGC TAC TCG AGA ACA AC - 3'		
TMER1-min-XhoI-F	5' - TCG AGT AGC TCA ATT GGT AGA GCA ACA GGT CAC CGA TCC TGG TGG TTC TCG TT CAA GTC C - 3'		Oligonucleotides annealed to create the TMER1-minimal promoter, then inserted into pNLP3
TMER1-min-HindIII-R	5' - AGC TTG GAC TTG AAC CGA GAA CCA CCA GGA TCG GTG ACC TGT TGC TCT ACC AAT TGA GCT A - 3'		
TMER4-min-XhoI-F	5' - TCG AGT AGC TCA ATT GGT AGA GCG GCA GGC TCA TCC CCT GCA GGT TCT CGG TTC AAT CCC - 3'		Oligonucleotides annealed to create the TMER4-minimal promoter, then inserted into pNLP3
TMER4-min-HindIII-R	5' - AGCTTG GGA TTG AAC CGA GAA CCT GCA GGG GAT GAG CCT GCC GCT CTA CCA ATT GAG CTA - 3'		
TMER5-min-XhoI-F	5' - TCG AGT AGC TCA ATT GGT AGA GCA TCA GGC TAG TAT CCT GTC GGT TCC GGT TCA AGT CC - 3'		Oligonucleotides annealed to create the TMER5-minimal promoter, then inserted into pNLP3
TMER5-min-HindIII-R	5' AGC TTG GAC TTG AAC CGG AAC CGA CAG GAT ACT AGC CTG ATG CTC TAC CAA TTG AGC TA - 3'		

Transfecting cells. For transfections, HEK 293 cells were cultured in 5% FBS DMEM without penicillin

or streptomycin for approximately 24 hours. Transfection solutions contained Opti-MEM (Thermo Fisher

Scientific), NanoLuc plasmid (pNLP3 with inserted pol III promoters), and the Firefly control plasmid (pGL3-

Control; Promega). After plasmids were added to the Opti-MEM, solutions were incubated with X-tremeGENE

HP DNA Transfection Reagent (Sigma-Aldrich) for at least 15 minutes at room temperature. Transfections

were performed in several plate formats, depending on the downstream use; transfection solutions were added

dropwise to the appropriate wells (10uL of solution for 96-well plate, 100uL of solution in 12-well plates, 200uL

of solution in 6-well plates). For all transfections, the molar ratio of NanoLuc plasmid to Firefly plasmid was

kept at 10:1; the total amount of DNA transfected per well was adjusted depending on the plate size

(approximately 10ng for 96-well, 100ng for 12-well, and 200ng for 6-well). Some experiments involved

transfecting 2ug of total plasmid per well in a 6-well format. Cells were incubated with transfection solution for

24 hours prior to downstream applications, unless otherwise stated.

To analyze how promoter activity and pol III transcription was affected by γHV68 infection, transfected

HEK 293 (for NanoLuc experiments) or 3T12 (for RNA flow) cells were infected with WT, TMER-TKO, or

EBER-KI γHV68 at a multiplicity of infection (MOI) of 1 or 5 plaque forming units per cell. Cells were cultured

for approximately 24 h prior to infection. Cell counts were determined by treating with 0.05% Trypsin-EDTA

(Life Tech, #25300-054) to remove and cells. These cells were mixed with Trypan Blue dye (Bio-Rad, #145-

0021) to obtain a live cell count using the TC20 Automated Cell Counter (Bio-Rad). Virus stocks were mixed

with 5% complete DMEM, then added to cells and incubated with virus for 1 hour. Viral inoculum was then

removed and replaced with complete 5% DMEM. For samples treated with phosphonoacetic acid (PAA, 200

µg/mL, Sigma-Aldrich #284270), inoculum was removed and replaced with 5% complete DMEM containing the

drug. Inhibition of RNA polymerase III was achieved by treating cells with 40 µM of CAS 577784-91-9 (Sigma,

#557404-M) immediately prior to transfection and again following viral inoculation.

Dual luciferase assays. Following transfection and infection, HEK 293 cell lysates were collected for

analysis of luciferase activity. All luciferase assays were performed using the Nano-Glo® Dual-Luciferase®

Reporter Assay System (Promega). For experiments performed in the 12-well format, supernatant was removed

and cells were scraped, collected into 1.5mL tubes, then pelleted. Cell pellets were resuspended in 250 µL of

Passive Lysis Buffer, per the manufacturer's protocol. 80 µL of the cell suspensions were used for assays. For

luciferase assays performed in the 96-well format, 80 µL of ONE-Glo™ EX Reagent was added directly to the

cells and supernatant, as recommended by the manufacturer. Samples were incubated at room temperature

while on a shaker for three minutes. This solution was then transferred from a transparent 96-well culture plate

to a white-wall luminescence plate prior to reading Firefly luciferase activity on the Tecan Infinite® 200 PRO

plate reader. Then, 80 µL of the NanoDLR™ Stop & Glo® Reagent was added to the solution and incubated at

room temperature on a shaker for at least 10 minutes. Samples were read on the Infinite® 200 PRO again for

the NanoLuc luminescence measurement.

RT-qPCR of NanoLuc RNA. RNA was isolated from transfected HEK 293 cells with TRIzol® Reagent

(Thermo Fisher Scientific, #15596026) and DNase-treated with TURBO™ DNase (Invitrogen, #AM2238)

following the manufacturer's protocols. RNA amplification and removal of DNA was confirmed by RT-PCR or

PCR amplification of a control host gene, 18S. RT-PCR was performed using the OneStep RT-PCR Kit (Qiagen,

#210212), with the following conditions: (i) 50°C for 30 m, (ii) 95°C for 15 m, (iii) 40 cycles of 94°C for 30 s, 52°C

for 30 s, and 72°C for 30 s, (iv) 72°C for 10 m, (v) hold at 4°C. PCR was performed using Taq DNA polymerase

(Qiagen, #201205) with the following conditions: (i) 95ºC for 5 min, (ii) 40 cycles of 94ºC for 30 s, 52ºC for 30 s,

72ºC for 30 s, (iii) 72ºC for 10 m, (iv) hold at 4ºC. RNA samples that showed no product following PCR

amplification were deemed DNA-free, and converted to cDNA using SuperScript III Reverse Transcriptase

(Invitrogen, #18080093) following the manufacturer's protocol. 100ng of the cDNA was then used for qPCR

analysis of the NanoLuc and 18S genes using the iQ™ SYBER® Green Supermix (Bio-Rad, #1708880) with the

following conditions: i) 95°C for 3 m, ii) 40 cycles of 95°C for 15 s, 60°C for 1 m, iii) 95°C for 15 s, 60°C for 1 m,

95°C for 15 s. Amplification of NanoLuc was normalized to 18S expression to calculate the relative difference of

NanoLuc expression using the Pfaffl method: $\frac{NanoLuc\ Primer\ Efficiency^{Target\ \Delta Ct}}{18S\ Primer\ Efficiency^{18S\ \Delta Ct}}$ (53).

PrimeFlow RNA assay. The PrimeFlow RNA assay kit (Thermo Fisher Scientific, # 88-18005-210) was

used to analyze the endogenous expression of non-coding RNAs in mock and γHV68-infected 3T12 cells. Cells

were infected at an MOI of 1 for 16 h, then processed with the PrimeFlow kit following the manufacturer's

protocol. Cells were treated with probes for TMERs (Type 4/AF488 or Type 1/AF647), EBERs (Type 1/AF647),

ORF18 (Type 4/AF488), and U6 snRNA or 4.5 rRNA (Type 6/AF750), with compatible probes used depending

on the experiment. Samples were collected on an LSR II Flow Cytometer (BD Biosciences), and included single

stain and "full minus one" controls. Flow cytometry data were analyzed using FlowJo software (version 10.6.1),

with compensation based on single stained beads and cells. Compensated flow cytometry data were

subsequently analyzed for singlet events (based on doublet discrimination), with positive and negative

populations based on control samples as shown in Supplementary Figures 1 and 2.

Software and statistical analysis. Statistical analysis and graphing were done in GraphPad Prism

(Version 8.0d). Statistical significance was tested by unpaired t test (comparing two conditions), one-way ANOVA

(comparing three or more conditions), or two-way ANOVA (comparing grouped data) and subjected to multiple

corrections tests using recommended settings in Prism. All flow cytometry data were analyzed in FlowJo (version

10.6.1) with flow cytometry data shown as pseudo-color dot plots.

ACKNOWLEDGEMENTS

This work was supported by the NIH grants R01AI121300 to LFvD and R21 AI134084 to LFvD and ETC,

T32 AI052066 to ANK and CO RNA Biosciences summer internship support to AM.

We thank the members of the Clambey and van Dyk lab for helpful discussions, members of the Colorado

RNA Bioscience Initiative for their insights, the Colorado ClinImmune core for flow cytometry services, and the

Molecular Biology Core Facility at Anschutz Medical Campus for sequencing services. We also thank the

Functional Genomics Facility at University of Colorado and the lab of Dr. Joan Steitz lab for their generous gifts

of the pLKO.1-puro plasmid (used to clone the U6 promoter) and the pSP73-EBER plasmid (used to clone the

EBER promoters), respectively.

REFERENCES

1. Longnecker R, Kieff E, Cohen JI. 2013. Epstein-Barr Virus, Fields Virology, 6th ed. Wolters Kluwer Health Adis (ESP).
2. Ganem D. 2006. KSHV infection and the pathogenesis of Kaposi's sarcoma. Annu Rev Pathol 1:273-96.
3. Barton E, Mandal P, Speck SH. 2011. Pathogenesis and host control of gammaherpesviruses: lessons from the mouse. Annu Rev Immunol 29:351-97.
4. Cesarman E. 2011. Gammaherpesvirus and lymphoproliferative disorders in immunocompromised patients. Cancer Lett 305:163-74.
5. Samols MA, Hu J, Skalsky RL, Renne R. 2005. Cloning and identification of a microRNA cluster within the latency-associated region of Kaposi's sarcoma-associated herpesvirus. J Virol 79:9301-5.
6. Pfeffer S, Sewer A, Lagos-Quintana M, Sheridan R, Sander C, Grasser FA, van Dyk LF, Ho CK, Shuman S, Chien M, Russo JJ, Ju J, Randall G, Lindenbach BD, Rice CM, Simon V, Ho DD, Zavolan M, Tuschl T. 2005. Identification of microRNAs of the herpesvirus family. Nat Methods 2:269-76.
7. Diebel KW, Smith AL, van Dyk LF. 2010. Mature and functional viral miRNAs transcribed from novel RNA polymerase III promoters. RNA 16:170-85.
8. Sun R, Lin SF, Gradoville L, Miller G. 1996. Polyadenylylated nuclear RNA encoded by Kaposi sarcoma-associated herpesvirus. Proc Natl Acad Sci U S A 93:11883-8.
9. Cai X, Lu S, Zhang Z, Gonzalez CM, Damania B, Cullen BR. 2005. Kaposi's sarcoma-associated herpesvirus expresses an array of viral microRNAs in latently infected cells. Proc Natl Acad Sci U S A 102:5570-5.
10. Zhu JY, Strehle M, Frohn A, Kremmer E, Hofig KP, Meister G, Adler H. 2010. Identification and analysis of expression of novel microRNAs of murine gammaherpesvirus 68. J Virol 84:10266-75.
11. Reese TA, Xia J, Johnson LS, Zhou X, Zhang W, Virgin HW. 2010. Identification of novel microRNA-like molecules generated from herpesvirus and host tRNA transcripts. J Virol 84:10344-53.
12. Kim DN, Lee SK. 2012. Biogenesis of Epstein-Barr virus microRNAs. Mol Cell Biochem 365:203-10.
13. Rosa MD, Gottlieb E, Lerner MR, Steitz JA. 1981. Striking similarities are exhibited by two small Epstein-Barr virus-encoded ribonucleic acids and the adenovirus-associated ribonucleic acids VAI and VAII. Mol Cell Biol 1:785-96.
14. Ahmed W, Philip PS, Tariq S, Khan G. 2014. Epstein-Barr virus-encoded small RNAs (EBERs) are present in fractions related to exosomes released by EBV-transformed cells. PLoS One 9:e99163.
15. Baglio SR, van Eijndhoven MA, Koppers-Lalic D, Berenguer J, Lougheed SM, Gibbs S, Leveille N, Rinkel RN, Hopmans ES, Swaminathan S, Verkuijlen SA, Scheffer GL, van Kuppeveld FJ, de Gruijl TD, Bultink IE, Jordanova ES, Hackenberg M, Piersma SR, Knol JC, Voskuyl AE, Wurdinger T, Jimenez CR, Middeldorp JM, Pegtel DM. 2016. Sensing of latent EBV infection through exosomal transfer of 5'pppRNA. Proc Natl Acad Sci U S A 113:E587-96.
16. Iwakiri D, Zhou L, Samanta M, Matsumoto M, Ebihara T, Seya T, Imai S, Fujieda M, Kawa K, Takada K. 2009. Epstein-Barr virus (EBV)-encoded small RNA is released from EBV-infected cells and activates signaling from Toll-like receptor 3. J Exp Med 206:2091-9.
17. Lerner MR, Andrews NC, Miller G, Steitz JA. 1981. Two small RNAs encoded by Epstein-Barr virus and complexed with protein are precipitated by antibodies from patients with systemic lupus erythematosus. Proc Natl Acad Sci U S A 78:805-9.
18. Iwakiri D. 2016. Multifunctional non-coding Epstein-Barr virus encoded RNAs (EBERs) contribute to viral pathogenesis. Virus Res 212:30-8.
19. Li Z, Duan Y, Cheng S, Chen Y, Hu Y, Zhang L, He J, Liao Q, Yang L, Sun LQ. 2015. EBV-encoded RNA via TLR3 induces inflammation in nasopharyngeal carcinoma. Oncotarget 6:24291-303.
20. Samanta M, Iwakiri D, Kanda T, Imaizumi T, Takada K. 2006. EB virus-encoded RNAs are recognized by RIG-I and activate signaling to induce type I IFN. EMBO J 25:4207-14.
21. Samanta M, Iwakiri D, Takada K. 2008. Epstein-Barr virus-encoded small RNA induces IL-10 through RIG-I-mediated IRF-3 signaling. Oncogene 27:4150-60.
22. Diebel KW, Claypool DJ, van Dyk LF. 2014. A conserved RNA polymerase III promoter required for gammaherpesvirus TMER transcription and microRNA processing. Gene 544:8-18.

23.   Bowden RJ, Simas JP, Davis AJ, Efstathiou S. 1997. Murine gammaherpesvirus 68 encodes tRNA-like sequences
which are expressed during latency. J Gen Virol 78 (Pt 7):1675-87.
24.   Feldman ER, Kara M, Coleman CB, Grau KR, Oko LM, Krueger BJ, Renne R, van Dyk LF, Tibbetts SA. 2014. Virus-
encoded microRNAs facilitate gammaherpesvirus latency and pathogenesis in vivo. MBio 5:e00981-14.
25.   Diebel KW, Oko LM, Medina EM, Niemeyer BF, Warren CJ, Claypool DJ, Tibbetts SA, Cool CD, Clambey ET, van Dyk
LF. 2015. Gammaherpesvirus small noncoding RNAs are bifunctional elements that regulate infection and
contribute to virulence in vivo. MBio 6:e01670-14.
26.   Bullard WL, Kara M, Gay LA, Sethuraman S, Wang Y, Nirmalan S, Esemenli A, Feswick A, Hoffman BA, Renne R,
Tibbetts SA. 2019. Identification of murine gammaherpesvirus 68 miRNA-mRNA hybrids reveals miRNA target
conservation among gammaherpesviruses including host translation and protein modification machinery. PLoS
Pathog 15:e1007843.
27.   Chiu YH, Macmillan JB, Chen ZJ. 2009. RNA polymerase III detects cytosolic DNA and induces type I interferons
through the RIG-I pathway. Cell 138:576-91.
28.   Yoneyama M, Fujita T. 2007. Function of RIG-I-like receptors in antiviral innate immunity. J Biol Chem 282:15315-
8.
29.   White RJ. 2005. RNA polymerases I and III, growth control and cancer. Nat Rev Mol Cell Biol 6:69-78.
30.   Amort M, Nachbauer B, Tuzlak S, Kieser A, Schepers A, Villunger A, Polacek N. 2015. Expression of the vault RNA
protects cells from undergoing apoptosis. Nat Commun 6:7030.
31.   Nandy C, Mrazek J, Stoiber H, Grasser FA, Huttenhofer A, Polacek N. 2009. Epstein-barr virus-induced expression
of a novel human vault RNA. J Mol Biol 388:776-84.
32.   Felton-Edkins ZA, Kondrashov A, Karali D, Fairley JA, Dawson CW, Arrand JR, Young LS, White RJ. 2006. Epstein-
Barr virus induces cellular transcription factors to allow active expression of EBER genes by RNA polymerase III. J
Biol Chem 281:33871-80.
33.   Karijolich J, Abernathy E, Glaunsinger BA. 2015. Infection-Induced Retrotransposon-Derived Noncoding RNAs
Enhance Herpesviral Gene Expression via the NF-kappaB Pathway. PLoS Pathog 11:e1005260.
34.   Karijolich J, Zhao Y, Alla R, Glaunsinger B. 2017. Genome-wide mapping of infection-induced SINE RNAs reveals a
role in selective mRNA export. Nucleic Acids Res 45:6194-6208.
35.   Schaller AM, Tucker J, Willis I, Glaunsinger BA. 2020. Conserved Herpesvirus Kinase ORF36 Activates B2
Retrotransposons during Murine Gammaherpesvirus Infection. J Virol 94.
36.   Oko LM, Kimball AK, Kaspar RE, Knox AN, Coleman CB, Rochford R, Chang T, Alderete B, van Dyk LF, Clambey ET.
2019. Multidimensional analysis of Gammaherpesvirus RNA expression reveals unexpected heterogeneity of gene
expression. PLoS Pathog 15:e1007849.
37.   White RJ. 2004. RNA polymerase III transcription and cancer. Oncogene 23:3208-16.
38.   Rollins J, Veras I, Cabarcas S, Willis I, Schramm L. 2007. Human Maf1 negatively regulates RNA polymerase III
transcription via the TFIIB family members Brf1 and Brf2. Int J Biol Sci 3:292-302.
39.   Bianco C, Mohr I. 2019. Ribosome biogenesis restricts innate immune responses to virus infection and DNA. Elife
8.
40.   White RJ. 2011. Transcription by RNA polymerase III: more complex than we thought. Nat Rev Genet 12:459-63.
41.   Schramm L, Pendergrast PS, Sun Y, Hernandez N. 2000. Different human TFIIIB activities direct RNA polymerase III
transcription from TATA-containing and TATA-less promoters. Genes Dev 14:2650-63.
42.   Summers WC, Klein G. 1976. Inhibition of Epstein-Barr virus DNA synthesis and late gene expression by
phosphonoacetic acid. J Virol 18:151-5.
43.   Li D, Fu W, Swaminathan S. 2018. Continuous DNA replication is required for late gene transcription and
maintenance of replication compartments in gammaherpesviruses. PLoS Pathog 14:e1007070.
44.   Gunnery S, Mathews MB. 1995. Functional mRNA can be generated by RNA polymerase III. Mol Cell Biol 15:3597-
607.
45.   Gunnery S, Maivali U, Mathews MB. 1997. Translation of an uncapped mRNA involves scanning. J Biol Chem
272:21642-6.

46. Rumi M, Ishihara S, Aziz M, Kazumori H, Ishimura N, Yuki T, Kadota C, Kadowaki Y, Kinoshita Y. 2006. RNA
polymerase II mediated transcription from the polymerase III promoters in short hairpin RNA expression vector.
Biochem Biophys Res Commun 339:540-7.
47. Orioli A, Pascali C, Quartararo J, Diebel KW, Praz V, Romascano D, Percudani R, van Dyk LF, Hernandez N,
Teichmann M, Dieci G. 2011. Widespread occurrence of non-canonical transcription termination by human RNA
polymerase III. Nucleic Acids Res 39:5499-512.
48. Muller M, Hutin S, Marigold O, Li KH, Burlingame A, Glaunsinger BA. 2015. A ribonucleoprotein complex protects
the interleukin-6 mRNA from degradation by distinct herpesviral endonucleases. PLoS Pathog 11:e1004899.
49. Glaunsinger BA. 2015. Modulation of the Translational Landscape During Herpesvirus Infection. Annu Rev Virol
2:311-33.
50. Owen TJ, O'Neil JD, Dawson CW, Hu C, Chen X, Yao Y, Wood VH, Mitchell LE, White RJ, Young LS, Arrand JR. 2010.
Epstein-Barr virus-encoded EBNA1 enhances RNA polymerase III-dependent EBER expression through induction
of EBER-associated cellular transcription factors. Mol Cancer 9:241.
51. Song W, Filonov GS, Kim H, Hirsch M, Li X, Moon JD, Jaffrey SR. 2017. Imaging RNA polymerase III transcription
using a photostable RNA-fluorophore complex. Nat Chem Biol 13:1187-1194.
52. Adler H, Messerle M, Wagner M, Koszinowski UH. 2000. Cloning and mutagenesis of the murine
gammaherpesvirus 68 genome as an infectious bacterial artificial chromosome. J Virol 74:6964-74.
53. Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res
29:e45.
54. Dumay-Odelot H, Durrieu-Gaillard S, El Ayoubi L, Parrot C, Teichmann M. 2014. Contributions of in vitro
transcription to the understanding of human RNA polymerase III transcription. Transcription 5:e27526.
55. van Zon A, Mossink MH, Schoester M, Scheffer GL, Scheper RJ, Sonneveld P, Wiemer EA. 2001. Multiple human
vault RNAs. Expression and association with the vault complex. J Biol Chem 276:37715-21.
56. Vachon VK, Conn GL. 2016. Adenovirus VA RNA: An essential pro-viral non-coding RNA. Virus Res 212:39-52.
57. Howe JG, Shu MD. 1989. Epstein-Barr virus small RNA (EBER) genes: unique transcription units that combine RNA
polymerase II and III promoter elements. Cell 57:825-34.
58. Howe JG, Shu MD. 1993. Upstream basal promoter element important for exclusive RNA polymerase III
transcription of the EBER 2 gene. Mol Cell Biol 13:2655-65.
59. Niller HH, Salamon D, Ilg K, Koroknai A, Banati F, Bauml G, Rucker O, Schwarzmann F, Wolf H, Minarovits J. 2003.
The in vivo binding site for oncoprotein c-Myc in the promoter for Epstein-Barr virus (EBV) encoding RNA (EBER)
1 suggests a specific role for EBV in lymphomagenesis. Med Sci Monit 9:HY1-9.

FIGURE LEGENDS

Figure 1. RNA Polymerase III promoters contain distinct motifs and respond to viral infection. **(A)** Pol III

transcribes RNA from gene-internal (type 1 and type 2) or gene-external (type 3) promoters. Each promoter type

contains distinct motifs, which are in turn bound by specific transcription factors (TF) that recruit pol III to the

promoter. Viral promoters can have canonical type 2 promoters, such as adenovirus; however, the γHV68

TMERs contain a triplicated, overlapping A box motif, and the EBV EBER promoters include upstream elements.

Gray boxes indicate the gene, while the "TTTT" marks the end of the gene where pol III transcription is

terminated. Diagrams are not to scale. A = A box, B = B box, C = C box, IE = intermediate element, "TTTT" =

pol III termination signal, TF = transcription factor, TATA = TATA box, PSE = proximal sequence element, DSE

= distal sequence elements, ETAB = EBER TATA-like box, ATF = activating transcription factor binding

sequence, Sp1 = Sp1-like binding sequence. **(B)** Optimization of the NanoLuc reporter vector. Mutations were

introduced into the pNL1.1 vector to remove the pol III termination signal (TTTT) from the NanoLuc coding region,

resulting in the pNLP3 vector. The human U6 promoter was cloned into each of these reporters, and dual

luciferase assays were performed to compare luciferase output (left panel). Data shown is representative of two

independent experiments with biological triplicates. Additionally, RNA was isolated from cells transfected with

these two constructs. Cells were mock or WT γHV68-infected for 16 h, then cellular RNA was used as a template

for primers targeting the entire NanoLuc gene (top gel, 534 nt), or targeting just the NanoLuc sequence upstream

of the termination sequence (bottom gel, 234 nt) (right panel, with primers as indicated). Data shown is from one

experiment with biological triplicates. NTC = Non-template control, black triangles = primers, * = location of

termination sequence in pNL1.1. **(C)** Experimental design. HEK 293 cells are transfected with a control Firefly

reporter (pGL3) and a pNLP3 reporter expressed from a pol III promoter of interest. 24 h post-transfection, cells

are infected with wild-type (WT) γHV68 at an MOI of 1, or mock treated. Cell lysates are collected 24 h post-

transfection for the dual luciferase assay. Promoter activity was measured for the **(D)** U6 and **(E)** TMER1

promoters using a dual luciferase assay. Cells were harvested every 4 hours post-infection up to 24 h. Each

time-point was repeated for biological triplicates. Raw values for the Firefly and NanoLuc activity for each

promoter are shown as luminescence units (LUs). Promoter activity was analyzed as average relative

luminescence units (RLUs = NanoLuc LUs / Firefly LUs), and as the fold change in NanoLuc luminescence units

compared to the 4 h uninfected samples (NanoLuc fold change = Infected Sample NanoLuc LU / Mock Sample

NanoLuc LU). Error bars = SEM.

Figure 2. γHV68 infection induces activity of multiple pol III promoter types. Promoter activity was

measured for the U6 (n = 8), tRNA (n = 3), EBER1 (n = 3), EBER2 (n = 4), TMER1 (n = 18), TMER4 (n = 4), and

TMER5 (n = 6) promoters during infection. Luciferase assays were performed as previously described, with cell

lysates collected 24 h post-infection. Each independent experiment (n) contained biological triplicates or

duplicates. All promoter activity changes are analyzed as the fold change in NanoLuc activity normalized to
uninfected samples (Infected NanoLuc LU / Mock NanoLuc LU). Error bars = SEM. Significant differences
analyzed by t-test and indicated as asterisks. P-values are indicated as follows: * = P ≤ 0.05, **** = P ≤ 0.0001.
695

Figure 3. Inhibition of RNA polymerase III decreases γHV68 infection induction of luciferase activity from
most pol III promoter types. Promoter activity was measured for the indicted promoters during infection. Prior
to transfection with luciferase constructs, samples were treated with an RNA polymerase III inhibitor (CAS
577784-91-9), marked as P3I, or left untreated. Cells were co-transfected with the Firefly luciferase construct
expressed by the SV40 promoter and NanoLuc constructs expressed by the indicated promoters. Inhibited cells
were treated again immediately following infection. Luciferase assays were performed as previously described,
with cell lysates collected 24 h post-infection. Activities of the **A)** pol II promoters (orange, SV40-expressed Firefly
luciferase and TK-expressed NanoLuc) and **B)** NanoLuc luciferase expressed from pol III promoters (red) are
shown. All luciferase activity changes are analyzed as the fold change in NanoLuc activity normalized to
uninfected samples (Infected NanoLuc LU / Mock NanoLuc LU). NanoLuc RNA was also measured from **A)** a
pol II promoter (light blue, TK) and **B)** a pol III promoter (dark blue, TMER1). RT-qPCR was performed for
NanoLuc RNA as previously described. Cells were treated as described with cells transfected with a total of 200
ng or 2 µg of DNA. RNA was collected 24 hpi and analyzed for expression of NanoLuc RNA. Firefly activity from
SV40 was measured from two independent experiments with eight samples per experiment. RT-qPCR data for
200 ng are from two independent experiments with biological triplicates, and data for 2 µg are from three
independent experiments with biological triplicates or duplicates. Error bars = SEM. Significant differences
analyzed by t-test and indicated as asterisks. P-values are indicated as follows: * = P ≤ 0.05, ** = P ≤ 0.01, *** =
P≤ 0.001, **** = P ≤ 0.0001.

Figure 4. Sequence external to the minimal promoter elements alters promoter activity. (A) Several pol III
promoters were cloned into the pNLP3 NanoLuc-expressing vector. Promoter schematics are not to scale. The
canonical minimal promoter elements for each promoter are shown in blue and notated with a blue line labeled
"min", with the intermediate sequence in light grey. Upstream elements are green, and external/extended
sequence is shown in dark grey. Nucleotide positions are shown above each schematic in relation to position
"0", which indicates the start of the minimal promoter sequence. Genomic coordinates are shown below each
schematic in red, and refer to the following reference sequences; Human U6 (RNU6-1) = NC_000015.10, Human
tRNA-Tyr (TRY-GTA11-1) = NC_000007.14, EBERs in Human Herpesvirus 4/EBV Whole Genome =
NC_007605.1, TMERs in MHV68 Whole Genome = NC_001826.2. The coordinates for human U6 snRNA and
tRNA are reversed to reflect their orientation as coded on the complement strand. The predicted structure for
the minimal and full TMER1 promoter using the RNAfold Webserver are shown. **(B)** Cloned reporter constructs

were transfected into HEK293 cells at a 10:1 molar ratio with a Firefly luciferase control expressed by the SV40

promoter. TK-expressed NanoLuc was used as a positive control. Cells were not infected. Following 48 hours of

transfection, dual luciferase assays were performed as previously described. Luciferase activity is expressed as

the ratio of NanoLuc activity to control Firefly activity (relative luminescence units, RLUs). **(C)** Luciferase assays

were performed as previously described, with cell lysates collected 24 h post-infection. All promoter activity

changes are analyzed as the fold change in NanoLuc activity normalized to uninfected samples. Each

experiment (n) included biological triplicates. Error bars = SEM. Significant differences analyzed by t-test and

indicated as asterisks. P-values are indicated as follows: * = P ≤ 0.05, ** = P ≤ 0.01, *** = P≤ 0.001, **** = P ≤

0.0001.

Figure 5. Effect of γHV68 infection on NanoLuc transcript levels. Promoter activity was measured for **(A)**

the U6, TMER1, and EBER full promoters, or **(B)** the TMER1, TMER4, and TMER5 minimal vs. full promoters.

Cells were treated as previously described and RNA was purified from cells at 16 h post-infection (U6 n = 2,

TMER1, n = 5) or 24hpi (EBER1 n = 2, EBER2 n = 2, TMER4 n = 2, and TMER5, n = 2). RNA was converted to

cDNA, then LightCycler real-time PCR using Syber Green was performed with primers targeting the NanoLuc

gene and a host control gene (18S). The relative difference of NanoLuc was calculated using the Pfaffl method

(2001, *Nucleic Acids Research*), where the ratio = $(E_{target})^{\Delta CP \ target \ (control-sample)}$ / $(E_{ref})^{\Delta CP \ ref \ (control-sample)}$. Each

experiment (n) included biological triplicates or duplicates. Error bars = SEM. Significant differences analyzed

by t-test and indicated as asterisks. P-values are indicated as follows: * = P ≤ 0.05, ** = P ≤ 0.01, *** = P≤ 0.001,

**** = P ≤ 0.0001.

Figure 6. Effect of viral and NF-κB inhibitors and an NF-κB activator on luciferase activity during γHV68

infection. A) Promoter activity was measured via luciferase activity for the following promoters during infection,

including with concurrent PAA treatment: TMER1 (n = 5), TMER4 (n = 2), TMER5 (n = 2), EBER1 (n = 2), EBER2

(n = 3), U6 (n = 2), and tRNA (n = 1). Each experiment contained biological duplicates or triplicates. Luciferase

assays were performed as previously described, with cell lysates collected 24 h post-infection. NanoLuc fold

change for infected samples is from previous figures for comparison to PAA-treated samples. All promoter activity

changes are analyzed as the fold change in NanoLuc activity normalized to uninfected samples. **B)** Promoter

activity was measured via RT-qPCR of the NanoLuc transcript expressed from the TMER1 promoter during

infection and concurrent phosphonoacetic acid (PAA) treatment. RT-qPCR was performed as previously

described. Data is shown as NanoLuc relative difference to a host gene (18S). N = 2 with biological triplicates.

C) Using the same RNA from experiments shown in panel B, RT-qPCR was performed for a viral late gene (gB)

to confirm the efficacy of PAA treatment, which resulted in an approximately 32-fold decrease in gB compared

to WT infected samples. gB was not detected (N.D.) in mock samples. **D)** NF-κB activation with 50 ng/µL TNFα

(NF-κB reporter n = 1 for each time point, U6 n = 2, TMER1 n = 3) or **E)** NF-κB inhibition by 10 μM BAY 11-7082

(U6 n = 3, TMER1 n = 4). NF-κB activation is shown in Firefly luminescence units, while NanoLuc activity from

the U6 or TMER1 promoters is expressed as fold change over mock. Cells included in the TNFα experiments

were transfected for 12 h, while all others were transfected for 24 h as previously described. Each experiment

(n) contains biological triplicates. Error bars = SEM. Significant differences analyzed by t-test (two conditions) or

one-way ANOVA (three conditions) and indicated as asterisks. P-values are indicated as follows: * = P ≤ 0.05,

** = P ≤ 0.01, *** = P≤ 0.001, **** = P ≤ 0.0001.

Figure 7. Endogenous expression of murine U6 snRNA and 4.5S rRNA during γHV68 infection. Murine

fibroblast cells (3T12) were infected with mock, WT γHV68, or an EBER knock-in (EBER-KI) γHV68 at an MOI=1,

harvested at 16 hpi and subjected to the PrimeFlow RNA Assay and flow cytometric analysis. **(A,B)** Analysis of

U6 snRNA expression comparing background fluorescence ("No probe", left) versus U6 probe hybridization in

Mock (middle), or WT γHV68 or EBER-KI.γHV68 infected cells (right). **(C,D)** Analysis of 4.5S rRNA expression

comparing background fluorescence ("No probe", left) versus 4.5S rRNA probe hybridization in Mock (middle),

or WT γHV68 or EBER-KI.γHV68 infected cells (right). **(E)** Identification of virally-infected cells that express either

the TMERs or the EBERs, following infection with mock, WT γHV68 or EBER-KI.γHV68. Gates define cells based

on positive or negative expression as indicated, with these populations used for subsequent quantitation in

panels F and G **(F,G)** The geometric mean fluorescence intensity (gMFI) was calculated for **(F)** U6 snRNA or

(G) 4.5S rRNA in mock infected, or virus infected samples, comparing cells that differed in expression of either

the TMERs or the EBERs in WT or EBER-KI infections respectively. Data depict flow cytometric analysis of

singlets, defined by sequential discrimination using SSC and FSC parameters. Gating is shown in Supplemental

Figure 1. Data are from 3 biological replicates, with statistical significance assessed using an unpaired t-test.

Statistical significance as follows: **, p<0.01 and ***, p<0.001.

Figure 8. Endogenous expression of murine U6 snRNA during γHV68 infection. Murine fibroblast cells

(3T12) were infected with mock, WT γHV68, or an EBER knock-in (EBER-KI) γHV68 at an MOI=5, harvested at

16 hpi and subjected to the PrimeFlow RNA Assay and flow cytometric analysis. **(A,B)** Analysis of viral gene

expression indicating viral ncRNA (TMER or EBER) expression and/or ORF18 expression comparing

background fluorescence ("No probe", left) versus TMER or EBER hybridization in Mock (middle), or WT γHV68

or EBER-KI.γHV68 infected cells (right). **(C,D)** Analysis of viral gene expression indicating ORF18 expression

(top) or viral ncRNA (TMER or EBER, bottom) expression comparing background fluorescence ("No probe", left)

versus probe hybridization in Mock (middle), or C) WT γHV68 or D) EBER-KI.γHV68 infected cells (right). These

plots indicate the gating used for further analysis in E-F, and gating is duplicating from Supplemental Figure 2.

(E) The geometric mean fluorescence intensity (gMFI) was calculated for viral ncRNA (TMERs or EBERs) in

mock infected or virus infected samples, comparing cells that differed in expression of ORF18. **(F,G)** The gMFI

was calculated for U6 snRNA in mock or virus infected samples, comparing cells that differed in expression of

either **(F)** ORF18 or **(G)** TMERs or EBERs in WT or EBER-KI infections. Data depict flow cytometric analysis of

singlets, defined by sequential discrimination using SSC and FSC parameters. Gating is shown in Supplemental

Figure 2. Data are representative of two experiments, with 3 biological replicates in each. Statistical significance

assessed using a one-way ANOVA with multiple comparisons. Statistical significance as follows: * = P ≤ 0.05, **

= P ≤ 0.01, *** = P≤ 0.001, **** = P ≤ 0.0001.

Supplemental Figure 1. Viral ncRNA expression serves as an indicator of virus infection and correlates

with abundance of host pol III RNAs. Murine 3T12 fibroblasts were infected with mock, WT γHV68, or an

EBER knock-in (EBER-KI) γHV68 at an MOI=1, harvested at 16 hpi and subjected to the PrimeFlow RNA Assay

and flow cytometric analysis. **(A)** Sequential gating strategy used to identify singlets, with representative

examples from mock (top), WT γHV68 (middle) or EBER-KI.γHV68 infected samples. **(B)** Analysis of U6 snRNA

expression as a function of TMER expression comparing background fluorescence ("No probe", first panel) with

U6 probe hybridization in Mock (second panel), WT γHV68 (third panel) or EBER-KI.γHV68 infected cells (fourth

panel). **(C)** Analysis of U6 snRNA expression as a function of EBER expression comparing background

fluorescence ("No probe", first panel) with U6 probe hybridization in Mock (second panel), WT γHV68 (third

panel) or EBER-KI.γHV68 infected cells (fourth panel). **(D)** Analysis of 4.5S rRNA expression as a function of

TMER expression comparing background fluorescence ("No probe", first panel) with U6 probe hybridization in

Mock (second panel), WT γHV68 (third panel) or EBER-KI.γHV68 infected cells (fourth panel). **(E)** Analysis of

4.5S rRNA expression as a function of EBER expression comparing background fluorescence ("No probe", first

panel) with U6 probe hybridization in Mock (second panel), WT γHV68 (third panel) or EBER-KI.γHV68 infected

cells (fourth panel). Data depict flow cytometric analysis analyzing singlets, defined by sequential discrimination

using SSC and FSC parameters in panel A. Data are from 3 biological replicates.

Supplemental Figure 2. Viral RNA expression serves as an indicator of virus infection and correlates

with abundance of host and viral pol III RNAs. Murine 3T12 fibroblasts were infected with mock, WT γHV68,

or an EBER knock-in (EBER-KI) γHV68 at an MOI=5, harvested at 16 hpi and subjected to the PrimeFlow RNA

Assay and flow cytometric analysis. **(A)** Sequential gating strategy used to identify singlets, with representative

examples from mock with TMERs probe (top), WT γHV68 (second row), mock with EBERs probe (third row), or

EBER-KI.γHV68 infected samples (fourth row). **(B)** Analysis of U6 snRNA expression as a function of TMER or

EBER expression comparing background fluorescence ("No probe", first and second panel) with U6 probe

hybridization in Mock (third panel), WT γHV68 (fourth panel) or EBER-KI.γHV68 infected cells (fifth panel). **(C)**

Analysis of U6 snRNA expression as a function of ORF18 expression comparing background fluorescence ("No

probe", first panel) with U6 probe hybridization in Mock (second panel), WT γHV68 (third panel) or EBER-
KI.γHV68 infected cells (fourth panel). Data depict flow cytometric analysis analyzing singlets, defined by
sequential discrimination using SSC and FSC parameters in panel A. Data are from 3 biological replicates.

Table 1. Promoters analyzed in this study. Pol III promoters were selected from several organisms and cloned

into a luciferase reporter to analyze promoter activity during infection. Parenthetical additions specify promoter

features ("3A" = triplicated A box; for EBERs, "+ upstream" = includes upstream elements, "no upstream" =

excludes upstream elements).

Origin	Promoter	Promoter Type	Promoter Insert Length (nt)	Reference
Human	5S rRNA	Type 1	42	(54)
Human	tRNA-Tyr	Type 2	53	Gene ID: 100189507
Human	Vault RNA1-1	Type 2	65	(31, 55); GenBank: AF045143.1
Human	U6 snRNA	Type 3	227	GenBank: M14486.1
Adenovirus	VA1	Type 2	58	(56)
γHV68	TMER1 full	Type 2 (3A)	91	(22)
γHV68	TMER1 minimal	Type 2 (3A)	55	(22)
γHV68	TMER4 full	Type 2 (3A)	93	(22)
γHV68	TMER4 minimal	Type 2 (3A)	55	(22)
γHV68	TMER5 full	Type 2 (3A)	94	(22)
γHV68	TMER5 minimal	Type 2 (3A)	53	(22)
EBV	EBER1	Type 2 (+ upstream)	157	(57-59)
EBV	EBER1 minimal	Type 2 (no upstream)	70	(57-59)
EBV	EBER2	Type 2 (+ upstream)	156	(57-59)
EBV	EBER2 minimal	Type 2 (no upstream)	68	(57-59)

Fig 1

Knox et al

Pol II Induction

A)

Pol III Induction

B)

Fig 3
Knox et al

Fig 4
Knox et al

NanoLuc RNA

A)

B)

Fig 5
Knox et al

Fig 6

Knox et al

Endogenous U6 RNA expression

A) No probe | Mock | WT γHV68

AF750 / TMER (AF488) ; U6 (AF750) / TMER (AF488)

B) No probe | Mock | EBER-KI.γHV68

AF750 / EBER (AF647) ; U6 (AF750) / EBER (AF647)

Endogenous 4.5S RNA expression

C) No probe | Mock | WT γHV68

AF750 / TMER (AF488) ; 4.5S (AF750) / TMER (AF488)

D) No probe | Mock | EBER-KI.γHV68

AF750 / EBER (AF647) ; 4.5S (AF750) / EBER (AF647)

E) Mock | WT γHV68 | EBER-KI.γHV68

SSC-A / TMER (AF488)

SSC-A / EBER (AF647)

F) U6 RNA gMFI

**

Phenotype: Total | TMER- TMER+ | EBER- EBER+
Infection: Mock | WT gHV68 | EBER-KI. gHV68

G) 4.5S RNA gMFI

*** ***

Phenotype: Total | TMER- TMER+ | EBER- EBER+
Infection: Mock | WT gHV68 | EBER-KI. gHV68

Fig 7

Knox et al

Endogenous U6 RNA expression

Fig 8

Knox et al