Parallel and Population-specific Gene Regulatory Evolution in Cold-Adapted Fly Populations

Yuheng Huang[1*@], Justin B. Lack[1**], Grant T. Hoppel[1], and John E. Pool[1@]

[1]Laboratory of Genetics, University of Wisconsin-Madison, Madison, Wisconsin 53706

[*]Current address: Department of Ecology and Evolutionary Biology, University of California, Irvine, Irvine, CA 92697

[**]Current address: Advanced Biomedical Computational Science, Frederick National Laboratory for Cancer Research, Frederick, MD 21701

[@]e-mail: yuhenh3@uci.edu; jpool@wisc.edu.

Abstract

Changes in gene regulation at multiple levels may comprise an important share of the molecular

changes underlying adaptive evolution in nature. However, few studies have assayed within- and

between-population variation in gene regulatory traits at a transcriptomic scale, and therefore

inferences about the characteristics of adaptive regulatory changes have been elusive. Here, we

assess quantitative trait differentiation in gene expression levels and alternative splicing (intron usage)

between three closely-related pairs of natural populations of *Drosophila melanogaster* from

contrasting thermal environments that reflect three separate instances of cold tolerance evolution.

The cold-adapted populations were known to show population genetic evidence for parallel evolution

at the SNP level, and here we find evidence for parallel expression evolution between them, with

stronger parallelism at larval and adult stages than for pupae. We also implement a flexible method to

estimate *cis*- versus *trans*-encoded contributions to expression or splicing differences at the adult

stage. The apparent contributions of *cis*- versus *trans*-regulation to adaptive evolution vary

substantially among population pairs. While two of three population pairs show a greater enrichment

of *cis*-regulatory differences among adaptation candidates, *trans*-regulatory differences are more

likely to be implicated in parallel expression changes between population pairs. Genes with

significant *cis*-effects are enriched for signals of elevated genetic differentiation between cold- and

warm-adapted populations, suggesting that they are potential targets of local adaptation. These

findings expand our knowledge of adaptive gene regulatory evolution and our ability to make

inferences about this important and widespread process.

Introduction

Different species or populations often evolve similar phenotypes when adapting to similar

environments (Schluter 2000; Losos, 2011). Although such parallel phenotypic evolution can be

caused by amino acid changes, there is increasing evidence that regulatory mutations altering gene

expression underlie many cases of phenotypic evolution (Wittkopp & Kalay, 2012; Jones et al. 2012;

Stern 2013; Sackton et al. 2019). Most studies on gene regulatory evolution focus on expression

abundance (the number of transcripts for a whole gene). However, alternative splicing changes

resulting in modified transcript proportions can also contribute to adaptation (Barbosa-Morais et al.

2012; Gamazon and Stranger 2014; Smith et al. 2018), and yet splicing evolution has received far

less study.

The level of parallelism for gene expression evolution varies across study systems. In some taxa and

natural conditions, significantly more genes show parallel changes (repeatedly up- or down-regulated

in one ecotype relative to the other among independent population pairs) than anti-directional

changes (Zhao et al. 2015; Hart et al. 2018; Kitano et al. 2018; McGirr and Martin. 2018). However,

some other cases did not show significant parallel patterns, or they even showed anti-parallel patterns

(Derome et al. 2006; Lai et al. 2008; Hanson et al. 2017). The varying degree of parallelism may

partly be explained by the level of divergence among ancestors: more closely related ancestors are

expected to show a higher degree of parallel genetic evolution underlying similar phenotypic

evolution (Conte et al. 2012; Rosenblum et al. 2014).

Furthermore, gene expression evolution can be caused by the same or different molecular

underpinnings. Because of the difficulties of mapping expression quantitative trait loci (eQTLs), a

first step is to classify the expression evolution into two regulatory classes. *Cis*-regulatory changes

are caused by local regulatory mutations and result in allele-specific expression in a hybrid of

divergent parental lines (Singer-Sam et al. 1992; Cowles et al. 2002; Yan et al. 2002; Wittkopp et al.

2004). *Trans*-regulatory changes are caused by mutations at other loci. They modify the expression

of both alleles in hybrid diploids and do not result in allele-specific expression (Yvert et al. 2003;

Wittkopp et al. 2004; Wang *et al.* 2007). The relative importance of *cis*- and *trans*-effects to parallel

evolution varies among different studies systems (Wittkopp et al. 2008; McManus et al. 2010;

Wittkopp and Kalay 2012; Coolon et al. 2014; Lemmon et al. 2014; Nandamuri et al. 2018). Many

previous studies have focused on regulatory evolution between relatively distantly related lineages

such as different species, from which population genetic evidence of adaptive evolution may not be

available. Some studies have investigated the *cis*- vs. *trans*-regulatory variation within or between

recently diverged populations but are limited to one or two populations (Chen et al. 2015; Osada et al.

2017; Glaser-Schmitt et al. 2018). To our knowledge, the only two cases comparing *cis*- and *trans*-

regulatory changes for repeated adaptive divergence between populations are from threespine

stickleback fish and they revealed contrasting patterns (Hart et al. 2018; Verta and Jones 2019).

Hence, the relative contributions of *cis*- and *trans*-effects to recent parallel adaptation remain mostly

unknown.

In part driven by interest in the evolutionary response to climate change, *Drosophila* has been used as

a model system to study the genetic basis of thermal adaptation (Hoffmann et al. 2003). Because

temperature is an important environmental variable along latitudinal clines, clinal populations of

Drosophila melanogaster have been studied for decades (Adrion et al. 2015). Along these clines,

populations exhibit different degrees of cold tolerance in the expected direction, suggesting spatially

varying selection related to temperature (Hoffmann and Weeks 2007; Schmidt and Paaby 2008). The

recent development of genomics has allowed identification of clinal genomic variants, which are

candidates for thermal adaptation (e.g., Kolaczkowski et al. 2011; Fabian et al. 2012; Bozicevic et al.

2016; Mateo et al. 2018). There is also evidence of parallel evolution at the genomic and

transcriptomic level (Reinhardt et al. 2014; Bergland et al. 2015; Machado et al. 2015; Zhao et al.

2015; Juneja et al. 2016; Zhao and Begun 2017). Some of these studies compared clines between

species (which may have somewhat distinct biology), while others compared clines between

Australia and North America (which both feature primarily European ancestry with clinally variable

African admixture). Other transcriptomic studies have identified genes showing differential

expression between sub-Saharan African and European populations (e.g., Catalan et al. 2012;

Huylmans and Parsch 2014), which are separated by moderately strong neutral genetic differentiation

associated with the out-of-Africa bottleneck.

More broadly, populations of *D. melanogaster* from contrasting environments offer an excellent

opportunity to study parallel gene regulatory evolution and its underlying mechanisms. Originating

from a warm sub-Saharan ancestral range (Lachaise et al. 1988; Pool et al. 2012), *D. melanogaster*

has occupied diverse habitats, including environments with contrasting temperature ranges. There are

at least three instances in which the species expanded to cold environments: from Africa into higher

latitude regions in Eurasia, from Ethiopia lowland to higher altitudes, and from South Africa lowland

to higher altitudes. Populations were collected from these six regions, representing three warm-cold

population pairs: Mediterranean pair (MED), collected in Egypt (EG, warm) and France (FR, cold);

Ethiopian pair (ETH) collected in Ethiopia lowland (EA, warm) and highland (EF, cold); and South

Africa pair (SAF), collected in South Africa lowland (SP, warm) and highland (SD, cold).

Importantly, each of these population pairs has the advantage of low genetic differentiation between

its warm- and cold-adapted members compared to the differentiations among pairs (Pool et al. 2017).

Although the cold populations have invaded colder habitats for only ~1000-2000 years (~15k-30k

generations) (Sprengelmeyer et al. 2020) and different habitats have distinct selective pressures

besides cold (e.g., air pressure, ultraviolet radiation, food resources), the cold-dwelling populations

have shown signals of parallel adaptation for cold tolerance and allele frequency changes (Pool et al.

2017). In the present study, this unique system allows us to assess the degree of parallelism for

transcriptomic changes underlying parallel adaptation to colder environments.

Because the selection environments can vary drastically across life stages of *Drosophila*, we may
expect to see different patterns of local adaptation and parallelism in gene expression across stages.
For *D. melanogaster*, the larvae are mostly located within fruit and their primary role is feeding. The
pupae are located on or near the fruit and are immobile. The adults are mobile; their primary role is
mating and reproduction (Powell, 1997; Sokolowski et al. 1986), and it is thought to be the primary
overwintering stage in seasonally cold environments (Lzquierdo 1991). And a recent study using *D.
melanogaster* populations across the globe found local adaptation to thermal environments at egg,
larval and adult stages but not the pupal stage (Austin and Moehring 2019). Therefore, we may
expect a different level of parallel gene expression evolution for thermal environments for the pupal
stage.

Here, we generate RNA sequencing (RNA-seq) data for multiple outbred genotypes from each of the
six population samples listed above, from larval, pupal, and adult stages. We estimate gene
expression and alternative intron usage levels for each sample, then identify cases of unusually high
quantitative trait differentiation between each pair of warm- and cold-adapted populations and
compare their genomic locations across developmental stages. We find that genes with highly
differentiated expression are enriched on the X chromosome in the adult stage relative to the larval
stage. We find evidence for parallel evolution for expression for both the larval and the female adult
stages, but less parallel signal for the pupal stage. We further tease out the *cis*- and *trans*-regulatory
effect at the adult stage by sequencing the transcriptomes of the parental lines from different
populations and their F1 offspring. Applying our resampling approach to study *cis*- and *trans*-
regulatory effects, we find that the relative contributions of these effects to adaptive expression
differentiation is quite variable across population pairs, with *trans*-effects showing greater
parallelism. Finally, we observe enrichments of genes with high F_{ST} among those that showed *cis*-
effects and identify several candidate genes with both *cis*-effects and high F_{ST}, as potential targets of
local adaptation.

Methods and Materials

Ecologically and phenotypically differentiated populations

The three *Drosophila melanogaster* cold-warm population pairs used in this study, France-Egypt
(MED), Ethiopia (ETH) and South Africa (SAF), were described in previous publications (Pool et al.

2012; Lack et al. 2015; Pool 2017). Previous study has shown that female adults from the cold

populations (FR, EF and SD) were more likely to recover after 96 hours at 4°C than the respective

warm populations (Pool et al. 2017). To extend these results, three inversion-free strains from each of

the cold populations as well as an ancestral warm adapted population (ZI) were used to measure egg

to adult viability at different temperatures. Viability was assayed at 15°C as the cold environment

and 25°C as the warm control environment. 40 mated female flies were allowed to lay eggs in a half

pint glass milk bottle with a standard medium at room temperature for 15 hours. Each strain occupied

~8 bottles. After the flies were removed and the numbers of eggs were counted, about half of the

bottles were incubated at 25°C and the other half 15°C. The numbers of adult flies that emerged from

each bottle were counted after 14 days and 42 days from warm and cold environments respectively.

Viability for each strain was measured as the average emergence proportion among bottles, which is

the number of emerged adults divided by the number of eggs. To determine significance, unpaired t-

tests between strains from each cold population and those from the ZI population were performed for

both temperature conditions.

RNA sample collection and sequencing

Within each population of the three warm/cold pairs (six populations in total), we selected 16 strains

and assigned them into eight crosses (Fig. 1). Before the crossing, all the strains had been inbred for

eight generations. The criterion for choosing parental strains for a cross was based on minimal

genomic regions of overlapping heterozygosity. Among the strains chosen within each population,

we used similar criteria to select four strains to perform crosses between the warm and the respective

cold populations. Two of the four strains were used as the maternal lines and the other two were used

as paternal lines in the between-population crosses. One cross between SD and SP populations was

lost. We also collected adult female samples from the parental inbred lines used in the between-

population crosses. Inversion frequencies are known to differ between these populations (Pool et al.

2017) and inversions have been associated with expression differences (Lavington & Kern 2017;

Said et al. 2018). While inversions are not an explicit focus of our study, they may contribute to

population expression differences. The inversion information for the strains used can be found in

Table S1.

Fig 1. Illustrations of the geographic origins of the three population pairs and the crossing design. (A) A map of average year-round temperature (°C) showing the geographic origins of each population sample studied, and their groupings into pairs of closely-related warm- and cold-derived population samples. The Mediterranean (MED) pair comprises a cold-derived France population (FR) and a warm-derived Egypt population (EG). The Ethiopian (ETH) pair comprises a cold-derived high-altitude population (EF) and a warm-derived low-altitude population (EA). Likewise, the South African (SAF) pair comprises a cold-derived high-altitude population (SD) and a warm-derived low-altitude population (SP). The location of an additional warm-derived population from Zambia (ZI), within the species' putative ancestral range, is also indicated. (B) Schematic figure showing the crossing design for one population pair (MED) as an example. Within-population crosses generated controlled outbred offspring for estimating P_{ST} to quantify population differentiation in gene expression; samples at three developmental stages (third instar larva, pupa and female adult) were collected from each cross. Parental inbred strains from warm- and cold-adapted populations and inter-population crosses between them were studied to estimate *cis*- and *trans*-regulatory effects that underlie the expression divergence; samples from female adults were collected.

All the flies were reared at 15°C, which approximated the derived cold condition. 20 virgin females and 20 males were collected from maternal and paternal lines respectively for each cross and allowed to mate and lay eggs for a week in half-pint bottles. Each bottle contained standard *Drosophila* medium (containing molasses, cornmeal, yeast, agar, and antimicrobial agents). For the within-population crosses, samples at three developmental stages were collected: larva, pupa and female adult. Third-instar larvae were collected on the surface of the medium. For pupa, new yellow pupae were collected within one day of pupation. For adult, female flies were collected 4-5 days after eclosion. For samples from between-population crosses and parental lines, only female adults were collected. All the samples were shock-frozen in liquid nitrogen immediately after collection.

Approximate 50 larvae or 50 pupae or 30 female adults were used for RNA extraction for each

sample. Total mRNA was extracted using the Magnetic mRNA Isolation Kit (New England Biolabs,

Ipswich, MA) and RNeasy MinElute Cleanup Kit (Qiagen, Hilden, Germany). Strand-specific

libraries were prepared using the NEBNext mRNA Library Prep Reagent Set for Illumina. Libraries

were size-selected for approximately 150 bp inserts using AMPureXP beads (Beckman Coulter, CA,

USA). The 179 libraries were quantified using Bioanalyzer and manually multiplexed for

sequencing. All libraries were sequenced on a HiSeq2500 (V4) with 75bp paired-end reads in two

flow cells. Numbers of paired-end reads generated for each library can be found in Table S2.

Quantifying gene expression and exon usage frequency

The paired-end sequence reads for the within-population cross samples were mapped to the

transcribed regions annotated in the *D. melanogaster* reference genome (release 6, BDGP6.84) using

STAR with parameters from ENCODE3's STAR-RSEM pipeline (Li and Dewey 2011; Dobin et al.

2013). We note that cold- and warm-derived members of each population pair are expected to have

very similar genome-wide reference sequence divergence (Lack et al. 2016a). For gene expression,

the numbers of reads mapped to each gene were quantified using RSEM (Li and Dewey 2011). Reads

mapped to the rRNA were excluded in the analysis. The expression abundance for each gene was

standardized by the numbers of reads mapped to the total transcriptome of the sample.

To quantify exon usage, we used Leafcutter (Li et al. 2018) to estimate the excision frequencies of

alternative introns. This phenotype summarizes different major splicing events, including skipped

exons, and 5' and 3' alternative splice-site usage. Leafcutter took the alignment files generated by

STAR as input to quantify the usage of each intron. Then Leafcutter formed clusters that contain all

overlapping introns that shared a donor or accept splice site. The default parameters were used: ≥ 50

reads supporting each intron cluster and $\leq 500kb$ for introns length. The exon usage frequency is the

number of intron excision events divided by the total events per cluster. It is worth noting that

Leafcutter only detects exon-exon junction usage and it is unable to quantify 5' and 3' end usage and

intron retention (Alasoo et al. 2019), which were not examined here.

Principal component analysis

To visually assess the overall patterns of variation in the transcriptomes among samples, we first

performed principal component analysis (PCA) for the within population cross samples across three

developmental stages using *DESeq2* (Love et al. 2014). The *DESeq* dataset object was constructed

from the matrix of the count data outputted from *RSEM*. After the variance stabilizing transformation

(vst), the top 5000 genes with highest variance across samples at the transformed scale were used for

PCA. The principal component value for each sample was obtained by the function *plotPCA* (Fig.

S2). We also performed principle component analysis for samples at each developmental stage. For

the adult stage, we included the F1 offspring from crosses within populations, F1 offspring from

crosses between populations and the inbred parental lines of the latter crosses.

Identifying outliers in gene expression and intron usage differentiation using within-population

crosses

To identify candidate genes under differential evolution between the warm and cold populations in

each pair, we first controlled for the potential transcriptome skew caused by very highly expressed

genes. For each expressed gene, we calculated the average expression of the cold samples

($AvgExp_{cold}$) and that of the warm samples ($AvgExp_{warm}$). Then we obtained the median of the ratio of

$AvgExp_{cold}/AvgExp_{warm}$ across all expressed genes for the population pair. Gene expression for the

warm samples was normalized by multiplying this median before subsequent analysis. This

correction was designed to avoid a scenario in which either the cold population or the warm

population had important expression changes in one or more highly expressed genes that caused the

relative expression of all other genes to shift, even if their absolute expression level did not.

We used P_{ST} statistics to quantify gene expression divergence between cold and warm populations in

each population pair using samples from within-population crosses:

$$P_{ST} = \frac{V_{between}}{V_{between} + 2V_{within}}$$

where $V_{between}$ is between-populations variance for expression abundance and V_{within} is the average

variance for expression abundance within populations. Although both within- and between-

population components of variance can be confounded by the environmental variance, P_{ST} is still a

useful statistic to quantify phenotypic differentiation (Lande 1992; Spitze 1993; Merila 1997;

Brommer 2011; Leinonen et al. 2013). Here, environmental variance should be reduced by the

common laboratory environment. To reduce sampling variance before calculating P_{ST}, for each gene,

we required the total mapped reads across all 48 within-population samples to exceed 200 for a given

developmental stage. Then for each population/stage, we excluded the crosses/samples with the

highest and lowest gene expression for each gene (to avoid high P_{ST} values being driven by single

anomalous values), resulting in six samples per population/stage. The P_{ST} quantile based on data

excluding extreme samples is concordant with the P_{ST} quantile calculated using all the crosses for

most cases (Fig. S3).

We chose the above P_{ST}-based approach instead of simply testing for differential expression in part

because our within-population samples reflect real variation as opposed to technical replicates. Also,

many alternative methods make assumptions about the data (e.g., negative binomial distribution for

transcript counts) which are difficult to apply to splicing, even if they hold for expression. P_{ST} and

the population genetic index F_{ST} are under the same theoretical framework, and are often directly

compared to search for evidence of adaptive trait differentiation. However, environmental and

measurement variance will downwardly bias P_{ST}, making targets of local adaptation less likely to

reach a threshold defined by genome-wide high F_{ST} outliers. Hence, in this study we simply focus on

the highest quantiles of P_{ST} for a given trait/population comparison, as detailed below.

As with gene expression, we used P_{ST} to estimate the intron usage differentiation between cold and

warm populations, with $V_{between}$ as the between-population variance for a given intron's usage

frequency, V_{within} as the average within-population variance for intron usage frequency. Before

calculating P_{ST}, for each exon-exon junction, we averaged the intron excision events (n_i) and the

alternative events (n_j) of the cluster across all samples in a developmental stage. The minimum for

both types of event had to be at least 5 ($n \in [n_i, n_j] \geq 5$). We also required that at least six samples

have intron usage count > 0 in each population for the exon-exon junction to be included in

subsequent analysis. Then for each exon-exon junction, we excluded the sample with highest and

lowest intron usage in a population/stage and calculated P_{ST}.

Examining the relative contribution of the X chromosome to population differentiation across

stages

For gene expression differentiation, we used the upper 5% quantile of P_{ST} as outlier cutoff to identify

candidate genes potentially under geographically differential selection. Then we calculated the

fraction of the outliers located on the X chromosome (f_x). To generate the null distribution of f_x, we

permuted the genes used in calculating P_{ST} and calculated f_x' for the top 5% of the permuted gene set.

This process was repeated 10,000 times to obtain a null distribution of f_x'. The upper 2.5% and lower

97.5% quantile of f_x' define the 95% confidence interval. To test whether the actual f_x is significantly

different from the null, the p-value is calculated as two times the proportion of the f_x' that were equal

or more extreme than the actual f_x (two-tailed test).

To test whether the developmental stage impacts the enrichment of outliers on the X chromosome,

we analyzed the fraction of genes on the X chromosome (f) using linear model (*lm* function) in R

(version 3.6.3):

$$f_x = \text{Stage} + \text{Type} + \text{Stage x Type} + \text{Pair}$$

where Stage is larvae/pupal/adult. Type is outliers or nonoutliers. Pair is the population pair. As we

are interested in whether the difference in *f* between outliers and the background depends on the

development stage, the model above was compared to a reduced model without the interaction term

Stage × Type using likelihood ratio test (*anova* function with test = "LRT"). If the Stage × Type for

the full dataset was significant, we performed the same analysis separately for larva-pupa, pupa-adult

and larva-adult datasets to determine which stages caused the significant Stage × Type effect.

Comparing P_{ST} outliers with published data for African and European populations

To study whether the adult fly in MED pair changes expression in similar ways as other African and

European population comparisons, we first obtained lists of candidate genes showing significant

differential expression between African and European populations for adult samples from Muller et

al. 2011 (padj < 0.05) and von Hecket et al. 2016. We calculated the numbers of P_{ST} outliers that are

shared with the published lists with between-populations differential gene expression (in the same

directions of expression changes). Then we permuted all the genes we tested in the MED pair and

selected the same number of genes as the true outliers randomly. We asked how many genes in the

randomly permuted list are shared with the published lists. We repeated the process 10,000 times to

obtain a null distribution of the shared numbers. The p-value was calculated as the proportion of the

null distribution that was equal or more than the actual number of shared genes (one-tailed test).

Another test was whether the shared genes for P_{ST} outliers were more likely to change expression co-

directionally with the published lists than the non-outliers, which was tested by a Chi-square test.

Examining co-directional change for outliers shared between population pairs

To study the degree of parallel evolution in gene expression, we identified outlier genes shared

between two population pairs and showing consistent changes in the cold populations relative to the

warm ones (co-directional). Whether the number of shared outliers with co-directional change was

significantly different from expected by chance was determined by a permutation-based test. For the

outlier genes in a certain pair, we calculated the number of these genes (N) that were shared and

changed expression in the same direction in the outliers from another pair. To generate the null

distribution of N, we permuted the genes used in calculating P_{ST} and obtained a set of genes that pass

a certain quantile in each pair. Then the N' was calculated based on the two permuted sets of outliers.

This process was repeated 10,000 times to obtain a null distribution of N'. To determine the

statistical significance, a p-value was calculated as two times the proportion of the N' that were equal

or more extreme than the actual N (two-tailed test). The statistics here and those below assume the

expression changes are independent among genes/introns, which is not always the case (genes can

interact with each other via regulatory networks). We performed similar tests for pairwise

comparisons between developmental stages for each population pair. The numbers for shared outliers

with consistent changes between pairwise stages were reported in Table S8.

The second approach used to examine parallelism of gene expression evolution was to focus on the

outlier genes for a specific population pair and examined whether the expression changes in another

pair followed the same directions. If cold adaptation causes similar evolution in gene expression,

those genes should tend to show changes in the same directions in both pairs. Each of the pairwise

population combinations had two comparisons: the outliers can come from either pair. For the outlier

genes in a certain pair, we calculated the fraction (F) of these genes changing expression in the same

direction in another pair. To generate the null distribution of F, we permuted the genes used in

calculating P_{ST} and calculated F' for the permuted genes that pass a certain quantile. This process

was repeated 10,000 times to obtain a null distribution of F'. The upper 2.5% and lower 97.5%

quantile of the distribution define the 95% confidence interval. The p-value was calculated as two

times the proportion of the F' that were equal or more extreme than the actual F (two-tailed test).

To identify intron usage outliers, a cutoff of the upper 5% P_{ST} was used. If multiple exon junctions

had P_{ST} above the top 5% cutoff, only the exon junction with the highest P_{ST} would be kept as an

outlier to control for nonindependence. Because the numbers of shared intron usage outliers in both

population pairs are small (<10), we only performed the second type of analysis described above. For

a certain developmental stage, we used the top 5% outlier intron usage in a particular pair and asked

what percentages of the intron usage changed co-directionally in another pair. To determine the

statistical significance, we used the permutation approach as described above.

GO enrichment test for P_{ST} outlier genes

The Gene Ontology enrichment tests were performed using the R package "clusterProfiler" (Yu et al.

2012) based on the fly genome annotation (Carlson 2018). The types of GO terms being tested

contained all three sub-ontologies: Biological Process (BP), Cellular Component (CC) and Molecular

Function (MF). Selection of overrepresented GO terms was based on adjusted p-value < 0.1 using

the "BH" method (Benjamini and Hochberg 1995) for each sub-ontology. This relaxed p-value

threshold (after accounting for multiple testing) was used in light of the hypothesis-generating goals

of this analysis. For gene expression, the upper 5% P_{ST} outliers were tested for GO enrichment

relative to all the expressed genes for each population pair for a certain stage. To determine whether

the shared significant GO terms between pairs were more than expected by chance, we randomly

sampled the same numbers of genes as the outliers and performed the GO test for both pairs and

identified the shared significant GO terms between pairs. We repeated the process 1000 times to get a

set of numbers for the shared significant GO terms and compared to the actual number of shared

significant GO terms to get a permuted p-value.

To access the functional categories of the differential intron usage, we calculated the quantile of P_{ST}

for each alternative intron's usage. To rank the differentiation for a gene, we used the highest

quantile (the most extreme differentiation) among the intron usages within the gene as the gene

quantile (q_{gene}). To account for the multiple testing of the intron usages for a gene, the adjusted total

numbers of testing is calculated as $n_{sum} = \sum_{i=1}^{i=j}(n_i - 1)$, where n_i is the number of testing for a

cluster and *j* is the number of clusters for the gene. Then, the adjusted gene quantile is $q'_{gene} = 1 - (1 -$

$q_{gene})$ x n_{sum}. The upper 5% q'_{gene} was used to identify the most differentiated genes for intron usage

and they were tested for GO enrichment as described above.

Estimating cis- and trans-effects of regulatory divergence using between-population crosses and

parental strains

To study the contributions of *cis-* and *trans*-regulatory effects on expression and intron usage

divergence, we focused our analysis on the upper 5% P_{ST} outliers for gene expression/intron usage.

For each gene/intron junction in each population pair, we selected a representative cross showing the

greatest difference between parental strains for this analysis. In addition, this difference needed to be

larger than the average difference between the cold and warm populations from the outbred crosses

for its pair.

To study allele-specific expression/intron junction usage, we obtained the genomic sequences of the

two parental strains aligned separately to the FlyBase *D. melanogaster* 5.77 assembly (Lack et al.

2015; 2016a). The SNP calling from the reference genome was done by samtools (Li et al. 2009). To

avoid mapping bias for the RNAseq reads (Degner et al. 2009; Stevenson et al. 2013), we updated

the reference based on the SNPs for the two parental stains by masking the SNPs as "N". The F1

female adult RNA-seq reads were mapped to the updated reference using STAR with options: --

chimFilter None --outFilterMultimapNmax 1 (Dobin et al. 2013). Because of the fairly high level of

heterozygosity within our inbred lines (Lack et al. 2015), we attempted to use polymorphic sites to

study the allele-specific expression instead of focusing on the fixed difference between parental

strains. However, based on the simulations we performed (see supplementary document), our new

method requires a large number of F1 offspring (>300 per cross) to reduce the random sampling of

parental alleles. For this experiment (only 30 F1 offspring per cross), we therefore used SNPs that

were fixed differences between the parental strains. SNPs were filtered with read counts ≥ 10 in the

F1 RNA-seq sample and the parental samples. Then the allele frequency in the RNA reads for the F1

sample was calculated to estimate allelic expression proportion. The allelic expression proportion for

each candidate gene p_{F1} was the median average allele frequency for all sites located in the gene

region.

We tested two null hypotheses corresponding to *cis*-only and *trans*-only regulatory differences using

a resampling approach. Under the null hypothesis that *cis*-regulatory effects are absent, the p_{F1} is

expected to be near 0.5 because the cold parental strain contributes half of the alleles to F1 offspring,

and alleles from different parents are expressed similarly in these F1s (Cowles et al. 2002; McManus

2010; Meiklejohn et al. 2014). Under the null hypothesis that *trans*-regulatory effects are absent, p_{F1}

is expected to approximate the ratio of the cold parental strain expression to the total expression of

both parental strains (Wittkopp et al. 2004): $r_{F0} = E_c/(E_c + E_w)$. However, sampling effects can cause

p_{F1} to deviate from the null expectations.

We accounted for different types of uncertainty on estimating p_{F1}. To account for the measurement

uncertainty in F1 expression, we sampled with replacement for the F1 reads mapped to each gene

until we reached the numbers of reads mapped to the gene. Then we recalculated the p_{F1}' for each

SNP and then averaged across sites for each gene. We repeated the above process 1000 times to get a

distribution of p_{F1}'. A 95% confidence interval of the distribution not overlapping with 0.5 suggested

the existence of a *cis*-effect.

To test for a *trans*-effect, the uncertainty when estimating the expression level in parental strains also

needs to be accounted for. For each gene/intron in a parental strain, we used binomial sampling based

on the expression level of the gene/intron. The sampling probability is the proportion of reads for that

gene/intron relative to total reads in a sample and the number of sampling events equals the total

reads of the sample. Then we had the updated expression for the cold strain E_c' and the warm strain

E_w'. The updated r_{F0}' is calculated as E_c'/(E_c'+ E_w'). The sampling and calculation were repeated

1000 times. Each time the r_{F0}' was paired with a p_{F1}' described above to calculate the difference D' =

r_{F0}' - p_{F1}'. A 95% confidence interval of D' not overlapping with 0 suggested the existence of a

trans-effect.

To test the specificity and sensitivity of this approach, we performed simulations to generate

expression read data and apply our method on the simulated data (supplementary document). We

found that our approach has good performance under reasonable conditions and can be adapted for

other traits, such as splicing. For splicing, the p_{F1} is the allele frequency for the diagnostic SNPs

located in the exon-junction and the r_{F0} is the ratio of the cold parental strain intron usage frequency

to the sum of the frequencies for both parental strains.

Based on the tests above, the set of candidate genes were classified into categories including no

significant *cis*- or *trans*-effect, cis only, and trans only (McManus 2010; Schaefke et al. 2013; Chen

et al. 2015). For genes showing both *cis*- and *trans*-effects, we further classified them based on

whether these two effects favored expression of the same (co-directional) or different parental allele

(anti-directional). For exon usage differentiation, we applied a similar approach to classify the

differentiated exons into the five categories, accounting for different sampling effects and

measurement errors. Instead of analyzing expression level of the parental strains (*E*), we analyzed

their intron usage frequency for the sets of outlier intron junctions.

Examining gene expression co-regulation among outliers for adult

To study the level of co-regulation among outliers related to cold adaptation, we focused on the

expressions of the outliers in the cold-derived populations. For each pairwise combination of two

outliers, we calculated the correlation coefficient of the expression values among the eight outbred

samples. To test whether the correlation coefficient is different from random expectation, we

permuted the eight outbred samples randomly for one gene for each gene combination, requiring at

least five of the eight samples to be changed. Then we calculated the correlation coefficient between

genes with the permuted samples. We repeated the process 10,000 times to obtain a null distribution

of the correlation coefficient. The p-value was calculated as the proportion of the null distribution

that was equal or more than the actual coefficient (one-tailed test). We used $p < 0.05$ as a cutoff to

identify significantly co-regulated outlier pairs. To compare the level of co-regulation between

populations, we calculated the proportion of co-regulated pairs were significantly co-regulated for

each population as well as the number of significant co-regulated partners for each gene in each

population.

Examining between-population genetic differentiation for genes with cis-effect

For the P_{ST} outliers identified with significant *cis*-effects, we hypothesized that causative *cis*-

regulatory elements may show elevated allele frequency differentiation between the warm and cold

populations. For expression abundance, the majority of cis-regulatory SNPs are located within 2kb

upstream of the transcription start site and downstream of the transcription end site (Massouras et al.

2012). Therefore, we used the interval from 2kb upstream to 2kb downstream as the focal region of a

gene for this analysis. We calculated window F_{ST} and SNP F_{ST} using sequenced genomes from

Drosophila Genome Nexus (Lack et al. 2015 & 2016a). For window F_{ST}, the division of windows

within a gene region was based on 250 non-singleton variable sites per window in the ZI population

(Pool et al. 2017). Each window needed to have at least five genotypes for each population. Before

assigning window F_{ST} to the focal genes, we confirmed that there is no large chromosomal scale of

differentiation between populations for each pair (Fig. S8). The highest F_{ST} for the windows

overlapping the focal region was assigned as its F_{ST_winmax}. To determinate the statistical significance

of F_{ST_winmax}, we calculated F_{ST_winmax} for all other blocks of the same number of windows (to account

for gene length) along the same chromosome arm where cross-over rates were above 0.5cM/Mb

(Comeron et al. 2012), but excluding those within 10 windows of the focal region. The specific non-

low recombination regions are: 2.3–21.4 Mb for the X chromosome, 0.5–17.5 Mb for arm 2L, 5.2–

20.8 Mb for arm 2R, 0.6–17.7 Mb for arm 3L, and 6.9–26.6 Mb for arm 3R. SNP F_{ST} was

calculated for sites with at least 10 alleles for each population. The highest value (F_{ST_SNPmax}) within

the focal region was thus obtained for the focal gene. Analogous to our F_{ST_winmax} permutation, we

also calculated F_{ST_SNPmax} for permuted regions with the same number of SNPs as the focal region,

along the non-low cross-over rate region on the same chromosome arm. For both F_{ST_winmax} and

F_{ST_SNPmax}, we then focused on regions in the upper 5% quantile of permuted values for further

analysis. To test whether the expression outliers with cis-effects are enriched for high F_{ST} outliers, we

analyzed the fraction of F_{ST} outliers (*f*) using a linear model (lm) in R:

$$f = \text{Type} + \text{Pair}$$

where Type is outliers with *cis*-effect (*cis*-outlier) or non-outliers, and Pair is the population pair. The

model above was compared to a reduced model without the Type term using likelihood ratio test

(anova function with test = "LRT"). For outlier genes that showed a *cis*-effect and high F_{ST}, location

and functional information were obtained from Flybase (Thurmond et al. 2019).

Results

Gene expression differentiations between warm- and cold-derived populations

In a cold environment (15°C), we found the FR and EF populations have significantly higher egg-to-

adult viability than an ancestral range population and the SD population follows the same trend (Fig.

S1, Table S3). In contrast, at a 25°C benign temperature all of the populations have relatively high

survival (75%). These findings were consistent with past results (Pool et al. 2017) in suggesting that

the cold-derived populations have adapted to low temperature.

We then surveyed the transcriptomes of larvae, pupae, and female adults for multiple genotypes from

each cold- and warm-adapted population using high-throughput RNA sequencing (RNA-Seq). To

focus on the transcriptomes of outbred genotypes, we generated eight within-population crosses from

each population under the derived cold environment (15 °C). We pursued this outbred strategy to

guard against inbreeding effects amplifying within-population regulatory variation and therefore

dampening estimates of regulatory differentiation between populations (estimated using the

quantitative genetic index P_{ST}). Comparing our inbred adult RNA-Seq data against outbred data from

a subset of the same strains, we found results that were mostly consistent with that expectation: five

out of six populations showed greater expression variance in inbred than outbred data, and two out of

three population pairs showed higher P_{ST} values from outbred than inbred data (Table S6). We

therefore focused on outbred data for subsequent population comparisons.

After performing PCA on normalized expression values, we found that the first and second principle

component of gene expression showed clear signals of developmental stages among samples, while

signals of transcriptome-wide population differentiation was more modest (Fig. S2). We then used

P_{ST} to quantify phenotypic differentiation of expression and splicing between populations in each

pair. P_{ST}, analogous to F_{ST} for genetic variation, measures the amount of trait variance between

populations versus total variance for a phenotype (Merila 1997; Brommer 2011; Leinonen et al.

2013). The genes/intron usages with highest P_{ST} quantiles are more likely to be under ecological

differential selection between populations than those with lower P_{ST} quantiles (Leder et al. 2015).

Genes were filtered for analysis based on ≥ 200 counts across all 48 within-population samples (8

samples per population, six populations in total for three pairs). The numbers of genes that passed the

filters for analysis were: 4699 genes for larva, 5098 genes for pupa and 6785 genes for adult. We

initially observed that in the ETH pair, the adult sample showed a general shift in transcriptome-wide

relative abundances between populations, caused by a few highly expressed genes (Fig. S4). Further

investigation suggested that the EF population was primarily responsible for the observed ETH

asymmetries in adults (Fig. S5). Many of the highly expressed genes in the cold-adapted and larger-

bodied EF population (Lack et al. 2016b) are related to muscle protein (Table S4). To correct for the

influence of such changes on relative expression levels, we standardized the expression values of

warm-derived populations by the median expression ratio between cold- and warm-derived

populations, resulting in about the same numbers of genes with increased and decreased expression

in the cold-derived population relative to the warm-derived one transcriptome-wide (Fig. S4). To

study gene expression divergence potentially under ecologically differential selection, we calculated

P_{ST} (Materials and Methods). The P_{ST} values for all genes for each population/stage are listed in

Table S4. We used the upper 5% quantile of P_{ST} as outliers for each population pair. For the outliers,

there is a strong directionality on the expression difference between populations for the ETH pair

(Fig. 2): a large majority of ETH P_{ST} outliers had higher expression in the cold-adapted EF

population in larvae and especially adults, with pupae showing a reversed pattern. These asymmetries

mirrored transcriptome-wide skews in expression proportion for this population pair, in that

substantially more genes had higher EF expression than the modal proportion (Fig S4). These

observations suggest unique regulatory features for the populations in the ETH pair, perhaps hinting

that many outlier genes might be co-regulated.

540
Figure 2. The relationship between expression differences between populations and P_{ST} illustrates
asymmetric expression differences in some population pairs / developmental stages. The x-axis is the
cold-derived population expression proportion, which is the ratio of mean expression of the cold-
derived population relative to the sum of mean expression values from the two populations.
Proportion higher than 0.5 (red vertical dashed lines) indicate a higher expression for the cold-
derived population and the warm-derived one. The blue horizontal dashed lines show the upper 5%
quantile of P_{ST}.

Comparing P_{ST} outliers with published studies

Since there are studies comparing whole-body transcriptomes between African and European

populations in this species (Muller et al. 2011; von Heckel et al. 2016), we examined whether our P_{ST}

outliers for adults from the MED pair overlapped with the candidates showing differential expression

between African and European populations from the published datasets. While each of these analyses

might detect regulatory evolution that occurred in Europe, we emphasize the distinctness of these

geographic comparisons, since our Egypt sample differs substantially from the Zimbabwe population

featured in those studies. For upper 5% quantile of P_{ST} outliers, we do not see more sharing with the

previous candidates from either study than random expectations (permutation test, p = 0.47 for

comparing with Muller et al. 2011 and p = 0.39 for comparing with von Heckel et al. 2016).

However, for the upper 10% of P_{ST}, the outliers of MED are more likely to be shared with the

candidates from Muller et al. 2011 than random expectations (permutation test, p = 0.0033) but not

with those from von Heckel et al. 2016 (permutation test, p = 0.47). Moreover, among all the shared

genes, our outliers were more likely to show gene expression change in the same directions as the

previous candidates than the non-outliers (χ^2 = 6.3, df = 1, p = 0.012 for comparing with Muller et al.

2011, χ^2 = 7.4, df = 1, p = 0.0065 for comparing with von Heckel et al. 2016).

X chromosomal and autosomal contributions to regulatory evolution

To investigate the contribution of autosomes versus the X chromosome to the expression

differentiation due to cold adaptation, we surveyed the locations of P_{ST} outliers for different

developmental stages (Fig. 3). At the larval stage, the proportions of P_{ST} outliers located on the X

chromosome were lower than the genome-wide level in each population pair (permutation test, p =

0.72 for MED; p = 0.0076 for ETH; p = 0.021 for SAF). In contrast, at the adult (female) stage, the

proportions of outliers located on the X tended to be higher than the background level (permutation

test, p = 0.30 for MED; p = 0.013 for ETH and p = 0.89 for SAF). For pupa, the X chromosome

enrichment was not significantly different between outliers and the background. Considering the

three population pairs together, the relative enrichments of outliers on the X chromosome were

significantly different among developmental stages (likelihood ratio test, p = 0.048). The patterns of

different enrichment among stages were caused by larva/adult differences (likelihood ratio test, p =

0.0058 for larva-adult; p = 0.25 for pupa-adult; p = 0.24 for larva-pupa). The larva/adult difference

might suggest either (1) a greater abundance of genes affecting adult fitness on the X chromosome,

as previously suggested (Gibson et al. 2001; Innocenti and Morrow 2010), (2) a greater influence of

female fitness on X chromosome evolution (e.g. Vicoso & Charlesworth 2006), in light of sex

differences between our adult (female) and larval (mixed sex) samples, or (3) differences in the

contributions of local adaptation and genetic drift to outlier sets at different stages, in combination

with differential effects of drift between the X chromosome and autosomes (e.g. Pool & Nielsen

2007).

Fig 3. The fraction of P_{ST} outliers located on the X chromosome varies between developmental stages.
The dashed error bar indicates the 95% confidence interval for the permuted data. If the real data
(orange bar) is outside the range of the error bar, it indicates the fraction is significantly different
from the genomic background (p < 0.05).
592

Co-directional evolution in gene expression between population pairs

Among the upper 5% of P_{ST} outliers, we found at least some significant signals of parallel expression

divergence in all three pairwise comparisons (MED vs. ETH; MED vs. SAF; ETH vs. SAF), where

the shared outliers with co-directional changes (i.e., expression difference between cold- and warm-

derived populations in the same direction for both pairs) were more abundant than expected by

random permutation (Table 1; Fig 4A). Averaged across three pairwise comparisons, 6.7% of the

outliers were shared and changed consistently for the adult stage while 4.7% of outliers were co-

directional for larva and only 2.6% were co-directional for pupa (for which one population pair

comparison for pupa showed significant anti-directional changes). Changing the P_{ST} outlier cutoffs to

2.5% or 10% produced qualitatively similar patterns (Table S7). We found one shared outlier with

co-directional changes among all three pairs for the adult stage: *Iml1*, a regulator of cell size,

starvation response, TOR signalling, and meiosis initiation (Bar-Peled et al. 2013; Wei et al. 2014).

No three-pair shared outliers were found for the larva and pupa stages. One shared outlier among

three pairs is not significantly more than the expectation by permutation. Still, it is worth noting that

changing the P_{ST} outlier cutoff to 10% results in 12 shared outliers with co-directional changes

among three pairs (permutation test, p < 0.0001), which suggests that meaningful regulatory

evolution may extend beyond our defined outliers. The names, P_{ST} values and quantiles of these 12

genes are shown in Table 2. Of these genes, we note the role of *CrebB* in circadian behavior, which

is known to differ between *D. melanogaster* populations from different thermal environments (Svetec

et al. 2015; Cao & Edery 2017; Helfrich-Förster et al. 2018). Overall, these analyses suggest that the

adult stage has the highest level of parallel evolution while pupa has the lowest.

stage	larva		pupa		adult	
Population pairs	MED	SAF	MED	SAF	MED	SAF
ETH	3.4% (3.0%)	3.0% (3.4%)	**5.5%** (2.7%)	2.0% (2.0%)	3.0% (2.4%)	**10.1%** (2.9%)
SAF	**7.7%** (3.0%)		**0.4%** (2.0%)		**5.3%** (2.4%)	

Table 1. Evidence for parallel expression evolution between population pairs. The percentages of P_{ST}
outlier with parallel expression abundance changes are shown. The random expectation is the median
of the permuted proportions (in brackets). The majority of proportions were higher than the
expectation, with larvae and adult stages showing stronger patterns than the pupa. Those that were
significantly different from the random expectation are in bold (permutation test, p < 0.01). We
found one case, the MED-SAF comparison at pupal stage, that showed a significantly lower level of
parallel evolution than the random expectation. Further detail regarding the numbers of shared and
non-shared outliers can be found in Figure 4A.

Pair	MED		ETH		SAF	
Gene name	P_{ST} value	quantile	P_{ST} value	quantile	P_{ST} value	quantile
Iml1	0.58	0.0177	0.85	0.0055	0.79	0.0001
larp	0.42	0.0532	0.70	0.0401	0.51	0.0074
Smox	0.49	0.0345	0.66	0.0607	0.49	0.0087
sky	0.38	0.0694	0.68	0.0529	0.47	0.0102
AGO1	0.36	0.0818	0.70	0.0423	0.45	0.0134
CG10365	0.53	0.0251	0.67	0.0553	0.39	0.0195
Nepl3	0.35	0.0840	0.64	0.0744	0.34	0.0270
CG42674	0.34	0.0948	0.84	0.0057	0.29	0.0383
CrebB	0.44	0.0461	0.74	0.0268	0.24	0.0566
Cka	0.56	0.0202	0.75	0.0249	0.23	0.0606
par-1	0.34	0.0883	0.70	0.0396	0.22	0.0713
CG5116	0.45	0.0432	0.78	0.0171	0.20	0.0796

Table 2. The P_{ST} values and quantiles for the 12 genes that passed the top 10% cutoff with consistent
expression changes across three population pairs at the adult stage.

The analysis above requires genes being outliers in both population pairs, which is quite restrictive

(the naïve expected proportion of sharing between two pairs under a 5% P_{ST} cutoff is 5%×5%×0.5 =

0.125%) and may miss some broader patterns of parallel changes. We therefore performed a

complementary analysis which only required genes being outliers in one population pair and

examined whether the expression for this set of genes changed in the same direction in another pair,

regardless of outlier status in the latter pair. For example, 235 genes were outliers in the MED pair at

the larval stage. Then in another pair, we calculated the fraction of the 235 genes with expression

differences between cold- and warm-derived populations in the same direction as for MED. There

were excesses of co-directional changes for the larval stage (Figure 4B). The patterns were weaker

for the adult stage and there were excesses of anti-directional changes for the pupal stage. Changing

the P_{ST} cutoff to 2.5% or 10% produce qualitatively similar patterns (Fig. S6).

Fig. 4. Extent and direction of parallel evolution in gene expression among population pairs. (A) Venn diagrams for the numbers of shared outliers with co-directional changes between pairs and the rest of genes. (B) Fractions of co-directional gene expression changes between pairs for the P_{ST} outliers identified in one pair of them. The name above is the pair used to identify outliers. The name below is the other pair in the comparison. The outliers used for each bar is the summed number of the pair in the Venn diagram. The dashed error bar indicates the 95% confidence interval for the fraction of co-directional expression changes in permuted data. If the real data is outside the range of the error bar, it indicates the fraction is significantly different from random expectation ($p < 0.05$, two-sided test based on permuted distribution).

We also performed similar analyses for P_{ST} outliers of alternative intron usage. The numbers of intron-excision junctions that passed the cutoffs for P_{ST} calculation were 4520 for larva, 5574 for pupa, and 7367 for adult. The adjusted gene quantiles for splicing are listed in Table S5. The patterns of co-directional changes were qualitatively similar to those for gene expression (Fig. S7). The fractions of co-directional changes were still highest for the larvae among the three stages; all of the comparisons except one showed an excess of co-directional changes relative to the background level, although only one comparison is significant based on the permutation test (outliers from MED being co-directionally expressed in SAF). Overall, the patterns for co-directional changes were weaker for splicing than those for gene expression. Table S9 lists the genes with the top 20 P_{ST} for both expression and intron usage for each population pair. An interesting example is *curled*, which is an extreme splicing outlier for ETH and MED, and an extreme expression outlier for ETH as well. Also known as *nocturnin*, one isoform of this gene is thought to have a dedicated role in circadian regulation (Nagoshi et al. 2010).

Enriched functional categories for the P_{ST} outliers

Significant Gene Ontology (GO) terms enriched in different sets of P_{ST} outliers for gene expression are listed in Table S10. Among the significant GO terms for different population pairs, we found five terms shared between the MED pair (24 significant terms) and ETH pair (47 significant terms) at the adult stage (mitochondrion, nucleoside metabolic process, ribonucleoside metabolic process, purine nucleoside metabolic process and oxidation-reduction process). The level of sharing was significantly more than expected by chance based on permuted outlier sets ($p < 0.001$, no shared GO terms were found in the permuted datasets), suggesting functional convergence for adult development to the cold environment for the MED pair and ETH pair. Further, similar (though non-identical) GO terms were identified from different pairs at different stages such as terms related to mitochondria, nucleoside metabolism, and oxidoreductase complex. However, the majority of GO

terms were unique for different pairs, suggesting that many functional changes for adaptation to cold

environments may be population-specific. For intron usage, we only found one significant GO term

for SAF pair at the larval stage (meiotic cell cycle).

Cis- and trans-acting contributions to differential gene expression at the adult stage

Gene expression differentiation can be caused by *cis-* or *trans*-regulatory effects. A *cis*-effect comes

from a local regulatory mutation and results in an allele-specific expression difference in a F1 hybrid;

while a *trans*-effect is caused by remote loci that modify both alleles in a diploid cell. Therefore, a

cis-effect can be estimated by the allelic expression in F1 hybrid (Cowles et al. 2002). We quantify

that effect based on the expression proportion in the F1 offspring of a between-population cross for

the allele from the cold population minus 0.5 (null expectation when *cis*-effect absent). A *trans*-effect

can be estimated by the expression difference between parents that was not attributed to the *cis*-effect

(Wittkopp et al. 2004), as described in the Materials and Methods.

First, we described the transcriptome-wide patterns of *cis-* and *trans*-effect sizes across all analyzed

genes at the adult stage. The magnitudes of *trans*-effect sizes were significantly larger than the *cis*-

effect sizes in all three population pairs (mean absolute *cis*-effects and *trans*-effects were: MED pair,

0.07 vs. 0.16, $p < 2.2e-16$; ETH pair, 0.07 vs. 0.16, $p < 2.2e-16$; SAF pair, 0.09 vs. 0.11, $p < 2.2e-16$.

'Mann-Whitney' paired test.). Moreover, we found strong negative relationships between *cis-* and

trans-effects within each population pair (Fig. S7), where the *cis-* and *trans*-effects were generally in

the opposite directions. Although the pattern can be biologically meaningful, it may also represent an

artifact from using the same F1 expression data for allele specific expression (ASE) estimation to

infer both *cis-* and *trans*-effects. Any measurement error on ASE will introduce an artifactual

negative correlation between *cis-* and *trans*-acting changes (see Discussion below).

Next, we used our flexible permutation approach (see Materials and Methods and supplementary

document) to study how many genes show a significant *cis*-effect, *trans*-effect, or both among the

three population pairs (Table 3). Fig. 5A shows a graphical depiction of *cis-* and *trans*-effects on F1

and parental samples, illustrating *cis*-effects influence both F1 allele-specific expression and the

parental expression ratio while *trans*-effects only influence the parental expression ratio. Averaged

across population pairs, about 60% of genes show significant *cis-* and/or *trans*-effects (62% for

outliers and 59% for non-outliers). We also found that 19% of genes show *cis*-regulatory effects

while 53% of them show *trans*-effects, consistent with *trans*-effects being stronger on average than

cis-effects. This apparently greater prevalence of *trans*-regulatory evolution was observed in spite of

our lesser power to detect *trans*- relative to *cis*-effects (Supplementary Text).

pair	expression type	Total tests	*Cis* only	*Trans* only	Both co-dir	Both anti-dir	Neither
MED	P_{ST} outliers	184	20 (11%)	74 (40%)	6 (3%)	14 (8%)	70 (38%)
	Non-outliers	4182	199 (5%)	2203 (53%)	99 (2%)	324 (8%)	1357 (32%)
ETH	P_{ST} outliers	231	12 (5%)	136 (59%)	10 (4%)	14 (6%)	59 (26%)
	Non-outliers	4569	297 (7%)	2224 (49%)	137 (3%)	379 (8%)	1532 (34%)
SAF	P_{ST} outliers	167	22 (13%)	47 (28%)	0 (0%)	15 (9%)	83 (50%)
	Non-outliers	3993	250 (6%)	691 (17%)	40 (1%)	738 (18%)	2274 (57%)

711   Table 3. The relative prevalence of significant *cis*- and *trans*-regulatory differences for P_{ST} outliers
712   versus non-outliers varies among population pairs. Numbers of gene expression abundance traits
713   showing different regulatory effects for P_{ST} outliers and non-outliers are shown. The percentage in
714   parentheses indicates the fraction of genes in each category relative to total genes in the tests.
715

Fig 5. Population pairs show distinct patterns of *cis*- and *trans*-regulatory evolution for expression
outliers. These plots depict evidence for *cis*- and *trans*-regulatory evolution based on the relative
expression proportion of cold strain alleles in parental and F1 datasets. (A) Conceptual depiction of
cis- and *trans*-effects. *Cis*-effects change the F1 allele-specific expression and the parental expression
proportion concordantly (along the one-to-one ratio line). *Trans*-effects only change the parental
expression proportion but not the F1 allele-specific expression (vertical line y = 0.5). Co-directional
cis- and *trans*-effects (*CisTransCodir*) locate in the two spaces within the 45° angle between the *Cis*
vector and the *Trans* vector. Anti-directional *cis*- and *trans*-effects (*CisTransAntidir*) locate in the
two spaces within 135° angle between *Cis* vector and *Trans* vector. (B-D) Evidence for *cis*- and
trans-regulatory evolution of putatively adaptive expression differences between warm- and cold-
derived populations (expression P_{ST} outliers).

To examine the *cis*- and *trans*-regulatory contributions to adaptive evolution of gene expression, we

compared genes showing *cis*- or *trans*-effects between P_{ST} outliers (Fig. 5B-D) and non-outliers.

Because of the potential artifact generating opposing *cis*- and *trans*-effects, we excluded genes

showing both significant *cis*- and *trans*-effects in opposite directions (Both anti-dir category in Table

3). For genes showing significant *cis*-effects (Fig. 6A), they were enriched in the outliers relative to

the non-outliers for the MED pair ($\chi2 = 11.6$, df = 1, p = 0.00066) and the SAF pair ($\chi2 = 4.8$, df = 1,

p = 0.029) but not for the ETH pair ($\chi2 < 0.01$, df = 1, p = 1). While for significant *trans*-effects

genes (Fig. 6B), they were enriched in the outliers relative to the non-outliers for the ETH pair ($\chi2 =$

9.6, df = 1, p = 0.0019) and the SAF pair ($\chi2 = 5.4$, df = 1, p = 0.020) but the enrichment is opposite

for the MED pair ($\chi2 = 10.2$, df = 1, p = 0.0014).

Because theory suggests that co-regulation of genes can amplify the contribution of *trans*-effects (Liu

et al. 2019), we examined the level of co-regulation between pairwise outliers by calculating the

correlation coefficient between the expression values among the eight outbred samples within each

cold-derived population. Indeed, the percentage of pairwise correlation with evidence of co-

regulation (p-value < 0.05) is much higher in EF population than that in FR and SD populations (Fig.

6C, Table S11). At the gene level, the median number of significant co-regulatory partners for EF is

about twofold that for FR and SD (Fig. 6D). This pattern supports the hypothesis that substantial co-

regulation of outlier genes results in more significant *trans*-effects for the ETH pair (Fig. 6B).

Further, the strong co-regulation in EF might be related to the asymmetry observed in the ETH

outliers (Fig. 2). However, the pattern could reflect variation in cell type content among strains rather

than co-regulation within cells (see Discussion).

751
Fig 6. The prevalence of *cis*- versus *trans*-regulatory P_{ST} outliers differs between population pairs,
while trans changes show greater parallelism. The upper panels show the fractions of genes with *cis*
regulatory effects (A) and those with *trans*-effects (B) for P_{ST} outliers and non-outliers in each
population pair. The middle panels show the proportion of co-regulated pairs among all pairwise
outliers (C) and the numbers of co-regulatory partners for outlier genes (D) in each cold-derived

population. The lower panels show the fractions of parallel outlier genes (*i.e.* shared and

codirectional between population pairs) and non-paralleled outlier genes (*i.e.* shared and anti-

directional between population pairs) showing *cis*-effect (E) and *trans*-effect (F) in different

population pairs. The number above the bar for the *cis*-effect shows the denominator of the fraction,

which applies to the *trans*-effect results as well. * indicates the fractions are significantly different

between two categories ($p < 0.05$).

To study the role of *cis*- and *trans*-regulatory effects on the parallel adaptation, we further focused on

the outlier genes. We categorized the top 10% P_{ST} outliers based on whether they were shared

between two pairs with consistent expression changes (parallel) or opposite changes (non-parallel).

Because genes with *cis*- (or *trans*-) effects in both population pairs in the parallel category would

indicate the contribution of *cis*- (or *trans*-) effects to parallel adaptive evolution, we identified shared

outlier genes with *cis*- (or *trans*-) effects in any two pairs in both parallel and non-parallel categories

(excluding genes showing both significant *cis*- and *trans*-effects in opposite directions). For *cis*-

effects, only one gene (*CG42788*) is significant in both ETH and SAF pairs (Fig. 6E). While for

trans-effects, the numbers of genes showing *trans*-effect are four for the MED and ETH pairs

(*CG8034*, *CG33981*, *kst* and *wdb*), two for the MED and SAF pairs (*scf* and *AP-2σ*) and four for the

ETH and SAF pairs (*CG7766*, *Dlg5*, *larp* and *Pink1*). Of these, *larp* had P_{ST} quantiles near or below

0.05 in all three population pairs (Table 2); its functions include mitochondrial regulation (Zhang et

al. 2019). There are significant enrichments of *trans*-effect genes in the parallel outlier category

relative to the non-parallel category in two out of three population pair comparisons (Fig. 6F, MED

& ETH: $p = 0.0097$; MED & SAF: $p = 0.014$; ETH & SAF: $p = 1$; Fisher's Exact Tests). As a

complementary analysis, we also included the non-shared outliers in the non-parallel category, and

we found qualitatively similar patterns for *cis*- and *trans*-effects (Fig. S9). Overall, there is stronger

evidence of *trans*-regulatory evolution contributing to parallel gene expression changes between

cold-adapted populations.

Cis- and trans-acting contributions to differential intron usage at the adult stage

For all intron usage traits, we found the magnitude of *trans*-effects on average to be higher than that

of *cis*-effects (mean absolute *cis*-effects and *trans*-effects are: MED pair, 0.13 vs. 0.17, $p = 0.0001$;

ETH pair, 0.30 vs. 0.32, $p = 0.04$; SAF pair, 0.17 vs. 0.19, $p = 0.0044$. 'Mann-Whitney' paired test).

Because of the limited diagnostic SNPs with enough read depth located in the intron junction regions,

there are few outlier introns tested for *cis*- and *trans*-regulatory effects (Table S12).

Elevated genetic differentiation at cis-regulated expression outliers

Since the *cis*-regulatory mutations contributing to local adaptation may show differentiation in allele

frequency between populations, we examined genetic differentiation for expression outliers with *cis*-

effects (including *cis*-only genes and genes with both *cis*- and *trans*-effects to increase sample sizes).

We examined whether each of these *cis*-outliers shows high F_{ST} between that pair of cold- and warm-

adapted populations – for both window F_{ST} (F_{ST_winmax}) and maximum SNP F_{ST} (F_{ST_SNPmax}). A gene

showing both significant *cis*-effect and higher F_{ST} quantile (here, the top 5% versus comparable

genomic regions) could reflect adaptive regulatory evolution targeting the surveyed sequences or

nearby sites. We first confirmed that there is little chromosomal scale differentiation between

populations (aside from known moderate X-autosome differences for the MED pair; Lack et al. 2015)

by plotting window F_{ST} across the major chromosome arms (Fig. S8). We observed no obvious

clustering of P_{ST} outliers along the chromosome arms. For window F_{ST} (Fig. 7A), high F_{ST} is

enriched in *cis*-effect outliers relative to the non-outliers for the MED pair (p = 0.022. Fisher's Exact

Test) and the other two pairs show the same trend (ETH: p = 0.22; SAF: p = 0.23). Although the

MED result would be only marginally significant if correcting for the three tests performed, a clearer

result is obtained when considering the three population pairs together. In this analysis, the fractions

of genes with high window F_{ST} were significantly higher in *cis*-outliers than that in non-outliers

(likelihood ratio test, p = 0.0056). For maximum SNP F_{ST} (Fig. 7B), the MED and SAF pairs showed

significant enrichments of high F_{ST} in *cis*-effect outliers versus the non-outliers, while the ETH pair

showed a weak trend in that direction (MED: p = 0.012; ETH: p = 0.49; SAF: p = 0.030). Similar

analysis combining the three pairs found that the fractions of high maximum SNP F_{ST} were

significantly higher in *cis*-outliers than that in non-outliers (likelihood ratio test, p = 0.0045). These

results reflected more than two-fold enrichment of F_{ST} outliers among *cis*-regulated P_{ST} outliers

across population pairs.

815

 Fig. 7. Enrichment of *cis-outliers* for genes with high window F_{ST} (A) and high SNP F_{ST} (B) for each
population pair. The number above the bar shows the denominator of the fraction. * indicates the
fractions are significantly different between *cis*-effect outliers and non-outliers (p < 0.05, Fisher's
Exact Test).

We further examined potential targets of *cis*-regulatory local adaptation, excluding genes with *cis*-

effects and trans-effects in the opposite directions. For the *cis*-genes of the MED pair, *Ciao1* and

Cyp6a22 showed high window F_{ST}, *CG42565* showed high F_{ST_SNPmax}, and *CG8034* and *Cyp6a17*

showed both. *CG8034* (a predicted monocarboxylic acid transporter) was also cited above as having

parallel *trans*-regulatory evolution between the MED and ETH pairs, and this gene's *cis-* and *trans-*

effects were both upregulated in FR relative to EG in MED. *CG42565* represents a differentially-

spliced product of a transcript that alternatively yields an isoform of *CG13510*, which is upregulated

in response to cold (Qin et al. 2005). Interestingly, *Cyp6a22* and *Cyp6a17* belong to the cytochrome

P450 protein family (*Cyp6a22* is 248 bp upstream of *Cyp6a17*). This region harbors a polymorphic

deletion of *Cyp6a17* which is associated with both colder temperature preference (Kang et al. 2011;

Chakraborty et al. 2018) and lower insecticide resistance (Duneau et al. 2018). Based on the

diagnostic SNPs for *Cyp6a17* (Good et al. 2014), we found the France-enriched allele for *Cyp6a22* is

likely linked to the *Cyp6a17* deletion. Likewise, at the population level, the frequency of intact

Cyp6a17 copy is 0.44 in France and 0.95 in Egypt. Hence, adaptive expression differences may

sometimes be driven by gene copy number differentiation between populations. For the *cis*-genes of

ETH, *RpL24* showed high window F_{ST} and *KrT95D* showed high F_{ST_SNPmax}. For SAF, *GXIVsPLA2*

showed high F_{ST_SNPmax}, while *AGO2* showed both high window F_{ST} and high F_{ST_SNPmax}. *AGO2* is

involved with antiviral defense and developmental regulation (Deshpande et al. 2005; Nayak et al.

2010) and was previously found to contain fixed differences between European and African

populations (Pool 2015). For the genes showing high F_{ST_SNPmax} in any pair, we plotted the SNP F_{ST}

along the gene region and nearby 20kb to show the sites that may be the most likely targets of

selection (Fig. 8). For *CG42565, CG8034* and *KrT95D*, we observed that the highest F_{ST} sites were

located within the gene regions. While for *GXIVsPLA2*, the highest SNP F_{ST} was 1758 bp

downstream of the gene. For *AGO2*, the highest SNP F_{ST} was 6359 bp downstream but the third

highest SNP F_{ST} was within the gene. Overall, the genetic differentiations between cold- and warm-

derived populations around these candidate genes can be quite local, but the linked signal of natural

selection can extend further, and there are often multiple SNPs that could represent plausible targets

of local adaptation within and outside a given gene region.

Fig 8. Local peaks of F_{ST} between warm- and cold-derived populations are observed at some
candidate genes for adaptive cis-regulatory evolution. SNP F_{ST} along the candidate genes with
flanking regions of 20kb is shown. The top diagram depicts the coding (orange) and non-coding
(gray) exon, captured from GBrowse 2 of *D. melanogaster* (R5.57) from FlyBase (St. Pierre et al.
2014). *Cyp6a17* is not plotted because of the known gene-scale polymorphic deletion.
857

Discussion

Parallel evolution has often been studied at the population genetic and trait levels, but it has less

frequently been analyzed at the transcriptome level (Stern 2013; Juneja et al. 2016). In this study, we

used three recent instances of adaptation to colder climates in *Drosophila melanogaster* to study the

evolution of gene expression and alternative splicing. We found a unique pattern of transcriptomic

evolution in the high altitude EF sample, involving elevated expression of highly expressed muscle

proteins and many other genes. We found the locations of differentially expressed genes on the X

chromosome versus the autosomes varies among developmental stages, with the adult female stage

having relatively more differentially expressed X-linked genes than the larval stage. We then saw

signals of parallel evolution in expression that were higher for larval and adult stages than for pupa.

Further, we studied *cis-* and *trans-*regulatory evolution in the context of this ecological adaptation,

finding that the relative roles of these regulatory mechanisms differ strongly among population pairs.

And we found a signal of *trans-*regulation contributing more predictably to the parallel evolution

between population pairs. Finally, those outliers showing *cis-*effects were enriched for high genetic

differentiation between populations, suggesting that some of them were the direct targets of selection

in cold environments.

Previous comparative transcriptomic studies on multiple pairs of *Drosophila* populations/species

have found parallel gene expression differences between high and low latitude populations (Zhao et

al. 2015; Juneja et al. 2016). However, because both clines in Australia and North America came

from admixtures between European and African ancestry and tropical populations in both clines have

a greater proportion of African ancestry (Bergland et al. 2016), it is hard to disentangle adaptive

divergence between high and low latitude populations from demographic effects. The demographic

influences may be smaller when comparing clines between species pairs (Zhao et al. 2015).

Interestingly, the latter study found 10% to 20% differential expressed genes between high and low

latitude populations were shared and changed co-directionally between two species *D. melanogaster*

and *D. simulans*. The percentages are even higher than what we found between population pairs at

adult stages (3% to 10%).

One important reason for the milder patterns of parallelism among our population pairs may be the

different selection agents in their unique habitats. In Zhao et al. 2015, samples from the high and low

latitude populations from both species were collected in the same areas. While for us, the cold-

derived FR population from the MED pair colonized a higher latitude environment than the related

warm population, whereas the other two cold-derived populations colonized higher altitude

environments where the selection agents may include air pressure, desiccation and ultraviolet

radiation (Pool et al. 2017). Although EF and SD have both adapted to higher altitudes (EF at 3,070

meters above sea level, SD at 2,000), SD is seasonally cold (like FR) whereas EF is perpetually cool.

We should also note that the standing variation available for adaptation may have differed between

our cold-derived populations due to their distinct demographic histories, including the trans-Saharan

bottleneck affecting the MED pair and a milder bottleneck in the history of the ETH pair, but no

meaningful bottleneck involving the SAF pair (Sprengelmeyer et al. 2020). Hence, although we do

significant evidence of parallel gene regulatory adaptation, we suggest that there are both ecological

and population genetic reasons to expect substantial non-parallel adaptation as well.

Notably, the EF population exhibits distinct phenotypic evolution such as darker pigmentation

(Bastide et al. 2014), larger body size (Pitchers et al. 2013; Lack et al. 2016b), and reduced

reproductive rate (Lack et al. 2016b). This distinct evolution in the EF population may explain the

strong directionality in gene expression changes between EF and EA for the outliers (Fig. 2), the

greater abundance of *trans*-regulatory outliers, and the elevated levels of co-regulation among

outliers (Fig. 6). Therefore, the underlying transcriptomic evolution for EF may partly reflect its

unique phenotypic evolution, not just adaptation to lower temperature. For example, the upregulation

of muscle proteins could reflect the differential abundance of tissues between these Ethiopian

populations that differ in size. This size evolution may have also altered the relative proportions of

different cell types, which may have driven some of the population differences in gene regulation

observed from our whole-organism samples. Future tissue-specific or cell type-specific expression

studies involving the EF population can help to examine these possibilities.

We found some evidence of parallel expression evolution between our cold-adapted populations.

Developmental stage has a strong effect on the levels of this parallelism, with adult and larva

showing significant parallelism while pupa showed a much weaker pattern (Table 1; Figure 4). This

is consistent with the observations that larval and adult stages show local adaptation to native

temperature but not the pupal stage (Austin and Moehring 2020). It is possible that pupal

metamorphosis might reflect a relatively constrained developmental program that limits opportunities

for thermal adaptation. The high level of detected parallelism in larvae could also reflect a higher

detection power due to less tissue diversity and hence broader spatiotemporal expression of relevant

differences. The intriguing pattern of anti-parallelism for some combinations of the pupal stages

might suggest that other selective agent is more important than cold (e.g., oxygen level, ultraviolet

radiation) for certain population pairs and the direction of selection on gene expression is opposite to

the cold. Further, the anti-directional pattern at the pupal stage could be caused by different rates of

development for cold-derived populations relative to the warm-derived ones in different pairs.

Evidence for such differences is mixed: rates were found to differ between high and low latitude

populations in Australia (James & Partridge 1995) but not between our EF and ZI populations (Lack

et al. 2016b). Because tissues at different days can generate a wide range of gene expression

difference (Hsu et al. 2019), if the cold-derived population develops faster than the warm-derived

one in one pair but in another pair the cold-derived population develops slower than the respective

warm-derived one, many of the expression differences will be anti-directional between the two pairs.

Moreover, because the pupal and larval samples were mixed sex, different rates of development for

males and females could led to a biased sex ratio in a sample, especially for pupae (Testa et al. 2013).

If the sex ratio bias happened in the cold-derived population in one pair but the warm-derived

population in another, it could conceivably result in anti-directional patterns for sex-biased genes.

Compared to the expression abundance, the pattern of parallelism is much weaker for intron usage

(Fig S7), which may partly stem from lower power to detect intron usage change (only a small

proportion of reads are informative for exon junctions). However, we still found the MED pair and

SAF pair show more parallel changes than the combinations with the ETH pair, which is consistent

with results for expression abundance. Given the increasing evidence for alternative splicing

contributing to environmental response and adaptation (e.g., Singh et al. 2017; Signor and Nuzhdin

2018; Smith et al. 2018), we need to study both expression abundance and splicing to fully

understand the evolution at the transcriptome level. The development of sequencing approaches with

long reads that cover the entire transcripts (e.g., Iso-Seq) will enable us to quantify isoforms

frequency directly and broaden the scope of alternative splicing variation that can readily be

quantified. Since splicing changes during development and among tissues (Brown et al. 2014;

Gibilisco et al. 2016), a detailed sampling throughout development of different tissues will also be

necessary to understand the role of splicing on ecological adaptation.

We found *trans*-effects on expression were more common than the *cis*-effects across the

transcriptome (Table 3), which is consistent with some previous studies (e.g., McManus et al. 2010;

Coolon et al. 2014; Albert et al. 2018; Glaser-Shmitt et al. 2018) but not with others (e.g., Lemmon

et al. 2014; Mack et al. 2016). The transcriptome-wide prevalence of *trans*-effects may be caused by

random regulatory changes biased toward *trans*-regulation because of the larger trans-mutational

target size (Landry et al. 2007; Metzger et al. 2016). Or, *trans*-regulatory changes may have higher

potential for coordinate regulation of multiple genes in networks (Metzger et al. 2016; Liu et al.

2019). To focus on the evolved changes potentially related to adaptation, we compared the

proportion of genes with *cis-/trans*-effects for P_{ST} outliers and to those for non-outliers and saw both

effects were enriched in outliers in certain pairs (Fig. 6A & B). These results indicate that the

mechanisms of adaptive gene regulatory evolution are highly population-specific, and that either

regulatory mechanism has the potential to play a disproportionate role in ecological adaptation.

Moreover, we found a predominance of *trans*-effects associated with parallel outliers than the non-

parallel outliers (Fig. 6E & F). In part because of the larger mutational target size of *trans*-regulatory

variation for a given target gene, the standing genetic variation for *trans*-regulatory variants may be

higher than the *cis*-ones in the ancestral population and therefore the *trans* variants can respond to

selection in different population pairs. However, studies in *Arabidopsis thaliana* and *Capsella*

grandiflora find that *trans*-eQTLs tend to have lower minor allele frequencies than *cis*-eQTLs

(Zhang et al. 2011; Josephs et al. 2020) but it is unclear whether these populations represent the

ancestral state before experiencing environmental changes. Also, the potential capacity of *trans*-

regulatory factors to co-regulate many genes may amplify the probability of parallel changes between

population pairs. Furthermore, since we used whole-body adult samples, it is possible that some

trans-acting factors regulated genes similarly across tissues while the some *cis*-effects were tissue-

specific and were undetected in our mixed-tissue samples. Finally, we emphasize that our study

focuses on regulatory changes that may have relatively larger effects (in focusing on P_{ST} outliers, and

in basing *cis/trans* analysis on strains showing clearer differences); but small changes may be

important for regulatory evolution as well, and may be differentially represented between categories

(e.g. *cis* vs. *trans*, parallel vs. non-parallel).

When we considered genes/introns showing both *cis-* and *trans-*effects, we observed that the two

types of effects were generally in opposite directions (anti-directional; Table 3). This is consistent

with the idea that gene regulation is under stabilizing selection in general and gene regulatory

networks evolve negative feedback to buffer effects of regulatory changes (Denby et al. 2012;

Coolon et al. 2014; Bader et al. 2015; Fear et al. 2016). With regard to our P_{ST} outliers, it is possible

that *cis-*acting changes might have evolved to compensate for unfavorable pleiotropic impacts of

adaptive *trans-*regulatory evolution (or possibly vice-versa). However, negative correlations between

cis- and *trans-*effects can also be an artifact coming from the measurement error on F1 expression

data. Because the F1 data was used to estimate ASE and compared it to 0.5 (*cis-*effect null) and to

parental expression proportion (*trans-*effect null), measurement error will introduce artifactual

negative correlation between *cis-* and *trans-*acting changes. Therefore, whether the opposing effects

between *cis-* and *trans-*acting changes are biologically meaningful will require further study. As

Fraser (2019) and Zhang and Emerson (2019) proposed, using independent F1 replicates or other

approaches such as eQTL mapping to infer *cis-* and *trans-*effects separately is necessary to affirm

evidence of compensatory evolution.

We expect that the adaptive expression divergence caused by *cis-*regulatory changes should leave a

signal in the genetic variation of the nearby genomic region. Therefore, we used F_{ST} statistics to

quantify genetic differentiation for the region around the focal genes. Window F_{ST} is sensitive to

classic hard sweeps, and relatively useful for incomplete sweeps and moderately soft sweeps, but it is

less useful for soft sweeps with higher initial frequencies of the beneficial allele (Lange and Pool

2016), for which SNP F_{ST} may be more sensitive. Indeed, a previous genomic study on these same

populations found a stronger signal of parallel change for SNP F_{ST} than for window F_{ST} genome-wide

(Pool et al. 2017). Here, we found genes with outlier *cis-*effects are enriched for those that show high

F_{ST}, especially those with high SNP F_{ST} (Fig. 7A & B). Hence, standing genetic variants may have

contributed importantly to the *cis-*regulatory changes for adaptation in our populations. Genes with

both significant *cis-*effects and high F_{ST} are likely to be the direct targets of the environmental

selection and good candidate for future mechanistic studies.

1012

Using three natural fly population pairs with recent adaptive divergence, our study found intriguing

patterns of parallel evolution in gene expression and provided new insights on the underlying

regulatory effect. In the future, using other approaches to study *cis-* and *trans-*effects in these

populations would be necessary, such as eQTL mapping, which can provide more genetic

information about the *trans*-regulatory loci. Also, studying the gene expression in different tissues,

including different organs from males and females, would provide us a clearer and more

comprehensive picture about parallel gene expression evolution. It would also be informative to

study other phenotypes besides gene expression that are more related to thermotolerance such as the

metabolic pathways identified in this study or nervous system function related to chill coma.

Moreover, studying the phenotypic plasticity at different developmental stages could help to explain

the different patterns of parallelism in expression evolution across stages and allow us to better

understand the importance of local adaptation versus plasticity in thermotolerance.

1025

1026

1027

Data Availability

1029 The raw RNAseq reads are available from the Sequence Read Archive (SRA) under accessions
1030 SRR14179998-SRR14180176 and BioProject PRJNA720479. The population genomics data is
1031 from the *Drosophila* Genome Nexus (Lack et al. 2016; http://www.johnpool.net/genomes.html).
1032 Custom R and Perl scripts for the *cis*- and trans- effects simulation and analysis can be found at
1033 https://github.com/YuhengHuang87/simulation_cis_trans.
1034

1035

Acknowledgements
1037 We thank Colin Dewey for helpful discussions and the UW-Madison Center for High Throughput
1038 Computing (CHTC) for cluster usage. We thank Jeremy Lange for suggestions on *Cyp6a17* gene
1039 analysis. This work was funded by NSF DEB grant 1754745 to JEP and by NIH NIGMS grant
1040 F32GM106594 to JBL.
1041

1042

References
Adrion JR , Hahn MW , Cooper BS. 2015. Revisiting classic clines in *Drosophila melanogaster*
in the age of genomics. Trends Genet 8:434–444.

Alasoo K , Rodrigues J , Danesh J , Freitag DF , Paul DS , et al. 2019. Genetic effects on
promoter usage are highly context-specific and contribute to complex traits. Elife. 8.
doi:10.7554/eLife.41673

Albert FW, Bloom JS, Siegel J, Day L, Kruglyak L. 2017. Genetics of trans-regulatory variation
in gene expression. eLife 100:371–44.

Arthaud L, Ben Rokia-Mille S, Raad H, Dombrovsky A, Prevost N, et al. 2011. Trade-off
between toxicity and signal detection orchestrated by frequency- and density-dependent
genes. PLoS One 6: e19805.

Austin CJ, Moehring AJ. 2019. Local thermal adaptation detected during multiple life stages
across populations of Drosophila melanogaster. J. Evol. Biol. 32, 1342–1351.

Bader DM, Wilkening S, Lin SG, Tekkedil MM, Dietrich K, et al. 2015. Negative feedback
buffers effects of regulatory variants. Mol Syst Biol 11:785.

Barbosa-Morais NL, Irimia M, Pan Q, Xiong HY, Gueroussov S, et al. 2012. The evolutionary
landscape of alternative splicing in vertebrate species. Science 338:1587–93.

Bastide H, Yassin A, Johanning EJ, Pool JE. 2014. Pigmentation in *Drosophila melanogaster*
reaches its maximum in Ethiopia and correlates most strongly with ultra-violet radiation in
sub-Saharan Africa. BMC Evol Biol 14:222–14.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate—a practical and powerful
approach to multiple testing. J Royal Stat Soc B 57:289–300.

Bergland AO, Tobler R, González J, Schmidt P, Petrov D. 2016. Secondary contact and local
adaptation contribute to genome-wide patterns of clinal variation in *Drosophila*
melanogaster. Mol Ecol 25:1157–1174.

Božičević V, Hutter S, Stephan W, Wollstein A. 2016. Population genetic evidence for cold
adaptation in European *Drosophila melanogaster* populations. Mol Ecol 25:1175–1191.

Brommer JE. 2011. Whither P_{ST}? The approximation of Q_{ST} by P_{ST} in evolutionary and
conservation biology. J Evol Biol 24:1160–1168.

Brown JB, Boley N, Eisman R, May GE, Stoiber MH, Duff MO, et al. 2014. Diversity and
dynamics of the *Drosophila* transcriptome. Nature. 7515:393–9.

Cao W, Edery I. 2017. Mid-day siesta in natural populations of D. melanogaster from Africa
exhibits an altitudinal cline and is regulated by splicing of a thermosensitive intron in the
period clock gene. BMC Evol. Biol. 17: 32.

Catalan A, Hutter S, Parsch J. 2012. Population and sex differences in *Drosophila melanogaster*
brain gene expression. BMC Genomics 13:1–12.

Chakraborty M, VanKuren NW, Zhao R, Zhang X, Kalsow S, Emerson JJ. 2017. Hidden genetic
variation shapes the structure of functional elements in *Drosophila*. Nat Genet 50:20–25.

Chen J, Nolte V, Schlötterer C. 2015. Temperature Stress Mediates Decanalization and
Dominance of Gene Expression in *Drosophila melanogaster*. PLoS Genet 11:e1004883.

Comeron JM, Ratnappan R, Bailin S. 2012. The many landscapes of recombination in
Drosophila melanogaster. PLoS Genet 8:e1002905.

Conte GL, Arnegard ME, Peichel CL, Schluter D. 2012. The probability of genetic parallelism
and convergence in natural populations. Proc Royal Soc B 279:5039–5047.

Coolon JD, McManus CJ, Stevenson KR, Graveley BR, Wittkopp PJ. 2014. Tempo and mode of
regulatory evolution in Drosophila. Genome Res 24:797–808.

Cowles CR, Hirschhorn JN, Altshuler D, Lander ES. 2002. Detection of regulatory variation in
mouse genes. Nat Genet 32:432–437

Degner JF, Marioni JC, Pai AA, Pickrell JK, Nkadori E, Gilad Y, Pritchard JK. 2009. Effect of
read-mapping biases on detecting allele-specific expression from RNA-sequencing data.
Bioinformatics 25:3207–3212.

Denby CM, Im JH, Yu RC, Pesce CG, Brem RB. 2012. Negative feedback confers mutational
robustness in yeast transcription factor regulation. Proc Natl Acad Sci USA 109:3874–3878.

Deshpande, G., Calhoun, G., Schedl, P. 2005. Drosophila argonaute-2 is required early in
embryogenesis for the assembly of centric/centromeric heterochromatin, nuclear division,
nuclear migration, and germ-cell formation. Genes Dev. 19:1680-1685.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15–21.

Duneau D, Sun H, Revah J, San Miguel K, Kunerth HD, Caldas IV, Messer PW, Scott JG,
Buchon N. 2018. Signatures of insecticide selection in the genome of *Drosophila*
melanogaster. G3 (Bethesda) 8:3469–3480.

Emerson JJ. 2019. Inferring compensatory evolution of cis- and trans-regulatory variation.
Trends Genet 35:1–3.

Fabian DK, Kapun M, Nolte V, Kofler R, Schmidt PS, Schlötterer C, Flatt T. 2012. Genome-
wide patterns of latitudinal differentiation among populations of *Drosophila melanogaster*
from North America. Mol Ecol 21:4748–4769.

Fear JM, Leon-Novelo LG, Morse AM, Gerken AR, Van Lehmann K, Tower J, Nuzhdin SV,
McIntyre LM. 2016. Buffering of genetic regulatory networks in *Drosophila melanogaster*.
Genetics 203:1177–1190.

Fraser HB. 2019. Improving estimates of compensatory cis–trans regulatory divergence. Trends
Genet 35:3–5.

Gamazon ER, Stranger BE. 2014. Genomics of alternative splicing: evolution, development and
pathophysiology Hum Genet 133:679-687

Gibilisco L, Zhou Q, Mahajan S, Bachtrog D. 2016. Alternative splicing within and between
Drosophila species, sexes, tissues, and developmental stages. PLoS Genet 12: e1006464.

Gibson JR, Chippindale AK, Rice AM. 2002. The X chromosome is a hot spot for sexually
antagonistic fitness variation. Proc. Biol. Sci. 269: 499–505.

Glaser-Schmitt A, Zecic A, Parsch J. 2018. Gene regulatory variation in Drosophila
melanogaster renal tissue. Genetics 210:287–301.

Good RT, Gramzow L, Battlay P, Sztal T, Batterham P, et al., 2014. The molecular evolution of
cytochrome P450 genes within and between Drosophila species. Genome Biol. Evol. 6:
1118–1134.

Hanson D, Hu J, Hendry AP, Barrett RDH. 2017. Heritable gene expression differences between
lake and stream stickleback include both parallel and antiparallel components. Heredity
119:339–348.

Hart JC, Ellis NA, Eisen MB, Miller CT. 2018. Convergent evolution of gene expression in two
high-toothed stickleback populations. PLoS Genetics 6:e1007443.

Helfrich-Forster, C. 2018. Sleep in Insects. Annu Rev Entomol 63:69-86.

Hoffmann AA, Hallas RJ, Dean JA, Schiffer M. 2003. Low potential for climatic stress
adaptation in a rainforest *Drosophila* species. Science 301:100–102.

Hoffmann AA, Weeks AR. 2007. Climatic selection on genes and traits after a 100 year-old
invasion: a critical look at the temper-ate-tropical clines in *Drosophila melanogaster* from
eastern Australia. Genetica 129:133–147.

Hsu SK, Jaksic AM, Nolte V, Barghi N, Mallard F, et al. 2019. A 24 h age difference causes
twice as much gene expression divergence as 100 generations of adaptation to a novel
environment. Genes 10:89.

Huylmans AK, Parsch J. 2014. Population- and sex-biased gene expression in the excretion
organs of *Drosophila melanogaster*. G3 (Bethesda). 4:2307–15.

Izquierdo JI. 1991. How does Drosophila melanogaster overwinter? Entomol. Exp. Appl. 59:
51–58.

Innocenti P, Morrow EH. 2010. The sexually antagonistic genes ofDrosophila melanogaster.
PLoS Biol. 8: e1000335.

James AC, Partridge L. 1995. Thermal evolution of rate of larval development in Drosophila
melanogaster in laboratory and field populations. J. Evol. Biol. 8: 315–330.

Jones FC, Grabherr MG, Chan YF, Russell P, Mauceli E, et al. 2012. The genomic basis of
adaptive evolution in threespine sticklebacks. Nature 484:55–61.

Josephs, EB., Lee. YW, Wood CW, Schoen DJ, Wright SI, et al. 2020. The Evolutionary Forces
Shaping Cis- and Trans-Regulation of Gene Expression within a Population of Outcrossing
Plants. Molecular Biology and Evolution, 37: 2386–2393

Juneja P, Quinn A, Jiggins FM. 2016. Latitudinal clines in gene expression and cis-regulatory
element variation in Drosophila melanogaster. BMC Genomics:1–11.

Kang J, Kim J, Choi K-W. 2011. Novel cytochrome P450, cyp6a17, is required for temperature
preference behavior in *Drosophila*. PLoS ONE 6:e29800.

Kitano J, Ishikawa A, Kusakabe M. 2018. Parallel transcriptome evolution in stream threespine
sticklebacks. Devel Growth Differ 61:104–113.

Kolaczkowski B, Kern AD, Holloway AK, Begun DJ. 2011. Genomic differentiation between
temperate and tropical Australian populations of *Drosophila melanogaster*. Genetics
187:245–260.

Kurmangaliyev YZ, Favorov,AV, Osman NM, Lehmann K-V, Campo D, et al. 2015. Natural
variation of gene models in Drosophila melanogaster. BMC Genomics 16:198

Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE. 2016a. A thousand fly genomes: An
expanded Drosophila genome nexus. Mol Biol Evol 33:3308–3313.

Lack JB, Monette MJ, Johanning EJ, Sprengelmeyer QD, Pool JE. 2016b. Decanalization of
wing development accompanied the evolution of large wings in high-altitude *Drosophila*.
Proc Natl Acad Sci USA 113:1014–1019.

Lai Z, Kane NC, Zou Y, Rieseberg LH. 2008. Natural variation in gene expression between wild
and weedy populations of *Helianthus annuus*. Genetics 179:1881–1890.

Landry CR, Lemos B, Rifkin SA, Dickinson WJ, Hartl DL. 2007. Genetic properties influencing
the evolvability of gene expression. Science 317:118–121.

Lange JD, Pool JE. 2016. A haplotype method detects diverse scenarios of local adaptation from
genomic sequence variation. Mole Ecol 25:3081–3100.

Lavington, E., and A. D. Kern, 2017 The effect of common inver-sion polymorphisms In(2L)t
and In(3R)Mo on patterns of tran-scriptional variation in Drosophila melanogaster. G3
(Bethesda)7: 3659–3668.

Leder EH, McCairns RJS, Leinonen T, et al. 2015. The evolution and adaptive potential of
transcriptional variation in sticklebacks—signatures of selection and widespread heritability.
Mol Biol and Evol 32:674–689.

Leinonen T, McCairns RJS, O'Hara RB, Merilä J. 2013. Q_{ST}–F_{ST} comparisons: evolutionary and
ecological insights from genomic heterogeneity. Nat Rev Genet. 14:179–190.

Lemmon ZH, Bukowski R, Sun Q, Doebley JF. 2014. The role of cis regulatory evolution in
maize domestication. PLoS Genet 10:e1004745–15.

Li B & Dewey Colin N. 2011. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. BMC Bioinform 12:323

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, et al. 2009. The Sequence alignment/map
(SAM) format and SAMtools. Bioinformatics 25:2078-2079.

Li YI, Knowles DA, Humphrey J, Barbeira AN, Dickinson SP, et al. 2017. Annotation-free
quantification of RNA splicing using LeafCutter. Nat Genet 50:151–158.

Liu, X., Y. I. Li, and J. K. Pritchard, 2019 Trans Effects on Gene Expression Can Drive
Omnigenic Inheritance. Cell 177: 1022–1034.e6.

Losos JB. 2011. Convergence, adaptation, and constraint. Evolution 65:1827–1840.

Machado HE, Bergland AO, O'Brien KR, Behrman EL, Schmidt PS, Petrov DA. 2016.
Comparative population genomics of latitudinal variation in *Drosophila simulans* and
Drosophila melanogaster. Mol Ecol 25:723–740.

Mack KL, Campbell P, Nachman MW. 2016. Gene regulation and speciation in house mice.
Genome Res 26:451–461.

Massouras A, Waszak SM, Albarca-Aguilera M, Hens K, Holcombe W, et al. 2012. Genomic
Variation and Its Impact on Gene Expression in *Drosophila melanogaster*. PLoS Genet
8:e1003055.

Mateo L, Ullastres A, González J. 2014. A transposable element insertion confers xenobiotic
resistance in *Drosophila*. PLoS Genet 10:e1004560.

McGirr JA, Martin CH. 2018. Parallel evolution of gene expression between trophic specialists
despite divergent genotypes and morphologies. Evol Lett 2:62–75.

McManus CJ, Coolon JD, Duff MO, Eipper-Mains J, Graveley BR, Wittkopp PJ. 2010.
Regulatory divergence in *Drosophila* revealed by mRNA-seq. Genome Res 20:816–825.

McManus CJ, Coolon JD, Eipper-Mains J, Wittkopp PJ, Graveley BR. 2014. Evolution of
splicing regulatory networks in *Drosophila*. Genome Res 24:786–796.

Meiklejohn CD, Coolon JD, Hartl DL, Wittkopp PJ. 2014. The roles of cis- and trans-regulation
in the evolution of regulatory incompatibilities and sexually dimorphic gene expression.
Genome Res 24:84–95.

Merilä J. 1997. Quantitative trait and allozyme divergence in the greenfinch (*Carduelis chloris*,
Aves: Fringillidae). Biol J Linn Soc 61:243–266.

Metzger BP, Duveau F, Yuan DC, Tryban S, Yang B, et al., 2016 Contrasting frequencies and
effects of cis- and trans-reg-ulatory mutations affecting gene expression. Mol. Biol. Evol.
33:1131–1146.

Müller L, Hutter S, Stamboliyska R, Saminadin-Peter SS, Stephanet W, et al. 2011. Population
transcriptomics of Drosophila melanogaster females. BMC Genomics 12: 81.

Nagoshi E, Sugino K, Kula E, Okazaki E, Tachibana T, et al. 2010 Dissecting differential gene
expression within the circadian neuronal circuit of Drosophila. Nat. Neurosci. 13: 60-68.

Nandamuri SP, Conte MA, Carleton KL. 2018. Multiple trans QTL and one cis-regulatory
deletion are associated with the differential expression of cone opsins in African cichlids.
BMC Genomics. 19:945.

Nayak A, Berry B, Tassetto M, Kunitomi M, Acevedo A, et al. 2010. Cricket paralysis virus
antagonizes Argonaute 2 to modulate antiviral defense in Drosophila. Nat. Struct. Mol. Biol.
17: 547-554.

Osada N, Miyagi R, Takahashi A. 2017. Cis- and trans-regulatory effects on gene expression in a
natural population of *Drosophila melanogaster*. Genetics 206:2139–2148.

Pitchers W, Pool JE, Dworkin I. 2012. Altitudinal clinal variation in wing size and shape in
african *Drosophila melanogaster*: one cline or many? Evolution 67:438–452.

Pool JE, Braun DT, Lack JB. 2016. Parallel evolution of cold tolerance within *Drosophila*
melanogaster. Mol Biol Evol 34:349–360.

Pool JE, Corbett-Detig RB, Sugino RP, Stevens KA, Cardeno CM, Crepeau MW, Duchen P,
Emerson JJ, Saelao P, Begun DJ, et al. 2012. Population genomics of sub-Saharan
Drosophila melanogaster: African diversity and non-African admixture. PLoS Genet
8:e1003080–24.

Pool JE, Nielsen R. 2007 Population size changes reshape genomic patterns of diversity.
Evolution 61:3001–3006.

Pool JE. 2015. The Mosaic Ancestry of the Drosophila Genetic Reference Panel and the *D.*
melanogaster reference genome reveals a network of epistatic fitness interactions. Mol Biol
Evol 32:3236–3251.

Powell JR. 1997. Progress and prospects in evolutionary biology, The *Drosophila* model. New
York, NY: Oxford University Press.

Qin, W., S. J. Neal, R. M. Robertson, J. T. Westwood, and V. K. Walker. 2005. Cold hardening
and transcriptional change in Drosophila melanogaster. Insect Mol. Biol. 14:607–613.

Reinhardt JA, Kolaczkowski B, Jones CD, Begun DJ, Kern AD. 2014. Parallel geographic
variation in *Drosophila melanogaster*. Genetics 197:361–373.

Rosenblum EB, Parent CE, Brandt EE. 2014. The molecular basis of phenotypic convergence.
Annu Rev Ecol Evol Syst 45:203–226.

Sackton TB, et al. 2019. Convergent regulatory evolution and loss of flight in paleognathous
birds. Science 364:74–78.

Said I, Byrne A, Serrano V, Cardeno C, Vollmers C, and Corbett-Detig R. 2018. Linked genetic
variation and not genome structure causes widespread differential expression associated with
chromosomal inversions. Proceedings of the National Academy of Sciences 115:5492–5497.

Schaefke B, Emerson JJ, Wang T-Y, Lu M-YJ, Hsieh L-C, Li W-H. 2013. Inheritance of gene
expression level and selective constraints on trans- and cis-regulatory changes in yeast. Mol
Biol Evol 30:2121–2133.

Schluter D. 2000. The ecology of adaptive radiation. Oxford Univ. Press, New York.

Schmidt PS, Paaby AB. 2008. Reproductive diapause and life-history clines in North American
populations of *Drosophila melanogaster*. Evolution 62:1204–1215.

Singer-Sam J, LeBon JM, Dai A, Riggs AD. 1992. A sensitive, quantitative assay for
measurement of allele-specific transcripts differing by a single nucleotide. PCR Methods
Appl.1: 160–163.

Signor S, Nuzhdin S. 2018. Dynamic changes in gene expression and alternative splicing
mediate the response to acute alcohol exposure in *Drosophila melanogaster*. Heredity
121:342-360.

Singh P, Börger C, More H, Sturmbauer C. 2017. The role of alternative splicing and differential
gene expression in cichlid adaptive radiation. Genome Biol Evol 9:2764–2781.

Smith CCR, Tittes S, Mendieta JP, Collier-Zans E, Rowe HC, Rieseberg LH, Kane NC. 2018.
Genetics of alternative splicing evolution during sunflower domestication. Proc Natl Acad
Sci USA 115:6768-6773.

Sokolowski, MB, Bauer SJ, Wai-Ping V, Rodriguez L, Wong JL, Kent C.1986. Ecological
genetics and behavior of Drosophila melanogaster larvae in nature. Animal Behavior
34:403–408.

Sprengelmeyer QD, Mansourian S, Lange JD, Matute DR, Cooper BS, Jirle EV, Stensmyr MC,
Pool JE. 2018. Discovery of *Drosophila melanogaster* from Wild African Environments and
Genomic Insights into Species History. biorxiv 45:1153–13.

St Pierre SE, Ponting L, Stefancsik R, McQuilton P, the FlyBase Consortium. 2013. FlyBase
102—advanced approaches to interrogating FlyBase. Nucleic Acids Res 42:D780–D788.

Stern DL. 2013. The genetic causes of convergent evolution. Nat Rev Genet 14:751–764.

Stevenson KR, Coolon JD, Wittkopp PJ. 2013. Sources of bias in measures of allele-specific
expression derived from RNA-seq data aligned to a single reference genome. BMC
Genomics 14:536.

Svetec N, Zhao L, Saelao P, Chiu JC, and Begun DJ. 2015. Evidence that natural selection
maintains genetic variation for sleep in Drosophila melanogaster. BMC Evol. Biol. 15: 41.

Testa ND, Ghosh SM, and Shingleton AW. 2013. Sex-specific weight loss mediates sexual size
dimorphism in Drosophila melanogaster. PLoS One 8:e58936.

Thurmond J, Goodman JL, Strelets VB, Attrill H, Gramates LS, et al. 2019. FlyBase 2.0: the
next generation. Nucleic Acids Res. 47:D759–D765. doi:10.1093/nar/gky1003.

Venables JP, Tazi J, Juge F. 2011. Regulated functional alternative splicing in *Drosophila*.
Nucleic Acids Res 40:1–10.

Verta JP, Jones FC. 2019. Predominance of cis-regulatory changes in parallel expression
divergence of sticklebacks. eLife. 8:e43785.

Vicoso, B, and B. Charlesworth, 2006 Evolution on the X chromosome: unusual patterns and
processes. Nat. Rev. Genet. 7: 645–653.

von Heckel, K., Stephan W., Hutter, S. 2016. Canalization of gene expression is a major
signature of regulatory cold adaptation in temperate Drosophila melanogaster. BMC
Genomics, 17:574.

Wang D, Sung HM, Wang TY, Huang CJ, Yang P, *et al.* 2007. Expression evolution in yeast
genes of single-input modules is mainly due to changes in trans-acting factors. Genome
Research 17: 1161–1169. https://doi.org/10.1101/gr.6328907
1302

Wei Y, Reveal B, Reich J, Laursen WJ, Senger S, *et al*. 2014. TORC1 regulators Iml1/GATOR1
and GATOR2 control meiotic entry and oocyte development in Drosophila. Proc. Natl.
Acad. Sci. U.S.A. 111: E5670--E5677.

Wittkopp PJ, Kalay G. 2011. Cis-regulatory elements: molecular mechanisms and evolutionary
processes underlying divergence. Nat Rev Genet 13:59–69.

Wittkopp PJ, Haerum BK, Clark AG. 2004. Evolutionary changes in cis and trans gene
regulation. Nature 430:85–88.

Wittkopp PJ, Haerum BK, Clark AG. 2008. Regulatory changes underlying expression
differences within and between *Drosophila* species. Nat Genet 403:346–350.

Yan H, Yuan W, Velculescu VE, Vogelstein B, Kinzler KW. 2002. Allelic variation in human
gene expression. Science 297:1143

Yu G, Wang L, Han Y, He Q. 2012. clusterProfiler: an R package for comparing biological
themes among gene clusters. OMICS 16:284-287.

Yvert G, Brem RB, Whittle J, Akey JM, Foss E, et al. 2003. Trans-acting regulatory variation in
Saccharomyces cerevisiae and the role of transcription factors. Nat. Genet. 35:57–64.

Zhao L, Begun DJ. 2017. Genomics of parallel adaptation at two timescales in *Drosophila*. PLoS
Genet 13:e1007016.

Zhao L, Wit J, Svetec N, Begun DJ. 2015. Parallel gene expression differences between low and
high latitude populations of *Drosophila melanogaster* and *D. simulans*. PLoS Genet
11:e1005184.

Zhang X, Cal AJ, Borevitz JO. 2011. Genetic architecture of regulatory variation in
Arabidopsis thaliana. Genome Res. 21:725–733.

Zhang Y, Wang ZH, Liu Y, Chen Y, Sun N, *et al*. 2019. PINK1 Inhibits Local Protein Synthes is
to Limit Transmission of Deleterious Mitochondrial DNA Mutations. Mol Cell:1127-1137.

1327