<u>Assessment of phylo-functional coherence along the bacterial</u>
<u>phylogeny and taxonomy</u>

Marcos Parras-Moltó[a], Daniel Aguirre de Cárcer[a#]

[a]Departamento de Biología, Universidad Autónoma de Madrid, Madrid, 28049 Spain.
(daniel.aguirre@uam.es).

Running title: Functional cohesiveness of phylogeny and taxonomy

ABSTRACT

In this report we use available curated phylogenies, taxonomy, and genome annotations

to assess the phylogenetic and gene content similarity associated with each different

taxa and taxonomic rank.   Subsequently, we employ the same data to delimit the

frontiers of functional coherence along the bacterial phylogeny. Among other things,

our results show that within-group phylogenetic and gene content similarity of taxa in

the same rank are not homogenous, and that these values show extensive overlap

between ranks. On the other hand, functional coherence along the 16S rRNA gene-

based phylogeny was limited to 44 particular nodes presenting large variations in

phylogenetic depth. For instance, the deep subtree affiliated to class Actinobacteria

presented functional coherence, while the shallower family Enterobacteriaceae-affiliated

subtree did not.

IMPORTANCE

While bacterial taxonomy and phylogeny resources, as well as related bioinformatic

tools, continue to improve, the question remains as to how they should best be

employed in studies using 16S rRNA gene surveys to assess bacteria-ecosystem

relationships, a widespread approach. The results contained herein lead to the

recommendation that all ranks from genus to class/phylum be employed if using

taxonomic binning in the analysis of 16S rRNA gene surveys. With regards to the use of

phylogeny or clustering-based approaches, single or arbitrary tree topology or sequence

distance thresholds should not be employed. Instead, the results presented can be used

to obtain more meaningful results in many microbial ecology and evolution research

scenarios. Moreover, we provide dedicated scripts and files that can be used to continue

the exploration of functional coherence along the bacterial phylogeny employing

different parameters or input data.

INTRODUCTION

Our knowledge of microbial communities' composition has greatly expanded over the last decade thanks to the advent of high-throughput sequencing-based metagenomic approaches. The use of strategies based on the high-throughput sequencing of 16S rRNA gene amplicons is nowadays widespread in studies seeking to assess community structure. The use of such phylogenetic marker is often preferred over shot-gun metagenomic sequencing due to its associated reduced costs and ease of analysis and interpretation. Most commonly, these studies aim not only to provide snapshots of community composition on different samples, but also to assess how these communities change along environmental gradients or sample groups, and significantly, why the observed changes take place in relation to the different putative functional roles of bacterial groups in the ecosystem.

Within these approaches, the taxonomic binning of reads based on the use of dedicated training sets remains the most common strategy in the description of community composition. The use of taxonomic ranks to organize the bacterial tree of life on the basis of evolutionary relationships has a colossal value in scientific communication, and it seems difficult to imagine the progress of microbial ecology without its use. However, the link between taxonomy and phylogeny or ecological function is not clear-cut (1). Another common strategy in the analysis of 16S rRNA gene data is the use of sequence clusters obtained at predefined similarity thresholds (OTUs) or, less frequently yet more meaningfully, the search for study-oriented patterns along the complete 16S rRNA gene phylogeny (e.g. (2, 3)).

The present study analyzes i) the phylo-functional coherence of taxa and taxonomic ranks, and ii) the frontiers along the bacterial phylogeny where gene content similarity and shared phylogeny cease to correlate. Our main goal is to provide adequate guidance for future analyses employing 16S gene surveys aiming to improve our understanding of the role that Bacteria play in the ecosystem.

RESULTS

Our results are based on available curated phylogenetic trees and 16S rRNA gene sequence data from the Genome Taxonomy Database (http://gtdb.ecogenomic.org/downloads), as well as genome annotations from proGenomes (http://progenomes.embl.de/). Due to uneven sampling, the number of available members per taxa varied wildly (Suppl. Fig. 1, Suppl. Mat. 1). In all five ranks considered (genus, family, class, order, phylum) a large proportion of groups contained few members, while a small proportion presented a large number of members. This fact will likely impact the overall accuracy of classifiers using the available data as training set (4).

Phylogenetic similarity, assessed using 16S rRNA gene distances as proxy, was also quite variable among groups within the same taxonomic rank (Figure 1A, Suppl. Mat. 1). As expected, the per-rank averages of intra-taxa distance averages followed an ascending trend from genus to phylum (Figure 1A). While correlated, the observed values are markedly smaller than the controversial cut-offs often employed in the field; 0.05, 0.10 and 0.20, for genus, family, and phylum, respectively (5). Another interesting trend relates to the coefficients of variation associated with the per-rank distributions of intra-taxa distance averages (from genus to phylum; 1.08, 0.73, 0.59, 0.48 and 0.36).

The values seem to indicate that phylogenetic depth of taxa in a rank becomes more

stable from genus to phylum. Equally anticipated, per-rank distributions of intra-taxa

gene content distance averages also followed an ascending trend from genus to phylum

(Figure 1A, Suppl. Mat. 1).

When analyzing intra-taxa distance averages (Figure 1A) and standard deviations

(Suppl. Fig. 2) it becomes clearer that taxonomic ranks are not cohesive in terms of

phylogenetic depth and gene content similarity. That is, for all taxonomic ranks, the

different taxa within the same rank do not present a homogeneous level of either

phylogenetic depth or gene content similarity. In this regard, the phylogenetic and gene

content similarity values of many taxa was lower than those for taxa in superior ranks,

and *vice versa*. This circumstance most likely stems from the fact that available

genomes are not evenly distributed throughout the bacterial tree of life, that for

operational and communication purposes the number of ranks is small, and that the

main focus of modern taxonomy is related to the production of monophyletic taxa (6).

Our initial results showed the expected overall correlation between gene content and

16S-based distances (Figure 1B). However, a noticeable discontinuity appeared in the

pairwise 16S-based distance values at *ca.* 0.5 which, in turn, seems to mark the limit of

the observed correlation. In order to better explore the frontiers of functional coherence

along the bacterial phylogeny, we developed *FunCongr.R;* a script that estimates

functional coherence at each node of a tree by comparing within-node average gene

content distance against what could be expected from a random draw of genomes along

the phylogeny. The script also defines a node as non-functionally-coherent if any of its

descendant nodes is non-functionally-coherent.

The analysis of the 16S rRNA gene phylogeny (Figure 2, Suppl. Fig. 3, Suppl. Mat. 2)

returned 44 functionally-coherent nodes. These nodes present large variations in depth,

showing within-node average 16S rRNA gene distance values ranging from 0.03 to 0.25

(average 0.12 ± 0.05). Without the goal of being exhaustive, a subsequent analysis based

on the consensus (80%) taxonomy of leaves in each node, with the sole criterion of

whether a particular taxon appears assigned to a single coherent node, revealed various

deep-branching nodes affiliated to the Acidobacteriota (p), Alphaproteobacteria (c),

Bacilli (c), Actinobacteria (c), Cyanobateriia (c), Fusobacteriales (o) Campylobacterales

(o), and a node classified merely as Bacteria (d). On the other hand, the

Gammaproteobacteria-affiliated region of the tree was partitioned among many

different coherent nodes, hence indicating that such class is not functionally coherent in

itself. Here, the Betaproteobacterales (o) stood out as a large and coherent node, while

many different coherent nodes appeared affiliated to the Pseudomonadales (o),

Enterobacterales (o), and even Enterobacteriaceae (f).

DISCUSSION

While bacterial taxonomy and phylogeny resources, as well as related bioinformatic

tools, continue to improve, the question remains as to how they should best be

employed in studies using 16S rRNA gene surveys to assess bacteria-ecosystem

relationships, a widespread approach. With regards to the use of taxonomic binning, our

results show that within-group phylogenetic and gene content similarity of taxa in the

same rank are not homogenous, and that these values show extensive overlap between

ranks. Thus, we believe that most taxonomic ranks should be considered when assessing

bacteria-ecosystem relationships, since it is not possible to know *a priori* which sections

and depths of the bacterial phylogeny will be affected by the studied environmental

gradient or sample group. The possible exceptions are species and phylum, since the

16S rRNA gene presents low phylogenetic resolution at the highest and lowest

taxonomic ranks (7), and very few of the dereplicated genomes employed by Parks *et al.*

present species-level classification.

With regards to the use of phylogeny-based approaches, the 16S rRNA gene-based

phylogeny showed remarkable correlation with gene content similarity. However, the

breaking points of such relationship appeared at different depths along each different

branching path of the phylogeny. Thus, single, arbitrary tree topology or sequence

distance thresholds should not be employed regarding phylo-functional coherence. For

instance, while a deep Actinobacteria (c)-affiliated node passed our functional

coherence test, the Enterobacteriaceae (f)-affiliated shallower subtree did not.

The strategy employed could be biased by low quality genome annotations, which

prompted us to employ the curated proGenomes database. In this sense, the depth of the

frontiers of functional coherence described here could be understood as conservative

estimates; for instance within the Bacteroidota (p)-affiliated subtree, node85 failed to

pass the functional coherence test, and thus the subtree was split into several coherent

units (Figure 2, Suppl. Fig. 3). While reducing the p-value of the permutation test to

0.01 returned the exact same tree topology, node85 contains only five genomes, and

thus missannotation could potentially have artificially reduced the limits of phylo-

functional coherence in this subtree. On the other hand, the loss of functional coherence

within the Enterobacteraceae (f)-affiliated subtree, and hence that of the larger

Gammaproteobacteria (c)-affiliated subtree as well (Figure 2, Suppl. Fig. 3), was

supported by the analysis of 35 genome annotations (node24704).

In addition to our previous results-driven recommendation that all ranks from genus to

class/phylum be employed if using taxonomic binning, we argue that our functional

coherence proxy (i.e. more within-node gene content similarity than expected by

chance, and no descendant nodes failing to pass such test) is intuitive and useful. Thus,

the results presented here can be used by the community, together with Park *et al.*'s

provided phylogenetic trees, to obtain more meaningful results in many microbial

ecology and evolution research scenarios. Moreover, the associated scripts and files are

freely available (https://git.io/Jec5U) so that (e.g.) different phylogenies, genome

annotations, or significance thresholds can be used in the assessment of phylo-

functional coherence.

METHODS

Phylo-functional coherence of taxa and ranks. First, the 16S rRNA gene sequences

for the representative (dereplicated) genomes employed to produce Parks *et al.*'s

taxonomy   (bac_ssu_r86.1.fna)   were   obtained   from   their   repository

(http://gtdb.ecogenomic.org/downloads)(6). Then, *Mothur* (8) commands were used to

align the sequences against SILVA reference alignment (9), and trimmed to contain

only the sequence delimited by universal primers 27f and 1492R (10). Sequences not

spanning the region were removed, leaving a total of 15 186 sequences. Taxa not

including at least three members were not considered for subsequent analyses. Distances

between the remaining 16S rRNA gene sequences were obtained using *Mothur,* and

later analyzed on the basis of their taxonomic affiliation (bac_metadata_r86.tsv),

producing summary statistics for all taxa at all ranks.

NCBI´s TaxIDs for each of the representative genomes (included within

bac_metadata_r86.tsv) were linked to the gene content information obtained from the

proGenomes resource (11) (complete annotation table kindly provided by Daniel

Mende), resulting in 6 989 different annotations for representative genomes presenting

nearly full length 16S rRNA gene sequences. Pairwise Jaccard distances between these

genomes were derived from the shared presence of genes using the *R* package *vegan*,

again producing within-taxa summary statistics for all taxa at all ranks.

Frontiers of functional coherence within phylogenetic trees. We developed a

dedicated R script, *FunCongr.R*, to explore the limits of functional coherence along the

bacterial   phylogeny.   Our   approach   employs   a   phylogenetic   tree

(gtb_r86.ssu.bacteria.fasttree.tree (6)) and pairwise Jaccard distances between genomes

obtained as mentioned above. The trees were initially processed to include constant

node labels, and pruned to remove leaves without functional annotation.

The script traverses the tree from leaves to root in such a way that nodes are evaluated

only if its descendant nodes have already been evaluated. Only nodes with at least 5

leaves are analyzed. At each non-flagged node, the script compares the average value of

pairwise Jaccard distances between its leaves to an empirical cumulative distribution

(ECD) of 1000 average values of pairwise Jaccard distances between N random leaves

along the phylogeny, where N is the actual number of leaves of the node being

evaluated. If the node's average Jaccard distance value is not within the lowest 5% of

the ECD, the node is flagged. Once a node has been flagged, all its ascendant nodes are

automatically flagged.

In this manner, we delimit functional coherence along the tree by pinpointing nodes that

either i) fail to pass the random ECD test, or ii) present descendant nodes that fail to

pass the ECD test. Thus, here functional coherence is presumed to persist if within-node

average gene content distance is lower (i.e. more similar) than what could be expected

from a random draw of genomes along the phylogeny (with $p<0.05$). The results are

finally processed by recording the last functionally-coherent node along each branching

path, along with their summary statistics. All figures of this work were produced using

ggplot2 package in R.

Acknowledgements

We thank the Bioinformatics Unit at CBMSO for their support. This work was funded

by the Spanish Ministry of Science and Innovation grant BIO2016-80101-R. The

funders had no role in study design, data collection and interpretation, or the decision to

submit the work for publication

Data availability

The datasets analyzed during the current study are available from their original source

(as stated above) and at (https://git.io/Jec5U).

REFERENCES

1.   **Kunin V, Ahren D, Goldovsky L, Janssen P, Ouzounis CA.** 2005. Measuring genome
conservation across taxa: divided strains and united kingdoms. Nucleic Acids Res
33:616-621.
2.   **Aguirre de Cárcer D.** 2018. The human gut pan-microbiome presents a compositional
core formed by discrete phylogenetic units. Scientific Reports **8:**14069.
3.   **Washburne AD, Silverman JD, Leff JW, Bennett DJ, Darcy JL, Mukherjee S, Fierer N,**
David LA. 2017. Phylogenetic factorization of compositional data yields lineage-level
associations in microbiome datasets. PeerJ **9.**

4.   **Beiko RG.** 2015. Microbial Malaise: How Can We Classify the Microbiome? Trends in
Microbiology **23**:671-679.
5.   **Schloss PD, Handelsman J.** 2004. Status of the Microbial Census. Microbiology and
Molecular Biology Reviews **68**:686.
6.   **Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA,**
Hugenholtz P. 2018. A standardized bacterial taxonomy based on genome phylogeny
substantially revises the tree of life. Nat Biotechnol **27**.
7.   **Janda JM, Abbott SL.** 2007. 16S rRNA gene sequencing for bacterial identification in
the diagnostic laboratory: pluses, perils, and pitfalls. J Clin Microbiol **45**:2761-2764.
8.   **Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,**
Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ,
Weber CF. 2009. Introducing mothur: open-source, platform-independent,
community-supported software for describing and comparing microbial communities.
Appl Environ Microbiol **75**:7537-7541.
9.   **Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J,**
Ludwig W, Glöckner FO. 2014. The SILVA and "All-species Living Tree Project (LTP)"
taxonomic frameworks. Nucleic Acids Research **42**:D643-D648.
10.   **Weisburg WG, Barns SM, Pelletier DA, Lane DJ.** 1991. 16S ribosomal DNA
amplification for phylogenetic study. Journal of bacteriology **173**:697-703.
11.   **Mende DR, Letunic I, Huerta-Cepas J, Li SS, Forslund K, Sunagawa S, Bork P.** 2017.
proGenomes: a resource for consistent functional and taxonomic annotations of
prokaryotic genomes. Nucleic Acids Research **45**:D529-D534.

FIGURE LEGENDS

Figure 1. Panel A; Box plots describing average within-group phylogenetic and gene-

content distance values for each of the proposed taxa in all considered ranks. **Panel B;**

Correlation between 16S rRNA gene and gene content-based distances. Each point

represents both distance values for each of the possible pairwise comparisons among the

6 989 dereplicated genomes with genome annotations ($y = 1.32x + 0.16$, $R^2 = 0.37$).

Figure 2. Frontiers of functional coherence along the 16S rRNA gene phylogeny.

The tree represents Park *et al*'s 16S rRNA gene-based phylogeny, pruned to contain

only leaves with nearly full length 16S rRNA sequences and functional annotations.

Colored boxes mark limits of phylo-functional coherence. Only the most prominent

results are annotated (see Suppl. Fig. 3 for an annotated and vertical version of the tree).

*Nodes that failed to past the functional coherence test.