1   **Title Page**

6   Full title:

7   **Somatic genetics of CDR3 control TCR V-domain rotational probability**

8   **effecting germline CDR2 scanning of polymorphic MHC**

10 Running title:

11 **Genetic calculus of TCR binding class-II pMHC**

16 Joseph S. Murray†

17
18 jsmurray@xenolausgenetics.com

22 **†Xenolaüs Genetics LLC**
23 10880 Wilshire Blvd., Suite 1101, Los Angeles, CA, 90024, USA

Somatic genetics of CDR3 control TCR V-domain rotational probability effecting germline CDR2 scanning of polymorphic MHC

Joseph S. Murray†*

†*Xenolaüs Genetics LLC*
Los Angeles, CA   UNITED STATES

Email Author: Dr. J.S. Murray, Ph.D. (jsmurray@xenolausgenetics.com)

Abstract:

The mechanism which adapts the T-cell antigen receptor (TCR) within a given *major histocompatibility complex* (MHC; HLA, in humans) genotype is essential for protection against pathogens. Historically attributed to relative affinity, genetically vast TCRs are surprisingly focused towards a micromolar affinity for their respective peptide (p) plus MHC (pMHC) ligands. Thus, the somatic diversity of the TCR with respect to *MHC restriction*, and (ultimately) to pathogens, remains enigmatic. Here, we derive a *triple integral equation* (from fixed geometry) for any given V-domain in TCR bound to pMHC. We examine solved complexes involving HLA-DR and HLA-DQ, where genetic linkage to the TCR is most profound. Certain V-beta domains displayed rare geometry within this panel—specifying a very low (highly-restricted) rotational probability/volumetric density (dV). Remarkably, hydrogen (H)-bond *charge-relays* distinguished these structures from the others; suggesting that CDR3 binding chemistry dictates CDR2 contacts on the respective MHC-II alpha-helix.

Keywords:

T-cell receptor; Major Histocompatibility Complex; TCR Selection; MHC Restriction; Protein Chemistry; Mathematical Biology

Introduction:

For better and (for) worse all the biology of the adaptive immune system cascades from a stochastic genetic mechanism (*somatic DNA rearrangement*)—generating TCR and antibodies (sIg) individualized within each precursor of a given T-cell or B-cell clone [1–4]. Thus, an enormous repertoire of these receptors (*est*. $\approx 10^8$ different α / β TCRs) pre-exists *clonal selection* against a given pathogen in a given human [4]. TCR (here-in, refers to α/β-type only) are utilized against antigenic structures which are all partially similar—all are MHC (human, HLA) proteins transporting a single 8–25 amino acid (a.a.) peptide (*abbrv.*, pMHC). The peptide component is extremely variable however; derived from intra-cellular, proteolytic processing of virtually any *self*, or non-self protein [4, 5]. In-fact, TCR diversity exceeds Ig diversity, although antibodies historically defined a V-region, *e.g.,* their elicitation against hitherto not-synthesised

66 organic structures [6].   Classic MHC class-I (HLA-A, -B, -C) and -II (HLA-DR, -DQ, -DP) genes
67 include the most polymorphic in the human genome, but diversity is limited to the alleles on
68 maternal and paternal sixth chromosomes [5, 7, 8].
69
This paper seeks to understand two puzzles of TCR:pMHC, wherein a novel theory for the first
can be used to re-examine the second in the context of existing evidence. Firstly, compared to
Ig and indeed other protein:protein binding reactions, TCR are of quite low affinities [6, 7].
Nevertheless, like antibodies, TCR do have exquisite specificity, where single peptide or MHC
a.a. changes can dictate reactivity or non-reactivity [7]. Second, the nature of T-cell selection in
the thymus prior to seeding secondary lymphoid tissues is thought to operate on the basis of
this same relative TCR:pMHC affinity, where overt 'self' peptide plus self-MHC recognition
(*negative selection*) leads to deletion from the repertoire, as does a complete lack of the same
recognition (*positive selection*). This *intermediate* TCR:pMHC binding in the foetal thymus
forms the basis for crucial biology of the T-cells' co-recognition of self-MHC simultaneous with
any given pathogen in adulthood, *i.e.*, the *MHC-restricted* adaptive immune response [4].
Despite extreme MHC polymorphism within the population, and co-dominance within the
individual, overwhelmingly most of the diversity in a patient's pMHC structures stem from the
peptide possibilities for these ligands—indeed, the 'hyper-variant' CDR3 loops are positioned
(conspicuously) atop the peptide component in solved TCR:pMHC complexes [9–15]; although
recent investigations have challenged ubiquity of this structure/function [16-18].
86
For example, we previously reported that one of the CDR3 loops (CDR3α) also makes
consistently close contacts with a conserved HLA-A component, *i.e.,* the *R65-motif* (a.a., R65-X-
X-K68-A69-X-S71-Q72) [17]. *Burrows et al*., also found conserved CDR3 contacts directly with
MHC-I [19]. More recently, *Sharon et al* formally linked germline TCR usage to HLA, particularly
via the class-II alleles [15]. Here, the *STCR-Dab* (www. opig.stats.ox.ac.uk/webapps/stcrdab/)
was used to identify all current TCR complexes involving HLA-DR and HLA-DQ (3.0 Å resolution
cut-off). Note, also included are *PDB* 4H1L, at 3.3 Å, and 3T0E, at 4.0 Å (see below), and the
'unusual topology' complexes (*PDB* 1YMM, 4Y1A & 4Y19) are excluded based on this
investigation targeting characteristic features.
96
We established a simple relationship between the observed (*i.e.*, measured) CDR3–CDR2
"pitch" angle for a given V-domain and its predicted (*i.e.*, calculated) CDR3–CDR2 pitch angle
[17]. Here, we found a linear relationship between calculated pitch and a new measure, "*d*V"—
by multivariable calculus. As is discussed, *d*V interprets V-domain orientation into a rotational
probability involving an apparent CDR2–MHC α-helix scanning function ($d\theta$). While *d*V were
unique for each TCR on each pHLA-DR, one TCR displayed a dramatic restriction in *d*V for Vβ.
This was isolated to a charge-relay H-bonding mechanism for CDR3β; hence, the chemistry of
"somatic-TCR" might dictate positioning of "germline-TCR" [20–22]. Within the seven pHLA-DQ
structures, the two *highly-restricted dV* TCR (4OZG & 4OZH) displayed distinct, yet functionally
similar mechanisms that shifted the same TCR:MHC H-bond by one MHC a.a. position (relative
to the nominal *d*V structure, 4OZF). Crucially, this involves an additional H-bond (MHC:MHC) in
suitable charge-relay mechanisms of 4OZG and 4OZH. We will discuss how these data suggest
that TCR binding *specificity* could be controlled by *orientation* (dynamics).

Results:

PDB files of solved TCR:pHLA-DR, and TCR:pHLA-DQ structures were used to investigate V-
domain geometry amongst the available complexes, *i.e.*, involving similar (but different) TCR,
and/or similar (but different) pMHC. All of these structures share the canonical (diagonal)
orientation of the TCR over pMHC, which was one of the earliest observed similarities between
different complexes [10]. As indicated in the summary tables (Tables 1A–1D; separation into
different tables is simply for clarity) some of the structures share the same TCR, where different
peptide (2IAM *vs.* 2IAN), or different DRB1 (1J8H *vs.* 1FYT; 6CQL *vs.* 6CQQ *vs.* 6CQR) were
compared. Some structures share pMHC binding different TCR (2IAM *vs.* 4E41; 6CQL *vs.* 6CQN;
4OZF–I). All the DR structures share the DRA chain [8, 23] and in some cases, so, too, the DRB1
allotype (2IAM, 2IAN, 4E41, 1FYT, 6CQR; all DRB1-1), (1J8H, 3T0E; both DRB1-4), (6CQL, 6CQN;
both DRB1-11); 6CQQ is DRB1-15, 4H1L is DRB3 and 1ZGL is DRB5 (isotypes). The DQ structures
are DQ8 (5KSA, 5KS9, 4GG6), or DQ2.5 (4OZF–I) (*NCBI* annotation and *BLAST*;
www.ncbi.nlm.nih.gov). While bound peptides in the DR group are of diverse protein origins,
all the DQ peptides are wheat *gliadins*; either the glia-gamma-1, glia-alpha-2, or glia-MM1.
Interest in these particular DQ structures is due to *celiac disease*, an autoimmune-like condition
known to be mediated by predominant T-cell clonotypes against ingested wheat. Some 95% of
celiac patients express HLA-DQ2.5 (DQA*05/DQB1*02), while the rest are either HLA-DQ8
(DQA*03/DQB1*03:02), or DQ2.2 (DQA*02:01/DQB1*02) [24]. For our purposes, the crystal
structures of these nearly identical complexes facilitated isolating the chemistry associated with
V-domain rotational probability/volumetric density (*d*V). Note that all sequences for each
component of all of these structures are available under the appropriate *PDB* file name at the
NCBI (www.ncbi.nlm.nih.gov).

TCR-pMHC Geometry

Shown in Figures 1A and 1B is an example of the geometry analysis based on fixed a.a. positions
in the HLA-DR, -DQ grooves (*e.g.*, *PDB* 1J8H). We originally used *Euler* angles
(www.mathword.wolfram.com/EulerAngles.html) to establish the basic method (for a TCR-Vα
cohort in pHLA-A2 complexes) [17]; here, we modified the method for the available solved
TCR:pHLA-DR, -DQ structures (summarized in Tables 1A–1D). In brief, the analysis is based on
measuring three angles corresponding to the *twist* (ω), *tilt* (λ), and *sway* (σ) of each V-domain
over the pMHC.

146
147
Figure 1 **The twist/tilt/sway of TCR-Vα and -Vβ relative to pMHC-II (A & B, respectively).** Illustrated by
the example of 1J8H. This same analysis was performed on all 19 structures (Tables 1A–1D). All angle
measurements were from Cα in *VMD-1.9.1* (www.ks.uiuc.edu) used to examine the diversity through the
38 V-domains: (i) in-plane to the MHC-groove {*twist* = ω}, with (ii) displacements perpendicular to the
groove {*tilt* = λ}, including (iii) side-to-side variation {*sway* = σ}. The a.a. positions used as coordinates
for angular measures (*dotted orange* vectors) across structures were fixed and are labelled (see text).
The measured incline of a V-domain, or "pitch" {*pitch* = φ_m}, is shown by *white dotted* vectors and could
be approximated by the equation: $\varphi_c = [\sigma \div (\lambda + \sigma)]\omega$ (see text, Tables 1A–1D). Side chains of a.a. in
measurements are shown by *CPK licorice*; Cα-backbones are in *new cartoon* and labelled; peptides are in
black. Note from A to B the view of the structure rotates 180°.

158
For HLA-DR a vector from the DRA a.a. N62 alpha-carbon (N62:Cα) to DRB1/3/5 a.a. D66
(D66:Cα; Cα used for measurements unless otherwise noted) bisects the MHC-groove from the
DRA α-helix to the DRB1/3/5 α-helix, then the angle at D66 to the Vα central cysteine (C22) is
computed with the *VMD angle-label* tool. This ω-angle can be seen in Figure 1A (Vα) and 1B
(Vβ; where the vectors run D66:N62:C23) as *dotted orange* lines near the *melon*-coloured "ω"
symbol, 86.88° (Vα), 84.91° (Vβ). The E11 (beta-sheet floor) to N62 to C22 (*tilt*-angle) is
similarly shown (1A) near the "λ" symbol, 134.93°; for Vβ, D66 is used (1B), F40:D66:C23, at
109.62°. Finally the σ-angle is measured from N62:E11:C22 (Vα) or D66:F40:C23 (Vβ), here at
35.67° and 47.99°, respectively. Analogously for HLA-DQ, TCR:pMHC structures, Vα *twist* is
based on a DQA a.a. N62:Cα to DQB D66:Cα vector bisecting the groove, where the angle vertex
is at D66 to Vα C23:Cα. Here, N11:Cα (not E11 found in DR) to the N62 vertex to C23 defines
tilt, and the reverse N62:N11:C23 is *sway*. For the DQ Vβ's, angles are the same as in DR
structures. Specifically, *twist* is defined by D66:N62:C23, *tilt* by F40:D66:C23, and *sway* as
D66:F40:C23 (Table 1D). We formulated the equation:
173
174 $$\varphi_c = k\varphi_m = [\sigma \div (\lambda + \sigma)]\,\omega$$
175
that predicts the pitch of a given V-domain (*pitch* = angle φ) from the ω, λ, and σ. By contrast,
measuring the φ-angle is based on finding the closest contact between the domain CDR2 and
the α-helix that is opposite N62 (for Vα) or D66 (for Vβ) (*yellow* highlighted in Tables 1A–1D).
The vectors are directed from that CDR2 a.a. to the across-the-groove α-helix a.a., then back to

the α-helix a.a. that is in contact with the CDR2 a.a. (*white dotted* lines, Figs. 1A & 1B). In Tables
1A–1D the calculated (φ_c) and measured (φ_m) pitch angles are shown along with the *k* values.
Thus, to the extent *k* approaches 1.00 indicates correlation between the observed pitch and
calculated pitch values. **Statistics**: HLA-DR group: *mean k* = 0.99 ± 0.36 (*s*), *n* = 24, *t* = 0.915, μ_0
= 1.00, *p* = 0.376, 95% CI, 0.880–1.10; (paired *Student*'s *t*-test; www.graphpad.com/quickcalcs;
www.select-statistics.co.uk). HLA-DQ group: *mean k* = 0.88 ± 0.35 (s), *n* = 14, *t* = 1.24, μ_0 = 1.00,
p = 0.234, 95% CI, 0.750–1.01. Overall (DR & DQ): *mean k* = 0.95 ± 0.35 (s), *n* = 38, *t* = 0.85, μ_0 =
1.00, *p* = 0.402, 95% CI, 0.870–1.03.

Triple Integrals

We observed that the closest CDR2α contacts with the α2-helix 'across-the-groove' from the
conserved R65 contact with CDR3α in HLA-A2 structures specified a polymorphic region from
a.a. H/R151 to a.a. A158 [17]. Similarly, the CDR2α closest contacts range for the DR structures
here (Tables 1A–1C) implicate a DRB α-helix range from a.a. E69 to a.a. T77; for CDR2β, a DRA
α-helix contacts range: a.a. Q57 to a.a. K67. For the DQ complexes: CDR2α contacts range is
DQB α-helix a.a. E69 to a.a. D76; and for CDR2β, DQA: Q57 to H68 (Table 1D). Thus, we
hypothesized that if these α-helical a.a. are indeed swept/scanned by CDR2 of the V-domain,
that we should be able to model such with integration from *spherical coordinates* in each
structure [25–28]. To normalize the equation across structures and use the geometry, the
central cysteine (C22/23) was chosen as a point rotating from the fixed across-the-groove a.a.
position that defines *twist*, *tilt*, and *sway* (Figs. 1 & 2). As shown in Figure 2B, the coordinates
model a 'slice' of a cone [26], where for each of the 38 V-domains we measured the distance
between the C22/23 and said α-helix a.a. (*e.g.*, 2IAM Vβ measures 22.12 Å)—this is the *rho* (ρ)
distance in the equation. The other measure was the angle ϕ, which is simply the difference
from 180° of the previously determined *tilt* (angle λ). The other variables were derived by
trigonometry (see Fig. 2B).

208
209

Figure 2 **Derivation of the V-domain *rotational* volumetric-density equation.** The same a.a. positions
used for the tilt (λ) angles were used to define the phi (ϕ) angles (*e.g.*, 49.85° for this structure, 2IAM Vβ,
gold) by subtraction of λ from a theoretical 180° vector. This is illustrated with actual positions in 2IAM
(A), showing the 180° approximating, 176.32° vector; the ~ 90°, 88.91° angle; the ~ 14.27 Å, 14.14 Å line
segment; and the ~ 16.90 Å, 17.29 Å line segment, *i.e.,* to show the essential geometry within a
structure; these are more accurately calculated by trigonometry in the derivation (B). Additionally, the
rho (ρ) segment from the appropriate α-helix position (see λ definitions) to a given V-domain's C22/23
was measured (here, 22.12 Å). The cone (full rotation) includes a probability (slice volume) through a
scanning path (*d*θ) for CDR2.
219
This is integration of a volume element in spherical coordinates (all *d*V values in Tables 1A–1D).
For strictly compositional reasons throughout "*the triple integral of d*V" may be referred to as
simply "*d*V." As discussed above it is not surprising that the ($\pi \cdot$ r / 4) values were close to the
actual distances between CDR2 contacts in the aforementioned MHC α-helix ranges using this
approach, *e.g.*, 13.27 Å, Figure 2B. Thus, the volume of the cone slice for each V-domain was
determined upon integration for the three spherical coordinates, rho (ρ), theta (θ) and phi (ϕ),
where the upper limit of each integral is derived from the measured ρ and ϕ values of each V-
domain as shown:
228

$$\iiint_V dV = \int_0^{\sin^{-1}(r/\rho)} \int_0^{\pi \text{-} r/4} \int_0^{z/\cos\phi} \rho^2 \sin\phi \, d\rho \, d\theta \, d\phi$$

229
230
231 Since ρ and ϕ are measured for each V-domain, derivation of upper limits for the first and third
232 integrals simply provide trigonometric relationships.
233

Table 1A   Summary Geometry TCR:pHLA-DR

PDB	CDR2[1]	CDR3	σ [¶]	λ	ω	φ	k[‡]	ρ (Å)[◊]	dV[‡]
2IAMα[*]	K53NZ:E69OE2 2.77	G92O:G58N 3.34	50.75	116.86	101.05	30.73c 18.13m	1.69	29.55	1703
2IAMβ	V50CG2:Q57CB 3.96	T98OG1:D66OD2 2.53	30.97	130.15	66.21	12.73c 17.69m	0.719	22.12	297
2IANα	K53NZ:E69OE1 2.95	A93N:E55O 3.51	50.99	116.16	99.72	30.42c 19.32m	1.57	29.39	1710
2IANβ	Y48OH:Q57NE2 3.30	T98OG1:D66OD1 2.74	30.27	130.76	64.98	12.21c 29.69m	0.411	21.78	270
4E41α	K53NZ:E69OE2 2.63	T93OG1:E55OE1 3.17	48.39	119.79	100.80	29.00c 16.93m	1.71	29.51	1512
4E41β	N54ND2:E55OE2 3.24	T97OG1:D66OD2 2.51	32.08	129.37	65.97	13.11c 26.55m	0.494	23.67	406

[*]Same TCR (E8) and MHC (DRA1, DRB1*0104; pep I28) as 2ian (pep T28); same pMHC as 4e41 (TCR, G4).

[1]Contacts in angstroms (Å) measured with VMD software (www.ks.uiuc.edu); *yellow* highlighted contacts of the measured *pitch* angle (φ_m), see text. Colour-coded possible H-bonds, pH 7: red (strong), magenta (medium), green (weak); hydrophobic (blue). The calculated *pitch* angle (φ_c) of Vα and Vβ determined as before, using the formula:

$$^{\ddagger}\ \varphi_c = [\sigma \div (\sigma + \lambda)] \cdot \omega$$

[¶]Measured angles with VMD in degrees; [◊]The value for ρ as measured with VMD in Å; volume of the rotational density distribution (about N-term Cys of the intra-domain disulfide) for Vα and Vβ was modeled as a cone (see text) where[¥]:

$$\iiint_V dV = \int_0^{\sin^{-1}(r/\rho)} \int_0^{\pi \cdot r/4} \int_0^{z/\cos\phi} \rho^2 \sin\phi \, d\rho \, d\theta \, d\phi$$

And where, $\pi/180$ is the conversion factor for the angular component of the integrand to yield dV in $\overset{\circ}{A}{}^3$.

234
235
Two example solutions are given below (parameters as defined, Figs. 1 & 2; text)—*for all 38*
triple integral solutions, see **Suppl.1.I***. The ($\pi \cdot r / 4$) circumference segment is used as the
upper limit of the $d\theta$ integral because it is accurate to a *path*, *i.e.*, a *distance*; formally (for θ-
angle in degrees): $\pi \cdot r$ (θ) / 180 = *arc length* in Å [26]. Hence, $d\phi$ is the only integrand for
conversion (multiplying by $\pi \div 180$) to yield the *cubic angstrom* (Å³) unit of volume. The C22/23
cysteine is historically used as the center of any given V-domain [6] and each V-domain slice is
thus the volumetric-density through its CDR2 *scanning* (*e.g.*, Fig. 2B). The calculus interprets a
comprehensive geometry of the V-domain into a *probability* of scanning using only the *ground-*
state structure, *i.e.*, *without* crystallographic data on theoretical conformers—suggesting here,
a classic protein function, *allostery* [28, 29]; see also 'protein machinery' in **Discussion**.
246

Table 1B Summary Geometry TCR:pHLA-DR

PDB	CDR2[1]	CDR3	σ[¶]	λ	ω	φ	k[‡]	ρ (Å)[◊]	dV[‡]
1J8Hα[†]	A52CB:E69CD 3.54	E102CD:A61CB 4.07	35.67	134.93	86.88	18.17c 17.15m	1.06	29.77	740
1J8Hβ	D51OD1:K67NZ 2.54	P100CG:L67CD1 3.43	47.99	109.62	84.91	25.85c 25.95m	0.996	27.91	1734
1FYTα	A52CB:E69CD 4.42	K103NZ:Q57OE1 4.36	37.11	133.00	86.29	18.81c 16.90m	1.11	29.70	826
1FYTβ	D51OD1:K67NZ 2.67	P100CB:D66CB 4.96	47.87	109.97	85.82	26.03c 23.44m	1.11	28.04	1747
3T0Eα	L50CD1:T106CG2 3.65	K96NZ:Q57OE1 4.06	38.06	132.31	88.95	19.87c 21.36m	0.930	29.99	855
3T0Eβ[‡]	E50OE1:K67NZ 3.77	N100CB:Q99CD 4.08	44.41	110.49	76.75	22.00c 20.98m	1.05	24.63	1024
4H1Lα	F50CD2:Q70CG 3.17	T98OG1:E55OE2 4.32	44.40	125.30	94.81	24.75c 19.78m	1.25	32.62	1782
4H1Lβ	I49CD1:L60CB 3.30	Y97CZ:D66CB 3.77	41.83	117.27	75.97	19.98c 18.84m	1.06	27.16	1198
1ZGLα	K51NZ:D76OD1 6.91	S91OG:E55OE2 2.90	51.82	117.60	85.36	26.11c 22.11m	1.18	33.78	2831
1ZGLβ	N51CG:A64CB 3.70	R99NH2:D66OD2 2.57	46.60	108.58	69.79	20.96c 14.06m	1.49	25.67	1281

[†] Same TCR, peptide and DRA (DRB1-4) as 1fyt (DRB1*0104).

[‡] DRB1 numbering is +29 from conventional due to attached peptide; Q93=Q64; L96=L67; D95=D66; Q99=Q70; T106=T77.

[1] Same as Table 1A legend

247
248

Table 1C Summary Geometry TCR:pHLA-DR

PDB	CDR2[1]	CDR3	σ[¶]	λ	ω	φ	k[‡]	ρ (Å)[◊]	dV[‡]
6CQLα[¶]	L58CD1:A73CB 3.85	A105N:E55OE1 3.00	37.14	133.17	93.59	20.41c 23.24m	0.878	29.01	746
6CQLβ	S66OG:Q57NE2 3.01	R108NH2:Y60OH 2.93	32.68	129.70	70.67	14.22c 18.95m	0.750	25.56	542
6CQNα	L58CD2:T77CG2 3.73	A105N:E55OE1 2.89	36.31	134.45	92.61	19.69c 22.34m	0.881	28.98	690
6CQNβ	S66OG:Q57NE2 3.17	L109CD1:D66CB 3.73	33.40	128.57	70.09	14.45c 23.56m	0.613	25.68	586
6CQQα	L58CD2:A73CB 3.78	A105N:E55OE1 2.88	35.95	134.32	89.29	18.85c 24.02m	0.785	28.97	693
6CQQβ	S66OG:Q57OE1 3.64	R108NH2:Y60OH 2.97	34.88	127.12	72.65	15.64c 22.76m	0.687	26.34	800
6CQRα	T57OG1:Q70NE2 3.17	A105N:E55OE1 3.03	36.52	133.75	87.44	18.75c 23.44m	0.800	28.99	718
6CQRβ	E60OE2:K67NZ 3.07	R108NH2:Y60OH 3.11	36.34	124.36	69.70	15.76c 29.07m	0.542	25.95	746

[¶] Same TCR (F24) and peptide and DRA (DRB1-11) as 6cqq (DRB1-15) and 6cqr (DRB1-1); same pMHC as 6cqn (TCR, F5; CDR3β G108R).

[1] Same as Table 1A legend

249
250

Table 1D Summary of TCR:pHLA-DQ Geometry

PDB	CDR2[1]	CDR3	σ[¶]	λ	ω	φ	k[†]	ρ[◊]	dV[#]
5KSAα[†]	Y57CE1:R7OCG **3.71**	Y113OH:D55OD2 **2.64**	36.16	134.95	94.04	15.90c 19.87m	*1.25*	28.76	651
5KSAβ	Y57CE2:A64CB **4.13**	Y114CD1:E66CG **3.59**	40.26	119.15	71.38	18.03c 23.32m	*0.773*	26.02	939
5KS9α	S58OG:D76OD2 **2.65**	N114ND2:T61OG1 4.24	41.63	128.32	92.51	22.66c 16.96m	*1.34*	28.65	918
5KS9β	R66NE:Q57OE1 **3.14**	D114OD2:E66OE2 **3.67**	33.80	126.25	67.09	14.17c 18.03m	*0.786*	23.57	465
4OZFα[‡]	H55NE2:R70NH2 **2.89**	A113CB:F58C2 5.05	40.39	131.50	111.69	26.24c 29.44m	*0.891*	33.17	1397
4OZFβ	N63OD1:K67NZ **3.50**	D114OD1:R70NE **3.73**	37.47	120.69	66.87	15.84c 25.68m	*0.617*	23.16	554
4OZGα	H55NE2:E69OE1 **2.93**	A115CB:Q57CB 4.11	45.55	124.99	100.43	26.82c 26.64m	*1.01*	31.91	1657
4OZGβ	Y55OH:Q57OE1 **3.35**	Y117OH:D66OD2 **2.72**	30.93	128.29	61.35	11.92c 31.10m	*0.383*	21.22	276
4OZHα	H55NE2:E69OE1 **2.79**	A110CB:D55CB **3.83**	48.60	121.57	102.66	29.32c 26.64m	*1.10*	32.35	2030
4OZHβ	Y55OH:Q57OE1 **2.86**	Y117OH:D66OD2 **3.21**	29.94	130.19	61.67	11.53c 32.11m	*0.359*	21.47	267
4OZIα	Q55OE1:R70NH1 **2.78**	S111OG:D55OD1 **3.21**	44.59	125.38	110.06	28.87c 19.13m	*1.51*	31.72	1588
4OZIβ	T66OG1:Q57OE1 4.08	Y115OH:D66OD1 **2.56**	38.48	116.06	66.45	16.55c 23.76m	*0.700*	22.10	549
4GG6α[¶]	L57CD2:A73CB **3.98**	R110CD:F58CD1 **3.71**	38.77	132.70	98.93	22.37c 25.38m	*0.881*	30.87	982
4GG6β	N58ND2:H68ND1 **3.94**	T113OG1:R70NE **2.95**	35.96	124.32	68.59	15.39c 22.73m	*0.677*	25.02	645

[†] Bel602 TCR same germline (TRAV 20*01/TRBV 9*01) as 5ks9 (Bel502); MHC (DQ8.5) different from 5ks9 (DQ8); peptide different [glia-gamma-1 (1-11)] from 5ks9 [glia-alpha-1 (1-16)].

[‡] Same MHC (DQ2) and peptide [glia-alpha-2 (1-13)] as 4ozg and 4ozh; 4ozi has same MHC (DQ2) with different peptide [glia-alpha-1 (1-13)]; all have different TCR.

[¶] Same MHC (DQ8) as 5ks9.

[1] refer to Table 1A legend.

251
252
Accordingly, the *mean dV* of these TCR V-domains was 955 ± 507 Å^3, excluding the unusually
large *dV* of the 1ZGL Vα (see below). Importantly, with the exclusion of 1ZGL (an apparent
outlier) there is a *linear relationship* between calculated pitch (φ_c) and the *dV* triple integral (see
plotted values from Tables 1A–1D in the **Suppl.1.II**)—corresponding to the equation: *y = 83.60x*
− 719.40; where *R^2 = 0.900*, by *linear regression* analysis. The lowest *dV* structures have the
lowest calculated *pitch* values. Overall, it says something about TCR:pMHC that may not be
intuitive—that a "flush" V-domain geometry could limit its capacity to find a compatible binding
interface with side-chains involving the aforementioned α-helix regions.
261
262   **CDR3:pMHC Contacts**
263

264 Particularly close TCR contacts with an autologous MHC during thymic development could
265 theoretically enhance positive selection [30]. However, some of the CDR2/α-helix interface
266 measurements (Tables 1A–1D) could suggest such TCR are not selected. Alternatively, we know
267 close CDR2β contacts with monomorphic DRA must contribute to linkages between DRB1 and
268 autoimmunity [15], but considerations here cannot directly exclude negative selection that
269 deletes 'forbidden' clones specific for self-peptide:self-MHC complexes [4]. All these structures,
270 even those with the highest resolution, are subject to limitations of crystallography [15]; and
271 indeed, we compute the H-bonds based on these coordinates. Nevertheless, we sought a
272 correlation between relative dV and contact distances for hydrogen bonds of the five
273 components; see Tables 2A–2C. For example, asking if particularly strong (close) H-bonds are
274 linked to a restricted dV?

276 **2IAM-Vα**

$$\lambda = 116.86°$$
$$\phi = 180 - 116.86 = 63.14°$$
$$\rho = 29.55\,\text{Å}$$
$$r = \sin\phi\,(\rho) = 0.892\,(29.55) = 26.36\,\text{Å}$$
$$\pi \cdot r / 4 = 3.1\overline{4}\,(26.36)/4 = 20.70\,\text{Å}$$

$$\iiint_v dV = \int_0^{63.14}\int_0^{20.70}\int_0^{29.55} \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi = \int_0^{63.14} 20.70\left[\frac{(29.55)^3}{3}\sin\phi\right] d\phi \quad \text{where,} \int\sin\phi = -\cos\phi$$

$$= \int_0^{63.14}\int_0^{20.70} \frac{1}{3}\rho^3 \sin\phi \Big|_{\rho=0}^{\rho=29.55} d\theta\, d\phi = -20.70\left[\frac{(29.55)^3}{3}\right]\cos\phi \Big|_{\phi=0}^{\phi=63.14} \quad \text{where, } \cos 0 = 1$$

$$= \int_0^{63.14}\int_0^{20.70} \frac{(29.55)^3}{3}\sin\phi\, d\theta\, d\phi = .452\left[-20.70\left[\frac{(29.55)^3}{3}\right]\right] - 1\left[-20.70\left[\frac{(29.55)^3}{3}\right]\right]$$

$$= \int_0^{63.14}\left[\frac{(29.55)^3}{3}\sin\phi\right]\theta \Big|_{\theta=0}^{\theta=20.70} d\phi = -80,475 - \left[-178,042\right] = 97,567 \quad \text{where, multipl. } x\,\frac{\pi}{180}$$
$$\text{corrects for angular integrand of } \phi$$

$$\therefore \iiint_v dV = 1,702.87\,\text{Å}^3$$

278 **2IAM-Vβ**

$$\lambda = 130.15°$$
$$\phi = 180 - 130.15 = 49.85°$$
$$\rho = 22.12\,\text{Å}$$
$$r = \sin\phi\,(\rho) = 0.764\,(22.12) = 16.90\,\text{Å}$$
$$\pi \cdot r / 4 = 3.1\overline{4}\,(16.90)/4 = 13.27\,\text{Å}$$

$$\iiint_v dV = \int_0^{49.85}\int_0^{13.27}\int_0^{22.12} \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi = \int_0^{49.85} 13.27\left[\frac{(22.12)^3}{3}\sin\phi\right] d\phi \quad \text{where,} \int\sin\phi = -\cos\phi$$

$$= \int_0^{49.85}\int_0^{13.27} \frac{1}{3}\rho^3 \sin\phi \Big|_{\rho=0}^{\rho=22.12} d\theta\, d\phi = -13.27\left[\frac{(22.12)^3}{3}\right]\cos\phi \Big|_{\phi=0}^{\phi=49.85} \quad \text{where, } \cos 0 = 1$$

$$= \int_0^{49.85}\int_0^{13.27} \frac{(22.12)^3}{3}\sin\phi\, d\theta\, d\phi = .645\left[-13.27\left[\frac{(22.12)^3}{3}\right]\right] - 1\left[-13.27\left[\frac{(22.12)^3}{3}\right]\right]$$

$$= \int_0^{49.85}\left[\frac{(22.12)^3}{3}\sin\phi\right]\theta \Big|_{\theta=0}^{\theta=13.27} d\phi = -30,879 - \left[-47,875\right] = 16,996 \quad \text{where, multipl. } x\,\frac{\pi}{180}$$
$$\text{corrects for angular integrand of } \phi$$

$$\therefore \iiint_v dV = 296.64\,\text{Å}^3$$

280 *for all (n = 38) triple-integral solutions, see **Suppl.1.I**.

Table 2A Possible H-bonding networks in TCR:pHLA-DR complexes

PDB	CDR3α:pep	Nn MHC:pep§	CDR3β:pep	pep:pep	α:α, α:β, β:β	MHC:MHC	α/β:MHC
2IAM*	none	A1N62ND2:L29O 2.0 A31N:A1N62OD1 2.2 A1N69ND2:A32O 2.1 V34N:A1N69OD1 1.8 B1R71NH1:N30O 2.1 B1R71NH2:N30O 2.0 N30ND2:B1Q70OE1 2.4	T95OH:A31O 1.7 H96NE2:N30O 2.8 H96NE2:A31O 2.8	none	αA93N:αQ91O 2.3 αQ94NE2:αA93O 2.1 αQ94NE2:αI90O 2.8	B1R71NH1: Q70OE1 2.2	αQ91NE2:B1T77O 2.2 A1G58N:αG92O 2.4 βT98N:B1D66OD1 2.0
2IAN (no H's)	none	A1N62ND2:L29O 3.0 A1N69ND2:A32O 3.1 V34N:A1N69OD1 3.0 B1R71NH1:N30O 2.9 B1R71NH2:N30O 2.8	T95OH:A31O 2.5 H96NE2:N30O 3.8 H96NE2:A31O 3.8	none	αG92N:αI90O 3.3	B1R71NH1: Q70OE1 3.2	αQ91NE2:B1T77O 2.9 βT98OG1:D66OD1 2.7
4E41	I28N:G91O 1.9	A31N:A1N62OD1 1.7 A1N62ND2:L29O 2.0 A1N69ND2:A32O 2.1 V34N:A1N69OD1 2.0 B1R71NH1:N30OD1 2.1 B1R71NH2:N30O 1.8 B1Q70NE2:N30O 2.8	R95NH1:I28O 1.9	none	βR95NH2:αG91O 2.0 βR95NH2:αT93O 2.4 βR95NE:αD89OD1 2.5	none	βT97N:D66OD2 1.9

§ Nn, nearest neighbour; shown possible H-bonds, donor → acceptor; colour-code: ≤ 2.5, red; 2.6 to 2.9, mauve; ≥ 3.0, green.

* Same TCR (E8) and MHC (DRA , DRB1*0104; pep I28) as 2ian (pep T28); same pMHC as 4e41 (TCR, G4).

282
283

Table 2B   Possible H-bonding networks in TCR:pHLA-DR complexes

PDB	CDR3α:pep	Nn MHC:pep§	CDR3β:pep	pep:pep	α:α, α:β, β:β	MHC:MHC	Nn α/β:MHC
1J8H†	none	A1N62ND2:Q311O 1.9 / A1N69ND2:L314O 1.8 / L316N:A1N69OD1 1.7 / B1Q70NE2:Q311OE1 2.1 / N312ND2:B1Q70OE1 2.1	K315N:G980O 2.0	none	αN99ND2:βL99O 2.2 / βY104N:βL99O 1.9	none	none
1FYT	none	A1N62ND2:Q311O 2.0 / T313N:A1N62OD1 1.9 / A1N69ND2:L314O 2.1 / L316N:A1N69OD1 1.7 / B1R71NH1:N312OD1 2.1 / B1R71NH2:N312O 1.9 / B1Q70NE2:Q311OE1 2.3	N312ND2:T97O 2.0 / K315N:G98O 2.1	N312ND2:T313O 2.7 / Q311NE2:K310O 2.8	αN99ND2:βL99O 2.0 / βY104N:βL99O 2.1	B1R71NH1:Q70OE1 2.0	none
3T0E‡	A5N:G92O 2.4	A1N62ND2:E6O 2.4 / R9NH1:B1Q93OE1 2.3 / R9NH1:B1D95OD1 2.8 / R9NH1:B1D95OD2 2.7	S9BN:Q8O 2.5 / R9NH2:N100O 2.2	G7N:E6OE1 2.6	βR95NH2:αN95OD1 1.8 / βY99N:βS98OG 2.1	B1L96N:D95OD1 1.7	βS98OG:A1N62OD1 1.4 / B1Q99NE2:βN100O 2.6
4H1L	none	A1N62ND2:N4O 1.9	none	C3SG:N4O 2.8	αN97ND2:βT98OG1 2.8	none	βR94NH2:A1N62OD1 1.6
1ZGL (no H's)	V1N:D99OD1 3.0 / H2N:D99OD2 3.3	A1N62ND2:I7O 3.2 / N6ND2:A162OD1 3.0	none	none	αK103N:βT102OG1 3.3 / αS93OG:βT102O 3.2 / βR99NH1:βE103OE1 2.8	none	αS101OG:A1E55OE2 2.9 / αS101N:A1E55OE2 3.1 / βR99NH2:B5D66OD2 2.6 / βR99NH2:B5D66O 2.8 / βV100N:B5D70OD1 3.3

§Nn, nearest neighbour; shown possible H-bonds, donor → acceptor; colour-code: ≤ 2.5, red; 2.6 to 2.9, mauve; ≥ 3.0, green.

†Same TCR, peptide and DRA (DRB1-4) as 1fyt (DRB1*0104).

‡DRB1 numbering is +29 from conventional due to attached peptide; Q93=Q64; L96=L67; D95=D66; Q99=Q70.

284
285
We examined the structures in *Swiss-PDB Viewer/Deepview-v4.1* (www.wpdbv.vital-it.ch); in
Swiss H-bonds can be computed after hydrogens are added based on coordinates; complexity
of certain structures (2IAN, 1ZGL) precluded computing hydrogens, and the shown H-bonding
distances are thus ≈ 1 Å larger (Tables 2A, 2B). Here, an obvious feature of all the complexes is
that there are two principal foci of H-bonding; one involving N62 of DRA (A1) (shaded *orange* in
the tables), which may include a separate secondary grouping at N69 (*light-orange* shade), and
the DRB1 (B1) centre involving R71 (*green* shades). Note, R71 is *polymorphic*, so structures
involving allotypes/isotypes that have a different a.a. at pos. 71 (1J8H, 3T0E, 1ZGL, 6CQQ &
4H1L) involve a different a.a. (1J8H, 3T0E, 6CQQ), or simply do not have a corresponding beta-
chain H-bonding centre (4H1L & 1ZGL). Superimposed upon the H-bonding centres are the H-

bonds between the CDR3α and peptide, the CDR3β and peptide, any peptide to peptide H-bonding, any intra- or inter- bonds between the TCR V-domains, any MHC to MHC H-bonding, and direct H-bonds between TCR V-domains and the MHC (Tables 2A–2C, *columns*).

Table 2C   Possible H-bonding networks in TCR:pHLA-DR complexes

PDB	CDR3α:pep	Nn MHC:pep§	CDR3β:pep	pep:pep	α:α, α:β, β:β	MHC:MHC	Nn α/β:MHC
6CQL¶	R95NH1:A105O 2.0 R95NH1:G106O 2.6 R95NH1:N107OD1 2.3 R95NH2:N107OD1 2.0	A1N69ND2:E97O 2.2 A99N:A1N69OD1 1.8 B1R71NH1:R95O 2.2 B1R71NH2:R95O 2.0	A110N:E97OE1 1.9 G111N:E97OE2 1.9	Q98N:E97OE1 2.0	βG112N:βL109O 2.7	B1R71NH2:D70OD2 2.4	αA105N:A1E55OE1 2.1
6CQN	R95NH1:A105O 2.1 R95NH1:G106O 2.6 R95NH1:N107OD1 2.2 R95NH2:N107OD1 1.9	A96N:A1N62OD1 1.9 A1N62ND2:L94O 1.9 A1N69ND2:E97O 2.2 A99N:A1N69OD1 1.8 B1R71NH1:R95O 2.0 B1R71NH2:R95O 1.9	A110N:E97OE1 1.7 G111N:E97OE2 2.0	Q98N:E97OE1 2.2	βG112N:βL109O 2.5	B1R71NH2:D70OD2 1.9	αA105N:A1E55OE1 2.0
6CQQ	R95NH1:G106O 2.4 R95NH1:N107OD1 2.3 R95NH2:N107OD1 1.8	A96N:A1N62OD1 1.8 A1N62ND2:L94O 1.8 A1N69ND2:E97O 2.1 A99N:A1N69OD1 2.0 S100N:B1D57OD1 2.6	R95NH2:A110O 2.0 A110N:E97OE1 2.1 G111N:E97OE2 2.3	Q98N:E97OE1 2.2	βQ115NE2:αN114O 2.1	none	αA105N:A1E55OE1 2.0 βR108NH2:B1D66OD1 2.2 βR108NH2:B1D66OD2 2.8
6CQR	R95NH1:A105O 2.2 R95NH1:N107OD1 2.1 R95NH2:N107OD1 2.0	A96N:A1N62OD1 1.9 A1N62ND2:L94O 2.1 A1N69ND2:E97O 2.2 A99N:A1N69OD1 1.8 B1R71NH1:R95O 2.4 B1R71NH2:R95O 1.8	R95NH2:A110O 2.2 A110N:E97OE1 1.8 G111N:E97OE2 2.2	Q98N:E97OE1 2.4	βQ115NE2:αN114O 2.1	none	αA105N:A1E55OE1 2.0

§Nn, nearest neighbour; shown possible H-bonds, donor → acceptor; colour-code: ≤ 2.5, red; 2.6 to 2.9, mauve; ≥ 3.0, green.

¶Same TCR (F24) and peptide and DRA (DRB1-11) as 6cqq (DRB1-15) and 6cqr (DRB1-1); same pMHC as 6cqn (TCR, F5; CDR3β G10BR).

In all cases, the CDR3α:peptide bonds are either non-existent, or isolated from the MHC:peptide centres. In 4E41, 6CQQ and 6CQR, CDR3α may link to CDR3β through both bonding nearby with peptide (*blue* shading). For 2IAM, 1J8H, and 1FYT there are bonds between intra-V (2IAM), or intra- and inter-V (1J8H, 1FYT), but these are not linked to the peptide (different *blue* shade). N62 is connected to a peptide:peptide H-bond (4H1L), or to an inter-Vβ bond (3T0E); in both cases these potential networks extend to the Vβ directly binding with N62. 1ZGL is the only structure with an inter-V bond potentially linked to a direct V-domain interaction with MHC, while 4H1L is unique in not showing any DRB H-bonding involving the CDR3 (recall these two structures involve non-DRB1 isotypes). In 6CQL and 6CQN there is a potential DRB1:peptide link with an MHC:MHC bond, but this does not involve the

312   TCR. The 6CQ series all have CDR3β:peptide bonds also involving a peptide:peptide bond; this
313   is not found in any of the other complexes (*mauve* shading). Finally, and most significantly here,
314   in 2IAM, 2IAN, 1FYT and 3T0E the DRB1:peptide centre is potentially connected through a
315   CDR3β:peptide bond with an MHC:MHC H-bond (Table 2A & 2B, *green* shades).
316
317   **CDR3:pMHC Chemistry**
318
H-bonds computed in *Swiss* are *potential* H-bonds, and to assess a given network the chemistry
of each bond must be examined [28, 31]. The analyses of Tables 2A–2C indicated that a
potential H-bonding network where CDR3 β is connected through peptide-bonding to
DRB1:peptide and DRB1:DRB1 H-bonds might correlate with the *highly-restricted dV* of Vβ in
2IAM (297 Å^3) and 2IAN (270 Å^3). However, these networks would need to be distinct from this
same type of potential network observed in 1FYT and 3T0E (Table 2B), which have *d*V for Vβ of
1747 Å^3 and 1024 Å^3, respectively (Table 1B). Shown is our analysis of the 2IAM-Vβ versus
1FYT-Vβ for contacts made by each CDR3β (Fig. 3). For 2IAM-Vβ a potential H-bonding network
involving CDR3β a.a. Y95 and H96 focused on central peptide a.a. N30 and A31 is apparent; and
DRB1 a.a. Q70 and R71 appear in this same network, while the β-sheet a.a. E28 (with R71:NE
and R71:NH2 bonds of precisely the same distance) cancel out. More specifically, there are
three *potential* H-bonds with the N30:O (at R71:NH1, R71:NH2, and H96:NE2); Y95 appears to
H-bond to A31:O, and there is also a potential bond between Q70:OE1 and R71:NH1 (Figs. 3A &
D). By contrast, 1FYT-Vβ displays single H-bonds between T97:O and N312:ND2 (peptide), and
G98:O and K315:N (peptide). The DRB1, R71:NH2 is involved with H-bonding to N312:O, and
R71:NH1 has possible H-bonds with N312:OD1 and with Q70:OE1 (Figs. 3B & E). Since 1FYT and
1J8H only differ via DRB1 alleles, we compared H-bonding for the two (Fig. 3B *vs*. 3C; 3E *vs*. 3F).
Illustrating the importance of the *Swiss* computation, in Figure 3C we anticipated that the R71K
polymorphism would effect three H-bonds—involving K71 in a manner similar to R71. However,
as is shown in Figure 3F, R71K actually shifted peptide contacts to Q70; in-fact, K71 did not
show any H-bonds. Importantly, this confirms the change in peptide conformation between the
two structures as originally reported [10], and suggests that alloreactivity of 1FYT/1J8H involves
the single G98:0 to K315:N bond in maintaining the ≈ 1740 Å^3 *d*V of the two Vβ (Table 1B). Still,
we had to examine the chemistry of each possible bond to arrive at feasible mechanisms;
arguably, this is necessary to understand binding.
344

347 Figure 3 **CDR3 H-bonding networks of 2IAM-Vβ (A & D), 1FYT-Vβ (B & E) and 1J8H-Vβ (C & F).** *PDB* files
348 analysed with *VMD* (top) and *Swiss-Deepview* (bottom). In *Swiss*, hydrogens added and H-bond
349 distances computed; see *dotted* lines in *green*. Involved side chains shown by atom in *licorice* (top) or
350 *stick* (bottom); The 2IAM CDR3β in *gold* (A) and *red* (D); 1FYT/1J8H CDR3β in *mauve* (B, C, E, F); and
351 peptides in *green*.

H-Bonding Mechanisms

Shown in Figure 4A is a suitable reaction mechanism for the 2IAM CDR3β binding to the peptide
(triosephosphate isomerse, 15mer)-HLA-DR complex (*PDB* ref., 12). When considering the
probability of hydrogen bonds in a mechanism, we draw all the possibilities using *ChemSketch*
(ACD Labs; www.acdlabs.com/resources) and traced electron pathways using standard
evaluations of electron configuration [28, 31]. Thus DRB1 (B1) Q70:OE1 would be in an H-bond
with B1R71:NH1 in the absence of the peptide, with the downstream effect of relieving the
charge on the B1R71:NH2. Bound peptide has the same effect on R71:NH2, where the N30:O
attacks the R71:NH2 (*orange* arrows). Here, the peptide would indirectly break the H-bond
between Q70 and R71 (*orange* blocks). When the TCR binds, the β H96:NE2 charge is
preferentially attacked by the N30:O (*purple* arrows); this reverses the previous bond between
the peptide and DRB1 (*purple* block) and has the downstream effect of re-forming the intra-
MHC bond between Q70 and R71 (*purple* blocks on the *orange* blocks). Note, this is the only
obvious mechanism that relieves both the H96 and R71 charges, and would be favoured over an
A31:O attack on H96:NE2 by the neighbouring βY95 H-bond with A31:O (Fig. 4A). Therefore,
just three of the possible H-bonds are predicted; and the overall effect is a *charge-relay* [28, 32,
33] between the MHC, the peptide, and the TCR.

Physiologically, the charge on R71:NH2 could be initially neutralized via the neighbour
Q70:OE1—subsequently replaced during processing by CLIP, and at HLA-DM exchange, with
antigenic peptide (ref. 5). During T-cell conjugation with the antigen presenting cell, the TCR

would replace the A30:O to R71:NH2 attack with an A30:O to H96:NE2 attack, which *relays* the
charge back to R71:NH2—subsequently, neutralized again by Q70:OE1. While the hallmark
charge-relay network of the 'catalytic triad' within *serine proteases* stabilizes formation of a
covalent acyl-enzyme intermediate [28]; here, a relay is utilized in a similar (albeit, noncovalent)
role. Undoubtedly, this would still be a stabilised transition-state [31], *i.e.*, given that these H-
bonds are de-localised across 3/5 components of the structure [32, 33].

382
383
Figure 4   **H-bonding mechanism for CDR3 β binding to peptide and MHC.** *ChemSketch* drawn
mechanisms are shown for 2IAM (A) and 1FYT (B) and were based on analyses in *VMD* and *Swiss* (Fig. 3).
Not necessarily to scale. Electrons shown with arrows (*purple*, involving the TCR; *orange*, related to the
peptide). Possible H-bonds are shown in *blue* (darker shade for key TCR bond) and are crossed with a
color-coded *block* line if not probable (see text).

By contrast, Figure 4B shows a suitable mechanism for the analogous CDR3β binding reaction
with the influenza HA peptide-HLA-DR complex of 1FYT. The differences are the peptide and
TCR, as again both 2IAM and 1FYT are DRB1*0104 structures. Similarly the peptide would
replace the internal B1Q70:OE1 to B1R71:NH1 with an H-bond directly with the charged
R71:NH2 via the peptide N312:O (*orange* arrows). Again, the TCR *could* replace this bond with
an attack by the CDR3β (here, T97:O on N312:ND2; *purple* arrows). However, the downstream
effect of such an attack in this case is a partial charge on N312:OD1 and an attack on R71:NH1.
Note this would restore the charge on R71:NH2, effectively breaking the peptide-MHC bond.
Also, despite what looks like proximity in the *ChemSketch* diagram, T97 is actually far removed
from R71 (Fig. 3E) such that an attack directly on R71:NH2 is not possible. Thus, because of a
remaining charge on R71:NH2, the T97:O attack on N312:ND2 (with its downstream effects)
would *not* be favoured (*orange* blocks). While similarly to 2IAM there is a neighbouring H-bond
(in this case to Q311), the net loss of a CDR3β H-bond clearly distinguishes the two mechanisms,
and shows that the 1FYT CDR3β does not favour a charge-relay mechanism; *i.e.*, would *not* have
the stabilized CDR3β tether at B1R71—ostensibly, the chemistry driving restricted *d*V in 2IAM.
Note that the 2IAN-Vβ mechanism appears to be similar to that of 2IAM (structures differ by a
single, sufficiently distant, a.a. of the peptide).   Next, 3T0E also showed the
TCR:peptide:MHC:MHC type of potential network by contacts analysis (Table 2B). Figure 5A
shows the a.a. involved in this potential network using *Swiss*, and the proposed mechanism is
shown in 5B. Note that the DRB1-4 allotype of 3T0E has the R71K polymorphism, and that the

involved peptide a.a. is R9. In terms of distinguishing the mechanism from 2IAM, the question
is whether the CDR3β N100 H-bond to R9 could exist with a B1:peptide H-bond. In fact, if the
Vβ closest contact N100:O is to attack the positive charge of R9:NH2 (purple arrows), it would
have to compete (indirectly) with the H-bond of B1Q93:OE1 to R9:NH1 (orange arrows).
Together with the B1Q99:NE2 possible bond to the same N100:O, the B1Q93 bond would be
favoured; thus, the 3T0E mechanism is not similar to 2IAM, most notably because CDR3β is not
likely to form an H-bond with the peptide. Note that in the related structure, *PDB* 3O6F, this
conclusion holds as well, where B1D95:OD2 has a possible H-bond to R9:NH2 at 1.6 Å; again,
not favouring a CDR3β N100:O bond to the peptide [34].
419

420
421
Figure 5 **H-bonds and H-bonding mechanism for 3T0E CDR3β.** *Swiss* was used as before (non-involved
a.a. deselected for simplicity) to compute possible H-bonds for 3T0E (A). Mechanism for 3T0E TCR
binding via CDR3β to peptide and DRB1 (B) shows possible H-bonds in *blue*, with improbable bonds
blocked with *grey* lines (*ChemSketch*). Electron flow shown with *purple* arrows (TCR), or orange (MHC).
CDR3β in *yellow* (top) or a.a. labelled *yellow* (bottom); DRB1 in *mauve* (top) or labelled a.a. in *pink*
(bottom); peptide by atom (top) or a.a. labelled *grey* (bottom).
428

429   **DQ H-Bonding Mechanisms**
430

As shown in Table 1D, we noted the highly-restricted *d*V in Vβ for two of the TCR:pHLA-DQ
structures (4OZH & 4OZG; 267 Å^3 and 276 Å^3, respectively) [35]. Here, the control structure is
4OZF (Vβ *d*V, 554 Å^3), where the pMHC is the same; in-fact, these three structures only differ by
their "non-germline" TCR, *i.e.*, by their CDR3 regions (*all 38 IMGT-computed CDR3 junctions*
shown in **Suppl.1.III**). Accordingly, we analysed H-bonding with *Swiss-Deepview* as before (Fig.
6). While very similar, we noted the absence (Fig. 6A & B) of the DQA1, T61:OG1 role in H-
bonding to Vβ, R109:NH1 (Fig. 6C). This is accomplished in 4OZH by preferential H-bonding to
Vβ a.a. S110 (Fig. 6A), or sterically in 4OZG by Vβ a.a. F110 (Fig. 6B). Thus, only in 4OZF is
T61:OG1 free to H-bond to the pivotal arginine (CDRβ, R109); note, this shift from R109
contacting MHC at a.a. N62 to T61 indirectly disrupts the MHC:MHC component of the charge-
relay mechanism (Fig. 7C vs. 7A & B). Crucially, the chemistry of S110 versus F110 versus A110-
L111-A112-A113 in the these TCR is ultimately the result of somatic genetic construction of the
CDR3β loop [**Suppl.1.III**; ref. 4].

445
446
Figure 6 **H-bonds for three different TCR CDR3β, (from germline-identical TCR) on the same pHLA-DQ.**
(A) 4OZH, showing S110:T61 H-bonds (*orange* arrow), (B) 4OZG, showing bulky F110 (orange arrow);
both of which could preclude (C) , the conserved R109 H-bonds to T61 (*orange* star). *Swiss-Deepview*
analysis as previously; DQ and CDR3β backbones in *white* ribbon; peptides in *yellow* wire; involved side
chains in *CPK*; computed possible H-bonds and distances in *green*.

Again, we can envision the mechanism sequentially. First, there is internal MHC:MHC H-
bonding between N11 and N62 (*purple* arrows, Fig. 7A & B). This facilitates the TCR binding to
peptide via a P8:O attack on the R109:NE proton (*blue* arrows), and this leads to neutralizing
the charge on R109:NH2. Uniquely here (*purple* arrows, Fig. 7C), when the CDR3β is permissible
for the T61:OG1 to R109:NH1 H-bond, *i.e.*, the a.a., A110-L111-A112-A113 (4OZF), this appears
to preclude both N62:OD1 attack on the charged R109:NH2 and the MHC-MHC H-bond
(N11:OD1 to N62:ND1); compare in Fig. 7C to 7A & B). Thus, all three TCR (across DR and DQ)
which displayed highly-restricted *d*V, also shared the likelihood of a *charge-relay* mechanism of
CDR3β binding, where an MHC to MHC bond is involved. Interestingly, this finding may suggest
selection in *celiac disease* for CDR3β joints that can disrupt the N11-N62 H-bond, which is an
indirect result of T61 as the source of R109 neutralization. For example, via a Th1→Th2

functional (reduced inflammation) shift [21]; and it is interesting to consider under a thymic-
selection model based on a 'window' of TCR affinity [4, 21, 22, 30].

467
468

Figure 7 **H-bond mechanisms for three different CDR3β, (from germline-identical TCR) on the same**
pHLA-DQ. (A) 4OZH, showing S110:T61 H-bonds (*orange* arrows), (B) 4OZG, showing bulky F110; both
of which could preclude (C), the conserved R109 H-bonding to T61, as only observed for the nominal *d*V
V β of 4OZF. As previously, possible H-bonds in *blue*, with improbable bonds blocked with *colour-*
matched lines (*ChemSketch*). Electron flow shown with *blue* arrows (peptide:TCR), or *orange* arrows
(MHC:TCR), or *purple* arrows (MHC:MHC). CDR3β a.a. labels in *yellow*, peptide a.a. in *blue*, DRB1 a.a. in
mauve.

Discussion:

We used spherical coordinates to derive a unique solved triple integral for each of the 38 V-
domains (Figs. 1 & 2; **Suppl.1.I**). Using this approach the mean volumetric-density of a TCR V-
domain through the putative CDR2 scanning path was 1060 Å^3. As indicated, $\iiint_v d$V is a 'slice'
volume of a cone with the vertex at the 'across-the-groove' MHC α-helix, where maximal CDR2
scanning (*d*θ) calculates close to the actual range of α-helix distances between the most N-term.
and most C-term. CDR2 contacts (Tables 1A–1D). Also, overall geometry consistent with across-
the-groove *tethering* via CDR3 (first formulated for TCR-Vα:pHLA-A2; ref. 17) was again, broadly
apparent. Most importantly, here we found a linear relationship between V-domain *pitch*;
calculated by:

$$\varphi_c = k\varphi_m = [\sigma \div (\lambda + \sigma)]\, \omega$$

and V-domain "*d*V" calculated by:

$$\iiint_v dV = \int_0^{\sin^{-1}(r/\rho)} \int_0^{\pi \cdot r/4} \int_0^{z/\cos\phi} \rho^2 \sin\phi\ d\rho\ d\theta\ d\phi$$

491
492

This suggests that very-low pitch may "restrict" $d\,\theta$, *i.e.*, the CDR2-scanning function. In
retrospect, this confirms intuition (excluding the broader *TCR-CD3 complex,* of course) on
possibilities for a mechanism involving just the five components. Indeed, H-bonding chemistry
which effects relatively simple physics is a hallmark of quite diverse protein machinery [28, 29].
Also, TCR had *k* values that varied in either direction—indicating that conformation adjustments
might moderate flush and open pitch *without* much increasing or decreasing of the *d*V, *viz.*, $R^2 \approx$
0.900 (**Suppl.1.II**). Thus, pitch calculated from a given V-domain's *twist-tilt-sway* might be a

kind of "hidden" correlate of TCR-selection; although there is not available crystallography on
any thymic (selecting) ligands [36] to actually test this theory. For 2IAM, there are data for the
uncomplexed TCR, and the CDR3β backbone is displaced ≈ 3.4 Å upon binding pMHC, while
β Y95 moves ≈ 9.0 Å to form the H-bond shown (Fig. 3A; *PDB*, ref. 12); indeed, the key to
understanding the TCR seems to be in this relationship between two 'induced-fit' TCR
conformations [4, 28, 37]. What might be suggested here is a conservation/approximation of
*d*V for the V-domains when binding the *selecting* (thymus) and *activating* (peripheral) pMHC;
despite that a given pitch on the two (potentially) different ligands *can* change dramatically,
usually it does not. Obviously, rotation *per se* could change between the two structures,
wherein different pitch values could nevertheless lead to matched *d*V, and (perhaps) certain
shared thermodynamics for the two binding reactions across time and space with two different
ligands (see below).

Differences between MHC alleles with regard to transplantation, so-called 'permissive' versus
'non-permissive' haplotypes, could evoke an increased number of low-concentration peptides
that a non-permissive graft might present with a given (selecting) *d*V [38, 39].
Alternatively/historically, *intrinsic* TCR affinity (*i.e.*, equilibrium and rate binding constants
specified by the α/β TCR protein) has been used to explain thymic selection *and* MHC restriction
[4, 40–43]. However, as previously indicated, it is difficult to understand the intermediate
(window) nature of affinity that would result from thymic selection in the context of a
subsequent affinity-based, MHC-restriction mechanism; this is not mutually exclusive of
whether or not a physical force (from T-cell motility) is an obligate component of TCR activation
[7, 28, 29, 31, 42, 43]. Crucially, TCR affinity for pMHC is generally *micromolar* binding—weak
reactions by antibody standards [7, 44]. Related factors including the orientation of CDR2α, 3°-
structure of α-chain constant domains and inter-chain disulfides, differences in CDR3α packing
and the extra HV4, all likely contribute to pMHC-elicited monoclonal antibody Fab binding *not*
like TCR [4, 45, 46]. Simply put, if positive/negative selection is based on TCR-affinity, it would
be logical that the selected repertoire reflects an *optimum* TCR-affinity; and (it follows), that a
qualitatively different mechanism might drive MHC restriction. In these and other regards,
there is a *stereochemical* alternative to an affinity-limited binding reaction [31, 32].

Briefly, orientation is assumed to drive the TCR-pMHC reaction involving a high(er) energy
"scanning" conformer, because scanning (leading to a suitable CDR2:MHC interface) has the
effect of lowering the *transition-state* free energy, $\Delta G^{\ddagger}$ (*i.e.*, *Curtin-Hammett* control) [31, 32, 47,
48]. For example, in TCR like E8 (of 2IAM/2IAN) very little CDR2β scanning is apparently needed,
due to the effect [49] of the proposed *charge-relay* stabilised transition-state. Thus, *d*V is a
universal consequence of the CDR3's chemistry—but usually the reaction requires more CDR2-
scanning (*i.e.*, usually V-domains have larger *d*V values; Tables 1A–1D). One might envisage a
free-energy 'trough' for successful induced-fit rates [37]; one provocative corollary being its
establishment during thymic selection. Unfortunately, technology to directly investigate the
issue is also limiting [reviewed in ref. 50]. Recently, ensemble refinement of crystal data with
MD simulations has suggested conformational diversity in the microsecond range, but the
conformational changes implicated here-in would be on the millisecond to minutes scale; in this
regard, NMR of membrane-bound receptors may offer promise [50–54]

A dynamics mechanism driven by H-bonding networks is consistent with the knowledge that
different T-cell dose-response curves are found at the same TCR affinity for the same pMHC [40,
55–58]. For the three 4OZ structures, it appears that somatic genetics directly effects H-bonds
implicated in V-domain dynamics (*d*V), and (indirectly) germline:germline contacts between
CDR2 and the MHC α-helices (*d*θ) [24, 44, 49, 59–62]. Stabilization of H-bonding via charge-
relays is implicated for V-domains with very-low *pitch* angles, as specified by the *twist-tilt-sway*
(*viz.*, orientation) and *highly-restricted d*V. What can change is the *measured* pitch angle. Thus,
an orientation selected on a thymic pMHC-II could specify particular *d*V (characterized by the
calculated pitch); and *in theory*, this *d*V must be approximated on a peripheral pMHC-II, *i.e.*,
leading to 'recognition' of this pMHC-II during an immune response. The plasticity of pitch
during recognition (induced fit) ostensibly might allow certain *d*V recapitulations on pMHC-II
with different closest contacts resulting from a different (peripheral) peptide [4, 7, 15, 30].
Again, usually we see *k* values close to unity. While there are myriad downstream implications,
taken together, these data support that V-domain rotation and germline to germline contacts
between TCR and pMHC *both* depend upon CDR3 H-bonding with a highly-conserved, *across-*
the-groove MHC motif [17, 19, 24, 30]—in this case, the polymorphic a.a. Y60-W-N-X-Q-K/R-
D/E-X-L-E-X70 sequence spanning closest contacts of CDR3β's on DRB1/DQB1 (Tables 1A–1D).
Similar to the class-I, R65-motif [17, 63], this class-II motif is present in *Galago sp.* (NCBI Acc. No.
AAA96291) as well as both *tarsiers* and *lemurs*; suggesting conservation for at least 63 million
years [64].

Finally, while this paper was in review, Baker and co-workers used spherical coordinates to
calculate TCR *center of mass* variation in TCR:pMHC complexes [65]. Importantly, these
investigators do not consider Vα and Vβ independently, although previous work from Hoffmann
et al., had shown the angle between Vα and Vβ characterized different TCR in pMHC complexes;
allowing these researchers to group a panel of TCR into six different clusters linked to clonotype
[66]. Thus, as we have shown here, it is crucial to consider the two TCR domains independently
with regard to geometry related to binding. Moreover, we would argue it is the volume
element that most accurately describes "the probability of collisions" between each V-domain
and pMHC atoms [ref. 67]. We have developed our method based on *Euler*'s classic equations
for the direct purpose to use multi-variable calculus to predict *effective* V-domain binding
volumes; we feel this most directly approaches the basic math of dynamics in TCR Vα/Vβ
specificity.

Here, we propose standard biochemistry linked to the variation we observed in the volume
element (*d*V) between different V-domains. For example, we would suggest that one must
consider the second-order kinetics of H-bond forming reactions between CDR2β and DRA/DQA
α-helix, a.a. Q57–H68 (Tables 1A–1D). Where:

$$\text{rate of reaction} = -d[A]/dt = -d[B]/dt = k[A][B]$$

could be limited by the rotation dynamics of the Vβ domain (see earlier discussion, re: *Curtin-*
Hammett control). This might explain why force applied experimentally [54] influences TCR

specificity, *i.e.*, slowing dynamics simply allows for the concentration term of CDR2 β side-
chain(s) in 'collision' with α-helix side-chain(s). It is interesting to speculate that so-called,
'catch-bonds' are a possible outcome from this dynamics [7, 42, 43, 54]. Moreover, we have
introduced the idea here that rare Vβ (very-low *d*V/highly-restricted rotational volume) might
be somehow specialized by this dynamics. Obviously, whether V-domain *d*V influences TCR
signaling, downstream subunit associations, or indeed plays a role in TCR selections awaits
direct testing.
595

596 # Methods:
597

598 *Availability of data and materials:* *PDB* files are public and available at www.ncbi.nlm.nih.gov; all
599 analytics data are either in the paper, or supplementary materials; *Competing interests:* Dr. Murray and
600 *Xenolaüs Genetics LLC* declare no competing interests regarding the research, or its publication; *Author*
601 *Contributions:* J.S.M. did the research and wrote the manuscript.
602

603 ## *PDB* Analysis
604

VMD-1.9.1 software (www.ks.uiuc.edu) was used for *PDB* files downloaded via *NCBI*
(www.ncbi.nlm.nih.gov) from the *RSCB-PDB* (www.rscb.org); views normalized with the *VMD*
xyz-axis tool; alpha carbon (Cα) main-chains in *new cartoon*; all alleles named per *NCBI*
annotation. *Euler*'s methods (www.mathword.wolfram.com/EulerAngles.html) were the basis
for the specific angle analyses, as previously reported. Briefly, three angles corresponding to
the *twist*, *tilt*, and *sway* of each domain over the pMHC were measured from fixed Cα through
the 19 structures: (i) in-plane to the MHC-groove {*twist* = ω }, with (ii) displacements
perpendicular to the groove {*tilt* = λ}, including (iii) side-to-side variation {*sway* = σ}. The a.a.
positions used as coordinates for angular measures across structures were *fixed*; see previous.
The incline of a V-domain, {*pitch* = φ_m}, was approximated (calculated) by the equation: $\varphi_c = [\sigma$
$\div (\lambda + \sigma)] \omega$ (see also **Results**, Tables 1A–1D). Pitch was also measured by using the closest
determined CDR2 contact Cα for an angle across-the-groove to the V-domain closest CDR3
contact with an α-helix side-chain (vertex), then back to said CDR2 closest contact within the
opposite α-helix ($\approx$ 2-fold symmetry); angular value in degrees via the *VMD angle-label* tool.
Triple integrals for all 38 V-domains based on the fixed geometry were solved as described in
Results (**Suppl.1.I**). Linear regression analysis by *MS-Excel* (**Suppl.1.II**); and statistics by paired
two-tailed *Student*'s t-test (www.insilico-net/tools/statistics/ttest).
622

623 ## Contacts Analysis
624

All measures were performed with *Swiss* or *VMD-1.9.1* as is specified in **Results**. Closest
contacts in *angstroms* (Å) were determined by examining appropriate coordinates between
structures (computed in *Swiss*, or measured/computed in *VMD*). Individual atomic contacts
are named per software annotation. DRA chain contacts are *abbrv*. "A1" to avoid confusion
with single-letter a.a. code for alanine (A); potential H-bond networks are colour-coded as is
described in Table captions and the text.

CDR3 Joint Analysis

Nucleotide sequences for all CDR3 of TCR were specified from *PDB* files (**Suppl.1.III**). TCR a.a. sequences were reverse translated using the *SMS* tool at www.bioinformatics.org. These were then imported into *IMGT* algorithms for joint analysis (www.imgt.org). Amino acid sequences of resulting CDR3 joints were determined by the *IMGT* algorithm; consensus *IMGT* numbering.

H-Bonding Mechanisms

Hydrogen (H) bonds were estimated either with *VMD*, or (more accurately) with *Swiss*. In *Swiss*, H-bonds are computed after computing hydrogens to the structures. H-bond distances were used as a factor in determining suitable organic reaction mechanisms, where relevant side-chains were reproduced with *Chemsketch* (www.acdlabs.com/resources). Standard evaluations of electron configuration per relevant atoms were used to predict electron flows [28, 31].

Acknowledgements:

This work was possible because of dedicated crystallographers, molecular biologists and computer scientists providing public availability of coordinates, sequences and software; on the community of science, thank you.

Supplementary details:

[Suppl.1.pdf] **I.** Triple Integral Solutions for all V-domains
            II. Linear Regression Analysis of Calculated Pitch versus *d*V (all V-domains)
            III. *IMTG* CDR3 Joint Analysis (all V-domains)

References:

1. Allison JP, McIntyre BW, Bloch D. Tumor-specific antigen of murine T-lymphoma defined with monoclonal antibody. *J Immunol* (1982) **129**:2293–300.

2. Tonegawa S. Somatic generation of antibody diversity. *Nature* (1983) **302**:575–81. doi: 10.1038/302575a0

3. Agrawal A, Eastman QM, Schatz DG. Transposition mediated by RAG1 and RAG2 and its implications for the evolution of the immune system. *Nature* (1998) **394**:744–51. doi: 10.1038/29457

4. Janeway CA Jr. *Immunobiology*, 5th ed. New York: Garland (2001).

5. Wieczorek M, Abualrous ET, Sticht J, Álvaro-Benito M, Stolzenberg S, Noé F, Freund C. Major
Histocompatibility Complex (MHC) Class I and MHC Class II Proteins: Conformational Plasticity
in Antigen Presentation. *Front Immuno*l (2017) **8**:292. doi: 10.3389/fimmu.2017.00292

6. Nisonoff A, Hopper JE, Spring SB. *The Antibody Molecule*, 1st ed. New York: Academic Press
(1975).

7. Gee MH, Sibener LV, Birnbaum ME, Jude KM, Yang X, Fernandes RA, *et al*. Stress-testing the
relationship between T cell receptor/peptide-MHC affinity and cross-reactivity using peptide
velcro. *Proc Natl Acad Sci USA* (2018) **115**:E7369–78. doi: 10.1073/pnas.1802746115

8. Brown JH, Jardetzky T, Saper MA, Samraoui B, Bjorkman PJ, Wiley DC. A hypothetical model
of the foreign antigen binding site of class II histocompatibility molecules. *Nature* (1988)
332:845–50. doi: 10.1038/332845a0

9. Davis MM, and Bjorkman PA. T-cell antigen receptor genes and T-cell recognition. *Nature*
(1988) **334**:395–402. doi: 10.1038/334395a0

10. Hennecke J, Wiley DC. Structure of a complex of the human alpha/beta T cell receptor (TCR)
HA1.7, influenza hemagglutinin peptide, and major histocompatibility complex class II molecule,
HLA-DR4 (DRA*0101 and DRB1*0401): insight into TCR cross-restriction and alloreactivity. *J Exp*
Med (2001) **195**:571–81. doi: 10.1084/jem.20011194

11. Li Y, Huang Y, Lue J, Quandt JA, Martin R, Mariuzza RA. Structure of a human autoimmune
TCR bound to a myelin basic protein self-peptide and a multiple sclerosis-associated MHC class
II molecule. *EMBO J* (2005) **24**:2968–79. doi: 10.1038/sj.emboj.7600771

12. Deng L, Langley RJ, Brown PH, Xu G, Teng L, Wang Q, *et al*. Structural basis for the
recognition of mutant self by a tumor-specific, MHC class II-restricted T cell receptor. *Nat*
Immunol (2007) **8**:398–408. doi: 10.1038/ni1447

13. Yin L, Crawford F, Marrack P, Kappler JW, Dai S. T-cell receptor (TCR) interaction with
peptides that mimic nickel offers insight into nickel contact allergy. *Proc Natl Acad Sci USA*
(2012) **109**:18517–22. doi: 10.1073/pnas.1215928109

14. Yin Y, Wang XX, Mariuzza RA. Crystal structure of a complete ternary complex of T-cell
receptor, peptide-MHC, and CD4. *Proc Natl Acad Sci USA* (2012) **109**:5405–10.   doi:
10.1073/pnas.1118801109

15. Sharon E, Sibener LV, Battle A, Fraser HB, Garcia KC, Pritchard JK. Genetic variation in MHC
proteins is associated with T cell receptor expression biases. *Nat Genet* (2016) **48**:995–1002.
doi: 10.1038/ng.3625

16. Gras S, Chadderton J, Del Campo CM, Farenc C, Wiede F, Josephs TM, *et al*. Reversed T Cell
Receptor Docking on a Major Histocompatibility Class I Complex Limits Involvement in the
Immune Response. *Immunity* (2016) **45**:749–60. doi: 10.1016/j.immuni.2016.09.007

17. Murray JS. An old Twist in HLA-A: CDR3α Hook up at an R65-joint. *Front Immunol* (2015)
6:268. doi: 10.3389/fimmu.2015.00268

18. Beringer DX, Kleijwegt FS, Wiede F, van der Slik AR, Loh KL, Petersen J, *et al*. T cell receptor
reversed polarity recognition of a self-antigen major histocompatibility complex. *Nat Immunol*
(2015) **16**:1153–61. doi: 10.1038/ni.3271

19. Burrows SR, Chen Z, Archbold JK, Tynan FE, Beddoe T, Kjer-Nielsen L, *et al*. Hard wiring of T
cell receptor specificity for the major histocompatibility complex is underpinned by TCR
adaptability. *Proc Natl Acad Sci USA* (2010) **107**:10608–13. doi: 10.1073/pnas.1004926107

20. Ashwell JD, Chen C, Schwartz RH.   High frequency and nonrandom distribution of
alloreactivity in T cell clones selected for recognition of foreign antigen in association with self
class II molecules. *J Immuno*l (1986) **136**:389–95.

21. Murray JS, Madri J, Tite J, Carding SR, Bottomly K. MHC control of CD4+ T cell subset
activation. *J Exp Med* (1989) **170**:2135–40. doi: 10.1084/jem.170.6.2135

22. Gottschalk RA, Hathorn MM, Beuneu H, Corse E, Dustin ML, Altan-Bonnet G, Allison JP.
Distinct influences of peptide-MHC quality and quantity on in vivo T-cell responses. *Proc Natl*
Acad Sci USA (2012) **109**:881–6. doi: 10.1073/pnas.1119763109

23. Kasahara M, Vazquez M, Sato K, McKinney EC, Flajnik MF.   Evolution of the major
histocompatibility complex: class II A cDNA clones from the cartilaginous fish. *Proc Natl Acad*
Sci USA (1992) **89**:6688–92. doi: 10.1073/pnas.89.15.6688

24. Gunnarsen, KS, Hoydahl, LS, Risnes, LF, Dahal-Koirala, S, Neumann, RS, Bergseng, E, *et al*. A
TCRα frame-work centered codon shapes a biased T cell repertoire through direct MHC and
CDR3β interactions. *JCI Insight* (2017) **2**:95193. doi:10.1172jci.insight.95193

25. Chen J, Thirumalai D. Interface Residues That Drive Allosteric Transitions Also Control the
Assembly of l-Lactate Dehydrogenase. *J Phys Chem B* (2018) **122:**11195–205.   doi:
10.1021/acs.jpcb.8b06430

26. Stewart J. *Calculus*, 6th ed. Belmont, CA: Brooks/Cole Thomson (2007).

27. Das BB, Park SH, Opella SJ. Membrane protein structure from rotational diffusion. *Biochim*
Biophys Acta (2015) **1848**:229–45. doi: 10.1016/j.bbamem.2014.04.002

28. Berg JM, Tymoczko JL, Stryer L. *Biochemistry*, 5th ed. New York: WH Freeman (2002).

29. Brunori M. The Bohr effect before Perutz. *Biochem Mol Biol Educ* (2012) **40**:297–9. doi:
10.1002/bmb.20629

30. Marrack P, Krovi SH, Silberman D, White J, Kushnir E, Nakayama M, *et al*. The somatically
generated portion of T cell receptor CDR3α contributes to the MHC allele specificity of the T cell
receptor. e*life* (2017) **6. pii**:e30918. doi: 10.7554/eLife

31. Carey FA, Sundberg RJ. *Advanced Organic Chemistry, pt. A. Structure and Mechanism*, 2[nd]
ed. New York: Plenum Press (1984).

32. Hernández G, Anderson JS, LeMaster DM. Assessing the native state conformational
distribution of ubiquitin by peptide acidity. *Biophys Chem* (2010) **153**:70–82. doi:
10.1016/j.bpc.2010.10.006

33. Koch M, Camp S, Collen T, Avila D, Salomonsen J, Wallny HJ, *et al*. Structures of an MHC
class I molecule from B21 chickens illustrate promiscuous peptide binding. *Immunity* (2007)
27:885-99. doi: 10.1016/j.immuni.2007.11.007

34. Yin Y, Li Y, Kerzic MC, Martin R, Mariuzza RA. Structure of a TCR with high affinity for self-
antigen reveals basis for escape from negative selection. *EMBO J* (2011) **30**:1137–48. doi:
10.1038/emboj.2011.21

35. Petersen J, Montserrat V, Mujico JR, Loh KL, Beringer DX, van Lummel M, *et al*. T-cell
receptor recognition of HLA-DQ2-gliadin complexes associated with celiac disease. *Nat Struct*
Mol Biol (2014) **21**:480–8. doi: 10.1038/nsmb.2817

36. Zerrahn J, Held W, Raulet DH. The MHC reactivity of the T cell repertoire prior to positive
and negative selection. *Cell* (1997) **88**:627–36. doi: 10.1016/s0092-8674(00)81905-4

37. Pusuluk O, Farrow T, Cemsinan D, Burnett K, Vedral V. Proton tunneling in hydrogen bonds
and its implications in an induced-fit model of enzyme catalysis. *Proc R Soc A* (2018)
474:20180037. dx.doi.org/10.1098/rspa.2018.0037

38. Clement CC, Becerra A, Yin L, Zolla V, Huang L, Merlin S, *et al*. The dendritic cell major
histocompatibility complex II (MHC II) peptidome derives from a variety of processing pathways
and includes peptides with a broad spectrum of HLA-DM sensitivity. *J Biol Chem* (2016)
291:5576–95. doi: 10.1074/jbc.M115.655738

39. Arrieta-Bolaños E, Crivello P, Metzing M, Meurer T, Ahci M, Rytlewski J, *et al*. Alloreactive T
cell receptor diversity against structurally similar or dissimilar HLA-DP antigens assessed by
deep sequencing. *Front Immunol* (2018) **9**:280. doi: 10.3389/fimmu.2018.00280

40. Murray JS. How the MHC selects Th1/Th2 immunity. *Immunol Today* (1998) **19**:157–63.
doi: 10.1016/s0167-5699(97)01237-1

41. Faro J, Castro M, Molina-París C. A unifying mathematical framework for experimental
TCR-pMHC kinetic constants. *Sci Rep* (2017) **7**:46741. doi: 10.1038/srep46741

42. Wang JH. T cell receptors, mechanosensors, catch bonds and immunotherapy. *Prog*
Biophys Mol Biol (2020) **pii**: S0079–6107(20)30008-0. doi: 10.1016/j.pbiomolbio.2020.01.001

43. Limozin L, Bridge M, Bongrand P, Dushek O, van der Merwe PA, Robert P. TCR-pMHC
kinetics under force in a cell-free system show no intrinsic catch bond, but a minimal encounter
duration before binding. *Proc Natl Acad Sci U S A* (2019) **116**:16943–48. doi:
10.1073/pnas.1902141116

44. Murray JS, Brown JC. Measurement of association constants in ELISA. Reactions between
solid-phase antibody and fluid-phase biotinylated antigen. *J Immunol Methods* (1990) **127**:25–8.
doi: 10.1016/0022-1759(90)90336-t

45. Dunbar J, Knapp B, Fuchs A, Shi J, Deane CM. Examining variable domain orientations in
antigen receptors gives insight into TCR-Like antibody design. *PLOS Comput Biol* (2014)
10:e1003852. doi: 10.1371/journal.pcbi.1003852

46. Hülsmeyer M, Chames P, Hillig RC, Stanfield RL, Held G, Coulie PG, *et al*. A major
histocompatibility complex-peptide-restricted antibody and T cell receptor molecules recognize
their target by distinct binding modes. *J Biol Chem* (2005) **280**:2972–80. doi:
10.1074/jbc.M411323200

47. Lodola A, Mor M, Zurek J, Tarzia G, Piomelli D, Harvey JN, Mulholland AJ. Conformational
effects in enzyme catalysis: reaction via high energy conformation in fatty acid amide hydrolase.
Biophys J (2007) **92**:L20–2. doi: 10.1529/biophyj.106.098434

48. Pielak RM, O'Donoghue GP, Lin JJ, Alfieri KN, Fay NC, Low-Nam ST, Groves JT. Early T cell
receptor signals globally modulate ligand:receptor affinities during antigen discrimination. *Proc*
Natl Acad Sci U S A (2017) **114**:12190–5. doi: 10.1073/pnas.1613140114

49. Deng L, Langley RJ, Wang Q, Topalian SL, Mariuzza RA. Structural insights into the editing
of germ-line-encoded interactions between T-cell receptor and MHC class II by Vα CDR3. *Proc*
Natl Acad Sci USA (2012) **109**:14960–5. doi: 10.1073/pnas.1207186109

50. Natarajan K, Jiang J, May NA, Mage MG, Boyd LF, McShan AC, *et al*. The Role of Molecular
Flexibility in Antigen Presentation and T Cell Receptor-Mediated Signaling. *Front Immunol*
(2018) **9**:1657. doi: 10.3389/fimmu.2018.01657

51. Buckle AM,   Borg   NA.   Integrating Experiment and Theory to Understand TCR-pMHC
Dynamics. *Front Immunol* (2018) **9**:2898. doi: 10.3389/fimmu.2018.02898

52. Pfleger C, Minges A, Boehm M, McClendon CL, Torella R, Gohlke H. Ensemble- and Rigidity
Theory-Based Perturbation Approach To Analyze Dynamic Allostery. *J Chem Theory Comput*
(2017) **13**:6343–57. doi: 10.1021/acs.jctc.7b00529

53. Carlson GM, Fenton AW.   What Mutagenesis Can and Cannot Reveal About Allostery.
Biophys J (2016) **110**:1912–23. doi: 10.1016/j.bpj.2016.03.021

54. Sibener LV, Fernandes RA, Kolawole EM, Carbone CB, Liu F, McAffee D, *et al*. Isolation and
visualization of a structural trigger that uncouples TCR signaling from pMHC binding. *Cell*
(2018) **174**:672–87.e27. doi: 10.1016/j.cell.2018.06.017

55. De Oliveira DB, Harfouch-Hammoud E, Otto H, Papandreou NA, Stern LJ, Cohen H, *et al*.
Structural analysis of two HLA-DR-presented autoantigenic epitopes: crucial role of peripheral
but not central peptide residues for T-cell receptor recognition. *Mol Immunol* (2000) **37**:813–
25. doi: 10.1016/s0161-5890(00)00109-7

56. Holzer U, Kwok WW, Nepom GT, Buckner JH.   Differential antigen sensitivity and
costimulatory requirements in human Th1 and Th2 antigen-specific CD4+ cells with similar TCR
avidity. *J Immunol* (2003) **170**:1218–23. doi: 10.4049/jimmunol.170.3.1218

57. Lin JJY, Low-Nam ST, Alfieri KN, McAffee DB, Fay NC, Groves JT. Mapping the stochastic
sequence of individual ligand-receptor binding events to cellular activation: T cells act on the
rare events. *Sci Signal* (2019) **12**:pii: eaat8715. doi: 10.1126/scisignal.aat8715

58. Galvez J, Galvez JJ, Garcia-Penarrubia P. Is TCR/pMHC Affinity a Good Estimate of the T-cell
Response? An Answer Based on Predictions From 12 Phenotypic Models. *Front Immunol* (2019)
10:349. doi: 10.3389/fimmu.2019.00349

59. Scharf L, West AP Jr, Sievers SA, Chen C, Jiang S, Han G, *et al*.   Structural basis
for germline antibody recognition of HIV-1 immunogens. *eLife* (2016) **5**:e13783.   doi:
10.7554/eLife.13783

60. Lu J, Van Laethem F, Bhattacharya A, Craveiro M, Saba I, Chu J, *et al*. Molecular constraints
on CDR3 for thymic selection of MHC-restricted TCRs from a random pre-selection repertoire.
Nat Commun (2019) **10**:1019. doi: 10.1038/s41467-019-08906-7

61. Khosravi-Maharlooei M, Obradovic A, Misra A, Motwani K, Holzl M, Seay HR, *et al*.
Crossreactive public TCR sequences undergo positive selection in the human thymic repertoire.
J Clin Invest (2019) **129**:2446–62. doi: 10.1172/JCI124358

62. Frick R, Gunnarsen KS, Dahal-Koirala S, Risnes LF, Sollid LM, Sandlie I, *et al*. A TRAV26-1-
encoded recognition motif focuses the biased T cell response in celiac disease. *Eur J Immunol*
(2020) **50**:142–5. doi: 10.1002/eji.201948235

63. Murray JS. Transposon-mediated death of an ancestral A-23-like allele: evolution of TCR-
positioning   motifs   in   the   HLA-A   lineage.   *Immunogenetics*   (2015)   **67**:473–6.   doi:
10.1007/s00251-015-0852-3

64. Steiper ME, Seiffert ER. Evidence for a convergent slowdown in primate molecular rates
and its implications for the timing of early primate evolution. *Proc Natl Acad Sci U S A* (2012)
109:6006–11. doi: 10.1073/pnas.1119506109

65. Singh NK, Abualrous ET, Ayres CM, Noé F, Gowthaman R, Pierce, BG, *et al*. Geometrical
characterization of T cell receptor binding modes reveals class-specific binding to maximize
access to antigen. *Proteins (*2020) **88**:503-513. doi: 10.1002/prot.25829

66. Hoffmann T, Krackhardt AM, Antes I.   Quantitative Analysis of the Association Angle
between T-cell Receptor Vα/Vβ Domains Reveals Important Features for Epitope Recognition.
PLoS Comput Biol (2015) e1004244. doi: 10.1371/journal.pcbi.1004244.

67. Pauling L. *General Chemistry*, 3rd ed.   San Francisco: WH Freeman (1970).