Extended-spectrum beta-lactamase antibiotic resistance plasmids have diverse transfer rates

and can be spread in the absence of selection

Fabienne Benz[1][*][¶], Jana S. Huisman[1,2][¶], Erik Bakkeren[3][¶], Joana A. Herter[3], Tanja Stadler[2,4],

Martin Ackermann[5,6], Médéric Diard[3,7], Adrian Egli[8,9][&], Alex R. Hall[1][&], Wolf-Dietrich Hardt[3][&],

Sebastian Bonhoeffer[1][&]

[1]Institute of Integrative Biology, Department of Environmental Systems Science, ETH Zurich,
Zurich, Switzerland
11
12 [2]Swiss Institute of Bioinformatics, Basel, Switzerland
13
14 [3]Institute of Microbiology, Department of Biology, ETH Zurich, Zurich, Switzerland
15
16 [4]Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland
17
18 [5]Institute of Biogeochemistry and Pollutant Dynamics, Department of Environmental Systems
19 Science, ETH Zurich, Zurich, Switzerland
20
21 [6]Department of Environmental Microbiology, Eawag, Duebendorf, Switzerland
22
23 [7]Biozentrum, University of Basel, Basel, Switzerland
24
25 [8]Division of Clinical Bacteriology and Mycology, University Hospital Basel, Basel, Switzerland
26
27 [9]Applied Microbiology Research, Department of Biomedicine, University of Basel, Basel,
28 Switzerland
29
30 [*] Corresponding author

31 E-mail: fabienne.benz@env.ethz.ch (FB)

33 [¶] These authors contributed equally to this work

34 [&] These authors also contributed equally to this work

Abstract

Horizontal gene transfer, mediated by conjugative plasmids, is one of the main drivers of the global spread of antibiotic resistance. However, the relative contributions of different factors that underlie this plasmid spread are unclear, particularly for clinically relevant plasmids harboring antibiotic resistance genes. Here, we analyze nosocomial outbreak-associated plasmids that reflect the most relevant Extended Spectrum Beta-Lactamase (ESBL) mediated drug resistance plasmids to i) quantify conjugative transfer dynamics, and ii) investigate why some plasmid-strain associations are more successful than others, in terms of bacterial fitness and plasmid spread. We show that, in the absence of antibiotics, clinical *Escherichia coli* strains natively associated with ESBL-plasmids conjugate efficiently with three distinct *E. coli* strains and one *Salmonella* enterica Serovar Typhimurium strain. In more than 40% of the *in vitro* mating populations, ESBL-plasmids were transferred to recipients, reaching final transconjugant frequencies of up to 1% within 23 hours. Variation of final transconjugant frequencies was better explained by variation in conjugative transfer efficiency than by variable clonal expansion of transconjugants. We also identified plasmid-specific genetic factors, such as the presence/absence of particular transfer genes, that influenced final transconjugant frequencies. Finally, we validated the plasmid spread in a mouse model for gut colonization, demonstrating qualitative correlation between plasmid spread *in vitro* and *in vivo.* This suggests a potential for predictive modelling of plasmid spread in the gut of animals and humans, based on *in vitro* testing. Altogether, this may allow straightforward identification of resistance plasmids with high spreading potential and to implement quarantine or decolonization procedures to restrict their spread.

Author summary

Antibiotic resistance is a major obstacle to the treatment of bacterial infections in clinics. Plasmids encoding antibiotic resistance genes can spread between bacteria in a density-dependent manner and accelerate the rise of resistant bacterial strains. This is particularly important for densely inhabited ecological niches such as the guts of humans and animals, where many bacteria interact. Understanding the exact contribution plasmids make to the global spread of antibiotic resistance remains an obstacle, because we lack quantitative studies implementing large-scale experimental testing of conjugation rates between clinically relevant bacterial strains. To counteract this knowledge gap, we studied clinical *Escherichia coli* isolates from human patients that carry extended-spectrum beta-lactamase producing plasmids. We found that these plasmids spread extensively through different bacterial populations and that both bacterial- and plasmid-specific factors determined the extent of plasmid spread. Our study combines detailed bioinformatic analyses, high-throughput *in vitro* testing and validation in an animal model. It suggests a potential for laboratory testing to understand and predict the spread of clinically relevant plasmids, including in the human gut microbiota, and thereby generates insights into novel treatment strategies to manage antibiotic resistance spread mediated by plasmids.

Introduction

The prevalence of antibiotic resistant bacterial pathogens is rising and continues to challenge the clinical use of antibiotics [1]. Infections with bacteria resistant to antibiotics are increasingly common and can result in death [2,3]. Importantly, resistance determinants are often plasmid-encoded. Plasmids can be transferred horizontally between different bacterial cells by conjugation, allowing rapid spread through diverse bacterial communities. This includes transfer among and between clinically relevant bacterial pathogens and commensals found in the gut microbiota of humans, farm animals, or in the environment [4–6]. In some cases, the role of selection imposed by antibiotics may be overestimated, because certain resistance plasmids can increase in frequency even in the absence of antibiotics [7–9] .

Whether a plasmid persists within a population depends on the plasmid as well as its association with a bacterial host. Some of these associations, such as *Escherichia coli* sequence type ST131 and plasmids of the IncFII family, carrying the *bla CTX-M* gene (which encodes a cefotaximase, conferring resistance to extended-spectrum beta lactams; ESBLs), are very successful and have become global threats [10–12]. ESBLs provide their bacterial host with resistance to beta-lactam antibiotics, such as the widely used penicillins and cephalosporins, and are generally encoded on plasmids of the incompatibility groups IncF and IncI [13–15]. Other plasmids, such as those carrying the carbapenemase-producing gene blaOXA-48, remain a clinical problem despite no association with a specific strain. For instance, blaOXA-48 plasmids have been reported in *E. coli, Klebsiella spp., Citrobacter spp.,* and *Enterobacter spp*. [16]. The family of Enterobacteriaceae is a key public health concern, as it contains some of the most important nosocomial pathogens and is responsible for the pandemic spread of ESBLs [10,15,17,18].

To counteract resistant pathogens and their successful transmission in hospitals, communities, and the environment, we need to improve our understanding of the factors that determine successful plasmid-strain associations and to elucidate the drivers of resistance plasmid spread in the presence or absence of antibiotics. Recent *in vitro* studies revealed several key factors driving changes in resistance plasmid frequency over time. If a plasmid of the same incompatibility group is already established in a potential recipient, surface- or entry exclusion may prevent further plasmid acquisition [19]. Plasmids can also increase transfer rate of co-residing plasmids, or allow other (mobilizable) plasmids to co-transfer [20,21]. However, once a plasmid is taken up by a bacterium, the plasmid replication system and its interaction with host factors define whether it can be replicated and stably maintained in the lineage. For example, the efficiency of plasmid transfer decreases if a recipient strain encodes a restriction system, to which the incoming plasmid is susceptible [22–24]. In some cases, compensatory mutations have been shown to reduce plasmid cost in the presence and absence of selective pressure [25]. In others, it is the high rates of conjugation that allow a plasmid to persist despite a growth disadvantage [26]. With the recent discovery of several novel defense systems against foreign DNA in bacteria [24], interactions between bacteria, particularly during horizontal gene transfer, become increasingly interesting to study. This may allow us to find tools to prevent spread of plasmids carrying antibiotic resistance genes.

The relative contributions of the various factors that drive plasmid spread remain poorly understood, particularly for clinical plasmids [11]. Past studies have largely focused on types of plasmids not directly relevant for resistance in clinical settings, or on individual plasmids

after moving them into model strains [22,25–27] (notable exception [28]). Studies using

mouse models have contributed to our understanding of the processes that drive conjugation

within the gut, but these have mostly been limited to laboratory strains with conjugative

plasmids that are not clinically relevant [8,9,29,30]. Human-based studies that demonstrate

horizontal transfer of resistance plasmids in the human gut [5,31] are of strong clinical

relevance, but often rely only on genomic evidence and thus do not allow to observe the

factors driving plasmid spread. There is therefore a shortage of studies connecting

mechanisms of plasmid spread found *in vitro* with corresponding patterns *in vivo* and with

the effective spread of plasmids in the clinics [11,32,33]. To improve the predictability of

clinically relevant plasmid spread, we need a quantitative understanding of resistance plasmid

transfer from their clinical host strains to diverse recipient strains, the effect of plasmids on

the growth phenotypes of these recipient strains, subsequent transfer of newly acquired

plasmids to other recipients, and the impact of bacterial host factors on these properties.

Here, we use clinical *E. coli* strains with their natively associated plasmids carrying ESBL genes

(ESBL-plasmids) to investigate the drivers of plasmid spread, and to study the correspondence

between *in vitro* and *in vivo* spread. Based on *in vitro* liquid culture mating assays in the

absence of antibiotic selection, we found that the final frequency of ESBL-plasmid carrying

recipient strains (transconjugants) was determined by donor-, recipient-, and plasmid-specific

factors. The plasmid itself had the largest contribution to final transconjugant frequency, as

all plasmids that carry functional transfer (*tra*) genes were found to spread. Additionally, none

of the tested plasmids show evidence of a growth cost in any recipient strain. Using a murine

model for gut colonization, we validated that these plasmids can spread efficiently *in vivo* in

the absence of antibiotic selection. As *in vitro*, plasmid carriage of the tested plasmids was

not associated with a major cost or benefit in the gut, confirming the critical role of

conjugation in driving the spread of antibiotic resistance plasmids. We showed that the range

of transfer efficiencies *in vivo* are qualitatively predictable based on our *in vitro* testing.

Results

Diverse clinical ESBL-plasmids spread extensively through recipient populations in the absence of antibiotics. We investigated the rate at which ESBL-plasmids from eight clinical donor *E. coli* strains spread through four non-resistant recipient populations. The donor strains stem from hospitalized patients and are representative of clinically relevant ESBL-producing *E. coli* [34]. Of the four recipients, two *E. coli* strains were isolated from healthy human subjects [35], one *E. coli* strain was previously isolated from mice housed in our facility [8], and one is a pathogenic wild type strain of *Salmonella enterica* Serovar Typhimurium. The chosen donor and recipient strains spanned the phylogenetic diversity of *E. coli* [36,37] (Fig 1) and contained many plasmids of various incompatibility groups (S1 Fig, S1 Table). Each donor carried one plasmid containing resistance genes, either of the plasmid family IncI1 or IncF (Table 1, S2 Fig). We found various virulence genes, including type 6 secretion systems (T6SSs, involved in bacterial killing), in donors and recipients (S3 Fig). The only strain with a pathogenic virulence profile is D4 (S3 Fig). All strains carry numerous intact prophage sequences (S1 File).

Table 1: Clinical ESBL-plasmids in their native host strains (donors).

Strain (*E. coli* sequence type)	ESBL-plasmid	Plasmid size	Replicon (incompatibility group)	Resistance genes on the ESBL-plasmid†	Antibiotic resistance of strain*
D1 (ST 117)	p1B_IncI	111 kB	IncI1	aadA5; **blaCTX-M-1**; dfrA17; sul2	Amp, Ceftri, Cotri
D2 (ST 648)	p2A_IncF	165.7 kB	IncFIA; IncFIB; IncFII	mph(A); catB3; aadA5; aac(6')-Ib-cr; dfrA17; sul1; sul2; **blaOXA-1**; tet(A); **blaCTX-M-27**; aph(3'')-Ib; erm(B)	Amp, Amo/C; Ceftri; Tobra; Cotri; Cipro
D3 (ST 648)	p3B_IncI	59.8 kB	IncI1	**blaCMY-42**	Amp; Amo/C; Pip-T; Ceftaz; Ceftri;

					Tobra; Amika; Cotri; Cipro
D4 (ST 40)	p4A_IncI	88.9 kB	IncI1	**blaCTX-M-1**	Amp; Ceftri
D5 (ST 131)	p5A_IncF	160 kB	IncFIB; IncFIC(FII); IncFIA	mph(A); aadA5; aac(6')-Ib-cr; dfrA17; sul1; **blaCTX-M-15**; aac(3)-IIa; tet(A); catB3; **blaOXA-1**	Amp; Amo/C; Ceftaz; Ceftri; Cefe; Tobra; Cotri; Cipro
D6 (ST 131)	p6A_IncI/F	157 kB	IncI1; IncFIA; IncFIB; IncFII	**blaTEM-1B**, aac(3)-IId, **blaCTX-M-8**	Amp; Ceftri; Cefe; Tobra; Cipro
D7 (ST 131)	p7A_IncF/Col156	134.9 kB	IncFIA; IncFIB; IncFII; Col156	mph(A); aph(3'')-Ib; aadA5; dfrA17; sul1; sul2; tet(A); **blaCTX-M-27**	Amp; Ceftri; Cotri; Cipro
D8 (ST 69)	p8A_IncF	131 kB	IncFIA; IncFIB	dfrA14, mph(A), **blaCTX-M-14**, tet(B)	Amp; Ceftri; Cotri

†Genes encoding the ESBL phenotype are highlighted in bold.

*Antibiotic resistance is defined as being above the EUCAST defined minimum inhibitory concentration breakpoint (MIC; see S2 Table for all antibiotics tested and MIC information). Donor strains fulfilled criteria for ESBL production based on EUCAST recommendations. ESBL mechanism was phenotypically and genotypically confirmed. Abbreviations: Amp = Ampicillin; Amo/C = Amoxicillin/Clavulanic acid; Pip-T = Piperacillin-Tazobactam; Ceftaz = Ceftazidim; Ceftri = Ceftriaxone; Cefe = Cefepim; Tobra = Tobramycin; Amika = Amikacin; Cotri = Cotrimoxazol; Cipro = Ciprofloxacin.

First, we performed conjugation assays between all clinical donors and non-resistant

recipients (approximately 1:1 ratio, referred to below as the "1st gen *in vitro* experiment") and

measured conjugation as the final fraction of the recipient population that carried the plasmid

after 23 hours of growth (hereafter termed "final transconjugant frequency"). Five of the

eight ESBL-plasmids were transferred to more than one *E. coli* recipient (Fig 2) and transfer

stemming from the donor-plasmid pair D4-p4A_IncI led to the highest final transconjugant

frequencies of ~0.1%. Final transconjugant frequencies spanned 5 orders of magnitude (Fig

2A). For the donor-plasmid pairs where we observed plasmid spread to multiple recipients,

average final transconjugant frequency varied depending on the donor-plasmid pair (effect

of donor-plasmid pair in a two-way ANOVA, excluding D2, D3, D7: $F_{4,66}$ = 87.665, P < 0.01) and

among recipient strains (effect of recipient in the same two-way ANOVA: $F_{2,66}$ = 5.439, P <

0.01). Furthermore, variation among donor-plasmid pairs depended on the recipient (donor-

plasmid pair×recipient interaction: $F_{8,66}$ = 3.164, P < 0.01). These plasmids, natively associated

with *E. coli*, reached comparable maximal transconjugant frequencies in *S.* Typhimurium

recipient populations (RS, Fig 2B). In *S*. Typhimurium, variation across donor-plasmid pairs

was similar to *E. coli* recipients, with the exception of p6A_IncI, which did not spread to

recipient RS.

Qualitatively, phylogenetic relatedness (Fig 1) between donor strains did not explain

similarities in their plasmid transfer dynamics in the 1st gen *in vitro* experiment. For example,

D5, D6 and D7 have effectively the same host background (ST 131), yet transferred their

plasmids with very different efficiencies (Fig 2). By contrast, D1 and D8 are relatively

phylogenetically distant, yet showed very similar transfer dynamics across recipients (Fig 2).

Phylogenetic relatedness between donor strain and recipient was not strongly associated

with transfer efficiency either: D5, D6 and D7 are all equally closely related to recipients RE1

and RE2, yet we observed very different final transconjugants frequencies, spanning four

orders of magnitude, among these 6 donor-recipient pairs (Fig 2). Finally, the *S.* Typhimurium

(RS) recipient is roughly equally phylogenetically distant to all *E. coli* donors used, yet we

observed widely varying transfer efficiencies, similar to those observed across *E. coli*

recipients.

To investigate the effect of genomic factors on the observed plasmid spread, we related final

transconjugant frequencies to the presence or absence of *tra* (transfer) genes and toxin-

antitoxin systems (TA-systems) on plasmids and of restriction-modification systems and

CRISPR-Cas in donor and recipient strains (S2 File). We found that the presence of relevant

tra genes on a plasmid was the main genomic factor that affected whether plasmids were

able to spread. *Tra* genes encode the sex pilus and the proteins required for the conjugative

transfer, and thus, when missing, plasmids cannot conduct their own transfer [38–40]. All

three cases in which the plasmid showed no or almost no spread to any recipient (i.e.

p2A_IncF, p7A_IncF/Col156, p3B_IncI) could be explained by their lack of functional *tra* genes

(S2 File). All plasmids encode for numerous TA-systems, likely leading to plasmid addiction,

and thus we could not find a relation between these and final transconjugant frequencies.

Despite the large number of unique donor-recipient combinations, the analysis of CRISPR-Cas

and restriction-modification systems did not allow a clear association between their presence

and the observed plasmid spread (S2 File). *Tra* genes, were the only genomic factor for which

presence or absence could be consistently associated with the final transconjugant frequency.

ESBL-plasmids from clinical donors carry no significant cost after transfer to new hosts. The

final frequency of the ESBL-plasmids in different recipient populations spanned five orders of

magnitude after 23 hours. This variation in plasmid spread could be driven by differences in

horizontal transmission, or in clonal expansion of transconjugants. Thus, we investigated the

effects of plasmids on bacterial growth in the absence of antibiotics for ten strain-plasmid

combinations. We estimated growth rates of recipients and transconjugants over 24 hours

and expressed plasmid cost as the growth rate of transconjugants relative to their respective

plasmid-free strains (Fig 3A for *E. coli* recipients and 3B for *S.* Typhimurium). None of the novel

host-plasmid combinations had a significantly different growth rate compared to the original

host strain (Student's t-Test for *E. coli* hosts and Wilcoxon Rank Sum Test for *S.* Typhimurium,

$P > 0.05$ in all cases, before and after Holm's correction for multiple testing). Although non-

significant, the *S*. Typhimurium transconjugants showed a consistent trend towards higher

relative growth rates when carrying an ESBL-plasmid, and were not normally distributed.

Whether this trend can explain the final frequency of *S*. Typhimurium transconjugants (Fig

2B) depends on the magnitude of the growth advantage, and the duration of the conjugation

assay. We calculated whether, under the assumption of purely exponential growth, the mean

difference in growth rate between recipients and transconjugants would be large enough to

explain the observed final transconjugant frequencies (S3 File). This is the case for the

combination of donors D1 and D8 with *S.* Typhimurium recipients, but for *E. coli* recipients,

clonal expansion is unlikely to explain the observed variation in final transconjugant

frequencies.

Plasmid transfer rate drives the final transconjugant frequency. Because for most cases

clonal expansion alone did not explain the observed final transconjugant frequencies, we

investigated whether variable rates of horizontal transfer (conjugation) could explain this. A

widely used method to estimate plasmid transfer rates is the mass action model from

Simonsen *et al.* [41]. When we applied it to the data obtained in the 1st gen *in vitro* experiment

(S3 Table), we found that across all donor-recipient combinations, the estimated conjugative

transfer rates correlate strongly with the observed final transconjugant frequencies (Fig 4,

Pearson's test, r = 0.99, P < 0.001). This suggests that plasmid transfer rates dictate final

transconjugant frequencies.

A drawback of the Simonsen method is that it assumes that (i) the growth rates for donor,

recipient and transconjugants are all equal, (ii) the conjugation rate does not change over

time (i.e. that the rate of conjugation from the donors and newly formed transconjugants to

the recipients is the same), and (iii) mating takes place in well-mixed liquid cultures. However,

the transfer rate estimates of the experiments described here are not likely to be affected

significantly by violation of these assumptions. Firstly, the observed differences in growth rate

for donors, recipients, and transconjugants (Fig 3 and S3 File), are too small to drive shifts in

relative frequencies over the course of our conjugation assays. Secondly, transconjugants

reached maximally 0.1% of the donor population size, which is comparatively small (S4 Fig).

Therefore, the vast majority of transfer events results from donor-to-recipient conjugation

rather than transconjugant-to-recipient transfer. Thirdly, regarding assumption (iii), we

conducted a control experiment to estimate the number of transconjugants observed solely

due to surface mating during the plating step of our conjugation assay (S4 Table). For most

donor-recipient pairs we found that the number of transconjugants due to on-plate mating

was negligible. The pairs D1-RE1 and D1-RE3, however, reached high transconjugant CFU

counts due to surface-mating only (35% and 85% of the numbers observed in the 1^{st} gen *in*

vitro experiment). Overall, we interpret estimates from the Simonsen method as an accurate

reflection of the conjugation rate in our experiments, but for D1-RE1 and D1-RE3, it may

reflect the combined effect of conjugation in liquid and on the agar surface after plating.

Variation in plasmid spread stems from plasmid, donor and recipient factors. In the 1^{st} gen

in vitro experiment each plasmid was present in a single donor strain (Fig 2), preventing us

from disentangling the effects of donor strain and plasmid on plasmid spread. Thus, we next

performed an experiment under conditions identical to above, using 8 different plasmid

donors (2^{nd} gen *in vitro,* Fig 5). Each donor is a unique combination of strain (one of three

different recipient strains from above) and plasmid (one of three different plasmids from

above), isolated from among the transconjugants of the 1^{st} gen *in vitro* experiment. The final

transconjugant frequency varied among donor strains (effect of donor strain in a three-way

ANOVA with plasmid, donor strain and recipient as factors, excluding plasmid p1B_IncI, see

Materials and methods: $F_{2,90} = 150.133$, $P < 0.001$) and among plasmids ($F_{1,90} = 49.717$, $P <$

0.001). This observed variation among plasmids depended on both the recipient and the

donor strain. For instance, in cases where *S.* Typhimurium RS acted as the donor and recipient

simultaneously, both IncI1 ESBL-plasmids yielded remarkably high final transconjugant

frequencies of 40 % for p1B_IncI and 39 % for p4A_IncI (donor strain×plasmid interaction:

$F_{2,90}$ = 96.352, P < 0.001; recipient×plasmid interaction: $F_{2,90}$ = 29.610, P < 0.001). A second

analysis including plasmid p1B_IncI, but excluding donor RE2 (see Materials and methods)

supported variation among donor strains ($F_{1,93}$ = 560.269, P < 0.001) and plasmids ($F_{2,93}$ =

156.075, P < 0.001), and that variation among plasmids depended on recipient and donor

strain (recipient×plasmid interaction ($F_{4,93}$ = 26.104, P < 0.001; donor strain×plasmid

interaction: $F_{2,93}$ = 3.999, P = 0.022). As in our 1st gen *in vitro* experiment, average final

transconjugant frequencies also varied among recipients (P < 0.001 for effect of recipient in

both three-way ANOVAs). Thus, the final frequency of transconjugants depended on all three

factors of donor strain, plasmid, and recipient.

The final transconjugant frequencies in this 2nd gen *in vitro* experiment, where we replaced

the native donor strains with transconjugants stemming from the 1st gen *in vitro* assay,

differed strongly from those of the 1st gen *in vitro* experiment. They increased >10-fold in 11

and decreased >10-fold in 4 out of the 24 mating populations (Figs 2 and 5). Because the

transferred plasmids and recipient strains were equal to, and the absolute frequencies of

donor and recipient comparable to, those of the 1st gen in *vitro* experiment (S4-5 Figs), we

tested whether these differing transconjugant frequencies stem from a difference in transfer

from native host versus transfer from a secondary host (S7 Fig). As donors of plasmid

p1B_IncI, we used its native host D1 and its transconjugants stemming from conjugation with

recipients RE1 and RS, both carrying the marker plasmid pACYC184, and an additional D1

strain carrying pACYC184. First, we tested whether the presence of pACYC184 in donors could

affect the transfer of ESBL-plasmids. Indeed, pACYC184 in D1 had a significantly negative

effect on transfer of p1B_IncI to RE1 (Wilcoxon Rank Sum Test, P = 0.035 after Holm's

correction for multiple testing) and to RS (Student's t-Test, P = 0.042 after Holm's correction

for multiple testing). Although significant, the effect of pACYC184 was small compared to the

difference in transconjugant frequency that resulted from transfer from native versus 2nd gen

donor strain. For instance, conjugation of the donor strains RE1 and RS with recipient RE1

resulted in a 45-fold and a 112-fold increase respectively, compared to conjugation of the

native donor strain D1 with recipient RE1. When donor and recipient were RS, final

transconjugant frequency even increased 2800-fold (S7 Fig). Overall, this experiment

confirmed the significant difference in final transconjugant frequency resulting from transfer

by native versus secondary donor strains and highlighted the great increase in final

transconjugant frequency when donor and recipient are the same strain (self-self transfer).

A possible explanation for the altered transfer efficiency when transconjugants acted as

plasmid donors, are mutational changes, in either the plasmid or the recipients, that might

have accumulated during the 1st gen or the 2nd gen *in vitro* experiment. Thus, we analyzed

transconjugants from the 1st and 2nd gen *in vitro* experiments by whole-genome sequencing

(one clone for each transconjugant, except for RE3 carrying plasmid p4A_IncI and p8A_IncF,

for which we sequenced three clones). We did not find any mutational changes in ESBL-

plasmids but a range of chromosomal mutations in transconjugants (S5 Table). However,

these mutations were not consistently present across transconjugants and could not be

related to adaptation to plasmid carriage. Nevertheless, we expect that host factors play a

major role in plasmid spread, and it is simply their interplay and our small sample size that

prevent us from detecting one single crucial conjugation rate determinant.

ESBL-plasmids can spread rapidly *in vivo*, with efficiencies correlating with the *in vitro* trends. Sequence-based studies have shown that ESBL-plasmids can spread between bacteria in the human gut [5,31], but their actual conjugation dynamics have not yet been studied *in vivo* (exception using laboratory *S.* Typhimurium strains [9]). Therefore, we extended our study to a more realistic murine model. We performed conjugation assays in mice with a limited microbiota [42] with three of the clinical donor strains carrying ESBL-plasmids (D4, D8, and D7) and RE3 as a recipient (Fig 6). These mice allow colonization of approximately 10^8 *E. coli* per gram feces, densities of *E. coli* that can be found in the guts of humans and animals [37]. Donors and recipients were introduced at equal densities but at different timepoints, the recipient one day prior to the donor, and were allowed to grow and conjugate over 7 days (Fig 6A). In line with our *in vitro* conjugation assay, we observed transconjugants for p4A_IncI and p8A_IncF, but not for p7A_IncF/Col156 (Fig 6B). Similar to the *in vitro* results in Fig 2, transfer of p4A_IncI resulted in higher transconjugants frequencies than transfer of p8A_IncF (day 1-6, Fig 6B). Plasmid p4A_IncI reached maximal proportions of plasmid carrying recipients of ~1% within only 4-8 hours of colonization (Fig 6C). Overall, we found that the rank order of final transconjugant frequencies is conserved between *in vitro* and *in vivo* conjugation. Moreover, the speed at which these clinical plasmids can establish themselves in the gut in the absence of antibiotic selection is striking.

As was the case *in vitro*, the transconjugant population *in vivo* was relatively minor compared to the size of the donor population (S4-6 Figs). This suggests that most transfer events derived from donor-to-recipient transfer, rather than transconjugant-to-recipient transfer. Given this,

it is likely that the conjugation dynamics *in vivo* can be predicted by the 1st gen *in vitro*

experiments (Fig 2).

To determine whether the increase in transconjugant population size in the gut over the

seven-day experiment was driven by clonal expansion or by conjugation, we investigated the

competitive advantage of transconjugants versus recipients *in vivo*. For two transconjugants

from the 1st gen *in vitro* experiment, we deleted the ESBL-plasmid origin of transfer (*oriT*) and

created "locked", non-conjugative plasmids. These were competed 1:1 with RE3, the recipient

used in the *in vivo* conjugation assay. A change in fitness conferred by plasmid carriage would

be reflected in the competitive index, which is calculated as the ratio between the population

sizes of recipient and "locked" transconjugant. After seven days of competition, only

transconjugants carrying p8A_IncF outcompeted RE3, with a competitive index of ~0.1 (Fig

6D). Over the course of the experiment, this plasmid-borne fitness advantage led the initial

transconjugant frequency of 0.5 to increase to a final transconjugant frequency of 0.9.

Because this two-fold increase in transconjugant frequency is small compared to the

transconjugant frequencies observed in the conjugation experiment (increasing from the

detection limit of 10^{-6} to a frequency of 0.01; Fig 6B), we conclude that final transconjugant

frequencies in the conjugation assay were mainly driven by conjugative transfer, rather than

by clonal expansion of transconjugants. This is seen to a larger extent for p4A_IncI, where the

transconjugant frequency increased substantially faster than with p8A_IncF (Fig 6B), despite

a total lack of a growth advantage over recipient RE3 (Fig 6D). Altogether, we show that in

the murine gut, ESBL-plasmids can be transferred rapidly between *E. coli* in the absence of

antibiotic selection.

Plasmid interactions: the effect of co-residence, co-transfer and plasmid exclusion on plasmid transfer. It has been proposed that the transfer rate of plasmids can be modified by either the presence of other plasmids in the same bacterial host, by the co-transfer of other plasmids or by the presence of other plasmids in recipients [19,43]. All donor and recipient strains used in this study carry at least one plasmid (S1 Table). We investigated whether interactions with these plasmids might affect transfer of the ESBL-plasmids, using sequenced transconjugants from the 1st and 2nd gen *in vitro* experiments, and 8 clones per plasmid-donor pair from the *in vivo* conjugation assays on day seven.

We found multiple indications of plasmid interference in our conjugation experiments (Figs 2 and 5). Firstly, RE1, the only recipient without an IncFII-plasmid, received IncFII ESBL-plasmids (e.g. when mating with D5 and D6) at a higher rate than all other recipients. P6A_IncI/F of D6 did not transfer, and p5A_IncF of D5 only poorly, to recipient RS that carries a plasmid with IncFII/FIB replicons (Fig 2). On the other hand, plasmid p8A_IncF, carrying IncFIA/FIB-replicons, spread through RS recipient populations, despite RS having an IncFIB replicon (Figs 2 and 5). Because all but one of our conjugative plasmids carry multiple replicons [44], which is suggested to be a strategy to circumvent limitations to spread due to plasmid incompatibility, we argue that the plasmid incompatibility in our conjugation assays to might be permeable. Secondly, both *in vitro* and *in vivo,* transfer of ESBL-plasmid p8A_IncF to recipient RE3 resulted in the loss of the resident F-plasmid pRE3A_IncF (S8 Fig): in 6/11 re-sequenced transconjugants it was lost, retained in 2/11 and in 3 /11 it showed very low frequency in the sequencing reads. Similarly, pRE1A_IncI was lost by recipient RE1, irrespective of the donor strain and whether it acquired an IncI1 or IncF ESBL-plasmid (S8 Fig). The cause and effect we can not trace back: loss of resident plasmids could have allowed

subsequent acquisition of ESBL-plasmids, or the acquisition of ESBL-plasmids could have been

followed by the loss of resident plasmids. Either way, despite this incompatibility of resident

and newly-acquired ESBL-plasmids comparably high transconjugant frequencies could be

reached (Figs 2 and 5).

In all the investigated conjugations, we only observed a few isolated cases of co-travelling

plasmids, which we could not link to the observed variation in final transconjugant

frequencies. We found plasmids co-transferred from donors D1 and D8 but not from D4 (S8

Fig). Although all three donors D1, D4 and D8 carry ColRNAI plasmids, we observed its transfer

only from D1, alongside ESBL-plasmid p1A_IncF. *In vitro*, multiple plasmids were transferred

from D8 alongside the ESBL-plasmid p8A_IncF, depending on the recipient: plasmid

p8C_IncBOKZ was passed together with plasmid p8G_Col8282 to RE2, from which both

plasmids were passed across the species boundary to *S.* Typhimurium RS in the 2^{nd} generation

in vitro experiment. Plasmid p8C_IncBOKZ also co-travelled to recipient RE1, but from there

it was not passed to any 2^{nd} generation recipient. No mutations were found on any of the co-

transferred plasmids. The prophage p8B_p0111 (S8 Fig and S1 File) was passed to recipient

RE3 *in vitro* but not *in vivo* and showed a deletion in the side tail fiber proteins stfE_2, and

stfR in all three sequenced transconjugants. Despite the presence of their essential transfer

genes (S2 File), we did not detect transfer of resident plasmids pRE2B_IncF of recipient RE2

or pRSA_IncF of recipient RS, to any recipient of the 2^{nd} gen *in vitro* experiment.

Discussion

In this study, we demonstrated that transfer rates of ESBL-plasmids derived from clinical donors varied over five orders of magnitude and spread in the absence of antibiotics through various recipient populations. We confirm that, in the absence of external factors such as selection by antibiotics or competition against other bacterial species, conjugative transfer is the main determinant of the spread of ESBL-plasmids *in vitro* and *in vivo*. Like others, we found that ESBL-plasmids are not costly in our experiments [45,46]. When correcting for growth, we found that the increase in plasmid frequency was strongly correlated with plasmid transfer rate (Fig 4). This might hold true even for costly plasmids, as others have shown that transfer rates can surpass a fitness cost and allow plasmids to spread [26,47]. Foremost, we found that all plasmids carrying the required *tra* genes spread from various donors through a multitude of recipient populations. This suggests that conjugative plasmids have a strong drive to spread and that donor and recipient determinants play a lesser role. Since half of all plasmids can transfer via conjugation [20], it is tempting to speculate that the majority of these plasmids may spread in the absence of selection by antibiotics.

Although the estimates of our transfer rates are in line with results obtained with common laboratory strains [7,27], we have shown that it is crucial to use multiple and clinical strains to better understand the factors driving plasmid transfer. For instance, the pathogenic *S.* Typhimurium RS was able to receive 4 out of 5 conjugative plasmids. When acting as donor too, it could yield transconjugants at the highest efficiency compared to other recipients, particularly for IncI1 plasmids (Figs 2 and 5). This has important implications in the spread of antibiotic resistance plasmids, because the tissues of infected animals can serve as a reservoir for *S.* Typhimurium, which can lead to long-term potential for plasmid transfer into co-

colonizing Enterobacteriaceae [9]. An additionally important characteristic of clinical strains

is the multitude of plasmids they harbor. It has been shown that their presence in donors and

recipients may affect the transfer of the plasmid of interest. This has previously not been

taken into account [27]. Here, we observed variation in plasmid transfer rate depending also

on the recipient strain, and could recapitulate that some recipients are inherently more

permissive to plasmids. The frequently used lab strain *E. coli* K12 seems to be an exceptionally

good recipient, which may have led to a historical overestimation of transfer potential of

plasmids [7,27] .

We found that *in vitro* transfer dynamics are a good qualitative predictor of *in vivo* plasmid

transfer in a murine model, and discovered similar variation across donor and recipient

combinations in both methods. Furthermore, we highlight the previously underappreciated

speed with which ESBL-plasmids can spread through recipient populations *in vivo*. Plasmid

donors colonized the gut within 4 hours and transferred ESBL-plasmids within the first 24

hours in the absence of antibiotic selection. This is particularly important, because our *in vivo*

model recapitulates a natural situation in which *E. coli* is an abundant member of the

microbiota of the gut lumen [37], and interacts with an invading ESBL-bearing *E. coli* [48,49].

However, contrary to our mouse experiments, not all natural Enterobacteriaceae populations

reach densities of 10^8 CFU/g feces in farm animal or human guts [37]. This may lead to an

overestimation of actual plasmid spread in our study. For such cases, the role of inflammation

inducing pathogens such as *S.* Typhimurium may be especially important, because they can

induce blooms of Enterobacteriaceae, leading to increased population densities and thus

plasmid transfer [8]. The strains carrying clinical ESBL-plasmid rapidly colonized the mouse

gut and thereby outperformed the resident strain (S6 Fig). All but one of our strains associated

with ESBL-plasmids encode a T6SS in their genome (S3 Fig), which may explain their

competitive success. Thus, especially *in vivo*, the strain background of the plasmid-host

association is crucial to long-term plasmid persistence.

To obtain exact quantitative predictions of *in vivo* plasmid spread based on *in vitro* data,

detailed mathematical modelling will be needed. Fischer *et al*. [50] showed that a very simple

model that does not take into account varying plasmid transfer rates to host microbiota,

underestimates observed *in vivo* transfer rates. Given the high transfer proficiency of RS

observed in this study, a realistic mathematical model will likely require detailed

understanding of inter-species aspects of plasmid transfer. Furthermore, one will need to

consider the recently discovered role of long-term reservoirs formed by plasmid-carrying

invasive enteropathogens, such as *S.* Typhimurium, in plasmid spread [9].

Although our broad study includes many of the factors known to determine plasmid transfer,

the biological basis of the variation in plasmid transfer we found remains unknown. For

instance, we found large differences in plasmid spread between transfer from native and from

secondary bacterial hosts and a consistent 1000-fold increase in final transconjugant

frequency when donor and recipient were the same strain (self-self transfer). Dimitriu *et al.*

suggested compatibility of restriction-modification systems as a likely mechanism for

increased transfer rates in such cases [27]. Here, neither the presence of restriction-

modification systems nor of CRISPR-Cas showed a consistent connection to the observed

plasmid spread. Because donors of the 1st gen and 2nd gen *in vitro* experiment had the same

handling before the conjugation assay, i.e. thawing and growing overnight, it is unlikely that

there is a role of plasmid de-repression. Moreover, other factors such as surface exclusion or

adaptive mutations seem, based on our genomic analysis, at most minor. Although evolution experiments demonstrated extensive plasmid host co-adaptation upon plasmid acquisition [51,52], neither for the 24 hours *in vitro* nor for the 7 days *in vivo* conjugation assay, could we find any mutational changes that would suggest recipient-plasmid co-adaptation. Because these ESBL-plasmids are non-costly in all tested recipients, we think further co-adaptation might not be necessary for plasmid spread on a species level. An alternative explanation for the lack of genomic modification related to plasmid-host adaptation could be transcriptional changes in response to plasmid carriage [53]. To statistically test the relative significance of donor, plasmid, and recipient factors, as those investigated here, in determining plasmid transfer rates, even larger scale studies are needed. The feasibility of this has been demonstrated by Alamam et al. [54], who studied horizontal resistance transfer for more than 60.000 pairs of cell populations in parallel. We advocate for high-throughput screening of clinically relevant donor and recipient strains *in vitro*.

Efforts must be made to minimize the transmission of strains containing antibiotic resistance plasmids. Understanding the factors that make plasmid-host associations successful, can give insight into novel measures to limit their spread. For instance, previous studies have suggested to exploit systems like CRISPR-CAS, restriction modification or surface exclusion [19,22,55,56]. We found multiple cases of plasmid interference due to plasmid incompatibility, which may provide an important further avenue for future studies to block the transfer and persistence of plasmids within a population of bacteria [57]. The results presented here contribute to a growing scientific field aimed at determining fitness and plasmid persistence factors that could ultimately inspire the development of anti-plasmid spread strategies.

Materials and methods

Strains and growth conditions. As plasmid donor strains, we use 8 ESBL-plasmid positive clinical *E. coli* isolates, sampled from hospitalized patients at the University Hospital Basel, Switzerland [58]. Additionally, we worked with 4 ESBL-plasmid negative recipient strains: two clinical *E. coli* isolates from healthy patients, and mouse-derived *E.coli* and *Salmonella enterica* Typhimurium isolates. A comprehensive list of strains and associated plasmids can be found in S1 Table. Marker plasmids were introduced by electroporation, to mark recipients with either pACYC184 (New England Biolabs) encoding Chloramphenicol (Cm) resistance or pBGS18 [59] encoding Kanamycin (Kan) resistance. An exception was recipient RS, having chromosomal Cm resistance (*marT::cat*) [29]. Unless stated otherwise, we grew bacterial cultures at 37°C and under agitation (180 rpm) in lysogenic broth (LB) medium, supplemented with appropriates amounts of antibiotics (none, 100μg/mL Ampicillin (Amp) , 50μg/mL Cm and 50μg/mL Kan). We stored isolates in 25% glycerol at -80°C.

Antibiotic resistance profiling. We used microdilution assays with a VITEK2 system (bioMérieux, France) to determine the minimum inhibitory concentrations (MIC). MIC breakpoints for ESBL were interpreted according to EUCAST guidelines (v8.1). In addition, we confirmed the resistance mechanism of suspected isolates phenotypically using ROSCO disk assays (ROSCO diagnostica, Denmark) and/or genotypically with detection of CTX-M1 and -9 groups using the eazyplex Superbug assay (amplex, Germany).

Constriction of non-transferrable plasmids. We generated non transferrable plasmids, by deleting their origin of transfer (*oriT*) region, using the lambda red recombinase system with pKD4 as template for the Kan resistance marker[60]. The following primers were used (5' to

3'): For IncI1 plasmids (p4A_IncI) DIncI1_oriTnikA_f (GCA TAA GAC TAT GAT GCA CAA AAA

TAA CAG GCT ATA ATG GGT GTA GGC TGG AGC TGC TTC) and DIncI1_oriTnikA_r (CCT TCT CTT

TTT CGG AAT GAC TGC ATT CAC CGG AGA ATC CAT GGG AAT TAG CCA TGG TCC)[8] and for F

plasmids (p8A_IncF) D25_2_oriT-nikA-ko_vw (CCA TGA TAT CGC TCT CAG TAA ATC CGG GTC

TAT TTT GTA AGT GTA GGC TGG AGC TGC TTC) and D25_2_oriT-nikA-ko-rev (GTG CGG ACA

CAG ACT GGA TAT TTT GCG GAT AAA ATA ATT TAT GGG AAT TAG CCA TGG TCC). All mutants

were verified by PCR (IncI1_oriT _val_f: AGT TCC TCA TCG GTC ATG TC, IncI1_oriT _val_r: GAA

GCC ATT GGC ACT TTC TC, D25_oriT _val_fw: CAT ACA GGG ATC TGT TGT C and

D25_2_oriT_ver_rv: CAG AAT CAC TAT TCT GAC AC) and loss of transfer function.

Sequencing, assembly, annotation. We sequenced all donor and recipient strains on an

Illumina MiSeq machine (paired end, 2x250 bp), Oxford Nanopore MinION, and PacBio

Sequel. Unicycler [61] (v0.4.7) was used to produce hybrid assemblies (for D1,2,4,6,7,8 the

Oxford Nanopore – Illumina hybrid assembly proved most contiguous; D3,D5, RE1, RE2, RE3

were assembled using Pacbio Sequel – Illumina hybrid assembly). Manual curation involved

removing any contig smaller than 1kB, and sequences up to 5 kB that mapped to the own

chromosome. Quality control was performed by mapping the paired end Illumina reads to the

finished assemblies using samtools (v1.2) and bcftools (v1.7) [62,63]. The reference sequence

of RS was downloaded from NCBI Genbank under the accession numbers NZ_CP034230.1 and

NZ_CP034231.1.

We extracted an alignment of the concatenated core genome genes of all *E. coli* donor and

recipient strains, using chewBBACA for core genome Multi-Locus Sequence Typing (cgMLST)

against the Enterobase cgMLST scheme [64,65]. The phylogenetic tree was inferred using

BEAST2 [66], with an HKY substitution model and a strong prior on the mutation rate of *E. coli*

(10^{-4} mutations per genome per generation, as estimated by Wielgoss et al. [67]).

We performed bacterial annotation using Prokka [68], and ST calling using mlst (Torsten

Seemann, https://github.com/tseemann/mlst), which makes use of the PubMLST website

(https://pubmlst.org/) developed by Keith Jolley [69]. Phylogroups were assigned using

ClermonTyper [70].

We determined genomic features using a range of bioinformatic tools, and by BLAST

comparison against various curated databases. Plasmid replicons and resistance genes were

determined using abricate (Torsten Seemann, https://github.com/tseemann/abricate) with

the PlasmidFinder [71] and ResFinder [72] databases respectively. Phages were located using

PHASTER [73] (listing only those marked as "complete"), type 6 secretion systems using

SecReT6 [74], virulence genes using the Virulence finder database [75], toxin-antitoxin

systems using the database TADB 2.0 [76], and CRISPR-Cas loci using CRISPRCasFinder [77].

Restriction-modification systems were determined by using grep on the term 'restriction' in

the prokka general feature format (GFF) files.

To determine the presence/absence of IncF and IncI1 transfer genes, we constructed our own

database to compare against. IncF transfer genes were taken from the supplementary

material of Fernandez-Lopez et al. [78], IncI1 transfer genes were copied from R64 using the

annotations by Komano et al. [38], and R621a annotated by Takahashi et al. [39].

***In vitro* conjugation assays**. We determined the final frequency of the recipient population

that obtained an ESBL-plasmid (T/(R+T)) in a high throughput, 96-well plate-based assay, with

two replica blocks per experiment. The experiments with *E. coli* recipients and the *S.*

Typhimurium recipient RS were performed independently. Donor and recipient grew over

night with or without ampicillin (100μg/mL), respectively. We washed the independent

overnight cultures (n=5-6 per strain) by spinning down and resuspending and added ~1μL of

6.5-fold diluted donor and recipient cultures into 150μL fresh LB with a pin replicator (total

~1000-fold dilution, aiming to reach approximately a 1:1 ratio of donor and recipient). These

mating populations grew for 23 hours in the absence of antibiotics and were only shaken prior

to hourly optical density (OD) measurements (Tecan NanoQuant Infinite M200 Pro). To

determine the final cell densities, we plated the mating cultures at the end of the conjugation

assay on selective plates. In the 1st gen *in vitro* experiment, where the clinical strains

transferred their native plasmids to recipients, we selected for donors+transconjugants with

Amp, for recipients+transconjugants with Cm (*E. coli* recipients carried pACYC184-Cm and RS

chromosomal *marT::cat*) and for transconjugants with their combination. Four replicates had

to be excluded because the satellite colonies, a common phenomenon when plating on Amp,

could not be distinguished from true transconjugant colonies. For the 2nd gen *in vitro*

experiment, we chose a subset of transconjugants generated in the 1st gen *in vitro* experiment

that we used as plasmid donors. Transconjugants and recipients of the clone type EC3 were

omitted because of the size of the experiment, as well as transconjugant D1-EC2, carrying

p1B_IncI, because it could not be regrown from glycerol stock. We selected for donor with

Cm, for recipient with Kan (recipients carried pBGS18-Kan) and for transconjugants with

Kan+Amp. With this approach, we were able to detect transconjugant populations if they

were greater than 10^{-8} cells per mL.

Alongside the second replica block of the 1st gen *in vitro* experiment, we performed a plate-

mating control experiment, to assess the extent of surface mating in our conjugation assay.

After washing and diluting the independent overnight cultures of the second replica block,

we grew clinical donors and *E. coli* recipients in isolation for the duration of the conjugation

experiment. Donors and recipients were mixed only before plating 20µL of the undiluted

mixed population (n_{total} = 1-3, S4 Table). In conjugations involving donors D4-D6 and D8, the

transconjugant CFU from plate-mating, was maximally ~15 % of the CFU observed in the 1st

gen *in vitro* experiment. For D1 in combination with RE1 and RE3, we found substantial

amounts of plate-mating, as high as fractions of 35 and 85 % of the transconjugants resulting

from the 1st gen *in vitro* experiment, respectively.

***In vitro* plasmid cost and growth rate assays**. To investigate the effect of plasmid carriage on

bacterial growth, we measured the growth rate of transconjugants and corresponding

recipients. Per D and R combination, we used three clonal transconjugants (4 replicates each),

which were obtained from independent mating populations of the 1st gen *in vitro* experiment.

We grew bacterial cultures in absence of antibiotics overnight and diluted them 150-fold by

transfer with a pin replicator to a 96-well plate, containing 150µL fresh LB per well. The

cultures grew without shaking and we estimated growth rates of recipients and

transconjugants based on 10 OD measurements over 24 hours and expressed plasmid cost as

their relative growth rates. Because transconjugants have passed the 1st gen *in vitro*

experiment, we had to exclude that an effect of plasmid carriage could be obscured by

transconjugants having adapted to the LB growth conditions. Thus, we exposed recipients (n

= 4) to the same culture handling as a conjugation assay would and found their growth rates

not to be different from the naïve recipient strains used in the plasmid cost assay (S9 Fig). We

had to exclude RE2 carrying p1B_IncI because it could not be regrown from glycerol stock and

further excluded RE3 carrying p1B_IncI because we had not stored 3 independently arisen

transconjugants.

For other growth measurements, we followed a similar approach of 150-fold dilution of

overnight cultures by transfer with a pin replicator to a 96-well plate, containing 150µL fresh

LB per well. Growth assays were performed in the plate reader with hourly OD

measurements. We estimated growth rates using the R package Growthcurver [79].

Infection experiments. We have previously established a murine model for enterobacterial

pathogen infection (Barthel et al, 2003) that has been used to measure conjugative plasmid

transfer [8,9,29,30]. For conjugation experiments, 8-16 week old C57BL/6 mice containing an

oligo microbiota [42], which allows colonization of approximately 10^8 *E. coli* per gram feces,

densities of *E. coli* that can be found in the guts of humans and animals [37], were infected

orogastrically with ~5x10^7 CFU of RE3, carrying marker plasmid pACYC184. 24 hours later,

~5x10^7 CFU of either D4, D7, or D8 were introduced orogastrically (Fig 6A). Feces were

collected daily, homogenized in 1 ml of PBS with a steel ball by a Tissue Lyser (Qiagen) at 25

Hz for 1 min. Samples were enumerated for bacterial populations on MacConkey media

containing the appropriate antibiotics (selection for donors+transconjugants with Amp, for

recipients+transconjugants with Cm and for transconjugants with their combination) and final

transconjugant frequencies T/(R+T) calculated.

For competition experiments, 8-16 week old C57BL/6 oligo microbiota mice were infected

orogastrically with a 1:1 mixture of both competitor strains (~5×10^7 CFU total. Feces were

collected and bacterial populations were enumerated daily as above. Bacteria were

enumerated on MacConkey agar containing chloramphenicol and replica-plated on media

containing chloramphenicol, kanamycin, and ampicillin to enumerate the locked

transconjugants. Competitive index is calculated by dividing the recipient population by the

locked transconjugant population.

Prior to all infections, overnight cultures in LB containing the appropriate antibiotics were

subcultured for 4 hours at 37°C without antibiotics (1:20 dilution) to ensure equal densities

of bacteria. Cells were washed in PBS and introduced into mice.

All infection experiments were approved by the responsible authority

(Tierversuchskommission, Kantonales Veterinäramt Zürich, license 193/2016).

Resequencing. We re-sequenced isolates from the 2nd generation *in vitro* experiment as well

as the *in vivo* transfer experiment, to study the genetic contribution to the observed transfer

rates. Resequencing was performed on an Illumina MiSeq (paired end, 2x150 bp). Reads were

mapped to the closed assemblies of respective donor and recipient strains using the breseq

pipeline (v 0.32.0). Mutations or indels shared by all re-sequenced strains were treated as

ancestral (S5 Table).

Statistical analyses. We performed statistical analyses using the software R (version 3.4.2).

The effect of donor, recipient and plasmids on the final transconjugant frequency was

analyzed with either a two-way ANOVA (1st gen *in vitro* experiment, factors donor-plasmid

pair and recipient) or a three-way ANOVA (2nd gen *in vitro* experiment, factors donor, plasmid,

recipient). For the 1st gen *in vitro* experiment, we excluded from this analysis strain-plasmid

pairs which did not result in transconjugants in the 1st gen *in vitro* experiment (D2, D3, D7)

and *S.* Typhimurium recipient RS. When single replicates for a given donor-recipient

combination lacked transconjugants (D5 and D6), we assigned these replicates a final

transconjugant frequency at the detection limit of 10^{-8}. The same data was used to perform

the correlation of transconjugant frequency and transfer rate. Data of the 2nd gen *in vitro*

experiment was not fully factorial. To enable testing of higher-order interactions here, we

therefore performed two 3-way ANOVAs: one excluding plasmid p1B_IncI and one excluding

donor RE2, for which we had to take the two replica blocks into account: $P < 0.001$). For two

replica populations of donor RE2 transferring p4A_IncI to RE1, we were not able to detect

donors on plates and assigned them a donor population size at the detection limit of 1 colony

per plate (10^8 cells/mL). For two replica populations (RS self-self transfer with p1B_IncI), we

had higher counts on plates selecting for transconjugants than on plates selecting for

recipients+transconjugants and replaced the resulting negative CFU/mL for recipients with 1

CFU/mL.

To express the final fraction of transconjugants as a plasmid transfer rate, we used the cell

densities of donor, recipient, transconjugant and their sum (CFU/mL), and growth rates of

mating populations, both estimated in the 1st gen *in vitro* experiment. Introducing these

values to the mass action model as previously described by Simonsen *et al.* [41] yields the

transfer rate constant γ $(mL(CFU * t)^{-1})$. For statistical comparisons derived from *in vivo*

experiments, Kruskal-Wallis tests were performed with Dunn's multiple test correction using

GraphPad Prism Version 8 for Windows.

Acknowledgements

704   We would like to acknowledge the members of the Bonhoeffer, Hall, Hardt, Diard, Egli,

705   Stadler, and Ackermann labs for helpful discussions. We would also like to thank the staff at

706   the RCHCI and EPIC animal facilities at ETH Zurich, particularly Sven Nowok; as well as the NGS

707   team and Dr. Daniel Wuethrich of the Clinical Bacteriology and Mycology department at the

708   University Hospital Basel for their excellent technical support.

References

1.   Sommer MOA, Munck C, Toft-Kehler RV, Andersson DI. Prediction of antibiotic

resistance: Time for a new preclinical paradigm? Nat Rev Microbiol. 2017;15(11):689–

96.

2.   Pitout JD, Laupland KB. Extended-spectrum β-lactamase-producing

Enterobacteriaceae: an emerging public-health concern. Lancet Infect Dis. 2008

Mar;8(3):159–66.

3.   Cassini A, Högberg LD, Plachouras D, Quattrocchi A, Hoxha A, Simonsen GS, et al.

Attributable deaths and disability-adjusted life-years caused by infections with

antibiotic-resistant bacteria in the EU and the European Economic Area in 2015: a

population-level modelling analysis. Lancet Infect Dis. 2019;19(1):56–66.

4.   F Wozniak RA, Waldor MK. Integrative and conjugative elements: mosaic mobile

genetic elements enabling dynamic lateral gene flow. Nat Publ Gr. 2010;8.

5.   Knudsen PK, Gammelsrud KW, Alfsnes K, Steinbakk M, Abrahamsen TG, Müller F, et

al. Transfer of a bla CTX-M-1-carrying plasmid between different Escherichia coli

strains within the human gut explored by whole genome sequencing analyses. Sci Rep

[Internet]. 2018;8(1):280. Available from: http://www.nature.com/articles/s41598-

017-18659-2

6.   Liebana E, Batchelor M, Hopkins KL, Clifton-Hadley FA, Teale CJ, Foster A, et al.

Longitudinal farm study of extended-spectrum beta-lactamase-mediated resistance. J

Clin Microbiol. 2006 May;44(5):1630–4.

7.   Lopatkin AJ, Huang S, Smith RP, Srimani JK, Sysoeva TA, Bewick S, et al. Antibiotics as

a selective driver for conjugation dynamics. Nat Microbiol [Internet]. 2016 Jun

11;1(6):16044. Available from: http://www.nature.com/articles/nmicrobiol201644

8.   Stecher B, Denzler R, Maier L, Bernet F, Sanders MJ, Pickard DJ, et al. Gut

inflammation can boost horizontal gene transfer between pathogenic and commensal

Enterobacteriaceae. Proc Natl Acad Sci [Internet]. 2012;109(4):1269–74. Available

from: http://www.pnas.org/cgi/doi/10.1073/pnas.1113246109

9.   Bakkeren E, Huisman JS, Fattinger SA, Hausmann A, Furter M, Egli A, et al. Salmonella

persisters promote the spread of antibiotic resistance plasmids in the gut. Nature

[Internet]. 2019 Sep 4;573(7773):276–80. Available from:

http://www.nature.com/articles/s41586-019-1521-8

10.   Davies J, Davies D. Origins and Evolution of Antibiotic Resistance. Microbiol Mol Biol

Rev. 2010;74(3):1092–2172.

11.   San Millan A. Evolution of Plasmid-Mediated Antibiotic Resistance in the Clinical

Context. Trends Microbiol. 2018;26(12):978–85.

12.   Dunn SJ, Connor C, McNally A. The evolution and transmission of multi-drug resistant

Escherichia coli and Klebsiella pneumoniae: the complexity of clones and plasmids.

Curr Opin Microbiol. 2019 Oct;51:51–6.

13.   Paterson DL, Bonomo RA. Extended-spectrum beta-lactamases: a clinical update. Clin

Microbiol Rev. 2005 Oct;18(4):657–86.

14.   Rozwandowicz M, Brouwer MSM, Fischer J, Wagenaar JA, Gonzalez-Zorn B, Guerra B,

et al. Plasmids carrying antimicrobial resistance genes in Enterobacteriaceae. J

Antimicrob Chemother. 2018;73(5):1121–37.

15.   Carattoli A. Resistance plasmid families in Enterobacteriaceae. Antimicrob Agents

Chemother. 2009;53(6):2227–38.

16.   Poirel L, Potron A, Nordmann P. OXA-48-like carbapenemases: The phantom menace.

J Antimicrob Chemother. 2012;67(7):1597–606.

17.   Bajaj P, Singh NS, Virdi JS. Escherichia coli β-Lactamases: What Really Matters. Front

758   Microbiol. 2016 Mar;7:417.

759   18.   Tacconelli E, Carrara E, Savoldi A, Kattula D, Burkert F. Global priority list of antibiotic-

760   resistant bacteria to guide research, discovery, and development of new antibiotics.

761   19.   Frost LS, Ippen-Ihler K, Skurray RA. Analysis of the sequence and gene products of the

762   transfer region of the F sex factor. Vol. 58, Microbiological Reviews. 1994. p. 162–210.

763   20.   Smillie C, Garcillan-Barcia MP, Francia M V., Rocha EPC, de la Cruz F. Mobility of

764   Plasmids. Microbiol Mol Biol Rev. 2010;74(3):434–52.

765   21.   Dionisio F, Zilhão R, Gama JA. Interactions between plasmids and other mobile

766   genetic elements affect their transmission and persistence. Vol. 102, Plasmid. 2019.

767   p. 29–36.

768   22.   Purdy D, O'Keeffe TAT, Elmore M, Herbert M, McLeod A, Bokori-Brown M, et al.

769   Conjugative transfer of clostridial shuttle vectors from Escherichia coli to Clostridium

770   difficile through circumvention of the restriction barrier. Mol Microbiol. 2002

771   Oct;46(2):439–52.

772   23.   Thomas CM, Nielsen KM. Mechanisms of, and barriers to, horizontal gene transfer

773   between bacteria. Vol. 3, Nature Reviews Microbiology. Nature Publishing Group;

774   2005. p. 711–21.

775   24.   Doron S, Melamed S, Ofir G, Leavitt A, Lopatina A, Keren M, et al. Systematic

776   discovery of antiphage defense systems in the microbial pangenome. Science (80-).

777   2018;359(6379):0–12.

778   25.   Harrison E, Guymer D, Spiers AJ, Paterson S, Brockhurst MA. Parallel Compensatory

779   Evolution Stabilizes Plasmids across the Parasitism-Mutualism Continuum. Curr Biol.

780   2015;25(15):2034–9.

26. Lopatkin AJ, Meredith HR, Srimani JK, Pfeiffer C, Durrett R, You L. Persistence and

reversal of plasmid-mediated antibiotic resistance. Nat Commun. 2017;8(1).

27. Dimitriu T, Marchant L, Buckling A, Raymond B. Bacteria from natural populations

transfer plasmids mostly towards their kin.

28. Hardiman CA, Weingarten RA, Conlan S, Khil P, Dekker JP, Mathers AJ, et al.

Horizontal Transfer of Carbapenemase-Encoding Plasmids and Comparison with

Hospital Epidemiology Data. 2016;

29. Diard M, Bakkeren E, Cornuault JK, Moor K, Hausmann A, Sellin ME, et al.

Inflammation boosts bacteriophage transfer between *Salmonella* spp. Science (80-).

2017;355(6330):1211–5.

30. Moor K, Diard M, Sellin ME, Felmy B, Wotzka SY, Toska A, et al. High-avidity IgA

protects the intestine by enchaining growing bacteria. Nature [Internet].

2017;544(7651):498–502. Available from: http://dx.doi.org/10.1038/nature22058

31. Conlan S, Park M, Deming C, Thomas PJ, Young AC, Coleman H, et al. Plasmid

Dynamics in KPC-Positive Klebsiella pneumoniae during Long-Term Patient

Colonization. MBio. 2016 Jun;7(3):e00742-16.

32. Hagbø M, Ravi A, Angell IL, Sunde M, Ludvigsen J, Diep DB, et al. Experimental

support for multidrug resistance transfer potential in the preterm infant gut

microbiota. Pediatr Res. 2019;

33. MacLean RC, San Millan A. The evolution of antibiotic resistance. Science (80-)

[Internet]. 2019 Sep 13;365(6458):1082–3. Available from:

http://www.sciencemag.org/lookup/doi/10.1126/science.aax3879

34. Lahlaoui H, Ben Haj Khalifa A, Ben Moussa M. Epidemiology of Enterobacteriaceae

producing CTX-M type extended spectrum β-lactamase (ESBL). Med Mal Infect

[Internet]. 2014;44(9):400–4. Available from:

http://dx.doi.org/10.1016/j.medmal.2014.03.010

35.   Wotzka SY, Kreuzer M, Maier L, Zünd M, Schlumberger M, Nguyen B, et al. Microbiota

stability in healthy individuals after single-dose lactulose challenge—A randomized

controlled study. PLoS One. 2018 Oct;13(10):e0206214.

36.   Clermont O, Christenson JK, Denamur E, Gordon DM. The Clermont Escherichia coli

phylo-typing method revisited : improvement of specificity and detection of new

phylo-groups. 2013;5:58–65.

37.   Tenaillon O, Skurnik D, Picard B, Denamur E. The population genetics of commensal

Escherichia coli. Vol. 8, Nature Reviews Microbiology. Nature Publishing Group; 2010.

p. 207–17.

38.   Komano T, Yoshida T, Narahara K, Furuya N. The transfer region of Incl1 plasmid R64:

Similarities between R64 tra and Legionella icm/dot genes. Mol Microbiol.

2000;35(6):1348–59.

39.   Takahashi H, Shao M, Furuya N, Komano T. The genome sequence of the

incompatibility group Iγ plasmid R621a: Evolution of IncI plasmids. Plasmid [Internet].

2011;66(2):112–21. Available from: http://dx.doi.org/10.1016/j.plasmid.2011.06.004

40.   Koraimann G. Spread and Persistence of Virulence and Antibiotic Resistance Genes: A

Ride on the F Plasmid Conjugation Module. EcoSal Plus. 2018;8(1).

41.   Simonsen L, Gordon DM, Stewart FM, Levin BR. Estimating the rate of plasmid

transfer: an end-point method. J Gen Microbiol [Internet]. 1990;136(11):2319–25.

Available from:

http://mic.microbiologyresearch.org/content/journal/micro/10.1099/00221287-136-

11-2319

42.   Brugiroux S, Beutler M, Pfann C, Garzetti D, Ruscheweyh HJ, Ring D, et al. Genome-

guided design of a defined mouse microbiota that confers colonization resistance

against Salmonella enterica serovar Typhimurium. Nat Microbiol [Internet].

2016;2(2):1–12. Available from: http://dx.doi.org/10.1038/nmicrobiol.2016.215

43.   Gama JA, Zilhão R, Dionisio F. Conjugation efficiency depends on intra and

intercellular interactions between distinct plasmids: Plasmids promote the

immigration of other plasmids but repress co-colonizing plasmids. Plasmid [Internet].

2017;93(August):6–16. Available from:

http://dx.doi.org/10.1016/j.plasmid.2017.08.003

44.   Villa L, García-Fernández A, Fortini D, Carattoli A. Replicon sequence typing of IncF

plasmids carrying virulence and resistance determinants. J Antimicrob Chemother.

2010;65(12):2518–29.

45.   Ranjan A, Scholz J, Semmler T, Wieler LH, Ewers C, Müller S, et al. ESBL-plasmid

carriage in E. coli enhances in vitro bacterial competition fitness and serum resistance

in some strains of pandemic sequence types without overall fitness cost. Gut Pathog.

2018 Dec;10(1):24.

46.   Cottell JL, Webber MA, Piddock LJ V. Persistence of Transferable Extended-Spectrum-

β-Lactamase Resistance in the Absence of Antibiotic Pressure. Antimicrob Agents

Chemother [Internet]. 2012 Sep;56(9):4703–6. Available from:

http://aac.asm.org/lookup/doi/10.1128/AAC.00848-12

47.   Tazzyman SJ, Bonhoeffer S. Fixation probability of mobile genetic elements such as

plasmids. Theor Popul Biol [Internet]. 2013;90:49–55. Available from:

http://dx.doi.org/10.1016/j.tpb.2013.09.012

48.   Tängdén T, Cars O, Melhus Å, Löwdin E. Foreign travel is a major risk factor for

853   colonization with Escherichia coli producing CTX-M-type extended-spectrum β-

854   lactamases: A prospective study with Swedish volunteers. Antimicrob Agents

855   Chemother. 2010;54(9):3564–8.

856 49. Kuenzli E, Jaeger VK, Frei R, Neumayr A, DeCrom S, Haller S, et al. High colonization

857   rates of extended-spectrum β-lactamase (ESBL)-producing Escherichia coli in Swiss

858   Travellers to South Asia- a prospective observational multicentre cohort study looking

859   at epidemiology, microbiology and risk factors. BMC Infect Dis. 2014;14(1):1–10.

860 50. Fischer EAJ, Dierikx CM, van Essen-Zandbergen A, Mevius D, Stegeman A, Velkers FC,

861   et al. Competition between Escherichia coli with and without

862   Extended-Spectrum-Beta-Lactamase-carrying plasmids in the broiler chicken gut. Appl

863   Environ Microbiol [Internet]. 2019;(June):AEM.00892-19. Available from:

864   http://aem.asm.org/content/early/2019/06/24/AEM.00892-19.abstract

865 51. Stalder T, Rogers LM, Renfrow C, Yano H, Smith Z, Top EM. Emerging patterns of

866   plasmid-host coevolution that stabilize antibiotic resistance. Sci Rep. 2017;7(1):1–10.

867 52. De Gelder L, Williams JJ, Ponciano JM, Sota M, Top EM. Adaptive plasmid evolution

868   results in host-range expansion of a broad-host-range plasmid. Genetics.

869   2008;178(4):2179–90.

870 53. Buckner MMC, Saw HTH, Osagie RN, McNally A, Ricci V, Wand ME, et al. Clinically

871   relevant plasmid-host interactions indicate that transcriptional and not genomic

872   modifications ameliorate fitness costs of Klebsiella pneumoniae carbapenemase-

873   carrying plasmids. MBio. 2018;9(2).

874 54. Alalam H, Graf FE, Palm M, Abadikhah M, Zackrisson M, Mattsson M, et al.

875   Conjugation factors controlling F-plasmid antibiotic resistance transmission. bioRxiv

876   [Internet]. 2018; Available from:

http://biorxiv.org/content/early/2018/03/05/271254.abstract

55.   Price VJ, Mcbride SW, Hullahalli K, Chatterjee A, Duerkop BA, Palmer KL. Enterococcus

faecalis CRISPR-Cas Is a Robust Barrier to Conjugative Antibiotic Resistance

Dissemination in the Murine Intestine. 2019;

56.   Marraffini LA, Sontheimer EJ. CRISPR interference limits horizontal gene transfer in

staphylococci by targeting DNA. Science. 2008 Dec;322(5909):1843–5.

57.   Kamruzzaman M, Shoma S, Thomas CM, Partridge SR, Iredell JR. Plasmid interference

for curing antibiotic resistance plasmids in vivo. PLoS One. 2017;12(2):1–20.

58.   Tschudin-Sutter S, Frei R, Schwahn F, Tomic M, Conzelmann M, Stranden A, et al.

Prospective Validation of Cessation of Contact Precautions for Extended-Spectrum

beta-Lactamase-Producing Escherichia coli. Emerg Infect Dis. 2016;22(6):1094–7.

59.   Spratt BG, J. Hedge P, Heesen S te, Edelman A, Broome-Smith JK. Kanamycin-resistant

vectors that are analogues of plasmids pUC8, pUC9, pEMBL8 and pEMBL9. Gene.

1986 Jan;41(2–3):337–42.

60.   Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in Escherichia

coli K-12 using PCR products. Proc Natl Acad Sci U S A. 2000;97(12):6640–5.

61.   Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome

assemblies from short and long sequencing reads. PLoS Comput Biol. 2017;13(6):1–

22.

62.   Li H. A statistical framework for SNP calling, mutation discovery, association mapping

and population genetical parameter estimation from sequencing data. Bioinformatics.

2011;27(21):2987–93.

63.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078–9.

64.   Silva M, Machado MP, Silva DN, Rossi M, Moran-Gilad J, Santos S, et al. chewBBACA:

902   A complete suite for gene-by-gene schema creation and strain identification. Microb

903   genomics. 2018;4(3):1–7.

904   65.   Alikhan NF, Zhou Z, Sergeant MJ, Achtman M. A genomic overview of the population

905   structure of Salmonella. PLoS Genet. 2018;14(4):1–13.

906   66.   Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu CH, Xie D, et al. BEAST 2: A Software

907   Platform for Bayesian Evolutionary Analysis. PLoS Comput Biol. 2014;10(4):1–6.

908   67.   Wielgoss S, Barrick JE, Tenaillon O, Cruveiller S, Chane-Woon-Ming B, Médigue C, et

909   al. Mutation Rate Inferred From Synonymous Substitutions in a Long-Term Evolution

910   Experiment With Escherichia coli. G3 Genes, Genomes, Genet [Internet].

911   2011;1(3):183–6. Available from:

912   http://g3journal.org/lookup/doi/10.1534/g3.111.000406

913   68.   Seemann T. Prokka: Rapid prokaryotic genome annotation. Bioinformatics.

914   2014;30(14):2068–9.

915   69.   Jolley KA, Maiden MCJ. BIGSdb: Scalable analysis of bacterial genome variation at the

916   population level. BMC Bioinformatics [Internet]. 2010; Available from:

917   http://www.biomedcentral.com/1471-2105/11/595

918   70.   Beghain J, Bridier-Nahmias A, Le Nagard H, Denamur E, Clermont O. ClermonTyping:

919   an easy-to-use and accurate in silico method for Escherichia genus strain phylotyping.

920   Microb Genomics. 2018;4(7):1–8.

921   71.   Carattoli A, Zankari E, Garcia-Fernandez A, Larsen MV, Lund O, Villa L, et al. In Silico

922   detection and typing of plasmids using plasmidfinder and plasmid multilocus

923   sequence typing. Antimicrob Agents Chemother. 2014;58(7):3895–903.

72.   Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al.

Identification of acquired antimicrobial resistance genes. 2012;(July):2640–4.

73.   Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster

version of the PHAST phage search tool. Nucleic Acids Res. 2016;44(W1):W16–21.

74.   Li J, Yao Y, Xu HH, Hao L, Deng Z, Rajakumar K, et al. SecReT6: A web-based resource

for type VI secretion systems found in bacteria. Environ Microbiol. 2015;17(7):2196–

202.

75.   Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, et al. Real-time

whole-genome sequencing for routine typing, surveillance, and outbreak detection of

verotoxigenic Escherichia coli. J Clin Microbiol. 2014;52(5):1501–10.

76.   Xie Y, Wei Y, Shen Y, Li X, Zhou H, Tai C, et al. TADB 2.0: An updated database of

bacterial type II toxin-antitoxin loci. Nucleic Acids Res. 2018;46(D1):D749–53.

77.   Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Néron B, et al.

CRISPRCasFinder, an update of CRISRFinder, includes a portable version, enhanced

performance and integrates search for Cas proteins. Nucleic Acids Res.

2018;46(W1):W246–51.

78.   Fernandez-Lopez R, de Toro M, Moncalian G, Garcillan-Barcia MP, de la Cruz F.

Comparative Genomics of the Conjugation Region of F-like Plasmids: Five Shades of F.

Front Mol Biosci. 2016 Nov;3:71.

79.   Sprouffske K, Wagner A. Growthcurver: an R package for obtaining interpretable

metrics from microbial growth curves. BMC Bioinformatics. 2016 Dec;17(1):172.

Fig 1. Phylogenetic tree of the *E. coli* donor and recipient strains, inferred using Bayesian

inference on a core genome alignment. The names at the tips are colored by *E. coli*

phylogroup. The *S.* Typhimurium recipient RS was not included in the phylogeny, but listed

here to allow comparison of the plasmid content. Blue rectangles indicate that a plasmid

carrying the given IncF or IncI incompatibility marker (replicon) was present in the strain. The

intensity of the shade of blue reflects the percent identity and coverage of the BLAST match

to the listed genes. A red dot indicates the ESBL-plasmid. If multiple replicons were found on

the same ESBL-plasmid this is indicated with open red circles.

Fig 2. ESBL-plasmids spread at variable rates in the absence of antibiotics (1st gen *in vitro*

experiment). Plasmid spread was measured as the final frequency of the recipient population

carrying the ESBL-plasmid (transconjugants), for three different *E. coli* (A) and one *S.*

Typhimurium (B) recipient populations. Circles represent independent replicates (n=4-6) and

the beams are mean values ± standard error of the mean (SEM). The detection limit was at ~10^{-}

^{8}.

Fig 3. No evidence for cost of plasmid carriage for transconjugants. Plasmid cost was measured

for ten strain-plasmid combinations, with three independently isolated transconjugants each (n

= 4; beams are mean values ± SEM). The growth rate of transconjugants and their plasmid free

complements was determined from independent cultures. Relative growth rate was calculated

by dividing the transconjugant growth rates by the mean growth-rate of plasmid-free strains.

Fig 4. Correlation of plasmid transfer rate and final transconjugant frequency estimated using the Simonsen formula [41]. Each data point results from the same liquid mating culture shown in Fig 2. Transconjugant frequencies and transfer rates can be found in the S3 Table.

Fig 5. Final transconjugant frequency depends on donor, recipient, and plasmid (2nd gen *in vitro* experiment). Eight transconjugants isolated from mating assays in the 1st gen *in vitro* experiment (Fig 2), used here as plasmid donor strains, transferred their plasmid to three different recipients. Circles represent independent replicates (n= 6), the beams are mean values ± SEM and different plasmids are indicated in color. The detection limit was at ~10^{-8}.

Fig 6*.* ESBL-plasmids can spread in the gut in the absence of selection. (A) Mouse model used for *in vivo* conjugation assays. These mice have a minimal microbiota that offers intermediate colonization resistance. Recipient RE3 was added one day before the plasmid donor. Populations were enumerated in faeces by selective plating. (B) Conjugation assay for three clinical donors and recipient RE3. (C) A subset of mice used in panel B were monitored for conjugation within the first 24 hours of infection. (B-C) Plasmid spread is reported as final transconjugant frequency. Dotted lines indicate the detection limit for selective plating. (D) Competition assay performed by infecting mice with a 1:1 mix of a "locked" transconjugant (*oriT* knockout) and recipient RE3. The competitive index was calculated by dividing the recipient population by the locked transconjugant population. (B, D) Kruskal-Wallis test $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), $p < 0.0001$ (****).

Fig 1.

Fig 2.

Fig 3.

Fig 4.

Fig 5.

Fig 6.

Supporting information

S1 Fig. Plasmid replicons. Presence (blue) / absence (white) of plasmid replicons. The intensity of the shade of blue reflects the percent identity and coverage of the BLAST match to the listed genes.

S2 Fig. Resistance genes. Presence (blue) / absence (white) of resistance genes. The intensity of the shade of blue reflects the percent identity and coverage of the BLAST match to the listed genes.

S3 Fig. Type 6 secretion systems and virulence genes. Presence (blue) / absence (white) of T6SS (defined by the presence of more than one prodigal T6SS-encoding gene in an operon) and virulence genes (defined by BLAST match to the listed genes; the shades of blue indicate the percent identity and coverage). The only strain with a pathogenic virulence profile (Fig 1.3) is D4, which carries chromosomally encoded intimin genes (*eae*, *tir*) commonly associated with Enteropathogenic *E. coli*. All strains, except RE3 and D1, encode for a type 6 secretion system (T6SS).

S4 Fig. Absolute population sizes of donors, recipients, and transconjugants for 1st gen in vitro experiment. Conjugation experiments were performed with natural donor strains bearing the indicated plasmid, and recipients (A) RE1 (B) RE2 (C) RE3 and (D) RS. Dotted lines indicate the detection limit by selective plating. Beams are mean values ± SEM.

S5 Fig. Absolute population sizes of donors, recipients, and transconjugants for 2nd gen *in vitro* experiment. Conjugation experiments were performed with 2nd gen donor strains bearing the indicated plasmid, and recipients (A) RE1 (B) RE2 and (C) RS. Dotted line indicates detection limit by selective plating. Beams are mean values ± SEM.

S6 Fig. Absolute population sizes of donors, recipients, and transconjugants for *in vivo* experiments in Fig 6. Each population is indicated by a unique colored symbol (see figure

legend). (A, B, C) Fecal loads of donor, recipients, and transconjugants in Figure 6B determined by selective plating. Dotted line indicates the detection limit.

S7 Fig. Plasmid transfer from native versus secondary host. We marked D1 with the same Cm-resistance plasmid (pACYC184) as the *E. coli* recipients in the 1^{st} gen conjugation experiment, to exclude pACYC184 having a major effect on transconjugants' ability to donate plasmids. Circles represent independent replicates (n= 5-6) and the beams are mean values ± SEM. The detection limit was at $\sim10^{-8}$.

S8 Fig. Overview of co-transferring plasmids, showing the changing plasmid composition across both *in vitro* conjugation assays (1^{st} gen is denoted as F_1, 2^{nd} gen as F_2). Plasmids are indicated in italics. Because all the resident plasmids of recipients are listed in F_1, for simplicity in F_2 only changing resident plasmids are shown. Plasmids that transfer or get lost are highlighted (blue = ESBL-plasmids, green = changes originating from F_0, red = changes originating from F_1). Plasmids transferred across generations are only indicated with lines and not listed as resident plasmids. Skulls mark the loss of resident plasmids.

S9 Fig. Control for transconjugant growth rates. Recipient strains (n = 4) that were used in the 1^{st} gen conjugation assay did not increase their growth rate compared to ancestral recipient strains. Thus, we can exclude that transconjugants have a growth advantage over ancestral recipients based on adaptation to laboratory conditions.

S1 Table. Strain overview. Overview of all strains used in this study, including their natural plasmid content and detected resistance genes. Replicon and resistance gene hits were recorded in this table if they had a coverage and percent identity of at least 70%.

S2 Table. Phenotypic resistance profile of donor strains. Minimum inhibitory concentration (µg/mL) measurements of ESBL donors used in this study.

S3 Table. Average transconjugant frequencies and average transfer rates in 1ˢᵗ gen *in vitro*

experiment.

S4 Table. Conjugation on plate.

S5 Table. Mutational changes that occurred during conjugation assays.

S1 File. Prophages and plasmid cointegration.

S2 File. Analyses to elucidate genomic effects on plasmid spread.

S3 File. Growth rates of all strains in the 1ˢᵗ generation *in vitro* experiments.