Intersubject consistent dynamic connectivity during natural vision revealed by functional MRI

Xin Di [1, 2], Bharat B Biswal [1, 2] *

1. Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, NJ, 07029, USA

2. School of Life Sciences and Technology, University of Electronic Science and Technology of China,

Chengdu, China

* Corresponding author:

Bharat B. Biswal, PhD

607 Fenster Hall, University Height

Newark, NJ, 07102, USA

bbiswal@yahoo.com

Abstract

The functional communications between brain regions are thought to be dynamic. However, it is usually

difficult to elucidate whether the observed dynamic connectivity is functionally meaningful or simply due

to noise during unconstrained task conditions such as resting-state. During naturalistic conditions, such

as watching a movie, it has been shown that local brain activities, e.g. in the visual cortex, are consistent

across subjects. Following similar logic, we propose to study intersubject correlations of the time courses

of dynamic connectivity during naturalistic conditions to extract functionally meaningful dynamic

connectivity patterns. We analyzed a functional MRI (fMRI) dataset when the subjects watched a short

animated movie. We calculated dynamic connectivity by using sliding window technique, and quantified

the intersubject correlations of the time courses of dynamic connectivity. Although the time courses of

dynamic connectivity are thought to be noisier than the original signals, we found similar level of

intersubject correlations of dynamic connectivity to those of regional activity. Most importantly, highly

consistent dynamic connectivity could occur between regions that did not show high intersubject

correlations of regional activity, and between regions with little stable functional connectivity. The

analysis highlighted higher order brain regions such as the default mode network that dynamically

interacted with posterior visual regions during the movie watching, which may be associated with the

understanding of the movie.

Keywords: default mode network; dynamic connectivity; intersubject correlation; movie connectome;

naturalistic condition; supramarginal gyrus

Highlights

- Intersubject consistency may provide a complementary approach to study brain dynamic connectivity

- Widespread brain regions showed highly consistent dynamic connectivity during movie watching, while these regions themselves did not show highly consistent regional activity

- Consistent dynamic connectivity often occurred between regions from different functional systems

1. Introduction

The functional communications between spatially remote brain regions, especially the dynamics of connectivity, is a key to understand brain functions (Bullmore and Sporns, 2012; Friston, 2011; Park and Friston, 2013). Recently, the dynamics of connectivity has drawn increasing interests of research, especially in resting-state (Allen et al., 2014; Fu et al., 2019, 2018; Hutchison et al., 2013). However, due to the unconstrained nature of resting-state, it is difficult to elucidate whether the observed changes of connectivity across sliding windows are due to real fluctuations of functional communications, or simply due to random fluctuations (Lindquist et al., 2014). Moreover, the blood-oxygen-level dependent (BOLD) signals measured by fMRI are sensitive to physiological noises, such as respiration, heartbeat (Teichert et al., 2010), and head motion (Power et al., 2012), which may give rise to spurious correlation estimates for short window.

One way to capture meaningful dynamic functional connectivity is to manipulate subjects' mental states during the course of scan, so that there is known reference for the changes of connectivity. For example, in a typical task-based fMRI study with blocked design, different task conditions are assigned as blocks. Therefore, the time courses of dynamic connectivity can be correlated with the task design to identify task related connectivity changes (Di et al., 2015; Rosenthal et al., 2017). An alternative approach is to expose the subjects with naturalistic stimuli, such as a short movie. Although there is no predefined references of dynamic connectivity changes, one may take advantage of the phenomenon of intersubject correlation to capture changes that are consistent across different subjects (Hasson et al., 2004; Nastase et al., 2019).

In the seminal study, Hasson and colleagues calculated intersubject correlations of the time series of BOLD signal (Figure 1A) when the subjects were watching a movie (Hasson et al., 2004). They demonstrated that several brain regions, especially the visual cortex, are highly correlated across subjects during the movie watching. We propose that similar approach can be applied to the time courses of dynamic connectivity to capture meaningful functional communication dynamics during natural vision. Specifically, dynamic connectivity is usually calculated using a sliding window approach, so that a time

series of dynamic connectivity can be obtained. The time courses of dynamic connectivity can then be

correlated across-subjects (Figure 1B). If the dynamic connectivity reflects real time functional

communications between regions that are caused by the viewing of natural stimuli, then the time courses

of dynamic connectivity from different subjects should somehow correlated. Therefore, we can apply

intersubject correlation method to identify meaningful dynamic communications between regions.

Figure 1 Illustration of the calculations of intersubject correlations of the time series of regional activity

(A) and the time courses of dynamic connectivity between two regions (B).

In the current study, we analyzed an fMRI dataset where the subjects were scanned when viewing

a short animated movie. The aim was to identify dynamic connectivity that were shared cross subjects

during the movie watching. In order to do so, we first performed regular intersubject correlation analysis

to identify brain regions that showed consistent regional activity. Given these regions, we adopted a seed-

based strategy to calculate dynamic connectivity between a seed region and every voxels in the brain. We

then evaluated and identified regions whose connectivity with the seed were consistent cross subjects.

Even though higher order association regions did not typically show high intersubject correlations of

regional activity (Hasson et al., 2004), their functional communications with lower order regions may be

consistent across subject following the narrative of the movie. We therefore hypothesized that

intersubject correlations of dynamic connectivity may be able to identify more widespread regions and

functional dynamics that are associated with the watching of the movie.

2. Materials and methods

2.1. Data and task

The fMRI data were obtained through openneuro (https://openneuro.org/; accession #: ds000228). Only

the data from adult subjects were analyzed. There were originally 33 adult subjects. Two subjects' data

were discarded because of poor brain coverage (subject #: sub-pixar123 and sub-pixar124), and two were

discarded due to large head motions (sub-pixar149 and sub-pixar150). As a result, a total of 29 subjects

were included in the current analysis (17 females). The mean age is 24.6 years old (18 to 39 years).

During the fMRI scan, the subjects watched a silent version of Pixar animated movie "Partly

Cloudy", which is 5.6 minutes long (https://www.pixar.com/partly-cloudy#partly-cloudy-1). Brain MRI

images were acquired on a 3-Tesla Siemens Tim Trio scanner using the standard Siemens 32-channel

head coil. Functional images were collected with a gradient-echo EPI sequence sensitive to BOLD

contrast in 32 interleaved near-axial slices (EPI factor: 64; TR: 2 s, TE: 30 ms, flip angle: 90°). The

voxel size were 3.13 mm isotropic, with 3 subjects with no slice gap and 26 subjects with 10% gap. 168

functional images were acquired for each subject, with four dummy scans collected before the real scans

to allow for steady-state magnetization. T1-weighted structural images were collected in 176 interleaved

sagittal slices with 1 mm isotropic voxels (GRAPPA parallel imaging, acceleration factor of 3; FOV:

256 mm). For more information about the dataset please refers to (Richardson et al., 2018).

2.2. FMRI data analysis

2.2.1. Preprocessing

FMRI data processing and analyses were performed using SPM12 and MATLAB (R2017b) scripts. A subject's T1 weighted structural image was first segmented into gray matter, white matter, cerebrospinal fluid, and other tissue types, and was normalized into standard Montreal Neurological Institute (MNI) space. The T1 images were then skull stripped based on the segmentation results. Next, all the functional images of a subject were realigned to the first image of the session and coregistered to the skull stripped T1 image of the same subject. Framewise displacement was calculated for the translation and rotation directions for each subject (Di and Biswal, 2015). Subjects who had maximum framewise displacement greater than 1.5 mm or 1.5^{o} were discarded from further analysis. See supplementary materials section S1 for additional analysis on the head motion effects. The functional images were then normalized to MNI space using the parameters obtained from the segmentation step with resampled voxel size of 3 x 3 x 3 mm^{3}. The functional images were then spatially smoothed using a Gaussian kernel of 8 mm. Lastly, a voxel-wise general linear model (GLM) was built for each subject to model head motion effects (Friston's 24-parameter model) (Friston et al., 1996), low frequency drift (1/128 Hz), and constant offset. The residuals of the GLM were saved as a 4-D image series, which were used for further intersubject correlation analysis. The residual time series were all mean centered because of the constant term included in the GLM.

Removing low frequency drifts in BOLD signals is a critical step for dynamic connectivity analysis (Leonardi and Van De Ville, 2015). Leonardi and Van De Ville have suggested a high-pass filter of 1/W Hz to avoid spurious dynamic connectivity caused by low-frequency fluctuations, where W represents the window length in the sliding window analysis. The high-pass filter of 1/128 Hz is the default in the GLM module in SPM. Given the window length of 60 s (30 TR) in the current analysis, we also applied high-pass filtering of 1/64 Hz in a supplementary analysis. The results are very similar to what using the 1 / 128 Hz high-pass filtering (see supplementary materials section S3).

2.2.2. Intersubject correlation analysis

The correlations of time series of either brain activity or dynamic connectivity are calculated between

pairs of subjects. If there are N subjects, then there will be N x (N-1) / 2 correlation coefficients. The

statistics of these correlations become tricky, because they are calculated from only N subjects, therefore

not independent. An alternative approach is leave-one-out (Nastase et al., 2019), where the time series of

one hold-out subject were correlated with the averaged time series of the remaining N – 1 subjects. The

averaged time series of N – 1 subjects were thought to reflect the consistent component rather than noisy

individual's time series. Therefore, the resulting correlations should be higher than the pair-wise

correlations. Another benefit is that this approach estimates one correlation for each subject, making

group level statistics easier. Therefore, we adopt the leave-one-out approach in the current analysis.

We first performed intersubject correlation analysis on regional activity time series. The

preprocessed BOLD time series were extracted for each voxel and subject in a gray matter mask. For a

given voxel, the time series of one subject was held out, and the averaged time series of the remaining

subject were calculated. Then the time series of the hold-out subject were correlated with the averaged

time series. This process was performed for every voxel and every subject, resulting in one correlation

map for one subject. The correlation maps were transformed into Fisher's z maps. Group level one

sample t test was then performed to identify regions whose intersubject correlations were consistently

greater than 0. However, the null hypothesis statistical significance testing may not provide much

information of the effect size. There may be only small but consistent correlations for each subject,

which could give rise to very high statistical significance in a one sample t test. Indeed, when doing such

null hypothesis statistical significance testing for intersubject correlation analysis, usually almost all the

brain regions will show somehow significant correlations (Chen et al., 2016). We are more interested and

focused on the real effect size, i.e. correlation coefficients, in our analysis. We therefore averaged the

Fisher's z maps, and transformed them back into r maps. The continuous r maps were shown in the

results section.

We next performed intersubject correlation analysis on dynamic connectivity using a seed-based

approach. Given that a set of brain regions showed high intersubject correlations of regional activity, we

defined these regions as seeds. We adopted a relatively high threshold of $r > 0.45$ for the averaged

intersubject correlation map of regional activity to isolate four visual related seeds. Two of them were

located in the medial and posterior portion of the occipital lobe, which mainly covered the lingual gyrus

and calcarine sulcus. The other two seeds were located bilaterally in the middle occipital gyrus and

extended to the middle temporal gyrus. We labeled them as left and right medial visual and lateral visual

seeds, respectively. In addition, we adopted a relatively low threshold of $r > 0.35$ to isolate the left and

right supramarginal gyrus seeds. The maps of the six seeds are available at:

https://identifiers.org/neurovault.collection:6245.

For each seed, we performed voxel-wise correlation analysis, i.e. calculating intersubject

correlations of dynamic connectivity between the seed and every voxel in the gray matter mask. For two

given time series from a seed and a voxel, we first applied sliding window technique to calculate dynamic

connectivity for each subject. The window length was set as 30 time points (60 s) (Nastase et al., 2019),

and the time step was set as 2 time point (4 s). Therefore, the time course of dynamic connectivity had 70

window steps. Next, we calculated correlations between the time courses of dynamic connectivity of a

given subject with the averaged dynamic connectivity of the remaining subjects for a given voxel. As a

result, there was one correlation map for each seed and each subject.

The r maps of correlations of dynamic connectivity were transformed into Fisher's z maps for

group level statistical analysis. Again, we simply calculated an averaged z map for a seed, and

transformed it back into r map. In addition, we performed group-level analysis to identify regions that

showed different dynamic connectivity patterns with different levels of seeds. Specifically, we calculated

contrast images from the Fisher's z maps for each subject representing the differences between specific

levels of seeds compared with the other seeds. For example, we calculated a contrast image using [1, 1, -

0.5, -0.5, -0.5, -0.5] on the six z maps of a subject to define the differences between the two medial visual

seeds and the remaining four seeds. The contrast images were entered into a one sample t test GLM using

nonparametric statistics in SnPM13 (Statistical NonParametric Mapping, http://warwick.ac.uk/snpm).

Resulting clusters were first formed at $p < 0.001$, and the cluster extend was thresholded using family-

wise error (FWE) corrected p < 0.0167 (0.05 / 3). The cluster level FWE threshold (0.0167) was chosen

to further account for the three levels of seeds (medial visual, lateral visual, and supramarginal seeds).

In addition to the voxel-based analysis, we also performed region of interest (ROI)-based analysis

for in-depth examinations of the dynamic connectivity effects. In addition to the six seeds, we included

three more regions that showed different intersubject correlations of dynamic connectivity with different

seeds. Specifically, they were the left precentral gyrus that showed higher intersubject correlations of

dynamic connectivity with the medial visual seeds, and the posterior cingulate cortex and medial

prefrontal cortex that showed higher intersubject correlations of dynamic connectivity with the

supramarginal gyrus seeds. The regions were defined based on the statistical significant clusters from the

group-level analysis. The maps of the three regions are available at:

https://identifiers.org/neurovault.collection:6245. The calculations of intersubject correlations of dynamic

connectivity were the same as the seed-based analysis.

The selections of sliding window length is nontrivial (Fu et al., 2014; Zhang et al., 2013). In

addition to the 30-TR window length, we also explored other window length of 10 TRs (20 s), 20 TRs (40

s), 40 TRs (80 s), 50 TRs (100 s), and 60 TRs (120 s). For each window length, we calculated

intersubject correlations of dynamic connectivity among the 9 ROIs.

2.2.3. Relations with other measures

We first compared the intersubject consistent dynamic connectivity with stable functional connectivity.

For each subject, we calculated correlation coefficients across the 9 ROIs, and transformed them into

Fisher's z. Then the z matrices were averaged across the 29 subjects, and transformed back into r values.

In addition, we calculated the consistent component of each ROI, i.e. averaging the time series across the

29 subjects. And then one single correlation matrix among the 9 ROIs was calculated.

Given the consistent component of the 9 ROIs, we also calculated dynamic connectivity between

pairs of ROIs using the same sliding window approach. The time courses of dynamic connectivity

calculated from the consistent component were compared with the averaged dynamic connectivity that

was calculated from each subject.

We further asked whether the observed intersubject consistent dynamic connectivity was driven

by the consistent component of regional activity, or by the subject-specific idiosyncratic component. To

do so, for each ROI, we regressed out the consistent component from each subject's time series, and

calculated dynamic connectivity from the residual time series for each subject. Intersubject correlations

of dynamic connectivity calculated from the residual time series were compared with those from the taw

time series.

Lastly, we calculated intersubject correlations of regional activity using the same sliding window

approach for the 9 ROIs. That is, for each ROI, intersubject correlation was calculated at each window,

resulting in a time course of intersubject consistency of regional activity in each of the ROI.

3. Results

3.1. Intersubject correlations of regional activity

We first calculated intersubject correlations of regional activity for every voxel in the brain during the

video watching (Figure 2A). The highest correlations were around 0.5. The major regions that had high

intersubject correlations were the visual cortex extending anterior to the fusiform gyrus and middle

temporal lobe. The bilateral supramarginal gyrus also showed high intersubject correlations. The

bilateral precentral gyrus also showed intersubject correlations, but the effect sizes were much smaller.

Figure 2A shows all the voxels with positive correlation values. It is noteworthy that many regions

showed very small intersubject correlations, including largely the prefrontal cortex and anterior temporal

lobe.

Figure 2 Intersubject correlation maps of regional activity (A) and dynamic connectivity with different

seeds (B through G). The seed regions were depicted in blue or green in respective rows. All voxels with

positive correlations are shown. The numbers at the bottom represent z or x coordinates in Montreal

Neurological Institute (MNI) space. LMV, left medial visual; RMV, right medial visual; LLV, left lateral

visual; RLV, right lateral visual; LSMG, left supramarginal gyrus; RSMG, right supramarginal gyrus. All

the maps are available at: https://identifiers.org/neurovault.collection:6245.

3.2. Intersubject correlations of dynamic connectivity

3.2.1 Seed-based analysis

We defined seed regions where there were high intersubject correlations of regional activity, which

included bilateral medial visual regions, lateral visual regions, and supramarginal gyrus. We next

calculated voxel-wise intersubject correlations of dynamic connectivity with the six seeds, respectively

(Figure 2B through 2G). There were widespread brain regions that showed intersubject consistent

dynamic correlations with different seeds. First of all, the effect sizes of the intersubject correlations of

dynamic connectivity, i.e. the correlation coefficients, were comparable to those in the intersubject

correlations of regional activity. Secondly, regions with intersubject correlations of dynamic connectivity

turned out to be more widespread and extended to the frontal and parietal regions that did not show high

intersubject correlations of regional activity. See supplementary materials section S2 for direct

comparisons between the intersubject correlations of dynamic connectivity and those of regional activity.

Thirdly, the left and right corresponding seeds showed similar dynamic connectivity patterns, but there

were substantial differences in the patterns of dynamic connectivity among medial visual, lateral visual,

and supramarginal gyrus seeds. In order to highlight specific brain regions that showed dynamic

connectivity with different seeds, we compared each pair of seeds with the remaining seeds using

nonparametric group-level model (Figure 3 and Table 1). The medial visual seeds showed more

consistent dynamic connectivity with the left precentral gyrus and occipital regions compared with other

seeds. The lateral visual seeds showed more consistent dynamic connectivity with several visual regions

compared with the other seeds. In contrast, the supramarginal seeds showed consistent dynamic

connectivity with the precuneus/posterior cingulate gyrus and medial prefrontal cortex compared with the

other seeds, which basically formed the default mode network.

Figure 3 Differential intersubject correlations of dynamic connectivity among the medial visual, lateral

visual, and supramarginal gyrus seeds (depicted on the left). All maps were thresholded at $p < 0.001$, and

cluster thresholded at $p < 0.0167$ (0.05 / 3) after family-wise error (FWE) correction using nonparametric

method. MV, medial visual; LV, lateral visual; and SM, supramarginal gyrus. Unthresholded statistical

maps are available at: https://identifiers.org/neurovault.collection:6245.

3.2.2. Relations with stable functional connectivity

In order to better understand and interpret the dynamic connectivity and regional functions, we further

calculated different types of connectivity measures among a set of regions of interest. In addition to the

six seeds, we defined left precentral gyrus, posterior cingulate cortex, and medial prefrontal cortex ROIs

that showed different dynamic connectivity with different seeds. Among the 9 regions, we calculated

regular mean functional connectivity (Figure 4A) and connectivity derived from the consistent

components across the 29 subjects (Figure 4B). These two correlation matrices looked similar, and

clearly showed three clusters of brain regions. The first four regions were all visual. The fifth to seventh

regions were the bilateral supramarginal gyrus, and lateralized frontal region, which were all high order

association brain regions. And the last two regions were part of the default mode network, which showed

negative correlations with the association regions in the consistent component correlations.

Figure 4 Correlation matrices among the 9 regions of interest (ROI) using different methods. A) Mean

functional connectivity across the 29 subjects. B) Correlations of the consistent component of each ROI

(averaged time series across the 29 subjects). C) Intersubject correlations of dynamic connectivity. LMV,

left medial visual; RMV, right medial visual; LLV, left lateral visual; RLV, right lateral visual; LSMG,

left supramarginal gyrus; RSMG, right supramarginal gyrus; LPCG, left precentral gyrus; PCC, posterior

286   cingulate cortex; and MPFC, medial prefrontal cortex.

The intersubject consistent dynamic connectivity matrix (Figure 4C) was largely different from

the two stable correlation matrices. Some high consistent dynamic connectivity was observed within the

visual regions. The highest correlation was between the left and right medial visual regions ($r = 0.70$). In

contrast, many consistent dynamic connectivity were shown between different functional networks, where

there were virtually none or even negative stable correlations. Specifically, the medial visual regions

showed high consistent dynamic connectivity with the left precentral gyrus ROI. The highest intersubject

correlation was 0.56 between left medial visual region and left precentral gyrus. The default mode

regions and supramarginal regions also showed high consistent dynamic connectivity. The highest

correlation was 0.64 between the posterior cingulate cortex and right supramarginal gyrus. It is

noteworthy that these regions generally showed negative stable correlations in Figure 4B.

Lastly, we analyzed the time courses of dynamic connectivity for the above mentioned pairs of

regions (Figure 5). The dynamic connectivity between left and right medial visual regions was in general

high, which is consistent with the results of stable connectivity. But it can be seen that the connectivity

level went down during the first half of windows, and continued with two cycles of up and down

fluctuations. The fluctuations rather than a monotonic linear trend suggest that the dynamics of

connectivity is not simply due to sensory habituations. The left medial visual region and left precentral

gyrus did not show high level of correlations in general. But it had small positive correlations at the

beginning of the run, went down to around zero, and then went back to small positive correlations. What

is more interesting is the dynamic connectivity between the right supramarginal gyrus and posterior

cingulate cortex, where the connectivity switched between positive and negative values during the whole

course.

Figure 5 Time courses of dynamic connectivity (Fisher's z) for three pairs of regions of interest. Each

thinner line represents the time course of one subject, and the thicker red lines represent the averaged time

courses. LMV, left medial visual; RMV, right medial visual; LPCG, left precentral gyrus; RSMG, right

313   supramarginal gyrus; PCC, posterior cingulate cortex.

3.2.3. Relations with the consistent component

A subsequent question is that whether the observed intersubject consistent dynamic connectivity is driven

by the consistent component of regional activity across subject, or by subject-specific idiosyncratic

components. We then regressed out the consistent component for each subject's time series and

calculated intersubject correlations of dynamic connectivity from the residual time series (Figure 6B).

Compared with the intersubject correlations of dynamic connectivity from the original analysis (Figure

6A), the consistency of dynamic connectivity from the residual time series were largely diminished.

Figure 6D and 6E illustrate the changes of dynamic connectivity time courses after the regression

between a representative ROI pair, i.e. right supramarginal gyrus and posterior cingulate cortex (see

supplementary Figure S4 for other ROI pairs). The intersubject correlation reduced from 0.64 to 0.29.

Figure 6C illustrates the dynamic connectivity of the consistent components of regional activity between

these two ROIs. The fluctuating pattern was similar to those calculated from individual subject's original

time series (Figure 6D), further confirmed that the consistent dynamic connectivity across individuals was

driven by the consistent component of regional activity.

Figure 6 A) and B) Intersubject correlations (ISC) of dynamic connectivity calculated from raw time

series (A) and residual time series after regressing out the intersubject consistent components (B). C)

Dynamic connectivity of the consistent component of regional activity between right supramarginal gyrus

(RSMG) and posterior cingulate cortex (PCC). D) and E) Time courses of dynamic connectivity (Fisher's

z) between RSMG and PCC calculated from raw time series (D) and the residual time series after

regressing out the intersubject consistent component. F) The time courses of intersubject correlations of

regional activity in RSMG and PCC.

Lastly, we calculated intersubject correlations of regional activity in every sliding window

(supplementary Figure S5). Figure 6F shows the time courses of intersubject correlations of regional

activity in the right supramaginal gyrus and posterior cingulate cortex ROIs. Both regions showed

similarly but reversed time courses as the dynamic connectivity between them. That is, during the two

periods of dips of dynamic connectivity, there were elevated intersubject correlations of regional activity

in both regions. But this kind of close relations cannot be observed in the other two pairs of ROIs

(supplementary Figure S4).

3.2.4. Effects of sliding-window length

We repeated the ROI-based intersubject correlation analysis of dynamic connectivity using different
window length from 10 TRs to 60 TRs. The intersubject correlation matrices were in general weaker
when the window was shorter, especially for the 10-TR window (Figure 7). As the window went longer,
the correlations matrices became similar to the 30-TR window results. But for even longer window, there
were two different trends. First, some of the intersubject correlations kept increasing, usually between
regions that involved in one or two visual ROIs (Figure 7B). On the other hand, some of the intersubject
correlations decreased after peaked at the 30-TR window, usually between regions that involved
supramarginal gyrus or posterior cingulate cortex. Figure 7C illustrated the time courses of dynamic
connectivity between right supramarginal gyrus and posterior cingulate cortex. It can be seen that the
variability of dynamic connectivity time courses were larger in short window. When using 10-TR
window, the dynamic connectivity changed fast, and were not aligned across subjects. When using 30-
TR window, the dynamic connectivity time courses became smoother, and the fluctuations were more
aligned across subjects, which in turn gave rise to higher intersubject correlations. But when using 50-TR
window, the time courses of dynamic connectivity become too smooth, so that the subject averaged trend
become less apparent. It's noteworthy that for the dynamic connectivity between the left and right medial
visual ROIs and between left medial visual and left precentral gyrus ROIs, there were linear trends of
dynamic connectivity across subjects, which gave rise to high intersubject correlations in longer windows
(Figure S6).

Figure 7 A) The effects of sliding-window length on the intersubject correlations of dynamic connectivity.

B) Intersubject correlations of dynamic connectivity of three pairs of regions of interest: left medial visual

(LMV) and right medial visual (RMV), LMV and left precentral gyrus (LPCG), and right supramarginal

gyrus (RSMG) and posterior cingulate cortex (PCC). C) The time courses of dynamic connectivity

between RSMG and PCC in three window lengths. TR, repetition time.

4. Discussion

In the current study, we proposed intersubject correlation analysis on the time courses of dynamic

connectivity during natural vision. We were able to identify intersubject consistent dynamic connectivity

at similar levels as the intersubject correlations of regional activity, although the time courses of dynamic

connectivity were thought to be nosier than the original time series. By using seed regions from the

visual cortex and supramarginal gyrus, we demonstrated widespread brain regions that showed high

intersubject consistent dynamic connectivity with these seeds, although these regions themselves did not

show intersubject correlations of regional activity. These regions included high order association regions

such as frontal and parietal regions, as well as the default mode network. The intersubject consistent

patterns of dynamic connectivity support the functional meaningfulness of dynamic connectivity during

movie watching, and suggest that dynamic connectivity could be a complementary avenue to characterize

the functions of a brain region.

The brain regions that had the highest intersubject correlations of regional activity are mainly in

the posterior visual related regions, which are consistent with previous studies (Hasson et al., 2004;

Nummenmaa et al., 2012). In addition, the current study found dynamic connectivity among different

levels of visual areas that were highly consistent across subjects. This is interesting because although the

overall functional connectivity among the visual areas are very high (Figure 4A), there are still

functionally meaningful fluctuations of interactions among the visual regions. The observable dynamics

of connectivity among visual areas are in line with previous studies showing task modulated connectivity

among visual areas in different task conditions (Di et al., 2019, 2015; Di and Biswal, 2017). It is

interesting to note that the dynamics of intersubject correlations of regional activity in the visual areas

also showed decreased trends at the beginning of the session (Figure S4 and S5). Therefore, the

decreased connectivity in the beginning may reflect adaptations effects in the visual areas. However,

during the latter half of the session, the dynamics of intersubject correlations of regional activity kept at a

stable level, which cannot explain the dynamics of connectivity between them (Figure S4).

The bilateral supramarginal gyrus regions are major regions outside the visual cortex that showed

high intersubject correlations of regional activity. The involvements of supramarginal gyrus of

intersubject correlations are inconsistent in the literature (Hasson et al., 2004; Kauppi et al., 2010), which

probably due to different movies the participants watched. Given their critical role in empathy (Silani et

al., 2013), it is reasonable to observe high intersubject correlations in the supramarginal gyrus during the

watching of the animated movie, which involves the understanding the intentions of different animated

characters. Interestingly, the intersubject correlations of regional activity in the supramarginal gyrus also

showed dynamics, with two periods of high correlations roughly between the 20^{th} and 30^{th} windows and

between the 50^{th} and 60^{th} windows (Figure 6F and S4B). The first may correspond to the scene when

Peck the stork and Gus the cloud first met, where their interactions appeared to be different from the other

storks and clouds. The second may coincide with the scene when Peck flew away, and Gus thought Peck

had abandoned him. These scenes require active inferences of the intentions of the characters, and may

involve mismatches between predictions and the actual story development. Therefore, it is reasonable to

see high cross-subject similarities in the supramarginal gyrus during these two periods.

In addition to regional activity, we also found that the default mode network showed highly

consistent dynamic connectivity with the supramarginal gyrus regions. Similar to the supramarginal

gyrus ROIs, the regional activity in the posterior cingulate cortex showed two periods of high consistent

regional activity (Figure 6F). But interestingly, during these two periods the posterior cingulate cortex

and supramarginal gyrus showed strong anti-correlation (Figure 6C). The default mode network involves

high-order representation of the world (Carhart-Harris and Friston, 2010). And the functional

communications between the default mode network and supramarginal gyrus may reflect the prediction

error between inner representation and the input from the video. Similar to a previous study using

dynamic intersubject connectivity analysis (Simony et al., 2016), both of the studies highlighted the

critical role of the default mode network in understanding of the narratives of a movie.

Generally speaking, the intersubject consistent connectivity and stable functional connectivity

showed disassociations. Specifically, the ROI pairs that showed highly consistent dynamic connectivity

may have high stable functional connectivity or very low overall connectivity. The latter case may be

more interesting, because it suggests transient functional communications that cannot be observed in

traditional functional connectivity analysis. The 9 ROIs used in the current analysis are from three

functional modules, i.e. unimodal visual, higher order task positive, and default mode networks. The

three functional modules can be confirmed in the matrix of stable connectivity (Figure 4A), where there

are high within-module functional connectivity but weak between-module connectivity. The matrix of

consistent dynamic connectivity, on the other hand, showed that there are more between-module dynamic

connectivity. These results are in line with the economy account of brain network organizations, which

suggests that the functional communications between modulates are costly in terms of energy

consumption, therefore are more transient (Bullmore and Sporns, 2012). It is also consistent with the

findings that the connectivity between modules are more variable and context dependent (Di and Biswal,

2019; Fu et al., 2017).

By calculating dynamic connectivity time courses from individual's time series, the proposed

method focused on the consistency of the dynamic connectivity time courses across subjects. Our

additional analysis showed that the consistent dynamic connectivity time courses was driven by the

dynamic connectivity of the consistent component of the regional time series, at least for the current video

watched. The latter method provides a simple approach to reveal the dynamics of connectivity, and is

closely related to the dynamic intersubject functional connectivity approach proposed by Simony et al.

(Simony et al., 2016). Our method, on the other hand, can not only reveal the time course of dynamic

connectivity, but can also provide a quantity of a connection about how the dynamic connectivity is

consistent across subjects. Eventually, we will be able to obtain a matrix of the consistency of dynamic

connectivity among ROIs from the whole brain. This is important because the seed-based approach used

in the current analysis may miss dynamic connectivity between regions that do not have consistent

regional activity. The connectome-based approach can provide a comprehensive mapping of dynamic

communications across the brain during the watching of a movie, and can be seen a special form of task

connectome (Di and Biswal, 2019).

The selection of window length for dynamic connectivity analysis is nontrivial (Fu et al., 2014;

Zhang et al., 2013). The shorter the window length, the finer the temporal resolution for dynamic

connectivity could be. However, less time points for each window would also mean noisier estimates of

connectivity. In the current analysis, 10-TR window gave very noisy estimate of functional connectivity,

thus making intersubject correlations very low. On the other hand, longer window will make connectivity

estimate accurate, but at a cost of losing temporal resolution. In the context of the current video watched,

30-TR window seems a balance. However, this time scale of dynamic connectivity fluctuations may not

be easily generalized to other videos or to resting-state. But it certainly can provide some insight to the

chosen of window length in future studies. In addition, some computational method may be used to avoid

the window length issue, e.g. using adaptive covariance estimates (Fu et al., 2014; Zhang et al., 2013) or

window-free method such as Kalman filtering (Kang et al., 2011) or instantaneous phase synchronization

(Glerean et al., 2012).

462 5. Conclusion

In the current study, we proposed intersubject correlation analysis on dynamic connectivity. The results

revealed widespread brain regions that showed consistent intersubject correlations of dynamic

connectivity. The consistent correlations support the functional significance of dynamic connectivity

during natural vision. The method may provide a complementary approach to understand the dynamic

nature of brain functional integrations.

Acknowledgement:

This study was supported by grants from (US) National Institute of Health (R01 AT009829; R01

DA038895).

474 Conflict of interest

The authors declared that there is no conflict of interest.

References:

Allen, E.A., Damaraju, E., Plis, S.M., Erhardt, E.B., Eichele, T., Calhoun, V.D., 2014. Tracking whole-
brain connectivity dynamics in the resting state. Cereb. Cortex N. Y. N 1991 24, 663–76.
https://doi.org/10.1093/cercor/bhs352
Bullmore, E., Sporns, O., 2012. The economy of brain network organization. Nat. Rev. Neurosci. 13,
336–349. https://doi.org/10.1038/nrn3214
Carhart-Harris, R.L., Friston, K.J., 2010. The default-mode, ego-functions and free-energy: a
neurobiological account of Freudian ideas. Brain J. Neurol. 133, 1265–83.
https://doi.org/10.1093/brain/awq010
Chen, G., Shin, Y.-W., Taylor, P.A., Glen, D.R., Reynolds, R.C., Israel, R.B., Cox, R.W., 2016.
Untangling the relatedness among correlations, part I: Nonparametric approaches to inter-subject

correlation analysis at the group level. NeuroImage 142, 248–259.
https://doi.org/10.1016/j.neuroimage.2016.05.023

Di, X., Biswal, B.B., 2019. Toward Task Connectomics: Examining Whole-Brain Task Modulated
Connectivity in Different Task Domains. Cereb. Cortex 29, 1572–1583.
https://doi.org/10.1093/cercor/bhy055

Di, X., Biswal, B.B., 2017. Psychophysiological Interactions in a Visual Checkerboard Task:
Reproducibility, Reliability, and the Effects of Deconvolution. Front Neurosci 1–36.
https://doi.org/10.3389/fnins.2017.00573

Di, X., Biswal, B.B., 2015. Characterizations of resting-state modulatory interactions in the human brain.
J. Neurophysiol. 114, 2785–96. https://doi.org/10.1152/jn.00893.2014

Di, X., Fu, Z., Chan, S.C., Hung, Y.S., Biswal, B.B., Zhang, Z., 2015. Task-related functional
connectivity dynamics in a block-designed visual experiment. Front. Hum. Neurosci. 9, 1–11.
https://doi.org/10.3389/fnhum.2015.00543

Di, X., Zhang, Z., Biswal, B.B., 2019. Understanding psychophysiological interaction and its relations to
beta series correlation. bioRxiv 322073. https://doi.org/10.1101/322073

Fox, M.D., Snyder, A.Z., Vincent, J.L., Corbetta, M., Van Essen, D.C., Raichle, M.E., 2005. The human
brain is intrinsically organized into dynamic, anticorrelated functional networks. Proc. Natl. Acad.
Sci. U. S. A. 102, 9673–8.

Friston, K.J., 2011. Functional and effective connectivity: a review. Brain Connect. 1, 13–36.
https://doi.org/10.1089/brain.2011.0008

Friston, K.J., Williams, S., Howard, R., Frackowiak, R.S., Turner, R., 1996. Movement-related effects in
fMRI time-series. Magn. Reson. Med. Off. J. Soc. Magn. Reson. Med. Soc. Magn. Reson. Med.
35, 346–55. https://doi.org/DOI 10.1002/mrm.1910350312

Fu, Z., Chan, S.-C., Di, X., Biswal, B., Zhang, Z., 2014. Adaptive covariance estimation of non-stationary
processes and its application to infer dynamic connectivity from fMRI. IEEE Trans. Biomed.
Circuits Syst. 8, 228–39. https://doi.org/10.1109/TBCAS.2014.2306732

Fu, Z., Tu, Y., Di, X., Biswal, B.B., Calhoun, V.D., Zhang, Z., 2017. Associations between Functional
Connectivity Dynamics and BOLD Dynamics Are Heterogeneous Across Brain Networks. Front.
Hum. Neurosci. 11. https://doi.org/10.3389/fnhum.2017.00593

Fu, Z., Tu, Y., Di, X., Du, Y., Pearlson, G.D., Turner, J.A., Biswal, B.B., Zhang, Z., Calhoun, V.D., 2018.
Characterizing dynamic amplitude of low-frequency fluctuation and its relationship with dynamic
functional connectivity: An application to schizophrenia. NeuroImage 180, 619–631.
https://doi.org/10.1016/j.neuroimage.2017.09.035

Fu, Z., Tu, Y., Di, X., Du, Y., Sui, J., Biswal, B.B., Zhang, Z., de Lacy, N., Calhoun, V.D., 2019.
Transient increased thalamic-sensory connectivity and decreased whole-brain dynamism in
autism. NeuroImage 190, 191–204. https://doi.org/10.1016/j.neuroimage.2018.06.003

Glerean, E., Salmi, J., Lahnakoski, J.M., Jääskeläinen, I.P., Sams, M., 2012. Functional Magnetic
Resonance Imaging Phase Synchronization as a Measure of Dynamic Functional Connectivity.
Brain Connect. 2, 91–101. https://doi.org/10.1089/brain.2011.0068

Hasson, U., Nir, Y., Levy, I., Fuhrmann, G., Malach, R., 2004. Intersubject synchronization of cortical
activity during natural vision. Science 303, 1634–40. https://doi.org/10.1126/science.1089506

Hutchison, R.M., Womelsdorf, T., Allen, E. a., Bandettini, P. a., Calhoun, V.D., Corbetta, M., Della
Penna, S., Duyn, J.H., Glover, G.H., Gonzalez-Castillo, J., Handwerker, D. a., Keilholz, S.,
Kiviniemi, V., Leopold, D. a., de Pasquale, F., Sporns, O., Walter, M., Chang, C., 2013. Dynamic
functional connectivity: Promise, issues, and interpretations. NeuroImage 80, 360–378.
https://doi.org/10.1016/j.neuroimage.2013.05.079

Kang, J., Wang, L., Yan, C., Wang, J., Liang, X., He, Y., 2011. Characterizing dynamic functional
connectivity in the resting brain using variable parameter regression and Kalman filtering
approaches. NeuroImage 56, 1222–34. https://doi.org/10.1016/j.neuroimage.2011.03.033

Kauppi, J.-P., Jääskeläinen, I.P., Sams, M., Tohka, J., 2010. Inter-subject correlation of brain
hemodynamic responses during watching a movie: localization in space and frequency. Front.
Neuroinformatics 4. https://doi.org/10.3389/fninf.2010.00005
Leonardi, N., Van De Ville, D., 2015. On spurious and real fluctuations of dynamic functional
connectivity during rest. NeuroImage 104, 430–436.
https://doi.org/10.1016/j.neuroimage.2014.09.007
Lindquist, M.A., Xu, Y., Nebel, M.B., Caffo, B.S., 2014. Evaluating dynamic bivariate correlations in
resting-state fMRI: A comparison study and a new approach. NeuroImage 101, 531–546.
https://doi.org/10.1016/j.neuroimage.2014.06.052
Nastase, S.A., Gazzola, V., Hasson, U., Keysers, C., 2019. Measuring shared responses across subjects
using intersubject correlation. Soc. Cogn. Affect. Neurosci. 14, 667–685.
https://doi.org/10.1093/scan/nsz037
Nummenmaa, L., Glerean, E., Viinikainen, M., Jääskeläinen, I.P., Hari, R., Sams, M., 2012. Emotions
promote social interaction by synchronizing brain activity across individuals. Proc. Natl. Acad.
Sci. U. S. A. 109, 9599–9604. https://doi.org/10.1073/pnas.1206095109
Park, H.-J., Friston, K., 2013. Structural and Functional Brain Networks: From Connections to Cognition.
Science 342, 1238411–1238411. https://doi.org/10.1126/science.1238411
Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but systematic
correlations in functional connectivity MRI networks arise from subject motion. NeuroImage 59,
2142–2154. https://doi.org/10.1016/j.neuroimage.2011.10.018
Raichle, M.E., MacLeod, A.M., Snyder, A.Z., Powers, W.J., Gusnard, D.A., Shulman, G.L., 2001. A
default mode of brain function. Proc. Natl. Acad. Sci. U. S. A. 98, 676–82.
https://doi.org/10.1073/pnas.98.2.676
Richardson, H., Lisandrelli, G., Riobueno-Naylor, A., Saxe, R., 2018. Development of the social brain
from age three to twelve years. Nat. Commun. 9, 1–12. https://doi.org/10.1038/s41467-018-
03399-2
Rosenthal, G., Sporns, O., Avidan, G., 2017. Stimulus Dependent Dynamic Reorganization of the Human
Face Processing Network. Cereb. Cortex 27, 4823–4834. https://doi.org/10.1093/cercor/bhw279
Silani, G., Lamm, C., Ruff, C.C., Singer, T., 2013. Right Supramarginal Gyrus Is Crucial to Overcome
Emotional Egocentricity Bias in Social Judgments. J. Neurosci. 33, 15466–15476.
https://doi.org/10.1523/JNEUROSCI.1488-13.2013
Simony, E., Honey, C.J., Chen, J., Lositsky, O., Yeshurun, Y., Wiesel, A., Hasson, U., 2016. Dynamic
reconfiguration of the default mode network during narrative comprehension. Nat. Commun. 7,
12141. https://doi.org/10.1038/ncomms12141
Teichert, T., Grinband, J., Hirsch, J., Ferrera, V.P., 2010. Effects of heartbeat and respiration on macaque
fMRI: Implications for functional connectivity. Neuropsychologia 48, 1886–1894.
https://doi.org/10.1016/j.neuropsychologia.2009.11.026
Zhang, Z.G., Fu, Z.N., Chan, S.C., Hung, Y.S., Motta, G., Di, X., Biswal, B.B., 2013. Adaptive window
selection in estimating dynamic functional connectivity of resting-state fMRI, in: 2013 9th
International Conference on Information, Communications & Signal Processing. IEEE, pp. 1–4.
https://doi.org/10.1109/ICICS.2013.6782935
579
580

581 **Table 1** Clusters with differential intersubject correlations of dynamic connectivity among the medial

582 visual, lateral visual, and supramarginal gyrus seeds. All clusters were thresholded at p < 0.001, and

583 cluster thresholded at p < 0.0167 (0.05 / 3) after family-wise error (FWE) correction using nonparametric

584 method.

		MNI Coordinates				
cluster FWE	Voxel	x	y	z	Peak t	Label
Medial visual > (lateral visual + supramarginal)						
< 0.001	514	-12	-100	-4	7.85	Occipital pole
< 0.001	372	24	-88	-4	6.80	Right inferior occipital gyrus
0.001	106	-51	11	23	5.87	Left precentral gyrus
Lateral visual > (Medial visual + supramarginal)						
0.016	78	48	-70	8	7.03	Right inferior occipital gyrus
0.013	86	15	-79	-4	5.64	Lingual gyrus
Supramarginal > (medial visual + lateral visual)						
< 0.001	1168	3	-40	44	9.12	Precuneus
0.002	237	-12	50	14	7.41	Medial superior frontal gyrus
0.006	122	-30	-4	44	6.90	Left middle frontal gyrus
0.003	175	18	50	44	6.44	Right middle frontal gyrus

586 MNI, Montreal Neurological Institute