Intersubject consistent dynamic connectivity during natural vision revealed by functional MRI

Xin Di [1,2], Bharat B Biswal [1,2] *

1. Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, NJ, 07029, USA

2. School of Life Sciences and Technology, University of Electronic Science and Technology of China,

Chengdu, China

* Corresponding author:

Bharat B. Biswal, PhD

607 Fenster Hall, University Height

Newark, NJ, 07102, USA

bbiswal@yahoo.com

Abstract

The functional communications between brain regions are thought to be dynamic. However, it is usually

difficult to elucidate whether the observed dynamic connectivity is functionally meaningful or simply due

to noise during unconstrained task conditions such as resting-state. During naturalistic conditions, such

as watching a movie, it has been shown that brain activities in the same region, e.g. visual cortex, are

consistent across subjects. Following similar logic, we proposed to study intersubject correlations of the

time courses of dynamic connectivity during naturalistic conditions to extract functionally meaningful

dynamic connectivity patterns. We analyzed a functional MRI (fMRI) dataset when the subjects watched

a short animated movie. We calculated dynamic connectivity by using sliding window technique, and

further quantified the intersubject correlations of the time courses of dynamic connectivity. Although the

time courses of dynamic connectivity are thought to be noisier than the original signals, we found similar

level of intersubject correlations of dynamic connectivity. Most importantly, highly consistent dynamic

connectivity could occur between regions that did not show intersubject correlations of regional activity,

and between regions with little stable functional connectivity. The analysis highlighted higher order brain

regions such as the lateral prefrontal cortex and the default mode network that dynamically interact with

posterior visual regions during the movie watching, which may be associated with the understanding of

the movie.

Keywords: default mode network; dynamic connectivity; intersubject correlation; naturalistic condition;

supramarginal gyrus

Highlights

- Intersubject shared time courses may provide a complementary approach to study dynamic connectivity

- Widespread regions showed highly shared dynamic connectivity during movie watching, while these regions themselves did not show shared regional activity

- Shared dynamic connectivity often occurred between regions from different functional systems

1. Introduction

The functional communications between spatially remote brain regions, especially the dynamics of connectivity, is a key to understand brain functions (Bullmore and Sporns, 2012; Friston, 2011; Park and Friston, 2013). Recently, the study of dynamic connectivity has drawn increasing research interest, especially in resting-state (Allen et al., 2014; Fu et al., 2019, 2018; Hutchison et al., 2013). However, due to the unconstrained nature of resting-state, it is difficult to elucidate whether the observed changes of connectivity across sliding windows are due to real fluctuations of functional communications, or simply due to random fluctuations (Lindquist et al., 2014). Moreover, the blood-oxygen-level dependent (BOLD) signals measured by fMRI are sensitive to physiological noises, such as respiration, heartbeat (Teichert et al., 2010), and head motion (Power et al., 2012), which may give rise to spurious correlation estimates for short window.

One way to capture meaningful dynamic functional connectivity is to manipulate subjects' mental states during the course of scan, so that there is known reference for the changes of connectivity. For example, in a typical task-based fMRI study with blocked design, different task conditions are assigned as blocks. Therefore, the time courses of dynamic connectivity can be correlated with the task design to identify task related connectivity changes (Di et al., 2015; Rosenthal et al., 2017). An alternative approach is to expose the subjects with naturalistic stimuli, such as a short movie. Although there is no predefined references of dynamic connectivity changes, one may take advantage of the phenomenon of intersubject correlation to capture changes that are consistent across different subjects (Hasson et al., 2004; Nastase et al., 2019).

In the seminal study, Hasson and colleagues calculated intersubject correlations of the time series of BOLD signal (Figure 1A) when the subjects were watching a movie (Hasson et al., 2004). They demonstrated that several brain regions, especially the visual cortex, are highly correlated across subjects during movie watching. We propose that similar approach can be applied to the time courses of dynamic connectivity to capture meaningful functional communication dynamics during natural vision. Specifically, dynamic connectivity is usually calculated using a sliding window approach, so that a time

series of dynamic connectivity can be obtained. The time courses of dynamic connectivity can then be

correlated across-subjects (Figure 1B). If the dynamic connectivity reflects real time functional

communications between regions that are caused by the viewing of natural stimuli, then the time courses

of dynamic connectivity from different subjects should somehow correlated. Therefore, we can apply

intersubject correlation method to identify meaningful dynamic communications between regions.

Figure 1 Illustrations of the calculations of intersubject correlations of the time series of regional activity

76   (A) and the time courses of dynamic connectivity between two regions (B).

In the current study, we analyzed an fMRI dataset where the subjects were scanned when viewing

a short animated movie. The aim was to identify dynamic connectivity that were shared cross subjects

during the movie watching. In order to do so, we first performed regular intersubject correlation analysis

to identify brain regions that showed consistent regional activity. Given these regions, we adopted a seed-

based strategy to calculate dynamic connectivity between a seed region and every voxels in the brain. We

then evaluated and identified regions whose connectivity with the seed were consistent cross subjects.

Even though higher order association regions did not typically show high intersubject correlations of

regional activity (Hasson et al., 2004), their functional communications with lower order regions may be

consistent across subject following the narrative of the movie. We therefore hypothesized that

intersubject correlations of dynamic connectivity may be able to identify more widespread regions and

functional dynamics that are associated with the watching of the movie.

2. Materials and methods

2.1. Data and task

The fMRI data were obtained through openneuro (https://openneuro.org/; accession #: ds000228). Only

the data from adult subjects were analyzed. There were originally 33 adult subjects. Two subjects' data

were discarded because of poor brain coverage (subject #: sub-pixar123 and sub-pixar123), and two were

discarded due to large head motions (sub-pixar149 and sub-pixar150). As a result, a total of 29 subjects

were included for the current analysis (17 females). The mean age is 24.6 years old (18 to 39 years).

97   During the fMRI scan, the subjects watched a silent version of Pixar animated movie "Partly

Cloudy", which is 5.6 minutes long (https://www.pixar.com/partly-cloudy#partly-cloudy-1). Brain MRI

images were acquired on a 3-Tesla Siemens Tim Trio scanner using the standard Siemens 32-channel

head coil. Functional images were collected with a gradient-echo EPI sequence sensitive to BOLD

contrast in 32 interleaved near-axial slices (EPI factor: 64; TR: 2 s, TE: 30 ms, flip angle: 90°). The

voxel size were 3.13 mm isotropic, with 3 subjects with no slice gap and 26 subjects with 10% slice gap.

168 functional images were acquired for each subject, with four dummy scans collected before the real

scans to allow for steady-state magnetization. T1-weighted structural images were collected in 176

interleaved sagittal slices with 1 mm isotropic voxels (GRAPPA parallel imaging, acceleration factor of

3; FOV: 256 mm). For more information for the dataset please refers to (Richardson et al., 2018).

2.2. FMRI data analysis

2.2.1. Preprocessing

FMRI data processing and analyses were performed using SPM12 and MATLAB (R2017b) scripts. A

subject's T1 weighted structural image was first segmented into gray matter, white matter, cerebrospinal

fluid, and other tissue types, and was normalized into standard Montreal Neurological Institute (MNI)

space. The T1 images were then skull stripped based on the segmentation results. Next, all the functional

images of a subject were realigned to the first image of the session and coregistered to the skull stripped

T1 image of the same subject. Framewise displacement was calculated for the translation and rotation

directions for each subject (Di and Biswal, 2015). Subjects who had maximum framewise displacement

greater than 1.5 mm or 1.5^{o} were discarded from further analysis. The functional images were then

normalized to MNI space using the parameters obtained from the segmentation step with resampled voxel

size of 3 x 3 x 3 mm^{3}. The functional images were then spatially smoothed using a Gaussian kernel of 8

mm. Lastly, a voxel-wise general linear model (GLM) was built for each subject to model head motion

effects (Friston's 24-parameter model) (Friston et al., 1996), low frequency drift (1/128 Hz), and constant

offset. The residuals of the GLM were saved as a 4-D image series, which were used for further

intersubject correlation analysis.

2.2.2. Intersubject correlation analysis

The correlations of time series of either brain activity or dynamic connectivity are calculated between

pairs of subjects. If there are N subjects, then there will be N x (N-1) / 2 correlation coefficients. The

statistics of these correlations become tricky, because they are calculated from only N subjects, therefore

not independent. An alternative approach is leave-one-out (Nastase et al., 2019), where the time series of

one hold-out subject were correlated with the averaged time series of the remaining N – 1 subjects. The

averaged time series of N – 1 subjects were thought to reflect the consistent component rather than the

noisy individual's time series. Therefore, the resulting correlations should be higher than the pair-wise

correlations. Another benefit is that this approach estimates one correlation for each subject, making

group level statistics easier. Therefore, we adopt the leave-one-out approach in the current analysis.

We first performed intersubject correlation analysis on regional activity time series. The

preprocessed BOLD time series were extracted for each voxel and subject in a gray matter mask. For a

given voxel, the time series of one subject was hold out, and the averaged time series of the remaining

subject were calculated. Then the time series of the hold-out subject were correlated with the averaged

time series. This process was performed for every subject and every voxel, resulting in one correlation

map for one subject. The correlation maps were transformed into Fisher's z maps. Group level one

sample t test was then performed to identify regions whose intersubject correlations were consistently

greater than 0. However, the null hypothesis statistical significance testing may not provide much

information of the effect size. There may be only small but consistent correlations for each subject,

which could give rise to very high statistical significance in a one sample t test. Indeed, when doing such

null hypothesis statistical significance testing for intersubject correlation analysis, usually almost all the

brain regions will show somehow significant correlations (Chen et al., 2016). We are more interested and

focused on the real effect size, i.e. correlation coefficients, in our analysis. We therefore averaged the

Fisher's z maps, and transformed them back into r maps. The continuous r maps were shown in the

results section.

We next performed intersubject correlation analysis on dynamic connectivity using a seed-based

approach. Given that a set of brain regions showed high intersubject correlations of regional activity, we

defined these regions as seeds. We adopted a relatively high threshold of $r > 0.45$ for the averaged

intersubject correlation map of regional activity to isolate four visual related seeds. Two of the seeds

were located in the medial and posterior portion of the occipital lobe, which mainly covered the lingual

gyrus and calcarine sulcus. The other two seeds were located bilaterally in the middle occipital gyrus and

extended to the middle temporal gyrus. We labeled them as left and right medial visual and lateral visual

seeds, respectively. In addition, we adopted a relatively low threshold of $r > 0.35$ to isolate the left and

right supramarginal gyrus seeds. For each seed, we performed a voxel-wise correlation analysis, i.e.

calculating intersubject correlations of dynamic connectivity between the seed and every voxel in the gray

matter mask. For two given time series from a seed and a voxel, we used sliding window technique to

calculate dynamic connectivity. The window length was set as 30 time points (60 s) (Nastase et al., 2019),

and the time step was set as 2 time point (4 s). Therefore, the time course of dynamic connectivity had 70

window steps. Similarly, we calculated correlations between the time courses of dynamic connectivity of

a given subject with the averaged dynamic connectivity of remaining subjects for a given voxel. As a

result, there was one correlation map for each seed and subject.

The r maps of correlations of dynamic connectivity were transformed into Fisher's z maps for

group level statistical analysis. Again, we also simply calculated averaged z map for a seed, and

transformed it back into r map. In addition, we performed a voxel-wise repeated measure one way

analysis of variance (ANOVA) to identify regions that showed specific dynamic connectivity patterns

with different seeds.

In addition to the voxel-based analysis, we also performed region of interest (ROI)-based analysis

for in-depth examinations of the dynamic connectivity effects. In addition to the six seeds, we included

four more regions that showed high intersubject correlations of dynamic connectivity with the seeds.

They were left precentral gyrus and left inferior frontal gyrus, which showed high intersubject

correlations of dynamic connectivity with the medial visual seeds, and posterior cingulate cortex and

medial prefrontal cortex, which showed high intersubject correlations of dynamic connectivity with the

supramarginal gyrus seeds. The calculations of intersubject correlations of dynamic connectivity were the

same as the seed-based analysis.

The selections of sliding window length is nontrivial (Fu et al., 2014; Zhang et al., 2013). In

addition to the 30-TR window length, we also explored other window length of 10 TRs (20 s), 20 TRs (40

s), and 40 TRs (80 s). For each window length, we calculated intersubject correlations of dynamic

connectivity among the 10 ROIs.

2.2.3. Stable functional connectivity

We also calculated stable functional connectivity among the 10 ROIs to compare them with the dynamic

connectivity. First, for each subject, we calculated correlation coefficients across the 10 ROIs, and

transformed them into Fisher's z. Then the z matrices were averaged across the 29 subjects, and

transformed back into r values. Second, we calculated the consistent component of each ROI, i.e.

averaging the time series across the 29 subjects. And then one single correlation matrix among the 10

ROIs was calculated. This connectivity of the consistent component is essentially the same as

intersubject functional connectivity proposed by Simony and colleagues (Simony et al., 2016).

3. Results

3.1. Intersubject correlations of regional activity

We first calculated intersubject correlations of regional activity for every voxel in the brain during the

video watching (Figure 2A). The highest correlations were around 0.5. The major regions that had high

intersubject correlations were the visual cortex extending anterior to the fusiform gyrus and middle

temporal lobe. The bilateral supramarginal gyrus also showed high intersubject correlations. The

bilateral precentral gyrus also showed intersubject correlations, but the effect sizes were much smaller.

Figure 2A shows all the voxels with positive correlation values. It is noteworthy that many regions

showed very small intersubject correlations, including largely the prefrontal cortex and anterior temporal

lobe.

Figure 2 Intersubject correlation maps of regional activity (A) and dynamic connectivity with different

seeds (B through G). The seed regions were depicted in blue in respective rows. All voxels with positive

correlations are shown. The numbers at the bottom represent z and x coordinates in Montreal

Neurological Institute (MNI) space.

3.2. Intersubject correlations of dynamic connectivity

3.2.1 Seed-based analysis

We defined seed regions where there were high intersubject correlations of regional activity, which

included bilateral medical visual regions, lateral visual regions, and supramarginal gyrus. We next

calculated voxel-wise intersubject correlations of dynamic connectivity with the six seeds, respectively

(Figure 2B through 2G). There were widespread brain regions that showed intersubject consistent

dynamic correlations with different seeds. First of all, the effect sizes of the intersubject correlations of

dynamic connectivity, i.e. the correlation coefficients, were comparable to those in the intersubject

correlations of regional activity. Secondly, regions with intersubject correlations of dynamic connectivity

turned out to be more widespread and extended to the frontal and temporal regions that did not show high

intersubject correlations of regional activity. Thirdly, the left and right corresponding seeds showed

similar dynamic connectivity patterns, but there are substantial different patterns of dynamic connectivity

among medial visual, lateral visual, and supramarginal gyrus seeds. In order to highlight specific brain

regions that showed dynamic connectivity with different seeds, we performed repeated measure ANOVA

and compared the maps of different level of seeds with other seeds (Figure 3 and Table 1). The medical

visual seeds showed consistent dynamic connectivity with mainly lateral brain regions, including the left

inferior frontal gyrus/precentral gyrus, bilateral supramarginal gyrus, and left orbital gyrus/inferior frontal

gyrus. The lateral visual seeds showed consistent dynamic connectivity with several visual regions. In

contrast, the supramarginal seeds showed consistent dynamic connectivity with the precuneus/posterior

cingulate gyrus, medial prefrontal cortex, and bilateral angular gyrus, which basically formed the default

mode network.

Figure 3 Differential intersubject correlations of dynamic connectivity among the medial visual, lateral

visual, and supramarginal gyrus seeds. Red, medial visual seeds greater than lateral visual and

supramarginal gyrus seeds. Blue, lateral visual seeds greater than medial visual and supramarginal gyrus

seeds. Green, supramarginal gyrus seeds greater than medial and lateral visual seeds. All maps were

thresholded at $p < 0.001$, and cluster thresholded at $p < 0.0167$ (0.05 / 3) after false discovery rate (FDR)

233   correction.

3.2.2. Relationships with stable functional connectivity

In order to better understand and interpret the brain regions and connectivity relationships, we further

calculated different types of connectivity measures among a set of regions of interest. In addition to the

six seeds, we defined left precentral gyrus and left inferior frontal gyrus ROIs that showed higher

dynamic connectivity with the medial visual seeds, and posterior cingulate cortex and medial prefrontal

cortex ROIs that showed high dynamic connectivity with the supramarginal seeds. Among the 10 regions,

we calculated regular mean functional connectivity (Figure 4A) and connectivity derived from the

consistent components (Figure 4B). These two correlations matrices look similar, and clearly showed

three clusters of brain regions. The first four regions were all visual. The fifth to eight regions were the

bilateral supramarginal gyrus, and lateralized frontal regions, which were all high order association brain

regions. The last two regions were part of the default mode network, which also showed negative

correlations with the association regions in the consistent component correlations.

Figure 4 Correlation matrices among the 10 regions of interest (ROI) using different methods. A) Mean

 functional connectivity across the 29 subjects. B) Correlations of the consistent component of each ROI

  (averaged time series across the 29 subjects). C) Intersubject correlations of dynamic connectivity.

The intersubject consistent dynamic connectivity matrix (Figure 4C) was largely different from

the two stable correlation matrices. Some high consistent dynamic connectivity was observed within the

visual regions. The highest correlation was observed between the left and right medial visual regions ($r =$

0.70). In contrast, many consistent dynamic connectivity were shown between different functional

networks, where there were virtually none or even negative stable correlations. Specifically, the medial

visual regions showed high consistent dynamic connectivity with the left frontal regions. The highest

intersubject correlation was 0.56 between left medial visual region and left precentral gyrus. The default

mode regions and supramarginal regions also showed high consistent dynamic connectivity. The highest

correlation was 0.60 between the posterior cingulate cortex and right supramarginal gyrus. It is

noteworthy that these regions generally showed negative stable correlations in Figure 4B.

3.2.3. The time courses of dynamic connectivity

Lastly, we analyzed the time courses of dynamic connectivity for the above mentioned pairs of regions

(Figure 5). The consistent dynamic connectivity between left and right medial visual regions was in

general high, which is consistent with the results of stable connectivity. But it can be seen that the

connectivity level went down during the first half of windows, and continued with two cycles of up and

down fluctuations. The fluctuations rather than a monotonic linear trend suggest that the dynamics of

connectivity is not simply due to sensory habituations. The left medial visual region and left precentral

gyrus did not show high level of correlations in general. But it had small positive correlations at the

beginning of the run, went down to around zero, and then went back to small positive correlations. What

is more interesting is the dynamic connectivity between the right supramarginal gyrus and posterior

cingulate cortex, where the connectivity switched between positive and negative values during the whole

course.

Figure 5 Time courses of dynamic connectivity (Fisher's z) for three pairs of brain regions. Each thinner

line represents the time course of one subject, and the thicker red lines represent the averaged time

courses. LMV, left medial visual; RMV, right medial visual; LPCG, left precentral gyrus; RSMG, right

supramarginal gyrus; PCC, posterior cingulate cortex.

3.2.4. The effects of sliding-window length

We repeated the ROI-based intersubject correlation analysis of dynamic connectivity using different

window length from 10 TRs to 40 TRs. The overall patterns of intersubject correlations of dynamic

connectivity looked similar (Figure 6), especially for the 30-TR window and 40-TR window results. But

the correlation values were weaker for shorter window lengths, especially for 10-TR window length. The

bottom row of Figure 6 illustrated the time courses of dynamic connectivity of a typical pair of regions,

i.e. the right supramarginal gyrus and posterior cingulate cortex. It can be seen that for short window, e.g.

10-TR window, the time course of dynamic connectivity were very noisy. Moreover, the fluctuations of

dynamic connectivity changed fast, and it seems that different subjects had different delays of certain

increases or decreases of connectivity. For longer window, the dynamic connectivity time courses

became smoother, and the fluactuations across subjects were more aligned across subjects, which in turn

gave rise to higher intersubject correlations.

Figure 6 The effects of sliding-window length on the intersubject correlations of dynamic connectivity

(top row), and on the time courses of dynamic connectivity between the right supramarginal gyrus and

posterior cingulate cortex (bottom row).

4. Discussion

In the current study, we proposed intersubject correlation analysis on the time courses of dynamic

connectivity during natural vision. We were able to identify intersubject consistent dynamic connectivity

at similar level as the intersubject correlations of regional activity, although the time courses of dynamic

connectivity were thought to be nosier than the original time series. By using seed regions from the

visual cortex and supramarginal gyrus, we demonstrated widespread brain regions that showed high

intersubject consistent dynamic connectivity with these seeds, although these regions themselves did not

show intersubject correlations of regional activity. These regions included high order association regions

such as the left precentral cortex and left inferior frontal gyrus, as well as the default mode network. The

intersubject consistent patterns of dynamic connectivity support the functional meaningfulness of

dynamic connectivity during movie watching, and suggest that dynamic connectivity could be a

complementary avenue to characterize the functions of a brain region.

　　　The brain regions that had the highest intersubject correlations of regional activity are mainly in

the posterior visual related regions, which are consistent with previous studies (Hasson et al., 2004;

Nummenmaa et al., 2012). In addition to this, the current study also found high intersubject correlations

among different levels of visual areas, and the highest intersubject correlation of dynamic connectivity

was between the left and right medial visual regions. This is interesting because the dynamic connectivity

is on top of overall high level of stable functional connectivity (Figure 4A). The observable dynamics of

connectivity among visual areas are also consistent with previous studies showing task modulated

connectivity among visual areas in different task conditions (Di et al., 2018, 2015; Di and Biswal, 2017).

　　　In addition to the dynamic connectivity among visual regions, the visual regions also showed

consistent dynamic connectivity with regions outside the visual network. Specifically, the medial visual

regions showed high intersubject correlations of dynamic connectivity with the bilateral supramarginal

gyrus, left precentral gyrus, and left inferior frontal gyrus, which are all high order association areas. The

stable connectivity results also showed that there were high stable connectivity among these association

regions, but weak stable connectivity with the visual regions, further confirmed that they were two

functional modules in the brain. In addition, the supramarginal gyrus, which is a part of the task positive

network (Fox et al., 2005), showed high intersubject correlations with the default mode network. The

stable connectivity results also showed negative correlations between the supramarginal gyrus and default

mode network regions, which further confirmed they were from two different functional modules. These

results suggest that the functional communications between the regions from different functional

modulates are not stable, but highly depend on task contexts (Bullmore and Sporns, 2012). It is also

consistent with findings that the connectivity between regions from different functional modules are more

dynamic and context dependent (Di and Biswal, 2019; Fu et al., 2017).

The supramarginal gyrus is the major region outside the visual cortex that showed high

intersubject correlations of regional activity. The involvements of supramarginal gyrus of intersubject

correlations are inconsistent in the literature (Hasson et al., 2004; Kauppi et al., 2010), which probably

due to different movies used. Given its critical role in empathy (Silani et al., 2013), it is reasonable to

find high intersubject correlations in the supramarginal gyrus during the watching of the animated movie,

which involves the understanding the intentions of different animated characters. More interestingly, we

also found that the default mode network showed high intersubject correlations of dynamic connectivity

with the supramarginal gyrus seeds. Similar to a previous study on the dynamics of intersubject

connectivity (Simony et al., 2016), both of the studies highlighted the critical role of the default mode

network in understanding of the narratives of a movie.

The selection of window length for dynamic connectivity analysis is nontrivial (Fu et al., 2014;

Zhang et al., 2013). The shorter the window length, the finer the temporal resolution for dynamic

connectivity could be. Nevertheless, there would also be less number of time points for each window,

resulting in noisier estimates of connectivity. The current results showed that the overall patterns of

intersubject correlation matrices of dynamic connectivity were similar across different window length.

But the effect sizes changed dramatically. From Figure 6, it seems that in shorter window length, e.g. 10-

TR, the dynamic connectivity time courses were not only noisier, but also with intersubject variability of

delays. The offset of connectivity fluactuations across subjects might compromise the intersubject

correlations. But when longer window length was used, the time courses become smoother, therefore

becoming more intersubject correlated. In other words, the consistent component of dynamic

connectivity reflects a slow and averaged pattern of dynamic communication between brain regions.

Further studies may use other correlations methods that were not sensitive to phase delays, e.g. coherence,

to capture dynamic patterns of connectivity in a finer temporal scale.

5. Conclusion

In the current study, we proposed intersubject correlation analysis on dynamic connectivity. The results

revealed widespread brain regions that showed consistent intersubject correlations of dynamic

connectivity. The consistent correlations support the functional significance of dynamic connectivity

during natural vision. The method may provide complementary approach to understand the dynamic

nature of brain functional integrations.

Acknowledgement:

This study was supported by grants from (US) National Institute of Health (R01 AT009829; R01

DA038895).

Conflict of interest

The authors declared that there is no conflict of interest.

References:

Allen, E.A., Damaraju, E., Plis, S.M., Erhardt, E.B., Eichele, T., Calhoun, V.D., 2014. Tracking whole-
374   brain connectivity dynamics in the resting state. Cereb. Cortex N. Y. N 1991 24, 663–76.
375   https://doi.org/10.1093/cercor/bhs352
376   Bullmore, E., Sporns, O., 2012. The economy of brain network organization. Nat. Rev. Neurosci. 13,
377   336–349. https://doi.org/10.1038/nrn3214
378   Chen, G., Shin, Y.-W., Taylor, P.A., Glen, D.R., Reynolds, R.C., Israel, R.B., Cox, R.W., 2016.
379   Untangling the relatedness among correlations, part I: Nonparametric approaches to inter-subject
380   correlation analysis at the group level. NeuroImage 142, 248–259.
381   https://doi.org/10.1016/j.neuroimage.2016.05.023
382   Di, X., Biswal, B.B., 2019. Toward Task Connectomics: Examining Whole-Brain Task Modulated
383   Connectivity in Different Task Domains. Cereb. Cortex 29, 1572–1583.
384   https://doi.org/10.1093/cercor/bhy055
385   Di, X., Biswal, B.B., 2017. Psychophysiological Interactions in a Visual Checkerboard Task:
386   Reproducibility, Reliability, and the Effects of Deconvolution. Front Neurosci 1–36.
387   https://doi.org/10.3389/fnins.2017.00573
388   Di, X., Biswal, B.B., 2015. Characterizations of resting-state modulatory interactions in the human brain.
389   J. Neurophysiol. 114, 2785–96. https://doi.org/10.1152/jn.00893.2014
390   Di, X., Fu, Z., Chan, S.C., Hung, Y.S., Biswal, B.B., Zhang, Z., 2015. Task-related functional
391   connectivity dynamics in a block-designed visual experiment. Front. Hum. Neurosci. 9, 1–11.
392   https://doi.org/10.3389/fnhum.2015.00543
393   Di, X., Zhang, Z., Biswal, B.B., 2018. Psychophysiological interaction and beta series correlation for task
394   modulated connectivity: modeling considerations and their relationships. bioRxiv 322073.
395   https://doi.org/10.1101/322073
396   Fox, M.D., Snyder, A.Z., Vincent, J.L., Corbetta, M., Van Essen, D.C., Raichle, M.E., 2005. The human
397   brain is intrinsically organized into dynamic, anticorrelated functional networks. Proc. Natl. Acad.
398   Sci. U. S. A. 102, 9673–8.
399   Friston, K.J., 2011. Functional and effective connectivity: a review. Brain Connect. 1, 13–36.
400   https://doi.org/10.1089/brain.2011.0008
401   Friston, K.J., Williams, S., Howard, R., Frackowiak, R.S., Turner, R., 1996. Movement-related effects in
402   fMRI time-series. Magn. Reson. Med. Off. J. Soc. Magn. Reson. Med. Soc. Magn. Reson. Med.
403   35, 346–55. https://doi.org/DOI 10.1002/mrm.1910350312
404   Fu, Z., Chan, S.-C., Di, X., Biswal, B., Zhang, Z., 2014. Adaptive covariance estimation of non-stationary
405   processes and its application to infer dynamic connectivity from fMRI. IEEE Trans. Biomed.
406   Circuits Syst. 8, 228–39. https://doi.org/10.1109/TBCAS.2014.2306732
407   Fu, Z., Tu, Y., Di, X., Biswal, B.B., Calhoun, V.D., Zhang, Z., 2017. Associations between Functional
408   Connectivity Dynamics and BOLD Dynamics Are Heterogeneous Across Brain Networks. Front.
409   Hum. Neurosci. 11. https://doi.org/10.3389/fnhum.2017.00593
410   Fu, Z., Tu, Y., Di, X., Du, Y., Pearlson, G.D., Turner, J.A., Biswal, B.B., Zhang, Z., Calhoun, V.D., 2018.
411   Characterizing dynamic amplitude of low-frequency fluctuation and its relationship with dynamic
412   functional connectivity: An application to schizophrenia. NeuroImage 180, 619–631.
413   https://doi.org/10.1016/j.neuroimage.2017.09.035
414   Fu, Z., Tu, Y., Di, X., Du, Y., Sui, J., Biswal, B.B., Zhang, Z., de Lacy, N., Calhoun, V.D., 2019.
415   Transient increased thalamic-sensory connectivity and decreased whole-brain dynamism in
416   autism. NeuroImage 190, 191–204. https://doi.org/10.1016/j.neuroimage.2018.06.003
417   Hasson, U., Nir, Y., Levy, I., Fuhrmann, G., Malach, R., 2004. Intersubject synchronization of cortical
418   activity during natural vision. Science 303, 1634–40. https://doi.org/10.1126/science.1089506
419   Hutchison, R.M., Womelsdorf, T., Allen, E. a., Bandettini, P. a., Calhoun, V.D., Corbetta, M., Della
420   Penna, S., Duyn, J.H., Glover, G.H., Gonzalez-Castillo, J., Handwerker, D. a., Keilholz, S.,
421   Kiviniemi, V., Leopold, D. a., de Pasquale, F., Sporns, O., Walter, M., Chang, C., 2013. Dynamic
422   functional connectivity: Promise, issues, and interpretations. NeuroImage 80, 360–378.
423   https://doi.org/10.1016/j.neuroimage.2013.05.079

Kauppi, J.-P., Jääskeläinen, I.P., Sams, M., Tohka, J., 2010. Inter-subject correlation of brain
hemodynamic responses during watching a movie: localization in space and frequency. Front.
Neuroinformatics 4. https://doi.org/10.3389/fninf.2010.00005
Lindquist, M.A., Xu, Y., Nebel, M.B., Caffo, B.S., 2014. Evaluating dynamic bivariate correlations in
resting-state fMRI: A comparison study and a new approach. NeuroImage 101, 531–546.
https://doi.org/10.1016/j.neuroimage.2014.06.052
Nastase, S.A., Gazzola, V., Hasson, U., Keysers, C., 2019. Measuring shared responses across subjects
using intersubject correlation. Soc. Cogn. Affect. Neurosci. 14, 667–685.
https://doi.org/10.1093/scan/nsz037
Nummenmaa, L., Glerean, E., Viinikainen, M., Jääskeläinen, I.P., Hari, R., Sams, M., 2012. Emotions
promote social interaction by synchronizing brain activity across individuals. Proc. Natl. Acad.
Sci. U. S. A. 109, 9599–9604. https://doi.org/10.1073/pnas.1206095109
Park, H.-J., Friston, K., 2013. Structural and Functional Brain Networks: From Connections to Cognition.
Science 342, 1238411–1238411. https://doi.org/10.1126/science.1238411
Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but systematic
correlations in functional connectivity MRI networks arise from subject motion. NeuroImage 59,
2142–2154. https://doi.org/10.1016/j.neuroimage.2011.10.018
Richardson, H., Lisandrelli, G., Riobueno-Naylor, A., Saxe, R., 2018. Development of the social brain
from age three to twelve years. Nat. Commun. 9, 1–12. https://doi.org/10.1038/s41467-018-
03399-2
Rosenthal, G., Sporns, O., Avidan, G., 2017. Stimulus Dependent Dynamic Reorganization of the Human
Face Processing Network. Cereb. Cortex 27, 4823–4834. https://doi.org/10.1093/cercor/bhw279
Silani, G., Lamm, C., Ruff, C.C., Singer, T., 2013. Right Supramarginal Gyrus Is Crucial to Overcome
Emotional Egocentricity Bias in Social Judgments. J. Neurosci. 33, 15466–15476.
https://doi.org/10.1523/JNEUROSCI.1488-13.2013
Simony, E., Honey, C.J., Chen, J., Lositsky, O., Yeshurun, Y., Wiesel, A., Hasson, U., 2016. Dynamic
reconfiguration of the default mode network during narrative comprehension. Nat. Commun. 7,
12141. https://doi.org/10.1038/ncomms12141
Teichert, T., Grinband, J., Hirsch, J., Ferrera, V.P., 2010. Effects of heartbeat and respiration on macaque
fMRI: Implications for functional connectivity. Neuropsychologia 48, 1886–1894.
https://doi.org/10.1016/j.neuropsychologia.2009.11.026
Zhang, Z.G., Fu, Z.N., Chan, S.C., Hung, Y.S., Motta, G., Di, X., Biswal, B.B., 2013. Adaptive window
selection in estimating dynamic functional connectivity of resting-state fMRI, in: 2013 9th
International Conference on Information, Communications & Signal Processing. IEEE, pp. 1–4.
https://doi.org/10.1109/ICICS.2013.6782935
459

461   **Table 1** Clusters with differential intersubject correlations of dynamic connectivity among the medial

462   visual, lateral visual, and supramarginal gyrus seeds. All clusters were thresholded at p < 0.001, and

463   cluster thresholded at p < 0.0167 (0.05 / 3) after false discovery rate (FDR) correction.

		MNI Coordinates				
cluster FDR	Voxel	x	y	z	Peak t	Label
Medial visual > (lateral visual + supramarginal)						
< 0.001	157	-51	8	23	6.00	Left precentral gyrus
< 0.001	439	-12	-100	-4	5.90	Occipital pole
< 0.001	87	-51	-49	50	5.23	Left supramarginal gyrus
< 0.001	86	-33	44	-10	5.15	Left lateral orbital gyrus
< 0.001	408	24	-88	-4	5.10	Right inferior occipital gyrus
0.007	46	-60	-58	5	5.05	Left middle temporal gyrus
< 0.001	95	66	-31	26	4.92	Right supramarginal gyrus
0.003	57	-48	-34	14	4.58	Left planum temporale
Lateral visual > (medial visual + supramarginal)						
< 0.001	165	15	-79	-4	6.09	Lingual gyrus
< 0.001	99	48	-70	8	5.98	Right inferior occipital gyrus
Supramarginal > (medial visual + lateral visual)						
< 0.001	1582	6	-40	44	6.92	Precuneus
< 0.001	142	-3	59	8	5.87	Medial superior frontal gyrus
< 0.001	254	27	11	47	5.50	Right middle frontal gyrus
< 0.001	105	-30	11	35	5.38	Left middle frontal gyrus
0.001	77	-45	-67	38	5.17	Left angular gyrus
< 0.001	90	48	-61	44	4.98	Right angular gyrus
< 0.001	85	-21	29	50	4.83	Left superior frontal gyrus
< 0.001	104	24	62	2	4.79	Right superior frontal gyrus
0.006	49	0	-91	11	4.40	Cuneus

465   MNI, Montreal Neurological Institute