1

Multi-level analysis of reproduction in the Antarctic midge, *Belgica antarctica*, identifies female and male accessory gland products that are altered by larval stress and impact progeny viability

Geoffrey Finch[1], Sonya Nandyal[1], Carlie Perrieta[1], Benjamin Davies[1], Andrew J. Rosendale[1,2], Christopher

J. Holmes[1], Josiah D. Gantz[3,4], Drew Spacht[5], Samuel T. Bailey[1], Xiaoting Chen[6], Kennan Oyen[1], Elise M.

Didion[1], Souvik Chakraborty[1], Richard E. Lee, Jr.[2], David L. Denlinger[4,5] Stephen F. Matter[1], Geoffrey M.

Attardo[7], Matthew T. Weirauch[6,8,9], and Joshua B. Benoit[1,*]

[1]Department of Biological Sciences, University of Cincinnati, Cincinnati, OH

[2]Department of Biology, Mount St. Joseph University, Cincinnati, OH, USA

[3]Department of Biology, Miami University, Oxford, OH

[4]Department of Biology and Health Science, Hendrix College, Conway, AR

[5]Departments of Entomology and Evolution, Ecology and Organismal Biology, The Ohio State University,

Columbus, OH

[6]Center for Autoimmune Genomics and Etiology, Cincinnati Children's Hospital Medical Center,

Cincinnati, OH 45229, USA;

[7]Department of Entomology and Nematology, University of California, Davis, Davis, CA 95616, USA;

[8]Divisions of Biomedical Informatics and Developmental Biology, Cincinnati Children's Hospital Medical

Center, Cincinnati, OH 45229, USA.

[9]Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH 45267, USA.

[*]Author for correspondence

Joshua B. Benoit

Department of Biological Sciences,

University of Cincinnati, Cincinnati, OH

Email: joshua.benoit@uc.edu

Phone: 513-556-9714

Abstract

The Antarctic midge, *Belgica antarctica*, is a wingless, non-biting midge endemic to Antarctica. Larval

development requires at least two years, but adult life lasts only two weeks. The nonfeeding adults

mate in swarms and females die shortly after oviposition. Eggs are suspended in a gel of unknown

composition that is expressed from the female accessory gland. This project characterizes molecular

mechanisms underlying reproduction in this midge by examining differential gene expression in whole

males, females, and larvae, as well as in male and female accessory glands. Functional studies were used

to assess the role of the gel encasing the eggs, as well as the impact of stress on reproductive biology.

RNA-seq analyses revealed sex- and development-specific gene sets along with those associated with

the accessory glands. Proteomic analyses were used to define the composition of the egg-containing gel,

which is generated during multiple developmental stages and derived from both the accessory gland

and other female organs. Functional studies indicate the gel provides a larval food source and thermal

and dehydration buffer, all of which are critical for viability. Larval dehydration stress directly reduces

production of storage proteins and key accessory gland components, a feature that impacts adult

reproductive success. Modeling reveals that bouts of dehydration may significantly impact population

growth. This work lays a foundation for further examination of reproduction in midges and provides new

information related to general reproduction in dipterans. A key aspect is that reproduction and stress

dynamics, currently understudied in polar organisms, are likely to prove critical for determining how

climate change will alter survivability.

Keywords: Antarctic midge, reproduction, stress biology, accessory glands, thermal buffering

1 INTRODUCTION

The Antarctic midge, *Belgica antarctica,* is long-lived, wingless, and the only insect endemic to maritime Antarctica(Convey & Block, 1996; Sugg, Edwards, & Baust, 1983). It has a patchy distribution along the western coast of the Antarctic Peninsula and South Shetland Islands, where it may form large aggregations into the thousands under favorable conditions (Convey & Block, 1996; SUGG et al., 1983). The larval period lasts two years; growth and development take place during the short austral summer, and larvae overwinter encased in ice (Usher & Edwards, 1984). Larvae commonly reside in damp areas and are non-selective feeders by consuming dead plant and animal matter, algae, and microorganisms (Edwards, & Baust, 1981; Strong, 1967). Larvae are extremely tolerant of numerous stresses including cold, dehydration, and UV exposure (JBenoit, Lopez-Martinez, Elnitsky, Lee, & Denlinger, 2009; Benoit, Lopez-Martinez, Michaud, et al., 2007; Lopez-Martinez et al., 2009; Lopez-Martinez, Elnitsky, Benoit, Lee, & Denlinger, 2008; Teets et al., 2008). Adult emergence is a fairly synchronous event occurring in early summer (Edwards, & Baust, 1981), and there is some evidence for protandry (Edwards, & Baust, 1981; Harada, Lee, Denlinger, & Goto, 2014). The wingless adults mate in swarms formed on rocks and other features of the substrate (Edwards, & Baust, 1981; Harada et al., 2014). Environmental stressors in Antarctica can be severe and highly variable over short distances and time periods (Convey, 1997; Kennedy, 1993), and swarming may play a role in locating and taking advantage of intermittent, favorable microhabitats (Convey & Block, 1996; Hahn & Reinhardt, 2006; Sugg et al., 1983). Mating swarms present a potential obstacle for establishing long-lasting colonies in a laboratory setting, as mating is likely facilitated by large-scale, synchronous adult emergence (Harada et al., 2014).

Adult females that emerge in the laboratory lay, with a few exceptions, a single egg mass, each containing about 40 eggs (Convey & Block, 1996; Edwards, & Baust, 1981; Harada et al., 2014; Sugg et al., 1983). Nevertheless, multiple matings are common, and multiple oviposition events have been

reported (Edwards, & Baust, 1981; Harada et al., 2014; Sugg et al., 1983). Specific underlying materials

transferred from the male to the female during copulation are unknown in *B. antarctica*. Eggs are

encased in a hygroscopic gel that has been suggested as a potential food source for developing larvae

(Edwards, & Baust, 1981; Harada et al., 2014). This gel is likely secreted by the female accessory gland

during oviposition, but the protein components of the gel are unknown. Little additional information is

available on reproduction in these extremophiles (Edwards, & Baust, 1981; Harada et al., 2014), and

chironomid reproduction in general is poorly studied beyond the basic descriptions of their reproductive

anatomy and impaired reproduction during exposure to toxic substances (Sibley, Ankley, & Benoit, 2001;

Vogt et al., 2007; Wensler & Rempel, 1962).

By contrast, reproduction in other dipteran species has been examined extensively (Alfonso-

Parra et al., 2016; Avila, Sirot, LaFlamme, Rubinstein, & Wolfner, 2011; J.B. Benoit, Attardo, Baumann.,

Michalkova, & Aksoy, 2015 ; A. G. Clark et al., 2007; Dottorini et al., 2007; Izquierdo et al., 2019; K. P. Lee

et al., 2008; McGraw, Clark, & Wolfner, 2008; Meier, Kotrba, & Ferrar, 1999; Papa et al., 2017; Polak et

al., 2017; Ravi Ram & Wolfner, 2007; Villarreal et al., 2018). In most Diptera, insemination does not

involve injection of a spermatophore; rather, male seminal fluid is usually transferred directly into the

female reproductive tract, often with the addition of a mating plug to reduce male-male competition as

is seen in mosquitoes and *Drosophila* (Giglioli & Mason, 1966; Lung & Wolfner, 2001; Mitchell et al.,

2015; Scolari et al., 2016). Some flies do utilize a spermatophore, which may facilitate quicker mating

while also creating a barrier to multiple inseminations (Kotbra, 1996). In the tsetse fly, *Glossina*

morsitans, a dipteran that uses a spermatophore, proteins making up the spermatophore are secreted

from the male accessory gland during copulation (Attardo et al., 2019; Scolari et al., 2016). The

spermatophore is deposited such that the spermatozoa are funneled efficiently to the openings of the

spermathecal ducts, allowing only one spermatophore to maintain this connection at a time (Attardo et

al., 2019; Scolari et al., 2016). In mosquitoes, accessory gland-specific proteins, along with the steroid

hormone, 20-hydroxy-ecdysone, are transferred to the female during copulation, producing a mating

plug (Dottorini et al., 2007; Mitchell et al., 2015; Rogers et al., 2008). In *An. gambiae* the mating plug

has multiple effects that promote copulation, enhance egg production, and even trigger egg laying

(Dottorini et al., 2007; Gabrieli et al., 2014; Mitchell et al., 2015; Thailayil, Magnusson, Godfray, Crisanti,

& Catteruccia, 2011). First-male precedence and last-male precedence have both been reported in

multiple species within the Diptera (Dixon, Coyne, & Noor, 2003; Gwynne, 2012; Price, 1997; Shutt,

Stables, Aboagye-Antwi, Moran, & Tripet, 2010), but it remains unknown how fertilization priority is

established in *B. antarctica*. Depending on the mating strategy of the species, contents of the

spermatophore may include a large amount of seminal fluid proteins (SFPs) and other factors or contain

primarily the sperm itself (Avila et al., 2011; Lung & Wolfner, 2001; Rogers et al., 2008; Scolari et al.,

2016). These seminal fluid proteins are suspected or demonstrated to have a variety of functions,

including induction of refractoriness in the female, counteracting protease activity of female secretions

with protease inhibitors, defending spermatozoa against microbial assault, and neutralizing reactive

oxidative species (Alfonso-Parra et al., 2016; Avila et al., 2011; Lung & Wolfner, 2001; Ravi Ram &

Wolfner, 2007; Shutt et al., 2010), all which can compromise sperm function and impair their ability to

fertilize. The particular cocktail and amounts of SFPs utilized reflects the selection pressure from life-

history differences, conspecific competition, and diverse reproductive strategies between species and

even within species (Hopkins et al., 2019; Hopkins, Sepil, & Wigby, 2017; Izquierdo et al., 2019; Papa et

al., 2017).

Secretions from the female accessory glands also play important roles in insemination and

oviposition in dipterans, as well as other insects. In the house fly, *Musca domestica,* female accessory

gland secretions enhance sperm viability (Degrugillier, 1985; Leopold & Degrugillier, 1973). In some

species, substances secreted from the female accessory gland are expelled with the eggs at oviposition

and often function as an adhesive, anchoring eggs to the substrate (Lococo & Huebner, 1980). Accessory

gland secretions may also provide protection from diverse biotic and abiotic stressors. In the medfly,

Ceratitis capitata, the primary components of the accessory gland secretion deposited during

oviposition are ceratotoxins, which act as potent antibacterial agents (Marchini, Bernini, Marri,

Giordano, & Dallai, 1991). Similarly, accessory gland secretions from the sand fly, *Phlebotomus papatasi*,

have antimicrobial effects that protect the eggs, developing embryos, and adult female reproductive

tract from microbial invasion (Rosetto et al., 2003). Some male seminal fluids also contain antimicrobial

peptides, probably for similar reasons as in the female (Lung, Kuo, & Wolfner, 2001; Poiani, 2006).

Female accessory gland proteins can also be a source of nutrition for developing progeny, either while

growth occurs in the mother or as a secreted food source to nourish external progeny (Benoit, Kölliker,

& Attardo, 2019; Benoit et al., 2015 ; Kaiwa et al., 2014; Kaulenas, 2012).

In this study, we use RNA-seq, proteomics, and functional analyses to examine the reproductive

physiology of *B. antarctica*. Specifically, male and female accessory glands are examined to identify

factors related to male accessory protein generation and synthesis of egg components during

oviposition. Proteomic analysis of the gel secretion is used to identify its components, while

comparative genomic analyses are used to identify orthologs of specific reproduction-associated genes

in mosquitoes and midges. Functional studies reveal that stress impinging on late instar larvae impacts

synthesis of gene products associated with reproduction, lowering both male and female reproductive

success. Furthermore, we determine that the gel likely acts to prevent egg dehydration and serves as

thermal buffer, preventing overheating of the eggs. Population growth modeling reveals that impaired

fecundity from larval stress may reduce reproduction below population replacement levels. Our analysis

shows that reproduction in the Antarctic midge is directly impacted by larval stress, and identifies novel

roles for products manufactured by the female accessory gland. These studies confirm that an

understanding of reproductive biology is critical for establishing how these Antarctic extremophiles are

able to survive and proliferate in the challenging polar environment.

2 MATERIALS AND METHODS

2.1 Midge collections

Antarctic midges were collected from islands near Palmer Station (64°46'S, 64°04'W) in January 2007

and January 2017. Males and females were separated based upon the major morphological characters

described previously (Convey & Block, 1996; Sugg et al., 1983; Usher & Edwards, 1984), homogenized

on-site, and stored in Trizol (Invitrogen) at -70 °C for shipment to the University of Cincinnati. Female

and male accessory glands were also dissected (N = 20-30) and stored in Trizol similar to whole body

stages.

Larvae were collected from the same location as adults. Larvae within organic debris were

returned to Palmer Station, and larvae were extracted into ice water with a modified Berlese funnel.

Following recovery, larvae were stored with substrate from their natural habitat (rocks, soils, moss, and

the alga *Prasiola crispa*, which serves as a food source for *B. antarctica*) at 2-4 °C. Larvae were shipped

to the University of Cincinnati and stored under similar conditions until they were used in studies

examining the impact of larval stress or gel presence on adult fertility or egg viability, respectively.

2.2 RNA extraction and processing

RNA was extracted from the midges by homogenization (BeadBlaster 24, Benchmark Scientific) in Trizol

reagent (Invitrogen), using manufacturer's protocols with slight modification based on other studies of

invertebrates (Hagan et al., 2018; Rosendale, Dunlevy, McCue, & Benoit, 2019). Extracted RNA was

treated with DNase I (Thermo Scientific) and cleaned with a GeneJet RNA Cleanup and Concentration

Micro Kit (Thermo Scientific) according to manufacturer's protocols. RNA concentration and quality

were examined with a NanoDrop 2000 (Thermo Scientific).

Poly(A) libraries were prepared by the DNA Sequencing and Genotyping Core at the Cincinnati

Children's Hospital Medical Center. RNA was quantified using a Qubit 3.0 Fluorometer (Life

Technologies). Total RNA (150-300 ng) was poly(A) selected and reverse transcribed using a TruSeq

Stranded mRNA Library Preparation Kit (Illumina). An 8-base molecular barcode was added to allow for

multiplexing and, following 15 cycles of PCR amplification, each library was sequenced on a HiSeq 2500

sequencing system (Illumina) in Rapid Mode. For each sample, 30-40 million paired-end reads at 75

bases in length were generated. Raw RNA-seq data have been deposited at the National Center for

Biotechnology Information (NCBI) Sequence Read Archive: Bio-project PRJNA576639. Along with the

RNA-seq samples collected for this study, larval (control, dehydration, and cryoprotective dehydration)

samples were acquired from Teets et al.(Teets et al., 2012) under the NCBI Bioproject PRJNA174315.

RNA-seq reads were trimmed for quality (Phred score limit of 0.05) and sequences with

ambiguities were removed. In addition, five and eight nucleotides were removed from the 5' and 3'

ends, respectively, and sequences shorter than 45 bases were removed. Reads before and after cleaning

and trimming were examined with FastQC for quality (S. Andrews

http://www.bioinformatics.babraham.ac.uk/projects/fastqc) to verify the quality of each set.

2.3 Gene expression analyses

RNA-seq analyses were conducted using two distinct pipelines. The first method utilized was CLC

Genomics (Qiagen), as previously described (Hagan et al., 2018; Rosendale et al., 2019).   Briefly, reads

were mapped to contigs with a cutoff of at least 80% of the read matching at 90% identity with a

mismatch cost of 2. Each read was permitted to align to only 20 contigs. Expression values were based

on total read counts in each sample calculated as transcripts per million reads mapped. EdgeR was used

to test significance among samples. A multiple comparison correction was performed (false discovery

rate, FDR). Genes were considered to be differentially expressed if the fold change was greater than 2.0

and the p-value was < 0.05. In addition, genes were required to have at least 5 mapped reads per

sample to be retained for further analyses. For whole-carcass expression analyses, genes were

considered sample-specific if enrichment was noted in relation to both of the other whole-carcass

datasets. For accessory gland specific analyses, genes were considered tissue specific if they were

enriched relative to the relevant whole-carcass dataset (female or male). In addition, a *de novo*

assembly was conducted on the female accessory gland RNA-seq datasets using Trinity (Grabherr et al.,

2011) based on standard methods to determine if bacterial symbionts were present in this organ and

could be transferred to the egg while in the gel.

The second method for examining transcript expression involved utilization of RNA-seq tools

available through the Galaxy software package (www.https://usegalaxy.org/) (Afgan et al., 2018;

Goecks, Nekrutenko, & Taylor, 2010) using Salmon with the suggested settings. Differential expression

between genes was examined with the DeSeq2 package (Love, Huber, & Anders, 2014). A general

linearized model assuming a binomial distribution followed by a false discovery rate (FDR) approach

were utilized to account for multiple testing. Cut-off values for significance, enrichment and sample-

specificity were the same as those used in analysis conducted with CLC Genomics.

Transcripts identified as sex- or development-specific were examined using the CLC-based

pipeline as there was over 95% overlap between each RNA-seq analysis method. Pathways enriched

within males, females, and larvae were identified with a combination of Database for Annotation,

Visualization and Integrated Discovery (DAVID (Huang da et al., 2009)), Blast2GO enrichment

analyses(Conesa et al., 2005), CLC gene set enrichment analysis (Clark & Ma'ayan, 2011), and g:Profiler

(Raudvere et al., 2019). Due to the taxonomic limitations of DAVID, sets of enriched transcripts were

compared by BLASTx to the *An. gambiae* and to the *D. melanogaster* RefSeq protein datasets to identify

homologous sequences. Blast hits (e-value < 0.001) from these two species were submitted to DAVID.

There was considerable overlap between the results, and only the CLC-based methods were used in

subsequent analyses.

Lastly, we utilized weighted correlation network analysis (WGCNA) to identify specific modules

of genes that have similar expression profiles across developmental stages and accessory glands

(Langfelder & Horvath, 2008). WGCNA was implemented in an R software package used to construct

correlation networks and describe correlation between gene expression across samples in RNA-seq or

microarray data. Genes sharing similar patterns of expression across samples were clustered into

modules to identify groups of biologically significant genes that were particular to one of the sample

groups. For this analysis, RNA-seq data were screened for genes of zero variance prior to WGCNA,

leaving 13,424 genes for signed network construction. The minimum module size allowed was 20 and

the soft power was set to 14 as determined by the package's scale-free topology function. Modules

exhibiting the highest Pearson correlation coefficient were selected for further analysis to determine

function and relationship to developmental stages and accessory glands. Modules identified as enriched

in a specific developmental stage or tissue were examined for enriched GO categories with the use of

g:Prolifer and DAVID, as described earlier.

2.4 Comparative analyses with other chironomid midges and *Anopheles* mosquitoes

This is the first study to examine genome-wide, sex- and stage-specific expression in a midge. However,

a recent study examines sex-specific expression in four species of anopheline mosquitoes, a clade not

distantly related to midges, along with expression specific to the male and female reproductive tracts

(MRT and FRT, respectively (Papa et al., 2017)). The species covered in this study were *An. gambiae, An.*

minimus, An. albimanus, and *An. arabiensis.* In addition, male accessory glands enriched genes for *B.*

antarctica were directly compared to those from Anopheles male accessory glands(Izquierdo et al.,

2019). Predicted gene sets from this study that had enriched expression in specific stages and organs

were used in comparative analyses with *B. antarctica.* The genomes of four species of chironomid

midges were also acquired: *Clunio marinus* (Kaiser et al., 2016)*, Parochlus steinenii* (Kim et al., 2017)*,*

Polypedilum vanderplanki (Gusev et al., 2014)*,* and *Polypedilum nubifer*(Gusev et al., 2014)*.* Studies that

resulted in these sequenced midge genomes did not include analyses of differential expression between

sexes.

For comparative analyses between midges, predicted gene sets for *B. antarctica* were compared

with genomes of each midge species, and results of these four analyses were pooled to establish

putative sets of genes common to all five species. This analysis also resulted in identification of

differentially expressed genes unique to *B. antarctica*. Similarly, predicted gene sets from *B. antarctica*

were compared with each species of *Anopheles* mosquito. Antarctic midge gene sets were compared to

Anopheles whole carcass gene sets, as well as reproductive tract-specific gene sets (Papa et al., 2017).

Lastly, we compared the expression of genes within the male accessory glands to orthologous gene sets

that are uniformly expressed in the male accessory glands of mosquitoes (Izquierdo et al., 2019). The

common gene sets produced by these analyses were then subjected to ontological analyses, using

gProfiler, to establish sex-specific enriched pathways. tBLASTp analyses (e-value < 0.001) were

performed using CLC Genomics Workbench (CLC bio Qiagen). Protein sequences were defined as

orthologs if they were reciprocal-best BLASTp hits having an e-value < 10^{-10}. Overlap was compared

between these analyses to produce putative sex-specific transcript sets.

Transcription factors (TFs) and their predicted DNA binding motifs were identified based on

methods used for other invertebrate genomes (Attardo et al., 2019; Benoit et al., 2016; Olafson et al.,

2019). In brief, putative TFs were identified by scanning the amino acid sequences of all proteins for

putative DNA binding domains using the HMMER software package (Eddy, 2009) and a compilation of

Pfam DNA binding domain models (Weirauch & Hughes, 2011). Experimentally determined DNA binding

motifs were then inferred from other species (*e.g., Drosophila*) based on amino acid identity, using

previously established rules (Weirauch et al., 2014). Using this collection of inferred DNA binding motifs,

we examined enrichment of each motif within the 500 and 2000 bp promoter regions of genes with

increased expression in each sex, larvae, and accessory glands. These results were then compared to the

expression profiles of each TF to determine specific TF candidates that might regulate sex and

reproduction associated genes.

2.5 PCR and qPCR analyses

Select genes of interest that were highly enriched within males and females were examined by PCR.

Total RNA was extracted from males, females, larvae, female accessory glands, and male accessory

glands as described previously in the RNA-seq section and used as a template for Superscript III reverse

transcriptase according to the manufacturer's protocols (Invitrogen). PCR was performed with gene-

specific primer pairs (Table S15) using a DNA polymerase kit (Promega). The PCR conditions were 95 °C

for 3 min, 35 cycles of 30 sec at 95 °C, 52-56 °C for 1 min, and 1 min at 70 °C using an Eppendorf

Mastercycler Pro Series. Three independent (biological) replicates were conducted for each sex or

tissue stage.

qPCR analyses were conducted based on previously developed methods (Rosendale, Romick-

Rosendale, Watanabe, Dunlevy, & Benoit, 2016). RNA was extracted as described previously for

independent biological replicates. Complementary DNA (cDNA) was generated with a DyNAmo cDNA

Synthesis Kit (Thermo Scientific). Each reaction used 250 ng RNA, 50 ng oligo (dT) primers, reaction

buffer containing dNTPs and 5 mmol•l^{-1} MgCl$_2$, and M-MuLV RNase H+ reverse transcriptase. KiCqStart

SYBR Green qPCR ReadyMix (Sigma Aldrich, St Louis, MO, USA) along with 300 nmol l^{-1} forward and

reverse primers, cDNA diluted 1:20, and nuclease-free water were used for all reactions. Primers were

designed using Primer3 based on contigs obtained from the transcriptome analysis (Table S15). qPCR

reactions were conducted using an Illumina Eco quantitative PCR system. Reactions were run according

to previous studies(A. J. Rosendale et al., 2016). Four biological replicates were examined for each sex,

and three biological replicates were examined for each accessory gland. Expression levels were

normalized to *rpl19* using the ΔΔCq method as previously described(Joshua B Benoit et al., 2018; A. J.

Rosendale et al., 2016). Fold change was compared between larvae, males, females, and accessory

glands followed by a Pearson correlation coefficient (r).

2.6 Proteomics analysis of accessory gland-derived gel

Samples were analyzed at the Proteomics and Metabolomics Laboratory at the University of Cincinnati.

Two proteomic samples were collected by removing eggs from the gel with a micropipette and

dissolving the gel in 1X PBS with 0.1% Tween. Samples (4 µg) were run on a 1D SDS PAGE gel and silver

stained to confirm the presence of proteins; at least 15 distinct proteins could be visualized. Based on

this initial characterization, gel proteins (6 µg) were run 2 cm into a 1D 4-12% Bis-Tris Invitrogen NuPage

gel using MOPS buffer. Lanes were excised, reduced with 10 mM dithiothreitol, alkylated with

Iodoacetamide and digested with trypsin according to the standard protocol (Heaven et al., 2016;

Turnier et al., 2018). The resulting peptides were concentrated with a speed vac centrifuge and

resuspended in 0.1% formic acid. Each sample (2 µg) was used in subsequent analyses. Nanoscale LC-

electrospray ionization-MS/MS (nanoLC-ESI-MS/MS) analyses were performed on a TripleTOF 5600

(Sciex, Toronto, ON, Canada) coupled to an Eksigent (Dublin, CA) nanoLC ultra nanoflow system. Protein

from each gel sample was loaded and analyzed as described (Heaven et al., 2016; Turnier et al., 2018).

The data were recorded using Analyst-TF (v.1.6) software and searched against the *B. antarctica* genome

(Kelley et al., 2014) using the Protein Pilot program (Sciex). Gel proteins were compared to those with

differential expression in specific tissues from our RNA-seq studies. Protein, carbohydrate, and lipid

content were examined through spectrophotometric assays based upon methods described in

Rosendale et al. (2019)

2.7 Thermal buffering by the accessory gland gel

The gel has been suggested to serve as a source of nutrients for newly emerged larvae (Convey, 1992;

Convey & Block, 1996; Harada et al., 2014). In addition to this role, we examined whether the gel

increases thermal buffering capabilities of the egg compared to eggs directly deposited on the local

substrate. To examine the effect, we placed an Omega thermocouple within six gels and immediately

adjacent to these six gels at a field location near Palmer Station. Temperature was measured every

minute over the course of three days.

To establish whether the gel prevents egg death caused by thermal stress, eggs with and

without gels were exposed to 20 °C for two hours. Females were allowed to lay their eggs onto filter

paper disks (Whatman) which were placed in 50 ml centrifuge tubes before being transferred to a 4 °C

water bath. Temperature was then ramped up to 20 °C over the course of four hours 4 °C per hour

before slowly being reduced back to 4 °C over the same course of time. These samples were compared

to those that were held statically at 4 °C without the gel. Following treatment, all eggs were maintained

at 4 °C and monitored for larval emergence.

2.8 Role of the accessory gland gel in dehydration

To determine if the accessory gland gel could prevent egg dehydration, we subjected midge eggs to

dehydrating conditions with and without the gel's presence. All eggs were removed from the gel with a

fine metal probe and half were carefully reinserted into the gel. Three groups of eggs with or without

the gel were moved to 75% RH at 4 °C for 12 hours. Following this treatment, eggs with no gel were

placed back into the gel. Viability was determined by counting the number of larvae that emerged from

the total number of eggs.

2.9 Impact of larval dehydration stress on fecundity

To determine if larval dehydration stress impacts adult fecundity, we performed combined RNA-seq

analysis (described previously) and exposure of larvae to dehydration stress, both standard and

cryoportective (Teets et al., 2012). RNA-seq studies were conducted on larvae that were quickly

dehydrated (30% water loss) and on those that had undergone a slower form of dehydration,

cryoprotective dehydration (30% water loss). The resulting data sets were then examined to find genes

with increased expression in male or female accessory glands.

360   To determine whether mating is directly impacted by dehydration stress, groups of 100 fourth

instar larvae (final larval instar) were held at 75% RH until they lost 40% of their water content.

Following dehydration, larvae were returned to the standard rearing conditions and monitored every 12

hours for the presence of pupae or newly emerged adults. Each adult was removed and stored

separately at 98% RH, 4°C until mating, which occurred no later than four days after treatment. The

following mating pairs were examined: males from dehydrated larvae vs. control females, females from

dehydrated larvae vs. control males, and both males and females from dehydrated larvae. Males and

females that failed to copulate were removed from the experiment.

2.10 Statistics

Replicates are independent biological samples. Sample sizes are listed in each method section or the

figure legend. Significance is indicated within each figure and/or in the figure legend. Statistical tests are

listed within the respective section in the methods or in the figure legends. All statistical analyses were

performed using JMP version 11 (SAS) or R-based packages.

2.11 Population level effects

To explore population level effects of dehydration, gel surrounding the egg mass, and thermal stress we

used a Leslie matrix approach (Lefkovitch, 1965). Here, the dominant eigenvalue of the matrix is the

population growth rate (λ). We simplified life history to egg to larvae to adult despite there being four

larval instars, potentially occurring over several years. For control populations we used a mean fecundity

of 42.9 (eggs laid), an egg survival rate of 0.82, and larval survival rate of 0.78. To determine the effects

of larval dehydration on population growth, we used fecundities (eggs laid) of 34.6, 35.6, and 21.5 for

male, female, and male and female dehydration, respectively; egg and larval survivorship were assumed

to be the same as control populations. To determine the effects of gel on population growth we used all

parameters (as for control populations), but reduced larval survival to 0.5, based on the experimental

data from our studies. Similarly, to investigate effects of thermal stress we reduced egg survivorship to

0.63, but left all other parameters at control values. We also investigated a worst case scenario with

male and female dehydration, no gel, and thermal stress. All values were derived from the previously

described experiments. We determined the dominant eigenvalue of each matrix using the function

"eigen" in R.

3 RESULTS

3.1 Description of mating and reproductive organs in *B. antarctica*

Beyond copulation and sex ratios (Edwards, & Baust, 1981; Harada et al., 2014; Sugg et al., 1983), little is

known about mating and reproductive aspects for *B. Antarctica.* In this study, the reproductive organs

were observed by dissection. Following copulation (Fig. 1A), a spermatophore containing sperm and

other, likely accessory gland products, are transferred as a bundle to the females (Fig. 1A, inset).

Females deposit a gel around the fertilized eggs, and it is within this gel that embryogenesis occurs (Fig.

1B). The source of the gel appears to be the female's accessory gland (Fig. 3C) because the gland is

depleted followed egg laying (Fig. 1D,E). The male reproductive organs, including the testes and

accessory glands, are depicted in Fig. 1F. The general organization and structure of the male and female

reproductive anatomy is similar to that of another midge, *Chironomus plumosus* (Wensler & Rempel,

1962) and mosquitoes (Masci et al., 2015), with the exception that the female accessory gland of *B.*

antarctica is dramatically enlarged.   It is this enlarged gland that we consider to be the source of the

gel deposited around the eggs.

3.2 RNA-seq analyses of *Belgica* reproduction

Differential transcript levels were determined for males, females, and larvae using two independent

pipelines using the *B. antarctica* genome as a reference (Kelley et al., 2014). The two pipelines yielded

95% overlap at our significance cut-off (two-fold difference, combined transcripts per million (TPM)

among all samples of at least 5, and a correction-based P-value < 0.05). Between 60% and 71% of the

reads from each RNA-seq set mapped to predicted genes. At the 4-fold or greater level for stage-

specificity, overlap between the two methods was over 99%; thus we used the CLC Genomics (Qiagen)

pipeline (Attardo et al., 2019; Rosendale et al., 2019) for the remaining analyses (Table S1). When

expression differences were compared among all developmental stages, each sex and associated

reproductive organs were most similar, followed by similarity between adults, while the larval stages

were the most unique (Fig. 2). When the *de novo* assembly was examined for presence of putative

microbial symbionts, we detected no substantial matches (Table S2), indicating the gel does not likely

serve as a source of potential microbial symbionts as in other insects (Kaiwa et al., 2014).

Individual comparisons between stages revealed 392 male-enriched genes, 1825 female-

enriched genes, and 862 genes enriched in larvae (Fig. 2, Table S3-S5). Specific gene ontology (GO)

categories were associated with each stage including carboxylic acid biosynthesis in males, DNA repair in

females, and aminoglycan metabolism in larvae (Fig. 3; Table S3-S5). When accessory glands were

specifically examined, 20 genes were enriched in the female accessory gland and had significantly higher

expression in females compared to other stages (Fig 4A; Table S6). A similar analysis for the male

accessory gland identified 25 enriched genes. GO categories associated with the female accessory gland

were associated with glycosylation and mucin biosynthesis. Notably, similar GO categories were

associated with gene products of the male accessory gland (Fig. 4B; Table S7).

Beyond this first analysis, weighted correlation network analysis (WGCNA, Langfelder &

Horvath, 2008) was used to identify clusters (modules) of highly correlated genes between

developmental stages and within specific reproductive organs (Fig. 5; Table S8-S12). Specific modules of

enriched expression were identified in each tissue and developmental stage (Fig. 5B,C). When GO

analyses were conducted for each developmental stage, unique GO categories were associated with

larvae, females, and males (Fig. 5). Female accessory glands had few enriched GO categories based on

WGCNA results; enriched categories included phosphatase binding, response to stimulus, and ion

channel activity. Male accessory glands also had a low number of GO categories associated with

enriched modules; these included metallopeptidase activity and integral components of the membrane.

These results provide correlated stage- and tissue-specific expression modules. For larvae, males, and

females, GO categories from stage specific modules showed overlap with our previous analysis (Fig. 3),

but unique categories were also identified (Fig. 5).

3.3 Proteomic analysis of accessory gland derived gel

Proteomic analysis of the gel surrounding the eggs revealed 24 associated proteins (Fig. 6). Three

proteins comprised a majority of the gel (Fig. 6B; Table S13), including a vitellogenin-like protein

(IU25_12621), larval serum protein (IU25_03947), and an apolipophorin (IU25_03809). The two

analyzed gel samples had a highly-correlated protein content (Pearson correlation = 0.92, Fig. 3C).

Expression analyses revealed that transcripts for most of the gel proteins are not directly generated in

the female accessory gland (Fig. 3D), but instead are produced in other female organs (e.g. IU25_12621)

or within the larvae (e.g. IU25_03947 and IU25_03809). qPCR validation of the protein gel components

confirmed that transcripts for each of the gel proteins are expressed in tissues identified in the RNA-seq

studies (Fig. 5E). Additional qPCR analyses for genes not involved in producing major components of the

gel served as validation for results obtained from other RNA-seq samples (Fig. 5E).

3.4 Transcriptional regulation of reproductive-associated factors

To compare potential mechanisms regulating gene expression between samples, we predicted putative

transcription factors (TFs, Fig. 7A, Table S14) and their DNA binding motifs. We next performed TF

binding site motif enrichment analysis in the upstream regulatory regions of our differentially expressed

gene sets (see Methods). This analysis revealed that fourteen TFs had significantly enriched binding

sites in at least one gene set (Fig. 7B). Of these, one TF showed enhanced binding in the 500bp

regulatory regions of male-enriched genes. The remaining thirteen showed enhanced binding in female-

enriched genes; we limited our analysis to the set of seven TFs with enhanced binding in both 500 and

2000bp regions of the female- and/or FeAG-enriched gene sets. Five members of this set showed higher

transcript levels in either the female or female accessory gland compared to other tissues and

developmental stages (Fig. 7C). The increased gene expression in the respective tissues or stage along

with enriched binding sites suggest that these TFS are key regulatory elements for female reproduction

in *B. antarctica*. Transcription factors include forkhead box protein 1 and mothers against

decapentaplegic (Mad) homolog 1. In *Drosophila* Mad has been identified as a participant in the

signaling pathway of *decapentaplegic* (dpp), a morphogenetic protein that plays a role in regulating the

development of egg polarity and early embryonic development (Xie & Spradling, 1998). Several

transcription factors identified in our screening remain uncharacterized with no assigned biological

functions.

3.5 Comparative analyses between anopheline mosquitoes and chironomid midges

Sex- and tissue-specific gene sets from *B. antarctica* were compared with predicted gene sets from four

species of *Anopheles* mosquitoes and four species of chironomid midges (Figs. 8 and 9). Additionally, *B.*

antarctica enriched gene sets were compared with gene sets enriched in male and female reproductive

tracts of the same four *Anopheles* mosquitoes (Papa et al., 2017) (Fig. 8 and 9) and in comparison to the

male accessory gland enriched genes from five *Anopheles* species (Izquierdo et al., 2019)

Genes enriched in females of *B. antarctica* showed the highest degree of conservation among all

species examined by a large margin. All five species of midges had 1396 genes in common, 1401 were

shared with all four *Anopheles* species, and 1267 were common to all 9 species examined (Fig. 8). Gene

ontology analysis of this set revealed enrichment in functions related to RNA, DNA, chromatin binding,

nuclease activity, chromosome and organelle organization, transport of RNA and proteins in and out of

the nucleus, RNA processing, and cell cycle regulation. No female-enriched genes detected in *B.*

antarctica were unique when compared to the other eight species examined. *Belgica* females have 122

genes with no orthologues in any of the other four midge species. However, this gene set did not show

significant enrichment in any GO functional categories.

Among males, 345 genes were common among midge species, and 337 genes were shared

between *B. antarctica* and all four mosquito species (Fig. 8). Genes common to all midge species were

enriched in functions associated with anion transport, alpha-amino acid catabolism, and carboxylic acid

transmembrane transport. As with females, no genes were identified in the male-enriched gene set that

were unique to *B. antarctica*. There were 16 genes unique to *B. antarctica* among the five midge

species, but, again, they were not significantly enriched in any single functional category.

The larva-enriched gene set was compared to the midge and mosquito gene sets. Over 600

genes were identified with orthologues common to all examined midge and mosquito species (Fig. 8).

The gene set common to all midge species was enriched in functional categories associated with chitin

metabolism, iron ion binding, and cytochrome P450-driven metabolism. Two identified genes were

unique to larvae of *B. antarctica* (Fig. 8; Table S5): one is uncharacterized, and the other putatively

encodes polyubiquitin B. Gene ontology analysis comparing larvae of *Belgica* with other midges revealed

88 unique genes, but no functional categories were significantly enriched for this gene set.

Gene sets specifically enriched in male and female reproductive tracts of each mosquito species

were compared to sex-specific gene sets in *B. antarctica* (Fig. 9; Table S6-7). Comparison of the female-

enriched gene set with the female reproductive tract (FRT)-specific gene set yielded 241 genes with

putative FRT orthologues. This set was particularly enriched in GO categories related to mitotic cell cycle

processes and regulation of gene expression. For example, several orthologues in this set code for

cyclins, cyclin dependent kinases, zinc-finger proteins, transcription factors, and transcription

termination factors. Additionally, a large number of genes in this set (~44%) mapped to apparently

uncharacterized orthologs. The comparison between male-enriched and Male Reproductive Tract (MRT)

specific genes yielded a common set of 48 genes. This set is dominated by CLIP-domain serine proteases

and cytochrome p450s, and also contains three glucose-dehydrogenases (Fig. 9). When *B. antarctica*

MAG-enriched genes are compared to those for mosquitoes, only four overlapping genes were

identified based on moderate or high expression reported in a previous study(Izquierdo et al., 2019) of

the mosquito MAG (Fig. 9C); these were identified as venom allergen, carboxypeptidase, serine

protease, and a gene of unknown function.

3.6 Dehydration reduces larval serum protein and possibly subsequent egg production

Since specific female accessory gland components are expressed in larvae, we asked whether larval

dehydration impacts expression of adult female accessory gland components (Fig. 10). To do so, we re-

analyzed results from a previous RNA-seq study (Teets et al., 2012) on dehydration and cryoprotective

dehydration of *B. antarctica* (Fig. 10). One of the major female accessory gland proteins (IU25_03947)

had significantly reduced expression when larvae experienced either dehydration or cryoprotective

dehydration (water loss specifically induced by cold temperatures) (Fig. 10A-B). This protein is a

component of the accessory gel and, as a hexamerin, serves as a protein reserve (Burmester, 1999;

Burmester, Massey Jr, Zakharkin, & Benes, 1998; Telfer & Kunkel, 1991). The dehydration-evoked

suppression in transcripts associated with reproductive-associated factors possibly results in reduced

amounts of materials invested in the progeny at birth and reduced egg output (Fig. 10D-F). These results

suggest that stress experienced as larvae may have a direct impact on female reproduction, most likely

acting through reduced expression of larval serum protein.

3.7 Dehydration reduces specific male accessory gland components and impacts fertilization

We next examined the impact of larval dehydration stress on male fertility. To do so, we analyzed

expression of accessory gland components in adult males after larval dehydration or cryoprotective

dehydration (Fig. 11A). Thirteen genes enriched in male accessory glands were differentially expressed

in the adult male following larval dehydration compared to fully-hydrated individuals (Fig. 11A).

Enrichment of glutathione transferase activity and phosphatidylcholine metabolic processes were

detected. Male body mass was not altered following dehydration stress, but mating was substantially

compromised (Fig. 11B-C). When males dehydrated as larvae mated with unstressed females, fewer

eggs were deposited, and additive reductions were noted if dehydrated males mated with females that

were dehydrated as larvae (Fig. 11C). These results indicate an impact on male fertility brought about

by dehydration stress during larval life.

3.8 Role of gel as a nutrient source

Analyses of major gel protein components (over 3% of the total proteins of the gel) was performed.

Amino acid content of the major protein components revealed that all essential amino acids can be

provided within the gel (Fig. 12A). Predicted glycosylation and phosphorylation sites of the proteins

suggest that these proteins could provide phosphate and sugar resources for the developing larva as the

gel is consumed (Fig. 12). Spectrophotometric analysis of the gel revealed that 81%, 14%, and 5% of the

caloric content consisted of proteins, carbohydrates, and lipids, respectively. Larvae were denied access

to the gel to determine its impact on larval growth (Fig. 12B). Fewer of the gel-less larvae were alive

after one month (Fig. 12B), and those remaining were nearly 30% smaller than larvae that hatched from

eggs with free access to the gel (Fig. 12C). Importantly, these larvae were provided access to algae and

other organic debris with and without gel, indicating that non-gel food resources do not fully

compensate for loss of the gel as a food source. These results underscore the value of the gel for

successful larval development.

3.9 Thermal and dehydration buffering by the gel

We performed experiments to determine whether the gel impacts dehydration resistance in developing

eggs. When gel-less eggs were held in desiccating conditions (75% relative humidity) at 4 °C for 12 h and

subsequently treated with water, no viable eggs were detected , even when the eggs were

subsequently covered with gel to promote growth (0% for all three replicates). By contrast, viability of

eggs encased in gel exceeded 80% under in the same conditions, suggesting that the accessory gel

protects eggs against dehydration-induced mortality. These results indicate that the gel is likely critical

for maintaining water homeostasis within the egg and possibly for the developing embryo as well.

We also examined whether the gel acts as a thermal buffer. First, temperature changes were

examined within the gel (next to the eggs) and on the surface immediately adjacent to the eggs (Fig.

13A). The gel buffered temperature changes during the course of the day by reducing both the

maximum temperature and the rate of temperature change (Fig. 13). Gel-less eggs exposed to 20 °C

displayed reduced viability when compared to eggs encased in gel and to those held constantly at 4 °C

(Fig. 13). These results indicate that the accessory gland gel provides both thermal and dehydration

protection to the eggs.

3.10 Population modeling

Dehydration of males, females, and both males and females together reduced population growth

relative to control populations (Fig. 14). The combined effect of male and female dehydration had the

largest effect, reducing population growth by nearly 21%. Lack of gel also negatively impact population

growth, resulting in a 13% reduction relative to controls. The effect of thermal stress on egg viability

would reduce population growth by 8%. In spite of these scenarios, populations would still be expected

to increase in abundance under each specific stress. Under the "worst case scenario" where fecundity is

reduced by male and female dehydration, larval survival is reduced by thermal stress, and there is a lack

of gel reducing egg survival, population growth rate is negative (λ = -0.95), indicating declining

population size. We note that all these growth rates are fairly liberal because of the relatively short time

over which larval survival was observed (12 days) relative to the time spent as larvae in nature. In

addition, it is likely that other factors, such as starvation, freezing, or pathogen attack, could occur,

resulting in additive impacts on population growth.

4 DISCUSSION

This study examined reproductive biology of the Antarctic midge with the goal of establishing key

molecular mechanisms associated with male and female biology. Combined RNA-seq and proteomics

established the transcriptional components of reproduction and protein constituents deposited in the

gel surrounding the egg. Specifically, we examined whether the gel that encases the eggs alters egg

viability and larval survival and examined the impact of larval dehydration exposure on adult fertility.

Little is known about midge reproductive biology and we have summarized the major findings of this

study in Figure 15. Thus, this study will hopefully provide a foundation for the fields of Antarctic and

midge reproductive biology. Lastly, we modeled the impact of multiple reproductive factors on

population growth, each of which exert minor effects but in combination could yield a negative growth

rate. Our results highlight the importance of understanding the reproductive biology of this Antarctic

insect, a species restricted to a limited geographic region and a specific habitat.

The large proportion of apparently conserved, female-specific or female-enriched genes that are

uncharacterized (≈44%) suggests there are many aspects of *Belgica* female physiology that remain

poorly understood. However, the results of the GO analysis of the female-enriched gene set are

consistent with results reported for *Drosophila* and other insect systems (Attardo et al., 2019; Olafson et

al., 2019; Panfilio et al., 2019; Swanson, Wong, Wolfner, & Aquadro, 2004). In an analysis of genes

differentially expressed between germline-naive and gonadectomized females in *Drosophila*, Parisi *et al*.

found enrichment in terms associated with metabolism (Parisi et al., 2004) . In fact, genes associated

with energy storage and utilization, such as digestive proteases or lipid storage genes, showed increased

expression in germline-naive males and females. This suggests that enrichment of such genes would be

expected in the larval gene set, even if adult *Belgica* were active eaters which they are not. Indeed, the

larva enriched gene set is replete with putative trypsins, chymotrypsin like proteins, and lipases (Cao &

Jiang, 2017; Graveley et al., 2011; McKenna et al., 2016; Pauchet et al., 2010; Venancio, Cristofoletti,

Ferreira, Verjovski-Almeida, & Terra, 2009). Oogenic gene expression has also been linked to nutrient

sensing in *Drosophila* (Shim, Gururaja-Rao, & Banerjee, 2013; Terashima & Bownes, 2005). Enrichment

of terms associated with metabolism, including components of ribosomes as protein-building

machinery, identified in *B. antarctica* is most likely associated with oogenesis-related biosynthesis. This

could include mobilizing lipids and proteins from the adult fat body during vitellogenesis, as well as

synthesis of components of the egg gel excreted from the accessory gland. Transcription factor analyses

revealed a subset of genes likely to have critical roles in female biology and thus warrant further more

directed studies. One identified was mothers against decapentaplegic (Mad) homolog 1, which likely is

likely critical for development (Newfeld, Chartoff, Graff, Melton, & Gelbart, 1996; Xie & Spradling, 1998).

Many enriched GO terms can also be tied to gametogenesis and embryogenesis or *Belgica*

females. Terms associated with cell cycle control and DNA/RNA/chromatin handling are among the main

examples. Beyond meiotic divisions that result in formation of gametes, which of course takes place in

the male as well, female may have to exercise more precise control over the timing of oocyte

maturation, deploying cell-cycle modulators, transcription regulation machinery, and/or RNA silencing

factors (Lefebvre & Lécuyer, 2018; Qazi, Heifetz, & Wolfner, 2003; Soller, Bownes, & Kubli, 1999). TF

analyses revealed a NF-kappa relish-like transcription factor as a potential regulator of transcript levels

for the female-enriched gene set. Indeed, NF-kappaB transcription factors play well-established roles in

transcriptional regulation of cytokines, molecules involved in cell cycle control, and development

(Gilmore, 2006; Kim & Kim, 2005; Sosic & Olson, 2003). Curiously, this TF has its highest expression in

males despite a paucity of binding sites in the regulatory regions of the male-enriched gene set.

However, expression in the male accessory gland is limited and lower than that of the female or female

accessory gland. The meaning of these findings remains unclear, but indicate a potential critical role

related to accessory gland function in *Belgica*.

The female also produces developmental factors that play a role in successful embryogenesis,

such as mRNAs, hormone biosynthesis precursors, and general cytoplasmic components (sperm are

quite cytoplasm-poor relative to eggs) (Swevers, Raikhel, Sappington, Shirk, & Iatrou, 2005). Among the

most highly enriched genes in females relative to males (>1000 fold higher) are a handful of genes

known to be involved in embryonic patterning in insects, such as *nanos*, *oskar*, and multiple innexins,

components of invertebrate gap junctions involved in intercellular communication during

embryogenesis (Bauer et al., 2005; De Keuckelaere, Hulpiau, Saeys, Berx, & Van Roy, 2018; Güiza, Barria,

Saez, & Vega, 2018; Quinlan, 2016; Richard & Hoch, 2015). This lends support to the notion that the

female-enriched gene set is comprised largely of genes regulating and supporting oogenesis and

embryogenesis. In addition, paternal DNA must be de-compacted after fertilization, meaning that sperm

nucleotide binding proteins must be degraded (Doyen et al., 2015). The enzymes/proteins that

participate in this process are likely already present in the egg, and therefore maternally contributed

(Tirmarche, Kimura, Dubruille, Horard, & Loppin, 2016). This is followed by histone/chromatin assembly

which is dependent on maternally provided histones and nucleosome assembly factors (Loppin,

Dubruille, & Horard, 2015; Tirmarche et al., 2016). Male-contributed mitochondrial DNA must also be

degraded (DeLuca & O'Farrell, 2012; Sato & Sato, 2011). This further accounts for the prevalence of

terms associated with histones and chromosome organization. Finally, in *Drosophila,* nurse cells

surrounding the oocyte are known to polyploidize regions of their nuclear DNA to enhance transcription

and support provisioning of the oocyte (Bastock & St Johnston, 2008; Kaulenas, 1992; Orr-Weaver,

2015). This provides yet another physiological link to DNA replication and chromosome organization, but

chironomids lack nurse cells. It is uncertain whether other cells of the ovary act in a similar manner to

Drosophila nurse cells (Bastock & St Johnston, 2008).

The dominant GO terms associated with the male-enriched gene set are 'anion transport' and

'carboxylic acid biosynthesis.' These are both terms that can be linked to spermatogenesis and may also

be linked to the differing metabolic needs of males, females, and larvae: larvae eat, digest, and grow;

females conduct vitellogenesis, oogenesis, and embryogenesis; and males likely devote the bulk of their

energy store to spermatogenesis and mating (Perry, Harrison, & Mank, 2014). The two terms may be

interrelated. For example, genes involved in synthesis and transport of pyruvate, a carboxylate anion

and the starting substrate of the tricarboxylic acid (TCA) cycle, could be associated with either GO term.

The functional role of pyruvate metabolism in sperm viability is not known, but mitochondrial activity is

demonstrated to be important for fertility, and a mitochondrial pyruvate transporter that is almost

uniquely expressed in the male germline has been identified in placental mammals (Vanderperre et al.,

2016). However, inhibition of pyruvate transporters in the mitochondrial membrane does not appear to

affect sperm motility, lending support to the notion that sperm motility is fueled primarily by glycolysis

(not import of pyruvate into mitochondria) (Vanderperre et al., 2016); alternatively, amino acid

metabolism may play a role in motility, as it has been associated with sperm motility in *An. gambiae*

(Izquierdo et al., 2019). TCA-based energy production must play a role in some other aspect of sperm

viability (Vanderperre et al., 2016). The fact that a mitochondrial pyruvate transporter is important in

sperm viability suggests that pyruvate transport and metabolism contribute to the significance of the GO

terms 'tricarboxylic acid cycle,' 'monocarboxylic acid metabolic process,' and 'anion transport.'

Additionally, sperm are thought to take up lactate and pyruvate through general monocarboxylic acid

transporters (MCTs) in humans (Ramalho-Santos et al., 2009; Rato et al., 2012; Vanderperre et al.,

2016). Perhaps there is an analogous uptake of energetically important organic acids in the seminiferous

tubules of *Belgica* and other insects. Finally, pyruvate may be useful in supplying Acetyl-CoA for the

histone acetylation that is essential for chromatin condensation during spermatogenesis (Ramalho-

Santos et al., 2009; Rato et al., 2012; Vanderperre et al., 2016).

Overall, males showed far fewer genes with enriched expression relative to the female than vice

versa. Of the 12 genes that are >1000 fold higher in the male compared to other stages, three are

putative metalloendopeptidases. These may be involved in the proteolytic processing of amyloid

precursor proteins (APPs), which are integral components of sperm membranes in humans, though their

specific roles have not been determined (Silva et al., 2015). Also of interest in this small set of highly

enriched genes are a few genes commonly, though not exclusively, associated with immunity: a leucine

rich immune protein TM, a toll protein, and an apparent homolog of the transcription factor NF-X1

(Brucker, Funkhouser, Setia, Pauly, & Bordenstein, 2012; De Gregorio, Spellman, Rubin, & Lemaitre,

2001; Irving et al., 2001; Palmer & Jiggins, 2015). This transcription factor binds X-box motifs to regulate

many eukaryotic genes and is generally thought to be a negative regulator of transcription (Stroumbakis,

Li, & Tolias, 1996). The homolog of this transcription factor identified in *D. melanogaster*, named

shuttlecraft (stc), is essential for normal embryonic development and is expressed most highly in the

embryo CNS. Moreover, expression of this gene is highest in the ovary, and the promoters of many

maternally contributed genes known to be crucial in embryonic development contain X-box motifs, such

as *oskar, torpedo, pumilio,* and vitelline membrane proteins (Stroumbakis et al., 1996). Therefore it is

tempting to suggest that, unlike in *Drosophila*, it is for some reason the male who produces this critical

TF, transferring it to the female during copulation in order to trigger or facilitate proper development of

their progeny. In support of the notion that the NF-X1 gene product is transferred to the female during

copulation, expression of the gene is roughly 24-fold higher in the male accessory gland than in the

whole carcass.

Gene ontology analysis of the larvae-enriched gene set revealed a preponderance of terms

related to peptidase, hydrolase, and detoxification activity. These GO terms are related to digestion and

detoxification of ingested materials by the growing larvae – neither of which would be relevant issues

for the non-feeding adults. One cytochrome p450s (CYP6Z2), up-regulated in larvae, has been implicated

in chemical resistance in *Anopheles* mosquitoes(Malik et al., 2016) and therefore may be critical for

larval survival in their potentially toxic microhabitat, such as seabird guano (Rial et al., 2016).

Besides the ingestion of food, larvae of *B. antarctica* are also longer-lived and must face the

seasonal challenges of permanent residence in Antarctica (Edwards, & Baust, 1981; Sugg et al., 1983). In

particular, larvae must survive freezing and desiccation during the austral winter along with potential

thermal stress during summer (Benoit et al., 2009; Benoit, Lopez-Martinez, Michaud, et al., 2007; Lopez-

Martinez et al., 2008; Michaud et al., 2008; Rinehart et al., 2006; Teets, Kawarasaki, Lee, & Denlinger,

2013). Genes enhancing survival under these conditions, such as heat shock proteins (Benoit et al.,

2009; Benoit, Lopez-Martinez, Michaud, et al., 2007; Lopez-Martinez et al., 2008; Michaud et al., 2008;

Rinehart et al., 2006; Teets et al., 2013), are expected to be up-regulated by larvae. Based on previous

studies in *Belgica* and other species that are desiccation and freeze tolerant, genes of importance could

include oxido-reductase related enzymes (e.g. cytochrome p450s), Late Embryogenesis Abundant (LEA)

proteins, protein repair methyltransferases, hemoglobins, aquaporins, and enzymes involved in

trehalose metabolism (Benoit et al., 2009; Benoit, Lopez-Martinez, Michaud, et al., 2007; Dunning et al.,

2013; Lopez-Martinez et al., 2008; Lv et al., 2010; Michaud et al., 2008; Rinehart et al., 2006; Ronges,

Walsh, Sinclair, & Stillman, 2012; Teets et al., 2013; J. Zhang, Marshall, Westwood, Clark, & Sinclair,

2011). Many of these categories are indeed enriched in the larvae and are likely substantial factors in

the stress resistance abilities of *B*. *antarctica*. Note that the presence of hemoglobin in insect

hemolymph is unique to chironomids (Gusev et al., 2014; Kaiser et al., 2016; S.-M. Lee, Lee, Park, & Choi,

2006) and furthermore it is thought to be unique to the larval stage. This could partially explain the

prevalence of iron binding as a GO term enriched in the larvae and conserved among the midge species

analyzed (Gusev et al., 2014; Kaiser et al., 2016; Lee et al., 2006).

Most prevalent among GO terms enriched in the larval gene set are those associated with

aminoglycan metabolism. This set includes, for example, multiple putative N-

acetylgalactosaminyltransferases, which catalyze the initial glycosylation of serine and threonine

residues (Tran & Ten Hagen, 2013). Mucin type O-glycosylation occurs commonly on proteins with

extracellular domains, comprising a portion of the extracellular matrix, and it is thought to be important

in intercellular communication and adhesion (Tran & Ten Hagen, 2013). In *Drosophila*, enzymes

responsible for building glycosylated proteins, such as mucins, are critical for embryonic development,

particularly for the CNS (Tian & Hagen, 2007; Tran & Ten Hagen, 2013; Zhang, Zhang, & Ten Hagen,

2008). Maternal and zygotic O-linked glycans have also been implicated in proper respiratory

development. These O-glycans are termed *pgant* in *Drosophila* (Tian & Hagen, 2007; Tran & Ten Hagen,

2013; Zhang et al., 2008). One essential gene, *pgant4*, is involved in regulating gut acidification, but this

is not the only such gene required in the digestive system (Tran & Ten Hagen, 2013; Tran et al., 2012).

The larva enriched gene set in *Belgica* includes putative homologs of *pgant4* and *pgant6*, as well as the

essential genes CG30463 and C1GalTA, both involved in mucin type O-linked glycosylation. However,

larva-enriched GO terms, as mentioned above, are dominated by hydrolysis and catabolic processes,

including glycosaminoglycan catabolism. This may be a sign of the breakdown of extracellular materials

associated with growth and development or an indication of a diet rich in aminoglycans.

Composition of male accessory glands has been a major focus in numerous insects. These

products influence fertilization rates, subsequent female receptivity to courtship, and are critical in the

composition of the spermatophore (Alfonso-Parra et al., 2016; Avila et al., 2011; Dottorini et al., 2007;

Gabrieli et al., 2014; Izquierdo et al., 2019; McGraw et al., 2008; Mitchell et al., 2015; Ravi Ram &

Wolfner, 2007; Rogers et al., 2008; Sirot, Wong, Chapman, & Wolfner, 2015; Thailayil et al., 2011;

Villarreal et al., 2018). Many of the male accessory gland genes from *Belgica* have orthologs based on

predicted genes from midge and mosquito genomes, but few overlapping orthologs were identified that

are expressed in the male accessory gland of *B. antarctica* compared to the male reproductive tissues of

mosquitoes (Izquierdo et al., 2019; Papa et al., 2017). This is not surprising since a similar lack in

orthology among male accessory gland products has been observed between *Drosophila* and *Glossina*

and other higher flies (Attardo et al., 2019; Scolari et al., 2016). We did not conduct biological

examination of specific roles for accessory gland proteins from *B. antarctica*, such as whether their

transfer during mating impacts refractoriness of females, as noted in many species (Avila et al., 2011;

Baldini, Gabrieli, Rogers, & Catteruccia, 2012; Ravi Ram & Wolfner, 2007; Sirot et al., 2015). There is an

enrichment for genes associated with glycoprotein synthase, which is unsurprising as glycoproteins are

common constituents of seminal fluid (Avila et al., 2011; Poiani, 2006). In addition, there were specific

serine proteases, immune factors, and products involved in response to oxidative stress. These are

similar to those observed in other fly species(Avila et al., 2011; Baldini et al., 2012; Ravi Ram & Wolfner,

2007; Sirot et al., 2015; Tian et al., 2017) and likely serve to preserve sperm viability but could impact

other aspects such as female biology (Abraham et al., 2016; Denis et al., 2017). These factors are likely

critical to male success, as this midge will mate on multiple occasions. It is important to note that

females will deposit eggs with and without fertilization (Harada et al., 2014), thus suggesting that factors

supplied in the spermatophore are not essential for ovulation and oviposition as in other fly species

(Avila et al., 2011; Baldini et al., 2012; Ravi Ram & Wolfner, 2007; Sirot et al., 2015).

Our gene expression analysis is one of the few that has explicitly examined gene expression

within the female accessory gland. Many, but not all, products present within the gel were expressed in

the female's accessory gland. A comparison of our results to expression pattern in mosquito female

tracts (Papa et al., 2017) revealed few genes with overlapping expression(Papa et al., 2017). Few studies

in other flies explicitly examined products generated by the female's accessory gland. Even in

Drosophila, the female accessory gland (paraovaria) remains one of the most understudied organs

(Attardo et al., 2014). In tsetse flies, this organ provides nourishment to developing intrauterine larvae

(Attardo et al., 2019; Benoit et al., 2015 ; Attardo et al., 2014), but there are no similarities, beyond

standard housekeeping genes, between this organ in tsetse flies and *B. antarctica*. Of interest,

regulation of transcript expression within the female accessory gland seems to be conserved (Attardo et

al., 2014), suggesting that regulatory aspects may be similar but drive expression of specific genes

related to unique functions of this organ in different flies. Transcription factor analyses identified a

single gene (IU25_08656) that has enriched levels of both binding sites upstream of female accessory

gland enriched genes and is itself expressed in the female accessory gland. This is an uncharacterized

zinc finger proteins, but is likely to have a critical role in female reproductive function for *B*. *antarctica*.

784   The three major protein components of the gel are vitellogenin, larval serum protein, and

apolipophorin, all which are reasonable components of a gel whose primary function includes fueling

the development of larvae upon hatching. *Belgica* larvae exhibit "drinking behavior" shortly before

hatching, suggesting that up to this point embryonic development is fueled by nutrient reserves present

in the egg at the time of oviposition (Harada et al., 2014). Upon hatching, the gel is ingested by the larva,

making it the first meal fueling their development. Larval serum protein likely acts as an amino acid

reservoir, while vitellogenin and apolipoproteins could provide fatty acid, sugar, and protein reserves.

The gel may also contribute other elements in smaller amounts, such as pre-synthesized developmental

hormones in an inactive form, hormone precursors, proenzymes, and enzyme cofactors that are

important for continued development (Harada et al., 2014). Larval serum protein is a storage protein

belonging to the family of hexamerins (Burmester, 1999; Burmester et al., 1998; Telfer & Kunkel, 1991).

Genes for such proteins are often highly expressed in the final larval instar preceding pupation

(Burmester, 1999; Burmester et al., 1998; Telfer & Kunkel, 1991) and serve as a nutrient reserve for

developing pupae and newly emerged adult. During the sweeping morphological changes that

accompany metamorphosis in holometabolous insects like *Belgica*, such storage proteins are extracted

from the larval fat body, transferred into the hemolymph, and subsequently re-sequestered in the newly

formed adult fat body(Burmester, 1999; Burmester et al., 1998; Keeley, 1978; Larsen, 1976; Telfer &

Kunkel, 1991). The following scenario is the likely fate of the larval serum protein found in the egg gel:

Initially re-sequestered by the adult fat body and shuttled to the female accessory gland with

vitellogenin, or possibly sequestered in the female accessory gland during adult development after

pupation without a layover in the fat body. Importantly, lack of expression in the female adult indicates

these proteins must be synthesized in the larva for subsequent use by the adult.

Along with establishing components of the female accessory gland derived gel, a major goal of

this study was to identify functional roles of this gel and to determine if the gel components are

impacted by stress in the developing larvae. Based on our results, one of the main functions of the gel is

to provide a nutritional resource, which leads to a higher larval survival. Products of the accessory

glands of females have been documented as food sources for insect species among many orders(Attardo

et al., 2019; Benoit et al., 2015 ; Denlinger & Ma, 1974; Kaiwa et al., 2014; Ma, Denlinger, Jarlfors, &

Smith, 1975). Production of gel substances surrounding eggs have been noted in other species, such as

in stink bugs, where the gel serves as a source of nutrition, protects the eggs, and provides a vehicle to

transfer microbial symbionts (Kaiwa et al., 2014). Similar to our study, removal of the gel inhibits

juvenile development in stink bugs, but in contrast, removal of the gel around the stink bug eggs has

little impact on hatching success. In addition to providing a critical nutritional source and preventing

dehydration, we show that the egg gel acts as a thermal buffer that limits temperature extremes. Our *de*

novo assembly of the female accessory gland failed to detect signatures of microbial symbionts within

the gel, indicating that the gel does not serve as a mechanism of bacterial transfer to the developing

larvae. This lack of microbial presence could very well limit microbial exposure of the midge until after

larvae emerge from the gel. Accessory gland products also have been demonstrated to possess

antimicrobial properties(Avila et al., 2011; Otti, McTighe, & Reinhardt, 2013; Otti, Naylor, Siva-Jothy, &

Reinhardt, 2009); this is a possibility for *B. antarctica* because immune peptides are present in the gel.

Larval dehydration stress had a major impact on fertility of both adult males and females. The

most likely cause of reduced fecundity in males and females is a direct reduction in larval serum protein,

a hexamerin that acts as a storage protein in larvae (Burmester, 1999; Burmester et al., 1998; Telfer &

Kunkel, 1991). This hexamerin represents one of the highest expressed transcripts in developing larvae,

so its accumulation during the juvenile stages is likely a critical resource for production of eggs and

accessory gland components. This effect is likely even more pronounced in *B. antarctica* because adults

do not feed or even readily drink water (Benoit, Lopez-Martinez, Elnitsky, Lee, & Denlinger, 2007). Along

with acting as a nutritional source, the larval serum protein generated in juvenile stage of females is

likely incorporated into the accessory gland gel, suggesting that a reduction in this product may directly

impact gel composition. Larval nutritional status has a direct impact on fecundity in numerous insect

systems (Aguila, Hoshizaki, & Gibbs, 2013; Aguila, Suszko, Gibbs, & Hoshizaki, 2007; Rosa &

Saastamoinen, 2017), including other midges (Sibley et al., 2001). To the best of our knowledge, larval

dehydration has not previously been examined in relation to subsequent adult fecundity. Other stressful

conditions, such as chemical exposure, have impaired both larval development and subsequent adult

reproduction in a non-biting midge, *Chironomus riparius*(Vogt et al., 2007).

This study provides an encompassing view of reproductive biology of the Antarctic midge, from

molecular mechanisms to the impact of larval stress exposure on adult fecundity. Key findings are

summarized in Figure 15. This is followed by population growth modeling to establish how these factors

directly impact persistence of this insect in its limited Antarctic range. Population modeling revealed

that each factor (dehydration stress, lack of gel, thermal buffering), by itself, has a small impact on

population growth but combined factors likely result in negative population growth. The limited

reproductive window of 2-3 weeks makes understanding both male and female reproduction critical for

understanding how this midge survives in Antarctica. Studies on the reproductive biology of flies have

been limited largely to *Drosophila* and disease vectors (sand flies, mosquitoes, etc.), and our results

expand into Chironomidae to provide the groundwork for future studies with this dipteran system.

ACKNOWLEDGEMENTS

This work was supported by the National Science Foundation grant DEB-1654417 (partially) and United

States Department of Agriculture 2018-67013 (partially) to J.B.B., National Science Foundation grant

OPP-1341393 to D.L.D., and National Science Foundation grant OPP-1341385 to R.E.L. We thank the

staff at Palmer Station, Antarctica for assistance in logistics, experiments, and helping make the field

season a pleasant experience.

AUTHOR CONTRIBUTION

This study was designed and conceived by J.B.B. The manuscript was written by G.F. and J.B.B. RNA-seq

studies were conducted by G.F., A.J.R., B. D., and S.T.B. Molecular and physiological studies were

conducted and interpreted by S. N., C. P., G. F., C.J.H, E.M.D., and J.B.B. Transcription factor analyses

were contributed by X.C. and M.T.W. and interpreted by J.B.B. and G.F. Additional data and statistical

analyses were conducted by K.J.O., G.M.A., and J.B.B. Samples were collected by J.D.G., D.S., R.E.L., and

D.L.D. Population modeling was conducted by S.F.M. Figures were prepared by G.F. and J.B.B.

Illustrations were by G.M.A. All authors were responsible for editing the manuscript and have approved

publication.

DATA AVAILABILITY STATEMENT

All data generated for this project have been submitted to NCBI or made available in the supplemental

files.

Figure 1: Antarctic midge, *Belgica antarctica*, during reproduction. A. Mating pair, male on left. Inset,

spermatophore transferred to females immediately after copulation. Image is posterior end of female

and white material is the spermatophore. B. Female depositing eggs and accessory gland-derived gel. C.

Accessory gland (left middle, b) and ovaries (top left and right, a) of gravid females 3 days after adult

eclosion. D. Female accessory glands (left, a) and ovaries (top and right, b) following egg and gel

deposition. E. Egg mass following the completion of deposition. F. Male reproductive tract, a. testes, b,

accessory gland, and c, common duct.

Figure 2: Gene expression heat map of Antarctic midge, *Belgica antarctica*, during development,

between sexes, and for accessory glands. Hierarchical clustering of RNA-seq gene expression patterns

for males, females, larvae, and accessory glands based on sample distance (Euclidean distance matrix) of

differentially expressed contigs.

Figure 3: Genes uniquely enriched for the Antarctic midge, *Belgica antarctica* in males, females, and

larvae and associated gene ontology enrichment. A. Gene enriched in males (left) and gene ontology

(right), B. Gene enriched in females (left) and gene ontology (right), C. Gene enriched in larvae (left) and

gene ontology (right). Each box represents a specific category and color represent major GO groups.

Figure 4: Genes uniquely enriched in the Antarctic midge, *Belgica antarctica*, female and male

accessory glands and associated gene ontology enrichment. A. Genes enriched in female accessory

glands (left) and gene ontology (right), B. Genes enriched in male accessory gland (left) and gene

ontology (right). GO conducted with g:Prolifer (Raudvere et al., 2019).

Figure 5: Weighted gene co-expression network analysis (WGCNA) across developmental stages and

for male and female accessory glands of the Antarctic midge, *Belgica antarctica*. A. Average linkage

hierarchical clustering dendrogram of the genes. Modules, designated by color code, are branches of the

clustering tree. B. Correlation of module eigengenes to developmental and accessory gland traits. Each

row corresponds to a module eigengene and columns are traits. *, represents values with a significant

positive correlation for Pearson r ($P < 0.05$). C. Unsupervised hierarchical clustering heatmap (bottom)

and dendrogram (top) of module eigengenes and traits. Gene ontology (GO) analysis of eigengenes

associated with larvae (D), males (E), and females (F). GO conducted with g:Prolifer(Raudvere et al.,

2019) and visualized with REVIGO(Supek, Bošnjak, Škunca, & Šmuc, 2011).

Figure 6: Proteomic analysis of female accessory gland derived gel material from the Antarctic midge,

***Belgica antarctica*.** A. Female depositing eggs with gel and protein components of two gel samples

without eggs. B. Identification of proteins that represent at least 3% of total protein composition of the

accessory gland gel. C. Congruence of protein abundance and content between two gel samples. D.

Heatmap for transcript levels of gel-specific genes among larvae, females, males, and accessory glands.

E. qPCR validation of RNA-seq data. All tested genes have Pearson correlation coefficients over 0.85. Gel

specific genes have a Pearson correlation of 0.87.

Figure 7: Transcription factors (TFs) and TF binding sites associated with reproduction in the Antarctic

midge, *Belgica antarctica*. A. Relative abundance of transcription factors encoded by genomes of

midges and mosquitoes. TF families towards the top contain more TFs in *B. antarctica*. B. Enrichment

for specific TF binding site motifs in regulatory regions (2000 bp; 500 bp data not shown due to overlap

with 2000 bp) of genes expressed highly in specific stages and accessory glands. Groups of TFs are

separated by their motif enrichment profiles across samples. Those highlighted in orange are

significantly enriched within the specific stage or tissue. Scale for heatmap is set at relative abundance

on a Z scale of -2 to 2 across each row. C. Transcript levels of select TFs with significant motif enrichment

in the promoters of genes expressed in specific tissues (orange font color in panel B).Scale for heatmap

is set at relative abundance on a Z scale of -2 to 2 across each row. MAG, male accessory gland; FeAG,

female accessory glands.

Figure 8: Comparative analysis of female, male, and larvae-specific gene sets with mosquitoes and

midges to Antarctic midge, *Belgica antarctica*. A. Female-specific genes compared to midges (left),

mosquitoes (middle), and genes with enriched expression in the female reproductive tract (FRT) of

mosquitoes (right) (Papa et al., 2017). B. Male-specific genes compared to midges (left), mosquitoes

(middle), and genes with enriched expression in the male reproductive tract (MRT) of mosquitoes

(right)(Papa et al., 2017). C. Larvae-specific genes compared to midges (left) and mosquitoes (right).

Protein sequences were defined as orthologs if they had reciprocal-best BLASTp hits with an e-value <

10^{-10}.

Figure 9: Comparative analysis of accessory gland gene sets with mosquitoes and midges to the

Antarctic midge. A. Female accessory gland genes compared to midges (left), mosquitoes (middle), and

genes with enriched expression in the female reproductive tract of mosquitoes (right) (Papa et al.,

2017). B. Male accessory gland genes compared to midges (left), mosquitoes (middle), and genes with

enriched expression in the male reproductive tract of mosquitoes (right) (Papa et al., 2017). C. Overlap

between genes expressed in male accessory glands between mosquitoes and *B. antarctica*. Left, highly

enriched in *Anopheles* male accessory gland. Right, enriched in *Anopheles* male accessory gland.

Enrichment for *Anopheles* male accessory gland genes is based on values from Izquierdo et al. (2019).

Protein sequences were defined as orthologs if they had reciprocal-best BLASTp hits with an e-value <

10^{-10}.

Figure 10: Expression changes in gel-associated proteins in larvae following dehydration stress. A.

Transcript level changes in larvae for gel proteins following dehydration stress. RNA-seq studies were

acquired from Teets et al. (2012). Orange denotes significance between control and dehydrated larvae

based on RNA-seq analyses. B. Heat map of transcript levels for gel-associated proteins during

dehydration (D) and cryoprotective dehydration (CD) compared to control (C) that are components of

the gel proteome and significantly altered by dehydration. C. Total mass before eclosion, D. post-

eclosion total mass, E. mass change after eclosion, and total egg production in females when control

(non-dehydrated) and dehydrated larvae were allowed to complete development. Analysis of variance

was utilized to examine statistical differences with the use of R statistic packages. Bars above indicate

significance at $P < 0.05$.

Figure 11: Impact of larval dehydration stress on male fertility. A. Expression profiles of male-

associated genes with expression differences after larval dehydration (D, dehydration, C, control, and

CD, cryoprotective dehydration, MAG, male accessory gland). B. Mass of males used in mating

experiments from dehydrated or control larvae. C. Female fertility (control or dehydrated) following

copulation with dehydrated or control males. Analysis of variance was utilized to examine statistical

differences with the use of R statistic packages. Bars above indicate significance at $P < 0.05$ unless

otherwise noted.

Figure 12: Accessory gland gel is critical for larval development. A. Amino acid composition and

putative phosphate and glycosylation sites of gel proteins based on sequence information. Relative

amounts are based on comparison levels between columns. B. Survival of developing larvae with (black)

and without (gray) gel presence at larval ecdysis. Open circles are the average and filled circles are each

replicate. C. Larvae length after 20 days with and without gel at larval ecdysis. Bar indicates significance

at $P < 0.05$. Analysis of variance was utilized to examine statistical differences with the use of R statistic

packages. Bars above or beside indicate significance at $P < 0.05$.

Figure 13: Role of accessory gland gel in relation to thermal buffering of eggs. A. Thermal profile within

(orange) and outside (blue) gel-egg mixture under field conditions, B. Egg viability following exposure to

$20°C$ for three hours. Eggs without gel (constant) were held at $4°C$ for the duration of the trial. Maximum

temperature change during 24 hour period. D. Rate of temperature change (minimum to maximum).

Analysis of variance was utilized to examine statistical differences with the use of R statistic packages. *

above indicates significance at $P < 0.05$ unless otherwise noted.

Figure 14: Population growth is impacted by dehydration and thermal stress in developing larvae. A. Population growth following altered egg production due to dehydration exposure as larvae in males, females, and both sexes combined compared to control (no dehydration of larvae). B. Growth based on the presence or absence of the gel under favorable conditions. C. Impact of thermal stress on egg viability with and without accessory gland gel.

Figure 15: Summary of Antarctic midge reproduction. Larval development (four stages) is condensed into a single representation or all stages. Adults live approximately two-three weeks. Egg development occurs over thirty days. Impact of specific conditions are highlighted based on experimental evidence from this study.

Supplemental table 1. RNA-seq results for complete *Belgica antarctica* gene set. Expression values are in transcripts per million. RNA-seq datasets are available under the following NCBI Bioprojects PRJNA174315 and PRJNA576639.

1000

Supplemental table 2- BLAST results for female accessory gland specific *de novo* transcriptome against bacterial sequences from the NCBI nr database.

1003

1004 **Supplemental table 3.** RNA-seq results for female-enriched *Belgica antarctica* gene set. Expression

1005 values are in transcripts per million. RNA-seq datasets are available under the following NCBI Bioprojects

1006 PRJNA174315 and PRJNA576639.

1007

1008 **Supplemental table 4.** RNA-seq results for male-enriched *Belgica antarctica* gene set. Expression values

1009 are in transcripts per million. RNA-seq datasets are available under the following NCBI Bioprojects

1010 PRJNA174315 and PRJNA576639.

1011

1012 **Supplemental table 5.** RNA-seq results for larvae-enriched *Belgica antarctica* gene set. Expression

1013 values are in transcripts per million. RNA-seq datasets are available under the following NCBI Bioprojects

1014 PRJNA174315 and PRJNA576639.

1015

1016 **Supplemental table 6.** RNA-seq results for female accessory gland-enriched *Belgica antarctica* gene set.

1017 Expression values are in transcripts per million. RNA-seq datasets are available under the following NCBI

1018 Bioprojects PRJNA174315 and PRJNA576639.

1019

1020 **Supplemental table 7.** RNA-seq results for male accessory gland-enriched *Belgica antarctica* gene set.

1021 Expression values are in transcripts per million. RNA-seq datasets are available under the following NCBI

1022 Bioprojects PRJNA174315 and PRJNA576639.

1023

1024 **Supplemental table 8.** WGCNA module results for female-enriched *Belgica antarctica* gene set.

1025 Expression values are in transcripts per million. RNA-seq datasets are available under the following NCBI

1026 Bioproject PRJNA174315 and PRJNA576639.

1027

1028 **Supplemental table 9.** WGCNA module results for male-enriched *Belgica antarctica* gene set. Expression

1029 values are in transcripts per million. RNA-seq datasets are available under the following NCBI Bioprojects

1030 PRJNA174315 and PRJNA576639.

1031

1032 **Supplemental table 10.** WGCNA module results for larvae-enriched *Belgica antarctica* gene set.

1033 Expression values are in transcripts per million. RNA-seq datasets are available under the following NCBI

1034 Bioproject PRJNA174315 and PRJNA576639.

1035

1036 **Supplemental table 11.** WGCNA module results for female accessory gland-enriched *Belgica antarctica*

1037 gene set. Expression values are in transcripts per million. RNA-seq datasets are available under the

1038 following NCBI Bioproject PRJNA174315 and PRJNA576639.

1039

1040 **Supplemental table 12.** WGCNA module results for male accessory gland-enriched *Belgica antarctica*

1041 gene set. Expression values are in transcripts per million. RNA-seq datasets are available under the

1042 following NCBI Bioproject PRJNA174315 and PRJNA576639.

1043

1044   **Supplemental table 13.** Transcript levels for gel-derived proteins from the *Belgica antarctica* gene set.

1045   Expression values are in transcripts per million. RNA-seq datasets are available under the following NCBI

1046   Bioproject PRJNA174315 and PRJNA576639. Relative abundance of protein amounts were based on the

1047   number of protein fragments that match a specific genes.

1048

1049   **Supplemental table 14.** Transcript levels for transcription factors from the *Belgica antarctica* gene set.

1050   Expression values are in transcript per million. RNA-seq datasets are available under the following NCBI

1051   Bioproject PRJNA174315 and PRJNA576639.

1052

1053   **Supplemental table 15**- Quantitative PCR primers used for the validation of RNA-seq result.

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

REFERENCES

Abbott, K. C., & Dwyer, G. (2007). Food limitation and insect outbreaks: complex dynamics in plant–
herbivore models. *Journal of Animal Ecology, 76*(5), 1004-1014.
Abraham, S., Lara-Pérez, L. A., Rodríguez, C., Contreras-Navarro, Y., Nuñez-Beverido, N., Ovruski, S., &
Pérez-Staples, D. (2016). The male ejaculate as inhibitor of female remating in two tephritid
flies. *Journal of Insect Physiology, 88*, 40-47.
Afgan, E., Baker, D., Batut, B., Van Den Beek, M., Bouvier, D., Čech, M., . . . Grüning, B. A. (2018). The
Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018
update. *Nucleic acids research, 46*(W1), W537-W544.
Aguila, J. R., Hoshizaki, D. K., & Gibbs, A. G. (2013). Contribution of larval nutrition to adult reproduction
in *Drosophila melanogaster*. *Journal of Experimental Biology, 216*(3), 399-406.
Aguila, J. R., Suszko, J., Gibbs, A. G., & Hoshizaki, D. K. (2007). The role of larval fat cells in adult
Drosophila melanogaster. *Journal of Experimental Biology, 210*(6), 956-963.
Alfonso-Parra, C., Ahmed-Braimah, Y. H., Degner, E. C., Avila, F. W., Villarreal, S. M., Pleiss, J. A., . . .
Harrington, L. C. (2016). Mating-induced transcriptome changes in the reproductive tract of
female *Aedes aegypti*. *PLoS Neglected Tropical Diseases, 10*(2), e0004451.
Attardo, G. M., Abd-Alla, A. M., Acosta-Serrano, A., Allen, J. E., Bateta, R., Benoit, J., . . . Christensen, M.
B. (2019). The Glossina genome cluster: comparative genomic analysis of the vectors of African
trypanosomes. *bioRxiv*, 531749.
Attardo, G. M., Benoit, J. B., Michalkova, V., Patrick, K. R., Krause, T. B., & Aksoy, S. (2014). Ladybird late
homeodomain factor regulates lactation specific expression of milk proteins during pregnancy in
the tsetse fly (*Glossina morsitans*). *PLoS neglected tropical diseases, 8*, e2645.
Avila, F. W., Sirot, L. K., LaFlamme, B. A., Rubinstein, C. D., & Wolfner, M. F. (2011). Insect seminal fluid
proteins: identification and function. *Annual review of entomology, 56*, 21-40.
Baldini, F., Gabrieli, P., Rogers, D. W., & Catteruccia, F. (2012). Function and composition of male
accessory gland secretions in *Anopheles gambiae*: a comparison with other insect vectors of
infectious diseases. *Pathogens and global health, 106*(2), 82-93.
Bastock, R., & St Johnston, D. (2008). *Drosophila* oogenesis. *Current Biology, 18*(23), R1082-R1087.
Bauer, R., Löer, B., Ostrowski, K., Martini, J., Weimbs, A., Lechner, H., & Hoch, M. (2005). Intercellular
communication: the *Drosophila* innexin multiprotein family of gap junction proteins. *Chemistry*
& biology, 12(5), 515-526.
Benoit, J., Kölliker, M., & Attardo, G. (2019). Putting invertebrate lactation in context. *Science (New York,*
NY), 363(6427), 593-593.
Benoit, J. B., Adelman, Z. N., Reinhardt, K., Dolan, A., Poelchau, M., Jennings, E. C., . . . Richards, S.
(2016). Unique features of a global human ectoparasite identified through sequencing of the
bed bug genome. *Nature Communications, 7*. doi:10.1038/ncomms10165
Benoit, J. B., Attardo, G. M., Baumann., A. A., Michalkova, V., & Aksoy, S. (2015). Adenotrophic viviparity
in tsetse flies: potential for population control and as an insect model for lactation. *Annual*
review of entomology, In press
Benoit, J. B., Lopez-Martinez, G., Elnitsky, M. A., Lee, R. E., & Denlinger, D. L. (2007). Moist habitats are
essential for adults of the Antarctic midge, *Belgica antarctica* (Diptera : Chironomidae), to avoid
dehydration. *European Journal of Entomology, 104*(1), 9-14.
Benoit, J. B., Lopez-Martinez, G., Elnitsky, M. A., Lee, R. E., & Denlinger, D. L. (2009). Dehydration-
induced cross tolerance of *Belgica antarctica* larvae to cold and heat is facilitated by trehalose
accumulation. *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology,*
152(4), 518-523. doi:DOI 10.1016/j.cbpa.2008.12.009

Benoit, J. B., Lopez-Martinez, G., Michaud, M. R., Elnitsky, M. A., Lee, R. E., Jr., & Denlinger, D. L. (2007).
Mechanisms to reduce dehydration stress in larvae of the Antarctic midge, *Belgica antarctica*.
Journal Insect Physiol, 53(7), 656-667. doi:10.1016/j.jinsphys.2007.04.006
Benoit, J. B., Michalkova, V., Didion, E. M., Xiao, Y., Baumann, A. A., Attardo, G. M., & Aksoy, S. (2018).
Rapid autophagic regression of the milk gland during involution is critical for maximizing tsetse
viviparous reproductive output. *PLoS neglected tropical diseases, 12*(1), e0006204.
Brucker, R. M., Funkhouser, L. J., Setia, S., Pauly, R., & Bordenstein, S. R. (2012). Insect Innate Immunity
Database (IIID): an annotation tool for identifying immune genes in insect genomes. *PLoS One,*
7(9), e45125.
Burmester, T. (1999). Evolution and function of the insect hexamerins. *European Journal of Entomology,*
96, 213-226.
Burmester, T., Massey Jr, H. C., Zakharkin, S. O., & Benes, H. (1998). The evolution of hexamerins and the
phylogeny of insects. *Journal of Molecular Evolution, 47*(1), 93-108.
Cao, X., & Jiang, H. (2017). An analysis of 67 RNA-seq datasets from various tissues at different stages of
a model insect, *Manduca sexta. BMC genomics, 18*(1), 796.
Clark, A. G., Eisen, M. B., Smith, D. R., Bergman, C. M., Oliver, B., Markow, T. A., . . . MacCallum, I. (2007).
Evolution of genes and genomes on the *Drosophila* phylogeny. *Nature, 450*(7167), 203-218.
doi:nature06341 [pii]10.1038/nature06341
Clark, N., & Ma'ayan, A. (2011). Introduction to statistical methods for analyzing large data sets: Gene
Set Enrichment Analysis (GSEA). *Science Signaling, 4*(190), tr4.
Conesa, A., Gotz, S., Garcia-Gomez, J. M., Terol, J., Talon, M., & Robles, M. (2005). Blast2GO: a universal
tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics,*
21(18), 3674-3676. doi:10.1093/bioinformatics/bti610
Convey, P. (1992). Aspects of the biology of the midge, *Eretmoptera murphyi* Schaeffer (Diptera:
Chironomidae), introduced to Signy Island, maritime Antarctic. *Polar Biology, 12*(6-7), 653-657.
Convey, P. (1997). How are the life history strategies of Antarctic terrestrial invertebrates influenced by
extreme environmental conditions? *Journal of Thermal Biology, 22*(6), 429-440.
Convey, P., & Block, W. (1996). Antarctic Diptera: ecology, physiology and distribution. *European Journal*
of Entomology, 93, 1-14.
De Gregorio, E., Spellman, P. T., Rubin, G. M., & Lemaitre, B. (2001). Genome-wide analysis of the
Drosophila immune response by using oligonucleotide microarrays. *Proceedings of the National*
Academy of Sciences, 98(22), 12590-12595.
De Keuckelaere, E., Hulpiau, P., Saeys, Y., Berx, G., & Van Roy, F. (2018). Nanos genes and their role in
development and beyond. *Cellular and Molecular Life Sciences, 75*(11), 1929-1946.
Degrugillier, M. E. (1985). In vitro release of house fly, *Musca domestica* L.(Diptera: Muscidae),
acrosomal material after treatments with secretion of female accessory gland and micropyle cap
substance. *International Journal of Insect Morphology and Embryology, 14*(6), 381-391.
DeLuca, S. Z., & O'Farrell, P. H. (2012). Barriers to male transmission of mitochondrial DNA in sperm
development. *Developmental cell, 22*(3), 660-668.
Denis, B., Claisse, G., Le Rouzic, A., Wicker-Thomas, C., Lepennetier, G., & Joly, D. (2017). Male accessory
gland proteins affect differentially female sexual receptivity and remating in closely related
Drosophila species. *Journal of Insect Physiology, 99*, 67-77.
Denlinger, D. L., & Ma, W.-C. (1974). Dynamics of the pregnancy cycle in the tsetse *Glossina morsitans*
Journal of insect physiology, 20, 1015-1026.
Dixon, S. M., Coyne, J. A., & Noor, M. A. (2003). The evolution of conspecific sperm precedence in
Drosophila. Molecular Ecology, 12(5), 1179-1184.

Dottorini, T., Nicolaides, L., Ranson, H., Rogers, D. W., Crisanti, A., & Catteruccia, F. (2007). A genome-
wide analysis in *Anopheles gambiae* mosquitoes reveals 46 male accessory gland genes, possible
modulators of female behavior. *Proceedings of the National Academy of Sciences, 104*(41),
16215-16220.
Doyen, C. M., Chalkley, G. E., Voets, O., Bezstarosti, K., Demmers, J. A., Moshkin, Y. M., & Verrijzer, C. P.
(2015). A testis-specific chaperone and the chromatin remodeler ISWI mediate repackaging of
the paternal genome. *Cell reports, 13*(7), 1310-1318.
Dunning, L. T., Dennis, A. B., Park, D., Sinclair, B. J., Newcomb, R. D., & Buckley, T. R. (2013).
Identification of cold-responsive genes in a New Zealand alpine stick insect using RNA-Seq.
Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 8(1), 24-31.
doi:10.1016/j.cbd.2012.10.005
Eddy, S. R. (2009). A new generation of homology search tools based on probabilistic inference. *Genome
Informatics 2009: Genome Informatics Series, 23*, 205-211.
J.Edwards, J.S., & Baust, J. (1981). Sex ratio and adult behaviour of the Antarctic midge *Belgica
antarctica* (Diptera, Chironomdae). *Ecological Entomology, 6*(3), 239-243.
Gabrieli, P., Kakani, E. G., Mitchell, S. N., Mameli, E., Want, E. J., Anton, A. M., . . . Catteruccia, F. (2014).
Sexual transfer of the steroid hormone 20E induces the postmating switch in *Anopheles
gambiae*. *Proceedings of the National Academy of Sciences, 111*(46), 16353-16358.
Giglioli, M., & Mason, G. (1966). *The mating plug in anopheline mosquitoes.* Paper presented at the
Proceedings of the Royal Entomological Society of London. Series A, General Entomology.
Gilmore, T. D. (2006). Introduction to NF-κB: players, pathways, perspectives. *Oncogene, 25*(51), 6680.
Goecks, J., Nekrutenko, A., & Taylor, J. (2010). Galaxy: a comprehensive approach for supporting
accessible, reproducible, and transparent computational research in the life sciences. *Genome
Biology, 11*(8), R86.
Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., . . . Regev, A. (2011). Full-
length transcriptome assembly from RNA-Seq data without a reference genome. *Nature
biotechnology, 29*(7), 644-652. doi:nbt.1883 [pii]10.1038/nbt.1883
Graveley, B. R., Brooks, A. N., Carlson, J. W., Duff, M. O., Landolin, J. M., Yang, L., . . . Booth, B. W. (2011).
The developmental transcriptome of *Drosophila melanogaster*. *Nature, 471*(7339), 473-479.
Güiza, J., Barria, I., Saez, J. C., & Vega, J. L. (2018). Innexins: Expression, Regulation and Functions.
Frontiers in physiology, 9, 1414.
Gusev, O., Suetsugu, Y., Cornette, R., Kawashima, T., Logacheva, M. D., Kondrashov, A. S., . . . Shimura, S.
(2014). Comparative genome sequencing reveals genomic signature of extreme desiccation
tolerance in the anhydrobiotic midge. *Nature Communication, 5*, 4784.
Gwynne, D.T. (2012). Male mating effort, confidence of paternity, and insect sperm competition. *Sperm
competition and the evolution of animal mating systems*, 117.
Hagan, R. W., Didion, E. M., Rosselot, A. E., Holmes, C. J., Siler, S. C., Rosendale, A. J., . . . Nine, G. A.
(2018). Dehydration prompts increased activity and blood feeding by mosquitoes. *Scientific
reports, 8*(1), 6804.
Hahn, S., & Reinhardt, K. (2006). Habitat preference and reproductive traits in the Antarctic midge
Parochlus steinenii (Diptera: Chironomidae). *Antarctic Science, 18*(2), 175-181.
Harada, E., Lee, R. E., Denlinger, D. L., & Goto, S. G. (2014). Life history traits of adults and embryos of
the Antarctic midge *Belgica antarctica*. *Polar Biology, 37*(8), 1213-1217.
Heaven, M. R., Funk, A. J., Cobbs, A. L., Haffey, W. D., Norris, J. L., McCullumsmith, R. E., & Greis, K. D.
(2016). Systematic evaluation of data-independent acquisition for sensitive and reproducible
proteomics—a prototype design for a single injection assay. *Journal of Mass Spectrometry,
51*(1), 1-11.

Hopkins, B. R., Sepil, I., Thézénas, M.-L., Craig, J. F., Miller, T., Charles, P. D., . . . Pizzari, T. (2019).
Divergent allocation of sperm and the seminal proteome along a competition gradient in
Drosophila melanogaster. *Proceedings of the National Academy of Sciences, 116*(36), 17925-
17933.
Hopkins, B. R., Sepil, I., & Wigby, S. (2017). Seminal fluid. *Current Biology, 27*(11), R404-R405.
Huang da, W., Sherman, B. T., Zheng, X., Yang, J., Imamichi, T., Stephens, R., & Lempicki, R. A. (2009).
Extracting biological meaning from large gene lists with DAVID. *Current Protocols in*
Bioinformatics, Chapter 13. doi:10.1002/0471250953.bi1311s27
Initiative, I. G. G. (2014). Genome sequence of the tsetse fly (*Glossina morsitans*): vector of African
trypanosomiasis. *Science, 25*(6182), 380-386.
Irving, P., Troxler, L., Heuer, T. S., Belvin, M., Kopczynski, C., Reichhart, J. M., . . . Hetru, C. (2001). A
genome-wide analysis of immune responses in *Drosophila*. *Proceedings of the National Academy*
of Sciences of the United States of America, 98(26), 15119-15124. doi:10.1073/pnas.261573998
Izquierdo, A., Fahrenberger, M., Persampieri, T., Benedict, M. Q., Giles, T., Catteruccia, F., . . . Dottorini,
T. (2019). Evolution of gene expression levels in the male reproductive organs of *Anopheles*
mosquitoes. *Life science alliance, 2*(1), e201800191.
Kaiser, T. S., Poehn, B., Szkiba, D., Preussner, M., Sedlazeck, F. J., Zrim, A., . . . Hummel, T. (2016). The
genomic basis of circadian and circalunar timing adaptations in a midge. *Nature, 540*(7631), 69.
Kaiwa, N., Hosokawa, T., Nikoh, N., Tanahashi, M., Moriyama, M., Meng, X.-Y., . . . Ito, M. (2014).
Symbiont-supplemented maternal investment underpinning host's ecological adaptation.
Current Biology, 24(20), 2465-2470.
Kaulenas, M. (1992). Structure and function of the female accessory reproductive systems. In *Insect*
Accessory Reproductive Structures (pp. 33-121): Springer.
Kaulenas, M. (2012). *Insect accessory reproductive structures: function, structure, and development* (Vol.
31): Springer Science & Business Media.
Keeley, L. (1978). Endocrine regulation of fat body development and function. *Annual Review of*
Entomology, 23(1), 329-352.
Kelley, J. L., Peyton, J. T., Fiston-Lavier, A.-S., Teets, N. M., Yee, M.-C., Johnston, J. S., . . . Denlinger, D. L.
(2014). Compact genome of the Antarctic midge is likely an adaptation to an extreme
environment. *Nature Communications, 5*, 4611.
Kennedy, A. D. (1993). Water as a limiting factor in the Antarctic terrestrial environment: a
biogeographical synthesis. *Arctic and Alpine Research, 25*(4), 308-315.
Kim, S., Oh, M., Jung, W., Park, J., Choi, H.-G., & Shin, S. C. (2017). Genome sequencing of the winged
midge, *Parochlus steinenii*, from the Antarctic Peninsula. *GigaScience, 6*(3), giw009.
Kim, T., & Kim, Y. (2005). Overview of innate immunity in *Drosophila*. *Journal of biochemistry and*
molecular biology, 38(2), 121.
Kotrba, M. (1996). Sperm transfer by spermatophore in Diptera: new results from the Diopsidae.
Zoological Journal of the Linnean Society, 117(3), 305-323.
Langfelder, P., & Horvath, S. (2008). WGCNA: an R package for weighted correlation network analysis.
BMC bioinformatics, 9(1), 559.
Larsen, W. J. (1976). Cell remodeling in the fat body of an insect. *Tissue and Cell, 8*(1), 73-92.
Lee, K. P., Simpson, S. J., Clissold, F. J., Brooks, R., Ballard, J. W. O., Taylor, P. W., . . . Raubenheimer, D.
(2008). Lifespan and reproduction in *Drosophila*: new insights from nutritional geometry.
Proceedings of the National Academy of Sciences, 105(7), 2498-2503.
Lee, S.-M., Lee, S.-B., Park, C.-H., & Choi, J. (2006). Expression of heat shock protein and hemoglobin
genes in *Chironomus tentans* (Diptera, chironomidae) larvae exposed to various environmental
pollutants: a potential biomarker of freshwater monitoring. *Chemosphere, 65*(6), 1074-1081.

Lefebvre, F. A., & Lécuyer, É. (2018). Flying the RNA nest: *Drosophila* reveals novel insights into the
transcriptome dynamics of early development. *Journal of developmental biology, 6*(1), 5.
Lefkovitch, L. (1965). The study of population growth in organisms grouped by stages. *Biometrics*, 1-18.
Leopold, R. A., & Degrugillier, M. E. (1973). Sperm penetration of housefly eggs: evidence for
involvement of a female accessory secretion. *Science, 181*(4099), 555-557.
Lococo, D., & Huebner, E. (1980). The ultrastructure of the female accessory gland, the cement gland, in
the insect *Rhodnius prolixus*. *Tissue and Cell, 12*(3), 557-580.
Lopez-Martinez, G., Benoit, J. B., Rinehart, J. P., Elnitsky, M. A., Lee, R. E., & Denlinger, D. L. (2009).
Dehydration, rehydration, and overhydration alter patterns of gene expression in the Antarctic
midge, *Belgica antarctica*. *Journal of Comparative Physiology B-Biochemical Systemic and
Environmental Physiology, 179*(4), 481-491. doi:DOI 10.1007/s00360-008-0334-0
Lopez-Martinez, G., Elnitsky, M. A., Benoit, J. B., Lee, R. E., Jr., & Denlinger, D. L. (2008). High resistance
to oxidative damage in the Antarctic midge *Belgica antarctica*, and developmentally linked
expression of genes encoding superoxide dismutase, catalase and heat shock proteins. *Insect
Biochem Mol Biol, 38*(8), 796-804. doi:10.1016/j.ibmb.2008.05.006
Loppin, B., Dubruille, R., & Horard, B. (2015). The intimate genetics of *Drosophila* fertilization. *Open
biology, 5*(8), 150076.
Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550.
Lung, O., Kuo, L., & Wolfner, M. F. (2001). *Drosophila* males transfer antibacterial proteins from their
accessory gland and ejaculatory duct to their mates. *Journal of Insect Physiology, 47*(6), 617-622.
Lung, O., & Wolfner, M. F. (2001). Identification and characterization of the major *Drosophila
melanogaster* mating plug protein. *Insect biochemistry and molecular biology, 31*(6-7), 543-551.
Lv, D. K., Bai, X., Li, Y., Ding, X. D., Ge, Y., Cai, H., . . . Zhu, Y. M. (2010). Profiling of cold-stress-responsive
miRNAs in rice by microarrays. *Gene, 459*(1-2), 39-47. doi:DOI 10.1016/j.gene.2010.03.011
Ma, W. C., Denlinger, D. L., Jarlfors, U., & Smith, D. S. (1975). Structural modulations in the tsetse fly milk
gland during a pregnancy cycle. *Tissue Cell, 7*(2), 319-330. doi:0040-8166(75)90008-7 [pii]
Malik, A., Khan, K. M., Asif, M., Arsalan, H. M., Zahid, S., Manan, A., & Rasool, M. (2016). Development of
Resistance Mechanism in Mosquitoes: Cytochrome P450, the Ultimate Detoxifier. *Journal of
Applied and Emerging Sciences, 4*(2), pp100-117.
Marchini, D., Bernini, L. F., Marri, L., Giordano, P. C., & Dallai, R. (1991). The female reproductive
accessory glands of the medfly *Ceratitis capitata*: antibacterial activity of the secretion fluid.
Insect Biochemistry, 21(6), 597-605.
Masci, V. L., Di Luca, M., Gambellini, G., Taddei, A. R., Belardinelli, M. C., Guerra, L., . . . Fausto, A. M.
(2015). Reproductive biology in Anophelinae mosquitoes (Diptera, Culicidae): fine structure of
the female accessory gland. *Arthropod structure & development, 44*(4), 378-387.
McGraw, L. A., Clark, A. G., & Wolfner, M. F. (2008). Post-mating gene expression profiles of female
Drosophila melanogaster in response to time and to four male accessory gland proteins.
Genetics, 179(3), 1395-1408. doi:10.1534/genetics.108.086934
McKenna, D. D., Scully, E. D., Pauchet, Y., Hoover, K., Kirsch, R., Geib, S. M., . . . Arsala, D. (2016).
Genome of the Asian longhorned beetle (*Anoplophora glabripennis*), a globally significant
invasive species, reveals key functional and evolutionary innovations at the beetle–plant
interface. *Genome biology, 17*(1), 227.
Meier, R., Kotrba, M., & Ferrar, P. (1999). Ovoviviparity and viviparity in the Diptera. *Biological Reviews
of the Cambridge Philosophical Society, 74*(3), 199-258.
Michaud, M. R., Benoit, J. B., Lopez-Martinez, G., Elnitsky, M. A., Lee, R. E., & Denlinger, D. L. (2008).
Metabolomics reveals unique and shared metabolic changes in response to heat shock, freezing

and desiccation in the Antarctic midge, *Belgica antarctica*. *Journal of insect physiology, 54*(4),
645-655. doi:DOI 10.1016/j.jinsphys.2008.01.003
Mitchell, S. N., Kakani, E. G., South, A., Howell, P. I., Waterhouse, R. M., & Catteruccia, F. (2015).
Evolution of sexual traits influencing vectorial capacity in anopheline mosquitoes. *Science,*
347(6225), 985-988.
Newfeld, S. J., Chartoff, E. H., Graff, J. M., Melton, D. A., & Gelbart, W. M. (1996). Mothers against dpp
encodes a conserved cytoplasmic protein required in DPP/TGF-beta responsive cells.
Development, 122(7), 2099-2108.
Olafson, P. U., Aksoy, S., Attardo, G. M., Buckmeier, G., Chen, X., Coates, C. J., . . . Friedrich, M. (2019).
Functional genomics of the stable fly, *Stomoxys calcitrans*, reveals mechanisms underlying
reproduction, host interactions, and novel targets for pest control. *bioRxiv*, 623009.
Orr-Weaver, T. L. (2015). When bigger is better: the role of polyploidy in organogenesis. *Trends in*
Genetics, 31(6), 307-315.
Otti, O., McTighe, A. P., & Reinhardt, K. (2013). In vitro antimicrobial sperm protection by an ejaculate-
like substance. *Functional Ecology, 27*(1), 219-226.
Otti, O., Naylor, R. A., Siva-Jothy, M. T., & Reinhardt, K. (2009). Bacteriolytic activity in the ejaculate of an
insect. *American Naturalist, 174*(2), 292-295. doi:Doi 10.1086/600099
Palmer, W. J., & Jiggins, F. M. (2015). Comparative genomics reveals the origins and diversity of
arthropod immune systems. *Molecular biology and evolution, 32*(8), 2111-2129.
Panfilio, K. A., Jentzsch, I. M. V., Benoit, J. B., Erezyilmaz, D., Suzuki, Y., Colella, S., . . . Ioannidis, P.
(2019). Molecular evolutionary trends and feeding ecology diversification in the Hemiptera,
anchored by the milkweed bug genome. *Genome Biology, 20*(1), 64.
Papa, F., Windbichler, N., Waterhouse, R. M., Cagnetti, A., D'Amato, R., Persampieri, T., . . . Papathanos,
P. A. (2017). Rapid evolution of female-biased genes among four species of *Anopheles* malaria
mosquitoes. *Genome research, 27*(9), 1536-1548.
Parisi, M., Nuttall, R., Edwards, P., Minor, J., Naiman, D., Lu, J., . . . Oliver, B. (2004). A survey of ovary-,
testis-, and soma-biased gene expression in *Drosophila melanogaster* adults. *Genome Biology,*
5(6), R40. doi:10.1186/gb-2004-5-6-r40
Pauchet, Y., Wilkinson, P., Vogel, H., Nelson, D., Reynolds, S., Heckel, D. G., & Ffrench-Constant, R.
(2010). Pyrosequencing the *Manduca sexta* larval midgut transcriptome: messages for digestion,
detoxification and defence. *Insect molecular biology, 19*(1), 61-75.
Perry, J. C., Harrison, P. W., & Mank, J. E. (2014). The ontogeny and evolution of sex-biased gene
expression in *Drosophila melanogaster*. *Molecular biology and evolution, 31*(5), 1206-1219.
Poiani, A. (2006). Complexity of seminal fluid: a review. *Behavioral Ecology and Sociobiology, 60*(3), 289-
310.
Polak, M., Simmons, L. W., Benoit, J. B., Ruohonen, K., Simpson, S. J., & Solon-Biet, S. M. (2017).
Nutritional geometry of paternal effects on embryo mortality. *Proceedings of the Royal Society*
B: Biological Sciences, 284(1864), 20171492.
Price, C. S. (1997). Conspecific sperm precedence in *Drosophila*. *Nature, 388*(6643), 663.
Qazi, M. C. B., Heifetz, Y., & Wolfner, M. F. (2003). The developments between gametogenesis and
fertilization: ovulation and female sperm storage in *Drosophila melanogaster*. *Dev Biol, 256*(2),
195-211.
Quinlan, M. E. (2016). Cytoplasmic streaming in the *Drosophila* oocyte. *Annual review of cell and*
developmental biology, 32, 173-195.
Ramalho-Santos, J., Varum, S., Amaral, S., Mota, P. C., Sousa, A. P., & Amaral, A. (2009). Mitochondrial
functionality in reproduction: from gonads and gametes to embryos and embryonic stem cells.
Human reproduction update, 15(5), 553-572.

Rato, L., Alves, M. G., Socorro, S., Duarte, A. I., Cavaco, J. E., & Oliveira, P. F. (2012). Metabolic regulation is important for spermatogenesis. *Nature Reviews Urology, 9*(6), 330.

Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., & Vilo, J. (2019). g: Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic acids research*.

Ravi Ram, K., & Wolfner, M. F. (2007). Seminal influences: *Drosophila* Acps and the molecular interplay between males and females during reproduction. *Integrative and comparative biology, 47*(3), 427-445.

Rial, D., Santos-Echeandía, J., Álvarez-Salgado, X. A., Jordi, A., Tovar-Sánchez, A., & Bellas, J. (2016). Toxicity of seabird guano to sea urchin embryos and interaction with Cu and Pb. *Chemosphere, 145*, 384-393.

Richard, M., & Hoch, M. (2015). *Drosophila* eye size is determined by innexin 2-dependent decapentaplegic signalling. *Developmental Biology, 408*(1), 26-40.

Rinehart, J. P., Hayward, S. A. L., Elnitsky, M. A., Sandro, L. H., Lee, R. E., & Denlinger, D. L. (2006). Continuous up-regulation of heat shock proteins in larvae, but not adults, of a polar insect. *Proceedings of the National Academy of Sciences of the United States of America, 103*(38), 14223-14227. doi:DOI 10.1073/pnas.0606840103

Rogers, D. W., Whitten, M. M., Thailayil, J., Soichot, J., Levashina, E. A., & Catteruccia, F. (2008). Molecular and cellular components of the mating machinery in *Anopheles gambiae* females. *Proceedings of the National Academy of Sciences, 105*(49), 19390-19395.

Ronges, D., Walsh, J. P., Sinclair, B. J., & Stillman, J. H. (2012). Changes in extreme cold tolerance, membrane composition and cardiac transcriptome during the first day of thermal acclimation in the porcelain crab *Petrolisthes cinctipes*. *Journal of Experimental Biology, 215*(Pt 11), 1824-1836. doi:10.1242/jeb.069658

Rosa, E., & Saastamoinen, M. (2017). Sex-dependent effects of larval food stress on adult performance under semi-natural conditions: only a matter of size? *Oecologia, 184*(3), 633-642.

Rosendale, A. J., Dunlevy, M. E., McCue, M. D., & Benoit, J. B. (2019). Progressive behavioural, physiological and transcriptomic shifts over the course of prolonged starvation in ticks. *Molecular Ecology, 28*(1), 49-65.

Rosendale, A. J., Romick-Rosendale, L. E., Watanabe, M., Dunlevy, M. E., & Benoit, J. B. (2016). Mechanistic underpinnings of dehydration stress in ticks revealed through RNA-seq and metabolomics. *Journal of Experimental Biology, Submitted.* .

Rosetto, M., Belardinelli, M., Fausto, A., Marchini, D., Bongiorno, G., Maroli, M., & Mazzini, M. (2003). A mammalian-like lipase gene is expressed in the female reproductive accessory glands of the sand fly *Phlebotomus papatasi* (Diptera, Psychodidae). *Insect molecular biology, 12*(5), 501-508.

Sato, M., & Sato, K. (2011). Degradation of paternal mitochondria by fertilization-triggered autophagy in *C. elegans* embryos. *Science, 334*(6059), 1141-1144.

Scolari, F., Benoit, J. B., Michalkova, V., Aksoy, E., Takac, P., Abd-Alla, A. M., . . . Attardo, G. M. (2016). The spermatophore in *Glossina morsitans morsitans*: insights into male montributions to reproduction. *Scientific reports, 6,* 20334.

Shim, J., Gururaja-Rao, S., & Banerjee, U. (2013). Nutritional regulation of stem and progenitor cells in *Drosophila*. *Development, 140*(23), 4647-4656.

Shutt, B., Stables, L., Aboagye-Antwi, F., Moran, J., & Tripet, F. (2010). Male accessory gland proteins induce female monogamy in anopheline mosquitoes. *Medical and veterinary entomology, 24*(1), 91-94.

Sibley, P. K., Ankley, G. T., & Benoit, D. A. (2001). Factors affecting reproduction and the importance of adult size on reproductive output of the midge *Chironomus tentans*. *Environmental Toxicology and Chemistry: An International Journal, 20*(6), 1296-1303.

Silva, J. V., Yoon, S., Domingues, S., Guimarães, S., Goltsev, A. V., e Silva, E. F. d. C., . . . Fardilha, M. (2015). Amyloid precursor protein interaction network in human testis: sentinel proteins for male reproduction. *BMC bioinformatics, 16*(1), 12.

Sirot, L. K., Wong, A., Chapman, T., & Wolfner, M. F. (2015). Sexual conflict and seminal fluid proteins: a dynamic landscape of sexual interactions. *Cold Spring Harbor perspectives in biology, 7*(2), a017533.

Soller, M., Bownes, M., & Kubli, E. (1999). Control of oocyte maturation in sexually mature *Drosophila* females. *Developmental biology, 208*(2), 337-351. doi:10.1006/dbio.1999.9210

Sosic, D., & Olson, E. N. (2003). A new twist on twist: Modulation of the NF-KappaB pathway. *Cell Cycle, 2*(2), 75-77.

Strong, J. (1967). Ecology of terrestrial arthropods at Palmer station, Antarctic Peninsula. *Entomology of Antarctica*, 357-371.

Stroumbakis, N. D., Li, Z., & Tolias, P. P. (1996). A homolog of human transcription factor NF-X1 encoded by the *Drosophila* shuttle craft gene is required in the embryonic central nervous system. *Molecular and cellular biology, 16*(1), 192-201.

Sugg, P., Edwards, J. S., & Baust, J. (1983). Phenology and life history of *Belgica antarctica*, an Antarctic midge (Diptera: Chironomidae). *Ecological Entomology, 8*(1), 105-113.

Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One, 6*(7), e21800.

Swanson, W. J., Wong, A., Wolfner, M. F., & Aquadro, C. F. (2004). Evolutionary expressed sequence tag analysis of *Drosophila* female reproductive tracts identifies genes subjected to positive selection. *Genetics, 168*(3), 1457-1465.

Swevers, L., Raikhel, A., Sappington, T., Shirk, P., & Iatrou, K. (2005). Vitellogenesis and post-vitellogenic maturation of the insect ovarian follicle. In L.I. Gilbert, S. Gill, & K. Iatrou (Eds.), *Comprehensive molecular insect science* (pp. 87-155). Elsevier, NY

Teets, N. M., Elnitsky, M. A., Benoit, J. B., Lopez-Martinez, G., Denlinger, D. L., & Lee, R. E. (2008). Rapid cold-hardening in larvae of the Antarctic midge *Belgica antarctica*: cellular cold-sensing and a role for calcium. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology, 294*(6), R1938-R1946. doi:DOI 10.1152/ajpregu.00459.2007

Teets, N. M., Kawarasaki, Y., Lee, R. E., Jr., & Denlinger, D. L. (2013). Expression of genes involved in energy mobilization and osmoprotectant synthesis during thermal and dehydration stress in the Antarctic midge, *Belgica antarctica*. *Journal of Comparative Physiology B, 183*(2), 189-201. doi:10.1007/s00360-012-0707-2

Teets, N. M., Peyton, J. T., Colinet, H., Renault, D., Kelley, J. L., Kawarasaki, Y., . . . Denlinger, D. L. (2012). Gene expression changes governing extreme dehydration tolerance in an Antarctic insect. *Proceedings of the National Academy of Sciences of the United States of America, 109*(50), 20744-20749. doi:10.1073/pnas.1218661109

Telfer, W. H., & Kunkel, J. G. (1991). The function and evolution of insect storage hexamers. *Annual Review of Entomology, 36*(1), 205-228.

Terashima, J., & Bownes, M. (2005). A microarray analysis of genes involved in relating egg production to nutritional intake in *Drosophila melanogaster*. *Cell Death Differ, 12*(5), 429.

Thailayil, J., Magnusson, K., Godfray, H. C. J., Crisanti, A., & Catteruccia, F. (2011). Spermless males elicit large-scale female responses to mating in the malaria mosquito *Anopheles gambiae*. *Proceedings of the National Academy of Sciences, 108*(33), 13677-13681.

1441   Tian, C.-B., Wei, D., Xiao, L.-F., Dou, W., Liu, H., & Wang, J.-J. (2017). Comparative transcriptome analysis
1442   of three *Bactrocera dorsalis* (Diptera: Tephritidae) organs to identify functional genes in the
1443   male accessory glands and ejaculatory duct. *Florida Entomologist, 100*(1), 42-51.
1444   Tian, E., & Hagen, K. G. T. (2007). O-linked glycan expression during *Drosophila* development.
1445   *Glycobiology, 17*(8), 820-827.
1446   Tirmarche, S., Kimura, S., Dubruille, R., Horard, B., & Loppin, B. (2016). Unlocking sperm chromatin at
1447   fertilization requires a dedicated egg thioredoxin in *Drosophila*. *Nature communications, 7*,
1448   13539.
1449   Tran, D. T., & Ten Hagen, K. G. (2013). Mucin-type O-glycosylation during development. *Journal of
1450   Biological Chemistry, 288*(10), 6921-6929.
1451   Tran, D. T., Zhang, L., Zhang, Y., Tian, E., Earl, L. A., & Ten Hagen, K. G. (2012). Multiple members of the
1452   UDP-GalNAc: polypeptide N-acetylgalactosaminyltransferase family are essential for viability in
1453   *Drosophila*. *Journal of Biological Chemistry, 287*(8), 5243-5252.
1454   Turnier, J. L., Brunner, H. I., Bennett, M., Aleed, A., Gulati, G., Haffey, W. D., . . . Witte, D. (2018).
1455   Discovery of SERPINA3 as a candidate urinary biomarker of lupus nephritis activity.
1456   *Rheumatology, 58*(2), 321-330.
1457   Usher, M. B., & Edwards, M. (1984). A dipteran from south of the Antarctic Circle: *Belgica antarctica*
1458   (Chironomidae) with a description of its larva. *Biological Journal of the Linnean Society, 23*(1),
1459   19-31.
1460   Vanderperre, B., Cermakova, K., Escoffier, J., Kaba, M., Bender, T., Nef, S., & Martinou, J.-C. (2016).
1461   MPC1-like is a placental mammal-specific mitochondrial pyruvate carrier subunit expressed in
1462   postmeiotic male germ cells. *Journal of Biological Chemistry, 291*(32), 16448-16461.
1463   Venancio, T., Cristofoletti, P., Ferreira, C., Verjovski-Almeida, S., & Terra, W. (2009). The *Aedes aegypti*
1464   larval transcriptome: a comparative perspective with emphasis on trypsins and the domain
1465   structure of peritrophins. *Insect molecular biology, 18*(1), 33-44.
1466   Villarreal, S. M., Pitcher, S., Helinski, M. E., Johnson, L., Wolfner, M. F., & Harrington, L. C. (2018). Male
1467   contributions during mating increase female survival in the disease vector mosquito *Aedes
1468   aegypti*. *Journal of insect physiology, 108*, 1-9.
1469   Vogt, C., Belz, D., Galluba, S., Nowak, C., Oetken, M., & Oehlmann, J. (2007). Effects of cadmium and
1470   tributyltin on development and reproduction of the non-biting midge *Chironomus riparius*
1471   (Diptera)—baseline experiments for future multi-generation studies. *Journal of Environmental
1472   Science and Health Part A, 42*(1), 1-9.
1473   Weirauch, M. T., & Hughes, T. (2011). A catalogue of eukaryotic transcription factor types, their
1474   evolutionary origin, and species distribution. In *A handbook of transcription factors* (pp. 25-73):
1475   Springer.
1476   Weirauch, M. T., Yang, A., Albu, M., Cote, A. G., Montenegro-Montero, A., Drewe, P., . . . Cook, K. (2014).
1477   Determination and inference of eukaryotic transcription factor sequence specificity. *Cell, 158*(6),
1478   1431-1443.
1479   Wensler, R. J., & Rempel, J. (1962). The morphology of the male and female reproductive systems of the
1480   midge, *Chironomus plumosus* L. *Canadian Journal of Zoology, 40*(2), 199-229.
1481   Xie, T., & Spradling, A. C. (1998). Decapentaplegic is essential for the maintenance and division of
1482   germline stem cells in the *Drosophila* ovary. *Cell, 94*(2), 251-260.
1483   Zhang, J., Marshall, K. E., Westwood, J. T., Clark, M. S., & Sinclair, B. J. (2011). Divergent transcriptomic
1484   responses to repeated and single cold exposures in *Drosophila melanogaster*. *Journal of
1485   Experimental Biology, 214*(Pt 23), 4021-4029. doi:10.1242/jeb.059535

1486 Zhang, L., Zhang, Y., & Ten Hagen, K. G. (2008). A mucin-type O-glycosyltransferase modulates cell
1487   adhesion during *Drosophila* development. *Journal of Biological Chemistry, 283*(49), 34076-
1488   34086.

1489

1490

1491

Figure 3

A.

B.

C.

A.

B.

Figure 5

Figure 6

Figure 7

A. Female

B. Male

C. Larvae

Figure 9

A. Female accessory gland

B. Male accessory glands

C. Male accessory glands enriched genes of mosquitoes

Figure 10

Figure 12

A.

B.

C.

Figure 13

A. Daily thermal changes

B. Exposure to 20°C

C. Maximum temperate shift

D. Rate of temperate change

Figure 1&

Figure 15

Mating, Fertilization and Seminal Fluid Transfer

Transfer of Male
Derived Gel Proteins

Female Accessory
Gland Derived Gel
Proteins

Spermatophore transfer
Multiple matings

Dehydration negativly
impacts male fertility

Cryoprotection and dehydration
alters yolk protein gene expression

**Egg Deposition and
Gel Matrix Secretion**

**Larval
Development**

Embryogenesis in protective gel

Gel Matrix
Prevents rapid egg dehydration
Functions as a thermal buffer
Nutritional resource for developing larvae