1

An experimentally-derived measure of inter-replicate variation in reference samples:

the same-same permutation methodology

David C. Handler and Paul A. Haynes*

Department of Molecular Sciences, Macquarie University, North Ryde, NSW 2109,

Australia

Email - david.handler@mq.edu.au

Email - paul.haynes@mq.edu.au

* Correspondence: Professor Paul A. Haynes, Department of Molecular Sciences,

Macquarie University, North Ryde, NSW 2109, Australia

Phone: 61-2-9850 6258

Fax: 61-2-9850 6200

Email - paul.haynes@mq.edu.au

Updated with clarifications 17th may 2021.

Abstract

The multiple testing problem is a well-known statistical stumbling block in high-

throughput data analysis, where large scale repetition of statistical methods introduces

unwanted noise into the results. While approaches exist to overcome the multiple testing

problem, these methods focus on theoretical statistical clarification rather than incorporating

experimentally-derived measures to ensure appropriately tailored analysis parameters. Here,

we introduce a method for estimating inter-replicate variability in reference samples for a

quantitative proteomics experiment using permutation analysis. This can function as a

modulator to multiple testing corrections such as the Benjamini-Hochberg ordered Q value

test. We refer to this as a 'same-same' analysis, since this method incorporates the use of six

biological replicates of the reference sample and determines, through non-redundant triplet

pairwise comparisons, the level of quantitative noise inherent within the system. The method

can be used to produce an experiment-specific Q value cut-off that achieves a specified false

discovery rate at the quantitation level, such as 1%. The same-same method is applicable to

any experimental set that incorporates six replicates of a reference sample. To facilitate

access to this approach, we have developed a same-same analysis R module that is freely

available and ready to use via the internet.

Keywords: Label-free shotgun proteomics, false discovery rates, data quality, data
validation, statistics

48
49 **1. Introduction**
50
Shotgun proteomics experiments that seek to compare 'reference' and 'treated' states of

a given sample will often contain thousands of individual comparisons, each requiring

statistical validation. In such circumstances, repeated use of the Student's t-test will

invariably introduce false discoveries into the results. A Student's t-test with a significance

cut-off threshold P value of 0.05 produces 95% confidence i.e. 5% of tests will have an equal

likelihood to be attributable to chance instead of experimental factors [1-4]. This is a follow-

on conclusion from the probability that at least one test in an experiment will be significant,

which is described as $1 – (1 - \alpha)^k$, where α is the significance cut-off and k is the number of

tests conducted [5].

Multiple Testing Corrections (MTCs) were introduced to help address this limitation [6],

including the use of Q values rather than P values [7], Bonferroni correction [8,9], Benjamini-

Hochberg (BH) adjusted t-test [10], Bonferroni-Holm test [11], and the Benjamini-Yoav

(BY) test [12]. MTCs, however, are often overly conservative and increase the false negative

rate by eliminating otherwise valid protein identifications. This is especially a problem at the

protein-quantitation level; MTCs, by their nature, contribute to a lessening of the protein

quantitation false discovery rate (PQ-FDR) at the expense of otherwise valid protein

identifications [13,14]. In the context of this article, PQ-FDR is defined as false discoveries

arising from comparative quantitative proteomics calculations between one or more samples.

There is always a balance to be struck between stringency and accuracy when controlling

false discoveries at the protein quantitation level [15]. A recent study in this area applied

Bayesian statistics to great effect, detecting a greater number of relevant protein quantitation

changes in previously published data sets [16]. There are also numerous software packages

available which incorporate various other MTC approaches, including Proteus [17], DAPAR

and ProStaR [18], MSqRob [19], UbiA-MS [20], ProteoSign [21], msVolcano [22], FDRtool

[23], MSstats [24], and limma [25].

Although the use of MTC correction methods in the proteomics field is not standardized

[9,26-28], MTCs are an important tool that researchers can employ for extracting the best

results from their dataset i.e. finding the balance between reducing noise without losing

signal. This desire to reduce the noise in the system led us to ask the question: is there a better

way to quantify variability between replicate analyses of a reference sample ?

One established approach for assessing variability across a sample set is to use

permutation analysis, based upon the Significance Analysis of Micro-array (SAM)

permutation methodology [29,30]. This is a similar theoretical framework to that used for

permutation analysis within Perseus [31], a well-established data analysis program in the

MaxQuant environment [32]. The SAM permutation analysis method assigns a score to each

gene on the basis of change in gene expression relative to the standard deviation of repeated

measurements. SAM then uses redundant permutations of repeated measurements to estimate

the percentage of genes identified by random chance as an artefact of the method, which is

used to calculate the false discovery rate. The permutations are performed across all of the

'reference' and 'treated' sample replicates within a given experimental data set. Those genes

with scores higher than the specified threshold are deemed potentially significant, and the

threshold can be adjusted to identify smaller or larger sets of genes, with FDR calculated for

each set.

The same – same method introduced in this study, in contrast, employs non-redundant

permutations of experimentally repeated measurements of protein abundance in replicate

analysis of a defined reference sample. The permutations are performed on data from the

reference samples only, isolated from the 'treated' samples. This is used to generate a single

average Q value indicative of the degree of variation of abundance across the reference

sample replicates. Proteins which reach a defined statistical significance threshold are

deemed to be false discoveries at the protein quantitation level, since comparing a reference

sample against itself should theoretically yield no changes in protein abundance. It is

important to emphasize that the underlying assumption is that the biological variability

between reference samples is zero, so this approach is accounting for the technical variability.

This facilitates subsequent assessment of induced biological variation between reference

samples and treated samples.

A specified false discovery rate in the same – same analysis of replicates of the reference

sample is used to generate a Q value threshold, and that value can then be carried forward to

the subsequent analysis of a reference sample versus 'treated' sample within the same larger

experimental data set. One of main the applications of this method for determining an

experimentally-derived measure of reference sample variability is that it can subsequently be

used to modify an existing MTC protocol for downstream analysis, thus minimising the PQ-

FDR without introducing false negatives. By performing a specific permutation analysis to

measure the variability inherent within reference sample replicates, we can produce an

experimentally modulated Q value threshold for use with MTCs when comparing the

reference sample to treated samples. In essence, rather than using a default Q value of .05, or

choosing a more stringent value, we are employing a Q value threshold that is experimentally

determined for each set of samples analyzed. The same-same method represents another tool

in the proteomics toolbox, and can be used to enable the extraction of additional biological

knowledge from large-scale datasets.

120
121   **2. Materials and Methods**
122
2.1 Label free quantitative proteomics data sets

To demonstrate the utility of the same-same approach we reanalyzed two sets of

previously published label free quantitative shotgun proteomics data. Protein identification

and Normalized Spectral Abundance Factor (NSAF) values [33,34] were sourced from

previously published studies from our laboratory on cultured Cabernet Sauvignon grape cells

grown at different temperatures [35] (ProteomeXchange identifier PXD000977) and leaf

tissue of IAC1131 rice plants exposed to drought stress [36] (ProteomeXchange identifier

PXD004096). The cultured Cabernet Sauvignon grape cell data consists of six biological

replicates of cell cultures maintained at 26°C, as the optimum, or control, temperature, and

biological triplicate analysis of cells maintained at 18°C and 10°C as moderate and extreme

cold stress conditions, and 34°C and 42°C as moderate and extreme heat stress conditions.

The rice leaf data consists of two sets of three biological replicates each of unstressed plants

as controls, and biological triplicate analysis of plants exposed to moderate drought stress,

extreme drought stress, and extreme drought stress followed by recovery.

2.2 Same – same permutation analysis of reference samples

For analysis using the same-same workflow, six replicates of a reference sample are run

through a PSM (peptide-to-spectrum matching) engine such as GPM or ProteomeDiscoverer

and protein identification lists are exported as csv files. At a minimum, protein identifier and

peptide count are needed. Next, these six replicates are grouped into two sets of triplicates

by the use of inner joins (dummy state 'control', dummy state 'treatment'), and a test array

is formed through a full join of the states [37]. One hundred Student's t-tests are conducted

on spectral counts from each identified protein comparison with significance cut-off values

from 0.01 to 1, stepped at 0.01 intervals. All proteins found at different quantitation levels

are considered false discoveries, since comparison between two data sets of the same sample

type would theoretically give identical quantitation with no observed changes. This process

is repeated for all ten combinations of non-redundant triplet pairs that six replicates can form.

The MTC analysis then begins by iterating over this 10x t-test array and applying one of five

user-specified MTC methods (BH, Benjamini-Yoav, Bonferroni, Hommel [38], and

Bioconductor Q [39]. The program then averages the MTC test results from all arrays

examined, and reports the point at which the significance cut-off corresponds to a user-

specified PQ-FDR.

The same-same methodology is automated through an R script. Source code is available

from https://bitbucket.org/peptidewitch/samesame/, and a freely accessible working web

version can be found at https://peptidewitch.shinyapps.io/samesame. The R Shiny web-app

provides three distinct outputs from the same-same analysis:

1) A series of Q value vs FDR bar plots (x axis 0.01 to 1, stepped at 0.01) from all ten

159   triplet paired combinations,

160   2) A series of P value histograms of these same combinations, and,

161   3) A numerical value that corresponds to the user-specified MTC cut-off that produces

162   the desired PQ-FDR (default 1%).

Input data types are not constrained to spectral counts, as in theory any data type that

consists of protein identifications coupled with abundance or intensity value measurements

can be used. However, the first generation of the analysis tool was designed and tested using

spectral counting-based data, so it is recommended that spectral counts or spectral abundance

factors be used initially.

2.3 Perseus permutation analysis of reference samples

To serve as a comparison against the same-same process, the same NSAF data from both

Grape and Rice samples as were reanalyzed using Perseus software [31]. Spreadsheet files

containing NSAF values for each set of samples were uploaded to Perseus through a generic

matrix upload. Using the two-sample module, we applied the Perseus permutation method as

a form of truncation using ungrouped (no grouping preserved), 250-count permutation

analysis on two-tailed Student's t-testing arrays with BH correction, comparing six reference

replicates with three replicates from each of the 'treated' sample states, with the specified

FDR thresholds ranging from 1-5%.

3. Results

The following section details how same-same approach was applied to the published data

from Grape and Rice cells. Figure 1 displays the outputs described above for the same-same

analysis conducted on the grape cell culture label-free data when specifying BH correction

and 1% PQ-FDR. The end-point of the same-same process is the modulated Q value, in this

example 0.054 (Figure 1C), produced from averaging the threshold values in Figure 1A at

the desired PQ-FDR value. This value can be used for downstream analysis on subsequent

control vs treatment samples as a modulator for the chosen MTC.

Figure 2 presents the subsequent downstream analysis of the grape cell cultures grown at

different temperatures. Figure 2A shows the number of proteins found to be significantly

differentially expressed in terms of protein fold change when comparing the set of six

reference replicates to the set of three replicates of cells grown at each temperature. These

are analysed using different statistical measures of significance: P values of 0.05 and 0.01 for

a student's t-test, BH Q value of 0.05, and BH using the same-same derived Q value (SS-Q),

and specifying PQ – FDR of 1%, 2% or 3%. It is evident that the same-same derived Q values

at 1% PQ-FDR produce results very similar to the use of default BH Q values, which is

expected given that the SS-Q value used is very close to the 0.05 BH-Q value threshold. The

two approaches give similar results, although it is noticeable that at a specified PQ – FDR of

3%, the comparison with the largest effect size (Figure 2E) shows significantly more

differentially expressed proteins.

Figure 1

202
203

Figure 2

Figure 3 presents the same type of analyses as shown in Figure 1 for the data derived

from comparative analysis of leaf tissue from IAC1131 rice plants exposed to different levels

of drought stress. Interestingly, in contrast to figure 2, it is clear that in this case there is a

direct correlation between observed effect size and number of differentially expressed

proteins identified using the SS-Q approach. In comparisons with greater effect size as

observed in P value histograms (Figure 3E,3F,3G), the same-same derived Q Values are able

to identify a greater number of differentially expressed proteins than were found using the

default BH Q value, and at 3% PQ-FDR are approaching the number of differentially

expressed proteins found using uncorrected P values.

Table 1 presents the results of analyzing the grape and rice cell NSAF data referred to

above using different analysis approaches, including Student t-tests with and without BH

correction, application of same – same derived Q values to a BH corrected t-test at specified

PQ-FDR values ranging from 1% to 5%, and t-tests using Perseus permutations at specified

PQ-FDR values ranging from 1% to 5%. The table shows the number of proteins which are

reported to be significantly differentially expressed when comparing the reference samples

against the grape cells grown at four different temperatures, and the rice cells grown under

three different watering regimes. It is clear from these comparisons that, as expected, the

uncorrected student's t-test gives a much greater number than any sort of correction. The BH

correction reduces the number of significant proteins by approximately 95%. The Perseus

permutation processing is even more strict and, for example, produces zero significant

identifiers in more than half of the grape sample comparisons. In contrast, the same-same-

modulated BH test is able to detect significantly differentially regulated proteins for every

test case for both tissue types while always remaining well below the results reported from

uncorrected Student's t-testing P values. Multiple testing correction still takes place, but the

experimentally derived Q value thresholds allow for the recovery of a greater number of

significant differences at the protein quantitation level.

Table 1 – Comparison of number of protein identifications retained using different

analysis approaches to assess protein quantitation false discovery rate

Comparisons	t-test[a]		Perseus permutations BH					Same-Same BH				
Grape	**.05**	**BH**	**1%**[b]	**2%**	**3%**	**4%**	**5%**	**1%**	**2%**	**3%**	**4%**	**5%**
Cont vs 10 C	735	42	0	0	0	0	0	48	54	72	91	99
Cont vs 18 C	560	29	0	0	0	9	9	29	34	43	51	65
Cont vs 34 C	432	21	0	0	0	0	6	24	28	36	42	48
Cont vs 42 C	775	83	0	0	0	0	44	108	116	133	166	199
Rice	**.05**	**BH**	**1%**	**2%**	**3%**	**4%**	**5%**	**1%**	**2%**	**3%**	**4%**	**5%**
Cont vs Ext	543	12	0	0	0	0	0	15	16	17	17	17
Cont vs Mod	481	13	0	0	0	0	0	13	17	20	20	20
Cont vs Recov	568	21	0	0	0	0	0	22	33	40	40	40

238   [a] .05 = standard 2-sample t-test, BH = Benjamin-Hochberg corrected 2 sample t-test

239   [b] protein quantitation false discovery rate assessed at 1%-5% using the approaches indicated

243   **Figure 3**

244
245

246
247   **4. Discussion**
248

The correlation observed between effect size and number of differentially expressed

proteins found in the dataset presented in Figure 3 has also been found in numerous other

datasets we have analysed. In general, SS-Q values are generally better suited to those

datasets that show a larger effect size. This may be due to the fact that not all quantitatively

different proteins in a small effect sample are false positives, or may be a consequence of

NSAFs overstating expression change ratios for protein identifications based on lower

spectral counts, which can help to increase the effect size [4]. While the use of higher Q value

thresholds raises the implicit question of whether or not the dataset contains too much noise,

it is important to remember why the same-same experiment is conducted in the first place. If,

in an experiment where we expect there to be minimal noise, we demonstrate that there is a

SS-Q threshold value that produces 1% PQ-FDR between sets of control or reference

replicates, then in a closely related experiment with the same reference sample using the same

threshold value, we can infer experimentally that the specified PQ-FDR has been achieved.

It is important to stress, however, that this method is suited more towards initial

discovery, and that follow-up experimentation must employ orthogonal validation protocols.

In order to obtain an experimentally-derived PQ-FDR of 1%, or other specified value, the

same-same method is a very useful tool, because inferring the PQ-FDR based on the Q value

cut-off alone does not yield corresponding PQ-FDR levels (i.e. a Q value of 0.05 does not

specifically produce either 5% or 1% PQ-FDR). Modifying the MTC significance value cut-

off so that it takes into account the experimental variability inherent within the replicates

helps to produce a more tailored list of differentially expressed protein identifications whilst

controlling for PQ-FDR. Also, compounding the same-same technique with another method

of filtering, such as fold change cut-offs, can reduce the number of false positives included

in the final dataset, further reducing the PQ-FDR [30,31].

In this research article, we have demonstrated a revised method for statistical analysis for

shotgun proteomics datasets. The same-same method facilitates the construction of post

analysis P value histograms and aids the researcher in choosing an appropriate statistical

testing protocol for their analysis. We have shown that in the right circumstances, using BH

Q value cut-offs derived from the same-same analysis yields a set of results that provide more

significantly differentially expressed proteins from a given dataset, while also determining

PQ-FDR at the experimental level. In the future, we hope to expand on this methodology so

that it can be applied equally well to other quantitative proteomics data types, and also

develop new tests to build onto the existing same-same architecture to further improve the

statistical rigour for all shotgun proteomics results.

----------||-----------

5. Acknowledgments

287   The authors acknowledge support from BioPlatforms Australia through the Australian

288   Government's National Collaborative Research Infrastructure Scheme, and the Australian

289   Research Council through Discovery Project DP190103140. The authors declare no conflict

290   of interest.

6. Author contributions

293   DCH and PAH contributed equally to the conceptualization and design of this study.

294   DCH was responsible for writing the software described. DCH prepared the original draft

295   of the manuscript, and PAH was responsible for revision and editing.

7. Figure Legends

Figure 1. Screenshots from the same/same shiny apps module (https://peptidewitch.shinyapps.io/SameSame), using the grape cell control samples and specifying BH correction at 1% PQ-FDR. (A) Q value vs PQ-FDR bar plots (x axis 0.01 to 1, stepped at 0.01) for all ten triplet paired permutations generated from six replicate analyses of a reference sample (see Figure 1) (B) P value histograms for each permutation, showing number of significantly expressed protein identifications sorted into P values bins in increments of 0.05. (C) displays a single numerical value which produces the desired PQ-FDR value (default BH at 1%, can be user specified).

Figure 2. Grape cell culture comparisons with application of different statistical significance measures. Cells grown at 26°C were designated as the reference sample, and compared with cells grown at 18°C (moderate cold), 10°C (extreme cold), 34°C (moderate

heat), and 42°C (extreme heat). Panel A displays the number of significantly differentially

expressed protein identifications found for each comparison using P values at .05 and .01,

Benjamini-Hochberg adjusted values at 0.05, and BH using the same-same derived Q value

(SS-Q), and specifying PQ – FDR of 1%, 2% or 3%. Panels B to E contain P value histograms

showing the number of significantly expressed protein identifications sorted into P value bins

in increments of 0.05, for each of the four experimental comparisons performed, as indicated.

Figure 3. IAC1131 rice samples drought stress comparisons with application of different

statistical significance measures. Control plants were unstressed, and compared with plants

exposed to moderate drought stress, extreme drought stress, or extreme drought stress

followed by recovery. Panel A displays the number of significantly differentially expressed

protein identifications found for each comparison using P values at .05 and .01, Benjamini-

Hochberg adjusted values at 0.05, and BH using the same-same derived Q value (SS-Q), and

specifying PQ – FDR of 1%, 2% or 3%. Panels B to G are P value histograms showing the

number of significantly expressed protein identifications sorted into P value bins in

increments of 0.05, for each of the six experimental comparisons performed, as indicated.

330
8. References

1.   Colquhoun, D. An investigation of the false discovery rate and the misinterpretation
of p-values. *R Soc Open Sci* **2014**, *1*, 140216.
2.   Diz, A.P.; Carvajal-Rodriguez, A.; Skibinski, D.O. Multiple hypothesis testing in
proteomics: A strategy for experimental work. *Mol Cell Proteomics* **2011**, *10*, M110
004374.
3.   Karp, N.A.; Lilley, K.S. Design and analysis issues in quantitative proteomics studies.
Proteomics **2007**, *7* 42-50.
4.   Sullivan, G.M.; Feinn, R. Using effect size-or why the p value is not enough. *J Grad*
Med Educ **2012**, *4*, 279-282.
5.   Streiner, D.L.; Norman, G.R. Correction for multiple testing: Is there a resolution?
Chest **2011**, *140*, 16-18.
6.   Bender, R.; Lange, S. Adjusting for multiple testing--when and how? *J Clin*
Epidemiol **2001**, *54*, 343-349.
7.   Armstrong, R.A. When to use the bonferroni correction. *Ophthalmic Physiol Opt*
2014, *34*, 502-508.
8.   Choi, H.; Nesvizhskii, A.I. False discovery rates and related statistical concepts in
mass spectrometry-based proteomics. *J Proteome Res* **2008**, *7*, 47-50.
9.   Benjamini, Y.; Hochberg, Y. Controlling the false discovery rate: A practical and
powerful approach to multiple testing. *J Royal Stat Society, Series B* **1995**, 289-300.
10.   Holm, S. A simple sequentially rejective multiple test procedure. *Scandinavian*
journal of statistics **1979**, *6*, 65-70.
11.   Benjamini, Y.; Yekutieli, D. The control of the false discovery rate in multiple testing
under dependency. *Annals of statistics* **2001**, *29*, 1165-1188.
12.   Carvajal-Rodriguez, A.; de Una-Alvarez, J. Assessing significance in high-
throughput experiments by sequential goodness of fit and q-value estimation. *PLoS*
One **2011**, *6*, e24700.
13.   Pascovici, D.; Handler, D.C.; Wu, J.X.; Haynes, P.A. Multiple testing corrections in
quantitative proteomics: A useful but blunt tool. *Proteomics* **2016**, *16*, 2448-2453.
14.   Noble, W.S. How does multiple testing correction work? *Nat Biotechnol* **2009**, *27*,
1135-1137.
15.   The, M.; Kall, L. Integrated identification and quantification error probabilities for
shotgun proteomics. *Mol Cell Proteomics* **2019**, *18*, 561-570.
16.   Gierlinski, M.; Gastaldello, F.; Barton, G.J. Proteus: An r package for downstream
analysis of maxquant output. *BioRxiv* **2018**, doi.org/doi10.1101/416511.
17.   Wieczorek, S.; Combes, F.; Lazar, C.; Gianetto, Q.G.; Gatto, L.; Dorffer, A.; Hesse,
A.M.; Coute, Y.; Ferro, M.; Bruley, C.*, et al.* Dapar & prostar: Software to perform
statistical analyses in quantitative discovery proteomics. *Bioinformatics* **2017**, *33*,
135-136.
18.   Goeminne, L.J.E.; Gevaert, K.; Clement, L. Experimental design and data-analysis in
label-free quantitative lc/ms proteomics: A tutorial with msqrob. *J Proteomics* **2018**,
171, 23-36.

19.   Zhang, X.; Smits, A.H.; van Tilburg, G.B.; Ovaa, H.; Huber, W.; Vermeulen, M.
Proteome-wide identification of ubiquitin interactions using ubia-ms. *Nat Protoc*
2018, *13*, 530-550.

20.   Efstathiou, G.; Antonakis, A.N.; Pavlopoulos, G.A.; Theodosiou, T.; Divanach, P.;
Trudgian, D.C.; Thomas, B.; Papanikolaou, N.; Aivaliotis, M.; Acuto, O.*, et al.*
Proteosign: An end-user online differential proteomics statistical analysis platform.
Nucleic Acids Res **2017**, *45*, W300-W306.

21.   Singh, S.; Hein, M.Y.; Stewart, A.F. Msvolcano: A flexible web application for
visualizing quantitative proteomics data. *Proteomics* **2016**, *16*, 2491-2494.

22.   Strimmer, K. A unified approach to false discovery rate estimation. *BMC*
Bioinformatics **2008**, *9*, 303.

23.   Choi, M.; Chang, C.Y.; Clough, T.; Broudy, D.; Killeen, T.; MacLean, B.; Vitek, O.
Msstats: An r package for statistical analysis of quantitative mass spectrometry-based
proteomic experiments. *Bioinformatics* **2014**, *30*, 2524-2526.

24.   Ritchie, M.E.; Phipson, B.; Wu, D.; Hu, Y.; Law, C.W.; Shi, W.; Smyth, G.K. Limma
powers differential expression analyses for rna-sequencing and microarray studies.
Nucleic Acids Res **2015**, *43*, e47.

25.   Gelman, A.; Hill, J.; Yajima, M. Why we (usually) don't have to worry about multiple
comparisons. *J Res Educ Effect* **2012**, *5*, 189-211.

26.   Kall, L.; Storey, J.D.; MacCoss, M.J.; Noble, W.S. Posterior error probabilities and
false discovery rates: Two sides of the same coin. *J Proteome Res* **2008**, *7*, 40-44.

27.   Rothman, K.J. No adjustments are needed for multiple comparisons. *Epidemiology*
1990, *1*, 43-46.

28.   Gierlinski, M.; Cole, C.; Schofield, P.; Schurch, N.J.; Sherstnev, A.; Singh, V.;
Wrobel, N.; Gharbi, K.; Simpson, G.; Owen-Hughes, T.*, et al.* Statistical models for
rna-seq data derived from a two-condition 48-replicate experiment. *Bioinformatics*
2015, *31*, 3625-3630.

29.   Tusher, V.G.; Tibshirani, R.; Chu, G. Significance analysis of microarrays applied to
the ionizing radiation response. *Proc Natl Acad Sci U S A* **2001**, *98*, 5116-5121.

30.   Tyanova, S.; Temu, T.; Sinitcyn, P.; Carlson, A.; Hein, M.Y.; Geiger, T.; Mann, M.;
Cox, J. The perseus computational platform for comprehensive analysis of
(prote)omics data. *Nat Methods* **2016**, *13*, 731-740.

31.   Tyanova, S.; Temu, T.; Cox, J. The maxquant computational platform for mass
spectrometry-based shotgun proteomics. *Nat Protoc* **2016**, *11*, 2301-2319.

32.   Neilson, K.A.; Ali, N.A.; Muralidharan, S.; Mirzaei, M.; Mariani, M.; Assadourian,
G.; Lee, A.; van Sluyter, S.C.; Haynes, P.A. Less label, more free: Approaches in
label-free quantitative mass spectrometry. *Proteomics* **2011**, *11*, 535-553.

33.   Neilson, K.A.; Keighley, T.; Pascovici, D.; Cooke, B.; Haynes, P.A. Label-free
quantitative shotgun proteomics using normalized spectral abundance factors.
Methods Mol Biol **2013**, *1002*, 205-222.

34.   George, I.S.; Pascovici, D.; Mirzaei, M.; Haynes, P.A. Quantitative proteomic
analysis of cabernet sauvignon grape cells exposed to thermal stresses reveals

alterations in sugar and phenylpropanoid metabolism. *Proteomics* **2015**, *15*, 3048-
3060.
35. Wu, Y.; Mirzaei, M.; Pascovici, D.; Chick, J.M.; Atwell, B.J.; Haynes, P.A.
Quantitative proteomic analysis of two different rice varieties reveals that drought
tolerance is correlated with reduced abundance of photosynthetic machinery and
increased abundance of clpd1 protease. *J Proteomics* **2016**, *143*, 73-82.
36. Mishra, P.; Eich, M. Join processing in relational databases. *CM Computing Surveys*
1992, *24*, 63-113.
37. Hommel, G.; Krummenauer, F. Improvements and modifications of tarone's multiple
test procedure for discrete data. *Biometrics* **1998**, *54*, 673-681.
38. Qian, H.R.; Huang, S. Comparison of false discovery rate methods in identifying
genes with differential expression. *Genomics* **2005**, *86*, 495-503.

427