Landscape of Dysregulated Placental RNA Editing Associated with Preeclampsia

Xiaoxue Yang[1], Jing Yang[1,2,3], Xiaozhen Liang[1], Qian Chen[1], Sijia Jiang[1], Haihua Liu[1], Yue Gao[1,2,3], Zhonglu Ren[1], Yanhong Yu[1], Mei Zhong[1*] and Xinping Yang[1,2,3*]

Author Affiliations

[1]Department of Obstetrics and Gynecology; Nanfang Hospital, Southern Medical University; Guangzhou, Guangdong, 510515; China.

[2]Key Laboratory of Mental Health of the Ministry of Education; Southern Medical University; Guangzhou, Guangdong, 510515; China.

[3]Department of Bioinformatics; School of Basic Medical Sciences, Southern Medical University; Guangzhou, Guangdong, 510515; China

Corresponding authors: xpyang1@smu.edu.cn, zhongmei@fimmu.com

Keywords: RNA editing, preeclampsia, differentially edited site, *LEP*, *miR-9-5p*

Abstract

Dysregulated RNA editing is well documented in several diseases such as cancer. The extent to which RNA editing might be involved in diseases originated in the placenta remains unknown, because RNA editing has rarely been studied in the placenta. Here, we have systematically profiled RNA editome on the placentae from 9 patients with early-onset severe preeclampsia (EOSPE) and 32 normal subjects, and a widespread RNA editing dysregulation in EOSPE has been identified. The mis-edited gene set is enriched with known preeclampsia-associated genes and differentially expressed genes in EOSPE. The "RNA editing events" at two microRNA binding sites in 3'-UTR of the *LEP* mRNA were generated, which could lead to increased expression level of *LEP* in trophoblast cells. Upregulation of *LEP* were also observed in the placentae of PE patients. These results suggest that widespread placental RNA editing may be involved in placental development and dysregulation of RNA editing in the placenta may contribute to the pathogenesis of preeclampsia.

Introduction

RNA editing was first discovered in the mitochondrial cytochrome c oxidase (COX) subunit II gene from trypanosomes in 1986 (Benne et al. 1986). It includes RNA sequence modifications, such as insertion or deletion of nucleotides and nucleotide conversion from cytidine (C) to uridine (U) (Chen et al. 1987; Powell et al. 1987; Benne 1994; Simpson et al. 2004), or from adenine (A) to inosine (I) (Bass and Weintraub 1987; Levanon et al. 2004). The C-to-U RNA editing involves site-specific changes in the mRNAs (Blanc and Davidson 2003), as are well documented in apolipoprotein B (*ApoB*) (Anant and Davidson 2001), neurofibromatosis type 1 (*NF1*) and N-Acetyltransferase 1 (*NAT1*) (Yamanaka et al. 1997; Mukhopadhyay et al. 2002). The most common RNA editing in mammals is A-to-I conversion, which is catalyzed by enzymes called adenosine deaminases acting on RNA (ADARs) (Polson et al. 1991; Savva et al. 2012). A-to-I RNA editing widely occurs in Alu transposons and noncoding regions such as introns and untranslated regions (UTRs) and in coding regions as well (Eisenberg and Levanon 2018).

RNA editing can lead to missense codon changes (Nishikura 2016), alternative splicing (Rueter et al. 1999), modification of microRNAs (miRNAs) and long noncoding RNAs (lncRNAs) (Kawahara et al. 2007; Gong et al. 2017), and miRNA binding sites (Nishikura 2010). The functions of RNA editing in noncoding regions are based on the site-specific modifications of RNA motifs, which mediate interactions with other molecules. Accumulating evidence on the function of RNA editing points to the regulation of RNA expression levels (Sommer et al. 1991; Nishikura 2010). RNA editing may regulate RNA expression levels through directly mediating RNA degradation (Kim et al. 2005), indirectly through alternative splicing (Rueter et al. 1999; Laurencikiene et al. 2006; Hsiao et al. 2018), or through modification of miRNA binding sites (Wang et al. 2013; Nakano et al. 2016). The suggested role for RNA-editing-mediated RNA degradation is supported by the observation that the editing enzyme ADAR interacts with UPF1, a mediator of nonsense-mediated decay

(NMD) (Agranat et al. 2008). RNA editing sites are enriched in intronic regions in sequencing data of RNA population purified by poly-d(T) (Huntley et al. 2016), suggesting that there might be an association between RNA editing and intron retention. Intron retention can affect gene functions in ways such as bringing in new functional elements (Buckley et al. 2011), down-regulating RNA expression through NMD (Belgrader et al. 1994), and interfering mRNA translation (Middleton et al. 2017). Several studies demonstrate that RNA editing in UTRs regulates mRNA degradation through disrupting miRNA binding (Wang et al. 2013; Nakano et al. 2016; Nakano et al. 2017).

RNA editing has been found in several human diseases (Maas et al. 2006), such as cancer and neurodegeneration (Dominissini et al. 2011; Singh 2012; Han et al. 2015; Hwang et al. 2016), involving in pathways of apoptosis (Wang et al. 2004; Liu et al. 2009), innate immune response (Mannion et al. 2014), cellular proliferation and DNA repair (Yeo et al. 2010). Genome-scale studies on RNA editing sites in cancer have revealed the critical roles of RNA editing in the global regulation of gene expression (Fumagalli et al. 2015; Han et al. 2015; Paz-Yaacov et al. 2015). Trophoblast cells in the placenta have some features similar to cancer cells (Holtan et al. 2009), and the placenta is a vital organ in human development, yet RNA editing in the placenta has been rarely documented. We do not know to what extent RNA editing might be involved in the pregnancy complications with placenta origin, such as preeclampsia (PE). Here, we carried out RNA editome profiling on placentae from 9 patients with early-onset severe preeclampsia (EOSPE) and 32 normal controls, and found 2,920 differentially edited sites (DESs) located in 1,229 genes. The genes with dysregulated RNA editing were enriched with known PE-associated genes, and also with differentially expressed genes (DEGs) in EOSPE. We further generated two mutations to mimic "RNA editing events" at two miRNA binding sites in 3'-UTR of *LEP*, leading to increased expression level of *LEP*. This is consistent

with the observed upregulation of *LEP* in the placenta of EOSPE patients. These results suggest that widespread RNA editing dysregulation in placentae might be involved in the pathogenesis of EOSPE.

Results

RNA editome profiling.

We performed RNA editome profiling on placental transcriptomes of 9 patients with EOSPE and 32 normal controls (**Supplemental Table S1**), using an in-house pipeline to search for PE-associated RNA editing sites (**Supplemental Fig. S1**). After removing adaptors and low quality reads, the cleaned reads were aligned to human reference genome (hg38 version) by tophat2 (**Supplemental Table S2**) and reliable variants were detected using Genome Analysis Toolkit (GATK) (McKenna et al. 2010; Kim et al. 2013). We identified 629,721 and 1,233,453 variants in EOSPE and normal samples respectively. After removing the variants that failed to pass the variant quality score recalibration (VQSR) filtering and/or had allele frequency higher than 0.05 in East Asian population (Eas_AF > 0.05) (See Methods), we detected 142,372 variants in 9 EOSPE samples and 399,565 variants in 32 normal samples respectively. Among these variants, 236,569 were identified as A/G or C/T mismatches, corresponding to A-to-I or C-to-U conversions in RNA (**Fig. 1A and 1B**), which are two main forms of RNA editing in mammals. Most of the potential RNA editing sites were located in transposons, especially in Alu transposons (**Fig. 1C**), consistent with previous studies in cancers (Bazak et al. 2014). About 89% (210,406/236,569) of these variants were located in noncoding regions (introns and intergenic regions) (**Fig. 1D**). The A-to-I RNA editing was the main editing form in most of the genomic regions, while C-to-U editing was found more often in exons.

Figure 1. RNA editome of placentae from patients with EOSPE and from normal subjects. (A) Flowchart for identification of RNA editing sites in placentae from EOSPE patients and normal subjects. Red boxes represent the analyses; blue boxes represent results of the analyses. EOSPE: early-onset severe preeclampsia; DES: differentially edited site; UTR: untranslated region. See Figure S1 for detailed flowchart. (B) RNA editing types in placentae. (C) Distribution of RNA editing sites in transposon elements (TE) and non-transposon regions. Transposon elements were extracted from RepeatMasker Database. (D) Distribution of RNA editing sites in exons, introns, 5'-UTRs, 3'-UTRs and intergenic regions. The genomic regions were defined by RefGene annotation. Numbers above the bars are the number of the sites. (E) Data overlaps and differences between our placental RNA editome dataset and RNA editing databases. (F) Validation of 89 RNA editing sites by Sanger sequencing and multiple alignment of the reads. Yellow bars represent the fraction of validated RNA editing sites and blue bars represent the fraction of genomic DNA mutations. Numbers within the bars are the number of the sites (all validation results of 89 sites were listed in **Supplemental Table S3**).

There are rarely studies on RNA editing in the placenta, and even the most comprehensive database REDIportal does not cover placenta (Picardi et al. 2017). Thus, the placenta-specific RNA editing sites are depleted in current RNA editing databases. We compared the RNA editing sites we discovered in the placenta with known RNA editing sites in the DARNED (Kiran and Baranov 2010), RADAR (Ramaswami and Li 2014) and REDIportal databases (Picardi et al. 2017). There are 3,149 RNA editing sites in our dataset overlapping with all of the three RNA editing databases, and about 48% (112,868/236,569) of placental RNA editing sites are recorded in REDIportal database. The remaining 121,749 RNA editing sites are not included in any of these three databases, indicating that these RNA editing sites may be placenta specific (**Fig. 1E**). To

empirically validate our dataset, we randomly picked 33 RNA regions with 89 RNA editing sites and amplified

the genomic DNA regions by PCR for Sanger sequencing (**Supplemental Table S3**). The validation rates were

80% for exonic editing sites, 71% for intronic editing sites and nearly 100% for editing sites in UTR regions

(**Fig. 1F**). These results demonstrated that our placental editome dataset is of high-quality.

Differentially edited sites (DESs) are associated with EOSPE and gene expression level.

For each RNA editing site, we calculated the editing ratio in every sample and evaluated the editing ratio

difference between EOSPE and normal placenta by one-tailed Wilcoxon test (See Methods). At the cutoff P

$= 0.05$, we obtained 2,920 differentially edited sites (DESs) which exhibited significant difference of editing

ratio between EOSPE and normal subjects (**Fig. 2A and Supplemental Table S4**).

The genes with DESs were significant enriched for PE-associated genes collected from literature (**Fig. 2B

and Supplemental Table S5**), and for differentially expressed genes (DEGs) in EOSPE (**Fig. 2C,

Supplemental Table S6** and related manuscript by Ren et al.), suggesting the DESs may be involved in PE.

To explore the biological functions and pathways of the genes with DESs, we performed enrichment analysis

of these genes in Gene Ontology (GO) terms and Kyoto Encyclopedia of Genes and Genomes (KEGG)

pathways using ClusterProfiler (Yu et al. 2012). We found that the genes with DESs are significantly enriched

in terms such as the female pregnancy, placenta development and autophagy (adjusted $P < 0.05$, **Supplemental

Fig. S2A and Supplemental Table S7**), and that the genes with DESs are significantly enriched in nine

pathways, including endocytosis, focal adhesion, ubiquitin mediated proteolysis and protein processing in

endoplasmic reticulum (**Fig. 2D and Supplemental Table S7**).

Figure 2. Overview of EOPSE differentially edited sites (DESs). (A) Distribution of DESs in genomic regions. Genomic regions were defined by RefGene annotation. Numbers above the bars are the numbers of the sites. See Table S4 for all DESs. (B-C) Fraction of known PE-associated genes (B) and fraction of differentially expressed genes (DEGs) of EOSPE (C) in the genes with DESs. P-values were calculated by Fisher-exact test in R. *: $P < 0.05$; **: $P < 0.01$; ***: $P < 0.001$; ****: $P < 0.0001$; n.s.: no significance. The standard error of the fraction was estimated using a bootstrapping method with 10,000 resamplings. PE-associated genes and EOSPE DEGs were listed in Supplemental Table S5 and S6. (D) The KEGG pathways enriched with genes harboring DESs. Color of bars: adjusted P-values; X-axis: the number of genes harboring DESs in the pathways. Enrichment analysis and P-value calculation were carried out using ClusterProfiler. (E) Gene-pathway-bipartite network for genes with DESs. Yellow diamonds represent the KEGG pathways, and red circles represent up-regulated genes in EOSPE, blue down-regulated genes and gray genes with no significant expression change. See Supplemental Table S7 for P-values for pathway enrichment (D) and gene-pathway interactions (E).

The endocytosis pathway contains 30 genes with DESs, including 5 EOSPE up-regulated genes (*ARPC2*, *PARD6B*, *EPS15*, *RHOA* and *EGFR*) and 3 EOSPE down-regulated genes (*ARAP3*, *RAB11FIP1* and *CYTH3*) (**Fig. 2E**), suggesting that the endocytosis pathway may be regulated by differential editing and may be involved in the pathogenesis of EOSPE. The focal adhesion pathway contains 23 genes with DESs, including 4 EOSPE up-regulated genes (*FLT1*, *DIAPH1*, *RHOA* and *EGFR*) (**Fig. 2E**). The gene *FLT1* is one of the few known potential biomarkers of PE (Thomas et al. 2007; Jebbink et al. 2011; McGinnis et al. 2017), and the genes *RHOA* and *EGFR* are also reported to be associated with PE (Hornsby et al. 1989; Faxen et al. 1998;

Zhou and Qiao 2006), suggesting that the focal adhesion function may be commonly affected in the placenta of EOSPE.

To investigate if these DESs were associated with gene expression levels, we classified all placenta samples into two groups, with vs. without RNA edited sites (edited group and not-edited group), for each DES (See Methods). One-tailed Wilcoxon test was applied to examine the difference in the expression levels of the gene containing the editing site between edited and not-edited group. We found significant differential expression for 352 genes with 542 DESs ($P < 0.05$, **Table S8**). We also calculated the correlation coefficient between editing ratios and the corresponding gene expression levels for DESs, and found that 305 DESs showed significantly correlation between editing ratio and gene expression level (**Supplemental Table S8**), suggesting that the editing ratio for some editing sites on the transcripts of a gene might regulate the gene expression level.

DESs in exons, introns and 3'-UTRs are associated with PE.

Of the 2,920 DESs, 225 were located in exons, 1,621 in introns, and 335 in 3'-UTRs. We calculated the enrichment of the literature-curated PE-associated genes (**Supplemental Table S5**) and DEGs in EOSPE (**Supplemental Table S6**) in the corresponding genes of these DESs. The genes harboring DESs in the three different regions were enriched with PE-associated genes respectively, suggesting the disease-association of DESs in each category (**Fig. 3A**). The genes with intronic DESs were also enriched with DEGs in EOSPE (**Fig. 3B**), suggesting that intronic DESs might be involved in the dysregulation of gene expression in this disease.

Figure 3. DESs in exons, introns and 3'-UTRs. (A-B) Fraction of known PE-associated genes (A) or DEGs of EOSPE (B) in genes with DESs in different regions. *P*-values were calculated by Fisher-exact test in R. *: *P* < 0.05; **: *P* < 0.01; ***: *P* < 0.001; ****: *P* < 0.0001; n.s.: no significance. The standard error of the fraction was estimated using a bootstrapping method with 10,000 resamplings. (C) Mutation types of exonic DESs. Mutation types were annotated based on RefGene using ANNOVAR. The numbers above the bars are the numbers of DESs. (D) Association between intronic DESs and gene expression. Red circle of the Venn diagram represents intronic DESs with significant gene expression difference, and yellow circle represents intronic DESs with significant correlation between editing ratio and gene expression level. See Supplemental Table S8 for all the intronic DESs associated with gene expression. (E) Bipartite network of miRNAs and genes with DESs in 3'-UTR. The miRNA-targeting regions harboring DESs in 3'-UTRs were predicted by TargetScan. Yellow: genes with DESs in 3'-UTRs; blue: miRNAs.

Among the 225 exonic DESs, 108 were located in protein-coding genes and 117 in noncoding RNAs. Of the 108 protein-coding exonic DESs, 73 (18 A-to-I and 55 C-to-U) were synonymous, 31 (16 A-to-I and 15 C-to-U) were nonsynonymous, 3 were stop-gain and 1 was unknown (**Fig. 3C**). Among these 34 nonsynonymous/stopgain DESs, 26 exhibited up-regulated editing levels in EOSPE samples (**Supplemental Fig. S3A**). Eight of the nonsynonymous DESs were predicted to have damaging effects on proteins according to the SIFT and Polyphen2 values by ANNOVAR (**Table 1**). For example, the DES at the position chr3:49360951 in exon of *RHOA* was predicted as a damaging mutation. *RHOA* showed increased editing ratio in EOSPE samples and had interactions with multiple DEGs of EOSPE (**Supplemental Fig. S3A and S3B**), suggesting that this DES in *RHOA* may play a role in the development of EOSPE.

Table 1. Exonic differentially edited sites that predicted as damaging/harmful

Chr[a]	Pos[b]	Type	Gene	Avsnp147	SIFT Pred	Polyphen2 Pred
chr1	111476845	C-to-U	C1orf162	rs150674974	D[c]	B[d]
chr1	155612848	C-to-U	MSTO1	rs622288	D	B
chr3	49360951	A-to-I	RHOA	-	D	-
chr3	122341557	C-to-U	CSTA	rs34173813	D	P[e]
chr3	195780494	C-to-U	MUC4	rs75737861	D	B
chr10	45826300	C-to-U	AGAP4	rs201626545	D	B
chr10	46383535	C-to-U	ANXA8L1	-	-	P
chr10	93681515	C-to-U	FRA10AC1	rs11187583	D	P

[a]Chr, chromosome; [b]Pos, position; [c]D, damaging; [d]B, benign; [e]P, possibly damaging

It is reported that RNA editing tends to occur in introns (Athanasiadis et al. 2004), and some intronic RNA editing sites are known to regulate gene expression by influencing circRNA biogenesis and alternative splicing (Rueter et al. 1999; Laurencikiene et al. 2006; Ivanov et al. 2015; Hsiao et al. 2018). We speculated that there might be an association between intronic DESs and intron retention (IR). We identified IR events by comparing the transcripts assembled by StringTie with the canonical transcripts from GENCODE (Pertea et al. 2016) (See Methods). For every IR event, we calculated the normalized percent spliced in (PSI) and compared the PSI between edited group and not-edited group (See Methods). We identified 359 intronic DESs in the retained introns, and found that 36 IR events with 42 intronic DESs which exhibited significant PSI difference between

edited and not-edited group by one-tailed Wilcoxon test (**Supplemental Fig. S3C**). Most of these DESs exhibited increased PSI in the edited group. Among these 42 DESs, 17 exhibited significantly difference of gene expression between edited and non-edited group. These results show that some intronic DESs are associated with intron retention.

To evaluate the association of the intronic DESs and the expression levels of corresponding genes, we compared the expression difference of the genes between two groups of placental samples with differential RNA editing (See Methods). We observed 390 intronic DESs located in genes which exhibit significant differential expression between placental samples with vs. without RNA edited sites (edited group vs. not-edited group) (**Fig. 3D and Supplemental Table S8**). We further calculated the correlation coefficient between RNA editing ratio and gene expression level, and found that 211 sites showed significant correlation between editing ratio and the expression level of corresponding genes (**Fig. 3D and Supplemental Table S8**), suggesting that these intronic DESs may regulate the gene expression level.

As a common regulatory strategy used by the cell, microRNA (miRNA) can bind to specific sequences in the 3'-UTRs of target mRNAs and regulate their stability (Wang et al. 2013; Nakano et al. 2016; Nakano et al. 2017). We predicted 2,219 miRNA-targeting regions in genes harboring DESs using TargetScan (Shi et al. 2017), and 51 DESs are located in 45 known miRNA-targeting regions (**Fig. 3e and Supplemental Table S9**). After discarding the DESs located in regions with null preferentially conserved targeting (PCT) values, we identified 31 DESs that may regulate miRNA binding (**Table 2**).

Table 2. 31 DESs in 3'-UTRs that located in miRNA-targeting regions with non-null PCT[a] values.

MiRNA	Chr[b]	Position	Gene	Transcript
miR-10-5p	chr1	35602159	PSMB2	ENST00000373237.3
miR-219-5p	chr1	52824220	ZYG11B	ENST00000294353.6
miR-24-3p	chr1	100024500	SLC35A3	ENST00000465289.1
miR-22-3p	chr1	202585129	PPP1R12B	ENST00000608999.1
miR-150-5p	chr1	204552462	MDM4	ENST00000391947.2
miR-455-3p.2	chr1	220058113	BPNT1	ENST00000322067.7
miR-10-5p	chr2	98623012	MGAT4A	ENST00000264968.3
miR-375	chr2	147931120	ORC4	ENST00000392857.5
miR-214-5p	chr7	17344771	AHR	ENST00000242057.4
miR-9-5p	chr7	128255810	LEP	ENST00000308868.4
miR-9-5p	chr7	128255811	LEP	ENST00000308868.4
miR-10-5p	chr8	1782334	CLN8	ENST00000331222.4
miR-184	chr8	43023708	HOOK3	ENST00000307602.4
miR-217	chr11	32853772	PRRG4	ENST00000257836.3
miR-217	chr11	78069179	NDUFC2	ENST00000281031.4
miR-214-5p	chr12	50930723	METTL7A	ENST00000332160.4
miR-214-5p	chr14	21460463	RAB2B	ENST00000397762.1
miR-214-5p	chr16	10529809	EMP2	ENST00000359543.3
miR-141-3p/200a-3p	chr16	13239082	SHISA9	ENST00000558583.1

miR-217	chr16	13239082	SHISA9	ENST00000558583.1
miR-490-3p	chr19	2836037	ZNF554	ENST00000317243.5
miR-196-5p	chr19	4654589	TNFAIP8L1	ENST00000327473.4
let-7-5p/98-5p	chr19	4654589	TNFAIP8L1	ENST00000327473.4
miR-214-5p	chr19	16550610	SLC35E1	ENST00000595753.1
miR-214-5p	chr19	51519173	SIGLEC6	ENST00000346477.3
miR-17-5p/20-5p/93-5p/106-5p/519-3p	chr19	52584380	ZNF701	ENST00000540331.1
miR-192-5p/215-5p	chr19	52839344	ZNF468	ENST00000595646.1
miR-214-5p	chr19	57861909	ZNF587	ENST00000339656.5
miR-216a-5p	chr20	33646143	CBFA2T2	ENST00000375279.2
miR-216b-5p	chr20	33646143	CBFA2T2	ENST00000375279.2
miR-10-5p	chrX	2906938	ARSD	ENST00000381154.1
miR-216a-5p	chrX	24077208	EIF2S3	ENST00000253039.4
miR-216b-5p	chrX	24077208	EIF2S3	ENST00000253039.4
miR-26-5p	chrX	77827544	MAGT1	ENST00000358075.6
miR-455-3p.2	chrX	77828504	MAGT1	ENST00000358075.6

a)PCT: preferentially conserved targeting; b)Chr: chromosome

RNA editing in 3'-UTR regulates *LEP* expression by disrupting miRNA binding.

Of all the DESs in 3'-UTRs, 24% (81/335) show significant correlation ($P < 0.05$) between editing ratio and gene expression level (**Table S8**), and five of them, including *EIF2S3*:chrX:24077208:A-to-I, *LEP*:chr7:128255810:A-to-I, *LEP*:chr7:128255811:A-to-I, MGAT4A:chr2:98623012:A-to-I and *ZNF554*:chr19:2836037:A-to-I, are located in the predicted miRNA-targeting regions (**Fig. 4A and Table 2**).

The *LEP* gene is known to be highly up-regulated in preeclamptic placenta (Enquobahrie et al. 2008), and the elevated circulating leptin level is an indicator for severe preeclampsia (Teppa et al. 2000). In our EOSPE placental transcriptome, the *LEP* expression exceeded normal level by more than 60 folds. We identified 11 DESs, and 10 of them are located in 3'-UTR of *LEP* (**Fig. 4B**). We evaluated the expression level of *LEP* and *miR-9-5p* in the placenta samples from 9 EOSPE patients and 9 randomly picked normal subjects using quantitative real-time PCR (qRT-PCR). The *LEP* was significantly up-regulated in the EOSPE compared to the normal controls (**Fig. 4C**), whereas *miR-9-5p* showed no significant expression difference (**Fig. 4D**). This missing correlation between *LEP* and *miR-9-5p* expression level indicated that the change of *LEP* expression level in patients was not due to a *miR-9-5p* expression change.

Figure 4. DESs in 3'-UTRs regulate gene expression by inhibiting miRNA binding. (A) Association of DESs in 3'-UTRs with gene expression level. Samples were classified into edited group or not-edited group based on the existence of the RNA editing event at a given editing site. The relative expression level was estimated as CPM (counts per million bases) using edgeR. One-tailed Wilcoxon test was used to evaluate the gene expression difference between edited group and not-edited group using R package. *LEP.1* referred to

'*LEP*:chr7:128255810:A-to-I'; *LEP.2* referred to: '*LEP*:chr7:128255811:A-to-I'. See Table S8 for CPM

values of these sites. (B) Two clusters of DESs (C1 and C2) in *LEP* 3'-UTR were shown in red letters and

labeled as S1-S10. S6 and S7 were located in the miRNA-targeting region and thus were chosen for testing

the effect of RNA editing on miRNA binding. (C-D) The qRT-PCR results of *LEP* (C) and *miR-9-5p* (D) on

placentae of 8 EOSPE patients and 8 randomly-chosen normal subjects. *ACTB* (C) or *U6* (D) was taken as

internal control. Wilcoxon test was performed to evaluate the difference of expression levels between

EOSPE and normal placentae using R package. Error bars represent the standard errors. (E) The qRT-PCR

results of *LEP* in HTR-8/SVneo cells with/without *miR-9-5p* transfection. *ACTB* was used as internal

control. Wilcoxon test was performed to evaluate the difference of *LEP* expression levels using R package.

Error bars represent the standard errors. (F) The results of luciferase assay on HTR-8/SVneo cells

transfected with plasmids carrying edited *LEP* 3'-UTR (*LEP$_{MUT2-UTR}$*) or not-edited *LEP* 3'-UTR (*LEP$_{WT-UTR}$*)

at S6 and S7 sites. Y-axis represents the relative luciferase activity. Wilcoxon test was used to evaluate the

difference of luciferase activity using R package. Error bars represent the standard errors. Significance in

(A) and (C-F): *: $P < 0.05$; **: $P < 0.01$; ***: $P < 0.001$; ****: $P < 0.0001$; n.s.: no significance. See

Supplemental Table S10 for numerical values in (C-F). (G) Schematic to illustrate the mechanism for 3'-

UTR RNA editing in the regulation of *LEP* expression.

　　Among the 10 DESs in the 3'-UTR of *LEP*, 2 sites (chr7:128255810:A-to-I, S6 and chr7:128255811:A-

to-I, S7) (**Fig. 4B**) are located in *miR-9-5p* targeting region, and the editing ratios of these two DESs are

significantly correlated with the expression level of *LEP* (**Supplemental Table S8**). These results suggest that

these two DESs in *LEP* 3'-UTR may alter *miR-9-5p* targeting motif and thus may inhibit miRNA-induced *LEP*

degradation, which may in turn lead to an upregulation of *LEP*.

To test this hypothesis, we first evaluated the regulation of *miR-9-5p* on *LEP* expression and then tested the effect of "RNA editing events" of these two editing sites on the expression of *LEP*. To evaluate the regulation of *miR-9-5p* on *LEP*, we over-expressed *miR-9-5p* in the HTR-8/SVneo cell line and quantified the expression level of *LEP* using qRT-PCR, and found *LEP* expression decreased significantly compared to control group (**Fig. 4E**), demonstrating that *miR-9-5p* could repress the expression of *LEP*. To test if *LEP* expression was regulated by the editing of the *miR-9-5p* binding sites, we constructed two reporter plasmids carrying a luciferase coding sequence fused to a 3'-UTR of *LEP* which had a *miR-9-5p* targeting region with the RNA editing sites S6 and S7. One plasmid contained wild-type (WT) 3'-UTR with normal S6 and S7 (*LEP$_{WT-UTR}$*), the other contained mutated 3'-UTR with A-to-G conversion (A-to-I editing) at S6 and S7 nucleotides (*LEP$_{MUT2-UTR}$*). These plasmids were co-transfected with *miR-9-5p* into the HTR-8/SVneo cell line derived from placenta trophoblast, and luciferase activity was examined to test the effect of miRNA on the expression of the reporter gene with edited or not-edited 3'-UTR of *LEP*.

As expected, the luciferase activity was down-regulated by 20% when *miR-9-5p* and *LEP$_{WT-UTR}$* were co-transfected into the cells, compared to transfection of *LEP$_{WT-UTR}$* alone (**Fig. 4F**), demonstrating that the luciferase activity was suppressed by *miR-9-5p* when the luciferase mRNA has wild type 3'-UTR of *LEP*. The luciferase activity was not changed when *miR-9-5p* and *LEP$_{MUT2-UTR}$* are co-transfected, compared to transfection of *LEP$_{MUT2-UTR}$* alone (**Fig. 4F**), suggesting that the luciferase expression was not suppressed by *miR-9-5p* when the luciferase mRNA has mutated 3'-UTR of *LEP*. These results suggest that RNA editing on the *miR-9-5p* binding sites in 3'-UTR may inhibit *LEP* mRNA degradation (**Fig. 4G**).

Discussion

Genome-scale RNA editing profiling has been done in a series of human normal tissues including brain, lung, kidney, liver, heart and muscle (Blow et al. 2004; Picardi et al. 2015). However, RNA editing has rarely been studied in placenta. Since RNA editing dysregulation is a common phenomenon in cancer, large-scale RNA editing studies on cancer tissues and cell lines have been well documented (Delva et al. 1987; Park et al. 2012; Xu et al. 2018). Recently, 85 genes were found to have differential RNA editing in two placental cell lines treated with hypoxia using transcriptome sequencing (preprint in bioRxiv) (Hasan et al. 2019). Since preeclampsia is thought to be caused by poor placentation, We have recently carried out transcriptome sequencing on placentae from 33 preeclampsia patients (including 9 early-onset severe, 15 late-onset severe and 9 late-onset mild PE patients) and 32 normal subjects, and found a large number of placental dysregulated genes in EOSPE (see related manuscript by Ren et al.). To further find out to what extent the RNA editing might be involved in gene dysregulation in preeclamptic placenta, we performed RNA editome profiling on the placental transcriptomes of the 9 patients with EOSPE and 32 normal controls, and identified a total number of 236,569 RNA editing sites, including 2,920 differentially edited sites (DESs) (**Fig. 1D and 2A**), which can be considered as potential disease-associated sites. The 2,920 DESs include 720 sites in intergenic regions and 2,200 in 1,229 genes, which contains 225 sites in exons, 1,621 in introns and 354 in UTRs (**Fig. 1A and Supplemental Table S4**). DESs in exons, introns or UTRs are enriched with known PE-associated genes respectively (**Fig. 3A**), suggesting that RNA editing sites in both coding and non-coding regions may be involved in PE.

The exonic DESs are mostly C-to-U site-specific changes. Some of these DESs might change the functions of the proteins through generating missense or stop-gain mutations, as reported in many studies on RNA editing in some diseases (Skuse et al. 1996; Anant and Davidson 2001; Mukhopadhyay et al. 2002).

Since the majority (73.7%) of the DESs in genes are located in introns, and since the intronic DESs were enriched with DEGs in EOSPE (**Fig. 3B**), we recognized their potential role in the preeclamptic gene dysregulation in the placenta (**Fig. 2A**). RNA editing in the introns affects alternative splicing and RNA degradation in cancer (Kim et al. 2005; Hsiao et al. 2018). The DESs in introns may lead to the gene dysregulation through the same mechanism. RNA editing in UTRs participates in the gene expression homeostasis during cancer development (Yang et al. 2017; Sagredo et al. 2018). Two DESs in the 3'-UTR of *LEP* gene were empirically verified to regulate the gene expression level (**Fig. 4**). We generated "RNA editing events" at miRNA-targeting sites in 3'-UTR of *LEP*, and confirmed that the RNA editing at these sites was able to increased expression level of *LEP* (**Fig. 4**), as observed in patients, suggesting that RNA editing is indeed involved in the gene expression regulation in the placenta of PE patients through modification of miRNA-editing site in 3'-UTRs. RNA editing on miRNA-targeting sites in 3'-UTRs is reported in previous studies in cancer (Yang et al. 2017; Sagredo et al. 2018).

Our comprehensive placental RNA editome dataset contains 236,569 RNA editing sites, 121,749 (51.5%) being placenta specific, which have not been reported in current RNA editome databases, 48.5% being common with other tissues, which have been reported in at least one database. We have identified 2,920 differentially edited sites (DESs) as potential PE-associated RNA editing sites. In conclusion, there is widespread RNA editing in the transcriptome of the placenta, and the dysregulated RNA editing on a group of editing sties may be involved in the development of preeclampsia.

Methods

Sample Collection.

This research has been approved by the Ethnics Board of Nanfang Hospital, Southern Medical University, China and all patients have signed the informed consent. We collected the tissue samples at the Department of

Obstetrics & Gynecology of Nanfang Hospital in China from January 2015 to July 2016. The clinical characteristics of each patient were extracted from the medical records, which strictly followed the American Board of Obstetrics and Gynecology, Williams Obstetrics 24th edition. Preeclampsia (PE) is characterized by new-onset hypertension (≥140/90 mmHg) and proteinuria at ≥20 weeks of gestation, or in the absence of proteinuria, hypertension with evidence of systemic disease (Chaiworapongsa et al. 2014). If patients with PE have systolic blood pressure ≥160 mmHg or diastolic blood pressure ≥110 mmHg, or other organ failures, we call them "preeclampsia with severe features" or briefly "severe PE". According to gestational age at diagnosis or delivery, PE can be divided as early-onset (< 34 weeks) and late-onset (≥ 34 weeks) (Redman and Sargent 2005). This study focused on the early-onset severe PE (EOSPE). Samples of placental tissues were collected from the mid-section of the placenta between the chorionic and maternal basal surfaces at four different positions, within 5 minutes after delivery. These mid-section placental tissues were placed into RNAlater® solution and stored at -80 ℃. The clinical information of the three clinical subtypes and comparisons between subtypes were listed in Supplemental Table S1.

RNA extraction and RNA-seq.

Total RNA was isolated using the RNeasy® Plus Universal Mini Kit (Qiagen, DEU) according to the manual manufacturer's instruction. Briefly, RNAs with poly(A) tails were isolated, and double-stranded cDNA libraries were prepared using TruSeq RNA Kit (Illumina, USA). The 125-base paired-end sequencing was carried out using Illumina Hiseq 2500 platform (Berry Genomics, CHN).

Reads alignment and variant calling.

Adaptor sequences were trimmed before read alignment. If any read in a pair exhibited more than 4 unidentified bases or more than 63 bases (half of the read length) with base quality score lower than 3, the paired reads would be judged as low quality and discarded. Then RNA-seq reads were indexed using bowtie2 (v2.2.8) and aligned to the genome using tophat2 (v2.1.1) (run with parameters '–p6 –r50 –b2 –very–sensitive') (Langmead and Salzberg 2012; Kim et al. 2013). The reference genome was downloaded from GATK (Genome Analysis Toolkit) (McKenna et al. 2010) hg38 bundles. Duplication levels were evaluated by Picard 'MakeDuplicates' (https://github.com/broadinstitute/picard; Broad Institute) (**Supplemental Table S2**). GATK (v3.5) recommended pipeline was modified to suit for RNA-seq data (**Supplemental Materials**). Only variants that passed GATK VQSR (Variant quality score recalibration) filtering were retained. ANNOVAR (v2016-02-01) was used to annotate pass-filtering variants (**Supplemental Materials**) and variants with allele frequency (estimated by 1000 genomes project) smaller than 0.05 in East Asian population (Eas_AF < 0.05) or without allele frequency annotation were selected for further analysis (Wang et al. 2010). Variants with ANNOVAR annotated as 'splicing', 'upstream' or 'downstream' would be collapsed into 'intergenic' category.

Detection of potential RNA editing sites.

Considering that our sequencing library was not strand specific, we extracted the variants with A/G and C/T mismatch or T/C and G/A mismatch corresponding to plus and minus strand. These detected variants were taken as potential as A-to-I and C-to-U RNA editing sites.

Validation of candidate RNA editing sites.

We collected 2,576,459 known RNA editing sites from database RADAR (version 2) (Ramaswami and Li 2014), 314,250 from DARNED (Kiran and Baranov 2010) and 4,668,508 from REDIportal (Picardi et al.

2017). Since these RNA editing databases were annotated using hg19 version genome, we converted coordinates of the RNA editing sites we detected to corresponding position in hg19 genome through LiftOver tool in UCSC (http://genome.ucsc.edu/cgi-bin/hgLiftOver) and 235,488 sites were successfully converted. The editing sites detected at mRNA level were re-evaluated by multiple sequence alignment of reads that covered the RNA editing sites.

For empirical validation, we randomly chosen 33 regions with 89 RNA editing sites, including 17 sites from exons, 37 from UTRs and 35 from introns (**Supplemental Table S3**). The regions with RNA editing sites were PCR-amplified from gDNA from placenta samples (9 EOSPE vs. 9 normal samples; Supporting Information) and then Sanger-sequenced for validation of the editing events (Sangon Biotech, CHN). Sites without gDNA mutations at the corresponding potential RNA editing sites (A-to-G or C-to-T on mRNA) were considered as validated RNA editing sites.

Detection of EOSPE differentially edited sites (DESs).

We evaluated the RNA editing level for a given site by RNA editing ratio, which is defined as the ratio of reads with editing to total reads. If there is no read covering the editing site in a sample, editing ratio for this site in this sample would be assigned as zero. To detect reliable differentially edited sites (DESs) in placenta, we set filtering criteria as: (i) having no less than three reads that supported the editing site and no less than five reads that covering the editing site in at least three samples; (ii) no more than three (~30%) EOSPE samples that had editing ratio = 1; (iii) no more than nine (~30%) normal samples that had editing ratio = 1. A total number of 23,645 editing sites were kept for DES detection. After classified editing sites into EOSPE up-editing and EOSPE down-editing groups, we compared the editing ratios between EOSPE and normal

placentae by one-tailed Wilcoxon test for these two groups. Then we selected RNA editing sites exhibited significantly editing ratio differences between EOSPE and normal placentae ($P < 0.05$) as DESs.

Detection of EOSPE differentially expressed genes (DEGs).

We mapped clean reads to human genome (hg38 version) by STAR (v2.5.2b) and called raw counts for every gene in every sample by HTSeq (v0.9.0) htseq-count separately (Dobin et al. 2013; Anders et al. 2015). Differentially expressed genes (DEGs) were detected using DESeq2 (v1.16.1) (Love et al. 2014) and edgeR (v3.18.1) (Robinson et al. 2010), and intersection of results from these two detection methods were subsequently taken as final DEGs (2,977 genes). Human reference genome and gene annotations were downloaded from Ensembl.

Collection of known PE-associated genes.

Known PE-associated genes were collected from eight published papers. After dataset merging, totally 1,177 genes were referred as PE-associated genes for further analysis (**Supplemental Table S5**).

Enrichment calculation of EOSPE DEGs and PE-associated genes in genes with DESs.

According to the enrichment calculation of hypergeometric distribution, we calculated enrichments for DEGs in EOSPE and PE-associated genes by comparing the fraction of these genes in foreground and background dataset. Genes with DESs and all expressed genes in EOSPE and normal placentae were referred as foreground and background dataset. A gene was defined as expressed gene if there is at least one read in all samples of both EOSPE and normal placentae. One tailed Fisher-exact test was used to calculate the P-value. The standard

error of the fraction was estimated using a bootstrapping method with 10,000 resamplings (randomly extracted 10% genes from foreground and background per resampling).

Functional enrichment of GO terms and KEGG pathways in genes with DESs.

Functional enrichment analysis was achieved by ClulsterProfiler (v3.10.1) through R (Yu et al. 2012). Threshold of adjusted P-values were set to 0.05 to find significantly enriched GO terms or KEGG pathways.

Association of differential RNA editing with differential gene expression.

For every differentially edited site (DES), we re-classified all 41 samples into two groups: edited group and not-edited group based on the existence of the editing event. DESs with less than three samples in edited or not-edited group were removed. Genes without expression in any of the samples were also removed. We then compared the gene expression between edited group and not-edited group by one-tailed Wilcoxon test. The DESs which had significant gene expression differences between edited group and not-edited group ($P < 0.05$) were considered as had association with the differential gene expression. Counts per million bases calculated by edgeR were used as gene expression here.

Correlation between differential RNA editing and gene expression level.

For a given DES, we calculated the correlation between editing ratio and gene expression level by Hmisc (https://cran.r-project.org/web/packages/Hmisc/index.html; v4.2-0) in R. Genes without expression in any of the samples were removed. Counts per million bases calculated by edgeR were used as gene expression. $P = 0.05$ was set as cutoff for significant correlation.

Protein-protein interaction (PPI) network of *RHOA* and EOSPE DEGs.

Human interactome data was from InWeb_InBioMap (https://www.intomics.com/inbio/map) and *RHOA*-centered PPI network was extracted from human interactome network containing *RHOA* and its interactors (Li et al. 2017). Then EOSPE DEGs were mapped onto *RHOA*-centered network and *RHOA*- subnetwork of DEGs in EOSPE was extracted. The network extraction was done using CytoScape (v3.4.0) (Shannon et al. 2003).

Identifying intron retention (IR) event.

We downloaded GENCODE canonical transcript annotation (v29) from UCSC Table Browser (Karolchik et al. 2004). The transcripts mapped to scaffolds were removed. If a gene had multiple transcripts mapped to different chromosomes, the gene would also be removed. Then we got a set of genes with one or more canonical transcripts. If the gene had more than one canonical transcripts, we merged them into one transcript by keeping the nonoverlapping exons and collapsing the overlapping exons. The final set of canonical transcripts were unique to their corresponding genes.

We assembled transcripts from RNA-seq reads by StringTie (v1.3.4) (Pertea et al. 2015), and compared them with known transcripts (ftp://ftp.ensembl.org/pub/release-84/gtf/homo_sapiens/Homo_sapiens.GRCh38.84.gtf.gz) by cuffcompare in Cufflinks (v2.2.1) (Trapnell et al. 2010). We further compared the assembled transcripts with corresponding final set of canonical transcripts, and identified the regions of the assembled transcripts that fell into the introns of the canonical transcripts. These regions were defined as IR events.

Calculation of percent spliced in (PSI).

For every IR event, we calculated the percent spliced in (PSI) according to the below formula (Schafer et al. 2015):

$$\text{PSI} = \frac{adjusted\ spliced-in\ reads}{(adjusted\ spliced-in\ reads + adjusted\ spliced-out\ reads)},$$

$$adjusted\ spliced - in\ reads = \frac{number\ of\ reads\ supproting\ IR\ events}{(IR\ event\ length\ +\ read\ length\ -\ 1)},$$

$$adjusted\ spliced - out\ reads = \frac{number\ of\ reads\ skipping\ introns}{(read\ length\ -\ 1)}.$$

Association of intronic differential RNA editing with intron retention.

We compared the percent spliced in (PSI) of each intron retention between edited group and not-edited group by one-tailed Wilcoxon test. For a DES, if RNA editing happened in less than three samples within either edited group or non-editing group, it was removed. The DESs which had significant PSI differences between edited group and not-edited group ($P < 0.05$) were considered as associated with the IR events.

Prediction of miRNA-targeting genes harboring DESs

Human miRNA-targeting information was downloaded from TargetScan database (v7.2) (Shi et al. 2017), and used for identifying DESs in miRNA-targeting regions.

Estimation of relative expression level of LEP and miR-9-5p in placentae.

For *LEP* and *miR-9-5p*, the mRNA expression level was evaluated by quantitative real-time quantitative PCR (qRT-PCR) in 8 EOSPE samples and 8 randomly-chosen normal samples. *ACTB* and U6 were used as internal control for *LEP* and *miR-9-5p* respectively. Expression levels of mRNAs were normalized to internal control

mRNA. The relative expression levels of the genes were calculated using the $2-\Delta\Delta Ct$ method. ΔCt was the difference of Ct values between quantification of *LEP* and *GAPDH* or between *miR-9-5p* and U6. The standard error of $2-\Delta\Delta Ct$ values was calculated to serve as error bar. The primers used for mRNA quantification were listed in Supporting Information. Wilcoxon test was used to calculate the significance of the expression difference between EOSPE and normal group using R package.

Cell culture.

HTR-8/SVneo cell line was purchased from ATCC (USA). Cells were grown in RPMI 1640 media (Gibco, Thermo Fisher Scientific, USA) supplemented with 10% (v/v) fetal bovine serum (FBS) and incubated at 37 °C and 5% (v/v) CO_2.

Quantitative real-time PCR (qRT-PCR) experiment.

The *miR-9-5p* mimic or negative control (NC) mimic (Shanghai GenePharma Co., Ltd, CHN) was transfected into HTR-8/SVneo cell line by Lipofectamine 2000 (Invitrogen, Thermo Fisher Scientific, USA) in six-well plates, and cells were harvested after 48 hours of transfection. RNAs were extracted using RNAeasy Mini Kit (Qiagen, USA) according to the manufacturer's instruction. The RNAs were reverse-transcribed into cDNAs using the PrimeScriptTM RT Kit (Takara, JPN) following the manufacturer's instruction. *ACTB* was used as the internal control. The qRT-PCR of *LEP* mRNA quantification in HTR-8/SVneo cell line was the same as described above. The relative expression levels of the genes were calculated using the $2-\Delta\Delta Ct$ method. Wilcoxon test was used to calculate the significance of the expression difference between the cells transfected with *miR-9-5p* mimic and with negative control mimic using R package. The standard error of $2-\Delta\Delta Ct$ values was calculated to serve as error bar.

Luciferase activity experiment.

Two luciferase-reporter plasmids were constructed, one carrying non-edited *miR-9-5p* target region (LEP_{WT-UTR}), another carrying S6 and S7 edited *miR-9-5p* target region of *LEP* ($LEP_{MUT2-UTR}$). The day before transfection, HTR-8/SVneo cells were plated at $1 \times 5,000$ cells per well in 96-well plates. Then reporter plasmids were co-transfected with *miR-9-5p* mimic or negative control (NC) into HTR-8/SVneo cell line and cultured in 96-well plates using TurboFect (Thermo Fisher Scientific, USA) according to the manufacturer's instruction. In each well, the cells were transfected with 100 ng of the tested constructs and 20 ng of pRL-TK. The activities of Renilla luciferase and Firefly luciferase were determined using the Dual-Luciferase Reporter Assay System (Promega, USA) 48 hours after transfection. Luminescence was detected by a GloMax 20/20 Luminometer (Promega, USA) according to the manufacturer's instructions. The error bars represent standard errors.

Data Access

The authors are committed to the release of data and analysis results, with the understanding that rapid and transparent data sharing will benefit the scientific research community. Detailed analyzed data are included in supplementary excel tables. RNA-seq data are going to be deposited at GEO and will be available. All renewable reagents and detailed protocols will be available upon request.

Acknowledgements

The work was supported by the National Natural Science Foundation of China grant No. 81571097 (X.Y.) and grant No. 81671466 (M.Z.), the Natural Science Foundation of Guangdong Province Grant No. 2016A030308020 (X.Y.), Science and Technology Projects of Guangzhou grant No. 201704020116 (X.Y.), and the Science and Technology Planning Project of Guangdong Province grant No. 2015B050501006 (M.Z.). The funding organizations had no role in design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript.

Author contributions:

Xinping Y. conceived the project; Xinping Y., M.Z. and Y.Y. supervised the project. Xiaoxue Y. performed computational analysis with contributions from H.L., Y.G., and Z.R.; J.Y., X.L. Q.C., S.J. performed the experiments. Xiaoxue Y. and Xinping Y. wrote the paper.

Disclosure Declaration

The authors declare no competing interests.

References

Agranat L, Raitskin O, Sperling J, Sperling R. 2008. The editing enzyme ADAR1 and the mRNA surveillance protein hUpf1 interact in the cell nucleus. *Proceedings of the National Academy of Sciences of the United States of America* **105**(13): 5028-5033.

Anant S, Davidson NO. 2001. Molecular mechanisms of apolipoprotein B mRNA editing. *Current opinion in lipidology* **12**(2): 159-165.

Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**(2): 166-169.

Athanasiadis A, Rich A, Maas S. 2004. Widespread A-to-I RNA editing of Alu-containing mRNAs in the human transcriptome. *PLoS biology* **2**(12): e391.

Bass BL, Weintraub H. 1987. A developmentally regulated activity that unwinds RNA duplexes. *Cell* **48**(4): 607-613.

Bazak L, Haviv A, Barak M, Jacob-Hirsch J, Deng P, Zhang R, Isaacs FJ, Rechavi G, Li JB, Eisenberg E et al. 2014. A-to-I RNA editing occurs at over a hundred million genomic sites, located in a majority of human genes. *Genome research* **24**(3): 365-376.

Belgrader P, Cheng J, Zhou X, Stephenson LS, Maquat LE. 1994. Mammalian nonsense codons can be cis effectors of nuclear mRNA half-life. *Molecular and cellular biology* **14**(12): 8219-8228.

Benne R. 1994. RNA editing in trypanosomes. *European journal of biochemistry* **221**(1): 9-23.

Benne R, Van den Burg J, Brakenhoff JP, Sloof P, Van Boom JH, Tromp MC. 1986. Major transcript of the frameshifted coxII gene from trypanosome mitochondria contains four nucleotides that are not encoded in the DNA. *Cell* **46**(6): 819-826.

Blanc V, Davidson NO. 2003. C-to-U RNA editing: mechanisms leading to genetic diversity. *The Journal of biological chemistry* **278**(3): 1395-1398.

Blow M, Futreal PA, Wooster R, Stratton MR. 2004. A survey of RNA editing in human brain. *Genome research* **14**(12): 2379-2387.

Buckley PT, Lee MT, Sul JY, Miyashiro KY, Bell TJ, Fisher SA, Kim J, Eberwine J. 2011. Cytoplasmic intron sequence-retaining transcripts can be dendritically targeted via ID element retrotransposons. *Neuron* **69**(5): 877-884.

Chaiworapongsa T, Chaemsaithong P, Yeo L, Romero R. 2014. Pre-eclampsia part 1: current understanding of its pathophysiology. *Nature reviews Nephrology* **10**(8): 466-480.

Chen SH, Habib G, Yang CY, Gu ZW, Lee BR, Weng SA, Silberman SR, Cai SJ, Deslypere JP, Rosseneu M et al. 1987. Apolipoprotein B-48 is the product of a messenger RNA with an organ-specific in-frame stop codon. *Science* **238**(4825): 363-366.

Delva E, Camus Y, Paugam C, Parc R, Huguet C, Lienhart A. 1987. Hemodynamic effects of portal triad clamping in humans. *Anesthesia and analgesia* **66**(9): 864-868.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**(1): 15-21.

Dominissini D, Moshitch-Moshkovitz S, Amariglio N, Rechavi G. 2011. Adenosine-to-inosine RNA editing meets cancer. *Carcinogenesis* **32**(11): 1569-1577.

Eisenberg E, Levanon EY. 2018. A-to-I RNA editing - immune protector and transcriptome diversifier. *Nature reviews Genetics* **19**(8): 473-490.

Enquobahrie DA, Meller M, Rice K, Psaty BM, Siscovick DS, Williams MA. 2008. Differential placental gene expression in preeclampsia. *American journal of obstetrics and gynecology* **199**(5): 566 e561-511.

Faxen M, Nasiell J, Blanck A, Nisell H, Lunell NO. 1998. Altered mRNA expression pattern of placental epidermal growth factor receptor (EGFR) in pregnancies complicated by preeclampsia and/or intrauterine growth retardation. *American journal of perinatology* **15**(1): 9-13.

Fumagalli D, Gacquer D, Rothe F, Lefort A, Libert F, Brown D, Kheddoumi N, Shlien A, Konopka T, Salgado R et al. 2015. Principles Governing A-to-I RNA Editing in the Breast Cancer Transcriptome. *Cell reports* **13**(2): 277-289.

Gong J, Liu C, Liu W, Xiang Y, Diao L, Guo AY, Han L. 2017. LNCediting: a database for functional effects of RNA editing in lncRNAs. *Nucleic acids research* **45**(D1): D79-D84.

Han L, Diao L, Yu S, Xu X, Li J, Zhang R, Yang Y, Werner HMJ, Eterovic AK, Yuan Y et al. 2015. The Genomic Landscape and Clinical Relevance of A-to-I RNA Editing in Human Cancers. *Cancer cell* **28**(4): 515-528.

Hasan S, Robbani SM, Afroze T, Ahsan GU, Hossain MM. 2019. Profiling Gene Specific Changes to RNA Editome in Preeclampsia. *bioRxiv*. 584672.

Holtan SG, Creedon DJ, Haluska P, Markovic SN. 2009. Cancer and pregnancy: parallels in growth, invasion, and immune modulation and implications for cancer therapeutic agents. *Mayo Clinic proceedings* **84**(11): 985-1000.

Hornsby PJ, Maghsoudlou SS, Cheng V, Cheng CY. 1989. Cyclic AMP-mediated cytoskeletal effects in adrenal cells are modified by serum, insulin, insulin-like growth factor-I, and an antibody against urokinase plasminogen activator. *Molecular and cellular endocrinology* **67**(2-3): 185-193.

Hsiao YE, Bahn JH, Yang Y, Lin X, Tran S, Yang EW, Quinones-Valdez G, Xiao X. 2018. RNA editing in nascent RNA affects pre-mRNA splicing. *Genome research* **28**(6): 812-823.

Huntley MA, Lou M, Goldstein LD, Lawrence M, Dijkgraaf GJ, Kaminker JS, Gentleman R. 2016. Complex regulation of ADAR-mediated RNA-editing across tissues. *BMC genomics* **17**: 61.

Hwang T, Park CK, Leung AK, Gao Y, Hyde TM, Kleinman JE, Rajpurohit A, Tao R, Shin JH, Weinberger DR. 2016. Dynamic regulation of RNA editing in human brain development and disease. *Nature neuroscience* **19**(8): 1093-1099.

Ivanov A, Memczak S, Wyler E, Torti F, Porath HT, Orejuela MR, Piechotta M, Levanon EY, Landthaler M, Dieterich C et al. 2015. Analysis of intron sequences reveals hallmarks of circular RNA biogenesis in animals. *Cell reports* **10**(2): 170-177.

Jebbink J, Keijser R, Veenboer G, van der Post J, Ris-Stalpers C, Afink G. 2011. Expression of placental FLT1 transcript variants relates to both gestational hypertensive disease and fetal growth. *Hypertension* **58**(1): 70-76.

Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, Kent WJ. 2004. The UCSC Table Browser data retrieval tool. *Nucleic acids research* **32**(Database issue): D493-496.

Kawahara Y, Zinshteyn B, Sethupathy P, Iizasa H, Hatzigeorgiou AG, Nishikura K. 2007. Redirection of silencing targets by adenosine-to-inosine editing of miRNAs. *Science* **315**(5815): 1137-1140.

Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. 2013. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome biology* **14**(4): R36.

Kim YK, Furic L, Desgroseillers L, Maquat LE. 2005. Mammalian Staufen1 recruits Upf1 to specific mRNA 3'UTRs so as to elicit mRNA decay. *Cell* **120**(2): 195-208.

Kiran A, Baranov PV. 2010. DARNED: a DAtabase of RNa EDiting in humans. *Bioinformatics* **26**(14): 1772-1776.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature methods* **9**(4): 357-359.

Laurencikiene J, Kallman AM, Fong N, Bentley DL, Ohman M. 2006. RNA editing and alternative splicing: the importance of co-transcriptional coordination. *EMBO reports* **7**(3): 303-307.

Levanon EY, Eisenberg E, Yelin R, Nemzer S, Hallegger M, Shemesh R, Fligelman ZY, Shoshan A, Pollock SR, Sztybel D et al. 2004. Systematic identification of abundant A-to-I editing sites in the human transcriptome. *Nature biotechnology* **22**(8): 1001-1005.

Li T, Wernersson R, Hansen RB, Horn H, Mercer J, Slodkowicz G, Workman CT, Rigina O, Rapacki K, Staerfeldt HH et al. 2017. A scored human protein-protein interaction network to catalyze genomic interpretation. *Nature methods* **14**(1): 61-64.

Liu F, Li B, Feng X, Wei YG, Li Y. 2009. [Analysis of the post-operative pulmonary infection in adult-to-adult living donor liver transplant recipients]. *Zhonghua gan zang bing za zhi = Zhonghua ganzangbing zazhi = Chinese journal of hepatology* **17**(8): 611-614.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology* **15**(12): 550.

Maas S, Kawahara Y, Tamburro KM, Nishikura K. 2006. A-to-I RNA editing and human disease. *RNA biology* **3**(1): 1-9.

Mannion NM, Greenwood SM, Young R, Cox S, Brindle J, Read D, Nellaker C, Vesely C, Ponting CP, McLaughlin PJ et al. 2014. The RNA-editing enzyme ADAR1 controls innate immune responses to RNA. *Cell reports* **9**(4): 1482-1494.

McGinnis R, Steinthorsdottir V, Williams NO, Thorleifsson G, Shooter S, Hjartardottir S, Bumpstead S, Stefansdottir L, Hildyard L, Sigurdsson JK et al. 2017. Variants in the fetal genome near FLT1 are associated with risk of preeclampsia. *Nature genetics* **49**(8): 1255-1260.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel S, Daly M et al. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome research* **20**(9): 1297-1303.

Middleton R, Gao D, Thomas A, Singh B, Au A, Wong JJ, Bomane A, Cosson B, Eyras E, Rasko JE et al. 2017. IRFinder: assessing the impact of intron retention on mammalian gene expression. *Genome biology* **18**(1): 51.

Mukhopadhyay D, Anant S, Lee RM, Kennedy S, Viskochil D, Davidson NO. 2002. C-->U editing of neurofibromatosis 1 mRNA occurs in tumors that express both the type II transcript and apobec-1, the catalytic subunit of the apolipoprotein B mRNA-editing enzyme. *American journal of human genetics* **70**(1): 38-50.

Nakano M, Fukami T, Gotoh S, Nakajima M. 2017. A-to-I RNA Editing Up-regulates Human Dihydrofolate Reductase in Breast Cancer. *The Journal of biological chemistry* **292**(12): 4873-4884.

Nakano M, Fukami T, Gotoh S, Takamiya M, Aoki Y, Nakajima M. 2016. RNA Editing Modulates Human Hepatic Aryl Hydrocarbon Receptor Expression by Creating MicroRNA Recognition Sequence. *The Journal of biological chemistry* **291**(2): 894-903.

Nishikura K. 2010. Functions and regulation of RNA editing by ADAR deaminases. *Annual review of biochemistry* **79**: 321-349.

-. 2016. A-to-I editing of coding and non-coding RNAs by ADARs. *Nature reviews Molecular cell biology* **17**(2): 83-96.

Park E, Williams B, Wold BJ, Mortazavi A. 2012. RNA editing in the human ENCODE RNA-seq data. *Genome research* **22**(9): 1626-1633.

Paz-Yaacov N, Bazak L, Buchumenski I, Porath HT, Danan-Gotthold M, Knisbacher BA, Eisenberg E, Levanon EY. 2015. Elevated RNA Editing Activity Is a Major Contributor to Transcriptomic Diversity in Tumors. *Cell reports* **13**(2): 267-276.

Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. 2016. Transcript-level expression analysis of RNA-seq

experiments with HISAT, StringTie and Ballgown. *Nature protocols* **11**(9): 1650-1667.

Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature biotechnology* **33**(3): 290-295.

Picardi E, D'Erchia AM, Lo Giudice C, Pesole G. 2017. REDIportal: a comprehensive database of A-to-I RNA editing events in humans. *Nucleic acids research* **45**(D1): D750-D757.

Picardi E, Manzari C, Mastropasqua F, Aiello I, D'Erchia AM, Pesole G. 2015. Profiling RNA editing in human tissues: towards the inosinome Atlas. *Scientific reports* **5**: 14941.

Polson AG, Crain PF, Pomerantz SC, McCloskey JA, Bass BL. 1991. The mechanism of adenosine to inosine conversion by the double-stranded RNA unwinding/modifying activity: a high-performance liquid chromatography-mass spectrometry analysis. *Biochemistry* **30**(49): 11507-11514.

Powell LM, Wallis SC, Pease RJ, Edwards YH, Knott TJ, Scott J. 1987. A novel form of tissue-specific RNA processing produces apolipoprotein-B48 in intestine. *Cell* **50**(6): 831-840.

Ramaswami G, Li JB. 2014. RADAR: a rigorously annotated database of A-to-I RNA editing. *Nucleic acids research* **42**(Database issue): D109-113.

Redman CW, Sargent IL. 2005. Latest advances in understanding preeclampsia. *Science* **308**(5728): 1592-1594.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**(1): 139-140.

Rueter SM, Dawson TR, Emeson RB. 1999. Regulation of alternative splicing by RNA editing. *Nature* **399**(6731): 75-80.

Sagredo EA, Blanco A, Sagredo AI, Perez P, Sepulveda-Hermosilla G, Morales F, Muller B, Verdugo R, Marcelain K, Harismendy O et al. 2018. ADAR1-mediated RNA-editing of 3'UTRs in breast cancer. *Biological research* **51**(1): 36.

Savva YA, Rieder LE, Reenan RA. 2012. The ADAR protein family. *Genome biology* **13**(12): 252.

Schafer S, Miao K, Benson CC, Heinig M, Cook SA, Hubner N. 2015. Alternative Splicing Signatures in RNA-seq Data: Percent Spliced in (PSI). *Current protocols in human genetics* **87**: 11 16 11-14.

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T. 2003. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome research* **13**(11): 2498-2504.

Shi Y, Yang F, Wei S, Xu G. 2017. Identification of Key Genes Affecting Results of Hyperthermia in Osteosarcoma Based on Integrative ChIP-Seq/TargetScan Analysis. *Medical science monitor : international medical journal of experimental and clinical research* **23**: 2042-2048.

Simpson L, Aphasizhev R, Gao G, Kang X. 2004. Mitochondrial proteins and complexes in Leishmania and Trypanosoma involved in U-insertion/deletion RNA editing. *Rna* **10**(2): 159-170.

Singh M. 2012. Dysregulated A to I RNA editing and non-coding RNAs in neurodegeneration. *Frontiers in genetics* **3**: 326.

Skuse GR, Cappione AJ, Sowden M, Metheny LJ, Smith HC. 1996. The neurofibromatosis type I messenger RNA undergoes base-modification RNA editing. *Nucleic acids research* **24**(3): 478-485.

Sommer B, Kohler M, Sprengel R, Seeburg PH. 1991. RNA editing in brain controls a determinant of ion flow in glutamate-gated channels. *Cell* **67**(1): 11-19.

Teppa RJ, Ness RB, Crombleholme WR, Roberts JM. 2000. Free leptin is increased in normal pregnancy and further increased in preeclampsia. *Metabolism: clinical and experimental* **49**(8): 1043-1048.

Thomas CP, Andrews JI, Liu KZ. 2007. Intronic polyadenylation signal sequences and alternate splicing generate human soluble Flt1 variants and regulate the abundance of soluble Flt1 in the placenta. *FASEB journal :*

official publication of the Federation of American Societies for Experimental Biology **21**(14): 3885-3895.

Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ, Pachter L. 2010. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nature biotechnology* **28**(5): 511-515.

Wang K, Li M, Hakonarson H. 2010. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic acids research* **38**(16): e164.

Wang Q, Hui H, Guo Z, Zhang W, Hu Y, He T, Tai Y, Peng P, Wang L. 2013. ADAR1 regulates ARHGAP26 gene expression through RNA editing by disrupting miR-30b-3p and miR-573 binding. *Rna* **19**(11): 1525-1536.

Wang Q, Miyakoda M, Yang W, Khillan J, Stachura DL, Weiss MJ, Nishikura K. 2004. Stress-induced apoptosis associated with null mutation of ADAR1 RNA editing deaminase gene. *The Journal of biological chemistry* **279**(6): 4952-4961.

Xu X, Wang Y, Liang H. 2018. The role of A-to-I RNA editing in cancer development. *Current opinion in genetics & development* **48**: 51-56.

Yamanaka S, Poksay KS, Arnold KS, Innerarity TL. 1997. A novel translational repressor mRNA is edited extensively in livers containing tumors caused by the transgene expression of the apoB mRNA-editing enzyme. *Genes & development* **11**(3): 321-333.

Yang CC, Chen YT, Chang YF, Liu H, Kuo YP, Shih CT, Liao WC, Chen HW, Tsai WS, Tan BC. 2017. ADAR1-mediated 3' UTR editing and expression control of antiapoptosis genes fine-tunes cellular apoptosis response. *Cell death & disease* **8**(5): e2833.

Yeo J, Goodman RA, Schirle NT, David SS, Beal PA. 2010. RNA editing changes the lesion specificity for the DNA repair enzyme NEIL1. *Proceedings of the National Academy of Sciences of the United States of America* **107**(48): 20715-20719.

Yu G, Wang LG, Han Y, He QY. 2012. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics : a journal of integrative biology* **16**(5): 284-287.

Zhou L, Qiao F. 2006. Expression of RhoA in placenta of preeclampsia. *Journal of Huazhong University of Science and Technology Medical sciences = Hua zhong ke ji da xue xue bao Yi xue Ying De wen ban = Huazhong keji daxue xuebao Yixue Yingdewen ban* **26**(6): 744-746.