1

The Intrinsically Disordered Region of ExbD is Required for Signal Transduction.

Dale R. Kopp[1] and Kathleen Postle[*]

Running title: *ExbD-TonB interactions*

Keywords: TonB, ExbD, intrinsic disorder, binding motif, energy transduction

[1]Department of Biochemistry and Molecular Biology, The Pennsylvania State University, University

Park, PA, 16802, USA

* Correspondence:

Kathleen Postle, Department of Biochemistry and Molecular Biology, The Pennsylvania State

University, University Park, PA 16802, USA. email: postle@psu.edu, Ph. 1-814-863-7568, Fx. 1-814-

863-7024

ABSTRACT:

In most Gram-negative bacteria, the TonB system is required to actively transport vital nutrients across their unenergized outer-membrane barriers. The cytoplasmic membrane proteins of the system ExbB, ExbD, and TonB work together to transduce the energy from the protonmotive force (PMF) of the inner membrane to the ligand-specific, outer membrane TonB-dependent transporters. However, the precise coordination and interactions between ExbB, ExbD, and TonB during energy transduction is unclear. Previously, deletions within a periplasmic disordered domain of ExbD had been shown to be important for TonB system activity and the essential ExbD-TonB PMF-dependent interaction. In this study, we have discovered a conserved motif, ΨXΨXLP (Ψ = hydrophobic-branched residues; X = non-hydrophobic residues) specifically within this domain that was required for both TonB system function and the ExbD-TonB PMF-dependent interaction. Alanine scanning mutagenesis found that the only functionally important residues within the ExbD disordered domain were those located in the motif. In addition, *in vivo* photo-cross-linking captured and identified five ExbD complexes from pBpa substitutions within the ExbD disordered domain: 2-ExbB-ExbD complexes, the ExbD homodimer, the PMF-dependent ExbD-TonB, and the first captured, ExbD-TonB *in*dependent interaction. Interestingly, many of these complexes were captured from single pBpa substitutions, which suggests that these residues are involved in multiple interactions. The data presented in this study indicate that the ExbD disordered domain is a highly dynamic region *in vivo* and a conserved motif within this domain is required for the TonB system to respond to protonmotive force.

Importance:

Within the context of pathogenicity, the TonB system is a virulence factor in many Gram-negative pathogens including E-S-K-A-P-E pathogenic species *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *and Pseudomonas aeruginosa*. Because the TonB system is unique to Gram-negative bacteria and is a periplasmically localized virulence factor, it is an appealing target for novel antibiotics. However, there is no current antibiotic against this system. Understanding the mechanism by which the TonB system functions will provide valuable information to design potential inhibitors targeting the system.

INTRODUCTION:

The TonB system is required for most Gram-negative bacteria to acquire vital nutrients such iron, vitamin B_{12}, nickel, cobalt, copper, heme, maltodextrin, and sucrose across their outer-membrane barriers (1), and is major virulence factor in many pathogenic Gram-negative bacteria (2, 3). Under iron-limiting conditions, such as the human serum (3), Gram-negative bacteria secrete iron-chelating compounds called siderophores that capture the iron with high affinity. However, because iron-siderophore complexes are too large, too scare, and too important to passively diffuse through the outer-membrane porins, they require customized 22-stranded beta-barrel transporters called TonB-dependent transporters (TBDTs) to transition across the outer-membrane (1, 4, 5). After the siderophore captures an iron atom extracellularly, the iron-siderophore complex binds to a TDBT with sub-nanomolar affinity, causing the ligand-loaded TBDT to signal on its periplasmic face that ligand is bound (6–9). Energy is required to release the iron-siderophore from the TBDT (1–3). Because the outer-membrane and periplasmic space lack a sufficient energy source, Gram-negative bacteria transduce the energy from the cytoplasmic membrane proton motive force (PMF) to the TBDTs using the three integral cytoplasmic membrane proteins of the TonB system--ExbB, ExbD, and TonB (12, 13).

In *Escherichia coli K-12*, there are six TBDTs that transport various sources of iron and one TBDT that transports vitamin B12 (4). With a cellular ratio of 7 ExbB: 2 ExbD: 1 TonB, ExbB and ExbD through an unclear mechanism harvest the PMF and transmit it to TonB (14). ExbB has three transmembrane domains (TMDs) with majority of the protein residing in the cytoplasm [Fig. 1, (15, 16)]. In contrast, ExbD and TonB are anchored by single N-terminal TMDs with the bulk of their residues occupying the periplasm (17, 18). ExbB, ExbD, and TonB also share homology through their TMDs with proteins TolQ, TolR, and TolA of the Tol system, which is necessary for outer membrane integrity and important for cell division (19–21). Both

systems are energized by PMF, suggesting that they have a common means of PMF utilization

that plays out as different roles in cell physiology due to different protein contacts made by

their soluble domains (22, 23).

Because the TBDTs greatly outnumber the molecules of TonB per cell (14) and there is

competition between TBDTs for "energized" TonB (24), TonB must undergo a cyclic pattern of

binding and release from TBDT to TBDT (8, 25). This pattern is referred to as the TonB

energization cycle (26). TonB binds to TBDTs both *in vitro* and *in vivo* whether or not it is

"energized", and therefore, it must be correctly configured during an energy transduction cycle

to enable productive interaction with the TBDT (6, 27–29). Both ExbB and ExbD are required

to correctly configure TonB to energize active transport across the outer membrane--ExbB

indirectly, and ExbD directly (16, 30, 31).

ExbB serves as a scaffolding protein for ExbD and TonB assembly, without which they

are both proteolytically unstable, and mediates ExbD-TonB interactions (16, 30–33). The

ExbD-TonB interactions, or lack thereof, define our model for the initial Stages of the TonB

energization cycle (26, 34). In Stage I, ExbD and TonB homodimers exist in separate

complexes with ExbB, but do not interact (26). To transition to Stage II, ExbB facilitates

bringing ExbD and TonB homodimers together in a single complex to form PMF-independent

ExbD-TonB interactions. The PMF-independent ExbD-TonB interaction has only been

detected indirectly by the protection from ExbD of the N-terminal ~155 residues of TonB in the

of proteinase K. A proteinase K-resistant conformation of TonB exists in spheroplasts that

requires a functional ExbD protein and accumulates in the absence of PMF (34, 35, 16). In the

presence of PMF, the ExbD and TonB homodimers swap partners to form PMF-dependent

ExbD-TonB heterodimers which is identified as Stage III. In this Stage, the PMF-dependent

ExbD-TonB interaction can be captured with formaldehyde *in vivo* (31). The only TMD among

the cytoplasmic membrane proteins of the TonB system that can participate in response to PMF is ExbD, with its D25 residue (16, 31, 36).

ExbD has five structurally distinct domains: a cytoplasmically localized N-terminus (residues 1-22), a predicted TMD (residues 23-43), a disordered periplasmic domain (44-63), a structured periplasmic domain (residues 64-133) and a flexible C-terminus (residues 134-141) (37). Experiments using *in vivo* disulfide cross-linking of engineered Cys residues in the ExbD and TonB distal periplasmic domains are consistent with the three-Stage model. The same ExbD residues in the region of 92-121 that form a homodimeric interface are almost the exact same residues that form ExbD-TonB heterodimers through the TonB periplasmic domain (38). Also, in a PMF-unresponsive ExbD mutant, ExbD (D25N), ExbD-TonB disulfide cross-links are either absent or significantly reduced (39).

A 10 residue-deletion scanning mutagenesis of ExbD revealed that deletions within ExbD periplasmic residues 62-141 stall TonB energization in Stage I, whereas deletions largely encompassing the disordered domain of ExbD (42-61) stall TonB energization in Stage II, unable to proceed to Stage III. However, it was unclear how the disordered domain of ExbD participated in energy transduction.

Here we discovered that a novel periplasmic motif, ΨXΨXLP (Ψ = hydrophobic-branched residues; X = non-hydrophobic residues) located very near the ExbD TMD, is essential for ExbD activity. This motif within the disordered domain of ExbD is required for transition from Stage II to Stage III and may account for crosstalk between the TonB and Tol systems because it also exists in TolR. More broadly, the disordered domain makes contacts with another ExbD and with ExbB through two different sites or conformations. Importantly, the disordered domain also forms the long-predicted PMF-*in*dependent complex between ExbD and TonB, visualized for the first time without requiring collapse of PMF, indicating it is part of

the normal energy transduction cycle. It also forms the PMF-dependent complex previously

visualized by formaldehyde cross-linking. This study thus identifies for the first time, at least

one of the ExbD domains and its contacts involved in TonB-dependent energy transduction.

RESULTS:

Within the ExbD disordered domain, residues 44-53 are essential for TonB system

activity.

In our previous study, deletions that encompassed most of the disordered domain of

ExbD protein, (Δ42-51) and (Δ52-61), were completely inactive because they eliminated the

ability of its partner protein, TonB, to undergo essential conformational changes in response to

change in PMF (35, 37). The effects of these deletions could have been due to structural

distortions. To circumvent that issue, here we engineered and characterized ExbD with blocks

of alanine substitutions to identify functionally important residues within the disordered domain

(Fig. 1B): ExbD (44-48ala), ExbD (49-53ala), ExbD (54-58ala), and ExbD (59-63ala). TonB

system activity was determined by measuring the initial transport rates of two different iron-

siderophores, ^{55}Fe-enterochelin and ^{55}Fe-ferrichrome (6, 40), to identify any TonB-dependent

transporter-specific activities.

While both ExbD (54-58ala) and ExbD (59-63ala) transported ^{55}Fe-enterochelin or ^{55}Fe-

ferrichrome at rates equivalent to the wild-type ExbD encoded on pKP999 (pExbD), neither

ExbD (44-48ala) nor ExbD (49-53ala) were able to transport either siderophore. ExbD (59-

63ala) required significantly more sodium propionate (20 mM) to achieve chromosomal ExbD

levels, suggesting that it was important for proteolytic stability (Fig. 2). These results excluded

residues 54-63 as functionally important for the TonB system and eliminated the possibility of

siderophore-specific effects.

Previously, ExbD (Δ42-51) and ExbD (Δ52-61) were unable to form the formaldehyde-cross-linked ExbD-TonB complex that typifies Stage III (35). Consistent with that, the inactive ExbD (44-48ala) and ExbD (49-53ala) had the same cross-linking profile in the presence and absence of TonB, whereas for the active ExbD (54-58ala), and ExbD (59-63ala), the absence of TonB led to the absence of the formaldehyde cross-linked ExbD-TonB complex (Fig. S1).

To test whether the absence of the PMF-dependent ExbD-TonB complex was due to the ability of ExbD (44-48ala) and ExbD (49-53ala) to somehow induce proton leakage, we measured the rate of PMF-dependent ethidium bromide efflux through the AcrA/AcrB/TolC complex (41, 42). None of the block ala mutants came close to PMF leakage caused by 15 μM CCCP, which nonetheless still supports 100% ^{55}Fe-ferrichrome transport (data not shown) (Fig. S2A). Because PMF was adequate for ^{55}Fe-ferrichrome transport, the reason for the inactivity of the block ala substitutions must lie elsewhere.

Expression of ExbD (44-48ala) did, however, cause a small but statistically significant increase in ethidium bromide accumulation compared to pExbD (Fig. S2B). The means by which ExbD (44-48ala) caused this low level of proton leakage was unclear, especially since the Δ*exbD* strain with *tolQR* present did not detectably leak protons (data not shown). Because there is crosstalk between the TonB system and the Tol-Pal system (43), and because RA1045 (Δ*exbD*, Δ*tolQR*) showed the same degree of proton leakage as ExbD (44-48ala), we speculate that protons could possibly have been leaking through TolA in the absence of TolQR or a functional ExbBD.

***In vivo* photo-cross-linking identifies characteristic PMF-dependent and novel PMF-independent ExbD-TonB complexes.**

Disordered domains of proteins are often regions of interaction with other proteins. In the case of ExbD and TonB, both of which contain regions of disorder, we know that they

formaldehyde-cross-link, but the residues involved in either protein are unknown. To identify

contacts made by the ExbD disordered domain, individual amber mutations were constructed

at codons 44-63 for *in vivo* photo-cross-linking. For *in vivo* photo-cross-linking, the

photoreactive, non-natural amino acid, *p*-benzoyl-L-phenylalanine (*p*Bpa) is incorporated at the

site of an amber mutation and can covalently capture protein interactions within 3 angstroms

upon exposure to ultraviolet (UV) light (44). The advantage of this approach over formaldehyde

cross-linking is that interactions through defined sites can be investigated, while leaving the

positions of residues in binding partners unconstrained.

To characterize ExbD-TonB interactions made through the ExbD disordered domain, we

chose ExbD (S54pBpa) and ExbD (R59pBpa) as representatives. Both variants were active in

iron transport assays and shared representative complexes with the rest of the pBpa

substitution mutants (shown below). On immunoblots developed with anti-TonB monoclonal

antibodies, two different ExbD-TonB complexes were captured by both variants, a doublet at

~55 kDa and a single band at ~46 kDa (Fig. 2, lanes 4 and 8). Because a C-terminally added

His6X epitope tag caused each of the complexes to migrate at a higher apparent mass

equivalent to that of the added His6X tag, all the bands consisted of TonB complexed with full-

length ExbD (Fig. 2, lanes 11 and 12). No complexes were detected in a strain where the *tonB*

gene was deleted (Fig. 2, lane 7).

The sole potentially proton-responsive amino acid in the TonB system, Asp 25, resides

in the TMD of ExbD [Fig. 1,(16, 36)]. ExbD (D25N) is a well-characterized inactive mutant

which causes the TonB energization cycle to stall at Stage II and is therefore useful for

differentiating between PMF- dependent and -independent ExbD interactions (45, 31, 34, 38).

Complexes decreased or eliminated by ExbD (S54pBpa) and ExbD (R59pBpa) in concert with

the D25N mutation (Fig. 2, lanes 5 and 9) or by treatment with the protonophore CCCP (Fig. 2,

lanes 6 and 10) identified the 55 kDa doublet as being PMF-dependent. Although the reason

for the doublet is not clear, it could represent two different PMF-dependent sites of contact

made by ExbD (S54pBpa) and ExbD (R59pBpa) with TonB.

In contrast, the abundance of 46 kDa ExbD-TonB complex was undiminished by the

presence of the ExbD D25N mutation or by CCCP treatment, indicating that is was a PMF-

independent complex. If anything, the ExbD D25N mutation generated higher levels of the 46

kDa complex, consistent with the idea that it stalls TonB at Stage II, where such complexes

could build up. These data were also consistent with our finding that in Stage II of TonB

energization, ExbD protects the first ~155 residues of TonB from degradation by proteinase K

in the absence of PMF (34). They also verified that loss of PMF causes a significant and, now

detectable, change in the sites through which ExbD disordered domain residues contact

TonB—enough difference that it results in significant change in apparent mass of the complex

from 55 kDa to 46 kDa.

While we knew that ExbD and TonB made *in vivo* contacts through their extreme C-

termini, many of them beyond the N-terminal ~155 residues of TonB (38, 39), we did not know

that additional ExbD contacts occurred so close to the ExbD (and, presumably, TonB) TMDs

(Fig. 1A,B).

Most of the pBpa substitutions in the ExbD disordered domain capture some form of

TonB contact.

To understand the full extent of interactions between the ExbD disordered domain and

TonB, *in vivo* photo-cross-linking was carried out through pBpa substitutions at ExbD codons

44-63. Most pBpa substitutions in the region from 44-52 supported significantly reduced TonB-

dependent ^{55}Fe transport activity (< 25% wild-type levels), consistent with the results from the

block ala substitutions which essentially eliminated activity (Fig. 4A). Substitutions at residues

53-63 all supported at least 50% activity except for ExbD (R59pBpa) which also supported only

25% activity. Several single pBpa substitution mutants from residues 54-63 were less active

than ExbD (54-58ala) and ExbD (59-63ala) (compare Fig. 2A with Fig. 4A, right panel). The

bulky pBpa side chain was likely responsible for the discrepancy, possibly alluding to the

importance of conformational flexibility or lock and key fit for these residues. Nonetheless,

except for ExbD (V45pBpa) and ExbD (V47pBpa), all the pBpa substitutions within the ExbD

disordered domain had detectable levels of activity. Therefore, the complexes captured by

these substitutions at least indicated TonB proteins within ~3angstroms of the pBpa moiety

(46) and likely reflected interactions that occurred during a TonB energization cycle.

The PMF-dependent ExbD-TonB complexes formed starting at approximately ExbD

residue A51(pBpa) and continuing through P63 (pBpa), with all the pBpa substitutions except

S52(pBpa) participating (Fig. 4B). Consistent with the predicted intrinsically disordered nature

of this region (35, 37), there was not a strong pattern of interaction suggestive of an α-helix or

β-sheet.

We had previously observed a PMF-dependent formaldehyde-cross-linked ExbD-TonB

complex, but did not know through which residues of the ExbD periplasmic domain it occurred

(31). The ExbD disordered domain contains only three formaldehyde-cross-linkable residues:

K46, R59, and K62 (47). Because ExbD (K46pBpa) was unable to capture the PMF-dependent

ExbD-TonB interaction via photo-cross-linking (Fig. 4B, left panel) and ExbD (59-63ala) still

captured the PMF-dependent formaldehyde-cross-linked ExbD-TonB complex (Fig. S1), the

idea that the ExbD disordered domain participated in the previously observed PMF-dependent

formaldehyde cross-link to TonB could be ruled out.

The PMF-independent ExbD-TonB interaction characterized in Fig. 3, was captured by

most of the residues within the ExbD disordered domain (Fig. 4B). Because the abundance of

complex was fainter at either end of the range of pBpa substitutions, it could be that the ExbD

disordered domain constituted the PMF-independent interface. Interestingly, the ExbD

disordered residues that trapped the PMF-independent interactions were similar to an ExbD

interface predicted to interact with TonB by phage-panning experiments (46-62) (48).

In addition to the ExbD-TonB PMF-dependent complex and ExbD-TonB PMF-

independent complex, there was also an unknown ~62 kDa complex trapped via photo-cross-

linking by ExbD (T53pBpa) and ExbD (Q57pBpa) (Fig. 4B). We do not know the identity of the

complex except that it certainly contains both ExbD and TonB.

The ExbD disordered domain interacts with ExbB and another ExbD.

ExbD formaldehyde- cross-links *in vivo* into a single homodimeric complex (~ 30 kDa)

and a single heterodimeric complex with ExbB (~41 kDa) through unknown residues (31, 34).

To determine if the ExbD disordered domain formed such complexes, we used ExbD

(D44pBpa) and (S54pBpa) as examples (Fig. 5A, B). In Fig. 5A, ExbD (D44 pBpa) formed

complexes with apparent masses of ~39 kDa and ~43 kDa, making them both, based on their

apparent masses, candidates to be potential ExbD-ExbB complexes. Surprisingly, the addition

of a His6X-tag caused both complexes to shift to a higher mass consistent with the added

His6X tag, indicating that both were ExbD-ExbB complexes. The presence of two different

ExbD-ExbB complexes indicated that either or both proteins were undergoing conformational

changes that influenced the nature of the contacts between them during the energy

transduction cycle.

In Fig. 5B, ExbD (S54pBpa) formed an ~31 kDa complex. A C-terminal His6X tag

caused the 31 kDa complex to migrate 1.2 kDa more slowly, twice the effect of the His 6X tag

on the apparent mass of the ExbD monomer (Fig. 5B). Thus the ~31 kDa complex was

identified as an ExbD homodimer. ExbD (S54pBpa) also formed the lower of the two ExbD-

286 ExbB complexes and, as expected, the ExbD-TonB PMF-dependent and -independent

287 complexes.

The majority of pBpa substitutions in the ExbD disordered domain capture contacts

with ExbB as well as ExbD.

To understand the full extent of interaction between the ExbD disordered domain and

itself or ExbB, we analyzed *in vivo* photo-cross-linking by the series of ExbD (pBpa)

substitutions by immunoblot with anti-ExbD antibody. As expected, the ExbD-TonB PMF-

dependent and -independent complexes could be discerned, but they were less confidently

identified due to the presence of other complexes in the same region of the immunoblot,

indicating that ExbD interacted with additional proteins, as seen in Fig. 5.

The ExbB-ExbD (upper, ~43 kDa) complex was prominently formed by residues from

the beginning of the ExbD disordered domain D44pBpa until P50pBpa, contacts that placed

the region just distal to the ExbD TMD close to ExbB (Fig.1). It is likely that ExbD pBpa 46-50

lanes (Fig. 6, lanes 5-10) also contain ExbD-TonB PMF-independent complex, which runs

similarly to the ExbB-ExbD (upper, ~43 kDa) complex (compare to Fig. 4 lanes 5-10) and is

therefore difficult to separately identify.

The ~ 39 kDa ExbD-ExbB (lower) complex was formed by pBpa substitutions

throughout the disordered domain from D44 through P63 at similar intensities. This indicates

that a site in ExbD ~20 residues from its predicted periplasmic TMD boundary [residues 32-43,

(15)] binds to ExbB. The fact that ExbB has only a 20 residue periplasmic loop and a 21-

residue periplasmic amino terminus (13), raises the possibility that the ExbD disordered

domain was condensed at some point during the TonB energization cycle such that it was

closer to the inner-membrane. Alternatively, the ExbB periplasmic terminus might extend,

structureless, into the periplasm.

In contrast to the ExbD-TonB complexes, the ExbD-ExbD homodimers were captured

more efficiently in the mutation-intolerant half of the ExbD disordered domain, residues 44-49

and 52-55, without emergence of a pattern indicative of structural elements.

The ExbD disordered domain contains a ΨXΨXLP motif.

A multiple sequence alignment of the ExbD disordered domain was generated and

displayed in a sequence logo using HHblits in the Gremelin.bakerlab program of all sequences

within an expect score of e^{-10} of *Escherichia coli K-12 strain* (NP_417478.1) (Fig. 7A). The

sequence logo identified a ΨXΨXLP motif where the psi symbol (Ψ) represents large aliphatic

side-chained residues, (11) and X represents any non-hydrophobic residue (Fig. 7B). The motif

includes residues V45, V47, L49, and P50, which reside in the mutation-intolerant segment of

the disordered domain and are also conserved in the *Escherichia coli* ExbD paralog, TolR of

the Tol system, but not in paralog MotB of the flagellar motor (Fig. 7B). Due to the similarities

between ExbB/ExbD TMDs of the TonB system and TolQ/TolR TMDs of the Tol system, there

is modest crosstalk between the two systems such that they can substitute functionally for one

another (43, 49). To our knowledge, MotAB has not been shown to account for any crosstalk. Its

conservation in TolR and ExbD suggests the ΨXΨXLP motif may have a shared function in

both proteins, and could account for the cross-talk between ExbB/ExbD and TolQ/R (43, 49, 50).

Only ExbD motif residues V45, V47, L49 and P50 are important for TonB system activity.

Assessment of the activities of the pBpa substitutions in the mutation-intolerant segment

showed that the least active ones [V45(pBpa) and V47(pBpa)], were part of the ΨXΨXLP motif

(Fig. 4A). To better define the motif, individual alanine substitutions were constructed in ExbD

from residues 44 through 53. ExbD variants carrying V45A, V47A. L49A, and P50A mutations

were the only ones with deficits in initial ^{55}Fe-ferrichrome transport rates significantly below

90% of pExbD, with activities of 32%, 21%, 34%, and 50%, respectively (Fig. S3). Thus V45,

V47, L49 and P50 were the only functionally important residues within the entire ExbD

disordered domain.

To further define the motif, each of the V45A, V47A, L49A, and P50A substitutions in

the motif region was combined with the others to create six double ala substitutions. Had these

mutations been simply additive in their effects, we would have seen that, for example, the

V45A, P50A double mutant should have ~16% activity (50% of 32%). Instead, this double

mutant, like all six double mutants, was inactive in ^{55}Fe-ferrichrome transport, suggesting a

degree of synergy between them (Fig. 8A). However, the iron transport assays are not

sensitive to very low levels of TonB system activity (51).

Colicins toxins are synthesized by some *E. coli* to kill sensitive strains. B-group colicins

all use the TonB system to cross the outer membrane of *E. coli* to gain access to their targets

in the periplasm, cytoplasmic membrane or cytoplasm (52) Colicin sensitivity assays, while not

as discriminative among higher levels of TonB system activity, can detect the activity

supported by as little as one molecule of TonB (51). To determine whether the double ala

substitutions could support any activity, their sensitivity to B-group colicins B, Ia, and M was

tested (Fig. 8B). All of the double-ala mutants were tolerant (insensitive) to virtually all colicins

tested, indicating that they do not work independently of each other and validating the

identification of this motif.

ExbD (V45A, V47A), ExbD (V47A, L49A), and ExbD (L49A, P50A) stall TonB at Stage II of

the energy transduction cycle.

ExbD regulates TonB conformational changes as TonB moves through an energy

transduction cycle (34). Stage II of the TonB energization cycle is a PMF-independent Stage

where the N-terminal ~ 155 residues of TonB (an ~23 kDa fragment) are protected from

proteinase K degradation if full-length ExbD is present, although an actual complex between

the two proteins had not been observed until this study (Fig. 5). Heretofore, Stage II had been detectable only when PMF was collapsed, preventing TonB from moving on to Stage III. When PMF, ExbB, ExbD, and TonB are all present and active, TonB transitions from Stage II to Stage III, rendering TonB fully sensitive to proteinase-K. Our observation that ExbD (44-48ala) and ExbD (49-53ala) failed to form a Stage III formaldehyde cross-link suggested that the effects of substitutions in the disordered domain motif might be to stall TonB at Stage II [Fig. S1,(34)]. Since only ExbD residues V45, V47, L49, and P50 were important for ExbD activity, we assessed whether the inactive double mutants (V45A, V47A), (V47A, L49A) and (L49A, P50A) were required for TonB to proceed beyond Stage II based on its sensitivity to exogenously added proteinase K in spheroplasts. For comparison, we assessed ExbD (D25N).

The hallmarks of the ExbD (D25N) mutant, where TonB is permanently stalled at Stage II are 1) that TonB forms a proteinase K-resistant fragment in both the presence and absence of CCCP and 2) that TonB is unable to remain in its stalled Stage II proteinase K-resistant conformation over time [Fig. 8, panel 4 +/- CCCP; (34)]. As expected, the ΔexbD, ΔtolQR strain did not support the ability of TonB to form the ~23 kDa proteinase K-resistant TonB fragment [Fig. 8, panel 3; (34)]. A ~28 kDa TonB fragment, seen previously and typified by ΔexbD, ΔtolQR and ExbD (D25N) in panels 3 and 4, does not appear to have relevance to the mechanism of TonB energization (34).

ExbD (V45A, V47A) and ExbD (V47A, L49A) largely mimicked the effect of ExbD (D25N) on TonB, leaving it stalled in Stage II (Fig. 9). ExbD (L49A, P50A) was too sensitive to proteinase K to be evaluated (Fig. 9 panel 7). Together the results suggest that the ExbD ΨXΨXLP motif was required for TonB to transition from Stage II to Stage III of the energization cycle.

The ExbD disordered domain ΨXΨXLP motif is required for correct configuration of the ExbD C-terminus.

To better understand why the ExbD ΨXΨXLP motif left TonB stalled in Stage II, we combined the V45A, V47A double mutation with pBpa substitutions along the length of ExbD: E14(pBpa)— presumably cytoplasmically localized based on the topology of ExbD; I24(pBpa)—in the TMD; S54(pBpa)—in the mutation-tolerant segment of the disordered domain S67(pBpa)—just beyond the disordered domain K97(pBpa)—a known point of ExbD-TonB contact and E136(pBpa)—near the C-terminus. With the exception of ExbD (I24pBpa), the other substitutions were active [ExbD E14pBpa, S67pBpa, K97pBpa, and E136pBpa substitutions were 72%, 24%, 90%, and 100% of ^{55}Fe-Ferrichrome transport of pKP999 + pEVOL in RA1045 (data not shown)].

Overall, the double V45A, V47A substitutions did not have much effect on ExbD photo-cross-links to TonB through distal pBpa substitutions, except for (S54pBpa) where it greatly diminished a PMF-dependent ExbD-TonB complex (Fig. 10A lanes 6,7). ExbD (E14pBpa), in the cytoplasm, did not form complexes with TonB (Fig. 10A, lanes 4,5), however, ExbD (I24pBpa), in the TMD, formed a PMF-independent TonB complex (10A lanes 6,7), indicating that at some point in its energy transduction cycle, TonB was in close proximity to ExbD through their TMDs. ExbD (K97pBpa) and ExbD (E136pBpa) also formed PMF-independent Stage II complexes with TonB (Fig. 10A, lanes 6,7,12-15). We do not know why the level of monomer was consistently reduced for ExbD (K97pBpa), but it clearly did not prevent capture of ExbD-TonB complexes.

Among the periplasmic pBpa substitutions tested, the presence of the V45A, V47A double substitution caused ExbD to photo-cross-link to a variety of unknown proteins, especially ExbD (S54pBpa) V45A,V47A, which captured at least six different complexes with

unknown proteins (Fig. 10B, lane 24). Furthermore, ExbD (E136pBpa) V45A,V47A captured

two prominent complexes with unknown proteins. These results showed that the disordered

domain of ExbD was important for regulating the ExbD C-terminal conformation and contacts.

The V45A, V47A mutations within ExbD show a protein-wide alteration of ExbD residue

contacts. In many cases, the mutations reduced distal ExbD residue interactions with TonB

and caused additional unknown complex formations. This demonstrates a tight connection

between the ΨXΨXLP motif and the distant ExbD C-terminal residues, and potentially

highlights a protein-wide functional role of the ΨXΨXLP motif in signal transduction.

DISCUSSION:

Understanding the molecular interactions among TonB system proteins in the

cytoplasmic membrane--ExbB, ExbD, TonB--and the TBDTs in the outer membrane is vital to

our understanding its mechanism. TonB-dependent energy transduction is cyclic, with TonB

engaging and disengaging from a TBDT as it energizes active transport across the OM.

Our current model has identified multiple stages that occur before the signal

transduction from the TonB system cytoplasmic membrane proteins to the TBDT is completed

[(26), Figure 12]. In Stage I, TonB and ExbD homodimers are each found in separate

complexes with ExbB homotetramers and they do not interact. In Stage II, the ExbB-TonB and

ExbB-ExbD complexes come together to form a multimeric heterocomplex in which ExbD

protects the first ~155 residues of TonB from digestion by proteinase K in spheroplasts. Prior

to this study, we had not been able to determine precisely through which residues the PMF-

independent Stage II complex forms. Nor had we been able to observe the TonB and ExbD

proteins complexed together in Stage II, although their existence was predicted by the model.

From Stage II the TonB system transitions to Stage III, characterized by a requirement for PMF

and a formaldehyde- cross-linked complex containing both proteins. The final Stage in our

current model is Stage IV in which reconfigured TonB binds the TBDT in such a manner that

active transport across the OM occurs. After Stage IV, the multimeric heterocomplex of ExbB,

ExbD, and TonB breaks down into isolated complexes of ExbB-ExbD, ExbB-TonB, and free-

floating ExbB (26).

The ExbD-TonB, ExbB-ExbD, and ExbD homodimeric complexes identified within the
ExbD disordered domain.

In vivo photo-cross-linking has the advantage of potentially capturing every interaction

within ~3 angstroms that a single residue, as a pBpa substitution, is involved with during the

TonB energization cycle (46). Throughout the ExbD disordered domain (residues 44-63

immediately distal to its TMD), each single ExbD residue captured multiple complexes

suggesting that it was a highly dynamic region. Disordered domains of proteins are known to

have a high degree of conformational plasticity that can dictate changes in protein

oligomerization and function (53). For example, the *E. coli* periplasmic protease/chaperone,

DegP, upon substrate binding to its intrinsically disordered domain, it changes from

proteolytically inactive hexamers to proteolytically active large polyhedrons consisting of 4, 6,

and,8 trimers (54, 55). Most of the ExbD pBpa substitutions within the disordered domain

captured at least four of the five identifiable complexes (ExbD homodimer, two ExbD-ExbB

complexes, and two ExbD-TonB complexes) through a single site. This indicated that residues

within the ExbD disordered domain must have undergone multiple partner swaps during the

TonB energization cycle and demonstrated the minimum number of disordered domain

contacts that occurred.

ExbD-TonB complexes – Residues within the ExbD disordered domain, as pBpa

substitutions, captured both PMF-independent and PMF-dependent ExbD-TonB interactions

using *in vivo* photo-cross-linking. The PMF-independent ExbD-TonB complex was captured by

ExbD disordered residues 45-63. It was the first time that this interaction was captured,

demonstrating the previously predicted existence of Stage II in the TonB energization cycle,

and identified residues through which this complex is formed. Previous residue-specific

disulfide experiments within the ExbD 92-121 residues did not trap the PMF-independent

interaction. It may be because the PMF-independent interaction mostly, albeit not entirely,

occurs with ExbD residues closest to the membrane which would explain why the TonB portion

that is protected by Proteinase K is that closest to its TMD. However, the number of residues

within ExbD disordered domain (20 amino acids) that trap this interaction is disproportional to

that of the periplasmic portion of TonB protected against proteinase K degradation (~123

amino acids). It is possible that the full ExbD periplasmic domain contributes to the protection

of TonB, but the PMF-independent interface is more localized. Consistent with this idea is that

the portion of TonB that is protected by proteinase K is similar to the length of the entire ExbD

periplasmic domain. Furthermore, the PMF-independent ExbD-TonB interaction is also

captured from distal ExbD C-terminal residues. Alternatively, the ExbD disordered domain may

induce a conformational change in the TonB predicted disordered domain (~118 amino acids)

to an ordered conformation, which in turn is protected from proteinase K (56). The idea that

ExbD could possibly influence TonB to transition from a disordered to ordered state has been

previously proposed (23, 57).

PMF-dependent ExbD-TonB complexes were captured from ExbD residue A51 to P63

(except for S52). Although deletions within the ExbD disordered domain confirmed its

importance for forming the ExbD-TonB PMF-dependent interaction, we showed here that the

ExbD disordered domain directly contacted TonB PMF-dependently. Thus, we identified

another ExbD region besides residues 92-121 can interact with TonB PMF-dependently.

However, like the ExbD region of 92-121 cysteine substitutions, alanine substitutions for the

ExbD disordered domain residues that captured the PMF-dependent ExbD-TonB interaction

did not significantly affect TonB system activity. Therefore, these specific residues are

dispensable and are not individually important TonB function. This may suggest that an ExbD

region(s) other than the disordered domain and the 92-121 residue region contains the

functionally important ExbD residues for the PMF-dependent ExbD-TonB interaction. It also

cannot be ruled out that some functionally important ExbD disordered residues (i.e. V45 and

V47) directly interact with TonB PMF-dependently, but due to low activity, the interaction was

not captured.

ExbD-ExbB complexes - Two distinct ExbD-ExbB complexes were captured via photo-

cross-linking: an ~ 39 kDa (lower) complex and an ~43 kDa (upper) complex. Previous

deletions within ExbD periplasmic residues, 92-121, prevented ExbD from formaldehyde cross-

linking to ExbB (35). Therefore, it was possible that a periplasmic ExbD-ExbB interaction *could*

exist. However, disulfide ExbD-homodimers within this region still formaldehyde cross-linked

with ExbB *in vivo*. The disulfide-bonded interface would have likely prevented a potential

switch to a ExbB-ExbD interface and thus, it seemed unlikely that the 92-121 region directly

interacted with ExbB (38). But in this study, ExbD-ExbB periplasmic interactions were identified

throughout the ExbD disordered domain in two distinct complexes.

Previously, only a single ExbD-ExbB complex of ~41 kDa had been identified with

formaldehyde cross-linking instead of two like what was captured in this study with photo-

cross-linking (31). Because formaldehyde cross-linking is non-residue specific, the two ExbB-

ExbD complexes may be distinguished due to the multiple crosslinks within a single complex.

This would potentially explain why the single formaldehyde cross-linked ExbB-ExbD complex

is the average molecular weight of the of the two ExbB-ExbD complexes seen with photo-

cross-linking.

The ExbB-ExbD (lower) complex was captured with similar intensities from all residues

throughout the ExbD disordered domain (~20 residues from its TMD). Because ExbB has very

little of its protein exposed to the periplasm, the ExbD-ExbB (lower) complex may suggest that

either the N-terminal tail of ExbB (~21 amino acid from its 1st TMD) is extended into the

periplasm or that the ExbD disordered domain is parallel to the membrane at some point

during the TonB energization cycle. The latter possibility would be consistent with one of the

three ExbB-ExbD structures isolated and mapped via 3D electron microscopy with ratios 4:2 in

which electron density of the ExbD periplasmic domain parallel with the membrane (58). It

would also be consistent with what was seen in the ExbD homolog, TolR of Tol system, in

which its C-terminus interacts with the ExbB homolog, TolQ (59). However, these correlations

would need to be further examined *in vivo* to determine if they are related observations.

The ExbD-ExbB (upper) complex was captured most strongly from residues

immediately distal to ExbD TMD and significantly waned after the first six N-terminal residues

of the ExbD disordered domain. Considering the presumed location of these residues, it seems

plausible that the ExbB-ExbD (upper) complex may be from where the initial periplasmic

residues of ExbD interacts with ExbB as it extends into the periplasm. This logic would

potentially explain why only ExbD residues nearest to its TMD captured this complex. The

ExbB-ExbD (upper) complex also may be suggestive of an ExbD extended conformation as

seen in a second ExbB-ExbD structure isolated and mapped via 3D electron microscopy (58).

However due to the lack of resolution, the ExbD disordered domain in the isolated ExbB-ExbD

structure from 3D electron microscopy could not be identified.

The addition of CCCP did not prevent ExbD disordered residues from the ExbB-ExbD

(upper) or ExbB-ExbD (lower) photo-cross-linked complexes. Therefore, neither ExbB-ExbD

complex was dependent of PMF. In a previous finding, the degree of Oregon Green® 488

maleimide labelling of cytoplasmic tail mutant TonB (L3C) was dependent its interactions with

the TBDTs. Because ExbB is presumably the only TonB system protein with large portion of its

protein in the cytoplasm, conformational changes within ExbB may influence the maleimide

labelling of the TonB cytoplasmic tail (60). Perhaps, the two photo-cross-linked ExbD-ExbB

complexes in this work are related to ExbB conformational changes that influence the Oregon

Green® 488 maleimide labelling of cytoplasmic tail mutant TonB (L3C). Further analysis would

be needed to determine if either ExbB-ExbD complex is influenced by the TonB-TBDT

interactions.

ExbD-ExbD complexes - The ExbD homodimer (~31 kDa) was also identified via *in vivo*

photo-cross-linking and had a similar molecular weight to the formaldehyde-cross-linked ExbD

homodimer (~30 kDa) (31). Except for ExbD (P50pBpa) or ExbD (A51pBpa), the intensity of

the ExbD homodimer complex gradually decreased with residues more distant from the TMD.

Because the residues that capture ExbD homodimers are immediately distal to its TMD, it

seems likely that the ExbD TMDs would be in close proximity. However this is not suggested in

a recent crystal structure (61).

Most of the ExbD disordered residues that captured ExbD homodimers did not capture

the PMF-dependent ExbD-TonB heterodimers. This indicates that most residues within its

disordered domain that have ExbD homodimeric interfaces do not share one with a PMF-

dependent ExbD-TonB interaction, which would differ from a mostly shared interface identified

within the ExbD 92-121 region (38). This suggests that transition from an ExbD homodimer to

PMF-dependent ExbD-TonB swap may be region specific. Similarly, TonB homodimers remain

active throughout the transduction cycle when they are tethered close to its TMD, but they are

inactive as a homodimer when they are tethered at the C-terminus, presumably because it

prevented the TonB C-terminus to form other necessary interactions (26). The observations in

this work may support the idea that the TonB system protein partner transition is instead likely

partner interface transitions that can occur different steps throughout the TonB energization

cycle.

It is mostly unclear about what enables the transition between the captured photo-cross-

linked ExbD complexes, many from which were captured by a single residue. Other than the

PMF-dependent ExbD-TonB interaction, none of the ExbD complexes were dependent on

PMF. Perhaps the complex transitions are influenced from ExbB, or a hypothesized mystery

protein(s) of the system (16, 42, 60). ExbB also has a predicted disordered region within its

cytoplasmic loop (16) in which deletions within the region cause growth arrest (42). It would be

interesting if ExbB has any role in dictating any of the disordered domain ExbD interaction

transitions or if signaling comes from the TBDT or from ligand-TBDT interaction.

The *in vivo* data as it relates to recent structures

562
Historically, purified structures within the TonB system have been controversial., The

ratio of the components of isolated TonB system complexes have not been internally

consistent [i.e. 4ExbB: 2ExbD and 4ExbB: 1ExbD: 1TonB from single particle EM studies (58,

48); 5ExbB:1ExbD and 6ExbB:3ExbD from crystal structure determinations (61, 62);

6ExbB:1ExbD from mass spectrometry (63); and most recently, 5ExbB: 2ExbD from cryo-EM

(64)], nor consistent with the system's *in vivo* cellular ratio of 7ExbB: 2ExbD: 1TonB (17).

Recently, two crystal structures and a cryo-EM structure have been published: 5 ExbB

with 1 ExbD TMD in the pore (crystal); 6 ExbB: 3 ExbD TMD in the pore (crystal); 5 ExbB: 2

ExbD TMD in the pore (cryo-EM) (61, 62, 64). While the ExbB monomers within the structures

make sense based on what is known about ExbB, the oligomeric structures diverge from what

is known *in vivo* about ExbB oligomers. 1) ExbB forms formaldehyde- cross-linked tetramers

as a dimer of homodimers *in vivo* and cross-links strongly to TonB (30). To date, TonB is

absent from all structures, but a single particle electron microscopy (EM) study, where it was

present in a complex of 4ExbB:1ExbD:1TonB (48). Consistent with that, our *in vivo* photo-

cross-linking data from this study indicated that ExbD (I24pBpa) in the ExbD TMD, captured

the ExbD-TonB PMF-independent complex, likely through direct interaction with the TonB

TMD. 2) Neither crystal nor the cryo-EM structure of ExbB oligomers permits ExbD and TonB

to interact through their transmembrane domains because the lumens of the ExbB pentamers

or hexamers are fully occupied by ExbD TMDs. Thus, TonB is proposed to bind on the

perimeter of the ExbB pentamer or hexamer (61, 62). 3) The TonB TMD interacts with ExbB

TMD1 *in vivo* (16), however, it is clear that TonB moves through its energization cycle stably

homodimerized near and probably through its TMD (26). While TMD1 is indeed on the outside

of the pentamer/hexamer structures, it is not possible to arrange TonB TMDs outside the ExbB

pentamer/hexamer TMDs such that two TonB TMDs can dimerize. 4) Since TonB moves

through its energy transduction cycle homodimerized through its N-terminus (26). TonB on the

periphery of ExbB oligomers results in a ratio of 5-6 ExbB:10-12 TonB, which conflicts with the

known cellular ratio of 7ExbB: 1TonB (17). One way to accommodate TonB homodimers on

the outside and at the known cellular ratios, is for one of the crystal structure subunits to

display asymmetry, such that it differed from the other members of the oligomer and could be

the only one to bind TonB. However, this was not observed.

It is well-established that TolR and MotB, paralogues of ExbD, form functional

homodimers *in vivo* (65, 66). In our study, several functionally active ExbD amber substitutions

immediately distal to the N-terminal ExbD TMD captured ExbD homodimers (but not

homotrimers). In addition, ExbD (I24pBpa) in the ExbD TMD, which is admittedly inactive, also

captured ExbD homodimers but not homotrimers. These data are contrary to the pentameric

ExbB crystal structure with only one ExbD TMD within its lumen and the hexameric ExbB

crystal structure with three ExbD TMDs within its lumen (61, 62). If ExbD homotrimers exist *in*

vivo, they must be asymmetrically arranged within the ExbB hexamer, and still without space

to accommodate contact with the TonB TMD (62). Although the cryo-EM structure had two

ExbD TMDs within its lumen, it also did not have space to accommodate the TonB TMD (64).

The discrepancies between *in vivo* results and *in vitro* ExbB structures raise questions

about the role they play in TonB-dependent energy transduction. Our current model suggests

stages in that process, and in this study, we identified important contacts among ExbD, TonB

and ExbB that occurred during those stages. Perhaps the *in vitro* structures represent as-yet-

unidentified intermediates. Nonetheless the absence of TonB from the ExbB *in vitro* structures,

and the skewed ratios that result when, as the crystal structure papers propose, TonB

homodimers are arranged on the peripheries of the ExbB oligomers, raise questions that need

to be resolved.

The functional role of ΨXΨXLP motif within the ExbD disordered domain

The most important function within ExbD disordered domain is the essential ΨXΨXLP

motif. Double-alanine mutations within this motif were completely inactive of nearly all TonB

system activity and stalled the TonB energization cycle in Stage II. The parallels with the

ΨXΨXLP motif and the PMF-unresponsive ExbD mutant, ExbD (D25N), suggest that the motif

is required for ExbD to respond to PMF. Supporting this idea, ExbD pBpa substitutions with

V45A and V47A substitutions, prevented ExbD (S54pBpa) and ExbD (K97pBpa) from forming

the PMF-dependent ExbD-TonB interaction. However, due to the inactivity of the pBpa

substitutions within this motif, it is difficult to determine if these residues are directly or

indirectly involved in the ExbD-TonB PMF-dependent interaction.

The V45A, V47A mutations also caused additional unknown ExbD complex formations.

Although these complexes are unidentified, it demonstrates that the ΨXΨXLP motif influences

distant ExbD C-terminal residue interactions. The motif may be important for conformational

changes in ExbD C-terminus to form the correct interactions and potentially highlights a

protein-wide functional role of the ΨXΨXLP motif in signal transduction.

The ExbD ΨXΨXLP motif is also conserved its paralog, TolR, but not in a more distant

paralog, MotB of the flagellar MotAB energy harvesting complex. Unlike MotAB, TolQR and

ExbBD complexes can account for detectable levels of TonB system and the Tol system

activity respectively (23, 43). However, the molecular mechanism of how these systems cross-

talk is unclear. Because the motif is important for the TonB system to response to PMF, it

seems likely this motif would be important for the Tol system to respond to PMF. Perhaps, the

conserved motif among both TolR and ExbD may provide molecular insight to how the

complexes cross-talk. It would be interesting if mutations in TolR ΨXΨXLP motif also are

important C-terminal interactions. Maybe the motif and its function are conserved in both ExbD

and TolR but, it triggers different system-specific C-terminal conformational changes.

Nonetheless, the role the ΨXΨXLP motif as it functions in TolR will need to be examined to

verify to the functional similarities.

Similarities and differences between the ExbD homodimers and the homolog, TolR,

dimeric crystal structure

The ExbB, ExbD, and TonB are homologous TolQ, TolR, and TolA of the outer-

membrane integrity complex called the Tol system (43, 50). The homology among each

corresponding protein occurs mostly through their TMDs, however, the ExbD periplasmic

disordered domain ΨXΨXLP motif is also conserved in TolR. Although ExbBD and TolQR

have functional cross-talk, TonB and TolA do not. This an important divergence of function of

each system. But considering that the ΨXΨXLP motif is conserved in both ExbD and TolR, it is

reasonable that the motif function is also conserved. In the ExbD NMR structure, ΨXΨXLP

motif exists as part of the ExbD disordered domain, however, a dimeric crystal structure of

TolR, periplasmic domains shows that a portion of the ΨXΨXLP motif in a beta strand (67).

The structure had a homodimeric interface that included residues 41, 43-47, 49 which

corresponded with the ExbD disordered domain residues 43,45-49, and 51 (Fig. S4). Our work

shows that ExbD homodimers were most abundant through residues 45-49 and 52-55

(residues 43-47 and 50-53 in TolR), which would be somewhat consistent with a similar

dimeric interface in ExbD as in the TolR structure. However, there were some differences

between the two interfaces. TolR (D49) was the final residue at its structural dimeric interface

but the ExbD position equivalent (A51) captured very little ExbD homodimer. Moreover, four

ExbD residues after A51 – S52, T53, S54, and T55 (TolR D49 - A50, T51, E52, and S53) –

captured significant amounts of homodimers, but were implicated as such in the TolR dimeric

structure (67). The conservation between TolR and ExbD wanes after a conserved proline

[ExbD (P50) and TolR (P48)] of the ΨXΨXLP motif until the final ~35 residues of ExbD (68).

Considering the difference between the physical properties of an aspartate and an alanine,

instead of ExbD A51, ExbD polar residues 52-55 may be the ExbD functionally equivalent to

TolR residue D49.

Although a similar dimeric interface as the TolR crystal structure may exist in ExbD, our

data support the Wojdyla et al. conclusions that this structure in ExbD cannot remain static *in*

vivo (67). Using the photo-cross-linking approach, a single residue could trap all contacts

throughout its transition of the TonB energization cycle which better demonstrates the dynamic

behavior ExbD and its signal transduction. We have identified multiple interacting partners with

single ExbD residues within the ExbD disordered domain. This indicates that the ExbD

disordered domain needs to interact with different partners as it proceeds through the TonB

energization cycle and therefore, it cannot remain static in a dimeric conformation.

Proposed function of the motif as it pertains to the Stages during the TonB energization cycle

The ΨXΨXLP motif is required to transition from Stage II to Stage III of the TonB energization cycle. Alanine mutations for the branched hydrophobic residues in the ΨXΨXLP motif are still able to convert TonB into a conformation that produces the TonB proteinase K-resistant fragment indicative of Stage II. Therefore, the hydrophobic-branched residues are not needed for initial association with TonB. The ExbD (S54pBpa) photo-cross-linked ExbD-TonB PMF-independent interaction is unaffected with the V45A and V47A mutations, yet distal ExbD pBpa substitutions, K97 and E136, with the mutations had reduced TonB interactions. This may suggest that TonB is initially recognized by residues within the disordered domain before the function of the motif is required. Perhaps, the initial association with TonB is what signals to the system to assume Stage II. And at this point, the ExbD disordered domain interacts with TonB in a way that either directly protects ~155 residues of TonB or influences TonB to assume Proteinase K, resistant conformation. Then, through an unclear mechanism, the ExbD ΨXΨXLP motif is required to transition the PMF-independent ExbD-TonB interaction in Stage II to a PMF-dependent ExbD-TonB interaction in Stage III.

Summary:

The data show that the ExbD disordered domain is a highly dynamic region and is involved in many system protein-protein interactions. The function of this domain is a highlighted by the ΨXΨXLP motif that is essential for TonB system activity and the ExbD-TonB PMF-dependent interaction. Because the ExbD disordered domain interacted with several partners and contains an essential motif for TonB activity, this would a promising target for a novel antibiotic against the TonB system in *Escherichia coli* and Gram-negative pathogens.

MATERIALS and METHODS:

Strains and plasmids. Strains and plasmids used in these studies are listed in Table S1.

Strain KP1526 was constructed by P1*vir transduction of ΔexbD::cam in RA1021 (W3110,

ΔexbD::cam) into RA1525 (W3110, ΔtolQR, ΔaroB)* [(35)*, RA1021 and RA1525* were kind gifts

from Ray Larsen]. The *tolQR* genes were deleted to eliminate crosstalk (43). Plasmids with

encoded alanine and amber substitution mutations were mostly derived from pKP999.Amber

substitutions in ExbD (V45A, V47A) encoding plasmids were derived from pKP2171. To

determine the identity of potential ExbB complexes, we added a His6X epitope tag to the C-

terminus of ExbB in the vector pKP1657 which carries the ExbBD operon and re-engineered

the ExbD pBpa substitutions on that plasmid. Plasmids pKP2159 and pKP2160 were derived

from pKP2145 and pKP2154, respectively. All plasmids were made using 25-cycle extralong

PCR. 5'-phosphorylated primers were used with the desired mutation encoded in the forward

primer and are available upon request. All mutants were verified by DNA sequencing at the

Penn State Genomics Core Facility, University Park, PA.

Media and growth conditions. All cultures were grown at 37°C with aeration. Saturated

cultures in Luria-Bertani broth were subcultured at 1:75 in M9 minimal media supplemented

with 0.4% glycerol, 0.2% Casamino Acids, 40 µg/ml tryptophan, 4 µg/ml vitamin B1, 1 mM

$MgSO_4$, 0.5 mM $CaCl_2$, and 37 µM $FeCl_3$. All cultures carrying plasmids with *exbD* were in

RA1045 (W3110, Δ*exbD*, Δ*tolQR*) unless noted and grown in 100 µg/mL ampicillin.

Subcultures were induced with sodium propionate (ranging from 0-45 mM) to approximate the

steady state levels of ExbD protein from W3110, the wildtype control (Table S1). All

subcultures were harvested at an A_{550} ranging from 0.40 to 0.60.

In vivo formaldehyde-cross-linking. Culture conditions were as described above. Culture
volumes were adjusted to harvest 0.5 A_{550} -mL of cells. Bacteria were resuspended in sodium
phosphate buffer (pH 6.8) and treated for 15 min at room temperature with 1% formaldehyde
(Electron Microscopy Sciences Cat #15710) as previously described (69). Following
formaldehyde treatment, bacteria were suspended in Laemmli sample buffer (70), with equal
cell numbers loaded in each well of 13% sodium dodecyl sulfate (SDS)-polyacrylamide gels.
ExbD complexes and TonB complexes were detected by immunoblotting with ExbD-specific
polyclonal antibodies and TonB-specific monoclonal antibodies, respectively (14, 71). Equal
loads were confirmed by Coomassie staining of the PVDF membrane used for the immunoblot.

In vivo photo-cross-linking. For _in vivo_ photo-cross-linking, plasmids with amber
substitutions in _exbD_ were co-expressed with pEVOL in RA1045 (W3110, Δ_exbD_, Δ_tolQR_),
which encodes an orthogonal tRNA synthetase and the corresponding orthogonal tRNA$_{CUA}$
that recognizes amber (UAG) codons (72). Cultures were grown to saturation in 100 µg/mL
ampicillin and 34 µg/mL chloramphenicol. Subcultures were grown in minimal M9 medium as
described above. At A_{550} of ~0.2, subcultures were treated with 0.004**%** arabinose (tRNA
synthetase/tRNA pair induction from pEVOL; Table S1), 0.8 mM the photo-cross-linkable
amino acid p-Benzoyl-L-phenylalanine (pBpa) and amounts of propionate required for ExbD
protein expression at near-chromosomal levels (Table S1). 2.0 A_{550}-ml were harvested at A_{550}
~ 0.5, pelleted at room temperature and resuspended in 2 mL supplemented M9 media (as
described above with 1.85 µM $FeCl_3$ and without antibiotics, pBpa, and propionate). Equal
volumes of each sample were transferred into single wells of each of two 24-well plates (NEST
Cell Culture Plate, cat: 702001). One plate was irradiated with 365-nm light for 30 min at room
temperature and the second plate was incubated in the dark simultaneously for 30 min at room
temperature. Samples were precipitated with 10% trichloroacetic acid (TCA) as described

previously and processed for immunoblot analysis with ExbD-specific polyclonal antibodies

and TonB-specific monoclonal antibodies (14, 71). Equal loads were confirmed by Coomassie

staining of the PVDF membrane used for the immunoblot.

Initial rates of ^{55}Fe-ferrichrome and ^{55}Fe-enterochelin transport. Cultures were grown as

described in media and growth conditions. Strains harboring plasmids with and without amber

substitutions in *exbD* were grown as described in the *in vivo* photo- cross-linking section. ^{55}Fe-

ferrichrome assay and ^{55}Fe-ferrichrome assay were performed in triplicate as previously

described (40). To exclude competition from native enterochelin, ^{55}Fe-enterochelin uptake was

assessed in *aroB-* strains that were subcultured with additional phenylalanine (40 µg/ml) and

tyrosine (40 µg/ml) to compensate for their inability to synthesize aromatic amino acids. To

determine steady state levels of the different ExbD variants, samples were precipitated with

10% TCA at the start of the assay and equivalent numbers of cells processed for immunoblot

analysis on 13% SDS-polyacrylamide gels. ExbD proteins were detected by immunoblotting

with ExbD-specific polyclonal antibodies (14, 71).

Ethidium bromide accumulation. Sub-cultures were grown as described above. At A_{550} of

0.5, 2 mL of cells were transferred to 4.5 mL disposable cuvettes (VWR Cat# 30620-276)

Ethidium bromide (6.25 µM) was added to the cells and fluorescence was detected using a LS-

55 fluorescence spectrophotometer. The excitation wavelength was 520 nm and the emission

wavelength was 590 nm. The width of the excitation and emission slits were 15 nm.

Fluorescence emission data was collected every second for 3 minutes using the Time Drive

application from the FLWINLAB software. For CCCP controls, either 15 µM or 60 µM CCCP

was added two minutes prior to ethidium bromide. The ethidium bromide accumulation rate for

each sample was determined by linear regression and recorded as Fluorescence Units per

second (FU/sec).

Proteinase K Accessibility. TonB sensitivity to proteinase K was performed as previously

described (8, 16, 35) Cultures were grown as described in media and growth conditions.

Spheroplasts were made as previously described (8). Samples were treated with carbonyl

cyanide *m*-chlorophenylhydrazone (60 μM), CCCP, where indicated 5 min prior to proteinase K

treatment (25 μg/mL). Samples were collected at 2 min, 5 min, 10 min, and 15 min following

proteinase K treatment with the addition of phenylmethylsulfonyl fluoride (1.0 mM) to inactivate

proteinase K at the specified times. Samples were precipitated with 10% TCA final

concentration and suspended in Laemmli sample buffer (70). Samples were loaded with equal

volumes on 13% SDS-polyacrylamide gels. Were detected via immunoblotting with ExbD-

specific polyclonal antibodies and TonB-specific monoclonal antibodies (14, 71).

Multiple Sequence Alignment. A sequence logo of the ExbD disordered region was

generated from a multiple sequence alignment by using HHblits in the Gremelin.bakerlab

program (73). Sequences aligned were all within an expect score of e^{-10} of *Escherichia coli K-*

12 strain (NP_417478.1). Positions were removed with >75% sequences alignment gaps. The

sequence for *Escherichia coli* ExbD (AAA69172.1), TolR (PSM33938.1), and MotB

(ACB03086.1) were aligned using clustal multiple sequence alignment generated by MUSCLE

(3.8).

Colicin Sensitivity Assay. The colicin sensitivity assay was performed as previously

described (40). Strains were grown to mid-exponential phase with expressing ExbD in T-broth.

The expression was confirmed by precipitation of cultures with trichloroacetic acid (TCA) and

were detected by immunoblot comparison to W3110. Cells were plated in T-top agar on T-

plates containing equal propionate concentrations as was in the sub-culture with 100 μg/ml

ampicillin. Five-fold dilutions of Colicins B, Ia, or M (Colicin B enters via FepA; Colicin Ia enters

via CirA; Colicin M enters FhuA**)** were spotted on the bacterial lawns and were incubated

791 overnight at 37°C. Scoring was reported as the highest dilution that produced clearing. The

792 assay was done in triplicate.

794 **ACKNOWLEDGEMENTS:**

795 Support from NIAID for grant AI11322 to K.P. is gratefully acknowledged.

REFERENCES:

1. Schauer K, Rodionov DA, Reuse H de. 2008. New substrates for TonB-dependent transport: do we only see the 'tip of the iceberg'? Trends in Biochemical Sciences 33:330–338.

2. Sheldon JR, Laakso HA, Heinrichs DE. 2016. Iron Acquisition Strategies of Bacterial Pathogens. Microbiology Spectrum 4.

3. Miethke M, Marahiel MA. 2007. Siderophore-Based Iron Acquisition and Pathogen Control. Microbiol Mol Biol Rev 71:413–451.

4. Noinaj N, Guillier M, Barnard TJ, Buchanan SK. 2010. TonB-dependent transporters: regulation, structure, and function. Annu Rev Microbiol 64:43–60.

5. Hickman SJ, Cooper REM, Bellucci L, Paci E, Brockwell DJ. 2017. Gating of TonB-dependent transporters by substrate-specific forced remodelling. Nature Communications 8:14804.

6. Gresock MG, Postle K. 2017. Going Outside the TonB Box: Identification of Novel FepA-TonB Interactions In Vivo. J Bacteriol 199.

7. Ferguson AD, Amezcua CA, Halabi NM, Chelliah Y, Rosen MK, Ranganathan R, Deisenhofer J. 2007. Signal transduction pathway of TonB-dependent transporters. PNAS 104:513–518.

8. Larsen RA, Thomas MG, Postle K. 1999. Protonmotive force, ExbB and ligand-bound FepA drive conformational changes in TonB. Molecular Microbiology 31:1809–1824.

9.   Moeck GS, Coulton JW, Postle K. 1997. Cell envelope signaling in Escherichia coli.

Ligand binding to the ferrichrome-iron receptor fhua promotes interaction with the energy-

transducing protein TonB. J Biol Chem 272:28391–28397.

10. Cornelissen CN, Anderson JE, Sparling PF. 1997. Energy-dependent changes in the

gonococcal transferrin receptor. Mol Microbiol 26:25–35.

11. Khursigara CM, Crescenzo GD, Pawelek PD, Coulton JW. 2004. Enhanced Binding of

TonB to a Ligand-loaded Outer Membrane Receptor ROLE OF THE OLIGOMERIC

STATE OF TonB IN FORMATION OF A FUNCTIONAL FhuA·TonB COMPLEX. J Biol

Chem 279:7405–7412.

12. Bradbeer C. 1993. The proton motive force drives the outer membrane transport of

cobalamin in Escherichia coli. J Bacteriol 175:3146–3150.

13. Krewulak KD, Vogel HJ. 2011. TonB or not TonB: is that the question? Biochemistry &

Cell Biology 89:87–97.

14. Higgs PI, Larsen RA, Postle K. 2002. Quantification of known components of the

Escherichia coli TonB energy transduction system: TonB, ExbB, ExbD and FepA. Mol

Microbiol 44:271–281.

15. Kampfenkel K, Braun V. 1993. Topology of the ExbB protein in the cytoplasmic

membrane of Escherichia coli. J Biol Chem 268:6050–6057.

16. Baker KR, Postle K. 2013. Mutations in Escherichia coli ExbB Transmembrane Domains

Identify Scaffolding and Signal Transduction Functions and Exclude Participation in a

Proton Pathway. J Bacteriol 195:2898–2911.

17.   Roof SK, Allard JD, Bertrand KP, Postle K. 1991. Analysis of Escherichia coli TonB

membrane topology by use of PhoA fusions. Journal of Bacteriology 173:5554–5557.

18.   Kampfenkel K, Braun V. 1992. Membrane topology of the Escherichia coli ExbD protein.

Journal of Bacteriology 174:5485–5487.

19.   Bernadac A, Gavioli M, Lazzaroni J-C, Raina S, Lloubès R. 1998. Escherichia coli tol-pal

Mutants Form Outer Membrane Vesicles. J Bacteriol 180:4872–4878.

20.   Masilamani R, Cian MB, Dalebroux ZD. 2018. Salmonella Tol-Pal Reduces Outer

Membrane Glycerophospholipid Levels for Envelope Homeostasis and Survival during

Bacteremia. Infect Immun 86.

21.   Gerding MA, Ogata Y, Pecora ND, Niki H, de Boer PAJ. 2007. The trans-envelope Tol-Pal

complex is part of the cell division machinery and required for proper outer-membrane

invagination during cell constriction in E. coli. Mol Microbiol 63:1008–1025.

22.   Cascales E, Lloubès R, Sturgis JN. 2001. The TolQ-TolR proteins energize TolA and

share homologies with the flagellar motor proteins MotA-MotB. Mol Microbiol 42:795–807.

23.   Brinkman KK, Larsen RA. 2008. Interactions of the Energy Transducer TonB with

Noncognate Energy-Harvesting Complexes. J Bacteriol 190:421–427.

24.   Kadner RJ, Heller KJ. 1995. Mutual inhibition of cobalamin and siderophore uptake

systems suggests their competition for TonB function. J Bacteriol 177:4829–4835.

25.   Kaserer WA, Jiang X, Xiao Q, Scott DC, Bauler M, Copeland D, Newton SMC, Klebba PE.

2008. Insight from TonB hybrid proteins into the mechanism of iron transport through the

outer membrane. J Bacteriol 190:4001–4016.

26.   Gresock MG, Kastead KA, Postle K. 2015. From Homodimer to Heterodimer and Back:

Elucidating the TonB Energy Transduction Cycle. Journal of Bacteriology 197:3433–3445.

27.   Chang C, Mooser A, Plückthun A, Wlodawer A. 2001. Crystal structure of the dimeric C-

terminal domain of TonB reveals a novel fold. J Biol Chem 276:27535–27540.

28.   Pawelek PD, Croteau N, Ng-Thow-Hing C, Khursigara CM, Moiseeva N, Allaire M,

Coulton JW. 2006. Structure of TonB in Complex with FhuA, E. coli Outer Membrane

Receptor. Science 312:1399–1402.

29.   Ghosh J, Postle K. 2004. Evidence for dynamic clustering of carboxy-terminal aromatic

amino acids in TonB-dependent energy transduction. Mol Microbiol 51:203–213.

30.   Jana B, Manning M, Postle K. 2011. Mutations in the ExbB cytoplasmic carboxy terminus

prevent energy-dependent interaction between the TonB and ExbD periplasmic domains.

J Bacteriol 193:5649–5657.

31.   Ollis AA, Manning M, Held KG, Postle K. 2009. Cytoplasmic Membrane Protonmotive

Force Energizes Periplasmic Interactions between ExbD and TonB. Mol Microbiol

73:466–481.

32.   Skare JT, Postle K. 1991. Evidence for a TonB-dependent energy transduction complex in

Escherichia coli. Mol Microbiol 5:2883–2890.

33.   Fischer E, Günter K, Braun V. 1989. Involvement of ExbB and TonB in transport across

the outer membrane of Escherichia coli: phenotypic complementation of exb mutants by

overexpressed tonB and physical stabilization of TonB by ExbB. J Bacteriol 171:5127–

5134.

34. Ollis AA, Postle K. 2012. ExbD mutants define initial stages in TonB energization. J Mol

Biol 415:237–247.

35. Ollis AA, Kumar A, Postle K. 2012. The ExbD Periplasmic Domain Contains Distinct

Functional Regions for Two Stages in TonB Energization. Journal of Bacteriology

194:3069–3077.

36. Swayne C, Postle K. 2011. Taking the Escherichia coli TonB Transmembrane Domain

"Offline"? Nonprotonatable Asn Substitutes Fully for TonB His20. Journal of Bacteriology

193:3693–3701.

37. Garcia-Herrero A, Peacock RS, Howard SP, Vogel HJ. 2007. The solution structure of the

periplasmic domain of the TonB system ExbD protein reveals an unexpected structural

homology with siderophore-binding proteins. Mol Microbiol 66:872–889.

38. Ollis AA, Postle K. 2011. The Same Periplasmic ExbD Residues Mediate In Vivo

Interactions between ExbD Homodimers and ExbD-TonB Heterodimers ▽. J Bacteriol

193:6852–6863.

39. Ollis AA, Postle K. 2012. Identification of Functionally Important TonB-ExbD Periplasmic

Domain Interactions In Vivo. Journal of Bacteriology 194:3078–3087.

40. Postle K. 2007. [12] - TonB System, In Vivo Assays and Characterization, p. 245–269. *In*

Simon, MI, Crane, BR, Crane, A (eds.), Methods in Enzymology. Academic Press.

41. Takatsuka Y, Nikaido H. 2009. Covalently Linked Trimer of the AcrB Multidrug Efflux

Pump Provides Support for the Functional Rotating Mechanism. J Bacteriol 191:1729–

1737.

42. Bulathsinghala CM, Jana B, Baker KR, Postle K. 2013. ExbB Cytoplasmic Loop Deletions
Cause Immediate, Proton Motive Force-Independent Growth Arrest. J Bacteriol
195:4580–4591.

43. Braun V, Herrmann C. 1993. Evolutionary relationship of uptake systems for biopolymers
in Escherichia coli: cross-complementation between the TonB-ExbB-ExbD and the TolA-
TolQ-TolR proteins. Mol Microbiol 8:261–268.

44. Ryu Y, Schultz PG. 2006. Efficient incorporation of unnatural amino acids into proteins in
Escherichia coli. Nat Methods 3:263–265.

45. Braun V, Gaisser S, Herrmann C, Kampfenkel K, Killmann H, Traub I. 1996. Energy-
coupled transport across the outer membrane of Escherichia coli: ExbB binds ExbD and
TonB in vitro, and leucine 132 in the periplasmic region and aspartate 25 in the
transmembrane region are important for ExbD activity. Journal of Bacteriology 178:2836–
2845.

46. Dormán G, Prestwich GD. 1994. Benzophenone photophores in biochemistry.
Biochemistry 33:5661–5673.

47. Metz B, Kersten GFA, Hoogerhout P, Brugghe HF, Timmermans HAM, Jong A de,
Meiring H, Hove J ten, Hennink WE, Crommelin DJA, Jiskoot W. 2004. Identification of
Formaldehyde-induced Modifications in Proteins REACTIONS WITH MODEL PEPTIDES.
J Biol Chem 279:6235–6243.

48. Sverzhinsky A, Chung JW, Deme JC, Fabre L, Levey KT, Plesa M, Carter DM,
Lypaczewski P, Coulton JW. 2015. Membrane Protein Complex ExbB4-ExbD1-TonB1
from Escherichia coli Demonstrates Conformational Plasticity. J Bacteriol 197:1873–1885.

49. Brinkman KK, Larsen RA. 2008. Interactions of the Energy Transducer TonB with

Noncognate Energy-Harvesting Complexes. Journal of Bacteriology 190:421–427.

50. Braun V. 1989. The structurally related exbB and tolQ genes are interchangeable in

conferring tonB-dependent colicin, bacteriophage, and albomycin sensitivity. Journal of

Bacteriology 171:6387–6390.

51. Larsen RA, Chen GJ, Postle K. 2003. Performance of Standard Phenotypic Assays for

TonB Activity, as Evaluated by Varying the Level of Functional, Wild-Type TonB. Journal

of Bacteriology 185:4699–4706.

52. Cascales E, Buchanan SK, Duché D, Kleanthous C, Lloubès R, Postle K, Riley M, Slatin

S, Cavard D. 2007. Colicin biology. Microbiol Mol Biol Rev 71:158–229.

53. van der Lee R, Buljan M, Lang B, Weatheritt RJ, Daughdrill GW, Dunker AK, Fuxreiter M,

Gough J, Gsponer J, Jones DT, Kim PM, Kriwacki RW, Oldfield CJ, Pappu RV, Tompa P,

Uversky VN, Wright PE, Babu MM. 2014. Classification of Intrinsically Disordered Regions

and Proteins. Chem Rev 114:6589–6631.

54. Jiang J, Zhang X, Chen Y, Wu Y, Zhou ZH, Chang Z, Sui S-F. 2008. Activation of DegP

chaperone-protease via formation of large cage-like oligomers upon binding to substrate

proteins. PNAS 105:11939–11944.

55. Krojer T, Sawa J, Schäfer E, Saibil HR, Ehrmann M, Clausen T. 2008. Structural basis for

the regulated protease and chaperone function of DegP. Nature 453:885–890.

56. Larsen RA, Deckert GE, Kastead KA, Devanathan S, Keller KL, Postle K. 2007. His20

Provides the Sole Functionally Significant Side Chain in the Essential TonB

Transmembrane Domain. J Bacteriol 189:2825–2833.

57. Vakharia-Rao H, Kastead KA, Savenkova MI, Bulathsinghala CM, Postle K. 2007.

Deletion and Substitution Analysis of the Escherichia coli TonB Q160 Region. Journal of

Bacteriology 189:4662–4670.

58. Sverzhinsky A, Fabre L, Cottreau AL, Biot-Pelletier DMP, Khalil S, Bostina M, Rouiller I,

Coulton JW. 2014. Coordinated Rearrangements between Cytoplasmic and Periplasmic

Domains of the Membrane Protein Complex ExbB-ExbD of Escherichia coli. Structure

22:791–797.

59. Lazzaroni JC, Vianney A, Popot JL, Bénédetti H, Samatey F, Lazdunski C, Portalier R,

Géli V. 1995. Transmembrane α-Helix Interactions are Required for the Functional

Assembly of theEscherichia coliTol Complex. Journal of Molecular Biology 246:1–7.

60. Gresock MG, Savenkova MI, Larsen RA, Ollis AA, Postle K. 2011. Death of the TonB

Shuttle Hypothesis. Front Microbiol 2.

61. Celia H, Noinaj N, Zakharov SD, Bordignon E, Botos I, Santamaria M, Barnard TJ,

Cramer WA, Lloubes R, Buchanan SK. 2016. Structural insight into the role of the Ton

complex in energy transduction. Nature 538:60–65.

62. Maki-Yonekura S, Matsuoka R, Yamashita Y, Shimizu H, Tanaka M, Iwabuki F, Yonekura

K. 2018. Hexameric and pentameric complexes of the ExbBD energizer in the Ton

system. eLife 7:e35419.

63. Pramanik A, Hauf W, Hoffmann J, Cernescu M, Brutschy B, Braun V. 2011. Oligomeric

structure of ExbB and ExbB-ExbD isolated from Escherichia coli as revealed by LILBID

mass spectrometry. Biochemistry 50:8950–8956.

64. Celia H, Botos I, Ni X, Fox T, De Val N, Lloubes R, Jiang J, Buchanan SK. 2019. Cryo-EM

structure of the bacterial Ton motor subcomplex ExbB–ExbD provides information on

structure and stoichiometry. Commun Biol 2:358.

65. Zhang XY-Z, Goemaere EL, Thomé R, Gavioli M, Cascales E, Lloubès R. 2009. Mapping

the Interactions between *Escherichia coli* Tol Subunits: ROTATION OF THE TolR

TRANSMEMBRANE HELIX. J Biol Chem 284:4275–4282.

66. Braun TF, Blair DF. 2001. Targeted disulfide cross-linking of the MotB protein of

Escherichia coli: evidence for two H(+) channels in the stator Complex. Biochemistry

40:13051–13059.

67. Wojdyla JA, Cutts E, Kaminska R, Papadakos G, Hopper JTS, Stansfeld PJ, Staunton D,

Robinson CV, Kleanthous C. 2015. Structure and Function of the Escherichia coli Tol-Pal

Stator Protein TolR. J Biol Chem 290:26675–26687.

68. Eick-Helmerich K, Braun V. 1989. Import of biopolymers into Escherichia coli: nucleotide

sequences of the exbB and exbD genes are homologous to those of the tolQ and tolR

genes, respectively. J Bacteriol 171:5117–5126.

69. Higgs PI, Myers PS, Postle K. 1998. Interactions in the TonB-Dependent Energy

Transduction Complex: ExbB and ExbD Form Homomultimers. J Bacteriol 180:6031–

6038.

70. Laemmli UK. 1970. Cleavage of Structural Proteins during the Assembly of the Head of

Bacteriophage T4. Nature 227:680–685.

71. Larsen RA, Myers PS, Skare JT, Seachord CL, Darveau RP, Postle K. 1996. Identification

of TonB homologs in the family Enterobacteriaceae and evidence for conservation of

TonB-dependent energy transduction complexes. J Bacteriol 178:1363–1373.

72. Young TS, Ahmad I, Yin JA, Schultz PG. 2010. An Enhanced System for Unnatural Amino

Acid Mutagenesis in E. coli. Journal of Molecular Biology 395:361–374.

73. Ovchinnikov S, Kamisetty H, Baker D. 2014. Robust and accurate prediction of residue–

residue interactions across protein interfaces using evolutionary information. eLife 3.

74.   Effect of loop deletions on the binding and transport of ferric enterobactin by FepA -

Newton - 1999 - Molecular Microbiology - Wiley Online Library.

75. Hill CW, Harnish BW. 1981. Inversions between ribosomal RNA genes of Escherichia coli.

Proc Natl Acad Sci U S A 78:7069–7072.

1000

1001

1002

1003

1004

1005
1006

FIGURES:

A.   **B.**

Figure 1: (A) TonB-dependent energy transduction. The iron-bearing siderophore enterochelin

(triangles) binds to the TBDT FepA with sub-nanomolar affinity (74). This high affinity

necessitates a mechanism to release the siderophore into the periplasmic space. In the

presence of cytoplasmic membrane proteins TonB, ExbB, ExbD, and the cytoplasmic

membrane proton motive force (PMF), the Fe-enterochelin is actively transported across the

outer membrane through FepA. Numbers indicate the ratio of the proteins in *E. coli* under iron-

limiting conditions (14). Because TonB is limiting, there must be a cycle during which TonB

binds and releases TBDTs. The Stages in the cycle have been studied, resulting in a model

where ExbB acts as a scaffold to manage the essential interaction between TonB and ExbD

that subsequently positions TonB for correct interaction with FepA. The interaction between

TonB and ExbD C-termini is dependent on PMF. The red box around ExbD identifies the ExbD

disordered region which is the topic of this study. Subsequent transport of enterochelin across

the cytoplasmic membrane does not require the TonB system. Adapted from Gresock et al.

2015, (26). (B) The NMR solution structure of ExbD periplasmic domain residues 44-141

[NCBI PDB ID: 2PFU, (37)]. The alpha helices and beta strands are colored in blue and green,

respectively. The ExbD disordered region is boxed in red. The first residue following the ExbD

TMD, D44, and last residue, P63 of the ExbD disordered region are labeled with the side chain

displayed. This image was generated using Swiss-PdbViewer [https://spdbv.vital-it.ch/(37)].

1044

Figure 2: Block alanine substitutions of ExbD residues 44-48 and 49-53 prevent TonB

system activity. (A) Initial rate of ^{55}Fe-enterochelin transport and corresponding steady-state

expression levels of the ExbD proteins. Strains were grown to mid-exponential phase and

assayed as described in Methods and Materials. The percentages of activity were calculated

relative to the initial rate of plasmid-encoded ExbD, pExbD (pKP999), expressed at

chromosomal levels in KP1526 (Δ*exbD*, Δ*tolQR, aroB*-). Assays were performed in triplicate

and carried out at least twice for each strain. The slopes were determined by a linear

regression fit of scintillation counts per minute. The error bars indicate mean ± SEM for a

representative triplicate set (above). The steady-state ExbD monomer levels at the time of

assay were determined by immunoblot analysis. To avoid competition from the only

synthesized siderophore, enterochelin, in *Escherichia coli K12*, *aroB*- strains were used.

cExbD designates chromosomally expressed ExbD from the wild-type strain, KP1270 (W3110,

aroB-). ExbD alanine substitution mutants are expressed from plasmids in KP1526 (Δ*exbD*,

Δ*tolQR, aroB*-). (B) Initial rate of [55]Fe-ferrichrome transport and corresponding steady-state

expression levels of ExbD proteins. The assay was performed and analyzed identically as in

(A) with using the siderophore, ferrichrome, which is transported through the TonB-dependent

transporter, FhuA. The percentages of activity were calculated relative to the initial rate of

plasmid-encoded ExbD, pExbD (pKP999) expressed at chromosomal levels in RA1045

(Δ*exbD*, Δ*tolQR*). cExbD designates chromosomally expressed ExbD from the wild-type strain,

W3110. All ExbD alanine substitution mutants are expressed from plasmids in RA1045

(Δ*exbD*, Δ*tolQR*). The plasmid identities and concentrations of propionate for chromosomal

expression are listed in Table S1.

Figure 3: Identification of ExbD-TonB PMF-dependent and PMF-independent complexes using *in vivo* photo-cross-linking. Plasmid-encoded ExbD (pBpa) substitutions were co-expressed with pEVOL in strain RA1045 (Δ*exbD*, Δ*tolQR*). At mid-exponential phase, strains were subjected to UV irradiation for photo-cross-linking as described in Materials and Methods.

Equivalent numbers of cells were precipitated with TCA and the ExbD-TonB complexes were

visualized on immunoblots of 13% SDS-polyacrylamide gels with anti-TonB monoclonal

antibodies. cExbD designates chromosomally expressed ExbD from the wild-type strain,

W3110. The parent plasmid strain pExbD (pKP999) and all ExbD (pBpa) substitutions were

expressed in RA1045 (Δ*exbD*, Δ*tolQR*), except for ExbD (S54pBpa) ΔtonB, which was

expressed in KP1509 (Δ*exbD*, Δ*tolQR, ΔtonB*). D25N is a mutation in the ExbD TMD. The

protonophore CCCP was added prior to UV exposure. His6X refers to ExbD (S54pBpa) with a

C-terminal His6X tag. On the left, positions of a PMF-dependent ExbD-TonB complex [ExbD-

TonB (PMF], a PMF-independent complex [ExbD-TonB (I)], monomeric TonB, and a TonB

degradation product are shown. Mass markers are indicated on the right. Steady state ExbD

and TonB monomer levels from the same experiment were determined by immunoblot analysis

of control samples that were not exposed to UV light. The plasmid identities and

concentrations of propionate for chromosomal expression are listed in Table S1.

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098　**A.**

1099

1100　**B.**

1101

1102

1103　**Figure 4: Virtually all the ExbD (pBpa) substitutions in the ExbD disordered region**

1104　**capture PMF-dependent or PMF-independent TonB complexes.** (A) Initial rate of [55]Fe-

1105　ferrichrome transport for ExbD (pBpa) substitutions at residues 44-54 (left panel) and residues

1106　54-63 (right panel) with the corresponding steady-state expression levels of ExbD proteins at

the time of assay above each panel. ExbD pBpa substitutions were plasmid-encoded and co-expressed with plasmid pEVOL in RA1045 ($\Delta exbD$, $\Delta tolQR$). Strains were grown to mid-exponential phase and assayed as described in Methods and Materials. The percentages of activity were calculated relative to the initial rate for [55]Fe-ferrichrome transport of plasmid-encoded ExbD, pExbD (pKP999). Assays were performed in triplicate and carried out at least twice for each strain. The slopes were determined by a linear regression fit of scintillation counts per minute. The error bars indicate mean ± SEM for a representative triplicate set. cExbD designates chromosomally expressed ExbD from the wild-type strain, W3110. The plasmid identities and concentrations of propionate for chromosomal expression are listed in Table S1. (B) *In vivo* photo-cross-linking of pBpa substitutions for ExbD residues 44-54 (left panel) and residues 54-63 (right panel) on an anti-TonB immunoblot. Note that ExbD S54)pBpa is included on both immunoblots in lanes 14 and 18 for comparison of levels of expression. All samples (identical as in A) were grown to mid-exponential phase, at which point photo-cross-linking was performed as described in Materials and Methods. Equivalent numbers of cells were precipitated with TCA and the ExbD-TonB complexes were visualized on immunoblots of 13% SDS-polyacrylamide gels with anti-TonB monoclonal antibodies. On the left and right, positions of a PMF-dependent ExbD-TonB complex [ExbD-TonB (PMF)], a PMF-independent complex [ExbD-TonB (I)], monomeric TonB, and a TonB degradation product are shown. Mass markers are indicated in the center. The steady-state ExbD and TonB monomer levels from the same experiment were determined by immunoblot analysis of control samples that were not exposed to UV light.

1132

1133

Figure 5: Two different ExbD-ExbB complexes and an ExbD homodimer are identified by

***in vivo* photo-cross-linking**. (A) *In vivo* photo-cross-linking of a representative ExbD photo-

cross-linkable mutant, ExbD (D44pBpa), in the presence of ExbB with (+) and without (-) a C-

terminal His6X-tag. At mid-exponential phase, strains were subjected to UV irradiation for

photo-cross-linking as described in Materials and Methods. Equivalent numbers of cells were

precipitated with TCA and the ExbD complexes were visualized on immunoblots of 13% SDS-

polyacrylamide gels with anti-ExbD polyclonal antibodies. cExbD designates chromosomally

expressed ExbB and ExbD from the wild-type strain, W3110. Plasmid pExbBD (pKP1657) and

its corresponding ExbD (D44pBpa) variants were expressed with pEVOL in RA1017 (W3110,

ΔexbBD::kan, ΔtolQRA). ExbD is overexpressed from plasmid pExbBD and its corresponding

ExbD (D44pBpa) variant in the absence of inducer relative to chromosomally ExbD

expression. Positions of ExbD-ExbB (upper)—blue asterisk on immunoblot, ExbD-ExbB

(lower)—yellow asterisk on immunoblot, likely ExbD-ExbD homodimers–green pound symbol

on immunoblot, and ExbD monomer are indicated on the left. pExbBD expressed with pEVOL

(lane 3) forms two unidentified complexes that are non-ExbD residue-specific (unlabeled).

Mass markers are indicated on the right. Steady-state ExbD monomer levels from the same

experiment were determined by immunoblot analysis of control samples that were not exposed

to UV light. The plasmid identities and concentrations of propionate for chromosomal

expression are listed in Table S1. (B) *In vivo* photo-cross-linking of ExbD (S54pBpa), with (+)

and without (-) a C-terminal His6X-tag using the same procedure as in (A). Plasmid pExbD

(pKP999) and a corresponding variant, ExbD (S54pBpa) were expressed with pEVOL in

RA1045 (Δ*exbD*, Δ*tolQR*). On the left, positions of a PMF-dependent ExbD-TonB complex

[ExbD-TonB (PMF], ExbD-ExbB, an ExbD homodimer, and monomeric ExbD are shown. Mass

markers are indicated on the right. Steady-state ExbD monomer levels from the same

experiment were determined by immunoblot analysis of control samples that were not exposed

to UV light. The plasmid identities and concentrations of propionate for chromosomal

expression are listed in Table S1.

1161

1162

1163

1164

Figure 6: The majority of pBpa substitutions in the ExbD disordered region capture

contacts with ExbB and ExbD. *In vivo* photo-cross-linking of ExbD pBpa substitutions in

residues 44-54 (left panel) and 54-63 (right panel) from Fig. 4 probed with anti-ExbD

antibodies. Note that ExbD (S54pBpa) is included on both immunoblots in lanes 14 and 18 for

comparison of levels of expression. Strains were grown to mid-exponential phase, at which

point photo-cross-linking was performed as described in Materials and Methods. Equivalent

numbers of cells were precipitated with TCA and the ExbD-TonB complexes were visualized

on immunoblots of 13% SDS-polyacrylamide gels with anti-ExbD polyclonal antibodies. cExbD

designates chromosomally expressed ExbD from the wild-type strain, W3110. Plasmid pExbD

(pKP999) and its corresponding ExbD pBpa substitutions were expressed in RA1045 (Δ*exbD*,

Δ*tolQR*) with plasmid pEVOL. Positions of the PMF-dependent ExbD-TonB complex identified

in Fig. 3, ExbD-ExbB (upper) and ExbD-ExbB (lower) from Figure 5A, the ExbD-ExbD from

Figure 5B and ExbD monomer are indicated on the right and left. The ExbD-TonB (PMF) is

differentiated from unknown complexes by a red asterisk (*). The ExbD-TonB PMF-

1181   independent complex is distinguished from ExbB-ExbD (upper) by a yellow pound symbol (#).

1182   Mass standards are in the center. The steady-state ExbD levels from the same experiment

1183   were determined by immunoblot analysis of control samples that were not exposed to UV light

1184   (bottom). The plasmid identities and concentrations of propionate for chromosomal expression

1185   are listed in Table S1.

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200 **A.**

1201   **Residues**   weblogo.berkeley.edu

1202

1203 **B.**

1204

```
1205  CLUSTAL multiple sequence alignment by MUSCLE (3.8)
1206
1207  ExbD (38)  ---APLATVD VKVNLP ASTSTPQPRPE-----------KPVYLSVKA--------(60)
1208  TolR (36)  ---APIITQS VEVDLP DATESQAVSSNDN---------PPVIVEVSG--------(60)
1209  MotB (61)  YFRTPLATAV TGGDRI SNSESPIPGGGDDYTQSQGEVNKQPNIEELKKRMEQSRLR (116)
1210    :*: * . : :.:   * .:.
1211
```

Figure 7: A conservation alignment of the ExbD disordered region identifies a ΨΧΨΧLP

motif. The psi symbol (Ψ) represents a residue with a large aliphatic side chain and X

represents any non-hydrophobic residue. (A) A WebLogo image of the ExbD disordered

region. The image was generated from an ExbD multiple sequence alignment by HHblits of

sequences within an E-value of e^{-10} of *Escherichia coli K-12* strain (NP_417478.1) using the

Gremelin.bakerlab program (gremlin.bakerlab.org/). Positions with >75% sequences alignment

gaps were removed. (B) A multiple sequence alignment was used to identify potential

ΨΧΨΧLP motifs in ExbD homologs TolR and MotB. A clustal multiple sequence alignment was

generated by MUSCLE (3.8) of *Escherichia coli* ExbD (AAA69172.1), TolR (PSM33938.1), and

MotB (ACB03086.1). A red box encompasses the *Escherichia coli* K12 ExbD ΨΧΨΧLP motif

containing conserved residues V45, V47, L49 and P50 and shows that these residues are

conserved in TolR but not MotB.

A.

B.

Mutant	Colicin B[+]	Colicin Ia[+]	Colicin M[+]
pExbD	8, 9, 8	8, 8, 8	8, 8, 8
ExbD (V45A, V47A)	T, T, T	T, T, T	T, T, T
ExbD (V45A, L49A)	T, T, T	T, T, T	T, T, T
ExbD (V45A, P50A)	T, T, T	0*, 0*, 0*	T, T, T
ExbD (V47A, L49A)	T, T, T	T, T, T	T, T, T
ExbD (V47A, P50A)	T, T, T	T, T, T	T, T, T
ExbD (L49A, P50A)	T, T, T	T, T, T	T, T, T

[+] T refers to tolerant to undiluted colicin lysate
* 0 refers to hazy clearing in with undiluted colicin

1225

Figure 8: The synergy of double-alanine substitutions within the ExbD ΨXΨXLP motif

establishes its essentiality for TonB-dependent energy transduction.

Initial rate of ^{55}Fe-ferrichrome transport for ExbD variants with the corresponding steady-state

expression levels of ExbD proteins at the time of assay above each panel. Strains were grown

to mid-exponential phase and assayed as described in Methods and Materials. The

percentages of activity were calculated relative to the initial rate for ^{55}Fe-ferrichrome transport

of plasmid-encoded ExbD, pExbD (pKP999). Assays were performed in triplicate and carried

out at least twice for each strain. The slopes were determined by a linear regression fit of

scintillation counts per minute. The error bars indicate mean ± SEM for a representative

triplicate set (above). Steady-state levels of ExbD-specific expression from the transport

experiments are indicated above the corresponding bars of the graph below them. cExbD

designates chromosomally expressed ExbD from the wild-type strain, W3110. The plasmid

identities and concentrations of propionate for chromosomal expression are listed in Table S1.

(B) Colicin sensitivity assay. Strains were grown to mid-exponential phase and assayed in

triplicate with Colicin B, Colicin Ia, and Colicin M as described in the Material and Methods.

pExbD (pKP999) and all ExbD alanine substitutions were expressed from plasmids in RA1045

(ΔexbD, ΔtolQR). The plasmid identities and concentrations of propionate for chromosomal

expression are listed in Table S1. Scoring was reported as the highest 5-fold colicin dilution

that produced clearing in three different trials. "T" refers to colicin tolerance (insensitivity) and

"0" refers to hazy clearing caused by the undiluted colicin lysate.

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

Figure 9: Mutations in the ExbD ΨXΨXLP motif leave TonB stalled at Stage II when PMF is present. Spheroplasts were generated from W3110 (cExbD), and RA1045 (Δ*exbD*, Δ*tolQR*), as well as RA1045 containing plasmids pKP999 (pExbD), pKP1194 [ExbD (D25N)], pKP2171 [ExbD (V45A, V47A)], pKP2201 [ExbD (V47A, L49A)], and pKP2202 [ExbD (L49A, P50A)] as described in the Materials and Methods. Each spheroplast was treated with proteinase K (PK) for 0 min, 2 min, 5 min, 10 min and 15 min in the presence of DMSO (-CCCP) or CCCP dissolved in DMSO (+CCCP). The samples were visualized on immunoblots of 13% SDS-polyacrylamide gels with anti-TonB monoclonal antibodies. Identities of the various strains assayed are indicated on the left. Positions of TonB monomers, the ~23 kDa

1267 PK-resistant TonB fragment indicative of Stage II of TonB energization and the ~28 kDa

1268 fragment with no known relevance to any Stage in the TonB energization cycle (34), are on the

1269 left. The appearance of the ~23 kDa TonB fragment upon lengthy exposure of ExbD (L49A,

1270 P50A) samples allowed confident identification of the 28 kDa TonB fragment generated in that

1271 strain. The concentrations of propionate for chromosomal expression are listed in Table S1.

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

A.

B.

1282

1283

Figure 10: The ExbD ΨXΨXLP motif controls ExbD interactions through other sites. *In*

vivo photo-cross-linking of ExbD (E14pBpa), ExbD (I24pBpa), ExbD (S54pBpa), ExbD

(S67pBpa), ExbD (K97pBpa), and ExbD (E136pBpa) with (+) and without (-) V45A,V47A

double alanine substitutions. Strains were grown to mid-exponential phase, at which point

photo-cross-linking was performed as described in Materials and Methods. Equivalent

numbers of cells were precipitated with TCA and the samples were visualized on immunoblots

of 13% SDS-polyacrylamide gels with either (A) anti-TonB monoclonal antibodies or (B) anti-

ExbD polyclonal antibodies. cExbD designates chromosomally expressed ExbD from the wild-

type strain, W3110. The parent plasmid pExbD (pKP999) and corresponding ExbD variants

were co-expressed with plasmid pEVOL in RA1045 (Δ*exbD, ΔtolQR*). (A) On the left, positions

of a PMF-dependent ExbD-TonB complex [ExbD-TonB (PMF)—red asterisk], a PMF-

independent complex [ExbD-TonB (I)], monomeric TonB, and a TonB degradation product are

shown (See Fig. 3). Mass markers are indicated on the right. Steady state ExbD and TonB

monomer levels from the same experiment were determined by immunoblot analysis of control

samples that were not exposed to UV light. (B) On the left, positions of ExbD-ExbB (upper)

and ExbD-ExbB (lower) (see Figure 5A), the ExbD-ExbD homodimer (see Figure 5B) and

ExbD monomer are shown. Unidentified complexes trapped by the ExbD pBpa variants

containing V45A,V47A are indicated on the immunoblot by a green (+) symbol. Mass markers

are indicated on the right. Steady-state ExbD levels from the same experiment were

determined by immunoblot analysis of control samples that were not exposed to UV light. The

plasmid identities and concentrations of propionate for chromosomal expression are listed in

Table S1.

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

	Stage I	Stage II	Stage III
Proteinase K- resistant TonB fragment	–	+	–
TonB-ExbD formaldehyde crosslink ★	–	–	+

1318

1319

Figure 11: Model: The PMF-dependent transition from Stage II to Stage III requires the

ExbD ΨXΨXLP motif. Initial Stages of TonB energization. The *in vivo* data suggest that the C-

terminus of TonB is an intrinsically disordered protein whose conformation is managed by

ExbD [(34), Fig. 1]. In Stage I, ExbD and TonB do not interact, although both are stabilized by

ExbB (not shown). TonB in untreated spheroplasts is degraded by exogenously added

proteinase K. In Stage II, the first ~155 residues of TonB are somehow protected from

proteinase K if the spheroplasts have been treated with the protonophore CCCP and if ExbD is

also present. The protonophore CCCP is required to prevent TonB transitioning to Stage III.

Although we hypothesized at that time that ExbD was protecting TonB, no Stage II complex

containing both proteins was detected.

In this study, we captured for the first time the ExbD-TonB PMF-independent

intermediate using *in vivo* photo-cross-linking as part of the normal energy transduction cycle,

without the need to add protonophore. Mutations in the ExbD ΨXΨXLP motif (gold box)

allowed capture of ExbD-TonB in Stage II, but prevented transition of TonB to Stage III. In

Stage III, ExbD and TonB can be formaldehyde cross-linked into a PMF-dependent complex

containing both proteins and in this work we also capture the Stage III, PMF-dependent form of

ExbD-TonB. This figure was adapted from Gresock et al, 2015 (26).

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

1350

1351

1352

SUPPLEMENTAL INFORMATION

Table S1, Kopp and Postle:

Strain	Genotype		Reference
W3110	F⁻ IN(*rrnD-rrnE*)*1*		(1)
KP1270	W3110, *aroB-*		(2)
RA1045	W3110, Δ*exbD*, Δ*tolQR*		(3)
KP1526	W3110, Δ*exbD*, Δ*tolQR, aroB-*		Present Study
RA1017	W3110, Δ*exbBD*::*kan*, Δ*tolQRA*		(4)
KP1509	W3110 Δ*exbD* Δ*tolQR*, Δ*tonB*::*kan*		(5)
Plasmid	**Genotype or Phenotype**	**ªPropionate Induction Concentration**	**References**
pEVOL-BpF	tRNA synthetase and tRNA for pBpa, camr , p15A	N/A	(6)
pKP999	*exbD* in pPro24	0.5 mM	(5)
pKP1657	*exbBD* in pPro24	0 mM	Present study
pKP1064	*ExbD (D25N)*	0 mM	(5)
pKP2073	ExbD (44-48ala)	0.25 mM	Present study
pKP2077	ExbD (49-53ala)	0.5 mM	Present study
pKP2153	ExbD (54-58ala)	0.75 mM	Present study
pKP2180	ExbD (59-63ala)	20 mM	Present study
pKP2140	ExbD (D44A)	0.25 mM	Present study
pKP2141	ExbD (V45A)	2.5 mM	Present study
pKP2142	ExbD (K46A)	0 mM	Present study
pKP2143	ExbD (V47A)	1 mM	Present study

pKP2144	ExbD (N48A)	0.25 mM	Present study
pKP2145	ExbD (L49A)	0.25 mM	Present study
pKP2154	ExbD (P50A)	0 mM	Present study
pKP2152	ExbD (S52A)	0.25 mM	Present study
pKP2146	ExbD (T53A)	0.25 mM	Present study
pKP2007	ExbD (D44Am)	0 mM	Present study
pKP2133	ExbD (V45Am)	0 mM	Present study
pKP2017	ExbD (K46Am)	0 mM	Present study
pKP2131	ExbD (V47Am)	0.5 mM	Present study
pKP2018	ExbD (N48Am)	2 mM	Present study
pKP2147	ExbD (L49Am)	0.5 mM	Present study
pKP1971	ExbD (P50Am)	0 mM	Present study
pKP2138	ExbD (A51Am)	45 mM	Present study
pKP1951	ExbD (S52Am)	1 mM	Present study
pKP2151	ExbD (T53Am)	1 mM	Present study
pKP1972	ExbD (S54Am)	1 mM	Present study
pKP2139	ExbD (T55Am)	8 mM	Present study
pKP1973	ExbD (P56Am)	5 mM	Present study
pKP2137	ExbD (Q57Am)	10 mM	Present study
pKP1974	ExbD (P58Am)	5 mM	Present study
pKP1898	ExbD (R59Am)	5 mM	Present study
pKP1975	ExbD (P60Am)	8 mM	Present study
pKP2027	ExbD (E61Am)	10 mM	Present study
pKP1949	ExbD (K62Am)	8 mM	Present study

pKP2136	ExbD (P63Am)	7 mM	Present study
pKP1926	ExbD (E14Am)	4 mM	Present study
pKP2064	ExbD (I24Am)	0 mM	Present study
pKP2167	ExbD (S67Am)	5 mM	Present study
pKP2227	ExbD (K97Am)	2 mM	Present study
pKP1907	ExbD (E136Am)	0 mM	Present study
pKP2171	ExbD (V45A, V47A)	1 mM	Present study
pKP2164	ExbD (V45A, L49A)	0.5 mM	Present study
pKP2182	ExbD (V45A, P50A)	0.25 mM	Present study
pKP2201	ExbD (V47A, L49A)	0.25 mM	Present study
pKP2190	ExbD (V47A, P50A)	0.25 mM	Present study
pKP2202	ExbD (L49A, P50A)	0.5 mM	Present study
pKP2213	ExbD (V45A,V47A, E14Am)	0 mM	Present study
pKP2214	ExbD (V45A,V47A, I24Am)	0 mM	Present study
pKP2161	ExbD (V45A,V47A, S54Am)	3 mM	Present study
pKP2215	ExbD (V45A,V47A, S67Am)	1.5 mM	Present study
pKP2228	ExbD (V45A,V47A, K97Am)	30 mM	Present study
pKP2220	ExbD (V45A,V47A, E136Am)	0 mM	Present study
pKP2063	ExbD (S54pBpa, D25N)	1 mM	Present study
pKP1932	ExbD (R59pBpa, D25N)	5 mM	Present study
pKP2259	ExbD (S54AM)-His(6x)	5mM	Present study
pKP2295	ExbB-ExbD (D44Am)	0 mM	Present study
pKP2272	ExbB-His(6x)-ExbD (D44Am)	0 mM	Present study

1355 [a] Propionate concentration required for plasmid induction in RA1045 background strain to
1356 achieve similar W3110 ExbD chromosomal protein levels.

SUPPLEMENTAL FIGURES Kopp and Postle:

1359

1360

1361

Figure S1: ExbD (44-48ala) and ExbD (49-53ala) fail to form a formaldehyde crosslinked ExbD-TonB PMF-dependent complex. *In vivo* formaldehyde-cross-linking of block alanine substitutions in the ExbD disordered region. Strains were grown to mid-exponential phase, at which point formaldehyde-cross-linking was performed as described in Materials and Methods. Equivalent numbers of cells were visualized on immunoblots of 13% SDS-polyacrylamide gels with (A) anti-ExbD polyclonal and (B) anti-TonB monoclonal antibodies. cExbD designates chromosomally expressed ExbD from the wild-type strain, W3110. The parent plasmid, pExbD (pKP999) and its corresponding ExbD block alanine substitutions were expressed in both RA1045 (ΔexbD, ΔtolQR) [+] and KP1509 (ΔexbD, ΔtolQR, ΔtonB) [-]. (A) Positions of the

previously characterized formaldehyde cross-linked PMF-dependent ExbD-TonB complex (red

asterisk on immunoblot), the ExbD-ExbB complex, the ExbD-ExbD homodimer and ExbD

monomer are shown on the left (5). Mass markers are shown on the right. (B) Positions of the

TonB-FepA complex, the TonB-ExbB complex, the PMF-dependent TonB-ExbD complex and

TonB monomer are shown. Mass markers are shown on the right. Below each immunoblot are

the levels of the corresponding monomer in each sample. The plasmid identities and

concentrations of propionate for chromosomal expression are listed in Table S1.

1378

1379

1380

1381

1382

1383

1384

1385

Figure S2: Failure to form the PMF-dependent ExbD-TonB formaldehyde crosslink is not

due to loss of PMF; a small, but significant difference is detected among the block ala

substitutions. The rate of ethidium bromide (EtBr) accumulation, an indirect assay for loss of

PMF, was measured for ExbD alanine block substitutions as described in the Materials and

Methods. Strains were grown to mid-exponential phase and assayed as described in Methods

and Materials. (A) The average EtBr accumulation rates (FU/sec) were calculated relative to

the rate of plasmid-encoded ExbD, pExbD (pKP999) expressed at chromosomal levels in

RA1045 (Δ*exbD*, Δ*tolQR*). The corresponding steady-state ExbD monomer levels determined

by immunoblot analysis from the experiment from one of the trials are shown above the bar

graph. cExbD designates chromosomally expressed ExbD from the wild-type strain, W3110.

For lanes 3 and 4, W3110 was treated with 60 µM CCCP or 15 µM (as labeled) was prior to

addition of EtBr as described in the Materials and Methods. ExbD alanine substitutions were

expressed from plasmids in RA1045 (Δ*exbD*, Δ*tolQR*). The plasmids and concentrations of

propionate for chromosomal expression are listed in Table S1. Numbers above the immunoblot

correspond to samples below them starting with (Δ*exbD*, Δ*tolQR*) on the left as #1.

1401

1402

Figure S3: ExbD residues V45, V47, L49 and P50 are important for TonB System activity.
Initial rate of [55]Fe-ferrichrome transport for ExbD single alanine substitutions in residues 44-53
and corresponding steady-state expression levels of the ExbD proteins. Strains were grown to
mid-exponential phase and assayed as described in Methods and Materials. The percentages
of activity were calculated relative to the initial rate of plasmid-encoded ExbD, pExbD
(pKP999), expressed at chromosomal levels in RA1045 ($\Delta exbD$, $\Delta tolQR$). Assays were

performed in triplicate and carried out at least twice for each strain. The slopes were

determined by a linear regression fit of scintillation counts per minute. The error bars indicate

mean ± SEM for a representative triplicate set. (above) Steady-state levels of ExbD-specific

expression from the transport experiments are indicated above the corresponding bars of the

graph below them. cExbD designates chromosomally expressed ExbD from the wild-type

strain, W3110. All ExbD alanine substitution mutants are expressed from plasmids in RA1045

(ΔexbD, ΔtolQR). The plasmid identities and concentrations of propionate for chromosomal

expression are listed in Table S1. The mean percent of the pExbD ^{55}Fe-ferrichrome transport

rates for ExbD (V45A), ExbD (V47A), ExbD (L49A), and ExbD (P50A) were 32%, 21%, 34%,

and 50% respectively. ExbD residue 51 is naturally an alanine in *Escherichia coli* W3110 and

is reflected in the pExbD initial rate.

1422

1423

1424

1425

1426

1427

1428

1429

1430

1431

1432

1433

1434

1435

1436

1437

1438

1439 **A.**   **B.**

E. coli TolR(36–142)

	Chain 1	Chain 2
1	Gln-41(N)	Thr-139(OG1)
2	Val-43(N)	Leu-137(O)
3	Val-45(N)	Val-135(O)
4	Asp-46(N)	Asp-46(O)
5	Leu-47(N)	Lys-133(O)
6	Tyr-117(N)	Lys-133(O)
7	Tyr-117(N)	Val-115(O)
8	Tyr-117(OH)	Ile-109(O)
9	Asp-118(N)	Lys-113(O)
10	His-128(NE2)	Leu-47(O)
11	Ser-134(OG)	Asp-46(OD1)
12	Val-135(N)	Glu-44(OE2)
13	Val-135(N)	Val-45(O)
14	Leu-137(N)	Val-43(O)
15	Met-138(N)	Tyr-117(OH)
16	Thr-139(N)	Gln-41(O)
17	Thr-139(OG1)	Gln-41(O)
18	Thr-139(OG1)	Gln-41(N)
19	Leu-137(O)	Val-43(N)
20	Val-135(O)	Val-45(N)
21	Asp-46(O)	Asp-46(N)
22	Lys-133(O)	Leu-47(N)
23	Val-115(O)	Tyr-117(N)
24	Lys-113(O)	Tyr-117(N)
25	Ile-109(O)	Tyr-117(OH)
26	Lys-113(O)	Asp-118(N)
27	Leu-47(O)	His-128(NE2)
28	Asp-46(OD1)	Ser-134(OG)
29	Val-45(O)	Val-135(N)
30	Glu-44(OE2)	Val-135(N)
31	Val-43(O)	Leu-137(N)
32	Tyr-117(OH)	Met-138(N)
33	Gln-41(O)	Thr-139(N)
34	Gln-41(O)	Thr-139(OG1)
35a	His-128(NE2)	Asp-49(OD1)
36a	Asp-49(OD1)	His-128(NE2)

β1

S-V-E-V-D-L-P

1440

1441

Figure S4: The hydrophobic residues in ΨXΨXLP motif are located at dimeric interface

of a TolR (36-142) dimeric crystal structure. (A) The table of the TolR residues at the

homodimeric interface. Chain 1 is one TolR and chain 2 is a second TolR. The red text

identifies the conserved hydrophobic residues in the ΨXΨXLP motif from chain 1 and the blue

text identifies the conserved hydrophobic residues in the ΨXΨXLP motif from chain 2. This

table was adapted from Wojdyla et. al, 2015 [(7), B, top image] The dimeric crystal structure of

TolR (36-142). The yellow rectangle identifies approximately where the motif lies from one

TolR of the homodimeric structure. This structure was adapted from Wojdyla et. al, 2015 [(7),

B, bottom image] The portion of one TolR where the motif residues lie. The conserved

hydrophobic valines are in a beta strand and the conserved leucine and proline lie in a turn immediately distal to that beta-strand.

Supplemental references:

1. Hill CW, Harnish BW. 1981. Inversions between ribosomal RNA genes of Escherichia coli. Proc Natl Acad Sci U S A 78:7069–7072.

2. Larsen RA, Thomas MG, Postle K. 1999. Protonmotive force, ExbB and ligand-bound FepA drive conformational changes in TonB. Molecular Microbiology 31:1809–1824.

3. Brinkman KK, Larsen RA. 2008. Interactions of the Energy Transducer TonB with Noncognate Energy-Harvesting Complexes. J Bacteriol 190:421–427.

4. Larsen RA, Deckert GE, Kastead KA, Devanathan S, Keller KL, Postle K. 2007. His20 Provides the Sole Functionally Significant Side Chain in the Essential TonB Transmembrane Domain. J Bacteriol 189:2825–2833.

5. Ollis AA, Manning M, Held KG, Postle K. 2009. Cytoplasmic Membrane Protonmotive Force Energizes Periplasmic Interactions between ExbD and TonB. Mol Microbiol 73:466–481.

6. Young TS, Ahmad I, Yin JA, Schultz PG. 2010. An Enhanced System for Unnatural Amino Acid Mutagenesis in E. coli. Journal of Molecular Biology 395:361–374.

7. Wojdyla JA, Cutts E, Kaminska R, Papadakos G, Hopper JTS, Stansfeld PJ, Staunton D, Robinson CV, Kleanthous C. 2015. Structure and Function of the Escherichia coli Tol-Pal Stator Protein TolR. J Biol Chem 290:26675–26687.