Positive and negative determinants of normal linear growth among pre-school children living in better-off households: an analysis of secondary data from sub-Saharan Africa

Dickson A. Amugsi[1*], Zacharie T. Dimbuene[2, 3], Elizabeth W. Kimani-Murage[1]

1. Maternal and Child Wellbeing Unit, Research Division, African Population and Health Research Center, Nairobi, Kenya

2. Department of Population Sciences and Development, University of Kinshasa, Democratic Republic of the Congo

3. Microdata Access Division, Statistics Canada, Ottawa, Canada

* Corresponding author

Email: damugsi2002@yahoo.com; damugsi@aphrc.org (DAA)

Abstract

This study examined the positive and negative determinants of normal linear growth among pre-

school children living in better-off households, using survey data from Ghana, Kenya, Nigeria,

Mozambique and Democratic Republic of Congo (DRC). The main outcome variable was child

height-for-age z-scores (HAZ), categorized into HAZ≥-2SD (normal growth/not stunted) and

HAZ<−2 (stunted). Logistic regression was utilized in the analysis. We estimated adjusted odd

ratios (aORs) of the determinants of normal growth. Higher maternal weight (measured by body

mass index) was associated with increased odds of normal growth in Mozambique, DRC, Kenya

and Nigeria. A unit increase in maternal years of education was associated with increased odds

in normal growth in DRC (aOR=1.06, 95% CI=1.03, 1.09), Ghana (aOR=1.08, 95% CI=1.04, 1.12),

Mozambique (aOR=1.08, 95% CI=1.05, 1.11) and Nigeria (aOR=1.07, 95% CI=1.06, 1.08). A year

increase in maternal age was associated positively with normal growth in all the five countries.

Breastfeeding was associated with increased odds of normal growth in Nigeria (aOR= 1.30, 95%

CI=1.16, 1.46) and Kenya (aOR=1.37, 95% CI=1.05, 1.79). Children of mothers who were working

had 25% (aOR=0.75, 95% CI=0.60, 0.93) reduced odds of normal growth in DRC. A unit change in

maternal parity was associated with 10% (aOR=0.90, 95% CI=0.84, 0.97), 23% (aOR=0.77, 95%

CI=0.63, 0.93), 25% (aOR=0.75, 95% CI=0.69, 0.82), 6% (aOR=0.94, 95% CI=0.89, 0.99) and 5%

(aOR=0.95, 95% CI=0.92, 0.99) reduced odds of normal growth in DRC, Ghana, Kenya,

Mozambique and Nigeria respectively. A child being a male was associated with 16% (aOR=0.82,

95% CI=0.68, 0.98), 40% (aOR=0.60, 95% CI=0.40, 0.89), 37% (aOR=0.63, 95% CI=0.51, 0.77) and

21% (aOR=0.79, 95% CI=0.71, 0.87) reduced odds of normal child growth in DRC, Ghana, Kenya

and Nigeria respectively. In conclusion, maternal education, weight, age, breastfeeding and

antenatal care are positive determinants of normal child growth, while maternal parity,

employment, and child sex and age are negative determinants. Interventions to promote child

growth should take into account these differential effects.

Key words: Normal growth, stunting, factors, sub-Saharan Africa, rich households

Introduction

Child health is a fundamental public health issue, for children's sake and because good child health sets one up for life-long health and functioning, and well-being. In sub-Saharan Africa (SSA), child physical health is of particular concern due to the high rates of illness and mortality in the region. Child normal growth, defined in this paper as children who are not stunted (short stature for age), is a foundation for optimal child health and wellbeing. This is because normal child growth is associated positively with cognitive development, higher school achievements, lower morbidity and mortality, higher economic productivity in adulthood and better maternal reproductive outcomes (1-3). This suggests the need for strong investment in nutrition interventions to promote child growth so as to ensure life-long benefits. Working with international partners such as WHO and UNCEF, governments of SSA have put in place various interventions to promote child growth by addressing stunting in the region (4, 5). However, the implementation of these programmes more often than not places more emphasis on child growth deficiencies and how to protect children against the risk factors contributing to these deficiencies (6, 7). This may make it difficult to directly attribute the effects of the programmes on child normal growth outcomes, except to infer that reduction in stunting implies increase in normal growth. The present study intends to fill this gap by providing robust evidence on the positive as well as negative determinants of normal growth among children living in better-off households. Indeed, many experts have called for this type of resource focused approach in promoting child health outcomes, as exemplified by the UNICEF childcare framework (8, 9).

There are a number of factors that affect child linear growth in low and middle incomes countries

(LMICs). These factors include maternal education, employment, household wealth index,

antenatal care (ANC), parity, maternal body mass index (BMI), urban place of residence,

breastfeeding, and maternal age among others. These factors affect child linear growth either

negatively or positively. There is substantial evidence that improvement in maternal education is

associated positively with child linear growth in many settings (10-14). Children whose mothers

are educated tend to have better nutritional status compared to children of mothers with no

education. Studies that examine the effect of maternal BMI have also elucidated the strong

positive association between maternal BMI and child growth (10, 11, 14-17). A study in Ethiopia

showed that maternal BMI was associated positively with children nutritional status (17).

Similarly, maternal age and parity are important factors that affect child growth. Fenske and

colleagues (11) observed that maternal age has significant effect on childhood stunting. Two

other studies established the relationship between maternal age and child nutritional status (18,

19). In India, undernourishment was more prevalent in children of 26-30 year age group mothers

than the other extremes of reproductive ages (18). Relatedly, mothers in their early 20s tend to

have children with poorer health outcomes, including nutrition, while children of older mothers

tend to suffer less from stunting (19, 20). Although, there is a paucity of literature on the effects

of maternal parity on child nutritional status, some few studies have observed negative effects

of parity on child growth (21-23). Further, Kuhnt and Vollmer (24) found in their study that having

at least four ANC visits is associated with reduced odds of stunting in pre-school children. Several

studies have observed a positive effect of breastfeeding on child growth (25-27). Household

wealth index also has a strong positive effect on child growth (28-30). Children in better-off

households tend to have better growth outcomes relative to those in poor households.

The literature reviewed above focus almost exclusively on child growth deficiencies

(undernutrition/abnormal growth) and the factors associated with the same. Statistical analyses

that simultaneously examine the positive as well as negative determinants of normal child

growth are, to the best of our knowledge, still missing. Protective factors against child growth

deficiencies may not necessarily be positive determinants of normal growth, it is significant to

conduct analysis, using normal growth as the key outcome to elucidate the direct relationship

between socio-demographic determinants and positive linear growth among children. It is widely

recognized that children in better-off households tend to have better growth and health

outcomes. However, stratified analysis to understand the key covariates responsible for the

positive growth outcomes in this sub-group is lacking. This study intended to fill this gap by

stratifying the analysis by better-off households, focusing on children who are growing normally

(rather than those that are not) and the factors that make them to grow well. We also

investigated the factors that poses a potential risk (negative determinants) to child growth in

better-off households. This will further our understanding of the key positive and negative

determinants of normal child growth. The objective of this study therefore is to examine the

positive and negative determinants of normal child growth in better-off households.

Methodology

Data sources and sampling strategy

This analysis used data from the Demographic and Health Surveys (DHS)) (31), conducted in

Ghana (2014), Kenya (2014), Nigeria (2013), Mozambique (2011) and Democratic Republic of

Congo (DRC) (2013-2014). The selection of these five countries was informed by our previous

work (12, 32). The DHS data are nationally representative, repeated cross-sectional household

surveys collected primarily in LIMCs every 5 years using standardised questionnaires to enable

cross-country comparisons (33, 34). The DHS utilises a two-stage sample design (35-39). The first

stage involves the selection of sample points or clusters from an updated master sampling frame

constructed from National Population and Housing Census of the respective countries. The

clusters are selected using systematic sampling with probability proportional to size. Household

listing is then conducted in all the selected clusters to provide a sampling frame for the second

stage selection of households. The second stage selection involves the systematic sampling of

the households listed in each cluster, and households to be included in the survey are randomly

selected from the list. The rationale for the second stage selection is to ensure adequate numbers

of completed individual interviews to provide estimates for key indicators with an acceptable

precision. All men and women aged 15-59 and 15-49 respectively, in the selected households

(men in half of the households) are eligible to participate in the surveys if they were either usual

residents of the household or visitors present in the household on the night before the survey.

Study participants

Our study population comprised children aged 0–59 months, born to mothers aged 15–49 years

living in better-off households. Information on children was obtained through face-to-face

interviews with mothers. Height was measured with an adjustable measuring board calibrated in

millimetres. Children younger than 24 months were measured lying down (recumbent length) on

the board, while standing height was measured for older children. The height data were

converted into Z-scores based on the 2006 WHO growth standards (40). The total samples used

in the current analysis were: Ghana, n= 1,247; Nigeria, n= 12,999; Kenya, n= 3,895; Mozambique,

n= 5,711; and DRC, n= 3,943.

Ethics statement

The DHS obtained ethical clearance from Government recognised Ethical Review

Committees/Institutional Review Boards of the respective countries as well as the Institutional

Review Board of ICF International, USA, before the surveys were conducted. Written informed

consent was obtained from the mothers of the children before participation. The authors of this

paper sought and obtained permission from the DHS program for the use of the data. The data

were completely anonymized and therefore the authors did not seek further ethical clearance

before their use.

Outcome and predictor variables

Outcome Variables

The main indicator of child linear growth used in this analysis was child height-for-age Z-scores

(HAZ). The HAZ scores were computed using the 2006 WHO growth standards (40) and classified

into normal growth (or not stunted) and stunted (or poor growth). In this paper, children who

have HAZ equal to or above -2 SD (HAZ ≥ -2SD) (40, 41) were described as having a normal growth,

while children with HAZ below −2 SD (HAZ < -2) from the median HAZ of the WHO reference

population (40) were considered stunted (chronically malnourished).

Stratification variable

We used the household wealth index (WI) as the stratification variable in our analysis. The WI

has been used in many DHS reports to measure inequalities in household characteristics, in the

use of health and other services, and in health outcomes (34, 35, 37, 42). It is an indicator of

wealth that is consistent with expenditure and income measurement among households (33, 34,

37). The index in the DHS dataset was created based on assets ownership and housing

characteristics of each household: type of roofing, and flooring material, source of drinking

water, sanitation facilities, ownership of television, bicycle, motorcycle, automobile among

others. Principal component analysis was employed to assign weights to each asset in each

household. The asset scores were summed up and individuals ranked according to the household

score. The WI was then divided into quintiles: poorest, poorer, middle, richer and richest (33, 34,

37). For the purpose of this analysis, we recoded middle, richer and richest households into

"better-off households". All the analyses were restricted to this sub-category.

Framework underpinning the analysis

The UNICEF conceptual framework (43), which outlines the causes of undernutrition

underpinned our empirical analysis. This is a socio-ecological framework encompassing factors

at the individual, household and societal levels. In the UNICEF framework, child malnutrition can

be analysed in terms of immediate, underlying and basic causes. The immediate causes are

inadequate dietary intakes and infectious disease, the underlying causes are inadequate

maternal and child care, inadequate health services and health environment and the basic causes

are institutional and socio-economic determinants and potential resources (43). However, the

present analysis was guided by the extended UNICEF conceptual framework for childcare,

survival, growth and development (9, 43). The extended UNICEF framework suggests that child

survival, growth and development are influenced by a web of factors, with three underlying

determinants being food security, healthcare and a healthy environment, and care for children

and women (9). These underlying determinants are in turn influenced by basic determinants.

These basic determinants may be described as "exogenous" determinants, which influence child

nutrition through their effect on the intervening proximate determinants (underlying

determinants). In effect, the underlying factors are therefore, endogenously determined by the

exogenous factors (44). In this analysis, we included only the basic factors (socio-demographic)

in our empirical models. We did this because there is evidence that in examining the association

between child growth outcomes and exogenous factors, the proximate factors are usually

excluded to prevent biased and uninterpretable parameters (44-46). Besides the basic factors,

we also included antenatal care (ANC) and breastfeeding practices, which relies mostly on

exogenous public health provisions rather than socio-demographic endowments of the

household (44). The significance of including these two variables in the model is that changes in

them are likely to be more responsive to policies, programmes and interventions rather than to

changes in socio-demographic endowments of the household (44). For example, there is

evidence that policy, institutional and contextual settings are key determinants of the prevalence

of breastfeeding practices (44, 47).

Data analysis

We built two empirical regression models for each of the five countries. In the first models, we

included maternal body mass index (BMI), education, age, work status, parity, breastfeeding

status, marital status, antenatal attendance, sex of household head, household size, number of

children under five years and place of residence. We adjusted for child dietary diversity (DD)—

the details of how the DD is created can be found elsewhere (12), age and sex in the second and

final model. The selection of the explanatory variables was informed by the UNICEF conceptual

framework of child care (9). We estimated adjusted odd ratios (aORs) of the positive and negative

determinants of normal child growth in rich households.

Results

Characteristics of study samples

The results showed that Ghana (87%) had the highest number of children with normal growth

followed by Kenya (80%), while in Mozambique, DRC and Nigeria, the prevalence ranged from

62% to 74%. Regarding dietary diversity intake, Mozambique had the highest prevalence of

children who consumed at least four food groups (21%), with DRC (12%) having the lowest

prevalence. Similarly, Mozambique had the highest number of women with normal weight (73%),

followed by DRC (69%). The prevalence ranged from 43% to 59% in Ghana, Kenya and Nigeria.

Ghana had the highest prevalence (59%) of women who had attained a secondary school

education, while Mozambique had the lowest prevalence (21%). Higher education was less than

15% among women across all countries, with Mozambique (1.3%) registering the lowest

prevalence. DRC had the highest prevalence (86%) ANC attendance among women followed by

Ghana (72%), while Nigeria had the lowest prevalence (19%). Also, Mozambique (35%) had the

highest number of women who were household heads, with Nigeria (14%) having the lowest

women headed households (Table 1).

Table 1: Characteristics of the study samples of the five countries

Variables	DRC		Ghana		Kenya		Mozambique		Nigeria	
Child-level covariates					%/mean	SD	%/mean	SD	%/mean	SD
Height-for-age (HAZ ≥-2)	62.0		87.0		80.0		66.0		74.2	
DD < 4 food groups	87.6		87.0		80.2		78.6		86.9	
DD >= 4 food groups	12.4		14.0		19.8		21.4		13.1	
Sex of child										
Female	50.4		48.0		49		49.2		49.5	
Male	49.6		52.0		51.0		50.8		50.5	
Mother-level covariates										
Body Mass Index (BMI)										
BMI <18.50	9.4		2.9		5.34		4.11		5.42	
BMI = 18.50-24.99	69.2		43.0		55.1		73.9		58.7	
BMI = 25-29.99	16.3		32.0		27.0		17.2		24.6	
BMI >= 30	3.9		22.0		12.6		4.54		10.9	
Education										
No education	12.6		15.0		6.0		24.5		20.6	
Primary education	38.7		18.0		50.8		53.4		24.3	
Secondary education	46.8		59.0		30.5		20.8		43.6	
Higher education	1.8		7.5		12.6		1.26		11.5	
Working status										
Not working	27.2		25.0		36.4		63.3		25.3	
IS working	72.5		74.0		63.4		36.7		74.3	
Parity	4.37	2.58	2.99	1.69	3.127	2.05	3.51	2.12	3.90	2.32
Is Breastfeeding	67.8		53.0		51.3		55.8		51.5	
Marital status										
Not in union	13.3		13.0		14.7		17.3		5.1	
Married	67.1		65.0		79.6		45.0		90.5	
Cohabiting	19.7		22.0		5.62		37.6		4.4	
Number of antenatal visits >=4	86.3		72.0		50.8		45.1		48.2	

Household-level covariates										
Sex of household head										
Household head is Female	19.7		29.0		29.6		34.5		13.9	
Household head is Male	80.3		71.0		70.4		65.5		86.1	
Household size	7.30	2.99	5.0	1.93	5.57	2.31	6.49	2.89	6.73	3.56
Number of children under 5	2.28		1.6	0.69	1.64	0.76	1.92	0.94	2.14	1.11
Community-level covariates										
Urban residence	49.2		73.0		52.3		45.2		55.1	

DD=Dietary diversity; DRC=Democratic Republic of Congo; SD=Standard deviati

Multivariable results of the factors promoting or inhibiting normal child growth

The results of the multivariable analysis of the positive and negative determinants of normal

growth among children under 5 years in better-off households are presented in Tables 2-6. The

results are presented as adjusted odd ratios (aORs).

Positive determinants of normal child growth

The results showed that maternal normal weight (measured by BMI) was associated with

increased odds of normal linear growth among children in Mozambique (aOR=1.82, 95% CI=1.33,

2.50), while overweight associated with increased odds of normal linear growth in DRC

(aOR=1.75, 95% CI=1.17,2.62), Kenya (aOR=2.01, 95% CI= 1.28, 3.16), Mozambique (aOR=2.19,

95%= 1.53, 3.14) and Nigeria (aOR=1.44, 95%=1.16,1.77) relative to children of underweight

mothers. Maternal obesity had similar effect on normal linear growth in Kenya, Mozambique and

Nigeria. One year increase in maternal years of education was associated with increased odds in

normal linear growth among children in DRC (aOR=1.06, 95% CI=1.03, 1.09), Ghana (aOR=1.08,

95% CI=1.04, 1.12), Mozambique (aOR=1.08, 95% CI=1.05, 1.11) and Nigeria (aOR=1.07, 95%

CI=1.06, 1.08). The results in Kenya did not reach statistical significance. An additional year in

maternal age was associated with increased odds of normal linear growth among children in all

the countries included in the analysis. Breastfeeding was associated with increased odds of

normal linear growth in Nigeria (aOR= 1.30, 95% CI=1.16, 1.46) and Kenya (aOR=1.37, 95%

CI=1.05, 1.79). In Ghana, Mozambique and DRC, breastfeeding was positively associated with

normal linear growth in the first model but this statistical significant association disappeared

after the child level covariates such as age, sex and dietary dieversity were included in the final

empirical model. Urban place of residence was associated with increased odds of normal linear

growth among children in DRC (aOR=1.32, 95% CI=1.06, 1.65) and Mozambique (aOR=1.18, 95%

CI=1.01, 1.38). The association did not reach statistical significance in the remaining three

countries.

Negative determinants normal child growth

This section examines the negative determinants of normal linear growth. The results showed

that children of mothers who were working had 25% (aOR=0.75, 95% CI=0.60, 0.93) reduced odds

of normal linear growth in DRC. A unit change in maternal parity was associated with 10%

(aOR=0.90, 95% CI=0.84, 0.97), 23% (aOR=0.77, 95% CI=0.63, 0.93), 25% (aOR=0.75, 95% CI=0.69,

0.82), 6% (aOR=0.94, 95% CI=0.89, 0.99) and 5% (aOR=0.95, 95% CI=0.92, 0.99) reduced odds of

normal linear growth among children in DRC, Ghana, Kenya, Mozambique and Nigeria

respectively. Similarly, a unit change in child's age was associated with reduced odds of normal

linear growth in DRC, Mozambique and Nigeria. Further, a child being a male was associated with

16% (aOR=0.82, 95% CI=0.68, 0.98), 40% (aOR=0.60, 95% CI=0.40, 0.89), 37% (aOR=0.63, 95%

CI=0.51, 0.77) and 21% (aOR=0.79, 95% CI=0.71, 0.87) reduced odds of normal linear growth

among children in DRC, Ghana, Kenya and Nigeria respectively. The association in Mozambique

did not reach statistical significance.

291   Table 2: Adjusted odd ratios (aOR) of positive and negative determinants of normal linear growth
292   among children living in better-off households, DRC

Variables	Model 1	Model 2
Mother-level covariates		
BMI (kg/m^2) = 18.50 - 24.99	1.093	1.124
	(0.782 - 1.526)	(0.796 - 1.589)
BMI (kg/m^2) = 25 - 29.99	1.685***	1.751***
	(1.138 - 2.493)	(1.172 - 2.617)
BMI (kg/m^2) >= 30	1.573	1.729
	(0.855 - 2.892)	(0.884 - 3.382)
Maternal education (in single years)	1.061***	1.060***
	(1.032 - 1.090)	(1.030 - 1.091)
Age of the mother (in years)	1.039***	1.051***
	(1.015 - 1.063)	(1.026 - 1.077)
Working status = Is working	0.741***	0.747**
	(0.597 - 0.920)	(0.597 - 0.934)
Parity	0.902***	0.899***
	(0.844 - 0.965)	(0.838 - 0.965)
Is Breastfeeding = YES	1.678***	1.184
	(1.370 - 2.055)	(0.943 - 1.487)
Marital Status = Married	1.034	1.076
	(0.756 - 1.415)	(0.786 - 1.473)
Marital Status = Cohabiting	0.976	0.994
	(0.687 - 1.389)	(0.697 - 1.418)
Number of antenatal visits = 4+ visits	1.589***	1.159
	(1.296 - 1.947)	(0.923 - 1.457)
Household-level covariates		
Head of HH is Male	0.865	0.868
	(0.681 - 1.098)	(0.678 - 1.111)
Household size	1.007	1.002
	(0.968 - 1.048)	(0.962 - 1.043)
Number of children under 5 years	0.945	0.954
	(0.844 - 1.059)	(0.850 - 1.071)
Community-level covariates		
Urban residence = Urban	1.322**	1.320**
	(1.066 - 1.640)	(1.059 - 1.646)
Child-level covariates		
Dietary Diversity (DD) >= 4		1.294*
		(0.971 - 1.724)
Age of the child (in months)		0.975***
		(0.969 - 0.982)
Sex of child = Male		0.815**
		(0.677 - 0.982)
Observations	**3,943**	**3,943**

95% Confidence Intervals (CIs) in parentheses; DD-Dietary diversity; HH-Household; BMI-Body mass index
*** p<0.01, ** p<0.05, * p<0.1

Table 3: Adjusted odd ratios (aOR) of positive and negative determinants of normal linear growth among children living in better-off households, Ghana

Variables	Model 1	Model 2
Mother-level covariates		
BMI (kg/m^2) = 18.50 - 24.99	0.692	0.725
	(0.224 - 2.136)	(0.241 - 2.181)
BMI (kg/m^2) = 25 - 29.99	1.230	1.270
	(0.385 - 3.929)	(0.410 - 3.931)
BMI (kg/m^2) >= 30	1.617	1.700
	(0.478 - 5.469)	(0.514 - 5.628)
Maternal education (in single years)	1.078***	1.081***
	(1.037 - 1.121)	(1.039 - 1.124)
Age of the mother (in years)	1.047*	1.052**
	(0.998 - 1.099)	(1.001 - 1.105)
Working status = Is working	1.276	1.264
	(0.821 - 1.983)	(0.804 - 1.987)
Parity	0.779**	0.766***
	(0.643 - 0.944)	(0.630 - 0.932)
Is Breastfeeding = YES	1.769**	1.748*
	(1.129 - 2.771)	(0.981 - 3.116)
Marital Status = Married	1.956**	2.104**
	(1.032 - 3.708)	(1.101 - 4.022)
Marital Status = Cohabiting	1.524	1.659
	(0.783 - 2.967)	(0.849 - 3.240)
Number of antenatal visits = 4+ visits	1.124	1.110
	(0.695 - 1.817)	(0.646 - 1.906)
Household-level covariates		
Head of HH is Male	1.250	1.200
	(0.783 - 1.996)	(0.753 - 1.911)
Household size	0.998	0.998
	(0.894 - 1.114)	(0.891 - 1.117)
Number of children under 5	0.834	0.843
	(0.555 - 1.253)	(0.559 - 1.274)
Community-level covariate		
Urban residence = Urban	1.020	0.988
	(0.640 - 1.628)	(0.623 - 1.565)
Child-level covariates		
Dietary Diversity (DD) >= 4		1.022
		(0.590 - 1.770)
Age of the child (in months)		0.998
		(0.982 - 1.014)
Sex of child = Male		0.596**
		(0.400 - 0.887)
Observations	**1,247**	**1,247**

95% Confidence Intervals (CIs) in parentheses; DD-Dietary diversity; HH-Household; BMI-Body mass index
*** p<0.01, ** p<0.05, * p<0.1

Table 4: Adjusted odd ratios (aOR) of positive and negative determinants of normal linear growth among children living in better-off households, Kenya

Variables	Model 1	Model 2
Mother-level covariates		
BMI (kg/m^2) = 18.50 - 24.99	1.245	1.313
	(0.824 - 1.881)	(0.872 - 1.976)
BMI (kg/m^2) = 25 - 29.99	1.880***	2.014***
	(1.195 - 2.960)	(1.284 - 3.158)
BMI (kg/m^2) >= 30	1.860**	1.991**
	(1.082 - 3.195)	(1.165 - 3.402)
Maternal education (in single years)	1.017	1.021
	(0.984 - 1.051)	(0.987 - 1.055)
Age of the mother (in years)	1.074***	1.075***
	(1.046 - 1.102)	(1.046 - 1.104)
Working status = Is working	0.969	0.982
	(0.775 - 1.211)	(0.786 - 1.227)
Parity	0.752***	0.752***
	(0.688 - 0.823)	(0.687 - 0.822)
Is Breastfeeding = YES	1.373***	1.367**
	(1.103 - 1.710)	(1.047 - 1.785)
Marital Status = Married	1.131	1.112
	(0.826 - 1.547)	(0.810 - 1.526)
Marital Status = Cohabiting	1.278	1.233
	(0.785 - 2.079)	(0.756 - 2.009)
Number of antenatal visits = 4+ visits	1.132	1.095
	(0.906 - 1.415)	(0.862 - 1.390)
Household-level covariates		
Head of HH is Male	0.986	0.987
	(0.781 - 1.245)	(0.780 - 1.248)
Household size	1.054*	1.047
	(0.991 - 1.122)	(0.984 - 1.114)
Number of children under 5	1.018	1.026
	(0.871 - 1.189)	(0.878 - 1.200)
Community-level covariate		
Urban residence = Urban	0.883	0.890
	(0.719 - 1.085)	(0.723 - 1.096)
Child-level covariates		
Dietary Diversity (DD) >= 4		0.784*
		(0.597 - 1.030)
Age of the child (in months)		0.996
		(0.989 - 1.003)
Sex of child = Male		0.625***
		(0.507 - 0.770)
Observations	**3,985**	**3,985**

95% Confidence Intervals (CIs) in parentheses; DD-Dietary diversity; HH-Household; BMI-Body mass index
*** p<0.01, ** p<0.05, * p<0.1

Table 5: Adjusted odd ratios (aOR) of positive and negative determinants of normal linear growth
among children living in better-off households, Mozambique

Variables	Model 1	Model 2
Mother-level covariates		
BMI (kg/m^2) = 18.50 - 24.99	1.780***	1.823***
	(1.303 - 2.430)	(1.329 - 2.500)
BMI (kg/m^2) = 25.00 - 29.99	2.144***	2.190***
	(1.502 - 3.060)	(1.528 - 3.137)
BMI (kg/m^2) >= 30	4.106***	4.268***
	(2.453 - 6.870)	(2.537 - 7.179)
Maternal education (in single years)	1.077***	1.079***
	(1.052 - 1.102)	(1.054 - 1.105)
Age of the mother (in years)	1.035***	1.039***
	(1.018 - 1.052)	(1.022 - 1.056)
Working status = Is working	0.941	0.950
	(0.813 - 1.088)	(0.821 - 1.098)
Parity	0.943**	0.942**
	(0.894 - 0.995)	(0.892 - 0.994)
Is Breastfeeding = YES	1.240***	1.070
	(1.068 - 1.439)	(0.906 - 1.263)
Marital Status = Married	0.977	1.009
	(0.787 - 1.213)	(0.811 - 1.255)
Marital Status = Cohabiting	1.278**	1.289**
	(1.035 - 1.578)	(1.043 - 1.593)
Number of antenatal visits = 4+ visits	1.055	0.942
	(0.911 - 1.222)	(0.807 - 1.100)
Household-level covariates		
Head of HH is Male	0.874*	0.887
	(0.746 - 1.024)	(0.757 - 1.039)
Household size	1.018	1.016
	(0.988 - 1.048)	(0.986 - 1.046)
Number of children under 5	0.998	1.009
	(0.908 - 1.098)	(0.916 - 1.111)
Community-level covariate		
Urban residence = Urban	1.185**	1.181**
	(1.014 - 1.385)	(1.010 - 1.381)
Child-level covariates		
Dietary Diversity (DD) >= 4		0.888
		(0.752 - 1.048)
Age of the child (in months)		0.991***
		(0.986 - 0.995)
Sex of child = Male		0.931
		(0.811 - 1.069)
Observations	**5,711**	**5,711**

95% Confidence Intervals (CIs) in parentheses; DD-Dietary diversity; HH-Household; BMI-Body mass index
*** p<0.01, ** p<0.05, * p<0.1

Table 6: Adjusted odd ratios (aOR) of positive and negative determinants of normal linear growth

among children living in better-off households, Nigeria

Variables	Model 1	Model 2
Mother-level covariates		
BMI (kg/m^2) = 18.50 - 24.99	1.135	1.129
	(0.934 - 1.379)	(0.931 - 1.371)
BMI (kg/m^2) = 25.00 - 29.99	1.434***	1.435***
	(1.157 - 1.777)	(1.159 - 1.777)
BMI (kg/m^2) >= 30	1.697***	1.706***
	(1.309 - 2.200)	(1.316 - 2.211)
Maternal education (in single years)	1.067***	1.068***
	(1.056 - 1.079)	(1.057 - 1.080)
Age of the mother (in years)	1.033***	1.039***
	(1.021 - 1.045)	(1.027 - 1.051)
Working status = Is working	1.072	1.106*
	(0.958 - 1.199)	(0.988 - 1.238)
Parity	0.957***	0.954***
	(0.925 - 0.989)	(0.922 - 0.986)
Is Breastfeeding = Yes	1.537***	1.303***
	(1.388 - 1.703)	(1.162 - 1.461)
Marital Status = Married	0.806*	0.803*
	(0.625 - 1.039)	(0.622 - 1.037)
Marital Status = Cohabiting	0.866	0.846
	(0.606 - 1.237)	(0.592 - 1.209)
Number of antenatal visits = 4+ visits	1.080	0.877**
	(0.975 - 1.197)	(0.776 - 0.990)
Household-level covariates		
Head of HH is Male	0.850**	0.856*
	(0.728 - 0.992)	(0.733 - 1.000)
Household size	0.980**	0.980**
	(0.963 - 0.998)	(0.962 - 0.998)
Number of children under 5	0.946*	0.941**
	(0.893 - 1.002)	(0.888 - 0.997)
Community-level covariate		
Urban residence = Urban	1.004	1.013
	(0.910 - 1.108)	(0.917 - 1.118)
Child-level covariates		
Dietary Diversity (DD) >= 4		1.055
		(0.904 - 1.231)
Age of the child (in months)		0.988***
		(0.984 - 0.991)
Sex of child = Male		0.785***
		(0.712 - 0.865)
Observations	**12,999**	**12,999**

95% Confidence Intervals (CIs) in parentheses; DD-Dietary diversity; HH-Household; BMI-Body mass index
*** p<0.01, ** p<0.05, * p<0.1

Discussion

This study investigated the positive and negative determinants of normal linear growth among

children living in better-off households in DRC, Ghana, Kenya, Mozambique and Nigeria. The

results shed new insights on both positive and negative determinants of child growth, and

country-specific variations of these effects in the five countries. In the current analysis, higher

maternal weight (measured by BMI) tends to have significant positive effect on normal growth

among children living in better-off households in all countries except Ghana. Thus, maternal

weight is a key determinant of positive child growth outcomes. Despite the fact that higher

maternal weight has positive effect on child normal growth, interventions should target at

increasing maternal weight qualitatively for the benefit of both the mother and child. This is

critical because of the negative effect of unhealthy weight on maternal health outcomes (48-50).

These findings are in line with the literature. Maternal nutrition was found to have a significant

positive effect on child linear growth in many settings (10-12). In India, BMI among other

variables, was found to have significantly effect on child linear growth (11). The preceding

discussion illuminated the crucial role maternal nutrition plays in improving child nutritional

status, even though the pathways through which this happens may be complex.

The importance role maternal education plays in promoting positive child health outcomes was

observed in the present study. Our results showed that maternal years of education have

significant positive effects on normal linear growth in all the five countries except Kenya. This

suggests that education has the potential to promote normal growth of children living in better-

off households. Interventions to promote child growth may have positive impact of normal linear

growth of children living in these households. The beneficial effects of maternal years of

education on child growth has previously been documented. An improvement in maternal

education was associated positively with dramatic change in linear growth among pre-school

children (10, 11). This may be the case because educated mothers tend to utilize both preventive

and curative health care more (51, 52). Educated mothers also tend to have more strongly

committed attitude towards good childcare than uneducated mothers (53, 54). Furthermore, the

more education the mother has the more the likelihood that she is sensitive and responsive to

caregiving duties (53, 54). Also, there is evidence that children seemingly engage more positively

with their mothers when maternal education is greater (54). All the above have positive effects

on child growth outcomes. The literature discussed above together with our study clearly

demonstrated the importance of maternal education for positive child health outcomes.

The health benefits of breastfeeding were illuminated in this study but only in two countries.

Breastfeeding was found to associate with the likelihood of normal linear growth among children

living in better-off households in Kenya and Nigeria. This suggests that mothers in these

households should be encouraged to practice breastfeeding because of its beneficial effects on

their children growth. These findings confirm the widely recognised benefits of breastfeeding

for improved health and developmental outcomes (25-27). In the contrary, breastfeeding

showed a significant positive effect on child normal linear growth in the models containing only

the socio-demographic factors in Ghana, Mozambique and DRC. This significant association

disappeared after child level covariates such as dietary diversity, age and sex were included in

the final empirical models. This suggests that whether breastfeeding will have positive effect on

normal child growth in better-off households or not is conditional on the inclusion or otherwise

of child level co-variates. This corroborates previous research, which suggests that child level

factors must be considered when evaluating the association of breastfeeding with

anthropometric outcomes (55). The non-significant positive effect of breastfeeding on child

growth has previously been documented (55-57). Thus, while breastfeeding is critical for positive

child health outcomes, it is not always the case that its effects would be statistically significant.

Our analysis also illuminated negative determinants of normal linear growth. Maternal work

status is inversely related to normal child growth in DRC. This implies that DRC mothers who are

engaged in any form of work tend to have children who have poor linear growth relative to

mothers who are not working. This negative effect may boil down to inadequate childcare due

to limited time available to working mothers. A study in India concluded that a mother's

employment compromises infant feeding and care, particularly so when mothers are not able to

get alternative caregivers (58). This study further reported that the compromises related to

childcare and feeding outweigh the benefits from employment (58). Other studies have shown

that mothers working away from home spend less time with their children compared to mothers

who are not working outside the home and therefore likely to have children who are

undernourished (59, 60). Although, it is a well-documented fact that women who are working

tend to have access to disposal income and therefore able to provide nutritious food as well as

other care services to their children (13, 17, 61, 62), the above discussion showed that maternal

employment, indeed, could have negative effect on child growth outcomes. It is worthy to note

that this analysis did not investigate the categories of employment and their effect on child

growth. We are therefore unable to tell the independent effect of the various occupational

groups on child linear growth. This is a limitation worth noting, as different occupations may have

different effects.

Another negative growth determinant is maternal parity (number of pregnancies reaching viable

gestational age). The results show that higher maternal parity has negative effect on normal

linear growth among children living in better-off households. The effect is largest in Kenya (25%)

followed by Ghana (23%), with the least effect being in Nigeria (5%). This suggests that higher

parity has a stronger negative effect on child normal growth in Kenya and Ghana relative to the

other countries. The negative effects of parity on child growth may be attributed to it being an

important factor that affect maternal depletion, particular so among high fertility mothers (63,

64), which has the potential to compromise the mothers ability to provide good care for their

children. The consequential effect of lack of good care is poor child growth. Secondly, women

with higher parity are likely to have many young children, who might compete for the available

care resources, which can have a negative effect on good care practices and consequently their

children growth outcomes. Previous studies have documented that the higher the maternal

parity, the less likelihood that their children will have positive growth outcomes (21, 22). For

example, children from multiparous mothers were found to have lower rates of accelerated

infant growth and lower levels of childhood body mass index than children of nulliparous mothers

(22). The preceding discussion clearly demonstrated that parity is a key negative determinant of

of child linear growth.

Similarly, two child biological factors (sex and age) were found to associate negatively with

normal linear growth. A year increase in child's age was associated negatively with normal linear

growth in three (DRC, Mozambique and Nigeria) of the five countries. This suggests that older

children living in better-off households have a less likelihood of achieving a normal linear growth.

These findings are consistent with previous research. Nshimyiryo and colleagues (65) observed

that increasing age of the child had a significant association with poor linear growth. For instance,

children aged 6–23 months were at lower risk of poor growth than those in the older age group

24–59 months (65). Also, being a male child affects the likelihood of children living in better-off

households to have a normal linear growth. Previous work has shown that poor linear growth

was higher among male children as compared to female children (66). Suggesting that male

children tend to be more vulnerable to poor growth than their female counterparts in the same

age group (67). This might be due to preferences in feeding practices or other types of exposures

(67). It could also be explained by the fact that boys are expected to grow at a slightly more rapid

rate compared to girls and their growth is perhaps more easily affected by nutritional deficiencies

or other exposures (68).

413 **Strengths and limitations of the study**

414 This study has some strengths. One important strength is the use of high quality large nationally

415 representative DHS data to investigate the positive and negative determinants of normal linear

416 growth among children living in rich households. This makes it possible for the findings to be

417 generalized to the population of young children in the respective countries. The large data also

418 help to produce more robust estimates of observed associations. The use of multi-country data

419 unmask differences and highlights commonalities in the effects of the correlates on child growth

420 across countries. This may not have been possible with single country data. Further, the height

421 data used for computing the HAZ indicator were objectively measured, reducing possible

422 misclassification. The novelty of this study is its focus on positive child growth outcomes rather

423 than child growth deficiencies. A limitation worth mentioning is the cross-sectional nature of the

424 data, which makes it difficult to disentangle potential reciprocal and otherwise complex causal

425 relationships. The conclusions in this paper are therefore meant to be restricted to statements

426 about the associations between the explanatory variables and the outcome variables.

427

428

429 **Conclusions**

430 Maternal weight (BMI) tends to have significant positive effects on normal linear growth among

431 children living in better-off households across all countries except Ghana. Interventions aimed at

432 increasing maternal weight qualitatively are likely to be effective in improving the linear growth

433 of children living in better-off households. Maternal years of education have significant positive

effects on normal linear growth in all the five countries except Kenya. This suggests that

education has the potential to promote normal linear growth among of children living in better-

off households. Breastfeeding was associated with the likelihood of normal linear growth in

Kenya and Nigeria. This implies that mothers in better-off households should be encouraged to

practice breastfeeding because of its beneficial effect on their children growth. Maternal work

status is inversely related to normal growth in DRC. Thus, in DRC, mothers who are engaged in

any form of work tend to have children with poor growth relative to mothers who are not

working. The results show that higher maternal parity associate negatively with normal linear

growth. The effect is largest in Kenya (25%) followed by Ghana (23%), with the least effect being

in Nigeria (5%). In effect, higher parity has a stronger negative effect on child normal linear

growth in Kenya and Ghana relative to the other countries.

Acknowledgements

We wish to express our profound gratitude to The DHS Program, USA for providing us access to

the data. We also wish to acknowledge institutions of respective countries that played critical

roles in the data collection process.

Competing Interest

The authors have no competing interests to declare.

Funding

This study did not receive funding from any source.

Data Sharing Statement

This study was a re-analysis of existing data that are publicly available from The DHS Program at

http://dhsprogram.com/publications/publication-fr221-dhs-final-reports.cfm.   Data   are

accessible free of charge upon a registration with the Demographic and Health Survey program

(The DHS Program). The registration is done on the DHS website indicated above.

Authors' Contribution

DAA conceived and designed the study, interpreted the results, wrote the first draft of the

manuscript, and contributed to revision of the manuscript. DAA and ZTD analysed the data. ZTD

and EWK contributed to study design, data interpretation, and critical revision of the manuscript.

All authors take responsibility of any issues that might arise from the publication of this

manuscript.

References

1.   Kowalski AJ, Georgiadis A, Behrman JR, Crookston BT, Fernald LCH, Stein AD. Linear Growth through 12 Years is Weakly but Consistently Associated with Language and Math Achievement Scores at Age 12 Years in 4 Low- or Middle-Income Countries. The Journal of nutrition. 2018;148(11):1852-9.

2.   Sudfeld CR, McCoy DC, Danaei G, Fink G, Ezzati M, Andrews KG, et al. Linear growth and child development in low- and middle-income countries: a meta-analysis. Pediatrics. 2015;135(5):e1266-75.

3.   Prendergast AJ, Humphrey JH. The stunting syndrome in developing countries. Paediatrics and International Child Health. 2014;34(4):250-65.

4.   WHO. Accelerating nutrition improvements in Sub-Saharan Africa: scaling up nutrition interventions: Final report 2012-2016. Geneva; 2017.

5.   Lartey A. Maternal and child nutrition in Sub-Saharan Africa: challenges and interventions. The Proceedings of the Nutrition Society. 2008;67(1):105-8.

6.   Hossain M, Choudhury N, Adib Binte Abdullah K, Mondal P, Jackson AA, Walson J, et al. Evidence-based approaches to childhood stunting in low and middle income countries: a systematic review. Archives of Disease in Childhood. 2017;102(10):903.

7.   Akombi BJ, Agho KE, Merom D, Renzaho AM, Hall JJ. Child malnutrition in sub-Saharan Africa: A meta-analysis of demographic and health surveys (2006-2016). PloS one. 2017;12(5):e0177338-e.

8.   Engle P. The role of caring practices and resources for care in child survival, growth, and developments: South and Southeast Asia Asian Development Review. 1999;17(1,2):132-67.

9.   Engle PL, Menon P, Haddadb L. Care and nutrition: concepts and measurements. Washington, DC: UNICEF/International Food Policy Institute. 1997.

10.   Uttamacharya U, Arokiasamy P, Verma R. Determinants of child Anthropometrics in India: A quantile regression analysis 2013 [Available from: http://iussp.org/en/event/17/programme/paper/3785.

11.   Fenske N, Burns J, Hothorn T, Rehfuess EA. Understanding Child Stunting in India: A Comprehensive Analysis of Socio-Economic, Nutritional and Environmental Determinants Using Additive Quantile Regression PloS one. 2013;8(11).

12.   Amugsi DA, Dimbuene ZT, Kimani-Murage EW, Mberu B, Ezeh AC. Differential effects of dietary diversity and maternal characteristics on linear growth of children aged 6-59 months in sub-Saharan Africa: a multi-country analysis. Public health nutrition. 2017;20(6):1029-45.

13.   Keino S, Plasqui G, Ettyang G, van den Borne B. Determinants of stunting and overweight among young children and adolescents in sub-Saharan Africa Food Nutr Bull. 2014;35(2):167-78.

14.   Rachmi CN, Agho KE, Li M, Baur LA. Stunting, Underweight and Overweight in Children Aged 2.0-4.9 Years in Indonesia: Prevalence Trends and Associated Risk Factors PloS one. 2016;11(5):e0154756.

15.   Felisbino-Mendes MS, Villamor E, Velasquez-Melendez G. Association of maternal and child nutritional status in Brazil: a population based cross-sectional study PloS one. 2014;9(1):e87486.

16.   Subramanian SV, Ackerson LK, Davey Smith G, John NA. Association of Maternal Height With Child Mortality, Anthropometric Failure, and Anemia in India JAMA2009. 2009;301(16):1691-701.

17.   Negash C, Whiting SJ, Henry CJ, Belachew T, Hailemariam TG. Association between Maternal and Child Nutritional Status in Hula, Rural Southern Ethiopia: Across Sectional Study PloS one. 2015;10(11):e0142301.

18.   Sonkaria L, Zafer A, Gaur KL, Manohar RK. Maternal Factors Associated with Nutritional Status of 1-5 years Children Residing in Field Practice Area of Rural Health Training Centre Naila, Jaipur (Rajasthan) India Natl J Community Med 2014;5(3):283-7.

19.   Finlay JE, Özaltin E, Canning D. The association of maternal age with infant mortality, child
anthropometric failure, diarrhoea and anaemia for first births: evidence from 55 low- and middle-income
countries BMJ Open. 2011;1(2).
20.   Fall CH, Sachdev HS, Osmond C, Restrepo-Mendez MC, Victora C, Martorell R, et al. COHORTS
investigators. Association between maternal age at childbirth and child and adult outcomes in the
offspring: a prospective study in five low-income and middle-income countries (COHORTS collaboration)
Lancet Glob Health. 2015;3(7):e366-77.
21.   Amugsi DA, Mittelmark MB, Lartey A. Dietary Diversity is a Predictor of Acute Malnutrition in Rural
but Not in Urban Settings: Evidence from Ghana British Journal of Medicine and Medical Research.
2014;4(25):4310–24.
22.   Gaillard R, Rurangirwa Akashi A, Williams Michelle A, Hofman A, Mackenbach Johan P, Franco
Oscar H, et al. Maternal Parity, Fetal and Childhood Growth, and Cardiometabolic Risk Factors.
Hypertension. 2014;64(2):266-74.
23.   Thuita FM, Mwadime RKN, Wang'ombe JK. Child nutritional status and maternal factors in an
urban slum in Nairobi, Kenya East African Medical Journal 2005;82(4):209-15.
24.   Kuhnt J, Vollmer S. Antenatal care services and its implications for vital and health outcomes of
children: evidence from 193 surveys in 69 low-income and middle-income countries. BMJ open.
2017;7(11):e017122-e.
25.   WHO. Continued breastfeeding for healthy growth and development of children 2019 [cited 2019
03.09]. Available from: https://www.who.int/elena/titles/continued_breastfeeding/en/.
26.   Yan J, Liu L, Zhu Y, Huang G, Wang PP. The association between breastfeeding and childhood
obesity: a meta-analysis. BMC public health. 2014;14:1267-.
27.   Kramer MS, Aboud F, Mironova E, Vanilovich I, Platt RW, Matush L, et al. Breastfeeding and child
cognitive development: new evidence from a large randomized trial. Archives of general psychiatry.
2008;65(5):578-84.
28.   Hong R, Banta JE, Betancourt JA. Relationship between household wealth inequality and chronic
childhood under-nutrition in Bangladesh. International journal for equity in health. 2006;5:15-.
29.   Hong R, Mishra V. Effect of wealth inequality on chronic under-nutrition in Cambodian children.
Journal of health, population, and nutrition. 2006;24(1):89-99.
30.   Mostafa Kamal SM, Jr., Rosliza AM, Aynul M. Effects of Wealth on Nutritional Status of Pre-school
Children in Bangladesh. Malaysian journal of nutrition. 2010;16(2):219-32.
31.   The   DHS   Program.   Demographic   and   Health   Surveys   [Available   from:
http://dhsprogram.com/data/available-datasets.cfm.
32.   Amugsi DA, Dimbuene ZT, Kyobutungi C. Correlates of the double burden of malnutrition among
women: an   analysis   of   cross   sectional   survey   data   from   sub-Saharan   Africa.   BMJ   Open.
2019;9(7):e029545.
33.   The DHS Program. Demographic and Health Surveys [cited 2018 06.09]. Available from:
https://dhsprogram.com/data/dataset_admin/download-datasets.cfm [
34.   The DHS Program. DHS Methodology   [Available from: http://dhsprogram.com/What-We-
Do/Survey-Types/DHS-Methodology.cfm.
35.   Ghana Statistical Service (GSS), Ghana Health Service (GHS), ICF Macro. GhanaDemographic and
Health Survey 2008 Accra, Ghana: GSS, GHS, and ICF Macro; 2009.
36.   Ministry of Health and Social Services (MoHSS), Macro International Inc. Namibia Demographic
and Health Survey 2006-07 Windhoek, Namibia and Calverton, Maryland, USA: MoHSS and Macro
International Inc. ; 2008.
37.   Ghana Statistical Service (GSS), Ghana Health Service (GHS), ICF International. Ghana
Demographic and Health Survey 2014. Rockville, Maryland, USA: GSS, GHS, and ICF International.; 2015.

38.   National Population Commission (NPC) [Nigeria], ICF International. Nigeria Demographic and Health Survey 2013. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF International. . 2014.

39.   National Bureau of Statistics-Kenya, ICF International. 2014 KDHS Key Findings. Rockville, Maryland, USA: KNBS and ICF International. 2015.

40.   WHO Multicentre Growth Reference Study Group. WHO Child Growth Standards based on length/height, weight and age Acta paediatrica (Oslo, Norway : 1992) Supplement. 2006;450:76-85.

41.   Faye CM, Fonn S, Levin J, Kimani-Murage E. Analysing child linear growth trajectories among under-5 children in two Nairobi informal settlements. Public health nutrition. 2019;22(11):2001-11.

42.   The Nambia Ministry of Health and Social Services (MoHSS) and ICF International, International. The Namibia Demographic and Health Survey 2013 Windhoek, Namibia, and Rockville, Maryland, USA: MoHSS and ICF. 2014.

43.   UNICEF. The State of World's Children: Causes of Child Malnutrition 1998 [Available from: https://www.unicef.org/sowc98/fig5.htm.

44.   Zanello G, Srinivasan CS, Shankar B. What Explains Cambodia's Success in Reducing Child Stunting-2000-2014? PLoS one. 2016;11(9):e0162668.

45.   Smith LC, Ruel MT, Ndiaye A. Why is child malnutrition lower in urban than in rural areas? Evidence from 36 developing countries World Devel. 2005;33(8):1285-305.

46.   Sastry N. What explains rural-urban differentials in child mortality in Brazil? Social science & medicine (1982). 1997;44(7):989-1002.

47.   Rollins NC, Bhandari N, Hajeebhoy N, Horton S, Lutter CK, Martines JC, et al. Why invest, and what it will take to improve breastfeeding practices? Lancet. 2016;387(10017):491-504.

48.   Abrha S, Shiferaw S, Ahmed KY. Overweight and obesity and its socio-demographic correlates among urban Ethiopian women: evidence from the 2011 EDHS BMC Public Health. 2016;636.

49.   Agyemang C, Owusu-Dabo E, de Jonge A, Martins D, Ogedegbe G, Stronks K. Overweight and obesity among Ghanaian residents in The Netherlands: how do they weigh against their urban and rural counterparts in Ghana? Public health nutrition2008.

50.   Amugsi DA, Dimbuene ZT, Mberu B, Muthuri S, Ezeh AC. Prevalence and time trends in overweight and obesity among urban women: an analysis of demographic and health surveys data from 24 African countries, 1991–2014. BMJ Open. 2017;7(10):e017344.

51.   Addai I. Determinants of use of maternal-child health services in rural Ghana. Journal of biosocial science. 2000;32(1):1-15.

52.   Zegeye EA, Mbonigaba J, Dimbuene ZT. Factors associated with the utilization of antenatal care and prevention of mother-to-child HIV transmission services in Ethiopia: applying a count regression model. BMC women's health. 2018;18(1):187.

53.   Richman AL, Miller PM, LeVine RA. Cultural and educational variations in maternal responsiveness. Developmental Psychology. 1992;28(4):614-21.

54.   NICHD. Early Child Care Research Network. Child care and mother-child interaction in the first 3 years of life. NICHD Early Child Care Research Network Dev Psychol 1999;35(6):1399-413. .

55.   Marquis GS, Habicht JP, Lanata CF, Black RE, Rasmussen KM. Association of breastfeeding and stunting in Peruvian toddlers: an example of reverse causality. International journal of epidemiology. 1997;26(2):349-56.

56.   Amugsi DA, Mittelmark MB, Lartey A, Matanda DJ, Urke HB. Influence of childcare practices on nutritional status of Ghanaian children: a regression analysis of the Ghana Demographic and Health Surveys. BMJ Open. 2014;4.

57.   Mulder-Sibanda M, Sibanda-Mulder FS. Prolonged breastfeeding in Bangladesh: indicators of inadequate feeding practices or mothers' response to children's poor health? Public Health. 1999;113(2):65-8.

58.   Nair M, Ariana P, Webster P. Impact of mothers' employment on infant feeding and care: a
qualitative study of the experiences of mothers employed through the Mahatma Gandhi National Rural
Employment Guarantee Act. BMJ Open. 2014;4(4):e004434.
59.   Abbi R, Christian P, Gujral S, Gopaldas T. The impact of maternal work on the nutrition and health
status of children Food and Nutrition Bulletin 1991;13(1):20-4.
60.   Popkin BM. Time allocation of the mother and child nutrition. Ecology of Food and Nutrition
1980;9:1-14.
61.   Sharaf MF, Rashadb AS. Economic and Socio-Demographic Determinants of Child Nutritional
Status in Egypt: A Comprehensive Analysis using Quantile Regression Approach. Working Paper No. 2016-
04: University of Alberta 2016 [Available from: http://www.ualberta.ca/~econwps/2016/wp2016-04.pdf.
62.   Abuya BA, Ciera J, Kimani-Murage E. Effect of mother's education on child's nutritional status in
the slums of Nairobi BMC Pediatr 2012;12:80.
63.   Zerihun T, Larson CP, Hanley JA. Anthropometric status of Oromo women of child bearing age in
rural south-western Ethiopia Ethiopian Journal of Health Development 1997;11(3):1-7.
64.   Teller H, Yimar G. Levels and determinants of malnutrition in adolescent and adult women in
southern Ethiopia Ethiopian Journal of Health Development 2000;14(1):57-66.
65.   Nshimyiryo A, Hedt-Gauthier B, Mutaganzwa C, Kirk CM, Beck K, Ndayisaba A, et al. Risk factors
for stunting among children under five years: a cross-sectional population-based study in Rwanda using
the 2015 Demographic and Health Survey. BMC Public Health. 2019;19(1):175.
66.   Mzumara B, Bwembya P, Halwiindi H, Mugode R, Banda J. Factors associated with stunting among
children below five years of age in Zambia: evidence from the 2014 Zambia demographic and health
survey. BMC Nutrition. 2018;4(1):51.
67.   Wamani H, Astrom AN, Peterson S, Tumwine JK, Tylleskar T. Boys are more stunted than girls in
sub-Saharan Africa: a meta-analysis of 16 demographic and health surveys. BMC Pediatr. 2007;7:17.
68.   Condo JU, Gage A, Mock N, Rice J, Greiner T. Sex differences in nutritional status of HIV-exposed
children in Rwanda: a longitudinal study. Tropical medicine & international health : TM & IH.
2015;20(1):17-23.