1

Virulence potential of a multidrug-resistant *Escherichia coli* strain belonging to the emerging clonal group ST101-B1 isolated from bloodstream infection

Ana Carolina de Mello Santos[1*], Rosa Maria Silva[1], Tiago Barcelos Valiatti[1], Fernanda Fernandes dos Santos[1], José Francisco Santos-Neto[1], Rodrigo Cayô[1,2], Ana Paula Streling[3], Carolina Silva Nodari[3], Ana Cristina Gales[3], Milton Yutaka Nishiyama-Junior[4], Eneas Carvalho[5], Tânia Aparecida Tardelli Gomes[1].

[1] Universidade Federal de São Paulo - Escola Paulista de Medicina (UNIFESP-EPM), Disciplina de Microbiologia, Departamento de Microbiologia, Imunologia e Parasitologia - DMIP, São Paulo - SP, Brazil.

[2] Universidade Federal de São Paulo - UNIFESP, Laboratório de Imunologia e Bacteriologia - LIB, Setor de Biologia Molecular, Microbiologia e Imunologia, Departamento de Ciências Biológicas - CDB, Instituto de Ciências Ambientais, Químicas e Farmacêuticas - ICAQF, Diadema - SP, Brazil.

[3] Universidade Federal de São Paulo - Escola Paulista de Medicina – (UNIFESP-EPM), Laboratório Alerta, Disciplina de Infectologia, Departamento de Medicina, São Paulo - SP, Brazil.

[4] Laboratório Especial de Toxinologia Aplicada – Instituto Butantan, São Paulo – SP, Brazil.

[5] Laboratório de Bacteriologia - Instituto Butantan, São Paulo - SP, Brazil.

*corresponding author: carolina.mello@unifesp.br (ACMS)

Authors summary

The phylogenetic origin of extraintestinal pathogenic Escherichia coli is mostly associated with phylogroup B2, and the majority of the studies regarding extraintestinal infection focus on the most virulent strains, which might also present multidrug-resistant (MDR) phenotype. Strains belonging to phylogroup B1 and isolated from extraintestinal infections are considered as opportunist pathogens and have their virulence neglected. We focus our study in one MDR strain isolated from bloodstream infection that belongs to phylogenetic group B1 to enlarge the knowledge about the virulence of this kind of strain. We demonstrated that the EC121 is capable of adheres to intestinal and bladder human cells, and invades the latter one; it survives to human serum bactericidal effects and produces biofilm. Additionally, the in vivo assay confirmed the EC121 virulence, showing that it should be considered a pathogenic strain. The genetic analyzes highlighted important aspects of EC121 which are typical from strains of sequence type 101, like its involvement in the spread of antimicrobial resistance genes and its relationship with extraintestinal infection from diverse sources. Information concerning the virulence of MDR strains is important for the development of global actions treating the spread of antimicrobial resistance, as well as to elucidate the pathogenesis of strains that were considered as an opportunist.

Abstract

Escherichia coli EC121 is a multidrug-resistant (MDR) strain isolated from bloodstream infection of an inpatient with persistent gastroenteritis and Zone T lymphoma, that died due to septic shock. Despite causing an extraintestinal infection, it harbors few known virulence factors and was assigned into phylogenetic group B1. To evaluate if the EC121 was pathogenic or opportunistic, its genome was sequenced, and

an *in vitro* characterization of some pathogenicity-associated properties was performed. The data retrieved from genome analysis showed that *E. coli* strain EC121 belongs to the O154:H25 serotype, and to ST101-B1, which was epidemiologically linked to extraintestinal infections and antimicrobial resistance spread as well. Moreover, it was closely related to Shiga-toxin producing *E. coli* (STEC). Besides, strain EC121 is an MDR strain harboring 14 antimicrobial resistance genes, including $bla_{CTX-M-2}$, and more than 50 complete virulence genetic clusters, which are reported to be associated either with DEC or ExPEC. The strain also displays the capacity to adhere to a variety of cell lineages, and invade T24 bladder cells, as well as the ability to form biofilms on abiotic surfaces, and survive the bactericidal serum complement activity. Additionally, it is virulent in the *Galleria mellonella* model. Altogether, *E. coli* EC121 unveiled to be a pathogen powered by its multi-drug resistance characteristic. Carry out studies providing accurate information about the virulence potential of all kinds of MDR strains are essential because these studies will help in the development of alternative therapies of infection management and spread control of MDR strains.

Keywords: Extraintestinal, intestinal, *Escherichia coli*, Multidrug resistance, hybrid pathogen, bloodstream infection, DEC, MDR, virulence, ST101.

Introduction

Escherichia coli is one of the most frequent pathogens isolated from bloodstream infections (BSI) around the world [1–5]. Despite all knowledge of extraintestinal infections due to pathogenic E. coli, the number of severe infections and outbreaks caused by these pathogens is rising [2,6]. Moreover, many of these infections are caused by multidrug-resistant (MDR) strains, leading to a higher burden of disease [7–10].

The term extraintestinal pathogenic Escherichia coli (ExPEC) is used to define strains recovered from any extraintestinal infection in humans or animals. Although many virulence factors are associated with the pathogenicity of this group, it is difficult to identify or classify ExPEC strains based on a specific group of virulence genes [11]. Some studies developed molecular virulence patterns that define strains that harbor intrinsic extraintestinal virulence potential [12] or are potentially capable of causing urinary tract infection [13]. These molecular patterns are useful tools to track ExPEC both in the gastrointestinal tract or environment (soil, water, food), enabling the search for ExPEC reservoirs.

Even though such methods may recognize the most virulent strains, they fail in identifying a considerable part of isolates recovered from clinical samples [13,14]. The reason is that infections take place as a result of an imbalance between the virulence potential of the pathogen and the immune defenses of the host what makes it sometimes unclear whether the infection is being caused by a true pathogen or by an opportunistic strain. Considering this, the use of epidemiologic data and multi-locus sequence typing (MLST) for the identification of strains belonging to major pathogenic clonal groups could help in the determination of the potential pathogenic role played by an E. coli strain [15–17].

The emergence of MDR *E. coli* strains calls attention to the spread of clones

carrying virulence along with resistance-encoding genes, making the control of these

pathogens potentially difficult [2,18–22]. In this context, ST131 is the MDR high-risk

clonal group widely disseminated and studied worldwide nowadays. Other clonal

groups, presenting MDR phenotype, like ST405, ST38, and ST648, have also emerged

and are already considered as of global risk [23]. On the other hand, some STs

presenting MDR phenotype, although being isolated around the world, have not their

pathogenic potential determined yet.

The spread of MDR pathogens is a major Public Health concern that needs to be

adequately addressed towards efficient control. Based on that, the World Health

Organization (WHO) called attention to this problem and the need for alternative

therapeutic options for the treatment of MDR infections. The development of vaccines

and anti-virulence compounds could be alternative approaches to combat MDR strains,

especially those showing pan drug resistance (PDR) phenotype [24]. However, for these

alternatives to be effective, advanced knowledge is necessary, since not all pathogenic

strains share the same virulence factors, and the use of the most prevalent virulence

factors as targets can be problematic as they can adversely affect the gut microbiota.

It is well accepted that the phylogenetic grouping of *E. coli* keeps a very good

correlation with the virulence potential of bacterial isolates. More recent

epidemiological data have shown that non-virulent strains are mostly classified in the

phylogenetic groups A, while diarrheagenic *E. coli* (DEC) are B1, and ExPEC are

mainly B2. However, the fact that most *E. coli* virulence factors are carried on mobile

genetic elements (e.g. plasmids and pathogenicity islands) may eventually cross these

phylogenetic boundaries and promote the appearance of potential pathogens in atypical

phylogroups.

Thus, a global analysis of virulence and resistance characteristics of ExPEC

isolates, especially those escaping typical classification, is essential for the

understanding of the infections they cause as well as for devising therapeutic

alternatives to cope with MDR *E. coli* strains.

Aiming to provide information about the virulence determinants in non-typical

ExPEC strains, we performed an extensive genotypic and phenotypic characterization of

the MDR *E. coli* EC121 strain, which was characterized as belonging to phylogenetic

group B1 and carrying only a few known virulence factors.

Material and Methods

Bacterial strain.

The *E. coli* strain EC121 was isolated from the blood of a patient diagnosed with T-zone lymphoma and persistent infectious gastroenteritis, who had been hospitalized in a tertiary hospital located in the city of São Paulo, Brazil, in 2007. The patient died due to septic shock two days after the isolation of the EC121. The EC121 strain was kept frozen in glycerol at -80ºC in ENTEROBACTERIALES-EXTRAINTESTINAL-EPM-DMIP collection nº A27A7C3. The initial virulence and resistance characterization showed that EC121 strain belonged to phylogroup B1, presented an MDR phenotype by routine susceptibility testing, and harbored few known virulence genes (*fim, hra, cvaC, ompA, ompT, sitA, iroN*). Furthermore, it was not considered a pathogenic strain because it harbored none of the virulence factors commonly involved in the characterization of ExPEC (presence of two of the following genes: *papA/C*, *sfaDE*, *afaBC*, *iuc/iut*, and *kpsMT*II) [25].

Total DNA extraction, whole-genome sequencing (WGS), and genome assembly.

The total bacterial DNA extraction was done using Wizard® Genomic DNA Purification Kit (Promega - USA) following the manufacturer's protocol. The extracted DNA was sequenced in an Illumina® Hiseq1500 (Illumina-USA), using the Rapid protocol to obtain 2x250 paired-end reads, according to the manufacturer´s recommendations. Raw data were processed with Trimmomatic, and then the paired-end reads were assembled using SPAdes (version 3.12.0), with default parameters, and careful mode on [26].

Genomic analyses and annotation.

The obtained draft genome was submitted to various online bioinformatics

platforms of the Center of Genomic Epidemiology (CGE) pipeline to determine (i)

sequence types [27] for both *E. coli* MLST schemes; (ii) serotype (SerotypeFinder 2.0)

[28]; (iii) presence and types of plasmid replicons (PlasmidFinder 2.0) [29]; (iv)

presence of resistance genes (ResFinder 3.1) [30]; and (v) STEC virulence factors

(VirulenceFinder 2.0) [31]. PHASTER [32] and PHAST [33] were used to detect

bacteriophage sequences in each contig of the draft genome.

The genome was annotated using Pathosystems Resource Integration Center

(PATRIC) Comprehensive Genome Analysis service that uses RASTtk [34]. Each

sequence that was assigned as a virulence factor in PATRIC's database was manually

submitted to BLAST/NCBI [35] and UniProt [36] to validate the virulence factors, to

get all information about the virulence genes detected, to evaluate the completeness of

the sequence and to determine its homology in relation to the RefSeq protein in Swiss-

Prot. PATRIC [34] service was also used to build a phylogenetic tree using RAxML-

VI-HPC or Fast tree 2, where all representative *E. coli* genomes from different

pathotypes were used to construct the tree, as well as the deposited genomes of *E. coli*

strains belonging to ST101 complex from diverse sources. The tree was built based on

the concatenated sequence of all shared proteins among all genomes using RAxML or

FastTree2. To construct the phylogenetic tree of EC121 and representative *E. coli*

pathotypes, two *Escherichia fergusonni* strains ATCC35469 and NCTC12128 were

used as outgroups**.** To build the phylogenetic trees, a total of 114 public genomes were

randomly selected among the published genomes from the ST101 complex (ST101,

ST359, ST2480, ST5957, and ST6388) using the PATRIC´s Genomes search tool. The

E. fergusonni strains ATCC35469, *E. coli* str IAI1, *E. coli* O157:H7 str Sakai, *E. coli*

O104:H4 str 2011c-3493 were used as outgroup. All phylogenetic tree final layout and

annotation were done using iTOL v.4 [37]. The annotated genome was submitted to

MacSyFinder from Galaxy@Pasteur [38] to detect CAS-CRISPR sequence type and the

presence of secretion systems [39,40].

Serum agglutination assay for typing the O and H antigens.

Serum agglutination assay was carried out following the standard methodology

as described by Orskov and Orskov [41] for serotyping, using O serum against O100

and O154, and H serum against H25 provided by the Centers for Disease Control and

Prevention (CDC, USA).

Plasmid DNA extraction and analysis.

Bacteria were cultivated in Tryptic soy broth (TSB - Difco, USA) at 37ºC, in a

static stove for approximately 18 h, and 1 mL of the culture was submitted to plasmid

alkaline extraction protocol [42]. The *E. coli* strain 39R861 was used as a plasmid mass

reference ladder and as control of extraction [43]. The plasmid extract was submitted to

electrophoresis in an agarose gel (0.8%) in Tris-Borate-EDTA (TBE) buffer, stained

with ethidium bromide solution (5 µg/mL), analyzed using Molecular Imager[®]Gel

Doc™ XR[+] with Image Lab™ Software System from Bio-Rad (USA). The molecular

weight of each plasmid was calculated as previously described [44], based on migration

in agarose gel of five different extraction assays followed by electrophoresis.

In silico **plasmid analysis.**

To accomplish the plasmid analysis, the following strategies were used. First,

the draft genome was submitted to CGE to identify the contigs that contained replicons;

subsequently, the assembled genome was analyzed by PlasFlow [45] to classify the

possible source of each contig (as chromosomal or plasmid). The contigs containing

replicons were analyzed using the Standard Nucleotide BLAST in NCBI.

Data availability.

The EC121 Whole Genome Shotgun project has been deposited at

DDBJ/ENA/GenBank under the accession VYQD00000000. The version described in

this paper is version VYQD01000000.

Antimicrobial susceptibility testing.

The minimum inhibitory concentration (MIC) was determined using the broth

microdilution method, following the European Committee on Antimicrobial

Susceptibility Testing (EUCAST) recommendations and breakpoints [46]. The

following antimicrobials (Sigma - USA) were tested: ampicillin,

piperacillin/tazobactam, ceftriaxone, ceftazidime, cefepime, aztreonam, ertapenem,

imipenem, meropenem, ciprofloxacin, amikacin, gentamicin, tigecycline, colistin,

polymyxin B, trimethoprim/sulfamethoxazole, and chloramphenicol. *E. coli* ATCC

25922 and *Pseudomonas aeruginosa* ATCC 27853 were used as quality control strains.

Conjugation assay

Conjugation assay was conducted to resolve the EC121 conjugative plasmids. In

the mating pair, strain EC121 was the donor strain whereas *E. coli* K-12 derived strains,

J53 [47] and C600 [48], resistant to sodium azide, were the recipient strains.

Conjugation was performed using overnight cultures of the donor and the recipient

strains grown in LB, in the proportion of 1:2, respectively. One milliliter of fresh LB

was added to the mating mixtures, following 3 hours incubation at 37 °C under static

condition. After the incubation period, 100 µl of each mating mixture was plated into

selective MacConkey agar plates supplemented with three different antibiotic

combinations (20 µg/mL gentamycin and 100 µg/mL sodium azide; 2 µg/mL

cefotaxime and 100 µg/mL sodium azide; and 30 µg/mL chloramphenicol and 100

µg/mL sodium azide). The colonies that grew in the selective medium

(Transconjugants) were purified in the same selective medium and then stored for

further characterization.

Characterization of conjugative plasmids

The transconjugants obtained were analyzed by PCR for the presence of

virulence encoding-genes, resistance $bla_{CTX-M-2}$ gene, and determination of the replicon

types. Additionally, the susceptibility profile of the transconjugants was determined by

the minimum inhibitory concentration (MIC) method [14,49,50].

Determination of the lowest bacterial inoculum which was resistant to human

serum complement.

To access the bacterial serum-resistance, the lowest bacterial inoculum resistant

to serum was assessed. Lyophilized human complement serum (Sigma, USA) was

reconstituted in sterile phosphate-buffered saline (PBS). The assay was performed in

96-wells plates, where complement serum was distributed in each well (90 uL per well).

Bacteria were grown overnight at 37°C, serially diluted (1:10) in complement serum

until 10^{-10} and incubated at 37°C. Aliquots of 10 µL of each well were seeded onto

MacConkey agar plates after 30 min, 1 h and 2 h of incubation. Simultaneously, another

assay was performed with previously heat-inactivated serum as control. The *E. coli*

strains J96 and C600 were used as resistant and susceptible controls, respectively [51].

The lowest bacterial inoculum resistant to human complement was determined by the

last bacterial dilution which had bacterial growth onto MacConkey after the challenge.

For each assay, the initial bacterial inoculum was determined by diluting bacteria in

PBS, plating in MacConkey agar and CFU counting. The data was reported in CFU/mL.

Biological assays were performed in triplicates.

Biofilm formation on abiotic surfaces.

Biofilm formation was evaluated on polystyrene and glass surfaces as described

by Lima et al [49] in a 24 hour-assay using the following media: DMEM high-glucose

and TSB. Each assay was performed in biological and experimental triplicates. The

EAEC 042 and laboratory *E. coli* HB101 strains were used as positive and negative

controls, respectively; in all assays, a non-inoculated well was used as control of dye

retention, and the prototype strain CFT073 as an ExPEC control.

Cell culture and maintenance.

HeLa (ATCC® CCL-2™), intestinal Caco-2 (ATCC® HTB-37™) and bladder T24

(ATCC® HTB-4™) cell lineages were used to evaluate the ability of strain EC121 to

interact with eukaryotic cells. HeLa and Caco-2 cells were cultured in DMEM, High

Glucose, GlutaMax™ (Gibco- ThermoFisher Scientific, USA), supplemented with 10%

bovine fetal serum (BFS) (Gibco, USA), 1% non-essential amino acids (Gibco, USA),

and 1x Penicillin-Streptomycin-Neomycin (PSN) antibiotic mixture (Gibco, USA),

while T24 cells (ATCC HTB-4) were cultured in McCoy 5A (modified) media (Gibco,

USA), supplemented with 10% of BFS and 1x PSN antibiotic mixture. All lineages

were kept at 37°C in an atmosphere of 5% CO_2. For all assays, cell suspensions

containing 10^5 cells/mL were seeded in 24-well plates, with or without glass coverslips

for qualitative or quantitative assays and cultured from two up to 10 days until.

Adherence assay in HeLa, Caco-2, and T24 cells.

The adherence properties of EC121 were evaluated qualitatively and

quantitatively. In both assays, the epithelial cells were washed three times with PBS and

1 mL of proper media, supplemented with 2% of BFS, was added. After the bacterial

inoculation, the qualitative and quantitative assays were incubated for 3 hours at 37 °C,

then washed three times with PBS and processed according to the type of the test. The

qualitative assays were performed with all cell lineages (HeLa, T24, and Caco-2) using

20 of inoculum, obtained from standardized bacterial cultures grown overnight, and

after the incubation and washes, were fixed and stained as described previously [52].

The quantitative assay was carried out using HeLa cells in full confluency to evaluate

the efficiency of the EC121 adherence in the presence and absence of 2% of D-

mannose. The multiplicity of infection (MOI) of 50, obtained from an overnight culture,

washed, and adjusted in PBS, confirmed by O.D. and bacterial count, was used as

inoculum. After incubation and washes, the cells were lysed by adding 1 mL of sterile

bi-distilled water, which was recovered, diluted, and plated onto MacConkey agar for

quantification as previously described [53]. The assays were performed in biological

duplicates and experimental triplicates, the data were expressed as SEM. The *E.*

coli strains C1845, CFT073, and C600 were used as controls.

Short period interaction and invasion assay in Caco-2, and T24 cells.

The invasion assays were carried out using Caco-2 and T24 cells in full

confluency, as described by Martinez et al. [54], with modifications, in two sets of

plates simultaneously. The infection was done using a MOI of 50, and the assays were

incubated for 2 h, at 37°C, in a normal atmosphere. After this period, one plate set was

washed three times with PBS and incubated again with PBS containing 100 µg/mL of

amikacin for 1 hour at 37°C, to kill all extracellular bacteria. After the incubation

period, the assay was washed three times to remove all antibiotics, cells were lysed with

water, and the well contents were collected, diluted and plated onto MacConkey agar to

obtain the number of internalized bacteria. The other set was washed with PBS three

times, the cells were lysed, and contents of each well were collected, diluted and plated

to obtain the total number of bacteria interacting with the cells in the period. An aliquot

of the PBS recovered from the last wash after incubation with amikacin was collected

and plated without dilution, to ensure that the treatment had killed all extracellular

bacteria. The total interaction value was the percentage of the total bacterial associated

with the cells in relation to the initial inoculum. The invasiveness was determined by the

ratio between the number of internalized bacteria and the initial inoculum, expressed in

percentage. The *Escherichia albertii* strain 1551::*eae* was used as adherent and non-

invasive control [55], the *E. coli* strain C600 as a negative control, and the CFT073 as

an ExPEC control. The assays were performed in biological and experimental

triplicates, and data were reported as SEM.

***In vivo* assay in *Galleria mellonella* virulence model**

The full virulence potential of EC121 was evaluated using the virulence model

of *G. mellonella* as previously described [56,57]. Briefly, 10 mL of logarithmic phase

cultures were washed twice and adjusted to the optical density of 0.7 (O.D.595) in

0.85% NaCl and serially diluted to obtain a bacterial concentration of $1x10^6$ CFU/mL.

Volumes of 10 µl of the bacterial suspension were injected in the left larvae proleg

using a Hamilton syringe (26S gauge, 50 µL capacity). The tests were performed in

three independent assays, each of them using 5 larvae per bacterial strain tested. The *E.*

coli strains CFT073 and C600 were used as positive and negative controls, respectively,

and saline injection as the procedure control.

Statistical analyses

Student t-test was applied to calculate statistical significance. For biofilm

production, the Wilcoxon matched-pairs test correction was used, and for the cell

interaction and invasion assays, the Wilcoxon-Mann-Whitney test was applied. The

Kaplan-Meier survival curve was done for survival analysis, and the difference between

the groups was determined by Log-rank (Mantel-cox) test and Gehan-Breslow-

Wilcoxon test. The threshold for statistical significance was a *P*-value < 0.05. The

analyses were performed in Prism 5.0 (GraphPad Prism Software, Inc).

Results

Genetic characterization and *in silic*o analysis.

The genome assembly of the strain EC121 generated 143 contigs, with a total

genome predicted size of 5,119,556 bp, and 50.43% of GC content. PlasFlow algorithm

identified 43 of 143 contigs as chromosomal and other 66 contigs as plasmidial. The

contigs assigned as plasmidial comprised at least 426,419 bp of the genome.

EC121 belongs to serotype O154:H25, ST101-B1, and is related to diarrheagenic *E.*

***coli*.** The MLST analyses showed that the EC121 strain belonged to the ST101/ST88,

according to the Warwick and Pasteur MLST schemes, respectively. The determination

of serotype type by sequencing analysis was inconclusive, because two possible

serotypes, O154:H25 and O100:H25, could be assigned. Using specific serum

agglutination assays it was determined that strain EC121 expressed the O154 antigen.

Thus, it was characterized as belonging to serotype O154:H25. Moreover, group IV

capsule encoding genes were identified on its genome.

A phylogenetic tree built using reference *E. coli* strains from all pathotypes

showed that EC121 was related to diarrheagenic *E. coli* (DEC) strains, since it was

positioned in a clade closely associated with Shiga toxin-producing *E. coli* (STEC)

(Figure 1).

A second phylogenetic tree was built with 95 strains available at the NCBI,

which belonged to the ST101 complex and were recovered from distinct sources (Figure

2, Table S1). Analysis of the tree showed that the majority of the strains of this ST are

MDR, many of them carrying *mcr*-1 (mobile colistin resistance gene), a variety of β-

lactamases (bla_{CTX-M}-like, bla_{OXA}-like, bla_{NDM}-like) and genes related to fosfomycin

resistance (*fosA3*). Interestingly, these strains were isolated from food, environment,

animals, and humans, as part of the microbiota or involved in both intestinal and

extraintestinal infections, in all continents. Among the strains isolated from human

infections (Figure S1 and Table S1), most were diagnosed as extraintestinal pathogens

(28 strains), while three were intestinal pathogens; among the latter, one was Shiga

toxin-producing *E. coli* (STEC), and one was enterotoxigenic *E. coli* (ETEC).

Regarding the isolates from food and animals, seven strains were identified as STEC

(Figure 2, Figure S1 and Table S1). Showing that ST101 is associated with intestinal

and extraintestinal infection and it is a clonal group associated with MDR phenotype.

The full resistome and isolation data (accession number, year and country of isolation,

etc.), related to strains from the ST101 complex used to build the phylogenetic trees,

were provided in Table S.

EC121 harbors genes involved in virulence and stress response. The EC121 genome annotation showed that strain EC121 contained 5,175 coding sequences (CDS), 82 tRNA, and 13 rRNA. One CRISPR *locus* was identified as type 1-IE and presented two arrays and 30 CRISPR-repeat regions (Figure 3A). Among the CDSs annotated, 702 corresponded to putative proteins designated as hypothetical proteins, and 4,473 CDS to putative proteins with functional assignments. Of interest, 221 genes were reported as belonging to systems involved in response to stress, virulence, and defense (Table 1 and Figures 3B-C).

Analysis using the MacSyFinder tool identified 10 type V secretion system proteins, nine of which were from type T5aSS and one from T5cSS (EhaG) (Table S2). One type III secretion system similar to *Salmonella* T3SS, and an incomplete type VI secretion system that carried only *tssB*, *tssD*, *tssE*, *tssH*, and *tssI* genes were also present (Table S2).

All genes that were reported by the PATRIC virulence factor database were manually curated to provide information about their full sequence. As shown in Tables 2, S3, and S4, the genome of the EC121 strain encodes multiple adhesins, invasins, iron uptake systems, and genes involved with the evasion of the host immune system. By the position of strain EC121 in the pathotype phylogenetic tree, some of these virulence factors are related to the pathogenesis of DEC, specifically of STEC and ETEC (Tables 2 and S3). Other virulence factors found were related to *Salmonella* spp. (PagN adhesin and systems associated with immune evasion and macrophage survival) and *Shigella* spp. (genes associated with intracellular survival and spread). Moreover, many accessory genetic clusters associated with the bacterial ability to cause extraintestinal infections, i.e., genes involved with biofilm formation, adherence to extraintestinal

cells, iron acquisition and immune evasion were also detected in the EC121 genome.

Furthermore, other clusters associated with Urinary Tract Infections (UTI) were

detected (Tables 2 and S3).

No complete phage sequences were detected in the EC121 strain. The search for

phage sequences in strain EC121 identified 12 regions containing genes from a variety

of different phages, ranging from 6 to 31 kb (Table 3, Table S4). Although the database

considered a predicted phage sequence in region 6 as intact, based on their score

criteria, it´s size (14,400 bp) was not compatible with the size of the predicted

"*Salmonella* phage Fels-2", whose complete genome sequence deposited in NCBI

database is 33,693 bp. Remarkably, parts of the cytolethal distending toxin (Cdt-I and

Cdt-V), and Shiga-like toxin (Stx1a and Stx2c) converting phages, as well as of *Shigella*

serotype-converting phages SfI, SfII, and SfV were detected among these regions

(Table 3 and Table S4). However, no toxin-encoding genes were identified in strain

EC121.

EC121 has multiple resistance genes, and several efflux-pumps compatible with its

antimicrobial susceptibility profile. The software ResFinder identified 15 resistance

genes in strain EC121, which are involved in reduced susceptibility to aminoglycosides,

β-lactams, macrolides, phenicols, sulphonamides, trimethoprim, and tetracyclines

(Table 4). Mutations in *parE*, *parC*, and in *gyrA* genes, which confer resistance to

fluoroquinolones, were also observed in the EC121 genome. Additionally, several

efflux-pumps related to resistance to heavy metals (copper and mercury), arsenic,

disinfectants (QacE), and antimicrobials (AcrAB-TolC, AcrAD-TolC, AcrEF-TolC,

AcrZ, EmrAB-TolC, EmrD, EmrKY-TolC, MacA, MacB, MdfA/Cmr, MdtABC-TolC,

MdtEF-TolC, MdtL, MdtM, and SugE) were also detected. The MDR phenotype profile

of strain EC121 assessed by the microdilution method was consistent with the genomic

findings, as can be seen in Table 4.

The EC121 strain harbors multiple plasmids. Four bands were detected by agarose

gel electrophoresis, suggesting that EC121 harbor multiple plasmids (Figure S2). Seven

different replicons were found in EC121 using PlasmidFinder 2.0 (IncHI2A, IncHI2,

IncQ1, IncFII, IncFIB, IncN, and IncM1).

Although the EC121 genome is still in draft, we analyzed the replicon containing

contigs identified by the PlasmidFinder to provide more information about the strain's

plasmidial content.

One contig of 128,478 bp in length bears both IncHI2A and IncHI2 replicons. The

contig's BLAST showed high identity with the pYps.F1 plasmid of the *Yersinia*

pseudotuberculosis strain Yps.F1 (cover: 99%, e-value: 0.0, identity: 99.77%) (Figure

S3).

The IncM1 replicon was identified in one contig of 65,565 bp in length, which was

circularized in the assembling, and showed high homology (100% of identity and

coverage) with the pASM2 plasmid (accession n° NZ_CP019841.1) of *Enterobacter*

roggenkampii strain R11 (Figure S4).

The other replicons were segregated into four different contigs, with IncFIB and IncFII

into contigs of 37,695 and 30,674 bp in length, respectively, and IncN and IncQ1 into

contigs with less than 3.5 kb each.

The IncFIB replicon's contig also contained the virulence genes *iroN, iss,* and *traT* as

identified in the sequence. The *in silico* analyzes suggested that the plasmid was similar

to the pAPEC plasmid that carries virulence and resistance genes simultaneously.

Although the IncFII replicon was identified into a different contig, the data indicated that both the IncFII and IncFIB replicons represent segments of the same plasmid. Further studies are required to unravel this plasmid genetic composition.

Conjugation assays were performed to evaluate the presence of conjugative plasmid in EC121, to this purpose using agar plates with different antimicrobial combinations (sodium azide [100 µg/mL] with gentamycin [20 µg/mL] or cefotaxime [2 µg/mL] or chloramphenicol [30 µg/mL]) to select the transconjugant strains. While no chloramphenicol-resistant transconjugant colony was detected, five transconjugants were recovered from the gentamycin selective agar plate and 20 from the cefotaxime selective agar plate. Some colonies were purified and then investigated regarding their replicon type and presence of virulence genes that are generally located in plasmids (*hlyF, sitA,* and *iroN*) by PCR. All cefotaxime-resistant transconjugants carried the IncHI2 replicon, $bla_{CTX-M-2}$ and lacked the virulence genes investigated, whilst all five gentamycin-resistant transconjugants carried multiple replicons (IncFIB, IncM1, IncN, and IncHI2), as well as the *hlyF, sitA*, and *iroN* genes (Table 5), and $bla_{CTX-M-2}$ resistance gene. These results suggested that the $bla_{CTX-M-2}$ gene was inserted into IncHI2A/IncHI2 plasmid.

One transconjugant strain of each type was submitted to the microdilution assay to verify the antimicrobial resistance phenotype. The *E. coli* strain ACC09 (*E. coli* J53 harboring the IncHI2/IncHI2A plasmid) obtained from the cefotaxime agar plate presented a resistant phenotype to ceftriaxone and co-trimoxazole and reduced susceptibility to all beta-lactams (Table 5), whereas the *E. coli* strain ACG04 (*E. coli* C600 carrying multiple plasmids) had reduced susceptibility to all the antimicrobial tested except for ciprofloxacin and tigecycline, and presented increased MIC values to all beta-lactams when compared to ACC09 (Table 5).

Although we have isolated only the IncHI2/IncHI2A plasmid, multiple plasmids were

transferred efficiently to the receptor strain in a short period of conjugation (3 hours)

indicating that they were all conjugative or mobilizable plasmids.

Virulence phenotype.

To evaluate the expression of the virulence-encoding genes detected, *in vitro*

assays were performed to analyze the ability of the strain to: (i) resist to the bactericidal

activity of the serum complement system; (ii) attach to abiotic surfaces and form

biofilms; (iii) adhere to and invade eukaryotic cells. Besides, *in vivo* assays were

performed using the *G. mellonella* infection model to evaluate the EC121 virulence

potential.

The EC121 strain was resistant to the bactericidal activity of the complement

system and adhere to abiotic surfaces. To disseminate in the host, extraintestinal

pathogenic bacteria must be able to survive the serum bactericidal activity. To identify

such a feature in EC121, we determined the lowest serum-resistant bacterial inoculum

using a pool of normal human sera (NHS). The lowest inoculum of EC121 strain that

resisted serum activity after two hours was 10^2 CFU/mL, which was similar to the one

obtained for the resistant control strain J96. *E. coli* strain C600, used as the susceptible

control, barely resisted to 30 min-exposure period in the highest inoculum tested (10^8

CFU/mL). To validate if the bacterial survival was associated with the resistance to

complement activity, assays were repeated with heat-inactivated serum. In this

condition, all strains survived the challenge with similar inoculum (Table 6).

The ability to adhere to the abiotic surface and form biofilm can confer many

advantages to any pathogen, including persistence in particular niches and tolerance

against antimicrobials and the host immune system. The EC121 strain was able to

adhere to borosilicate coverslips and polystyrene when grown in DMEM and TSB, as

shown in Figure 4. Although its adherence to the abiotic surface was not massive as the

adherence presented by the positive control strain (EAEC 042), it was significantly

more intense than that of the negative control strain (HB101) (Figures 4). Furthermore,

EC121 produced similar ($P > 0.05$) or higher ($P < 0.01$) biofilm masses as the ExPEC

prototype strain CFT073 in DMEM and TSB, respectively (Figure 4b).

The EC121 strain adheres to and invades eukaryotic cell lineages. At first, we

accessed, qualitatively, the ability of strain EC121 to adhere to HeLa, T24, and Caco-2

cells, using 3-h adherence assay without D-mannose. This assay showed that EC121

interact efficiently with all cells tested (Figure S5). Subsequently, a quantitative 3-h

adherence assay in HeLa cells using initial inoculum of 1×10^7 was performed in the

absence of D-mannose, except for EC121, which also was tested in the presence of 2%

of D-mannose. EC121 was able to adhere to, both in the presence or absence of D-

mannose, which abolishes the adherence mediated by type-1 fimbriae. However, the

presence of D-mannose reduced significantly the adherence ability of strain EC121,

lowing the adherence in about 90% (Figure 5), from 1.3×10^7 CFU/mL to 1.58×10^6

CFU/mL (Figure 5). Noteworthy, EC121 adherence levels in the absence of D-mannose

was higher than all controls tested in the same conditions ($P < 0.01$) (Figure 5).

We also investigated whether the EC121 strain was able to invade T24 and

Caco-2 cells. In a short-period invasion assay, the EC121 strain interacted with T24

cells as efficiently as CFT073 but invaded the cell lineage significantly more (P <

0.005) (Figure 6). On the other hand, its interaction with differentiated and polarized

Caco-2 cells was lower when compared with CFT073, although there was no difference

related to their capacity to invade this type of cell lineage on the conditions tested.

The capacity to interact and invade eukaryotic cells in the presence of D-mannose was

also assessed to evaluate whether the interaction and invasion abilities of EC121 were

dependent on type-1 fimbriae or another mannose-dependent adhesin. The EC121

interaction with both lineages reduced significantly in the presence of D-mannose, like

in HeLa cells showing that mannose-sensible adhesins contributed to its capacity to

interact with the cell lineages tested. Additionally, EC121 remained invasive in the

presence of D-mannose but with reduced bacterial counts in T24 cells ($P< 0.003$)

whereas there was no difference in the invasiveness in Caco-2 (Figure 6), showing that

mannose-sensible adhesins contributed to both adherence and invasion of EC121,

although it was not the only factor associated to these traits.

EC121 is virulent in the *Galleria mellonella* virulence model. The utilization of the

G. mellonella virulence model showed that EC121 promotes higher mortality rates than

the non-pathogenic *E. coli* strain C600 ($P< 0.005$) and the mock-injection ($P< 0.0001$).

Although the *E. coli* strain CFT073 killed more larvae than the EC121, there was no

significant difference among their survival curves (Figure 7), showing that EC121 is

virulent in the model used.

Discussion

Authors should discuss the results and how they can be interpreted in

perspective of previous studies and of the working hypotheses. The findings and their

implications should be discussed in the broadest context possible. Future research

directions may also be highlighted.

The *E. coli* EC121 strain was isolated in 2007 from a bloodstream infection of

an inpatient that presented persistent gastroenteritis and T-zone lymphoma. Since initial

analyses showed that it belonged to phylogenetic group B1 and carried few virulence

markers commonly related to extraintestinal pathogenic *E. coli,* it was classified as

ExPEC negative (ExPEC-) [25] and therefore considered as an opportunistic pathogen.

However, considering that about 40% of extraintestinal infections are caused by strains

devoid of virulence factors [13,14] and that EC121 was an MDR strain, its entire

genome was sequenced to further understand its virulence potential.

Interestingly, the EC121 strain belongs to ST101, which has been previously

reported to be involved in nosocomial outbreaks caused by Metallo-β-lactamases-

producing strains in many countries from Europe, Asia, and Oceania [58–62].

Furthermore, ST101 has also been detected among strains of non-outbreak related

extraintestinal infections [63–70], water [71], poultry infection [72], retail food [63,73–

75], and healthy human and animal intestinal microbiota [70,76–79], mostly presenting

an MDR phenotype. Shrestha et al. [80] drew attention to ST101 due to the PDR

phenotype presented by some strains of this ST, and mainly because it is not considered

a pandemic clone, although it has been isolated worldwide. We, therefore, analyzed data

about the infection type, isolation source, and resistance genetic markers presented by

the strains of the ST101 complex that were previously deposited in the NCBI (Table

S1). Such analysis evidenced that MDR strains of this complex were spread worldwide.

In addition, such ST complex is related to Shiga toxin-producing strains as well as

strains isolated from extraintestinal infections, human and animal microbiota, retail

food, and environment. Moreover, many strains simultaneously carry the $bla_{\text{CTX-M-55}}$,

mcr-1, *fosA3*, and *qnrS1* genes. Interestingly, one strain present simultaneously *aac(6')-*

Ib-cr, $bla_{\text{CTX-M-55}}$, $bla_{\text{NDM-5}}$, $bla_{\text{OXA-1}}$, *mcr-1*, *fosA3*, and other 19 resistance genes.

Likewise, the EC121 strain showed multiple antimicrobial resistance genes,

including genes that confer resistance to third-generation cephalosporins ($bla_{CTX-M-2}$). It

is worth mentioning that the EC121 strain was isolated in 2007, and its MDR phenotype

was relevant since at that period it was susceptible only to carbapenems, polymyxins,

and amikacin. Recently, the *E. coli* strain ICBEC72H, which belongs to ST101 and

carried only $bla_{CTX-M-8}$ and *mrc-1*[64] was isolated from a human extraintestinal

infection in Brazil. Similarly, the ST101 *E. coli* strain 200H (Table S1) was isolated

from a human urinary tract infection and carried bla_{OXA-9}, *mcr-1*, and *aac(6')-Ib-cr*.

These reports show that MDR strains belonging to the ST101 complex have been

circulating in Brazil for a long time.

Some authors showed that the *E. coli* strain 912 (ST101) was selected by the

usage of antimicrobial agents in animals and that it was able to colonize human and pig

gut and spread through the environment, reaching and colonizing animals that were not

under antimicrobial treatment [81,82]. These same authors have also shown that ST101

strains can naturally acquire and transfer plasmid-borne antimicrobial resistance genes

in the gut [81,82]. Such a feature is important for various reasons. First, strain EC121

carried multiple plasmids, which harbored different combinations of antimicrobial-

resistance genes, all of which were successfully transferred to *E. coli* K-12 derived

strains in a 3-hours conjugation assay. Besides, *E. coli* strains belonging to ST101 were

recovered from retail meat in Europe and Asia [74,75], and from extraintestinal

infections in Brazil and the USA in the same regions in which they were detected from

retail meat [63,72].

Additionally, strains belonging to the ST101 complex carrying multiple

resistance genes were recovered from the intestine of healthy humans and animals in

many countries. Therefore, even if these strains do not cause infection directly, they

could potentially transfer plasmids to other bacteria, even from distinct genera. Such

cross genera plasmid transfer could be easily identified in the plasmids reported in the

present study; IncM1 plasmid, for example, is closely related to plasmids found in

Klebsiella spp. and *Enterobacter* spp., while IncHI2/HI2A is related to *Salmonella* spp.

and *Yersinia* spp. plasmids. Together, these findings reinforce the high risks associated

with strains belonging to the ST101 complex due to their ability to colonize humans and

animals' gut, to easily disseminate via retail food and water, being able to acquire and

spread antimicrobial resistance-encoding genes.

Strains from the ST101 complex are included in the phylogenetic group B1,

which implies that they do not have all the classical virulence factors that are usually

associated with the most virulent ExPEC strains [83,84]. Many studies reported

phylogroup B1 *E. coli* strains as commensals or as intestinal pathogens, but not as

extraintestinal pathogens [83–85].

The genomic analysis of the EC121 strain showed a high number of virulence

genes, demonstrating that it presents all the traits necessary to be considered as an

extraintestinal pathogenic agent, like adhesins, iron acquisition systems, and genes

related to immune system evasion. Moreover, like other ExPEC strains, EC121

displayed multiple virulence genes related to each feature, reflecting the redundant

phenotype that ensures its pathogenicity. However, even considering the completeness

of each sequence and each operon, which was manually checked, the presence of

virulence genes "per se" does not guarantee that all of them are expressed. Therefore, to

evaluate the expression of such traits, distinct phenotypic assays were performed and

confirmed the virulence genetic background of EC121.

To test the bacterial ability to resist the serum complement activity that could be

conferred by the presence of *traT, iss,* and *ompT*, a two hours challenge assay was

performed, in which one particle can traverse all circulatory system at least twice, so a

pathogen that resists complement's activity during this period, even with a small

bacterial load, is, in theory, more capable of reaching different niches and spread

through the bloodstream or cause a bloodstream infection. The EC121 strain resisted the

NHS for two hours with an inoculum similar to the resistant *E. coli* J96 control strain,

thus confirming the EC121 serum resistant phenotype. Serum complement is the first

immunological barrier to control pathogens that reach the bloodstream. Complement

resistance confers the possibility to spread to different body sites through the

bloodstream. Hallström et al. [86] reported the relationship of bacterial resistance to

NHS with sepsis severity, and other authors have associated it with different kinds of

extraintestinal infections [87–89].

The ability to colonize and attach to surfaces is also an important trait for any

pathogenic bacteria; in this way, the assays carried out showed not only that EC121

strain was able to adhere to and invade different cell lineages, including bladder T24

cells but that it could also attach and produce biofilm on abiotic surfaces. Peirano et al.

[90] showed that ExPEC negative ST101 MDR strains isolated from extraintestinal

infections could interact with HEp-2 and Caco-2 cells more efficiently than strains

belonging to the epidemic clones ST131 and ST405, which are ExPEC positive [90].

The strain EC121 interacted with all cell lineages tested, but its interaction was

significantly higher in T24 cells. Besides, its interaction's capacity was similar to the

strain CFT073 (ExPEC prototype), and it invaded T24 cells more efficiently, suggesting

that EC121 might be capable to produce intracellular bacterial communities (IBC).

IBCs were related to bacterial persistence and recurrent infections in the host [91].

Overall, the EC121 invasiveness might be more related to its persistence ability than to

its capacity to transpose epithelial barriers, since persistence may help bacteria to evade

the immune system and confer protection against antimicrobials agents. However,

further studies are required to prove this hypothesis.

Ten virulence encoding-genes involved with biofilm production were identified

in the EC121 genome (Table S3). The capacity to produce biofilm might confer many

advantages to bacteria, like protection against the immunological system and

antibiotics, assisting its persistence and spread in the environment. However, biofilm

production depends on many factors, like temperature and presence of specific nutrients

in the media or environment [92–94]. Moreover, the ability to produce biofilm has been

reported to vary among ExPEC strains and that strains with this capacity are considered

to be more pathogenic [95].

Interestingly, many of the EC121 virulence factors detected in the draft genome

are related to diarrheagenic *E. coli*, even though none of them is implicated in DEC

pathotype definition. The presence of many genetic features related to Shiga-toxin

producing *E. coli* (STEC) strains, *e.g*., Hcp, EhaG, and Lpf-1$_{O26}$, as well as the

proximity of EC121 to the clade that contains STEC strains and *E. coli* O104:H4 strain

2011C-3493, draws attention to its potential diarrheagenic background.

Many features identified in the EC121 genome reinforce its linkage with STEC

strains. Phenotypically, EC121 expressed the O154:H25 serotype, but it possesses the

group IV capsule-encoding genes. This kind of capsular group is known to be

thermoresistant and expressed as K_{LPS} or O-antigen capsule. This could explain the

expression of the O154 instead of O100 antigen, despite the presence of all genes

related to the expression of the latter. Interestingly O100 is a STEC related serogroup.

Moreover, much of the phage remains detected in the EC121 strain were related

to Stx-converting phages; besides, ST101 strains carrying the *stx$_{1a}$* gene have been

reported in food sources [73,96]. In humans, ST101 strains were already reported in a

patient with Hemolytic Uremic Syndrome (HUSEC) [96–98] and in non-bloody diarrhea related to a Stx1a-producing *E. coli* strain [99]. Although only one of these strains had its genome sequenced, some ST101 *E. coli* strains recovered from animals and food were found to carry *stx1*. Interestingly, most Stx-converting phages remains found in EC121 were similar to those commonly related to Stx1a production, corroborating with the results presented here. The genome of three non-STEC strains from diarrheic patients was found in GenBank, one of which was devoid of DEC virulence factors. Likewise, EC121 was isolated from bloodstream infection of one inpatient with persistent infectious gastroenteritis which was probably the source of EC121 infection. Unfortunately, the *E. coli* isolated from stool was not stored not allowing further comparison between bloodstream and stool *E. coli* isolates.

Several studies evaluated *E. coli* pathogenicity in the surrogate *G. mellonella* model [56,100–104], these studies point to the model efficiency in differentiating pathogenic strains from non-pathogenic, especially using bacterium inoculum of 10^5 CFU per larvae or lower [103,104]. Moreover, Jonsson et al. [100] have shown that pathogenic *E. coli* needs an inoculum of at least 10^3 CFU to lead larvae to death. To evaluated EC121 virulence, a bacterial inoculum of 10^4 CFU per larvae was used, and in the assayed conditions EC121 was as virulent as the ExPEC prototype CFT073, thus corroborating with the *in silico* and *in vitro* data, showing that some strains from the ST101 are truly pathogenic.

In summary, our extensive *in silico, in vitro,* and *in vivo* analyses of virulence and resistance properties of *E. coli* strain EC121, an O154:H25 B1-ST101 strain isolated from a human bloodstream infection, confirmed its virulence potential and increased the knowledge on the complex scenario of virulence traits present in the MDR

E. coli ExPEC negative group, contributing to the potential development of strategies to

control the spread of such pathogens.

Acknowledgments

We are grateful to the Laboratório Especial de Microbiologia Clínica (LEMC) of the

Federal University of São Paulo (UNIFESP), São Paulo - SP, Brazil, for providing the

E. coli strain EC121.

Financial support

This study was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo

(FAPESP) that provided research grants to E.C. (Process Number: 16/01656-5), R.M.S.

(Process Number: 2009/00402-6), T.A.T.G. (Process Number: 2018/17353-7), and

master scholarship to T.B.V. (Process Number: 2017/21947-7). We are also grateful to

the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for

providing grants under finance code 001 to A.C.M.S. (PNPD), A.P.S. (PhD), C.S.N.

(PhD), and to the National Council for Science and Technological Development

(CNPq) for providing a grant to A.C.G. (Process number: 305535/2014-5).

Ethics

The strain EC121 used in this research was obtained from clinical routine after

laboratory procedures. No additional procedure was performed to acquire any bacterial

strain, so the consent form was not required as determinate by the Brazilian National

Health Council nº 466/12 and 510/16. All patient information was obtained from

medical records, and the research was done with the approval of the local Research

Ethics Committee of the Federal University of São Paulo - UNIFESP/São Paulo

Hospital (CEP 2031/08 and CEP N 7140160317).

Competing interests

A.C.G. has recently received research funding and/or consultation fees from Eurofarma, MSD, Pfizer, and Zambon. Other authors have nothing to declare. This study was not financially supported by any Diagnostic/Pharmaceutical company.

Supporting information captions

Table S1 – ST101 complex strains, NCBI accession, multidrug resistance genotype, source, and country of isolation.

Table S2 – Type V Secretion system protein identified by MaSyFinder.

Table S3 – Complete Genetic cluster identified in EC121 strain and their predicted association with virulence.

Table S4 -Virulence genes identification and validation in NCBI and Swiss-Prot.

Table S5 - Detailed information of all predicted phages detected in strain EC121.

Figure S1 – Phylogenetic relationship among 62 human isolates from the ST101 complex

Figure S2 - Plasmid content of the EC121strain.

Figure S3 – Phylogenetic relationship obtained from sequence alignment of pEC121_IncHI2/HI2A, using BLASTn

Figure S4 – Phylogenetic relationship obtained from sequence alignment of pEC121_IncM1 plasmid, using BLASTn

Figure S5 – EC121 adherence in eukaryotic cell linages HeLa, Caco-2, and T24.

References

1. Biedenbach DJ, Moet GJ, Jones RN. Occurrence and antimicrobial resistance pattern comparisons among bloodstream infection isolates from the SENTRY Antimicrobial Surveillance Program (1997–2002). Diagn Microbiol Infect Dis. 2004;50: 59–69. doi:10.1016/j.diagmicrobio.2004.05.003

2. Temkin E, Fallach N, Almagor J, Gladstone BP, Tacconelli E, Carmeli Y, et al. Estimating the number of infections caused by antibiotic-resistant *Escherichia coli* and *Klebsiella pneumoniae* in 2014: a modelling study. Lancet Glob Heal. 2018;6: e969–e979. doi:10.1016/S2214-109X(18)30278-X

3. Tu B, Liu SX, Wu D, Zhang X, Shi L, Xie YX, et al. [Clinical features of community-acquired bloodstream infection due to Gram-negative bacilli in patients with liver cirrhosis]. Zhonghua Gan Zang Bing Za Zhi. 2018;26: 23–27. doi:10.3760/cma.j.issn.1007-3418.2018.01.007

4. Satwani P, Freedman JL, Chaudhury S, Jin Z, Levinson A, Foca MD, et al. A Multicenter Study of Bacterial Blood Stream Infections in Pediatric Allogeneic Hematopoietic Cell Transplantation Recipients: The Role of Acute Gastrointestinal Graft-versus-Host Disease. Biol Blood Marrow Transplant. 2017;23: 642–647. doi:10.1016/j.bbmt.2017.01.073

5. Nivesvivat T, Piyaraj P, Thunyaharn S, Watanaveeradej V, Suwanpakdee D. Clinical epidemiology, risk factors and treatment outcomes of extended-spectrum beta-lactamase producing *Enterobacteriaceae* bacteremia among children in a Tertiary Care Hospital, Bangkok, Thailand. BMC Res Notes. 2018;11: 624. doi:10.1186/s13104-018-3729-3

6. GLASS | Global antimicrobial resistance surveillance system (GLASS) report. WHO. 2018 [cited 17 Jun 2019]. Available:

https://www.who.int/glass/resources/publications/early-implementation-

report/en/

7.   Landman WJM, van Eck JHH. The incidence and economic impact of the

Escherichia coli peritonitis syndrome in Dutch poultry farming. Avian Pathol.

2015;44: 370–378. doi:10.1080/03079457.2015.1060584

8.   Kaye KS, Marchaim D, Chen T-Y, Baures T, Anderson DJ, Choi Y, et al. Effect

of Nosocomial Bloodstream Infections on Mortality, Length of Stay, and

Hospital Costs in Older Adults. J Am Geriatr Soc. 2014;62: 306–311.

doi:10.1111/jgs.12634

9.   Barriere SL. Clinical, economic and societal impact of antibiotic resistance.

Expert Opin Pharmacother. 2015;16: 151–153.

doi:10.1517/14656566.2015.983077

10.   Ciorba V, Odone A, Veronesi L, Pasquarella C, Signorelli C. Antibiotic

resistance as a major public health concern: epidemiology and economic impact.

Ann Ig. 27: 562–79. doi:10.7416/ai.2015.2048

11.   Johnson JR, Russo TA. Extraintestinal pathogenic *Escherichia coli* : "The other

bad *E. coli* ." J Lab Clin Med. 2002;139: 155–162. doi:10.1067/mlc.2002.121550

12.   Johnson JR, Murray AC, Gajewski A, Sullivan M, Snippes P, Kuskowski MA, et

al. Isolation and molecular characterization of nalidixic acid-resistant

extraintestinal pathogenic *Escherichia coli* from retail chicken products.

Antimicrob Agents Chemother. 2003;47: 2161–8. doi:10.1128/aac.47.7.2161-

2168.2003

13.   Spurbeck RR, Dinh PC, Walk ST, Stapleton AE, Hooton TM, Nolan LK, et al.

Escherichia coli Isolates That Carry *vat* , *fyuA* , *chuA* , and *yfcV* Efficiently

Colonize the Urinary Tract. Infect Immun. 2012;80: 4115–4122.

doi:10.1128/IAI.00752-12

14.   Santos ACM, Zidko ACM, Pignatari AC, Silva RM. Assessing the diversity of

the virulence potential of *Escherichia coli* isolated from bacteremia in São Paulo,

Brazil. Brazilian J Med Biol Res. 2013;46: 968–973. doi:10.1590/1414-

431X20133184

15.   Zhao L, Zhang J, Zheng B, Wei Z, Shen P, Li S, et al. Molecular epidemiology

and genetic diversity of fluoroquinolone-resistant *Escherichia coli* isolates from

patients with community-onset infections in 30 Chinese county hospitals. J Clin

Microbiol. 2015;53: 766–770. doi:10.1128/JCM.02594-14

16.   Izdebski R, Baraniak A, Herda M, Fiett J, Bonten MJM, Carmeli Y, et al. MLST

reveals potentially high-risk international clones of *Enterobacter cloacae*. J

Antimicrob Chemother. 2015;70: 48–56. doi:10.1093/jac/dku359

17.   Brauner J, Hallin M, Deplano A, Mendonça R, Nonhoff C, Ryck R, et al.

Community-acquired methicillin-resistant *Staphylococcus aureus* clones

circulating in Belgium from 2005 to 2009: changing epidemiology. Eur J Clin

Microbiol Infect Dis. 2013;32: 613–620. doi:10.1007/s10096-012-1784-6

18.   Riley LW. Pandemic lineages of extraintestinal pathogenic *Escherichia coli*. Clin

Microbiol Infect. 2014;20: 380–90. doi:10.1111/1469-0691.12646

19.   Roer L, Overballe-Petersen S, Hansen F, Schønning K, Wang M, Røder BL, et

al. *Escherichia coli* Sequence Type 410 Is Causing New International High-Risk

Clones. mSphere. 2018;3: e00337-18. doi:10.1128/mSphere.00337-18

20.   Reineke K, Sevenich R, Hertwig C, Janßen T, Fröhling A, Knorr D, et al.

Comparative study on the high pressure inactivation behavior of the Shiga toxin-

producing *Escherichia coli* O104:H4 and O157:H7 outbreak strains and a non-

pathogenic surrogate. Food Microbiol. 2015;46: 184–194.

doi:10.1016/J.FM.2014.07.017

21. Kennedy CA, Walsh C, Karczmarczyk M, O'Brien S, Akasheh N, Quirke M, et

al. Multi-drug resistant *Escherichia coli* in diarrhoeagenic foals: Pulsotyping,

phylotyping, serotyping, antibiotic resistance and virulence profiling. Vet

Microbiol. 2018;223: 144–152. doi:10.1016/J.VETMIC.2018.08.009

22. WHO | Global action plan on antimicrobial resistance. WHO. 2017 [cited 17 Jun

2019]. Available: https://www.who.int/antimicrobial-

resistance/publications/global-action-plan/en/

23. Shaik S, Ranjan A, Tiwari SK, Hussain A, Nandanwar N, Kumar N, et al.

Comparative Genomic Analysis of Globally Dominant ST131 Clone with Other

Epidemiologically Successful Extraintestinal Pathogenic Escherichia coli

(ExPEC) Lineages. MBio. 2017;8: e01596-17. doi:10.1128/mBio.01596-17

24. Magiorakos A-P, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et

al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria:

an international expert proposal for interim standard definitions for acquired

resistance. Clin Microbiol Infect. 2012;18: 268–281. doi:10.1111/J.1469-

0691.2011.03570.X

25. Santos AC de M. Potencial de virulência e suscetibilidade a antimicrobianos de

amostras de *Escherichia coli* isoladas de bacteremia: sua relação com o estado de

imunocompetência do paciente e a origem da infecção. Universidade Federal de

São Paulo (UNIFESP). 2013. Available:

http://repositorio.unifesp.br/handle/11600/22975

26. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al.

SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-

Cell Sequencing. J Comput Biol. 2012;19: 455–477. doi:10.1089/cmb.2012.0021

27.   Larsen M V., Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, et al.

849   Multilocus Sequence Typing of Total-Genome-Sequenced Bacteria. J Clin

850   Microbiol. 2012;50: 1355–1361. doi:10.1128/JCM.06094-11

851   28.   Joensen KG, Tetzschner AMM, Iguchi A, Aarestrup FM, Scheutz F. Rapid and

852   easy in silico serotyping of *Escherichia coli* isolates by use of whole-genome

853   sequencing data. J Clin Microbiol. 2015;53: 2410–2426.

854   doi:10.1128/JCM.00008-15

855   29.   Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L,

856   et al. In silico detection and typing of plasmids using PlasmidFinder and plasmid

857   multilocus sequence typing. Antimicrob Agents Chemother. 2014;58: 3895–903.

858   doi:10.1128/AAC.02412-14

859   30.   Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al.

860   Identification of acquired antimicrobial resistance genes. J Antimicrob

861   Chemother. 2012;67: 2640–2644. doi:10.1093/jac/dks261

862   31.   Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, et al. Real-

863   time whole-genome sequencing for routine typing, surveillance, and outbreak

864   detection of verotoxigenic *Escherichia coli*. J Clin Microbiol. 2014;52: 1501–

865   1510. doi:10.1128/JCM.03617-13

866   32.   Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a

867   better, faster version of the PHAST phage search tool. Nucleic Acids Res.

868   2016;44: W16–W21. doi:10.1093/nar/gkw387

869   33.   Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. PHAST: A Fast Phage

870   Search Tool. Nucleic Acids Res. 2011;39: W347–W352. doi:10.1093/nar/gkr485

871   34.   Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, et al. Improvements

872   to PATRIC, the all-bacterial Bioinformatics Database and Analysis Resource

Center. Nucleic Acids Res. 2017;45: D535–D542. doi:10.1093/nar/gkw1017

35.   Altschul S, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped

BLAST and PSI-BLAST: a new generation of protein database search programs.

Nucleic Acids Res. 1997;25: 3389–3402. doi:10.1093/nar/25.17.3389

36.   Bateman A. UniProt: A worldwide hub of protein knowledge. Nucleic Acids Res.

2019;47: D506–D515. doi:10.1093/nar/gky1049

37.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. Nucleic Acids Res. 2019;47: W256–W259.

doi:10.1093/nar/gkz239

38.   Mareuil F, Doppelt-Azeroual O, Ménager H, Mareuil F, Doppelt-Azeroual O,

Ménager H. A public Galaxy platform at Pasteur used as an execution engine for

web services. F1000Research. 2017;6.

doi:10.7490/F1000RESEARCH.1114334.1

39.   Abby SS, Néron B, Ménager H, Touchon M, Rocha EPC. MacSyFinder: A

Program to Mine Genomes for Molecular Systems with an Application to

CRISPR-Cas Systems. Torres N V., editor. PLoS One. 2014;9: e110726.

doi:10.1371/journal.pone.0110726

40.   Abby SS, Cury J, Guglielmini J, Néron B, Touchon M, Rocha EPC.

Identification of protein secretion systems in bacterial genomes. Sci Rep. 2016;6:

23080. doi:10.1038/srep23080

41.   Orskov F, Orskov I. *Escherichia coli* O:H serotypes isolated from human blood.

Prevalence of the K1 antigen with technical details of O and H antigenic

determination. Acta Pathol Microbiol Scand Suppl. 1975;83: 595–600. Available:

http://www.ncbi.nlm.nih.gov/pubmed/1106116

42.   Birnboim HC, Doly J. A rapid alkaline extraction procedure for screening

recombinant plasmid DNA. Nucleic Acids Res. 1979;7: 1513–1523.

doi:10.1093/nar/7.6.1513

43.   Moran RA, Anantham S, Hall RM. An improved plasmid size standard,

39R861+. Plasmid. 2019;102: 6–9. doi:10.1016/j.plasmid.2019.01.002

44.   Meyers JA, Sanchez D, Elwell LP, Falkow S. Simple Agarose Gel

Electrophoretic Method for the Identification and Characterization of Plasmid

Deoxyribonucleic Acid. J Bacteriol. 1976. Available: http://jb.asm.org/

45.   Krawczyk PS, Lipinski L, Dziembowski A. PlasFlow: predicting plasmid

sequences in metagenomic data using genome signatures. Nucleic Acids Res.

2018;46: e35–e35. doi:10.1093/nar/gkx1321

46.   EUCAST. The European Committee on Antimicrobial Susceptibility Testing.

Breakpoint tables for interpretation of MICs and zone diameters. Version 9.0,

2019. http://www.eucast.org. Available: http://www.eucast.org.

47.   Matsumura Y, Peirano G, Pitout JDD. Complete genome sequence of Escherichia

coli J53, an azide-resistant laboratory strain used for conjugation experiments.

Genome Announc. 2018;6. doi:10.1128/genomeA.00433-18

48.   Allué-Guardia A, Nyong EC, Koenig SSK, Vargas SM, Bono JL, Eppinger M.

Closed Genome Sequence of Escherichia coli K-12 Group Strain C600 .

Microbiol Resour Announc. 2019;8. doi:10.1128/mra.01052-18

49.   Lima MP, Yamamoto D, Santos AC de M, Ooka T, Hernandes RT, Vieira MAM,

et al. Phenotypic characterization and virulence-related properties of *Escherichia*

albertii strains isolated from children with diarrhea in Brazil. Pathog Dis.

2019;77. doi:10.1093/femspd/ftz014

50.   Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification

of plasmids by PCR-based replicon typing. J Microbiol Methods. 2005;63: 219–

228. doi:10.1016/J.MIMET.2005.03.018

51.　Houdouin V, Bonacorsi S, Brahimi N, Clermont O, Nassif X, Bingen E. A

uropathogenicity island contributes to the pathogenicity of Escherichia coli

strains that cause neonatal meningitis. Infect Immun. 2002;70: 5865–9.

doi:10.1128/iai.70.10.5865-5869.2002

52.　Garcia BG, Ooka T, Gotoh Y, Vieira MAM, Yamamoto D, Ogura Y, et al.

Genetic relatedness and virulence properties of enteropathogenic *Escherichia coli*

strains of serotype O119:H6 expressing localized adherence or localized and

aggregative adherence-like patterns on HeLa cells. Int J Med Microbiol.

2016;306: 152–164. doi:10.1016/j.ijmm.2016.02.008

53.　Santos FF, Yamamoto D, Abe CM, Bryant JA, Hernandes RT, Kitamura FC, et

al. The Type III Secretion System (T3SS)-Translocon of Atypical

Enteropathogenic *Escherichia coli* (aEPEC) Can Mediate Adherence. Front

Microbiol. 2019;10. doi:10.3389/fmicb.2019.01527

54.　Martinez JJ, Mulvey MA, Schilling JD, Pinkner JS, Hultgren SJ. Type 1

pilus☐mediated bacterial invasion of bladder epithelial cells. EMBO J. 2000;19:

2803–2812. doi:10.1093/EMBOJ/19.12.2803

55.　Hernandes RT, Silva RM, Carneiro SM, Salvador FA, Fernandes MCDC,

Padovan ACB, et al. The localized adherence pattern of an atypical

enteropathogenic *Escherichia coli* is mediated by intimin omicron and

unexpectedly promotes HeLa cell invasion. Cell Microbiol. 2008;0:

071003010119002-??? doi:10.1111/j.1462-5822.2007.01054.x

56.　Alghoribi MF, Gibreel TM, Dodgson AR, Beatson SA, Upton M. Galleria

mellonella Infection Model Demonstrates High Lethality of ST69 and ST127

Uropathogenic E. coli. Chakravortty D, editor. PLoS One. 2014;9: e101547.

doi:10.1371/journal.pone.0101547

57.   Andrey DO, Pereira Dantas P, Martins WBS, Marques De Carvalho F, Gonzaga

L, Almeida P, et al. An Emerging Clone, Klebsiella pneumoniae Carbapenemase

2–Producing K. pneumoniae Sequence Type 16, Associated With High Mortality

Rates in a CC258-Endemic Setting. Clin Infect Dis. 2019 [cited 16 Mar 2020].

doi:10.1093/cid/ciz1095

58.   Ranjan A, Shaik S, Mondal A, Nandanwar N, Hussain A, Semmler T, et al.

Molecular Epidemiology and Genome Dynamics of New Delhi Metallo-β-

Lactamase-Producing Extraintestinal Pathogenic *Escherichia coli* Strains from

India. Antimicrob Agents Chemother. 2016;60: 6795–6805.

doi:10.1128/AAC.01345-16

59.   Pfeifer Y, Trifonova A, Pietsch M, Brunner M, Todorova I, Gergova I, et al.

Clonal Transmission of Gram-Negative Bacteria with Carbapenemases NDM-1,

VIM-1, and OXA-23/72 in a Bulgarian Hospital. Microb Drug Resist. 2017;23:

301–307. doi:10.1089/mdr.2016.0059

60.   Poirel L, Lagrutta E, Taylor P, Pham J, Nordmann P. Emergence of metallo-β-

lactamase NDM-1-producing multidrug-resistant Escherichia coli in Australia.

Antimicrob Agents Chemother. 2010;54: 4914–6. doi:10.1128/AAC.00878-10

61.   Poirel L, Savov E, Nazli A, Trifonova A, Todorova I, Gergova I, et al. Outbreak

caused by NDM-1- and RmtB-producing *Escherichia coli* in Bulgaria.

Antimicrob Agents Chemother. 2014;58: 2472–4. doi:10.1128/AAC.02571-13

62.   Yoo JS, Kim HM, Koo HS, Yang JW, Yoo JI, Kim HS, et al. Nosocomial

transmission of NDM-1-producing Escherichia coli ST101 in a Korean hospital. J

Antimicrob Chemother. 2013;68: 2170–2172. doi:10.1093/jac/dkt126

63.   Yamaji R, Friedman CR, Rubin J, Suh J, Thys E, McDermott P, et al. A

Population-Based Surveillance Study of Shared Genotypes of *Escherichia coli*

Isolates from Retail Meat and Suspected Cases of Urinary Tract Infections.

Bradford PA, editor. mSphere. 2018;3: e00179-18. doi:10.1128/mSphere.00179-

18

64. Fernandes MR, McCulloch JA, Vianello MA, Moura Q, Pérez-Chaparro PJ,

Esposito F, et al. First Report of the Globally Disseminated IncX4 Plasmid

Carrying the *mcr-1* Gene in a Colistin-Resistant *Escherichia coli* Sequence Type

101 Isolate from a Human Infection in Brazil. Antimicrob Agents Chemother.

2016;60: 6415–7. doi:10.1128/AAC.01325-16

65. Gerhold G, Schulze MH, Gross U, Bohne W. Multilocus sequence typing and

CTX-M characterization of ESBL-producing E. Coli: A prospective single-centre

study in Lower Saxony, Germany. Epidemiol Infect. 2016;144: 3300–3304.

doi:10.1017/S0950268816001412

66. Khan ER, Aung MS, Paul SK, Ahmed S, Haque N, Ahamed F, et al. Prevalence

and Molecular Epidemiology of Clinical Isolates of *Escherichia coli* and

Klebsiella pneumoniae Harboring Extended-Spectrum Beta-Lactamase and

Carbapenemase Genes in Bangladesh. Microb Drug Resist. 2018;24: 1568–1579.

doi:10.1089/mdr.2018.0063

67. Pfeifer Y, Witte W, Holfelder M, Busch J, Nordmann P, Poirel L. NDM-1-

producing Escherichia coli in Germany. Antimicrob Agents Chemother. 2011;55:

1318–9. doi:10.1128/AAC.01585-10

68. Williamson DA, Freeman JT, Roberts SA, Heffernan H, Dyet K, Paterson DL, et

al. Rectal colonization with New Delhi metallo- -lactamase-1-producing

Escherichia coli prior to transrectal ultrasound (TRUS)-guided prostate biopsy. J

Antimicrob Chemother. 2013;68: 2957–2959. doi:10.1093/jac/dkt266

69.   Potron A, Poirel L, Rondinaud E, Nordmann P. Intercontinental spread of OXA-

48 beta-lactamase-producing Enterobacteriaceae over a 11-year period, 2001 to

2011. Eurosurveillance. 2013;18: 20549. doi:10.2807/1560-

7917.ES2013.18.31.20549

70.   Samuelsen Ø, Overballe-Petersen S, Bjørnholt JV, Brisse S, Doumith M,

Woodford N, et al. Molecular and epidemiological characterization of

carbapenemase-producing Enterobacteriaceae in Norway, 2007 to 2014. Butaye

P, editor. PLoS One. 2017;12: e0187832. doi:10.1371/journal.pone.0187832

71.   Mantilla-Calderon D, Jumat MR, Wang T, Ganesan P, Al-Jassim N, Hong P-Y.

Isolation and Characterization of NDM-Positive Escherichia coli from Municipal

Wastewater in Jeddah, Saudi Arabia. Antimicrob Agents Chemother. 2016;60:

5223–31. doi:10.1128/AAC.00236-16

72.   Braga JFV, Chanteloup NK, Trotereau A, Baucheron S, Guabiraba R, Ecco R, et

al. Diversity of *Escherichia coli* strains involved in vertebral osteomyelitis and

arthritis in broilers in Brazil. BMC Vet Res. 2016;12: 140. doi:10.1186/s12917-

016-0762-0

73.   Koo H-J, Kwak H-S, Yoon S-H, Woo G-J. Phylogenetic group distribution and

prevalence of virulence genes in Escherichia coli isolates from food samples in

South Korea. World J Microbiol Biotechnol. 2012;28: 1813–1816.

doi:10.1007/s11274-011-0954-5

74.   Müller A, Jansen W, Grabowski NT, Monecke S, Ehricht R, Kehrenberg C.

ESBL- and AmpC-producing *Escherichia coli* from legally and illegally

imported meat: Characterization of isolates brought into the EU from third

countries. Int J Food Microbiol. 2018;283: 52–58.

doi:10.1016/J.IJFOODMICRO.2018.06.009

1023 75. Xie M, Lin D, Chen K, Chan EWC, Yao W, Chen S. Molecular Characterization

1024   of Escherichia coli Strains Isolated from Retail Meat That Harbor blaCTX-M and

1025   fosA3 Genes. Antimicrob Agents Chemother. 2016;60: 2450–2455.

1026   doi:10.1128/AAC.03101-15

1027 76. Wu C, Wang Y, Shi X, Wang S, Ren H, Shen Z, et al. Rapid rise of the ESBL

1028   and *mcr-1* genes in *Escherichia coli* of chicken origin in China, 2008–2014.

1029   Emerg Microbes Infect. 2018;7: 1–10. doi:10.1038/s41426-018-0033-1

1030 77. Boehmer T, Vogler AJ, Thomas A, Sauer S, Hergenroether M, Straubinger RK,

1031   et al. Phenotypic characterization and whole genome analysis of extended-

1032   spectrum beta-lactamase-producing bacteria isolated from dogs in Germany.

1033   Mokrousov I, editor. PLoS One. 2018;13: e0206252.

1034   doi:10.1371/journal.pone.0206252

1035 78. Zhou X, García-Cobos S, Ruijs GJHM, Kampinga GA, Arends JP, Borst DM, et

1036   al. Epidemiology of Extended-Spectrum β-Lactamase-Producing *E. coli* and

1037   Vancomycin-Resistant Enterococci in the Northern Dutch–German Cross-Border

1038   Region. Front Microbiol. 2017;8: 1914. doi:10.3389/fmicb.2017.01914

1039 79. Wang Y, Tian G-B, Zhang R, Shen Y, Tyrrell JM, Huang X, et al. Prevalence,

1040   risk factors, outcomes, and molecular epidemiology of *mcr-1*-positive

1041   *Enterobacteriaceae* in patients and healthy adults from China: an

1042   epidemiological and clinical study. Lancet Infect Dis. 2017;17: 390–399.

1043   doi:10.1016/S1473-3099(16)30527-8

1044 80. Shrestha B, Tada T, Miyoshi-Akiyama T, Shimada K, Ohara H, Kirikae T, et al.

1045   Identification of a novel NDM variant, NDM-13, from a multidrug-resistant

1046   Escherichia coli clinical isolate in Nepal. Antimicrob Agents Chemother.

1047   2015;59: 5847–50. doi:10.1128/AAC.00332-15

1048   81.   Herrero-Fresno A, Zachariasen C, Hansen MH, Nielsen A, Hendriksen RS,

1049   Nielsen SS, et al. Apramycin treatment affects selection and spread of a

1050   multidrug-resistant Escherichia coli strain able to colonize the human gut in the

1051   intestinal microbiota of pigs. Vet Res. 2016;47: 12. doi:10.1186/s13567-015-

1052   0291-z

1053   82.   Trobos M, Lester CH, Olsen JE, Frimodt-Moller N, Hammerum AM. Natural

1054   transfer of sulphonamide and ampicillin resistance between Escherichia coli

1055   residing in the human intestine. J Antimicrob Chemother. 2009;63: 80–86.

1056   doi:10.1093/jac/dkn437

1057   83.   Picard B, Garcia JS, Gouriou S, Duriez P, Brahimi N, Bingen E, et al. The link

1058   between phylogeny and virulence in *Escherichia coli* extraintestinal infection.

1059   Infect Immun. 1999;67: 546–53. Available: https://iai.asm.org/content/67/2/546

1060   84.   Johnson JR, Clermont O, Menard M, Kuskowski MA, Picard B, Denamur E.

1061   Experimental Mouse Lethality of *Escherichia coli* Isolates, in Relation to

1062   Accessory Traits, Phylogenetic Group, and Ecological Source. J Infect Dis.

1063   2006;194: 1141–1150. doi:10.1086/507305

1064   85.   Duriez P, Clermont O, Picard B, Denamur E, Bonacorsi S, Bingen E, et al.

1065   Commensal *Escherichia coli* isolates are phylogenetically distributed among

1066   geographically distinct human populations. Microbiology. 2001;147: 1671–1676.

1067   doi:10.1099/00221287-147-6-1671

1068   86.   Hallström T, Resman F, Ristovski M, Riesbeck K. Binding of complement

1069   regulators to invasive nontypeable Haemophilus influenzae isolates is not

1070   increased compared to nasopharyngeal isolates, but serum resistance is linked to

1071   disease severity. J Clin Microbiol. 2010;48: 921–7. doi:10.1128/JCM.01654-09

87.   Hughes C, Phillips R, Roberts AP. Serum resistance among Escherichia coli

strains causing urinary tract infection in relation to O type and the carriage of

hemolysin, colicin, and antibiotic resistance determinants. Infect Immun.

1982;35: 270–5. Available: http://www.ncbi.nlm.nih.gov/pubmed/7033137

88.   Adler NRL, Stevens MP, Dean RE, Saint RJ, Pankhania D, Prior JL, et al.

Systematic Mutagenesis of Genes Encoding Predicted Autotransported Proteins

of *Burkholderia pseudomallei* Identifies Factors Mediating Virulence in Mice,

Net Intracellular Replication and a Novel Protein Conferring Serum Resistance.

Wooten RM, editor. PLoS One. 2015;10: e0121271.

doi:10.1371/journal.pone.0121271

89.   Desroches M, Clermont O, Lafeuillade B, Rodriguez C, Darty M, Royer G, et al.

Genotypic and phenotypic characteristics of *Escherichia coli* involved in

transfusion-transmitted bacterial infections: implications for preventive

strategies. Transfusion. 2018;58: 1940–1950. doi:10.1111/trf.14812

90.   Peirano G, Mulvey GL, Armstrong GD, Pitout JDD. Virulence potential and

adherence properties of Escherichia coli that produce CTX-M and NDM β-

lactamases. J Med Microbiol. 2013;62: 525–530. doi:10.1099/jmm.0.048983-0

91.   Scott VCS, Haake DA, Churchill BM, Justice SS, Kim JH. Intracellular Bacterial

Communities: A Potential Etiology for Chronic Lower Urinary Tract Symptoms.

Urology. Elsevier Inc.; 2015. pp. 425–431. doi:10.1016/j.urology.2015.04.002

92.   Rivardo F, Martinotti MG, Turner RJ, Ceri H. Synergistic effect of lipopeptide

biosurfactant with antibiotics against Escherichia coli CFT073 biofilm. Int J

Antimicrob Agents. 2011;37: 324–331. doi:10.1016/j.ijantimicag.2010.12.011

93.   Wu H, Moser C, Wang HZ, Høiby N, Song ZJ. Strategies for combating bacterial

biofilm infections. International Journal of Oral Science. Sichuan University

Press; 2015. pp. 1–7. doi:10.1038/ijos.2014.65

94. Kjelstrup CK, Barber AE, Norton JP, Mulvey MA, L'Abée-Lund TM.

Escherichia coli O78 isolated from septicemic lambs shows high pathogenicity in

a zebrafish model. Vet Res. 2017;48. doi:10.1186/s13567-016-0407-0

95. Mittal S, Sharma M, Chaudhary U. Biofilm and multidrug resistance in

uropathogenic *Escherichia coli*. Pathog Glob Health. 2015;109: 26–29.

doi:10.1179/2047773215Y.0000000001

96. Hauser E, Mellmann A, Semmler T, Stoeber H, Wieler LH, Karch H, et al.

Phylogenetic and molecular analysis of food-borne shiga toxin-producing

Escherichia coli. Appl Environ Microbiol. 2013;79: 2731–40.

doi:10.1128/AEM.03552-12

97. Mellmann A, Bielaszewska M, Köck R, Friedrich AW, Fruth A, Middendorf B,

et al. Analysis of Collection of Hemolytic Uremic Syndrome–associated

Enterohemorrhagic *Escherichia coli*. Emerg Infect Dis. 2008;14: 1287–1290.

doi:10.3201/eid1408.071082

98. Bai X, Mernelius S, Jernberg C, Einemo IM, Monecke S, Ehricht R, et al. Shiga

toxin-producing *Escherichia coli* infection in Jönköping county, Sweden:

Occurrence and molecular characteristics in correlation with clinical symptoms

and duration of stx shedding. Front Cell Infect Microbiol. 2018;8: 125.

doi:10.3389/fcimb.2018.00125

99. Ferdous M, Friedrich AW, Grundmann H, de Boer RF, Croughs PD, Islam MA,

et al. Molecular characterization and phylogeny of Shiga toxin–producing

Escherichia coli isolates obtained from two Dutch regions using whole genome

sequencing. Clin Microbiol Infect. 2016;22: 642.e1-642.e9.

doi:10.1016/j.cmi.2016.03.028

100. Jønsson R, Struve C, Jenssen H, Krogfelt KA. The wax moth Galleria mellonella

as a novel model system to study Enteroaggregative Escherichia coli

pathogenesis. Virulence. Taylor and Francis Inc.; 2017. pp. 1894–1899.

doi:10.1080/21505594.2016.1256537

101.   Guerrieri CG, Pereira MF, Galdino ACM, Santos ALS dos, Elias WP, Schuenck

RP, et al. Typical and Atypical Enteroaggregative *Escherichia coli* Are Both

Virulent in the *Galleria mellonella* Model. Front Microbiol. 2019;10.

doi:10.3389/fmicb.2019.01791

102.   Heitmueller M, Billion A, Dobrindt U, Vilcinskas A, Mukherjee K. Epigenetic

mechanisms regulate innate immunity against uropathogenic and commensal-like

Escherichia coli in the surrogate insect model Galleria mellonella. Infect Immun.

2017;85. doi:10.1128/IAI.00336-17

103.   Ciesielczuk H, Betts J, Phee L, Doumith M, Hope R, Woodford N, et al.

Comparative virulence of urinary and bloodstream isolates of extra-intestinal

pathogenic Escherichia coli in a Galleria mellonella model. Virulence. 2015;6:

145–151. doi:10.4161/21505594.2014.988095

104.   Williamson DA, Mills G, Johnson JR, Porter S, Wiles S. In vivo correlates of

molecularly inferred virulence among extraintestinal pathogenic Escherichia coli

(ExPEC) in the wax moth Galleria mellonella model system. Virulence. 2014;5:

388–393. doi:10.4161/viru.27912

105.   Beghain J, Bridier-Nahmias A, Nagard H Le, Denamur E, Clermont O.

ClermonTyping: An easy-to-use and accurate in silico method for *Escherichia*

genus strain phylotyping. Microb Genomics. 2018;4: e000192.

doi:10.1099/mgen.0.000192

1146

1147 **TABLES AND CAPTIONS**

1148

1149 <p style="text-align:center">**Table 1 -** EC121 annotation overview.[a]</p>

Protein features	Occurrence
Hypothetical proteins	702
Proteins with functional assignments	4,473
Proteins with E.C.[b] number assignments	1,300
Proteins with G.O.[b] assignments	1,076
Proteins with Pathway assignments	911
Proteins with PLfam[b] assignments	5,063
Proteins with PGfam[b] assignments	5,064

1150 *a*. Results obtained using PATRIC annotation service;

1151 *b*. Abbreviations: E.C. number - Enzyme commission universal G.O. - Gene Ontology

1152 Consortium; PLfam- PATRIC genus-specific family; PGfam- PATRIC cross-genus

1153 family.

1154

Table 2 – Complete virulence factors identified

Virulence traits	Associated with virulence in			
	Other Genera	DEC	ExPEC	Various
Colonization and invasion	MlaA, MsbB, Sfm, MisL, IcsP	Lpf-1$_{O26,}$ EhaG, Hcp, Elf, EhaA, EhaB, CFA-I	FdeC, Pix, Ygi, Yad, Yeh, Yra, Yfc, YchO, IbeB, IbeC, EptC, OmpA	Type 1 Fimbriae (H190), Ecp, Curli
Immune evasion	SodB, TrxA, SirA, FpkA		Iss, OmpTp[a], OmpTc[a], Mig-14, HlyF	RelA
Iron acquisition				Sit, Iro, Fhu, Enterobactin
Regulators	DsbA, DegP, SlyA, CpxAR	EvgAS	DsbAB, PhoPQ	QseBC, RcsAB
Toxins and Bacteriocin			Microcin V, Colicin B, Colicin M	ClyA, Hly III
Non-LEE effectors		EspL1, EspL4, EspX1, EspX4, EspX5, EspR1		

 [a] OmpTc for chromosomal variant of OmpT protein, and OmpTp for plasmid variant of OmpT protein.

1157 **Table 3 -** Predicted phages detected in the EC121 genome.

Region	Length	Completeness[a]	Most Common Phage/accession number	GC %
1	6.3 kb	Incomplete	*Bacillus* phage G/NC_023719	51.27
2	30.9 kb	Incomplete	*Salmonella* phage 118970_sal3/NC_031940	49.82
3	14.2 kb	Questionable	Enterobacteria phage P88/NC_026014	49.75
4	20 kb	Incomplete	*Shigella* phage POCJ13/NC_025434	47.95
5	28.5 kb	Incomplete	*Shigella* phage SfII/NC_021857	45.37
6	14.4 kb	Intact	Enterobacteria phage Fels-2/NC_010463	49.56
7	9.4 kb	Incomplete	*Bacillus* phage Shanette/NC_028983	49.50
8	7.5 kb	Incomplete	Enterobacteria phage phi92/NC_023693	44.44
9	10.2 kb	Questionable	Enterobacteria phage P2/NC_001895	55.39
10	3.8 kb	Incomplete	Bacteriophage WPhi/NC_005056	50.16
11	8.4 kb	Questionable	Phage cdtI/NC_009514	46.05
12	6.7 kb	Incomplete	*Shewanella* sp. phage¼/NC_025436	47.92

1158 *a*. Phage completeness was determined by scores obtained in the search algorithm

1159 (PHASTER) based on the number of the phage's specifics CDSs detected in the

1160 analyzed region.

1161

1162 **Table 4 -**Antimicrobial resistance genotype and phenotype observed in strain EC121.

Antimicrobial Class	Genotype	Phenotype	
		Antimicrobial agent	MIC^b(µg /mL)
Aminoglycosides	*aph(3'')-Ib*, *aph(6)-Id*, *aph(4)-Ia*, *aac(3)-Iva*	Amikacin	2
		Gentamicin	**32**
β-lactams	*bla*$_{TEM-1B}$, *bla*$_{CTX-M-2}$	Ampicillin	**≥256**
		Ceftazidime	**64**
		Ceftriaxone	**≥512**
		Cefepime	**16**
		Piperacillin/Tazobactam	**>256/4**
		Meropenem	≤0.5
		Ertapenem	≤0.5
		Imipenem	≤0.5
		Aztreonam	**>32**
Sulfonamides/ Trimethoprim	*sul2*, *sul1*, *sul1'*, *dfrA14*, *dfrA7*	Trimethoprim/ Sulfamethoxazole	**>128/ 2,432**
Phenicols	*catA1*	Chloramphenicol	**>64**
Tetracyclines	*tet(A)*	Minocycline	**16**
		Tigecycline	**1**
Fluoroquinolones	*parE*(S$_{458}$A)*; parC* (S$_{80}$I); *gyrA* (S$_{83}$L and D$_{87}$Y)	Ciprofloxacin	**>64**
Polymyxins	-	Colistin	≤0.25
		Polymyxin B	≤0.25
Fosfomycin	-	Fosfomycin	0.25
Macrolides	*mph(A)*, *mdf(A)*	Azithromycin	NT

1163 *a*. chromosomal mutations and mobile genes related to antimicrobial resistance

1164 identified;

1165 *b*. Resistance is highlighted in bold, following EUCAST (2020) breakpoints.

1166 NT, not tested.

1167

1168

Table 5 - Resistance profile, replicon type, and virulence genes identified in the EC121's plasmids transferred to *E. coli* K-12 strains

Traits	EC121	J53	C600	ACG04	ACC09
Antimicrobial [a]					
Ceftazidime	64	0.25	0.25	**8**	**2**
Ceftriaxone	≥64	≤0.125	≤0.125	**>64**	**>64**
Cefepime	16	0.25	≤0.125	**16**	**2**
Piperacillin/ Tazobactam	>256	2	2	**8**	**2**
Aztreonam	>32	≤0.125	≤0.125	**16**	**4**
Amikacin	4	2	2	**16**	**2**
Gentamycin	16	0.5	0.5	**32**	0.5
Trimethoprim/ Sulfamethoxazole	>128	≤0.25	≤0.25	**>128**	**>128**
Ciprofloxacin	>64	≤0.125	≤0.125	≤0.125	≤0.125
Minocycline	16	1	1	**4**	1
Tigecycline	1	0.25	0.125	0.25	0.125
Replicon type [b]					
IncHI2	+	-	-	+	+
IncL/M	+	-	-	+	-
IncFIB	+	-	-	+	-
IncN	+	-	-	+	-
Virulence [c]					
hlyF	+	-	-	+	-
iroN	+	-	-	+	-
sitA	+	-	-	+	-
Resistance gene					
*bla*CTX-M	+	-	-	+	+

[a] The antimicrobial resistance phenotype of transconjugant strains was assessed by
minimum inhibitory concentration (MIC), performed using the broth microdilution
method according to EUCAST guideline, and was expressed in µg/mL. Bold values
indicate an increased MIC value when comparing strains bearing plasmids with its
receptor without plasmid.

[b] Replicons identified in EC121 by PlasmidFinder, and screened by PCR;

[c] Virulence genes identified in EC121 genome and screened by PCR;

+ for presence of the gene; - for the absence of the gene;

1177

1178

1179

1180

1181

 Table 6 -Estimated bacterial inoculum resistant to serum activity (CFU/mL).[a]

	Strains					
	J96		EC121		C600	
Challenge period	**NHS**	**inHS**	**NHS**	**inHS**	**NHS**	**inHS**
30 min	10^1	10^1	10^2	10^1	10^8	10^1
1 h	10^2	10^1	10^2	10^1	NG	10^1
2 h	10^2	10^1	10^2	10^1	NG	10^1

1183 *a.* Values represent the approximate relative mean of the lowest bacterial inoculum that

1184 remained viable after the challenge. All assays were performed in triplicate using 50%

1185 serum diluted in PBS (v/v);

1186 NHS - Normal human serum; inHS - inactivated human serum; NG - no growth after

1187 the challenge.

1188

 FIGURES CAPTIONS

 Figure 1 - EC121 Phylogenetic tree.

A phylogenetic tree was built with genomes of reference *E. coli* strains, of relevant

pathogenic strains from all *E. coli* pathotypes, and some strains from ST101, using

Maximum Likelihood-based algorithm (RAxML) in PATRIC. In parenthesis, the strain

pathotype (when known), the Sequence type, and phylogroup according to

ClermonTyping [105] are depicted. Diarrheagenic *E. coli* strains were in orange;

Extraintestinal pathogenic *E. coli* strains were in red, commensal or strains that the

origin was not described were in black. In bold and with purple label background, the

strain studied in present work. Bootstrap upper than 80 were informed in the tree.

1199

Figure 2 - ST101 complex phylogenetic tree.

The phylogenetic tree was built using 95 *E. coli* strains from ST101 complex (ST101,

ST359, ST2480, ST5957, and ST6388) from diverse source and countries using all

shared protein among them on FastTree2 to build the tree. *E. coli* strain IAI1, *E. coli*

O157:H7 str Sakai, and *E. coli* O104:H4 str 2011c-3493 are used as outgroups.

Bootstrap upper than 50 were informed in the tree. Label colors are related with isolates

origin or host diseases, being Shiga toxin-producing *E. coli* (STEC) in orange

independently of the origin; Diarrheagenic *E. coli* (DEC) strains in yellow;

Extraintestinal pathogenic *E. coli* (ExPEC) strains in red; isolates from microbiota in

brown; isolates from retail food or environment in green; strains that the origin was not

described in black. One strain isolated from Crohn's disease is in purple. In bold and

with purple label background, the strain of present work. * All strains that harbor AMR

for 3 or more antimicrobial classes were designed as MDR. Other FQ (Fluoroquinolone)

resistance genes detected were the mobile genes *qepA* and *aac(6')-Ib-cr*. Mutations that

confer resistance to FQ were not considered to build the AMR information in this tree.

1215

Figure 3 - CRISPR locus composition and biological systems assignment in EC121.

A – Genomic architecture representation of CRISPR *locus* in the EC121 genome; the

image was obtained in MacSyFinder from genome annotation; **B -** Representation of

EC121 genome composition in subsystems based on protein biological data obtained *in*

silico; **C -** EC121 genome schematic composition, based on annotation, ordered by

contig size. In circle from outer to the inner, forward strand, reverse strand, RNA related

genes, antimicrobial resistance, virulence factors, GC content, and GC skew. The colors

in forward and reverse strands correspond to the subsystems presented in B. Figures 2B

and 2C were obtained using the comprehensive genome analysis service at PATRIC.

1225

1226

Figure 4 - Biofilm formation on abiotic surfaces. A. Qualitative assay showing the

EC121 adhered to glass coverslips after 24h of incubation at 37 °C, in DMEM, EAEC

042 and HB101 were used as positive and negative controls, respectively. Bacteria were

stained with crystal violet, observed in optical microscopy (OM) 400 x **B.** Quantitative

biofilm assays were performed in DMEM and TSB, at 37°C, for 24 h, comparing

EC121 capacity to produce biofilm in polystyrene surface. EAEC 042 was used as

positive control, CFT073 as an ExPEC control and HB101 as negative control. *P*

values: ** <0.01; *** <0.001; ns >0.05.

.

Figure 5 – Quantitative adherence assay on HeLa cell. Quantitative assay performed

in HeLa cells (3-h of incubation at 37°C) in the absence of D-mannose, except when

indicated.; M+, assay performed in the presence of 2% D-Mannose. The CFT073, C1845, and C600 strains were used as controls. Experiments were done in biological duplicates and experimental triplicates. *P* values: * < 0.05, ** < 0.01.

Figure 6 – **Interaction and invasion in different eukaryotic cell lineages**. Short period quantitative invasion assay was performed in Caco-2 and T24 cells using a MOI of 50 in confluent cell cultures. All assays were carried out in the absence of D-mannose, except for EC121 M+, where 2% D-mannose was added in the media; *E. coli* CFT073 was used as ExPEC control and *E. albertti*1551-2::eae was used as adherent non-invasive control. The assays were performed in experimental triplicates and biological duplicates. *P* values: ** <0.008; ns > 0.05

Figure 7 – *Galleria mellonella* infection model. The virulence of EC121 was assessed using an inoculum of 10^4 CFU per *G. mellonella* larvae. The larvae survival rate, expressed by Kaplan-Meier plots indicated that EC121 was more virulent than the negative control (saline) and the non-pathogenic K-12 derived *E. coli* C600, while there was no significant difference between the survival rates of EC121 and the ExPEC prototype strain CFT073. *P* value: ** <0.005; **** <0.0001. The assay was performed three time with five larvae per group.

Tree scale: 0.001 ⊢—⊣

Escherichia fergusoni NCTC 12128
Escherichia fergusonii ATCC 35469
Escherichia coli IAI39 (ExPEC ST62-F)
Escherichia coli APEC O2-211 (ExPEC ST117-G)
Escherichia coli O24:H4 strain ECO1127 (ExPEC ST117-G)
Escherichia coli O127:H6 str. E2348/69 (EPEC ST15-B2)
Escherichia coli O25b:H4-ST131 str. EC958 (ExPEC ST131-B2)
Escherichia coli strain AR_0089 (ST131-B2)
Escherichia coli 536 (ExPEC ST127-B2)
Escherichia coli ABU 83972 (ST73-B2)
Escherichia coli CFT073 (ExPEC ST73-B2)
Escherichia coli J96 (ExPEC ST12-B2)
Escherichia coli APEC IMT5155 (ExPEC ST140-B2)
Escherichia coli S88 (ExPEC ST95-B2)
Escherichia coli APEC O1 (ExPEC ST95-B2)
Escherichia coli IHE3034 (ExPEC ST95-B2)
Escherichia coli APEC O18 (ExPEC ST95-B2)
Escherichia coli UTI89 (ExPEC ST95-B2)
Escherichia coli RS218 (ExPEC ST95-B2)
Escherichia coli 042 (EAEC ST414-D)
Escherichia coli UMN026 (ExPEC ST597-D)
Escherichia coli O145:H28 str. RM13514 (STEC ST32-D)
Escherichia coli O157:H7 str. Sakai (EHEC ST11-E)
Escherichia coli O157:H7 str. EDL933 (EHEC ST11-E)
Escherichia coli 53638 (EIEC ST6-A)
Escherichia coli strain H10407 (ETEC ST48-A)
Escherichia coli O162:H9 strain ECO0002 (ExPEC ST10-A)
Escherichia coli O29:H4 strain ECO1495 (ExPEC ST10-A)
Escherichia coli str. K-12 substr. MG1655 (ST10-A)
Escherichia coli BW25113 (K-12 ST10-A)
Escherichia coli BW2952 (Laboratorial strain ST10-A)
Escherichia coli APEC O78 (ExPEC ST23-C)
Escherichia coli O28ac:NM str. 02-3404 (EIEC ST4627-B1)
Escherichia coli IAI1 (commensal ST1128-B1)
Escherichia coli O111:H- str. 11128 (STEC ST16-B1)
Escherichia coli O26:H11 str. 11368 (STEC ST21-B1)
Escherichia coli O103 str. RM8385 (STEC ST723-B1)
Escherichia coli O104:H4 str. ON2010 (EAHEC ST678-B1)
Escherichia coli O104:H4 str. 2009EL-2050 (EAHEC ST678-B1)
Escherichia coli O104:H4 str. 2011C-3493 (EAHEC ST678-B1)
Escherichia coli O104:H4 strain LB226692 (EAHEC ST678-B1)
Escherichia coli strain 015PP2015 (ExPEC ST101-B1)
Escherichia coli strain 21_Esco_HA-NL (ST101-B1)
Escherichia coli strain H8Ecoli (ST101-B1)
Escherichia coli strain STEC 2359 (STEC ST101-B1)
Escherichia coli strain zlp18 (ST101-B1)
Escherichia coli strain B146 (ST101-B1)
Escherichia coli strain MOD1-EC1635 (ST101-B1)
Escherichia coli strain EC121 (ExPEC ST101-B1)
Escherichia coli strain Ecol_224 (ST101-B1)
Escherichia coli strain 50673720 (ST101-B1)

A

cas3 *cse1* *cse2* *cas7* *cas5* *cas6* *cas1* *cas2*

0   1000 aa

B

Subsystem (Subsystems, Genes)

- METABOLISM (115, 933)
- STRESS RESPONSE, DEFENSE, VIRULENCE (47, 221)
- PROTEIN PROCESSING (46, 274)
- ENERGY (36, 342)
- CELLULAR PROCESSES (25, 204)
- MEMBRANE TRANSPORT (25, 124)
- DNA PROCESSING (21, 121)
- RNA PROCESSING (14, 75)
- CELL ENVELOPE (13, 129)
- MISCELLANEOUS (12, 83)
- REGULATION AND CELL SIGNALING (7, 46)

C

A

EC121   EAEC 042   HB101

B

DMEM   **TSB**