Stress-mediated convergence of splicing landscapes in male and female Rock Doves

Andrew S. Lang[a][*], Suzanne H. Austin[b], Rayna M. Harris[b], Rebecca M. Calisi[b], Matthew

5   D. MacManes[a]

[a]Department of Molecular, Cellular, and Biomedical Sciences, University of New

Hampshire

[b]Department of Neurobiology, Physiology and Behavior, University of California, Davis

10   *Corresponding author

ABSTRACT

Background

The process of alternative splicing provides a unique mechanism by which eukaryotes are able to produce numerous protein products from the same gene. Heightened variability in the proteome has been thought to potentiate increased behavioral

30   complexity and response flexibility to environmental stimuli, thus contributing to more refined traits on which natural and sexual selection can act. While it has been long known that various forms of environmental stress can negatively affect sexual behavior and reproduction, we know little of how stress can affect the alternative splicing associated with these events, and less still about how splicing may differ between

35   sexes. Using the model of the rock dove (*Columba livia),* our team previously uncovered sexual dimorphism in the basal and stress-responsive gene transcription of a biological system necessary for facilitating sexual behavior and reproduction, the hypothlamic-pituitary-gonadal (HPG) axis. In this study, we delve further into understanding the mechanistic underpinnings of how changes in the environment can affect reproduction

40   by testing the alternative splicing response of the HPG axis to an external stressor in both sexes.

Results

This study reveals dramatic baseline differences in HPG alternative splicing between males and females. However, post submitting subjects to a restraint stress paradigm,

45   we found a significant reduction in these differences between the sexes. In both stress and control treatments, we identified a higher incidence of splicing activity in the

pituitary in both sexes as compared to other tissues. Of these splicing events, the core

exon event is the most abundant form of splicing and more frequently occurs in the

coding regions of the gene. Overall, we observed less splicing activity in the 3'UTR end

50 of transcripts than the 5'UTR or coding regions.

Conclusions

Our results provide vital new insight into sex-specific aspects of the stress response on

the HPG axis at an unprecedented proximate level. Males and females uniquely

respond to stress, yet exhibit splicing patterns suggesting a convergent, optimal splicing

55 landscape for stress response. This information has the potential to inform evolutionary

theory as well as the development of highly-specific drug targets for stress-induced

reproductive dysfunction.

70   **INTRODUCTION**

Organismal behavior and its mechanistic underpinnings have been consistent

quandaries for many biologists (Vermeij, 1991; Sutherland, 1998; Holway & Suarez,

1999; Wong & Candolin, 2015). Even among the few cases in which an adaptive

75   behavior is clearly understood, we have little information on how proximate mechanisms

have led to ultimate behavioral adaptations. One of the mechanisms by which animals

modulate their response to stimuli is by adjusting the levels of endogenous protein

products in physiological systems. (Sapolsky, Romero & Munck, 2000; Zhao et al.,

2017). This variable production may result in differential binding of hormones and

80   signaling peptides, enabling an organism to receive information more accurately about

external stimuli and effectively react (Ohama et al., 2017; Calisi et al., 2018). In addition

to adjusting the quantity of a given protein, organisms may further alter these proteins

by producing slightly modified versions (e.g., isoforms) of each transcript (Smith, Patton

& Nadal-Ginard, 1989; Tapial et al., 2017). This variable modification of gene products

85   is called alternative splicing.

Alternative splicing is a process unique to eukaryotes (Izquierdo & Valcárcel, 2006) that

involves cleavage of transcribed RNA at specific splice sites and varying inclusion or

exclusion of genomic elements (introns and exons). In the human genome,

90   approximately 80% of exons are >200bp in length (Krawczak et al., 2007;

Gudlaugsdottir et al., 2007); however, exon sizes identified in other species vary from a

single base to >17,000bp in length. Each human gene contains, on average, eight

exons (Sakharkar, Chow & Kangueane, 2004). This variable inclusion of genetic

sequences results in a dramatic increase in the number of potential transcript and

95   protein products that a single gene may produce. Alternative splicing presents an

additional mechanism by which mRNA levels and gene expression can be regulated,

while also greatly increasing proteome diversity. Splicing activity is thought to be

responsible for the majority of proteomic diversity in eukaryotes (Birzele, Csaba &

Zimmer, 2008) and, potentially, may be an underlying mechanism of functional genomic

100   evolution (Bush et al., 2017).

Numerous types of splicing events exist that occur at different frequencies in a given

genome and alter proteins in subtle to dramatically different ways (Kim, Magen & Ast,

2007). Cassette exon splicing (also referred to as exon skipping) is the most common

105   type of splicing event in vertebrates and invertebrates, while intron retention is more

common in plants. Additionally, alternative selection of 5' and 3' splice sites, coupled

with variable adenylation of the transcript, results in further modification of protein

products (Gueroussov et al., 2015; Tian & Manley, 2017). The splicing process consists

of two major steps: assembly of the spliceosome and the actual splicing of pre-mRNA

110   (Wang et al., 2015). In brief, the spliceosome is comprised of several small nuclear

ribonucleoproteins that positionally establish the 5' splice site, the branch point

sequence, and the 3' site. An assembly of spliceosome complexes and eight

evolutionarily-conserved RNA-dependent ATPases/helicases is then followed by the

execution of numerous splicing steps, ultimately resulting in exon excision, exon

115   ligation, or intron retention (Wang et al., 2015). The inclusion of an exon in the final

mRNA product is entirely driven by cis- and trans-acting elements/factors. The

interaction of these elements within the splicing process promotes or inhibits

spliceosome activity on various splice regions, resulting in alternative splicing (Wang et

al., 2006; Wang & Burge, 2008).

Alternative splicing mechanisms enable organisms to sense and react to minute

changes in the local environment, allowing both plants and animals to tailor their

responses to their surroundings with extreme precision (Staiger & Brown, 2013;

Preußner et al., 2017). Previous research has revealed unique roles for alternative

125   splicing in the immune response of chickens with avian pathogenic *E.coli* (Sun, 2017),

mediation of abiotic stress response pathways of plants (Laloum, Martín & Duque,

2018), and enhanced fear memory of mice (Nijholt et al., 2004). Alternative splicing has

also been implicated in various aspects of cancer, including oncogenesis (Climente-

Gonzalez et al., 2017) and cancer drug resistance (Chen & Weiss, 2015; Siegfried &

130   Karni, 2018). Some studies have identified a sex-bias in alternative splicing in

Drosophila (McIntyre et al., 2006; Telonis-Scott et al., 2009; Gibilisco et al., 2016), while

others have identified unique sex-specific splicing differences in human brains (Trabzuni

et al., 2013). The diverse roles of alternative splicing in biological processes and

behavioral responses inherently speak to the depth and breadth that alternative splicing

135   drives organismal physiology and behavior, at both local and global levels. By

identifying the splicing landscape that modulates gene expression and mRNA transcript

composition in both males and females, we increase the resolution at which we can

comprehend the proximate mechanisms underlying animal physiology and behavior.

140　In vertebrates, a symphony of physiological events is required to regulate sexual

behavior and reproduction, and these mechanisms are driven by an interconnected

biological system made up of the hypothalamus in the brain, the pituitary gland, and the

gonads (testes/ovaries) (Sower, Freamat & Kavanaugh, 2009; Nozaki, 2013;

MacManes et al., 2017; Calisi et al., 2018). This hypothalamic-pituitary-gonadal (HPG)

145　axis can be disrupted in multiple, complex ways (Johnson et al., 1992; Toufexis et al.,

2014; Geraghty & Kaufer, 2015; Calisi et al., 2018). However, we know little about how

stress affects the HPG axis at the level of alternative splicing, and we know even less

regarding its effects at this level in males versus females. Understanding how the

alternative splicing landscape of the reproductive axis changes in the face of stress will

150　not only offer more insight into how stress can affect reproduction, but deepen our

proximate knowledge of biological processes and sexaully-biased behavioral responses

in general.

　　　　Using the classic reproductive model (Darwin, 1860; Lehrman, 1955; Ball &

Silver, 1983; Buntin, Ruzycki & Witebsky, 1993) and rising genomics model (Shapiro et

155　al., 2013; Gillespie et al., 2013; Domyan et al., 2016; MacManes et al., 2017; Calisi et

al., 2018) of the rock dove, *Columba livia,* we have identified sexual dimorphism in both

basal (MacManes et al., 2017) and restraint stress-responsive (Calisi et al., 2018) HPG

gene expression at the level of RNA transcription. In this study, we traverse beyond the

level of transcription to test for sex-biased alternative splicing patterns in the HPG axis

160　of the rock dove in response to a restraint stress stimulus. Using a relatively highly-

replicated (n=12/sex) study design, we identify significantly similar and different splicing

events between the sexes and in response to restraint stress treatment. To our knowledge, this is the first report of sex-specific splicing events in the HPG axis in response to a stressor.

MATERIALS AND METHODS

Animal Collection

We housed birds at the University of California, Davis, in semi-enclosed aviaries (5' x 4'

170   x 7'), with 8 sexually reproductive adult pairs per aviary. Food and water were provided *ad libitum*. Birds were exposed to natural light, which was supplemented with artificial light on a 14L:10D cycle. We sampled sexually mature males and females that were without eggs or chicks so as to control for physiological changes that occur to facilitate parental care behaviors. Birds were sampled between 0900—1100 (PST) to also control

175   for potential circadian rhythm confounds. All handling and sampling procedures followed approved animal care and handling protocols (UC Davis IACUC #18895). Samples from the control group were taken within 5min of entering their cages. Birds in the restraint stress treatment group were captured in <1min upon entering their aviary and immediately and individually restrained in cloth bags for 30min prior to sampling,

180   replicating methods described in Calisi et al. 2018.

Prior to sampling, birds were anesthetized using isoflurane until unresponsive (<2min), at which time they were immediately decapitated. Trunk blood was collected for the detection of concentrations of the adrenal hormone, corticosterone; high concentrations

185   confirmed the effectiveness of the stress treatment protocol (Calisi et al., 2018). All

tissues (brains, pituitaries, and gonads) were flash frozen on dry ice and transferred to a

-80°C freezer for storage until additional processing, a process described in (Shapiro et

al., 2013; Gillespie et al., 2013; MacManes et al., 2017; Calisi et al., 2018). Briefly, a

cryostat (Leica CM 1860) was used to section the brains coronally at 100μm to enable

190   optimal visualization and biopsy of the hypothalamus, which included the adjoined

lateral septum. Hypothalamic sections, pituitaries, and gonads were preserved in

RNALater (Invitrogen, Thermo Fisher Scientific) and shipped on dry ice from UC Davis

to the University of New Hampshire for cDNA library preparation and sequencing. We

sequenced tissue from whole homogenized hypothalami, pituitaries, testes and

195   (homogenized tissue from the oviduct and ovarian follicles). In total, we processed and

sequenced hypothalami, pituitary glands, and gonads (testes/ovaries) from 48 birds (12

males and 12 females, each, per treatment —control and stress), resulting in 144 cDNA

libraries.

200   *cDNA Library Preparation and Sequencing*

Library preparation and sequencing has been previously described in (Shapiro et al.,

2013; Gillespie et al., 2013; MacManes et al., 2017; Calisi et al., 2018). In brief, all

tissue samples were thawed on ice in an RNAse-free work environment. Total RNA was

extracted with a standard Trizol extraction protocol (Thermo Fisher Scientific, Waltham,

205   MA). Total RNA quality and quantity was assessed with gel electrophoresis and a Broad

Range RNA Qubit assay (Thermo Fisher Scientific, Waltham, MA). Illumina sequence

libraries were prepared using the NEB Next Ultra Directional RNA Library Prep Kit (New

England Biolabs, Ipswich, MA). Quality and quantity of the cDNA libraries were

validated with a Tapestation 2200 Instrument (Agilent, Santa Clara, CA). Libraries were

210   diluted to 5nM and pooled. We sent multiplexed library pools to the New York Genome

Center for 125 base pair paired-end sequencing on a HiSeq 2500 platform. 16 samples

were re-sequenced at Novogene Sequencing Center to increase coverage of those

samples with the lowest read abundance.

215   *Pigeon Genome Annotation*

Many of the currently-available methods for identification of splicing events are heavily

dependent upon genome assembly completeness and annotation quality. The existing

annotated assembly for *C. livia* (GCF_000337935.1) did not provide the level of

contiguity required for accurate identification of splicing events. To address this, we

220   used an unannotated chromosome-level assembly for the rock dove

(GCA_001887795.1). We annotated this chromosome-level assembly with Maker

version 2.3.1(Cantarel et al., 2008). A custom script (available here:

https://github.com/AndrewLangvt/Dissertation/blob/master/cliv2_gff2gtf.py) was used to

generate the required GTF file from the resultant GFF.

Sequence QC, Read Processing, and Splicing Event Identification

Sequence data were downloaded to the Pittsburg Supercomputing Center network

"Bridges" and read quality confirmed with FastQC v0.11.5 (Andrews, 2011). Read data

from all 144 samples was first error corrected with Rcorrector v3 (Song & Florea, 2015),

230   then adapter sequences and corrected reads with PHRED<2 were removed from

sequence datasets.

To accurately identify splicing events in our study system, we used Whippet, an

algorithm that rapidly identifies splicing events from RNAseq data (Sterne-Weiler et al.,

235   2017). (Code available here:

https://github.com/AndrewLangvt/Scripts/tree/master/splicing_analysis/whippet)

Whippet uses annotated transcript features to generate splice graphs and transcript-

level mapping data to calculate Percent Spliced In (PSI) values for all paths through the

directed graph (Sterne-Weiler et al., 2017). Whippet v0.10.4 was used to quantify read

240   data and identify differential splicing events between treatments. Events with probability

>0.95 and abs(deltaPSI) ≥0.1 were deemed significant and considered to be true

splicing events.

Splicing Event Comparisons

245   To understand the sex-specific response to restraint stress, we asked (1) how similar

are males and females at the exonic level (i.e. assessing splicing events between

sexes, within treatments), and (2) how do males and females differ in their response to

stress (i.e., assessing splicing events within the sexes, between treatments). We did not

include analyses between different tissues (e.g., male gonads to female gonads, or

250   male pituitary to male hypothalamus) as numerous other studies have identified tissue-

specific splicing events; these comparisons would not aid in discerning unique splicing

patterns between sexes, or between treatments (Grosso et al., 2008; Saha et al., 2017;

Pineda & Bradley, 2018). Whippet identifies splicing events by type, including Core

Exon (CE), Alternate Acceptor (AA), Alternate Donor (AD), Retained Intron (RI),

255 Tandem Alternative Polyadenylation Site (TE), Tandem Transcription Start Site (TS),

Alternative First Exon (AF), and Alternative Last Exon (AL). Of all the event types

detected by Whippet, core exon, alternate acceptor, alternate donor, and retained intron

encompassed 95% of our identified splicing events. For simplicity, we did not represent

the TE, TS, AF, or AL events in our figure depicting splicing by type even though these

260 events are included in all of the analyses.

Gene Ontology Parent Term Analysis

We assessed abstracted (or parent) gene ontology (GO) terms of spliced genes to

identify patterns of enrichment or depletion of gene functions. Lists of Entrez Gene IDs

265 were analyzed through the "Gene List Analysis" tool at http://www.pantherdb.org to link

GO terms with each ID. These terms were further analyzed with go_abstract.py

(https://github.com/twestbrookunh), a python script that abstracts a provided ontology

term to its "parent" term from the Gene Ontology Consortium Database (Ashburner et

al., 2000; Mi et al., 2017; The Gene Ontology Consortium, 2019). We generated a

270 matrix of expected abundances for each tissue using this same analysis on the entire *C.*

livia genome. GO terms with counts deemed significantly different from expected counts

via Chi-Squared analyses were visualized in R. We calculated differences of counts

from expected values and did not include parent ontology terms of very low abundance

in the genome (<2%) in our visualization.

Exon Size, Location, and Motif Analysis

We used a custom Python script, available at

https://github.com/AndrewLangvt/Scripts/blob/master/splicing_analysis/exon_extract.py

to extract desired exons from the genome and translate them into amino acid

280   sequences based upon the phase and strand delineated in the GFF while

simultaneously determining the length of each exon and the transcript region from which

it was spliced (5'-UTR, CDS, or 3'-UTR). Distributions of exon lengths were compared

to the genomic distribution to determine statistical significance using a Wilcoxon test.

Expected counts of splicing events per region were determined from the genomic

285   proportions and difference from expected was normalized by expected values across

tissues. We further assessed the functional nature of alternatively spliced exons by

investigating these sequences for protein motifs. To accomplish this, we used MOTIF, a

GenomeNet tool (https://www.genome.jp/tools/motif/) which identifies protein motifs

within protein sequences by searching the PFAM database (Finn et al., 2016) and NCBI

290   databases COG (PMID: 10592175), SMART (PMID: 10592234), and TIGRFAM (PMID:

11125044).

RESULTS

295   *Sequencing Results, Read Data, and Code Availability*

Samples were sequenced to a read depth between 2.3 million and 24.5 million read

pairs, for a total of 144,363,425 paired-end reads (more fully described in (Calisi et al.,

2018). Read data corresponding to the control birds are available using the European

Nucleotide Archive project ID PRJEB16136; read data corresponding to the stressed

300   birds are available at PRJEB21082. All code for analyses in this manuscript can be

found at https://github.com/AndrewLangvt/Scripts/tree/master/splicing_analysis/.

Male vs. Female Splicing Comparison

Our first aim was to understand sex-typical splicing in the hypothalamus and pituitary by

305   assessing each tissue for alternative splicing events between males and females. We

counted the number, and type, of alternatively spliced loci between males and females

in each treatment state (control: male vs female; stress: male vs female). This approach

allowed us to determine how the splicing landscape changed between sexes in

response to restraint stress, and in which state the sexes shared a more similar splicing

310   profile. As previously stated, we did not include gonads in this comparison due to

inherent splicing differences between tissue types. Chi-squared tests (hereafter, ChiSq)

were used to determine statistical significance ($p < 0.05$) throughout our analyses; all p-

values, degrees of freedom, and sample sizes are included in Table 1.

Analysis		Description	p	df	n
MvF	tissue	more splicing in P than H	6.18E-04	1	338
		more splicing in HC than HS	1.46E-03	1	158
	Event Type	CE most abundant- H	2.22E-13	3	145
		CE most abundant- P	5.70E-43	3	212

		more RI in HC than HS	0.012	1	27
		more CE in HC than HS	0.053	1	77
		more RI in PC than PS	0.052	1	13
	region	more splicing in CDS- H	5.95E-43	2	197
		more splicing in CDS- P	6.52E-141	2	465
	compare to genome	more splicing in 5'UTR - H	0.015	1	25
		more splicing in 5'UTR - P	0.041	1	47
		>6% decrease in 3UTR- P	2.60E-07	1	12
CvS	tissue	more splicing in P than H or G	1.04E-06	1	490
		MH more active than FH in stress response	9.39E-04	1	139
		FG more active than MG	1.92E-04	1	133
	Event Type	CE most abundant- H	3.64E-07	3	129
		CE most abundant- P	1.34E-43	3	212
		CE most abundant- G	8.74E-15	3	119
		more CE in FG than MG	3.01E-03	1	71
		RI- H significant splicing between sexes	0.041	1	24

	region	more splicing in CDS - H	4.41E-40	2	166
		more splicing in CDS- P	9.60E-233	2	629
		more splicing in CDS- G	5.13E-67	2	221
	compare to genome	less splicing in 3'UTR- H	0.042	1	11
		less splicing in 3'UTR- P	2.87E-12	1	8
		less splicing in 3'UTR- G	8.77E-04	1	8
		less splicing in 5'UTR- P	2.04E-04	1	25
		more splicing in CDS- P	3.50E-04	1	596

Table 1. Statistics for all Chi-Square Tests. This table contains all Chi-Square values, degrees of freedom (df), and sample size (n) for every test of significant splicing events in this paper. H=Hypothalamus. P=Pituitary. G=Gonad. C=Control. S=Stress. CE=Core Exon. RI=Retained Intron. CDS=Coding Sequence. UTR=Untranslated Region.

Male vs. Female Splicing Comparison: Events by Type

In total, we identified 158 splicing events in the hypothalamus and 225 events in the pituitary. When compared to the hypothalamus, the 42% increase of splicing event abundance seen in the pituitary is significant (ChiSq p=6.18e-4). In both tissues, more

325   events were identified in the control state compared to the stress condition (hyp: 99

control/59 stress, pit: 123 control/102 stress), but only the relationship in the hypothalamus was statistically significant (Figure 1, ChiSq p: hyp=1.46e-3; pit=0.162).

330   These total counts were further broken down by event type (Figure 1). The core exon event was the most abundant event identified across sex in both the hypothalamus and pituitary, regardless of treatment (ChiSq p: hyp=2.22e-13, pit=5.70e-43). Of these core exon splice events, there were almost twice as many in the pituitary compared to hypothalamus (pit: 74 control/67 stress, hyp: 48 control/30 stress). Within these event types, we tested for statistical significance between splicing differences in males and

335   females of each treatment group. Retained intron events in the hypothalamus were the only event to differ significantly between treatments (ChiSq p=0.012), with nearly 3 times (280% increase) more splicing events in the control state than the stressed. Both core exon events in the hypothalamus and retained intron events in the pituitary reflected a similar increased abundance in splicing events of the control state, though

340   these relationships were not significant (ChiSq p: CE-Hyp=0.053, RI-Pit=0.052). The distribution of PSI values between males and females did not vary between treatments, indicating that the level of event inclusion/exclusion difference between the sexes was generally unaffected by treatment.

Splicing Event by Type

Figure 1. Splicing events by type for both the a) Male vs Female and b) Control vs Stress comparisons. Rows denote tissue type (labeled on the right), and counts of splicing events are further broken down by event type. Alternatively spliced genes in the male vs female analysis revealed, in both tissues, more events in the control versus

350   restraint stress condition. The core exon event was the most abundant regardless of

tissue or treatment. Light blue represents the control group; yellow is restraint stress. Hypothalamic retained intron events were the only event to differ significantly between treatments, represented by a red star (ChiSq p=0.012). In the control vs splicing comparison, more splicing occurred in the male hypothalamus; while in the gonad, more

355   splicing occurred in the female. Blue represents males; green represents females. Red stars represent statistical significance between abundances in males and females, with more core exon splice events occurring in the female gonad than male (ChiSq p=3.01e-3), and more hypothalamic retained intron events found in males than females (ChiSq p=0.041).

Male vs. Female Splicing Comparison: Genes of Interest
Using our comparison of male to female splicing patterns, we were able to identify sex-specific alternatively spliced genes in the stress response. We provide a full list of spliced genes within each comparison (Table S1). Some of these spliced genes are

365   involved in functional gene expression within the HPG axis. POU class 2 homeobox 1 (*POU2F1*), a transcription factor that regulates transcription of gonadotropin-releasing hormone (GnRH) (Chandran & DeFranco, 1999; Cheng et al., 2002), is alternatively spliced in the male pituitary stress response. GnRH is a primary regulator of the HPG axis (Pohl et al., 1983; Campbell, Dobson & Scaramuzzi, 1998; Takahashi et al., 2016).

370   Splicing of *POU2F1* likely affects the HPG axis, indirectly, by modulating transcription of GnRH (Rosen, Jameel & Barkan, 1988; Schneider, Tomek & Gründker, 2006; Jin & Yang, 2014). Through alternative splicing of the *POU2F1* gene in the pituitary, males may be altering signaling pathways within the HPG axis to optimize stress response.

375   Few genes were consistently alternatively spliced between males and females in both

treatments. Those that did exhibit consistent alternative splicing between the sexes

were often related to immune function. Rap1 GTP-ase activating protein 1 (*RAP1GAP*)

is consistently alternatively spliced between male and female hypothalami, in both the

control and stress treatments. Previous findings have shown *RAP1GAP* to be a putative

380   oncogene (Nellore et al., 2009). This gene mediates the strength of cell adhesions

through regulation of Rap1, thus modulating T-cell response (Katagiri et al., 2002).

Alternatively spliced between sexes in the pituitary, P-selectin (*SELP*) is known to

preserve immune function in mice (González-Tajuelo et al., 2017). The corresponding

ligand, P-selectin glycoprotein ligand-1 (PSGL-1), negatively regulates T-cell response

385   through binding of SELP (Matsumoto, Miyasaka & Hirata, 2009). Genes consistently

alternatively spliced between males and females may reflect splicing-level sexual

dimorphism, indicating that males and females inherently differ in their splicing

landscapes. Further, these differences appear to speak to a sex-specific stress

response through modification of genes related to immune processes. Through future

390   study of the splicing landscape of these genes of interest in additional tissues and

states, we will likely reveal additional inherent splicing differences between the sexes.

Male vs. Female Splicing Comparison: Gene Ontology

By observing abundances of parent ontology terms significantly deviating from genomic

395   expectation, we were able to gain a broader understanding of gene-types targeted by

alternative splicing. In the list of significant Molecular Function terms (Figure 2), splicing

of organic and heterocyclic compound genes is underrepresented in the pituitary, while small molecular and drug binding is overrepresented in the hypothalamus regardless of treatment. There was an interaction seen in the hypothalamus; splicing of organic and

400   heterocyclic compound genes was overrepresented in the stress treatment, though underrepresented in the control group. Biological Process terms suggest that males and females differ very little in their splicing profile of metabolic genes in either the hypothalamus or pituitary, given there were fewer spliced genes with metabolism GO terms in these tissues (Figure S1). Finally, splicing events in stressed males and

405   females are more abundant in genes related to cell/neuronal structure of the pituitary than the hypothalamus (Figure S2).

Figure 2. Molecular Function GO analysis, Male vs. Female (normalized counts of

410   observed-expected). Splicing in the pituitary is more prevalent in heterocyclic and organic cyclic compound binding genes, while splicing in the hypothalamus affects small molecule and drug binding loci. Counts for all terms in this figure were significantly

different from the expected value in at least one of the tissues. Parent ontology terms along the y-axis are in descending order from most frequent in the genome to less

415   frequent. The left panel depicts counts from hypothalamic spliced genes, and the right panel spliced genes from the pituitary. Blue represents control treatment, and orange is restraint stress. Abundances are *observed counts – expected* (based upon genomic predictions)/ *total events within that tissue*. We did not include any terms that were attributed to less than 2% of the genome.

Male vs. Female Splicing Comparison: Exon size, location, and motifs
To further characterize the splicing profile of the sexes, we visualized distributions of exon sizes, where these exons were located, and protein motifs contained therein. In terms of size distribution, our analyses revealed no significant difference between the

425   control (male vs female) and stresss (male vs female) comparisons in the hypothalamus or pituitary, suggesting that the size of alternatively spliced exons between males and females does not differ between control or stressed states (Figure 3). In all cases, spliced exons were smaller than predicted by the genomic distribution (Wilcoxon test p: hyp-control= 4.52e-12; hyp-stress= 2.16e-3; pit-control= 2.75e-8; pit-stress= 3.53e-10)

430   (Figure 3). Splicing events occurred in a variety of protein motifs, but no particular motif was significantly more spliced than the others (Figure S3).

Size of Core Exon Splicing Events

Comparison
- Control: Male vs Female
- Stress: Male vs Female
- Female: Control vs Stress
- Male: Control vs Stress
- Genome

Exon Length (Log^{10})

Figure 3. Distributions of core exon splicing event lengths for between-sex spliced loci in the control (light blue) and stress (orange) states as well as control-stress spliced loci in male (blue) and female (green) states. Neither the hypothalamus or pituitary showed significant difference between control (light blue) vs stress (orange) comparisons. In the pituitary of control vs stress comparison, lengths of spliced exons are significantly larger in the male pituitary than that of spliced exons in the female pituitary (Wilcoxon, p = 0.022). Distribution of genome exon sizes is colored in red. All comparisons with genome distribution, exon sizes were significantly smaller (Wilcoxon, p < 2.80e-3).

In both tissues, core exon splice sites occurred in protein coding sequences much more frequently than either of the untranslated regions (ChiSq p: hyp=5.95e-43, pit=6.52e-141). This, perhaps, is not surprising given that the CDS regions are more abundant in the genome and alteration to these regions will ultimately result in changes to the

protein sequence. The abundance of alternatively spliced exons present in the 5' & 3'

untranslated regions of the hypothalamus and pituitary was significantly different from

expected values (Figure 4). In the hypothalamus, more spliced exons occurred in the

450   5'UTR than genomic proportions would predict, with more dramatic shifts in the restraint

stress treatment than the control (ChiSq, p=0.015). In the pituitary, the control group

exhibited more spliced exons of the 5'UTR and both treatment groups presented more

than a 6% decrease of splicing events in 3'UTR regions than predicted from genomic

values (ChiSq p: 5'UTR=0.041, 3'UTR=2.60e-7).

Figure 4. Transcriptional regions of male-female splicing events. In both tissues, the 5'UTR region was a site of increased splicing activity, while the 3'UTR regions underwent less splicing. The upper half represents percentage of total splicing events (for each tissue) in each transcript region. There were significantly more CDS events than 5UTR or 3UTR (ChiSq p: hyp=5.95e-43, pit=6.52e-141). The lower portion depicts how these counts differ from the expected (i.e. a value of 0.5 here indicates a 50% deviation from genomic expectation). Blue represents control treatment, while orange is restraint stress. Red stars represent significance between observed counts and

465   expected from genomic predictions (ChiSq p: Hyp-5'UTR=0.015, Pit-5'UTR=0.041,Pit-

       3'UTR=2.60e-7).

       Control vs. Stress Splicing Comparison

       The second aim of this study was to observe splicing differences within each sex in

470   control and stress states, to identify how each sex individually responded to restraint

       stress. We compared, within each sex, control and stress treatments (female: control to

       stress; male: control to stress). We counted the number of alternative splicing events for

       these within-sex, across-treatment comparisons. Here, we include comparisons

       between gonads as the alternative splicing events identified are within-sex and thus, we

475   can observe how male and female gonads individually respond to restraint stress.

       ChiSq tests were used to determine statistical significance throughout our analyses; all

       p-values, degrees of freedom, and sample sizes are included in Table 1.

       Control vs. Stress Splicing Comparison: Events by Type

480   Similar to our comparison of male-female alternative splicing, control-stress splicing

       reveals more splicing in the pituitary than other tissues (ChiSq p=1.04e-6). The male

       hypothalamus is more active in stress-response splicing compared to the female

       hypothalamus (ChiSq p=9.39e-4), and the female gonad exhibits more activity than the

       male gonad (ChiSq p=1.92e-4). The male hypothalamus displays 59% more splicing

485   events than the female hypothalamus, and we identified 78% more splicing events in

       the ovaries than the testes (Figure 1).

Of all event types, the core exon event was most abundant in all tissues (ChiSq p: hyp=3.64e-7, pit=1.34e-43, gon=8.74e-15)(Figure 1). Of these core exon events, the

490   gonads were the only tissue to present statistical significance across sexes; more core exon splice events occurred in the ovaries than in the testes (ChiSq p=3.01e-3). This parallels our previous findings of elevated female gonadal gene expression in response to restraint stress (Calisi et al., 2018). The only other event that differed significantly between the sexes was hypothalamic retained intron events, with more found in males

495   than females (ChiSq p=0.041).

Control vs. Stress Splicing Comparison: Genes of Interest

Through assessment of alternative splicing events between control and stress states, we were able to identify genes spliced in response to stress within each sex. Estradiol

500   17-beta-dehydrogenase 11 (HSD17B11) was alternatively spliced between treatments in the male hypothalamus. HSD17B11 plays a role in hormone metabolism, through which it may mediate endogenous estrogen levels (Ariazi et al., 2011). Through feedback on the brain, estrogens control the pulsatile release of GnRH and can influence stress signaling (Acevedo-Rodriguez et al., 2018) and enhance HPA function

505   (Handa et al., 1994). HSD17B11 is also related to CEBPB (CCAAT/enhancer-binding protein beta), a transcription factor regulating the expression of genes involved in immune and inflammatory responses (Chinery et al., 1997; Pless et al., 2008). The splicing of this gene, and others in our list, suggest a sex-specific response to stress mediated via alternative splicing.

As in our male-female splicing analysis, we identified few genes consistently

alternatively spliced; however, the genes that were consistently spliced in both sexes

were quite interesting. Rho family GTPases function as molecular switches to regulate a

variety of cellular processes including cytoskeletal growth, gene transcription, and

515   cytokinesis (Moon & Zheng, 2003; Miller & Bement, 2009). Both males and females

displayed alternative splicing of the pituitary gene Rho GTPase Activating Protein 32

(ARHGAP32). Through response to GnRH pulsatile expression, Rho GTPases

regulates cell motility and cytoskeletal rearrangements(Naor, Benard & Seger, 2000;

Godoy, Nishimura & Webster, 2011). The shift from a control to stressed state likely

520   places different constraints on cellular motility, requiring expedited movement of

signaling molecules and energy transport throughout an organism; consistent splicing of

ARHGAP32 may be one means by which cellular movement is adjusted during the

stress response.

525   The hypothalamus of both males and females exhibit alternative splicing of the Retinitis

Pigmentosa GTPase Regulator Interacting Protein 1 Like (RPGRIP1L) gene.

RPGRIP1L is an adapter protein, localized at the ciliary transition zone, that interacts

with G-protein coupled thromboxane A2 receptor (TBXA2R) to negatively regulate

signaling at the ciliary base (Tokue, Sasaki & Nakahata, 2009; Gerhardt et al., 2015).

530   This ciliary protein governs both autophagy and proteasome activity (Gerhardt et al.,

2015; Struchtrup et al., 2018). As both sexes respond to stress by alternatively splicing

the RPGRIP1L gene, the alteration of this locus may enable organisms to respond more

effectively to a stressful environment through signal transduction and

processing/disposal of damaged cells. Additional observation of the functioning of these

535   gene products in control and stress states will provide further understanding as to the

role these loci play in stress response.

Control vs. Stress Splicing Comparison: Gene Ontology

In addition to characterizing splicing on a gene-by-gene basis, we assessed the gene

540   lists, as a whole, to determine if a particular type of gene or genes related to a specific

biological process were more readily spliced in the stress response. Our ontology

analysis of the stress vs. control splicing events revealed elevated levels of splicing in

genes with the following GO terms: small molecule binding, carb derivative binding, and

drug binding (Figure 5). These terms were more abundant than genomic predictions in

545   the male hypothalamus, female pituitary, and male gonad, while these same terms were

less abundant than expected in the female hypothalamus and female gonad. Of the

male tissues, the hypothalamus and testes appear more active in this area. There also

were more spliced genes related to drug binding than expected.

Figure 5. Molecular Function GO Analysis, Stress vs. Control (normalized counts of observed-expected). Binding genes were more frequently spliced in the male hypothalamus and the female pituitary. Counts for all terms in this figure were significantly different from the expected value in at least one of the tissues. Parent ontology terms along the y axis are in descending order from most frequent in the genome to least frequent. The three "panels" depict counts from the hypothalamus, pituitary, and gonads (respectively, from left to right). Abundances are observed counts – expected (based upon genomic predictions)/ total events within that tissue. We did not include any terms that were attributed to less than 2% of the genome.

Control vs. Stress Splicing Comparison: Exon size, location, and motifs
Analysis of distributions of alternatively spliced exon sizes reveal that all tissues, regardless of sex, exhibit distributions of control-stress splicing events smaller than the genomic distribution (Wilcoxon, all p < 2.80e-3)(Figure 3). The pituitary was the only tissue in which male and female distributions differed, with the male pituitary exons being longer than female (Wilcoxon, p = 0.022). Similar to the male-female splicing analysis of splicing in protein motifs, we found no significant difference in the distribution of events occurring in any one motif (Figure S6).

In analyzing locations of spliced exons, we again identified more events in coding regions than either 5' or 3' UTR, for all tissues (ChiSq p: hyp=4.41e-40, pit=9.60e-233, gon=5.13e-67)(Figure 6). Similar to our male-female analysis, we also identified a significantly lower abundance of splicing events in the 3'UTR regions of all tissues in the

control-stress analysis (ChiSq p: hyp=0.042, pit=2.87e-12, gon=8.77e-4). There were

575 fewer events in the 5'UTR of the pituitary (ChiSq p=2.04e-4), and more events occurred

in the pituitary CDS than expected (ChiSq p=3.50e-4).

Figure 6. Transcriptional regions of control-stress splicing events.

580 There were significantly fewer splicing events in the 3'UTR, agreeing with our previous

findings, indicating that splicing machinery may selectively avoid this transcript region.

The upper half represents percentage of total splicing events (for each tissue) in each

transcript region. There were significantly more CDS events than 5'UTR or 3'UTR

(ChiSq p: hyp=9.77e-23, pit=9.60e-233, gon=5.13e-67). The lower portion depicts how

585   these counts differ from the expected (i.e. a value of 0.5 here indicates a 50% deviation

from genomic expectation). Blue represents male, while green is female. Red stars

represent significance between observed and expected proportions of splice location

(ChiSq p: hyp-3'UTR=0.042, pit-CDS=3.50e-4, pit-5'UTR=2.04e-4, pit-3'UTR=2.87e-12,

gon-3'UTR=8.77e-4).

Treatment-specific and Sex-specific Isoforms

Because we know that treatment and sex affect gene expression documented (Calisi et

al., 2018), we assessed each tissue for treatment-specific, and sex-specific isoforms of

a gene. We did not identify any treatment-, nor sex-specific isoforms. This result

595   suggests that even though gene expression may shift, and splicing events can be

specific to an external stimulus, specific isoforms are not exclusive to one sex or

treatment. A potential shortcoming of this particular analysis is the heavy reliance upon

annotation. As the *C. livia* genome continues to improve with the progression of

technology and sequence analysis tools we encourage others to reassess this work.

Bimodal distribution of PSI

We assessed the distributions of PSI values for each splicing event type across all

analysis comparisons to determine if there existed a differential in the frequency that a

splicing event was present within a given state. The measure of "percent spliced in"

605   reflects the frequency that one splicing isoform was seen over another in an individual.

Ultimately, we find that PSI values do not show dramatic deviation for splicing events

between treatments, nor between sexes (Figure S7). The only two instances that there

appears to be a binary shift is in the case of alternate acceptors (AA), specifically in the

hypothalamus of the male vs female analysis, and in the hypothalamus of the control vs

610   stress analysis. This suggests that sex is not a driving factor for PSI values of stress-

response alternative splice sites.

We would like to note that the algorithm used to identify significant splice events

implicitly biases our results toward a bimodal distribution as any event with a deltaPSI

615   between -0.1 and 0.1 was discounted. To determine how dramatically this shifted our

distributions, we temporarily called all splicing events with $p \leq 0.05$ significant. The

additional 9 events this included for the male-female comparison were all found in the

pituitary (3 control, 6 stress). Sixty-one events in the control-stress comparison had

deltaPSI values between -0.1 and 0.1, 58 of which were also in the pituitary (53 male, 5

620   female) and 3 in the female gonad. Most significant splicing events have abs(deltaPSI)

values greater than 0.1 in the male-female comparison. This is perhaps not particularly

surprising as splicing differences are likely more biologically effective when included or

excluded at higher PSI levels due to a more dramatic shift in protein presence for a

given isoform. The 53 events in the male pituitary with low abs(deltaPSI) may suggest

625   the male pituitary responds to stress not only by splicing genic regions at higher PSI

values, but also by numerous, low-level splicing events. Further analyses using

additional splicing tools and datasets are needed to validate this hypothesis.

DISCUSSION

To survive an unanticipated environmental perturbation, or stressor, the body activates multiple physiological systems to mobilize needed resources, with the intent of returning the body back to a homeostatic level (Michael Romero & Wingfield, 2015). The stress response is considered relatively conserved across vertebrates (Clarke et al., 1994;

635 Tsigos & Chrousos, 2002; Leonard, 2005; Gutierrez-Triana et al., 2014; Taniuchi et al., 2016) and includes the limbic system, the central nervous system, the stress or hypothalamic-pituitary-adrenal (HPA) axis, and the thyroid or hypothalamic-pituitary-thyroid axis (HPT), all of which can directly or indirectly influence the HPG axis (Tsigos & Chrousos, 2002; Helmreich et al., 2005; Herman et al., 2005; Romero-Ramírez,

640 Nieto-Sampedro & Barreda-Manso, 2017). Selection should favor an optimized stress response in which an organism's physiological and behavioral responsiveness to perturbation promotes its survival and, ultimately, reproduction. However, we know little of how stress can affect the reproductive system at the genomic level of alternative splicing, and even less so about the similarities and differences experienced by males

645 and females. Using the rock dove model in a highly-replicated RNA-Seq study, we tested how an external environmental stressor could affect the alternative splicing response of the HPG axis in both males and females. We discovered a significant sex difference in the splicing profile of hypothalamic and pituitary tissues. When subjects underwent a restraint stress treatment, we observed a decrease in the total number of

650 significant splicing events between sexes as compared to controls (Figure 1). Two

potential explanations for this are (1) the "convergent landscape hypothesis": when males and females are stressed, they converge on a single splicing landscape, or (2) the "resource redirection hypothesis": the stress response shuts down non-essential biological processes, minimizing background splicing "noise" that differentiates

655   unstressed males and females, to redirect resources to bodily processes critical for survival.

Convergent Landscape Hypothesis

The decrease in alternative splicing events experienced by stressed males and females

660   as compared to their unstressed controls may suggest that both sexes are converging on an optimal splicing landscape. Previous reports show differential gene expression (DGE) between sexes in both an unstressed control state (Shapiro et al., 2013; Gillespie et al., 2013; MacManes et al., 2017), and in response to a stressor (Shapiro et al., 2013; Gillespie et al., 2013; MacManes et al., 2017; Calisi et al., 2018). Given these

665   previous findings, coupled with ours presented here, we suggest that future study of the HPG stress response should endeavor to further characterize sex-specific and sex-convergent responses at all levels - genomic, proteomic, and methylomic. If other response mechanisms between males and females showed a similar pattern where the sexes exhibit fewer differences in a stress state, it may suggest that there was an

670   optimal stress response phenotype being maintained at numerous levels, in addition to genomic ones currently understood. If this pattern of convergence is not seen at other physiological levels, it is possible that the decrease in splicing differences between males and females in the restraint stress treatment may reflect a stress response

unique to the splicing level that does not exist in the proteomic or methylomic levels or

675   additional variables unassessed in this study (e.g., stage of reproductive cycle).

Resource Redirection Hypothesis

A second hypothesis to explain the presence of decreased alternative splicing

differences between stressed males and females relative to controls is that the stress

680   response redirects resources from less essential biological processes, like reproduction,

to those more necessary for survival. Biological sex manifests in various distinct

physiological phenotypes, particularly concerning those associated with reproductive

processes (Ngun et al., 2011; Arnold & Itoh, 2011; Bundy, Vied & Nowakowski, 2017;

Lee, 2018)However, an organism's response to a stressor, no matter the sex, can

685   trigger the mobilization of internal bodily resources to promote its survival (Wendelaar

Bonga, 1997; Goldstein, 2010; Nicolaides et al., 2015; Smulders, 2017). Thus, an active

sex-specific physiologically-driven alternative splicing landscape under stress may

become reconciled in the HPG axis to shunt focus to other biological processes to

support survival. This reduction in sexually dimorphic splicing background noise would

690   result in fewer splicing differences between the sexes, as we observed. The remaining

splicing differences may lie in loci generally implicated in the stress response, though

future studies are needed to say this definitively. While our attempts to map spliced

genes to Kegg pathways did not yield a discernable pattern of splicing of a particular

pathway, we did find that several of the alternatively spliced genes may be involved in

695   DNA-damage response and immune system processes, both of which are intimately

linked with the stress response (Hara et al., 2011; Dhabhar, 2014). Additional study of

the splicing landscape of males and females in various conditions, environments, age

groups, and reproductive stages will shed further light on splicing variation between the

sexes, and if our results seen are indeed a reflection of decreased activity of peripheral

700   biological processes during stress.

Genomic Events by Type

As compared to the hypothalamus and gonads, the pituitary in both sexes exhibited the

most active splicing profile under stress as compared to controls. This pattern was

705   consistent with our previous report of increased differential gene expression (DGE) in

the pituitary following restraint stress (Calisi et al., 2018), suggesting this gland may be

a targeted site for mediating a stress response in the HPG axis through both gene

expression and alternative splicing mechanisms. Regarding overall splicing activity,

males exhibited greater responsiveness to stress than females in the hypothalamus and

710   pituitary, with a significant increase in response of the male hypothalamus compared to

that of the female. Contrarily, examination of the same samples taken from the same

birds showed that females exhibited greater DGE responsiveness to stress than males

in all three tissues (Calisi et al., 2018). Thus, while examination at a gene transcription

level may suggest the female reproductive axis as more responsive to the restraint

715   stress treatment, examination at the alternative splicing level suggests males to be the

more responsive of the sexes. The next steps would be to examine the protein products

produced, but for now suffice it to say caution should be taken when determining which

sex is more "responsive" to any stimuli, as this can vary by biological level of

examination.

720   Further investigation of the types of splicing events that occurred following

restraint stress revealed that core exon events as compared to all other event types

were vastly more abundant across all tissues, regardless of sex or treatment, a

phenomenon supported by findings in humans and mice (Koralewski & Krutovsky,

2011). Regarding the effects of stress treatment, more core exon events were present

725   in the female gonads as compared to the male gonads. The higher abundance of core

exon events could indicate that exon inclusion/exclusion is a more efficient approach to

modulate a resulting protein, thereby triggering a more specific phenotypic response to

a change in environment (Chen & Manley, 2009; Rodrigues, Grosso & Moita, 2013;

Wang et al., 2015; Yabas, Elliott & Hoyne, 2015). This may then signify that the stress

730   response of the female gonad is driven more heavily by core exon alterations to a

transcript, while transcripts in the male gonad are more readily modified with all types of

splicing events.

The only other splicing event type that differed significantly between the stress

response of males and females was the retention of introns, with more retained intron

735   events occurring in the male hypothalamus. Retained intron events are a means by

which gene expression can be modified, often by downregulation (Ge & Porse, 2014).

However, these events may also alter *cis* and *trans* elements that would otherwise

modify transcript stability or translational efficacy (Jacob & Smith, 2017). Perhaps

through differential utilization of the retained intron splicing event, the male

740   hypothalamus could further modulate gene expression, targeting a more optimal stress

response.

Gene Ontology

In an attempt to gain a better understanding of potential products being generated by
745   splicing activity, we examined our results using Gene Ontology (GO). Our inquiry of
genes undergoing sex-biased alternative splicing revealed many GO terms related to
"binding." The function of these genes is linked with how hormones bind to their
receptors, and by extension, how endocrine messages are received. With the increase
in abundance of spliced hypothalamic binding genes and a decrease of "binding" terms
750   in the pituitary spliced genes, it appears that males and females may differ in how cells
in the hypothalamus bind various protein products. While the function of genes in the
male and female pituitary related to "binding" exhibit fewer alternative splicing events
than expected by genomic prediction, the sexes showed greater difference in alternative
splicing of genes associated with "cell/neuronal structure". This may suggest that similar
755   functions can arise from sex-specific structures.

Alternative splicing exhibited in stressed versus control treatment subject HPG
tissues also reveal sexual-dimorphic functionality. Once again, many of the alternatively
spliced gene products within each sex in response to stress are related to "binding".
Specifically, we observed elevated splicing of these genes in the male hypothalamus,
760   female pituitary, and male gonad. This pattern seems to suggest that the female
pituitary is responding to restraint stress by altering how molecules/drugs are bound,
while the male hypothalamus is more active in responding to stress via splicing binding
genes. Thus, sexual differences in splicing landscapes in our samples may signify sex-
biased hypothalamic function in response to stress, but similar pituitary functions
765   between the sexes.

Exon size, location, and motifs

The exon size distribution of alternative splicing events was significantly smaller than

that predicted by the genomic distribution, agreeing with previous findings that the

770   likelihood of a core exon being spliced may be inversely related to its size (Song et al.,

2009). This inverse relationship of exon size and frequency of splicing may be a

reflection of the highly-specific and precise adjustments required of splicing machinery

(Black, 2000; Birzele, Csaba & Zimmer, 2008; Mittendorf et al., 2012). Inclusion or

excision of a larger genetic sequence could prove more biologically challenging due to

775   the physical distance between the two splice sites, and variable inclusion of a larger

exon may result in a more dramatic alteration to the protein product, negating the effect

of fine-tuning a response. Though the alternative splicing detected in both sexes

exhibited smaller than expected exon sizes, size distributions in the female pituitary

were smaller than those of the male pituitary. This may suggest that splicing in the

780   female pituitary is targeted towards minute alteration, while genes in the male pituitary

are modulated by movement/removal of larger sequence segments. Additional study of

splicing in pituitaries of both sexes will be required to clarify the biological implications of

this sex-bias.

785   In all comparisons, we found that the vast majority of splicing occurs in the protein

coding region, or coding sequence (CDS) of a gene. When we examine the frequency

of splicing location in relation to the genomic expectation (assuming an even distribution

of splicing across all genic regions), we see that variation from expectation actually lies

in the untranslated regions. Analyses comparing males and females, within each

790   treatment group, revealed increased splicing of 5'-UTR genes in both the hypothalamus

and pituitary, and a >6% decrease in splicing of the 3'-UTR genes in the pituitary.

Splicing differences identified within the HPG axis of each sex in stressed birds as

compared to controls displayed lower 3'UTR events in all tissues, and lower 5'UTR

events in the pituitary. The increased abundance of 5'UTR alternative splicing events in

795   males as compared to female subjects suggests that differences in the 5'UTR region of

transcripts exist between the sexes (regardless of control/stress state). Previous

research has found that the 5'UTR can mediate translation of proteins, regulate

expression, and even regulate some splicing events (Kramer et al., 2013; Álvarez et al.,

2016; Ohta et al., 2018). Thus, differential splicing of the 5'UTR between males and

800   females may be a sex-specific mechanism for modifying protein production. The one

instance in which we see a decrease in splicing of 5'UTR events is the pituitary, where

each sex exhibits fewer splicing events in response to stress. Males and females may

differ in the 5'UTR, but both sexes seem to minimize splicing of this region in the

pituitary in response to restraint stress. The decrease in splicing events in the 3'UTR

805   could signify that splicing events in the HPG axis do not favor the 3'UTR in response to

restraint stress. 3' ends of transcripts may be protected from alternative splicing, as

evidenced by this decrease observed in both sexes and treatments. Degradation of

mRNA transcripts can increase due to environmental stress (Canadell et al., 2015), and

it may be that splicing events in the 3'UTR would result in undesired degradation of

810   transcripts, or perhaps no degradation at all.

We did not identify any differences in the frequency of protein motif splicing between the sexes or treatments. As motifs combinations and conformations heavily impact protein function, we anticipated identifying specific motifs wherein splicing more readily

815   occurred. Our query did not reveal this, suggesting that splicing is not targeting protein motifs within the HPG axis in either sex or treatment. Instead, the genic region (CDS, 5'UTR, 3'UTR) and gene-type appear to be larger drivers of the location that splicing events occur.

820   **SUMMARY**

In both sexes and treatment groups, we identified the most alternative splicing activity in the pituitary as compared to the hypothalamus and gonads. Our previous research also revealed the pituitary to have an increased response to restraint stress at the level of

825   gene expression as compared to the hypothalamus or gonads (Calisi et al., 2018), suggesting a heightened stress response in this part of the HPG axis. Of genomic splicing events identified, core exon events are the most abundant form of splicing in all tissues/sexes/treatments, and splicing more frequently occurs in CDS regions. We found less splicing in the 3'UTR and more splicing in the 5'UTR than expected in both

830   sex and treatment comparisons, suggesting that the 3' end of transcripts is more biologically constrained than the 5' end in our samples. Another splicing constraint evidenced by our findings, is that of an inverse relationship of exon size and splicing frequency. The overall reduction in splicing differences between the sexes when animals experienced a restraint stress treatment may point to a conserved splicing

835   response to stress within the HPG axis. However, despite this reduction in sex-splicing

differences, the male hypothalamus and female ovary experienced increased splicing

activity in the face of stress as compared to the female hypothalamus and male testes,

respectively. Sex differences in alternative splicing within the HPG axis, as well as in

previously reported gene expression (Calisi et al., 2018) support sex-specific

840   mechanisms for the stress response of the reproductive axis. While females experience

increased differential gene expression in their HPG axis as compared to males in

response to restraint stress (Calisi et al., 2018), we found that males experience

increased alternative splicing. Our examination of the vertebrate stress response at

multiple levels of biological organization offers a more complete picture of its

845   mechanistic underpinnings between the sexes at an unprecedented proximate level.

These data inspire further integrative levels of investigation to inform and potentially

revolutionize evolutionary theory and the study of stress-induced reproductive

dysfunction.

DECLARATIONS

Ethics approval and consent to participate

860   Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Competing interests

The authors declare that they have no competing interests.

870   Funding

This work was funded by NSF IOS 1455960 (to RMC and MM).

Authors' contributions

RMC and MM designed the study. SA conducted avian manipulation and tissue

sampling. AL conducted RNA extraction, cDNA library preparation. AL conducted

875   computational analyses with input from RH. AL wrote the manuscript with input from all.

ACKNOWLEDGEMENTS

We thank the many undergraduate and graduate students of the MacManes and Calisi

Lab for their assistance on this project, including their involvement in avian husbandry,

conducting the experiment, tissue collection, RNA extraction, and cDNA library

880   preparation.

REFERENCES

Acevedo-Rodriguez A, Kauffman AS, Cherrington BD, Borges CS, Roepke TA, Laconi M. 2018. Emerging insights into hypothalamic-pituitary-gonadal axis regulation and interaction with stress signalling. *Journal of neuroendocrinology* 30:e12590.

Álvarez D, Voß B, Maass D, Wüst F, Schaub P, Beyer P, Welsch R. 2016.

905 Carotenogenesis Is Regulated by 5'UTR-Mediated Translation of Phytoene Synthase Splice Variants. *Plant physiology* 172:2314–2326.

Ariazi EA, Cunliffe HE, Lewis-Wambi JS, Slifker MJ, Willis AL, Ramos P, Tapia C, Kim HR, Yerrum S, Sharma CGN, Nicolas E, Balagurunathan Y, Ross EA, Jordan VC. 2011. Estrogen induces apoptosis in estrogen deprivation-resistant breast cancer

910 through stress responses as identified by global gene expression across time. *Proceedings of the National Academy of Sciences of the United States of America* 108:18879–18886.

Arnold AP, Itoh Y. 2011. Factors causing sex differences in birds. *Avian biology research* 4. DOI: 10.3184/175815511X13070045977959.

915 Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nature genetics* 25:25–29.

920 Ball GF, Silver R. 1983. Timing of incubation bouts by ring doves (Streptopelia risoria). *Journal of comparative psychology* 97:213–225.

Birzele F, Csaba G, Zimmer R. 2008. Alternative splicing and protein structure evolution. *Nucleic acids research* 36:550–558.

Black DL. 2000. Protein Diversity from Alternative Splicing: A Challenge for

925   Bioinformatics and Post-Genome Biology. *Cell* 103:367–370.

Bundy JL, Vied C, Nowakowski RS. 2017. Sex differences in the molecular signature of the developing mouse hippocampus. *BMC genomics* 18:237.

Buntin JD, Ruzycki E, Witebsky J. 1993. Prolactin receptors in dove brain: autoradiographic analysis of binding characteristics in discrete brain regions and

930   accessibility to blood-borne prolactin. *Neuroendocrinology* 57:738–750.

Bush SJ, Chen L, Tovar-Corona JM, Urrutia AO. 2017. Alternative splicing and the evolution of phenotypic novelty. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 372. DOI: 10.1098/rstb.2015.0474.

Calisi RM, Austin SH, Lang AS, MacManes MD. 2018. Sex-biased transcriptomic

935   response of the reproductive axis to stress. *Hormones and behavior* 100:56–68.

Campbell BK, Dobson H, Scaramuzzi RJ. 1998. Ovarian function in ewes made hypogonadal with GnRH antagonist and stimulated with FSH in the presence or absence of low amplitude LH pulses. *The Journal of endocrinology* 156:213–222.

Canadell D, García-Martínez J, Alepuz P, Pérez-Ortín JE, Ariño J. 2015. Impact of high

940   pH stress on yeast gene expression: A comprehensive analysis of mRNA turnover during stress responses. *Biochimica et biophysica acta* 1849:653–664.

Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, Holt C, Sánchez Alvarado A, Yandell M. 2008. MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. *Genome research* 18:188–196.

945   Chandran UR, DeFranco DB. 1999. Regulation of gonadotropin-releasing hormone
gene transcription. *Behavioural brain research* 105:29–36.

Cheng CK, Yeung CM, Hoo RLC, Chow BKC, Leung PCK. 2002. Oct-1 is involved in
the transcriptional repression of the gonadotropin-releasing hormone receptor
gene. *Endocrinology* 143:4693–4701.

950   Chen M, Manley JL. 2009. Mechanisms of alternative splicing regulation: insights from
molecular and genomics approaches. *Nature reviews. Molecular cell biology*
10:741–754.

Chen J, Weiss WA. 2015. Alternative splicing in cancer: implications for biology and
therapy. *Oncogene* 34:1–14.

955   Chinery R, Brockman JA, Dransfield DT, Coffey RJ. 1997. Antioxidant-induced Nuclear
Translocation of CCAAT/Enhancer-binding Protein β: A CRITICAL ROLE FOR
PROTEIN KINASE A-MEDIATED PHOSPHORYLATION OF Ser299. *The Journal
of biological chemistry* 272:30356–30361.

Clarke AS, Wittwer DJ, Abbott DH, Schneider ML. 1994. Long-term effects of prenatal
960   stress on HPA axis activity in juvenile rhesus monkeys. *Developmental
psychobiology* 27:257–269.

Climente-Gonzalez H, Porta-Pardo E, Godzik A, Eyras E. 2017. The functional impact
of alternative splicing in cancer. *bioRxiv*:076653. DOI: 10.1101/076653.

Dhabhar FS. 2014. Effects of stress on immune function: the good, the bad, and the
965   beautiful. *Immunologic research* 58:193–210.

Domyan ET, Kronenberg Z, Infante CR, Vickrey AI, Stringham SA, Bruders R,
Guernsey MW, Park S, Payne J, Beckstead RB, Kardon G, Menke DB, Yandell M,

Shapiro MD. 2016. Molecular shifts in limb identity underlie development of feathered feet in two domestic avian species. *eLife* 5:e12115.

970 Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A. 2016. The Pfam protein families database: towards a more sustainable future. *Nucleic acids research* 44:D279–85.

Ge Y, Porse BT. 2014. The functional consequences of intron retention: alternative
975 splicing coupled to NMD as a regulator of gene expression. *BioEssays: news and reviews in molecular, cellular and developmental biology* 36:236–243.

Geraghty AC, Kaufer D. 2015. Glucocorticoid Regulation of Reproduction. *Advances in experimental medicine and biology* 872:253–278.

Gerhardt C, Lier JM, Burmühl S, Struchtrup A, Deutschmann K, Vetter M, Leu T, Reeg
980 S, Grune T, Rüther U. 2015. The transition zone protein Rpgrip1l regulates proteasomal activity at the primary cilium. *The Journal of cell biology* 210:115–133.

Gibilisco L, Zhou Q, Mahajan S, Bachtrog D. 2016. Alternative Splicing within and between Drosophila Species, Sexes, Tissues, and Developmental Stages. *PLoS genetics* 12:e1006464.

985 Gillespie MJ, Crowley TM, Haring VR, Wilson SL, Harper JA, Payne JS, Green D, Monaghan P, Stanley D, Donald JA, Nicholas KR, Moore RJ. 2013. Transcriptome analysis of pigeon milk production - role of cornification and triglyceride synthesis genes. *BMC genomics* 14:169.

Godoy J, Nishimura M, Webster NJG. 2011. Gonadotropin-releasing hormone induces
990 miR-132 and miR-212 to regulate cellular morphology and migration in

immortalized LbetaT2 pituitary gonadotrope cells. *Molecular endocrinology* 25:810–820.

Goldstein DS. 2010. Adrenal responses to stress. *Cellular and molecular neurobiology* 30:1433–1440.

995 González-Tajuelo R, Silván J, Pérez-Frías A, de la Fuente-Fernández M, Tejedor R, Espartero-Santos M, Vicente-Rabaneda E, Juarranz Á, Muñoz-Calleja C, Castañeda S, Gamallo C, Urzainqui A. 2017. P-Selectin preserves immune tolerance in mice and is reduced in human cutaneous lupus. *Scientific reports* 7:41841.

1000 Grosso AR, Gomes AQ, Barbosa-Morais NL, Caldeira S, Thorne NP, Grech G, von Lindern M, Carmo-Fonseca M. 2008. Tissue-specific splicing factor gene expression signatures. *Nucleic acids research* 36:4823–4832.

Gudlaugsdottir S, Boswell DR, Wood GR, Ma J. 2007. Exon size distribution and the origin of introns. *Genetica* 131:299–306.

1005 Gueroussov S, Gonatopoulos-Pournatzis T, Irimia M, Raj B, Lin Z-Y, Gingras A-C, Blencowe BJ. 2015. An alternative splicing event amplifies evolutionary differences between vertebrates. *Science* 349:868–873.

Gutierrez-Triana JA, Herget U, Lichtner P, Castillo-Ramírez LA, Ryu S. 2014. A vertebrate-conserved cis-regulatory module for targeted expression in the main

1010 hypothalamic regulatory region for the stress response. *BMC developmental biology* 14:41.

Handa RJ, Burgess LH, Kerr JE, O'Keefe JA. 1994. Gonadal steroid hormone receptors and sex differences in the hypothalamo-pituitary-adrenal axis. *Hormones and*

behavior 28:464–476.

1015   Hara MR, Kovacs JJ, Whalen EJ, Rajagopal S, Strachan RT, Grant W, Towers AJ,
        Williams B, Lam CM, Xiao K, Shenoy SK, Gregory SG, Ahn S, Duckett DR,
        Lefkowitz RJ. 2011. A stress response pathway regulates DNA damage through
        β2-adrenoreceptors and β-arrestin-1. *Nature* 477:349–353.

        Helmreich DL, Parfitt DB, Lu X-Y, Akil H, Watson SJ. 2005. Relation between the
1020   hypothalamic-pituitary-thyroid (HPT) axis and the hypothalamic-pituitary-adrenal
        (HPA) axis during repeated stress. *Neuroendocrinology* 81:183–192.

        Herman JP, Ostrander MM, Mueller NK, Figueiredo H. 2005. Limbic system
        mechanisms of stress regulation: hypothalamo-pituitary-adrenocortical axis.
        Progress in neuro-psychopharmacology & biological psychiatry 29:1201–1213.

1025   Holway DA, Suarez AV. 1999. Animal behavior: an essential component of invasion
        biology. *Trends in ecology & evolution* 14:328–330.

        Izquierdo J-M, Valcárcel J. 2006. A simple principle to explain the evolution of pre-
        mRNA splicing. *Genes & development* 20:1679–1684.

        Jacob AG, Smith CWJ. 2017. Intron retention as a component of regulated gene
1030   expression programs. *Human genetics* 136:1043–1057.

        Jin J-M, Yang W-X. 2014. Molecular regulation of hypothalamus-pituitary-gonads axis in
        males. *Gene* 551:15–25.

        Johnson JE, Kalmar GB, Sohal PS, Walkey CJ, Yamashita S, Cornell RB. 1992.
        Comparison of the lipid regulation of yeast and rat CTP: phosphocholine
1035   cytidylyltransferase expressed in COS cells. *Biochemical Journal* 285 (Pt 3):815–
        820.

Katagiri K, Hattori M, Minato N, Kinashi T. 2002. Rap1 functions as a key regulator of T-cell and antigen-presenting cell interactions and modulates T-cell responses. *Molecular and cellular biology* 22:1001–1015.

1040 Kim E, Magen A, Ast G. 2007. Different levels of alternative splicing among eukaryotes. *Nucleic acids research* 35:125–131.

Koralewski TE, Krutovsky KV. 2011. Evolution of exon-intron structure and alternative splicing. *PloS one* 6:e18055.

Kramer M, Sponholz C, Slaba M, Wissuwa B, Claus RA, Menzel U, Huse K, Platzer M,

1045 Bauer M. 2013. Alternative 5' untranslated regions are involved in expression regulation of human heme oxygenase-1. *PloS one* 8:e77224.

Krawczak M, Thomas NST, Hundrieser B, Mort M, Wittig M, Hampe J, Cooper DN. 2007. Single base-pair substitutions in exon-intron junctions of human genes: nature, distribution, and consequences for mRNA splicing. *Human mutation*

1050 28:150–158.

Laloum T, Martín G, Duque P. 2018. Alternative Splicing Control of Abiotic Stress Responses. *Trends in plant science* 23:140–150.

Lee SK. 2018. Sex as an important biological variable in biomedical research. *BMB reports* 51:167–173.

1055 Lehrman DS. 1955. The Physiological Basis of Parental Feeding Behavior in the Ring Dove (Streptopelia Risoria). *Behaviour* 7:241–285.

Leonard BE. 2005. The HPA and immune axes in stress: The involvement of the serotonergic system. *European psychiatry: the journal of the Association of European Psychiatrists* 20:S302–S306.

1060   MacManes MD, Austin SH, Lang AS, Booth A, Farrar V, Calisi RM. 2017. Widespread patterns of sexually dimorphic gene expression in an avian hypothalamic-pituitary-gonadal (HPG) axis. *Scientific reports* 7:45125.

Matsumoto M, Miyasaka M, Hirata T. 2009. P-selectin glycoprotein ligand-1 negatively regulates T-cell immune responses. *Journal of immunology* 183:7204–7211.

1065   McIntyre LM, Bono LM, Genissel A, Westerman R, Junk D, Telonis-Scott M, Harshman L, Wayne ML, Kopp A, Nuzhdin SV. 2006. Sex-specific expression of alternative transcripts in Drosophila. *Genome biology* 7:R79.

Michael Romero L, Wingfield JC. 2015. *Tempests, Poxes, Predators, and People: Stress in Wild Animals and how They Cope*. Oxford University Press.

1070   Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D, Thomas PD. 2017. PANTHER version 11: expanded annotation data from Gene Ontology and Reactome pathways, and data analysis tool enhancements. *Nucleic acids research* 45:D183–D189.

Miller AL, Bement WM. 2009. Regulation of cytokinesis by Rho GTPase flux. *Nature cell*
1075   *biology* 11:71–77.

Mittendorf KF, Deatherage CL, Ohi MD, Sanders CR. 2012. Tailoring of membrane proteins by alternative splicing of pre-mRNA. *Biochemistry* 51:5541–5556.

Moon SY, Zheng Y. 2003. Rho GTPase-activating proteins in cell regulation. *Trends in cell biology* 13:13–22.

1080   Naor Z, Benard O, Seger R. 2000. Activation of MAPK cascades by G-protein-coupled receptors: the case of gonadotropin-releasing hormone receptor. *Trends in endocrinology and metabolism: TEM* 11:91–99.

Nellore A, Paziana K, Ma C, Tsygankova OM, Wang Y, Puttaswamy K, Iqbal AU, Franks SR, Lv Y, Troxel AB, Feldman MD, Meinkoth JL, Brose MS. 2009. Loss of Rap1GAP in papillary thyroid cancer. *The Journal of clinical endocrinology and metabolism* 94:1026–1032.

Ngun TC, Ghahramani N, Sánchez FJ, Bocklandt S, Vilain E. 2011. The genetics of sex differences in brain and behavior. *Frontiers in neuroendocrinology* 32:227–246.

Nicolaides NC, Kyratzi E, Lamprokostopoulou A, Chrousos GP, Charmandari E. 2015. Stress, the stress system and the role of glucocorticoids. *Neuroimmunomodulation* 22:6–19.

Nijholt I, Farchi N, Kye M, Sklan EH, Shoham S, Verbeure B, Owen D, Hochner B, Spiess J, Soreq H, Blank T. 2004. Stress-induced alternative splicing of acetylcholinesterase results in enhanced fear memory and long-term potentiation. *Molecular psychiatry* 9:174–183.

Nozaki M. 2013. Hypothalamic-pituitary-gonadal endocrine system in the hagfish. *Frontiers in endocrinology* 4:200.

Ohama N, Sato H, Shinozaki K, Yamaguchi-Shinozaki K. 2017. Transcriptional Regulatory Network of Plant Heat Stress Response. *Trends in plant science* 22:53–65.

Ohta S, Nakagawara S, Hirai S, Miyagishima K, Horiguchi G, Kodama H. 2018. The 5′ UTR intron-mediated enhancement of constitutive splicing of the tobacco microsome ω-3 fatty acid desaturase gene. *Plant biotechnology reports* 12:105–114.

On the Origin of Species by Means of Natural Selection, or the Preservation of

Favoured Races in the Struggle for Life. 1860. *The British and foreign medico-chirurgical review* 25:367–404.

Pineda JMB, Bradley RK. 2018. Most human introns are recognized via multiple and tissue-specific branchpoints. *Genes & development* 32:577–591.

Pless O, Kowenz-Leutz E, Knoblich M, Lausen J, Beyermann M, Walsh MJ, Leutz A. 2008. G9a-mediated lysine methylation alters the function of CCAAT/enhancer-binding protein-beta. *The Journal of biological chemistry* 283:26357–26363.

Pohl CR, Richardson DW, Hutchison JS, Germak JA, Knobil E. 1983. Hypophysiotropic signal frequency and the functioning of the pituitary-ovarian system in the rhesus monkey. *Endocrinology* 112:2076–2080.

Preußner M, Goldammer G, Neumann A, Haltenhof T, Rautenstrauch P, Müller-McNicoll M, Heyd F. 2017. Body Temperature Cycles Control Rhythmic Alternative Splicing in Mammals. *Molecular cell* 67:433–446.e4.

Rodrigues R, Grosso AR, Moita L. 2013. Genome-wide analysis of alternative splicing during dendritic cell response to a bacterial challenge. *PloS one* 8:e61975.

Romero-Ramírez L, Nieto-Sampedro M, Barreda-Manso MA. 2017. Integrated Stress Response as a Therapeutic Target for CNS Injuries. *BioMed research international* 2017:6953156.

Rosen H, Jameel ML, Barkan AL. 1988. Dexamethasone suppresses gonadotropin-releasing hormone (GnRH) secretion and has direct pituitary effects in male rats: differential regulation of GnRH receptor and gonadotropin responses to GnRH. *Endocrinology* 122:2873–2880.

Saha A, Kim Y, Gewirtz ADH, Jo B, Gao C, McDowell IC, GTEx Consortium, Engelhardt

1130   BE, Battle A. 2017. Co-expression networks reveal the tissue-specific regulation of transcription and splicing. *Genome research* 27:1843–1858.

Sakharkar MK, Chow VTK, Kangueane P. 2004. Distributions of exons and introns in the human genome. *In silico biology* 4:387–393.

Sapolsky RM, Romero LM, Munck AU. 2000. How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative

1135   actions. *Endocrine reviews* 21:55–89.

Schneider F, Tomek W, Gründker C. 2006. Gonadotropin-releasing hormone (GnRH) and its natural analogues: a review. *Theriogenology* 66:691–709.

Shapiro MD, Kronenberg Z, Li C, Domyan ET, Pan H, Campbell M, Tan H, Huff CD, Hu H, Vickrey AI, Nielsen SCA, Stringham SA, Hu H, Willerslev E, Gilbert MTP,

1140   Yandell M, Zhang G, Wang J. 2013. Genomic diversity and evolution of the head crest in the rock pigeon. *Science* 339:1063–1067.

Siegfried Z, Karni R. 2018. The role of alternative splicing in cancer drug resistance. *Current opinion in genetics & development* 48:16–21.

Smith CW, Patton JG, Nadal-Ginard B. 1989. Alternative splicing in the control of gene

1145   expression. *Annual review of genetics* 23:527–577.

Smulders TV. 2017. The Avian Hippocampal Formation and the Stress Response. *Brain, behavior and evolution* 90:81–91.

Song L, Florea L. 2015. Rcorrector: efficient and accurate error correction for Illumina RNA-seq reads. *GigaScience* 4:48.

1150   Song S, Huang Q, Guo J, Li-Ling J, Chen X, Ma F. 2009. Comparative component analysis of exons with different splicing frequencies. *PloS one* 4:e5387.

Sower SA, Freamat M, Kavanaugh SI. 2009. The origins of the vertebrate hypothalamic-pituitary-gonadal (HPG) and hypothalamic-pituitary-thyroid (HPT) endocrine systems: new insights from lampreys. *General and comparative*

1155   *endocrinology* 161:20–29.

Staiger D, Brown JWS. 2013. Alternative splicing at the intersection of biological timing, development, and stress responses. *The Plant cell* 25:3640–3656.

Sterne-Weiler T, Weatheritt RJ, Best A, Ha KCH, Blencowe BJ. 2017. Whippet: an efficient method for the detection and quantification of alternative splicing reveals

1160   extensive transcriptomic complexity. *bioRxiv*:158519. DOI: 10.1101/158519.

Struchtrup A, Wiegering A, Stork B, Rüther U, Gerhardt C. 2018. The ciliary protein RPGRIP1L governs autophagy independently of its proteasome-regulating function at the ciliary base in mouse embryonic fibroblasts. *Autophagy* 14:567–583.

Sun H. 2017. Deciphering alternative splicing and nonsense-mediated decay modulate

1165   expression in primary lymphoid tissues of birds infected with avian pathogenic E. coli (APEC). *BMC genetics* 18:21.

Sutherland WJ. 1998. The importance of behavioural studies in conservation biology. *Animal behaviour* 56:801–809.

Takahashi A, Kanda S, Abe T, Oka Y. 2016. Evolution of the Hypothalamic-Pituitary-

1170   Gonadal Axis Regulation in Vertebrates Revealed by Knockout Medaka. *Endocrinology* 157:3994–4002.

Taniuchi S, Miyake M, Tsugawa K, Oyadomari M, Oyadomari S. 2016. Integrated stress response of vertebrates is regulated by four eIF2α kinases. *Scientific reports* 6:32886.

1175 Tapial J, Ha KCH, Sterne-Weiler T, Gohr A, Braunschweig U, Hermoso-Pulido A, Quesnel-Vallières M, Permanyer J, Sodaei R, Marquez Y, Cozzuto L, Wang X, Gómez-Velázquez M, Rayon T, Manzanares M, Ponomarenko J, Blencowe BJ, Irimia M. 2017. An atlas of alternative splicing profiles and functional associations reveals new regulatory programs and genes that simultaneously express multiple
1180 major isoforms. *Genome research* 27:1759–1768.

Telonis-Scott M, Kopp A, Wayne ML, Nuzhdin SV, McIntyre LM. 2009. Sex-specific splicing in Drosophila: widespread occurrence, tissue specificity and evolutionary conservation. *Genetics* 181:421–434.

The Gene Ontology Consortium. 2019. The Gene Ontology Resource: 20 years and still
1185 GOing strong. *Nucleic acids research* 47:D330–D338.

Tian B, Manley JL. 2017. Alternative polyadenylation of mRNA precursors. *Nature reviews. Molecular cell biology* 18:18–30.

Tokue S-I, Sasaki M, Nakahata N. 2009. Thromboxane A2-induced signal transduction is negatively regulated by KIAA1005 that directly interacts with thromboxane A2
1190 receptor. *Prostaglandins & other lipid mediators* 89:8–15.

Toufexis D, Rivarola MA, Lara H, Viau V. 2014. Stress and the reproductive axis. *Journal of neuroendocrinology* 26:573–586.

Trabzuni D, Ramasamy A, Imran S, Walker R, Smith C, Weale ME, Hardy J, Ryten M, North American Brain Expression Consortium. 2013. Widespread sex differences in
1195 gene expression and splicing in the adult human brain. *Nature communications* 4:2771.

Tsigos C, Chrousos GP. 2002. Hypothalamic–pituitary–adrenal axis, neuroendocrine

factors and stress. *Journal of psychosomatic research* 53:865–871.

Vermeij GJ. 1991. When biotas meet: understanding biotic interchange. *Science*

1200   253:1099–1104.

Wang Z, Burge CB. 2008. Splicing regulation: from a parts list of regulatory elements to

an integrated splicing code. *RNA* 14:802–813.

Wang Y, Liu J, Huang BO, Xu Y-M, Li J, Huang L-F, Lin J, Zhang J, Min Q-H, Yang W-

M, Wang X-Z. 2015. Mechanism of alternative splicing and its regulation.

1205   *Biomedical reports* 3:152–158.

Wang Z, Xiao X, Van Nostrand E, Burge CB. 2006. General and specific functions of

exonic splicing silencers in splicing control. *Molecular cell* 23:61–70.

Wendelaar Bonga SE. 1997. The stress response in fish. *Physiological reviews* 77:591–

625.

1210   Wong BBM, Candolin U. 2015. Behavioral responses to changing environments.

Behavioral ecology: official journal of the International Society for Behavioral

Ecology 26:665–673.

Yabas M, Elliott H, Hoyne GF. 2015. The Role of Alternative Splicing in the Control of

Immune Homeostasis and Cellular Differentiation. *International journal of molecular*

1215   *sciences* 17. DOI: 10.3390/ijms17010003.

Zhao C, Wang P, Si T, Hsu C-C, Wang L, Zayed O, Yu Z, Zhu Y, Dong J, Tao WA, Zhu

J-K. 2017. MAP Kinase Cascades Regulate the Cold Response by Modulating

ICE1 Protein Stability. *Developmental cell* 43:618–629.e5.

1220

SUPPLEMENTARY TABLES & FIGURES

1225

Table S1. List of Spliced Genes. This list contains the Entrez ID for all genes in which

an alternative splicing event was found during this study. Column headers refer to the

comparison (M= male, F= female, H= hypothalamus, P= pituitary, G= gonad, C=

control, S= stress)

1230

Figure S1. Male vs. Female Biological Process GO Analysis

1235   Figure S2. Male vs. Female Cellular Component GO Analysis

Figure S3. Male vs. Female Exon Motifs

Figure S4. Control vs. Stress Biological Process GO Analysis

Figure S5. Control vs. Stress Cellular Component GO Analysis

Protein Motifs of Spliced Exons Stress vs. Control

Figure S6. Control vs. Stress Exon Motifs

Distribution of Percent Spliced In (PSI) Values

A)

Splicing PSI: Female vs Male

B)

Splicing PSI: Control vs Stress

Negative values: higher in females

Negative values: higher in control

Treatment ☐ control ☐ stress

Sex ☐ female ☐ male

1245

Figure S7. Distribution of Percent Spliced In (PSI) Values for the a) control vs stress analysis and also the b) male vs female analysis. The figure is divided into vertical panels, each representing the tissues labeled a the top The right two panels show the distribution of Percent Spliced In (PSI) values, with "|" symbols at the base of each ridge

1250   plot each corresponding to the PSI value of a single event. The heights of the ridge plots should not be compared from one event to another, but within an event between treatments. Splicing event types are displayed along the y-axis (CE: core exon, AA: alternate acceptor, AD: alternate donor, RI: Retained intron). The coloring scheme is the

same as previous figures (light blue: control, orange: restraint stress, green: female,

1255   blue: male).