Non-invasive multiphoton imaging of neural structure and activity in *Drosophila*

Max Jameson Aragon[1,5,#], Mengran Wang[2,#], Jamien Shea[1], Aaron T. Mok[2], Haein Kim[1], Kawasi

M. Lett[3], Nathan Barkdull[4], Chris B. Schaffer[3], Chris Xu[2] and Nilay Yapici[1,*]

[1]*Department of Neurobiology and Behaviour, Cornell University, 14853, Ithaca, NY, USA*
[2]*School of Applied and Engineering Physics, Cornell University, 14853, Ithaca, NY, USA*
[3]*The Meinig School of Biomedical Engineering, Cornell University, 14853, Ithaca, NY, USA*
[4]*Department of Physics, University of Florida, 32611, Gainesville, FL, USA*
[5]*Current address: Princeton Neuroscience Institute, Princeton University, 08544, Princeton, NJ, USA*

[#]*These authors contributed equally to this work*
[*]*Corresponding author: Nilay Yapici (ny96@cornell.edu)*

Abstract

A limitation of current methods for examining neural activity over long periods of time in the fly, *Drosophila melanogaster*, is the need to remove the head cuticle and the underlying tissue to gain optical access to the brain, a process that damages circulation and restricts the length of imaging time. Here, we developed a non-invasive preparation for structural and functional imaging of the fly brain through the intact head cuticle. We first showed that the head cuticle transmits long-wavelength laser light with surprisingly high efficiencies. In fact, the tissue that interferes with laser light during multiphoton imaging is the air sacs underneath the head cuticle. We developed a non-invasive imaging preparation that compresses the air sacs, and used it to image the mushroom body Kenyon cells and central complex ring neurons through the cuticle at cellular resolution using both 2-and 3-photon microscopy. We also performed non-invasive short-term and long-term functional imaging (for the first time for 12 consecutive hours) of odour-evoked calcium responses from the mushroom body Kenyon cells. Our results demonstrate that the non-invasive imaging preparation developed here extends the time limits of current *in vivo* imaging methods used in flies that require an invasive surgery, and opens up new ways to capture neural activity from the intact fly brain.

Main

Animal nervous systems across lineages have evolved to solve many of the same problems, such as foraging for food and water, finding mates to reproduce, and avoiding predators to stay alive. They navigate their environment via coordinated movements, and learn and remember the relative values of sensory stimuli around them in order to maximize their fitness and survival. At each instant in time, an animal must evaluate external sensory information and its current behavioural state to decide what to do next [1-6]. A major technological challenge to revealing how the brain encodes behavioural states in real-time is that even the simplest neural computation involves interactions across the nervous system at various time scales, while our tools for assessing neural activity are restricted in time and space because of the currently available imaging sensors, methods, and preparations [7]. Optical methods remain the most established and fruitful path for revealing population dynamics in neural circuits at long time scales (ranging from minutes to hours) by providing high temporal and spatial resolution measurements [8-10].

The fly, *Drosophila melanogaster*, offers an ideal experimental system to investigate neural correlates of behavioural states and decisions because of its compact nervous system and diverse state-dependent behaviours that it executes in response to sensory stimuli [4, 11, 12]. To understand how molecularly defined neural circuits evaluate sensory information in different behavioural states, it is critical to capture the activity of populations of neurons over long time scales as flies are changing their physiological needs [8, 13]. These functional imaging experiments require non-invasive imaging preparations, which should allow chronic neural activity imaging for at least 12 to 24 hours. Current methods used in fly optical physiology require the fly head cuticle, trachea, and fat body to be removed by microsurgery to provide optical access to the nervous system [14-18]. These preparations are limited in imaging duration because after some time, the brain tissue starts dying due to damaged circulation resulting from the cuticle removal surgery. For example, with current imaging preparations, fly olfactory neurons show reliable Ca^{2+} responses for about four to five

hours after surgery [18]. A non-invasive imaging method and a preparation in which the head cuticle

and underlying tissue are left intact, thereby eliminating the need for traumatic head surgery before

functional imaging, is essential for advancing fly neuroscience research in the direction of chronic

recordings of neural activity during ongoing behaviours. This includes being able to image the

same fly brain across multiple days. In mice, multi-day imaging experiments are achieved by

implanting a cranial window following removal of part of the skull [19, 20]. Similar imaging

preparations have been developed for flies [14, 16, 21]. However, because imaging window

implantation requires a tedious surgery with low success rates and complications that occur

afterwards, these methods are not commonly used.

A recent development in *in vivo* multiphoton imaging is the use of long wavelength lasers in

3-photon (3P) microscopy which improves the signal-to-background ratio by several orders of

magnitude compared to current 2-photon (2P) imaging methods [22-24]. For example, 3P excitation at

1300 nm allows the functional imaging of neurons labelled with a genetically encoded Ca^{2+}

indicator GCaMP6 in the CA1 hippocampal region in an intact mouse brain [24, 25]. Recently, 3P

microscopy at 1700 nm excitation has shown promising results in imaging the fly brain through the

cuticle [26].

Here, we developed a non-invasive imaging preparation for flies to image neural structure

and activity through the intact head cuticle with both 2P and 3P microscopy. We first measured the

ballistic and total optical transmission of long-wavelength lasers through the dorsal fly head cuticle

and surprisingly found that the cuticle has high transmission at the wavelengths that are used to

excite fluorophores in 2P and 3P microscopy. We showed that the tissue that interferes with the

laser light and limits imaging through the cuticle into the brain is not the head cuticle but the air

sacs and the trachea underneath the cuticle. Next, we developed a new imaging preparation that

mildly compresses the air sacs and allows non-invasive imaging of the fly brain without any

surgery. Using this imaging preparation, we captured neural structures of mushroom body Kenyon

cells and the central complex ring neurons with cellular resolution at 910 nm (2P) and 1320 nm

(3P) excitation. We also performed short-term and long-term functional imaging of odour-evoked

neural activity of Kenyon cells comprising the mushroom body γ-lobes using a genetically encoded

Ca^{2+} indicator, GCaMP6s [27]. Kenyon cells are the primary intrinsic neurons in the insect mushroom

body. Diverse subtypes of Kenyon cells (n=~2200) extend their axons along the pedunculus and in

the dorsal and medial lobes [28-30]. These neurons receive and integrate information from

heterogeneous sets of projection neurons which carry olfactory, gustatory, and visual sensory

information [31, 32]. In our chronic 3P functional imaging experiments, we found that odour evoked

Ca^{2+} responses in the mushroom body γ-lobes change with time.

Our non-invasive imaging preparation developed here for the first time allows 2P and 3P

structural and functional imaging of the fly brain without any microsurgery, opening a door to

capturing neural activity at long time scales and potentially through the entire lifespan of flies.

Ultimately, we expect that our results will push forward the current methods that are used for

anatomical and functional imaging of the adult fly nervous system *in vivo*.

Results

Fly head cuticle transmits long wavelength laser light with high efficiency

To develop a non-invasive imaging preparation using multiphoton microscopy, we first tested the

transmission of infrared wavelengths through the fly head cuticle. Previous experiments showed

that the dorsal head cuticle of *Drosophila melanogaster* transmits light with increased efficiency at

longer wavelengths [33]. We measured the total and ballistic transmission of infrared wavelengths

through the cuticle. Dissected head cuticle samples were mounted between two glass coverslips,

and placed in the beam path between the laser source and the photodetector (Fig. 1a). The total and

ballistic transmission through the cuticle samples were measured using a custom-built system (Fig.

1b). For ballistic transmission, light from a single-mode fibre was magnified and focused on the

cuticle with a ~25µm spot size. Fig. 1c illustrates the light path of ballistic transmission

experiments. The sample was moved around to obtain measurements at different locations on the

same sample. Ballistic transmission through the cuticle was measured at seven different

wavelengths (852 nm, 915 nm, 980 nm, 1054 nm, 1310 nm, 1510 nm, 1630 nm). Our results

showed that ballistic transmission of the laser light through the cuticle samples increased with laser

wavelength (Fig. 1d). At all of the laser wavelengths tested, the ballistic transmission through the

cuticle was surprisingly high, reaching up to >90% at 1310 nm (Fig. 1d). The fluorescence signal

within the focal volume in 2P- and 3P-microscopy is mostly generated by the ballistic photons.

Next, we measured the total transmission through the head cuticle. For these measurements,

laser light from a single mode fibre was magnified and focused on the cuticle sample with a ~50μm

spot size. An integrating sphere was placed immediately after the cuticle to measure the total laser

transmission. Fig. 1e illustrates the light path of total transmission experiments. Total transmission

through the cuticle was measured at nine different wavelengths (532 nm, 633 nm, 852 nm, 915 nm,

980 nm, 1054 nm, 1310 nm, 1510 nm, 1630 nm). Similar to the ballistic transmission experiments,

we found that the total transmission increases with wavelength, reaching a plateau around 915 nm

(Fig. 1f). The total transmission at shorter wavelengths, 532 nm and 633 nm, was also relatively

high (>60%) which is critical for collecting emitted fluorescence produced by green (532 nm) and

red (633 nm) fluorophores after 2P or 3P excitation. Our results demonstrated that absorption of

long wavelength laser light by the *Drosophila* head cuticle is minimal, suggesting non-invasive

excitation of green (GFP, GCaMP) and red fluorophores (RFP, RCaMP) through the cuticle is

possible in adult flies using 2P or 3P excitation *in vivo*.

2P and 3P excitation allows non-invasive structural imaging of the fly brain

Based on our results for cuticle transmission, we next used a non-invasive imaging preparation and

a custom-built multiphoton microscope equipped with a wavelength-tuneable 2P and 3P laser

system to image the fly brain in head-fixed flies *in vivo* (Fig. 2a, Methods). We developed three

types of non-invasive imaging preparations with different head compression levels: no

compression, semi-compression, and full-compression (Fig. 2b-2d, Methods). As a proof of principle experiment, we expressed membrane tagged GFP selectively in mushroom body Kenyon cells and imaged the fly brain through the cuticle using 2P and 3P excitation at 910 nm and 1320 nm laser wavelengths, respectively (Fig. 2e-2g). Mushroom body Kenyon cells are bilaterally symmetric groups of neurons that are subdivided according to their axonal trajectories. The cell bodies of these neurons are located around the calyx of the mushroom bodies. Kenyon cell dendrites arborize in the calyx, while their axons fasciculate into anatomically distinct structures called lobes, with the dorsal lobes forming α and α′ branches, and the medial lobes containing β, β′, and γ branches [29, 30, 34]. We used transgenic flies that specifically expressed GFP in Kenyon cells forming α, β and γ lobes [35]. Using both 2P and 3P excitation, we were able to visualize different anatomical compartments of the fly mushroom body. In non-compressed flies, the mushroom body lobes were barely visible in both 2P and 3P imaged flies. Compressing the head against the glass with forceps during the curing process drastically improved image quality (Fig. 2e-2g, Supplementary video 1 & 2). Flies behaved healthier (based on observations of the leg movement) in semi-compressed preps compared to full-compressed preps, thus we used semi-compressed preps in all further structural and functional imaging experiments with 2P and 3P excitation.

We also investigated why head compression improves image quality in intact flies. Our hypothesis is that head compression reduces the volume of the air sacs and surrounding tissue, allowing better transmission of the long wavelength laser light through the structures underneath the head cuticle including dorsal air sacs and other tracheal branches. To test our hypothesis, we surgically removed air sacs from one side of the fly head and imaged the brain using 2P and 3P excitation without any head compression. As predicted, in both 2P and 3P imaged flies, we were able to see the mushroom body lobes on the side where air sacs were removed but not on the side with intact air sacs (Supplementary Fig. 1a-1c, Supplementary video 3). Our results clearly showed that the tissue that interferes with 2P and 3P laser light is not the cuticle itself but the air sacs and

153 the other tissues that are underneath the head cuticle. In parallel to our findings, a recent study also

154 showed that deflating the trachea within the fly brain increases the maximal imaging depth of

155 multiphoton microscopy [36].

3P non-invasive imaging of mushroom body Kenyon cells and central complex ring neurons
at cellular resolution.

Our non-invasive imaging experiments showed that through–cuticle imaging is possible with both

2P and 3P excitation. For further detailed structural imaging experiments, we used 3P excitation at

1320 nm. We imaged different anatomical compartments of the mushroom body Kenyon cells and

central complex ring neurons at cellular resolution (Fig. 3). We started acquiring images from the

deepest structures we could detect in the brain (depth=~150 µm), then moved to shallower areas in

~0.5 µm depth increments and ended imaging outside of the head cuticle (Fig. 3a-3e). In the flies

expressing membrane tagged GFP specifically in the mushroom body Kenyon cells, we were able

to visualize the calyx and axonal tracts along the peduncle as well as the dorsal and medial lobes

containing α, β and γ branches, respectively (Fig. 3a-3b,). In addition to the fluorescence signal,

adipose, epithelial-like tissues, and tracheal branches surrounding the mushroom body lobes

produced a bright Third Harmonic Generation (THG) signal (yellow) [37]. The tracheal branches

were most apparent near the medial lobes at 96 µm depth (Fig. 3a, mid panel). When we focused on

the calyx region at higher magnification using a compressed imaging prep with the posterior brain

positioned closer to the objective lens, we were also able to distinguish single Kenyon cell bodies,

which are densely packed around the calyx (Fig. 3c).

Next, we imaged the central complex, another higher order processing centre, that is located

in the dorsal and medial part of the fly brain (Fig. 3d-3g). The insect central complex is highly

interconnected brain neuropil which processes sensory information and guides a diverse set of

behavioural responses which include navigation, walking initiation, and turning direction [38-40]. It is

composed of functionally distinct, but anatomically connected, compartments: the protocerebral

bridge, ellipsoid body, fan-shaped body, and the paired noduli [38]. The ellipsoid body consists of a

group of neurons, the ring neurons, that extend their axons to the midline and form a ring-like

structure that is composed of different layers [38, 39, 41]. Using an ellipsoid body-specific promoter, we

expressed membrane tagged GFP in the fly brain and imaged the ring neurons through the cuticle at

1320 nm (Supplementary video 4 & 5). Ring neuron cell bodies, as well as the lateral triangle and

the ellipsoid body layers, were clearly visible (Fig. 3d & 3e). Using ring neuron arbours and

tracheal branches, we estimated the lateral resolution of our 3P images. The full width at half

maximum (FWHM) of the lateral brightness distribution measured by a ring neuron's neurite cross

section was ~1.24 µm for the fluorescent signal (Fig. 3f) and ~0.79 µm for tracheal branches

captured by the THG signal (Fig. 3g). We were also able to image the ellipsoid body neuropil with

2P excitation through the cuticle. Compared with 3P excitation, the resolution and contrast in

images taken by 2P excitation is somewhat reduced in the semi-compressed preparation

(Supplementary Fig. 2a & 2b). We also repeated our 3P imaging experiments using a fibre laser-

amplifier system with an integrated infrared optical parametric amplifier (OPA, KM Labs). The

results using this low power laser system are comparable to what we have obtained previously

(Supplementary Fig. 3a & 3b), indicating that a more compact, less powerful laser source is

adequate for imaging fly brain structures with 3P microscopy.

Our results demonstrate that non-invasive 3P microscopy is able to capture single neurons

and fine anatomical structures within the mushroom body and central complex with comparable

resolution to the open cuticle preparations using 2P microscopy [18].

Simultaneous 2P and 3P imaging of odour-evoked responses of mushroom body Kenyon cells

through the head cuticle

Encouraged by our structural imaging results, we next tested the applicability of 2P and 3P

microscopy to capture neural activity in the fly brain through the intact head cuticle. In these

experiments, a custom odour delivery and behaviour system was used, where flies were head fixed

on a polymer ball under the microscope (Fig. 4a-4c). We expressed GCaMP6s selectively in Kenyon cells and stimulated the fly olfactory sensory organ, the antenna, with the food odour apple cider vinegar (Fig. 4b). Using a multiphoton scanning microscope, odour evoked Ca^{2+} responses of mushroom body γ-lobes were simultaneously captured with 2P (910 nm) and 3P (1320 nm) excitation using temporal multiplexing and the semi-compressed imaging prep (Fig. 4a-4e, methods). A brief 3s odour stimulus triggered a robust fluorescence increase in Kenyon cell axons comprising the mushroom body γ-lobes (Fig. 4f, Supplementary video 6). Kenyon cell axons are innervated by distinct subsets of dopaminergic neurons, which carry contextual information to the mushroom body depending on the fly's previous experiences. Based on dopaminergic innervation, γ-lobes can be subdivided into five anatomical compartments [42] (Fig. 4d & 4e). To determine whether the localized dopaminergic input along the Kenyon cell axons impacts the olfactory responses in the γ-lobes, we calculated the normalized peak fluorescence signal in each compartment of the γ-lobes. We found no differential responses in $\Delta F/F_0$ amplitude to food odour across different compartments of the mushroom body or between 2P and 3P excitation of GCaMP6s (Fig. 4g-4i, Supplementary Fig. 4a). Using a full-compressed imaging preparation, we also imaged the Kenyon cell body responses to odour stimulation and were able to manually distinguish single cell Ca^{2+} traces on a given imaging plane (Supplementary Fig. 4b & 4c).

Together, these results suggested that 2P and 3P microscopy with 910 nm and 1320 nm excitation, respectively, are effective in capturing odour evoked Ca^{2+} responses in the mushroom body lobes and cell bodies through the intact head cuticle, and achieve temporal resolution comparable to 2P microscopy with the open cuticle preparations [42].

3P imaging of neural activity captures odour evoked responses of Kenyon cells for 12 hours

Studying how neural circuits change activity during learning or in alternating behavioural states requires non-invasive chronic imaging methods that permit recording neural activity over long time scales. Leveraging the non-invasive nature of our preparation, we pushed the limits of functional

imaging of the fly brain in response to odour stimulation to longer time scales (12 hours). Using

our custom odour delivery system, we stimulated the fly antenna with apple cider vinegar in ~10

minute sessions every four hours while imaging through the head cuticle using our 3P microscope

(Fig. 5a). To determine the absolute signal level during long-term imaging, we first calculated the

photon counts for representative trials at each time point from the mushroom body $\gamma2$ compartment.

The photon counts per pixel per second had magnitudes above baseline similar to previously

reported levels for 3P excitation of GCaMP6s protein in the mouse brain [23] (Supplementary Fig. 5).

Next, we analysed the average odour responses per fly in each γ-lobe compartment and time

point (Fig. 5b-5e). We calculated the normalized peak fluorescent signal and the area under the

normalized fluorescence curve, as metrics representing the odour response strength and persistence

during chronic imaging, respectively. Our analysis showed that the evoked response to food odour

did not change with food and water deprivation in any of the imaged γ-lobe compartments (Fig. 5f).

However, the persistence of the odour evoked Ca^{2+} response in all of the γ-lobe compartments

increased as the experiment progressed (Fig. 5g). We especially noticed a change at 4 hours after

the first odour trial, which may reflect a critical state change in the nervous system. Flies continued

to show robust calcium responses for at least 4 hours after the change in dynamics (Fig. 5b-5e).

During these long-term imaging experiments, we captured the fly's behaviour in parallel

with the odour stimulation to assure that the fly stayed alive during long-term imaging

(Supplementary Fig. 6). We also checked for heat induced damage after 3P imaging in flies using

HSP70 protein as a marker for stress response in glial cells and neurons [43, 44]. We first tested

whether HSP70 protein levels reflect heat induced stress response in the fly brain by placing flies in

a 30°C incubator for 10 minutes and staining for HSP70 protein right after the incubation. Flies that

were kept at room temperature had low levels of the HSP70 protein in a very small number of

neurons in the brain (Supplementary Fig. 7a). In contrast, placing flies in a 30°C incubator for 10

minutes caused a significant increase in the number of HSP70 positive neurons and an elevation of

HSP70 protein levels across the fly brain (Supplementary Fig. 7b). Next, we tested whether 3P

imaging causes a heat induced stress response in fly neurons. Head-fixed flies in a semi-

compressed imaging prep were imaged for 10 minutes using 3P excitation at 1320nm. we chose to

use 10 minutes because this is the total amount of imaging time per time point during our chronic

functional imaging experiments. Our results showed that there was no obvious tissue damage or

heat-stress response detected by the HSP70 protein levels at the average power levels that we used

for functional imaging experiments (~20mW) (Supplementary Fig. 7c). However, increasing laser

power (~25mW) caused structural abnormalities and a significant increase in HSP70 protein levels

in the fly brain (Supplementary Fig. 7d). These results suggested that 3P non-invasive imaging of

the fly brain is safe at lower power levels we used in our experiments, and the non-invasive

imaging preparation developed here allows recording of neural activity within an individual fly

over long-time scales, which was previously not possible with commonly used open-cuticle

preparations.

Discussion

Plasticity in neural states and the ability to learn and remember relevant information are hallmarks

of nervous systems, and allow animals to adapt and respond to dynamic environments.

Investigating how physiological states, sleep, and learning change the function of neural circuits

requires tracking the activity of molecularly defined sets of neurons over long time scales. These

experiments require non-invasive imaging methods to record neural activity *in vivo*. In this study,

we extensively analysed the fly cuticle by measuring its ballistic and total transmission efficiency

across different laser wavelengths. Our results showed that the fly cuticle transmits long

wavelength light with high efficiency (Fig. 1). Surprisingly, the transmission at 915 nm

wavelength, which is used for 2P microscopy, is only slightly lower than 1320 nm, which is used

for 3P excitation of GFP and GCaMP. Based on these results, we investigated the use of 2P and 3P

microscopy for non-invasive anatomical imaging of the fly brain. We showed that both 2P and 3P

microscopy can be used for non-invasive imaging of fly brain structure and function (Fig. 2-4).

While 3P imaging appeared to have higher contrast than 2P imaging, we did not see noticeable

difference in the recorded activity traces when performing simultaneous 2P and 3P functional

imaging of the mushroom body. We also showed that the tissues interfering with the transmission

of the 2P and 3P excitation light during in-vivo imaging are the air sacs located between the head

cuticle and the brain, rather than the head cuticle itself (Supplementary Fig. 1). By compressing the

fly head to a glass coverslip, we flatten this heterogenous tissue, which increases the transmission

of laser light and therefore allows high resolution, non-invasive imaging of the fly brain structures.

The non-invasive imaging method developed here also significantly extends the time frame

of current *in vivo* imaging preparations used for anatomical and functional studies in fly

neuroscience. Non-invasive imaging of the fly brain will allow us to capture the activity of neural

populations during changing behavioural states; facilitate decoding of neural plasticity during

memory formation; and permit observation of changes in brain structures during development and

aging. Our first application of long-term functional imaging of the fly brain captures food odour

responses from mushroom body γ-lobes, for the first time up to 12 hours. Our results suggest that

odour evoked Ca^{2+} responses gradually change during the repeated odour stimulation. In these

chronic imaging experiments, we aimed to image odour responses specifically from γ-lobe

compartments. However, because γ and β lobes are very close to each other in depth, some of the

fluorescence Ca^{2+} signal we have captured might have been pooled from both of these mushroom

body medial lobes. Further experiments using more specific drivers for γ or β lobes will aim to

understand the specific location and origins of the change in Ca^{2+} signals and the mechanisms that

mediate it. While our focus here was to develop non-invasive in vivo structural and functional

imaging methods that can extend imaging length for fly neuroscience research, we anticipate that

there will be a wide variety of uses for this technology in *Drosophila* and in other insect models

including mosquitoes.

Methods

Fly stocks

Flies were maintained on conventional cornmeal-agar-molasses medium at 23-25°C and 60-70%

relative humidity, under a 12hr light: 12hr dark cycle (lights on at 9 A.M.). Fly strains and sources:

Mef2-GAL4.247 (Bloomington # 50742); *GMR15B07-GAL4* (Bloomington # 48678), *10XUAS-*

IVS-mCD8::GFP (Bloomington # 32186); *20XUAS-IVS-GCaMP6s* (Bloomington # 42746).

Detailed genotypes of all fly strains used in the paper are as follows

Fig. 1

Males, w^{1118}/Y; 10XUAS-IVS-mCD8::GFP; Mef2-GAL4.

Fig. 2

Males, w^{1118}/Y; 10XUAS-IVS-mCD8::GFP; Mef2-GAL4.

Fig. 3

Panel a, c: Males, w^{1118}/Y; 10XUAS-IVS-mCD8::GFP; Mef2-GAL4.

Panel d-g: Males, w^{1118}/Y; 10XUAS-IVS-mCD8::GFP; GMR15B07-GAL4.

Fig. 4 & 5

Males, w^{1118}/Y; 20XUAS-IVS-GCaMP6s; Mef2-GAL4.

Supplementary Fig. 1

Males, w^{1118}/Y; 20XUAS-IVS-GCaMP6s; Mef2-GAL4.

Supplementary Fig. 2

Males, w^{1118}/Y;10XUAS-IVS-mCD8::GFP;GMR15B07-GAL4.

Supplementary Fig. 3

Panel a, Males, w^{1118}/Y; 10XUAS-IVS-mCD8::GFP; Mef2-GAL4

Panel b, Males, w^{1118}/Y;10XUAS-IVS-mCD8::GFP;GMR15B07-GAL4.

Supplementary Fig. 4-7

Males, w^{1118}/Y; 20XUAS-IVS-GCaMP6s; Mef2-GAL4.

Optical transmission measurements of the fly head cuticle

Drosophila cuticle was dissected from the dorsal head capsule of flies that are age and gender controlled (male, 5 days old). The dissected cuticle was sandwiched between two #1.5 coverslips with UV curable resin to avoid dehydration of the sample (Fig. 1a). Measurements from each dissected cuticle was done within a day. The first few measurements were repeated at the end of all measurements to ensure that dehydration or protein degradation, which may vary the optical properties of the tissue, did not happen as the experiment progressed. The total transmission and ballistic transmission of cuticle samples were measured using a custom-built device (Fig. 1b). For ballistic transmission experiments, we assume collimated light passes through the sample since the Rayleigh range for a 25 µm ($1/e^2$) focus spot is ~1mm - 2mm (λ = 1630 nm – 852 nm), which is much larger than the thickness of the whole coverslip sandwich-prep. The transmitted light from the cuticle was coupled to a single mode fibre with identical focusing optics to measure its ballistic transmission. A photodetector was used to monitor the power fluctuation during the measurement and thus to ensure identical incident power (~10mW) on the cuticle in each measurement. An infrared camera was used to image the sample and incident beam to ensure the incident light spot is always on the cuticle and to avoid very dark pigments (usually at the edge of the cuticle), ocelli, and possible cracks introduced during dissection. The ballistic transmission of the cuticle was then calculated as the optical power ratio between the ballistic transmissions of the cuticle and the surrounding areas without the cuticle, containing only the UV curable resin (i.e., the reference) using the equation below (SMF = single mode fibre, T = transmission, T_0= reference transmission):

$$T_{ballistic} = \frac{T^{SMF}}{T_0^{SMF}}$$

For total transmission experiments, we assume collimated light passes through the sample since the Rayleigh range for a 50 µm ($1/e^2$) spot size is ~5mm - 10mm (λ = 1630 nm – 532 nm), which is much larger than the thickness of the whole coverslip sandwich-prep. A photodetector was used to monitor the power fluctuation during the measurement. The incident power on the sample is

~10mW. An infrared camera was setup to visualize the light spot and the cuticle when the

integrating sphere is removed. The cuticle was translated to obtain measurements at different

locations. The total transmission of the cuticle was then calculated as the optical power ratio

between the transmissions of the cuticle and the surrounding areas without the cuticle, containing

only the UV curable resin (i.e., the reference) measured by the integrating sphere (IS = integrating

sphere, T = transmission, T_o= reference transmission).

359
$$T_{total} = \frac{T^{IS}}{T_0^{IS}}$$

Laser sources

Excitation source (Fig. 2-5, Supplementary Fig 1,2 &3-5): The excitation source is a wavelength-

tuneable optical parametric amplifier (OPA, Opera-F, Coherent) pumped by a femtosecond laser

with a MOPA (Master Oscillator Power Amplifier) architecture (Monaco, Coherent). A prism pair

(10SF10, Newport) and a silicon wafer are used to compensate for the dispersion of the optics of

the light source and the microscope including the objective, for 1320 nm. The laser repetition rate

is maintained at 1 MHz for both structural and functional imaging.

Excitation source (Supplementary Fig. 3): The alternative laser used as the excitation source for 3P

microscopy is an optical parametric amplifier (OPA, KM Labs), pumped by the KM Labs Y-

Fi™HP. A two-prism glass compressor (SF 10 glass) is used to compensate for the normal

dispersion of the optics of the light source and the microscope. The Y-Fi™ OPA has an operating

wavelength centred at 1300 nm and outputs average power of ~400 mW (1 MHz repetition rate).

Pulse duration is less than 150 fs. Power control of the laser was achieved using a half wave plate

(HWP) (Thorlabs) and a polarizing beam splitter cube (PBS) (Thorlabs) and ScanImage module for

MATLAB (Mathworks).

Multiphoton microscopes

Imaging setup (Fig. 2-5, Supplementary Fig. 1,2 & 4,5): The images were acquired with a custom-

built multiphoton microscope. The scan lens is a C-coated achromat (AC254-030-C-ML, Thorlabs)

for high transmission at 1320 nm, and the transmission of the tube lens (Sutter Instrument) is 82%.

For imaging experiments that use GFP and GCaMP6s, we used a custom high numerical aperture

(NA) water immersion microscope objective (Olympus XLPLN25XWMP2, 25X, NA 1.05), which

is specially coated for high transmission (~70%) at 1320 nm. 2 detection channels were used to

collect the fluorescence signal and the third harmonic generation (THG) signal by photomultiplier

tubes (PMT) with GaAsP photocathode (H7422-40, Hamamatsu). For GFP and GCaMP6s imaging

at 1320 nm, fluorescence signal and THG signal were selected by a 520/60 nm band-pass filter

(FF01-520/60-25, Semrock) and a 435/40 nm band-pass filter (FF02-435/40-25, Semrock),

respectively. For signal sampling, the PMT current is converted to voltage and low-pass-filtered

(200 kHz) by a transimpedance amplifier (C7319, Hamamatsu). Analog-to-digital conversion is

performed by a data acquisition card (NI PCI- 6110, National Instruments). ScanImage 5.4 (Vidrio

Technologies) running on MATLAB (MathWorks) was used to acquire images and control a 3D

translation stage to move the sample (MP-285, Sutter Instrument Company). All imaging depths

and thickness are reported in raw axial movement of the motorized stage, unless otherwise stated.

Imaging setup (Supplementary Fig. 3): Images were acquired using a custom-built multiphoton

microscope with a pair of galvanometric scanners (Cambridge Technology). The objective used

was a high numerical aperture water immersion objective (Olympus XLPLN25XWMP2, 25X, NA

1.05). Signal from specimens was collected through the objective and then reflected by multiple

dichroic beam splitters. (FF520-Di02- 25 x 36, Semrock, and FF484-Di01- 25 x 36, Semrock) to

the detectors. The two-channel detection system consists of a green fluorescence signal and third

harmonic generation (THG). ScanImage was used to control the ILS100CC motorized translation

stage (Newport). High-resolution structural images were typically taken with 512x512

pixels/frame, 0.24 Hz frame rate, and multiple frame averages at each depth.

Non-invasive *in vivo* imaging preparation

All animals used for imaging experiments were male flies with indicated genotypes kept in 25°C incubators and maintained on conventional cornmeal-agar-molasses medium. Flies used for chronic functional experiments were 2-7 days old, and flies used for short term functional experiments were 1-4 days old. To perform intact brain imaging, flies were first head-fixed in a 40 mm weigh dish (VWR#76299-236) with a hole made with forceps. A drop of UV curable resin (Liquid plastic welder, Bondic®) was applied to the head and thorax, which was then cured with blue light (~470 nm) and fused to a cover glass. Fly proboscises were immobilized with blue light curable resin to minimize head motion caused by muscle contractions. Only flies in Supplementary Fig. 1 went through surgery to remove the air sacs using a sharp needle.

Odour delivery

Food odour, apple cider vinegar, was delivered using a custom built olfactometer as described previously [45]. Clean room air was pumped (Active Aqua Air Pump, 4 Outlets, 6W, 15 L/min) into the olfactometer, and the flow rate was regulated by a mass flow controller (Aalborg GFC17). Two Arduino controlled 3-way solenoid valves (3-Way Ported Style with Circuit Board Mounts, LFAA0503110H) controlled air flow. One valve delivered the odourised airstream either to an exhaust outlet or to the main air channel, while another valve directed air flow either to the stimulus or control channel. The stimulus channel contained a 50 ml glass vial containing undiluted apple cider vinegar (volume=10 ml) (Wegmans), while the control channel contained a 50 ml glass vial containing mineral oil (volume=10 ml). For odour stimulation experiments, flies were placed approximately 1 cm from a clear PVC output tube (OD = 1.3 mm, ID = 0.84 mm), which passed a ~1< L/min air stream to the antennae. The odour stimulus latency was calculated before the experiments using a photo ionization detector (PID) (Aurora Scientific). We sampled odour delivery using the PID every 20 ms and found average latency to peak odour amplitude was < 100ms across 34 measurements.

Simultaneous 2P and 3P Structural and Functional Imaging

Simultaneous imaging with 2P excitation and 3P excitation is achieved by temporal multiplexing of the 910- nm Ti:Sapphire laser and 1,300- nm Spirit-NOPA laser (Fig. 4 and Supplementary Fig.4): They were spatially overlapped at the same focal position after the objective. The 910- nm laser was intensity modulated with an electro-optic modulator (EOM), switching it on for $\sim$1 μs between two adjacent Spirit-NOPA laser pulses. The EOM is driven by a radio frequency signal triggered by the Spirit-NOPA pulse train so that the sample is alternatively illuminated either with 910- nm laser for 2P excitation or 1,320- nm laser for 3P excitation. By recording the NOPA pulses and the PMT signal at 5 MHz sampling rate (200 ns pixel time), the 2P and 3P excited fluorescence signals can be temporally demultiplexed with a postprocessing MATLAB script.

For short term functional imaging, flies were food deprived for 16-24 hours in vials with a wet Kim wipe. Each odour stimulation trial consisted of 50 seconds of clean mineral oil, 3 seconds of undiluted apple cider vinegar stimulus, and another 50 seconds of mineral oil. Between trials, scanning was stopped for 20 seconds to minimize the risk of imaging-induced tissue stress. Five trials were performed sequentially.

3P Chronic Functional Imaging

Flies used in long term functional imaging experiments were fed 100 mM sucrose just before the first trial to assure that the flies were in a fed state. Each odour stimulation trial consisted of 50 seconds of clean mineral oil, 3 seconds of undiluted apple cider vinegar stimulus, and another 50 seconds of mineral oil. Between trials, scanning was stopped for 20 seconds. The images were captured at 256x512 pixel resolution and 2.13 Hz frame rate. Five trials were performed sequentially and the five-trial block was repeated every four hours. Between trial blocks, scanning was stopped, and air passing through the stimulation tube was redirected to the exhaust valve to prevent desiccation. To further prevent desiccation, flies were placed on a water-absorbing polymer bead. Fly behaviour was videotaped from the side to capture any movement by a CCD camera

(FLIR, Blackfly S, BFS-U3-13Y3M-C) equipped with a 1.5-megapixel macro lens (Computar,

COM MLM3X-MP, 3X Macro Zoom). We captured fly behaviour with videos at 4.3 Hz speed

using the Spinnaker SDK (Flir) image acquisition software under IR light illumination.

Photon counting

To show the absolute signal strength in the unit of photon count rate, pixel values are multiplied by

a calibration factor, which is obtained by taking the ratio of the photon count rate with a photon

counter (SR400, Stanford Research Systems) and the average pixel value, simultaneously recorded

on a uniform fluorescein sample (Supplementary Fig. 5).

Immunohistochemistry for brain tissue damage assessment

To investigate laser-induced stress in the brain during 3-photon imaging through the intact head

cuticle, 1-2-day old male homozygous MB>GCaMP6s flies were imaged continually at 1320 nm

for 10 minutes. The laser was focused 70μm below the cuticle with ~20mW or ~25mW average

power. Flies were head compressed and fed 100mM sucrose before imaging to reproduce actual

experimental conditions. Laser exposed brains were dissected approximately 2 hours after

mounting. To test HSP70 antibody efficacy on fly brains, a positive control group of flies was first

exposed to 30°C for 10 minutes in an incubator to induce HSP70 expression. Negative control

group of flies was kept at room temperature. For whole mount staining, brains were dissected in

phosphate-buffered saline (PBS) and incubated in 4% paraformaldehyde (PFA) in PBS for 20-30

minutes at room temperature on an orbital shaker. Tissues were washed 3-4 times over 1 hour in

PBS (calcium- and magnesium-free; Lonza BioWhittaker #17-517Q) containing 0.1% Triton X-

100 (PBT) at room temperature. Samples were blocked in 5% Normal Goat Serum in PBT (NGS-

PBT) for 1 hour and then incubated with primary antibodies diluted in NGS-PBT for 24 hours at

4°C. Primary antibodies used were anti-GFP (Torrey Pines, TP40, rabbit polyclonal, 1:1000), anti-

BRP (DSHB, nc82, mouse monoclonal, 1:20), and anti-HSP70 (Sigma, SAB5200204, rat

monoclonal, 1:200). The next day, samples were washed 5-6 times over 2 hours in PBT at room

temperature and incubated with secondary antibodies (Invitrogen) diluted in NGS-PBT for 24

hours at 4°C. On the third day, samples were washed 4-6 times over 2 hours in PBT at room

temperature and mounted with VECTASHIELD Mounting Media (Vector Labs, Burlingame, CA,

USA) using glass slides between two bridge glass coverslips. The samples were covered by a glass

coverslip on top and sealed using clear nail polish. Images were acquired at 1024x1024 pixel

resolution at ~1.7 μm intervals using an upright Zeiss LSM 880 laser scanning confocal microscope

and Zeiss digital image processing software ZEN. Maximum projection images of Z-stacks were

generated in the ZEN image-processing software. The power, pinhole size and gain values were

kept the same for all imaged brains during confocal microscopy.

Data analysis

Resolution measurements: We measured the lateral brightness distribution of small features within

the GFP labelled fly brains (Fig. 3f) and the THG signal (Fig. 3g), which sets an upper bound on

the lateral resolution value. Lateral intensity profiles measured along the white lines were fitted by

a Gaussian profile for the estimation of the lateral resolution.

Image processing for structural imaging. TIFF stacks containing fluorescence data were converted

to 8 bits. When necessary, stacks were registered using the TurboReg plugin in ImageJ. Images

shown in Fig. 3 are an average of 3 z-stacks spanning 3 microns. 3D reconstructions of the z-stacks

were generated using the Imaris v9.2, Bitplane Inc., software available at http://bitplane.com.

Images shown in Supplementary Fig. 2 are an average of single z sections at different depths.

Image processing for activity recordings. TIFF stacks containing fluorescence data were converted

to 32 bits, and pixel values were left unscaled. When necessary, stacks were registered using the

TurboReg plugin in ImageJ. Images acquired during the multiplexed 2p-3p imaging sessions were

first median filtered with a filter radius of 10 pixels to reduce high amplitude noise. To compute

$\Delta F/F_0$ traces, γ-lobe ROIs were first manually selected using a custom Python script. F_0 was

computed as the average of 10 frames preceding stimulus onset. The F_0 image was then subtracted

from each frame, and the resulting image was divided by F_0. The resulting trace was then low-pass

filtered by a moving mean filter with a window size of 2 frames (8 frames for multiplexed activity

data). Data were analysed using Python (code available upon request) and plotted in Microsoft

Excel. Peak $\Delta F/F_0$ was determined by the peak value within 20 frames after the odour delivery. We

measured the persistence as the total area under the normalized curve after the peak $\Delta F/F_0$ value. To

prevent underestimating persistence as a result of negative values in the $\Delta F/F_0$ trace, values that

followed a zero-crossing were set to zero. Averaged traces were manually inspected for each lobe

and trial block, and rejected if peak $\Delta F/F_0$ was lower than the noise detected in the entire trial.

These traces had lower than 0.1 peak $\Delta F/F_0$ value. These accounted for less than 15% of the total

trials in 3 flies tested for chronic imaging.

Fly movement analysis: Fly motion traces were obtained using the DeepLabCut tracking program

[46]. To train the network, we manually annotated movement in the fly bodies and legs for 340

frames across multiple cropped and contrast-enhanced videos. The data shown in Supplementary

Fig. 6 shows the activity of front tarsi or hind legs.

Statistical analysis

All statistical analyses were performed using Prism V8.0.1 (GraphPad). Data collected in

functional imaging experiments are shown as mean ± SEM. Details of statistical methods are

reported in the Fig. legends. Statistical significance in chronic imaging across time points and in

different mushroom body compartments was determined with a two-way repeated measures

ANOVA. In all of the figures, plots labelled with different letters are significantly different from

each other.

Acknowledgements

We thank Joe Fetcho, Andy Bass, David Owald, and members of the Yapici Lab for comments on the manuscript. We acknowledge Bloomington Drosophila Stock Centre (NIH P40OD018537) and the Developmental Studies Hybridoma Bank (NICHD of the NIH, University of Iowa) for reagents. We thank Li Yan McCurdy (Yale University) and Matt Einhorn (Cornell University) for help with the design and construction of the custom built olfactometer, Elena Zhong for training the neural network for behavioural analysis in Deep Lab Cut, Joe Fetcho (Cornell University) for help with Imaris software, and Nancy M. Bonini (University of Pennsylvania) for her advice on the HSP70 antibody. Research in N.Y.'s laboratory is supported by a Cornell University Nancy and Peter Meinig Family Investigator Program, a Pew Biomedical Scholar Award, the Alfred P. Sloan Foundation Award, AFAR Research Grant for Junior Faculty, NSF NeuroNex Program Grant (DBI-1707312), and NIH R35 ESI-MIRA Grant (R35GM133698-01).

Author contributions

The authors have made the following declarations about their contributions: Conceived and supervised the study: C.X., and N.Y. Designed the experiments: M.J.A, M.W., J.S., A.M., C.X., and N.Y. Performed the experiments: J.S., N.B., and A.M. (Fig. 1, Supplementary Fig. 1), M.J.A., and M.W. (Fig. 2-5, Supplementary Fig. 2 -6), K.L. (Supplementary Fig. 3), H.K. (Supplementary Fig. 7). Analysed the data and prepared Figures: M.J.A., M.W, J.S., A.M. and N.Y. Wrote the paper: N.Y. with input from C.X., M.J.A., M.W., C.S., J.S., A.M., K.L., and H.K.

Competing financial interests

The authors declare no competing financial interests.

Data availability

All raw imaging data supporting the findings of this study are available from the corresponding author on request.

References

1.   Tinbergen, N. The study of instinct. (Oxford University Press, Oxford; 1969).

2.   Nevin, J.A. The study of behavior : learning, motivation, emotion, and instinct. (Scott, Foresman, Glenview, IL; Brighton; 1973).

3.   Hunt, L.T. & Hayden, B.Y. A distributed, hierarchical and recurrent framework for reward-based choice. *Nat Rev Neurosci* **18**, 172-182 (2017).

4.   Dickson, B.J. Wired for sex: the neurobiology of Drosophila mating decisions. *Science* **322**, 904-909 (2008).

5.   McFarland, D.J. Decision making in animals. *Nature* **269**, 15-21 (1977).

6.   Lutcke, H., Margolis, D.J. & Helmchen, F. Steady or changing? Long-term monitoring of neuronal population activity. *Trends Neurosci* **36**, 375-384 (2013).

7.   Lerner, T.N., Ye, L. & Deisseroth, K. Communication in Neural Circuits: Tools, Opportunities, and Challenges. *Cell* **164**, 1136-1150 (2016).

8.   Luo, L., Callaway, E.M. & Svoboda, K. Genetic Dissection of Neural Circuits: A Decade of Progress. *Neuron* **98**, 256-281 (2018).

9.   Svoboda, K. & Yasuda, R. Principles of two-photon excitation microscopy and its applications to neuroscience. *Neuron* **50**, 823-839 (2006).

10.   Ji, N., Freeman, J. & Smith, S.L. Technologies for imaging neural activity in large volumes. *Nat Neurosci* **19**, 1154-1164 (2016).

11.   Yapici, N., Zimmer, M. & Domingos, A.I. Cellular and molecular basis of decision-making. *EMBO Rep* **15**, 1023-1035 (2014).

12.   Barron, A.B., Gurney, K.N., Meah, L.F., Vasilaki, E. & Marshall, J.A. Decision-making and action selection in insects: inspiration from vertebrate-based theories. *Front Behav Neurosci* **9**, 216 (2015).

13.   Simpson, J.H. & Looger, L.L. Functional Imaging and Optogenetics in Drosophila. *Genetics* **208**, 1291-1309 (2018).

14.   Grover, D., Katsuki, T. & Greenspan, R.J. Flyception: imaging brain activity in freely walking fruit flies. *Nat Methods* **13**, 569-572 (2016).

15.   Minocci, D., Carbognin, E., Murmu, M.S. & Martin, J.R. In vivo functional calcium imaging of induced or spontaneous activity in the fly brain using a GFP-apoaequorin-based bioluminescent approach. *Biochim Biophys Acta* **1833**, 1632-1640 (2013).

16.   Sinha, S. et al. High-speed laser microsurgery of alert fruit flies for fluorescence imaging of neural activity. *Proc Natl Acad Sci U S A* **110**, 18374-18379 (2013).

17.   Seelig, J.D. et al. Two-photon calcium imaging from head-fixed Drosophila during optomotor walking behavior. *Nat Methods* **7**, 535-540 (2010).

18.   Wang, J.W., Wong, A.M., Flores, J., Vosshall, L.B. & Axel, R. Two-photon calcium imaging
reveals an odor-evoked map of activity in the fly brain. *Cell* **112**, 271-282 (2003).

19.   Trachtenberg, J.T. et al. Long-term in vivo imaging of experience-dependent synaptic
plasticity in adult cortex. *Nature* **420**, 788-794 (2002).

20.   Hefendehl, J.K. et al. Repeatable target localization for long-term in vivo imaging of mice
with 2-photon microscopy. *J Neurosci Methods* **205**, 357-363 (2012).

21.   Huang, C. et al. Long-term optical brain imaging in live adult fruit flies. *Nat Commun* **9**, 872
(2018).

22.   Wang, M. et al. Comparing the effective attenuation lengths for long wavelength in vivo
imaging of the mouse brain. *Biomed Opt Express* **9**, 3534-3543 (2018).

23.   Wang, T. et al. Three-photon imaging of mouse brain structure and function through the intact
skull. *Nat Methods* **15**, 789-792 (2018).

24.   Horton, N.G. et al. In vivo three-photon microscopy of subcortical structures within an intact
mouse brain. *Nat Photonics* **7** (2013).

25.   Ouzounov, D.G. et al. In vivo three-photon imaging of activity of GCaMP6-labeled neurons
deep in intact mouse brain. *Nat Methods* (2017).

26.   Tao, X. et al. Transcutical imaging with cellular and subcellular resolution. *Biomed Opt
Express* **8**, 1277-1289 (2017).

27.   Chen, T.W. et al. Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*
499, 295-300 (2013).

28.   Strausfeld, N.J., Hansen, L., Li, Y., Gomez, R.S. & Ito, K. Evolution, discovery, and
interpretations of arthropod mushroom bodies. *Learn Mem* **5**, 11-37 (1998).

29.   Crittenden, J.R., Skoulakis, E.M., Han, K.A., Kalderon, D. & Davis, R.L. Tripartite
mushroom body architecture revealed by antigenic markers. *Learn Mem* **5**, 38-51 (1998).

30.   Ito, K. et al. The organization of extrinsic neurons and their implications in the functional
roles of the mushroom bodies in Drosophila melanogaster Meigen. *Learn Mem* **5**, 52-77 (1998).

31.   Yagi, R., Mabuchi, Y., Mizunami, M. & Tanaka, N.K. Convergence of multimodal sensory
pathways to the mushroom body calyx in Drosophila melanogaster. *Sci Rep* **6**, 29481 (2016).

32.   Owald, D. & Waddell, S. Olfactory learning skews mushroom body output pathways to steer
behavioral choice in Drosophila. *Curr Opin Neurobiol* **35**, 178-184 (2015).

33.   Lin, Y.Y. et al. Three-wavelength light control of freely moving Drosophila Melanogaster
for less perturbation and efficient social-behavioral studies. *Biomed Opt Express* **6**, 514-523 (2015).

34.   Zheng, Z. et al. A Complete Electron Microscopy Volume of the Brain of Adult Drosophila
melanogaster. *Cell* **174**, 730-743 e722 (2018).

35.   Krashes, M.J., Keene, A.C., Leung, B., Armstrong, J.D. & Waddell, S. Sequential use of
mushroom body neuron subsets during drosophila odor memory processing. *Neuron* **53**, 103-115
(2007).

36.   Hsu, K.J., Lin, Y.Y., Chiang, A.S. & Chu, S.W. Optical properties of adult Drosophila brains
in one-, two-, and three-photon microscopy. *Biomed Opt Express* **10**, 1627-1637 (2019).

37.   Barad, Y., Eisenberg, H., Horowitz, M. & Silberberg, Y. Nonlinear scanning laser
microscopy by third harmonic generation. *SPIE milestone series* **175**, 47-49 (2003).

38.   Wolff, T., Iyer, N.A. & Rubin, G.M. Neuroarchitecture and neuroanatomy of the Drosophila
central complex: A GAL4-based dissection of protocerebral bridge neurons and circuits. *J Comp*
Neurol **523**, 997-1037 (2015).

39.   Pfeiffer, K. & Homberg, U. Organization and functional roles of the central complex in the
insect brain. *Annu Rev Entomol* **59**, 165-184 (2014).

40.   Seelig, J.D. & Jayaraman, V. Neural dynamics for landmark orientation and angular path
integration. *Nature* **521**, 186-191 (2015).

41.   Xie, X. et al. The laminar organization of the Drosophila ellipsoid body is semaphorin-
dependent and prevents the formation of ectopic synaptic connections. *Elife* **6** (2017).

42.   Cohn, R., Morantte, I. & Ruta, V. Coordinated and Compartmentalized Neuromodulation
Shapes Sensory Processing in Drosophila. *Cell* **163**, 1742-1755 (2015).

43.   Podgorski, K. & Ranganathan, G. Brain heating induced by near-infrared lasers during
multiphoton microscopy. *J Neurophysiol* **116**, 1012-1023 (2016).

44.   Lindquist, S. Varying patterns of protein synthesis in Drosophila during heat shock:
implications for regulation. *Dev Biol* **77**, 463-479 (1980).

45.   Raccuglia, D. et al. Presynaptic GABA Receptors Mediate Temporal Contrast Enhancement
in Drosophila Olfactory Sensory Neurons and Modulate Odor-Driven Behavioral Kinetics. *eNeuro*
3 (2016).

46.   Mathis, A. et al. DeepLabCut: markerless pose estimation of user-defined body parts with
deep learning. *Nat Neurosci* **21**, 1281-1289 (2018).

Fig. 1: Ballistic and total optical transmission of the fly head cuticle. a, Schematic of the cuticle transmission preparation. Fly head cuticle is dissected and placed in UV curable resin between two cover slips. **b,** Schematic of the cuticle optical transmission measurement setup. **c,** Schematic of the ballistic optical transmission through the cuticle. **d,** Results of the ballistic optical transmission experiments across various wavelengths (the mean and standard error of mean of all measurements (n = 56 at each wavelength) from 5 different cuticle samples). **e,** Schematic of the total optical transmission through the cuticle. **f,** Results of the total optical transmission experiments across various wavelengths (the mean and standard error of mean of all measurements (n = 20 at each wavelength) from 5 different cuticle samples). ANOVA with post-hoc Tukey's test was conducted to compare transmission at different wavelengths (p<0.001).

Fig. 2. **Non-invasive imaging of the fly brain with 2P and 3P excitation. a,** Schematic of the
multiphoton microscope setup. Fly head is fixed to a cover slip and placed under the objective
(HWP: half-wave plate, PBS: polarization beam splitter, PMT: photomultiplier tube). The imaging
window on the fly head is shown in the picture (lower left). **b-d**, The uncompressed and
compressed imaging preparations. The first column shows the side image of the fly that is head
fixed to the cover glass (scale bar =1mm). The second and third columns show the uncompressed
and compressed fly head visualized under a brightfield (top view) and fluorescent dissecting
microscopes (widefield-fluo), respectively. Arrows and the rectangle area in Widefield-Fluo
column indicate the imaging window (scale bar=200μm). **e-g,** Cross section imaging of the
mushroom body (MB) Kenyon cells expressing CD8-GFP through the head cuticle at 910 nm (2P)
and 1320 nm (3P) excitation. The Z projections of 2P (cyan, left) and 3P (green, right) imaging
stacks. For each imaging preparation, the same fly head is imaged with 3P excitation and 2P
excitation (scale bar=20μm).

Fig. 3. High resolution 3P structural imaging of the mushroom body and the central complex

neurons through the cuticle with 1320 nm excitation. a, Cross section images of the mushroom

body (MB) lobes through the cuticle with 3P excitation at 1320 nm (green). Third harmonic

generation (THG) imaging visualizes the tracheal arbours (yellow). White arrows indicate different

MB compartments that are identified (scale bars= 30 μm, semi-compressed prep). **b**, 3D

reconstruction of the z-stack images in 3A (scale bars= 30 μm). **c,** Kenyon cell bodies that are

located around the calyx (scale bar= 5 μm, compressed prep.) **d,** Cross section imaging of the

central complex (CC) ring neurons through the cuticle with 1320 nm excitation (green). Third

harmonic generation (THG) imaging visualizes the tracheal arbours (yellow). Arrows indicate

different CC compartments that are identified. **e,** 3D reconstruction of the z stack images in 3c,

GFP (upper panel) and THG (lower panel) (scale bars= 30 μm, semi-compressed prep). **f,** The GFP

profile of CC ring neurons (green) and **g,** the THG profile of surrounding trachea (yellow). Lateral

intensity profiles measured along the white lines are fitted by a Gaussian profile for the lateral

resolution estimation (scale bars=20 μm, semi-compressed prep).

Fig. 4. Simultaneous 2P and 3P functional imaging of short-term odour-evoked responses of the mushroom body Kenyon cells. a, Schematic of the custom made olfactometer. This olfactometer is used to puff odours to the fly that is placed on a ball for functional Ca^{2+} imaging using GCaMP6s and a semi-compressed imaging prep. An Arduino board synchronizes the FLIR Black Fly S Camera to capture fly behaviour. ScanImage software is used to capture images from temporally multiplexed 2P (910nm) and 3P (1320nm) excitation at 160x165 pixel resolution (described in detail in the methods). **b,** Stimulus timeline. Flies were stimulated with air (50s), before and after the odour stimulus (odour + air, 3s). Each trial lasted 103s. Flies were stimulated 5 times using the same odour stimulus scheme. **c,** Picture of the head fixed fly on the ball under the 3-photon microscope. The coloured dots show the body parts that were tracked and used to train the DeepLabCut neural network. **d,** Schematic of the mushroom body anatomy indicating the location of α, β, and γ lobes. **e,** γ-lobes have discrete anatomical compartments (shown as $\gamma2$–$\gamma5$). **f,** GCaMP6s is expressed in the mushroom body Kenyon cells. Normalized ($\Delta F/F_0$) GCaMP6s signal is shown before (left) and after (right) odour stimulus (scale bar= 20 μm). **g,** Odour-evoked responses of Kenyon cells captured by 2P excitation at 910 nm and **h,** 3P excitation at 1320 nm. **i,** Comparison of 2P and 3P average responses over time (n=3 flies, 4-5 trials per fly, data are presented as mean ± SEM in **g** and **h,** grey bar indicates when stimulus is present).

Fig. 5. Long-term 3P imaging of odour-evoked responses of the mushroom body Kenyon cells

a, Stimulus timeline for chronic odour imaging. GCaMP6s signal is captured from Kenyon cells

axons innervating mushroom body γ-lobes using the semi-compressed prep. Normalized $\Delta F/F_0$

signal is shown before (left) and after (right) the odour stimulus (scale bar= 20 μm). **b-e,**

Quantification of the normalized $\Delta F/F_0$ signal over time in each γ-lobe compartment. Each lobe's

response is colour coded differently. Grey bar indicates when the odour stimulus is present. Each

grey line indicates the average response of a fly over multiple trials in a given hour. The average

response of 3 flies is shown. Each compartment's response is labelled with a different colour. **f,**

Quantification of the peak amplitude ($\Delta F_{max}/F_0$). **g,** Area under the curve (AUC) after the odour

stimulus for each compartment of the γ-lobes (Two-way repeated measures ANOVA. Data are

presented as mean ± SEM, ns= not significant, ***=p<0.001, n=3 flies, 3-5 trials per time point).

Supplementary Information

Supp. Fig. 1. Removal of air sacs allows 2P and 3P imaging in an uncompressed prep.

Cross section imaging of the mushroom body (MB) Kenyon cells expressing GCaMP6s through the

head cuticle at 910 nm (2P) and 1320 nm (3P) excitation after removing air sacs on one side of the

head. Head is not compressed to the cover slip **a,** The Z projections of 2P (cyan) and **b,** 3P (green)

imaging stacks. For each imaging preparation, the same fly head is imaged with 3P excitation and

2P excitation. Mushroom body structures are visible on the side where air sacs are removed. The

dotted lines show the area where the air sacs block imaging. **c,** Higher resolution 3P images taken

from the brain side where the air sacs are removed. Mushroom body lobes and peduncle are clearly

visible without the head compression. Scale bars = 20μm.

Supp. Fig. 2. Non-invasive imaging of the central complex ring neurons with 2P and 3P excitation. Cross section imaging of the Central Complex (CC) ring neurons expressing CD8-GFP through the head cuticle at 910 nm (2P) and 1320 nm (3P) excitation. **a-b**, The Z projection (top-most) and single optical section images at ~10 μm step of ellipsoid body processes. **a,** 2P (cyan) and **b,** 3P (green) images are shown. For each imaging preparation, the same fly head is imaged with both 3P and 2P excitation. (scale bars=25μm, semi-compressed prep).

Supp. Fig. 3. 3P structural imaging of the fly brain through the cuticle with 1320 nm using an alternative laser source. Cross section imaging of the fly brain through the cuticle at 1320 nm. **a,** Compressed z-stack images of flies expressing membrane tagged GFP in the Mushroom Body (MB) Kenyon cells (green, right). **b,** Compressed z-stack images of flies expressing membrane tagged GFP in Central Complex (CC) ring neurons (green, right). Third harmonic generation (THG) imaging visualizes the tracheal arbours (yellow). Arrows indicate different MB and CC compartments that are identified with 3P imaging (scale bars= 25µm).

Supp. Fig. 4. Short term 2P and 3P functional imaging of odour-evoked responses of the mushroom body gamma lobes and cell bodies. a, Ca^{2+} imaging of odour responses in gamma lobe compartments. Left and right panels show single trial normalized $\Delta F/F_0$ signal over time using simultaneous 2P (cyan) and 3P (green) excitation from indicated gamma lobe compartments (semi-compression prep). **b,** 3P structural image of the Kenyon cell bodies and ROI selection. **c,** Quantification of the normalized $\Delta F/F_0$ signal over time in 5 ROIs, each corresponding to a Kenyon cell soma imaged with 3P excitation. Grey bar indicates when the odour stimulus is present (scale bars= 10μm, full-compression prep).

Supp. Fig. 5. Photon counts during 3-photon functional imaging. Odour-evoked neural activity in GCaMP6s-labeled Kenyon cells forming mushroom body γ-lobes, imaged at average power of ~20 mW and 1 MHz repetition rate. **a-d,** Plots show the activity of the γ2-compartment of the mushroom body in fly#2 in a single trial across different time points. The fluorescence intensity is converted to photon counts per frame in the selected ROI (~50x50 pixels, frame rate=2.13Hz.).

743 **Supp. Fig. 6. Fly motion traces during chronic imaging. a,** Picture of the head fixed fly on the

744 polymer ball under the 3-photon microscope. The coloured dots show the body parts that were

745 tracked and used to train the DeepLabCut neural network. **b-c,** The speed of indicated legs (blue

746 and green) during the odour stimulus trials. Grey bar indicates when the stimulus is present. Speed

747 is calculated as arbitrary units per second (AU/s).

MB>GCaMP6s (1320nm, anti-BRP, anti-GFP, anti-HSP70)

Supp. Fig. 7. HSP70 staining of fly brains shows no heat-induced stress responses after 1320 nm 3P excitation with average power (~20mW) used in functional imaging experiments. a, Without heat shock, there is minimal HSP70 protein expressed in the fly brain. **b,** When flies are exposed to heat (30ºC) for 10 minutes, the HSP70 protein level is significantly elevated across the brain. **c-d**, HSP70 protein expression in the fly brain after 3-photon imaging at 1320 nm for 10 minutes. **c,** There is no obvious change in HSP70 protein levels when power is 20mW. **d,** HSP70 protein expression is significantly elevated when power is 25mW (scale bars= 50 µm). The optical power is measured at the sample plane of the microscope using a power meter (n=2-4 flies per condition).

Supp. Video 1: Z stack of the mushroom body (MB) Kenyon cells expressing CD8-GFP.
Cross section imaging through the head cuticle at 1320 nm (3P) excitation (scale bar=20μm, compressed prep).

Supp. Video 2: Z stack of the mushroom body (MB) Kenyon cells expressing CD8-GFP.
Cross section imaging through the head cuticle at 910 nm (2P) excitation (scale bar=20μm, compressed prep).

Supp. Video 3: Z stack of the mushroom body (MB) Kenyon cells expressing GCaMP6s.
Imaging is done through the head cuticle using 1320 nm (3P) excitation after removing air sacs only on one side of the head (scale bar = 50μm, no compression).

Supp. Video 4: Z stack of the ellipsoid body (EB) ring neurons expressing CD8-GFP.
Cross section imaging through the head cuticle at 1320 nm (3P) excitation (scale bar=20μm, semi-compressed prep).

Supp. Video 5: 3D reconstruction of Z stacks of the ellipsoid body (EB) ring neurons expressing CD8-GFP. (Green= CD8-GFP, yellow=THG, scale bar= 40 μm, semi-compressed prep).

Supp. Video 6: Odour-evoked responses of mushroom body gamma lobes captured from an intact fly head through the cuticle. Functional imaging is performed using a food odour, apple cider vinegar, with 3P excitation at 1320 nm. Video is sped up 10x (semi-compressed prep).