Title:

Rest-task Modulation of fMRI-derived Global Signal Topography is Mediated by Transient Co-activation Patterns

Jianfeng Zhang [1,2,*], Zirui Huang [3], Shankar Tumati [4], Georg Northoff [1,4,5,6,*]

[1] Mental Health Center, Zhejiang University School of Medicine, Hangzhou, China

[2] College of Biomedical Engineering and Instrument Sciences, Zhejiang University, Hangzhou, China

[3] Center for Consciousness Science, Department of Anesthesiology, University of Michigan Medical School, Ann Arbor, Michigan, 48109, USA

[4] Institute of Mental Health Research, University of Ottawa, Ottawa, Canada

[5] Center for Cognition and Brain Disorders, Hangzhou Normal University, Hangzhou, China

[6] Graduate Institute of Humanities in Medicine, Taipei Medical University, Taipei, Taiwan

* jianfeng_zhang@zju.edu.cn (J.Z); georg.northoff@theroyal.ca (G.N)

Abbreviated title: Rest-task Modulation of Global Signal Topography

Abstract

Recent resting-state fMRI studies have revealed that the global signal (GS) exhibits a non-uniform spatial distribution across the gray matter. Whether this topography is informative remains largely unknown. We therefore tested rest-task modulation of global signal topography by analyzing static global signal correlation and dynamic co-activation patterns in a large sample of fMRI dataset (n=837) from the Human Connectome Project. The GS topography in the resting-state and in seven different tasks was first measured by correlating the global signal with the local timeseries (GSCORR). In the resting state, high GSCORR was observed mainly in the primary sensory and motor regions, while low GSCORR was seen in the association brain areas. This pattern changed during the seven tasks, with mainly decreased GSCORR in sensorimotor cortex. Importantly, this rest-task modulation of GSCORR could be traced to transient co-activation patterns at the peak period of global signal (GS-peak). By comparing the topography of GSCORR and respiration effects, we observed that the topography of respiration mimicked the topography of global signal in the resting-state whereas both differed during the task states; due to such partial dissociation, we assume that GSCORR could not be equated with a respiration effect. Finally, rest-task modulation of GS topography could not be exclusively explained by other sources of physiological noise. Together, we here demonstrate the informative nature of global signal topography by showing its rest-task modulation, the underlying dynamic co-activation patterns, and its partial dissociation from respiration effects during task states.

Keywords: global signal, human connectome project, resting state, task modulation, co-activation patterns

Introduction

One of the major confounds that limits the cognitive and clinical applications in fMRI is the global signal (GS), which is defined as the spatial average of time varying blood-oxygen-level-dependent signals [1,2]. The GS is often considered to represent physiological noise caused by respiratory and cardiac events [3], and has been recommended to be regressed out during data preprocessing prior to secondary analyses.

However, other studies have shown that the GS is not merely non-neuronal noise, and it contains important information about neuronal activity. For instance, the GS in fMRI exhibits a high correlation to electrophysiological measures on the cortical level [4,5]; its global fluctuation may partly stem from subcortical regions (e.g., basal forebrain) relating to arousal [6,7]; and its variance fluctuates across time of day [8]. Finally, abnormalities in the level of the GS have been observed in psychiatric disorders like schizophrenia [9] and bipolar disorder [10]. Together, these findings underscore the potentially informative nature of the GS.

One key feature of the GS is its topographical pattern [6,10-12]. Although the GS is generated by the shared activity across the gray matter, recent studies have demonstrated a non-uniform topographical distribution of GS across brain regions in both monkeys [7] and humans [6,11,13] during the resting state. Higher levels of GS correlation (GSCORR) are observed in the primary sensory regions such as the auditory and visual cortices, and lower levels of GS correlation are seen in the higher-order cortical regions including the prefrontal cortex during the resting state.

The informative nature and thus the functional relevance of resting state GS topography is still subject to debate. On the one hand, this topography seems to be functionally relevant as it correlates with cognitive performance in a healthy population [12] and shows abnormal in

psychiatric disorders like schizophrenia [11], autism spectrum disorder [14,15] or bipolar

disorder [10]. On the other hand, the resting state GS topography resembles the spatial pattern of

a respiration effect [3,16-18]; this raises the question whether the GS topography merely

represents physiological noise, i.e., respiration. One way to reconcile the two seemingly

contradictory observations seems to be investigating the modulation of GS during task state

relative to the resting state and how that stands in relation to respiration effect.

The overarching aim of this study is to probe the informative nature [19] of GS topography by

examining its modulation during various cognitive tasks relative to the resting state, i.e., rest-task

modulation. For that purpose, we utilize a large sample of fMRI dataset from the Human

Connectome Project (HCP) including two days' resting states and seven different tasks. First, we

quantified the GS topography and its modulation during different tasks. We sought to

demonstrate whether and how external tasks modulate GS topography, e.g., in a task-specific or

task-unspecific manner. Secondly, following a recent study showing how GS may govern the

brain's spatiotemporal dynamics [20], we explored the possible mechanism underlying rest-task

modulation of GS topography by analyzing its dynamic nature in terms of transient co-activation

patterns at the critical time points of GS (i.e., the peaks of GS) [6]. Thirdly, we tested whether

rest-task modulation of GS topography on the cortical level stems from the contributions of

subcortical regions [6] , or can be explained by the non-neuronal physiological noise, such as

respiration [3,21]. Finally, given the importance of reproducibility of fMRI studies in general

[22], we examined the rest-task modulation in an independent 7 Tesla dataset from the HCP with

two additional tasks.

Results

Global signal topography in the resting and task states

We first calculated the functional connectivity between the GS and time course in each

grayordinate (i.e., gray matter in cortical and subcortical regions) using Pearson correlation (i.e.,

global signal correlation (GSCORR)) with fisher's z transformation (Fig 1A). We quantified the

GSCORR topography at a group level by averaging the maps of GSCORR across 837 subjects

for resting states at two days and seven tasks, respectively. Consistent with previous findings

[6,11], a typical GS topography during resting state was observed, with higher levels of

GSCORR in the sensory regions (visual, auditory and somatosensory regions) and lower levels

of GSCORR in the association regions (prefrontal and parietal cortices) (Fig 1B). Interestingly,

the GSCORR topography in tasks differed from the resting state, where higher GSCORR was

persisted in visual but not in somatosensory regions (Fig 1B).

We further quantified the spatial similarity of the GS patterns among the resting states and tasks

using intraclass correlation coefficient (ICC) [23], which was widely used for investigating the

spatial similarity in test-retest reliability studies in rs-fMRI [22,24,25]. In reporting these

findings, we categorized the ICC into five common intervals[26]: $0 < ICC \leq 0.2$ (slight); $0.2 <$

$ICC \leq 0.4$ (fair); $0.4 < ICC \leq 0.6$ (moderate); $0.6 < ICC \leq 0.8$ (substantial); and $0.8 < ICC \leq 1.0$

(almost perfect). Additionally, given the spatial similarity analyses can be confounded by the

local spatial adjacency at the grayordinate level, we thus minimized this potential by performing

the spatial similarity analyses based on a parcellation template with 1000 ROIs [27]. Firstly, we

observed an almost identical GS topography during the resting states on the two days (ICC =

0.9986, 95% CI: [0.9985 0.9988]) (Fig 1C), suggesting a high reliability of GS topography across time. High ICCs were also observed across different tasks (ICC = 0.9356, 95% CI: [0.9296 0.9411]) (Fig 1C). The ICC was also high between resting state and tasks (ICC = 0.8666, 95% CI: [0.8504 0.8813]), however, the value was lower compared to both rest-rest and task-task similarities (p < 0.001 in both, Fisher's z test, Fig 1C). Together, the spatial pattern of GSCORR showed small but significant changes from resting state to task state.

Fig 1. **Spatial topography of global signal correlation**. **(A)** An illustration of global signal correlation (GSCORR) as a correlation between GS and time series in each region of gray matter. **(B)** Spatial patterns of GSCORR in two days' resting state and seven tasks. Top panel yielded the averaged spatial patterns for the resting states and task states, respectively. Bottom panel yielded the spatial pattern in each condition. **(C)** The spatial similarity was performed by using intraclass correlation coefficient (ICC), at the group level, based on a parcellation with 1000 ROIs. Top panel demonstrated the ICC matrix between two days' resting states and seven tasks. Bottom bar chart demonstrated the ICC ± 95% confidence interval for rest-rest, rest-task, and task-task respectively. *** p<0.001. Data available at DRYAD: https://doi.org/10.5061/dryad.xsj3tx9bw

Task modulation of GS topography

An approach to specify the informative nature of GS topography is rest-task modulation. If GSCORR changes from rest to task, the GSCORR must be assumed to contain some information. For that purpose, we first investigated the overall GSCORR across time (REST1 and REST2),

and state (rest and seven tasks). The comparison of GSCORR within the two rests showed no

significant difference despite the large sample size (n=837). In contrast, the comparisons of

GSCORR between rest and tasks showed a consistent reduction of GSCORR in all tasks (in all

contrasts between rest and tasks, t > 12, Cohen's d>0.4, Bonferroni corrected at α<0.01) (Fig 2A).

The reduction of GSCORR can occur uniformly across all regions in the brain, or, alternatively,

can be manifested in a non-uniform way across different regions. To address this issue, we

performed grayordinate-based comparison between rest and tasks. As the sample size was large

and therefore easily reached significance, we used a strict threshold at Bonferroni corrected at α

< 0.01 for further analyses, and the results were illustrated by Cohen's d as an index of effect

size which is insensitive to sample size [28]. We observed a non-uniform regional distribution of

task modulation of GSCORR. A large number of regions, centered in somatosensory, showed

GSCORR reduction during the task states, while only some regions exhibited task-related

increase or remained unchanged (Fig 2B).

We further tested whether the rest-task modulation of GSCORR (i.e., reduced, increased, or

unchanged), is task-unspecific or task-specific. We thereby generated the overlapping maps, and

calculated the number of task modulations (i.e., reduced, unchanged, or increase) across the

seven tasks for each grayordinate (Fig 2C top panel), and the percentages of grayordinates for the

number of each modulation (Fig 2C bottom panel). For those regions showing GSCORR

reduction, 54.9% regions showed reduction in at least 4 tasks (4-7 tasks), while only 15.5%

regions showed GSCORR reduction in 1-3 tasks. For those regions that remained unchanged,

37.8% exhibited similar GSCORR values across at least 4 tasks (4-7 tasks), while 33.4% of these

regions remained unchanged in 1-3 tasks. Finally, we refrained from making this calculation in

the regions that showed GSCORR increase as their number was rather small and inconsistent

(20.1% in 1-3 tasks and 4.5% in 4-7 tasks) across the various tasks (Fig 2C bottom panel).

Together, these findings suggest that the dominant rest-task modulation of GSCORR was task-

unspecific decrease as it mostly occurs across different tasks.

We lastly examined the regional specificity of GSCORR in rest-task modulation. GSCORR

reductions were observed mainly in the auditory and somatosensory regions as well as in the

regions belonging to the DMN. In contrast, the task-related regions exhibited unchanged (plus a

small set of increased) modulation, including the primary visual cortex (six of the seven tasks

included visual components/stimuli), and the regions belonging to frontoparietal executive

control network and ventral attention network (most of the tasks involved some executive and/or

attentional function) (Fig 2C top panel).

Taken together, we observed that the GSCORR topography was modulated during different tasks

in mostly task-unspecific ways. Such rest-task modulation was mainly manifested in GSCORR

reduction in a large set of regions while only some regions exhibited task-related GSCORR

increase (and others remained unchanged).

Fig 2. **Rest-task modulation of global signal topography.** **(A)** overall GSCORR in day1

(REST1) and day2 (REST2) resting state and seven tasks. **(B)** grayordinate-based group

comparison for GSCORR. The maps were tested with paired t-tests between task and rest, the

maps were thresholded at Bonferroni corrected $\alpha < 0.01$, and illustrated by Cohen's d to inspect

the effect size. **(C)** top panel: the overlapping map across the 7 tasks illustrated the counted

number of each grayordinate showing decrease (left), unchanged (middle), and increase (right) in

the tasks when comparing to rest at the level of Bonferroni corrected $\alpha < 0.01$. Bottom panel

illustrated the percentage of grayordinates for each counted number in the overlapping maps. ***

p<0.001. Data available at DRYAD: https://doi.org/10.5061/dryad.xsj3tx9bw

Instantaneous co-activation pattern of GS

Previous studies have shown that selective averaging of fMRI frames exhibiting regional peaks

of BOLD activity spatially mimics rsfMRI networks obtained via static seed-based correlation

analysis [29-31]. A recent study has shown that this relationship may also hold true for the GS

[6], suggesting that, instead of heightened static correlations between regions, global signal

topography may be traced to the dynamic co-activation patterns that occur over brief epochs [30].

Therefore, we hypothesized that rest-task modulation of GS topography may be traced to the

dynamics of transient co-activation at the peak time points of the GS.

The co-activation pattern of GS were extracted by simply averaging the time points at the top 17%

GS (GS-peak), a same threshold as used in a previous study [6] (Fig 3A). The topography of GS-

peak resembled that of GSCORR (Fig 3B) with both showing analogous rest-task modulation

pattern (Fig 3C). By inspecting the spatial similarity between GSCORR and GS-peak, we

observed that the diagonal value of the similarity matrix reached around 1, suggesting that these

two measurements (GSCORR, GS-peak) provide almost identical information about GS spatial

pattern. Moreover, our data suggest that rest-task modulation in GS topography is closely related

to the dynamic transition of co-activation pattern over brief epochs (Fig 3D).

Fig 3. **Instantaneous activation topography at the peak of GS.** **(A)** An illustration of

calculating the co-activation pattern at the peak (top 17%) of the GS (GS-peak). **(B)** Spatial

patterns of GS-peak in the two days' resting states and the seven tasks. Top panel yielded the

averaged spatial patterns for rest and task, respectively. Bottom panel yielded the spatial pattern

in each condition. **(C)** Grayordinate-based group comparisons for GS-peak. The spatial maps

were tested with paired t-tests, the maps were thresholded at Bonferroni corrected $\alpha < 0.01$, and

illustrated by Cohen's d to account for the effect size. **(D)** ROIs-based spatial similarity between

GSCORR and GS-peak. Data available at DRYAD: https://doi.org/10.5061/dryad.xsj3tx9bw

Decomposing GS-peak into a subset of co-activation patterns

Although we observed that the GS topography reflected an instantaneous co-activation at the

peak time points of GS, it is important to understand whether the GS topography is a single

united entity or a combination of different co-activation sets. To address this question, we

adopted a data driven approach (i.e., k-means clustering algorithm) that partitioned the whole-

brain frames into spatially congruent co-activation patterns (CAPs) (Fig 4A), and assigned each

fMRI frame to a cluster label [20,32]. The occurrence rate of these CAPs at the peak time point

of GS was measured by the number of CAP's occurrences divided by the total number of peak

time points (Fig 4A).

As previous studies have demonstrated that a few recurring dominant network states explain the

vast majority of rs-fMRI temporal dynamics in both mouse [20] and human [31,33], we here

single out the spatial patterns of 6 and 8 CAPs. We determined six CAPs as the optimized

number, because these six CAPs yielded previously partitioned large-scale rsfMRI networks into

three pairs of "mirror" motifs, and was dissociable across pairs (Fig 4C and see S1 Fig for

results of 8 CAPs) [20,32]. For example, the CAP1 and CAP4 clearly showed a pair of co-

(de)activation networks in somatosensory (CAP1) and frontoparietal network (CAP4). The

CAP2 and CAP5 showed a pair of visual (CAP2) and ventral attention network (CAP5). And the

CAP3 and CAP6 was a pair of default-mode network (CAP3) and dorsal attention network

(CAP6) (Fig 4B).

We next examined the difference of CAP occurrence rates at the time points of the GS peak

between resting state and tasks. During the resting state, the topography of GS-peak was mainly

constituted by CAP1 (18%) and CAP2 (22%). While during tasks, the contribution of CAP1

increased to 33% (Δrates = 15.6%, p < 0.001) on average and the contribution of CAP2

decreased to 8% (Δrates = -13.4%, p < 0.001) (Fig 4B and S2 Fig for percentages in each task).

Finally, we examined whether the rest-task modulation of GSCORR topography is associated

with a shift towards any particular CAPs. The spatial correlation between the ΔGSCORR and

CAP1 showed significant positive correlations (r = 0.33 on average, p < 0.001) and the CAP4 as

its opposite showed significant negative correlation (r=-0.36 on average, p < 0.001), whereas the

ΔGSCORR and CAP2 showed significant negative correlations (r = -0.66 on average, p < 0.001)

and its opposite (CAP5) showed significant positive correlations (r = 0.67 on average, p < 0.001).

Together, these findings suggested that the GS topography is constituted by a combination of

several co-activation patterns which were modulated in the frequency of their occurrence during

the tasks.

Fig 4. Transient co-activation patterns (CAPs) at GS-peak. (A) an illustration of the k-means

clustering algorithm that partitions the whole-brain frames into spatially congruent co-activation

patterns (CAPs) (step 1), and calculating the occurrence rate of each CAP at GS-peak (step 2).

(B) spatial topography of CAPs and their occurrence rates at the time points of GS-peak. Task

modulation denoted the difference in the occurrence rate of the CAP's between task and rest

(Δrate). **(C)** Spatial correlation between the CAPs. The six CAPs were composed of three pairs

of opposite co-activation patterns (i.e., CAP1 vs. CAP4, CAP2 vs. CAP5, and CAP3 vs. CAP6),

as shown by their spatially negative correlations. **(D)** Spatial correlation between the CAPs and

the rest-task difference in GSCORR, i.e., ΔGSCORR. Data available at DRYAD:

https://doi.org/10.5061/dryad.xsj3tx9bw

Spatial topography of respiration effect

Previous studies showed that the GS in the resting state is closely related to respiration

fluctuations [3,16,17]. We therefore investigated the relationship between the GS and respiration

timeseries through their respective spatial topographies and rest-task modulation. If the GS

indeed mirrors the fluctuations of respiration, one would expect to observe a comparable spatial

topography and rest-task modulation between the respiration and GS.

We first transformed the respiration timeseries into respiration volume per time (RVT) (see

details in Materials and Methods), which has been shown to be closely related with the GS

[3,16,17]. We performed cross-correlations over the time lags from -72 to 72 s between the GS

and RVT and then grouped into rest and task states. The highest negative correlation was

observed around 12 s lag, which was similar to the previous findings of the negative relationship

between GS and RVT [3,34]. However, the GS-RVT correlation showed differences between

resting and task states, where overall weaker correlations were seen in the task states around the

time interval ranging from -38 s to 18 s (paired t-test, Bonferroni corrected $\alpha < 0.01$) (Fig 5A).

We next calculated the RVT topography by correlating the flipped (reversing the negative into a

positive value, to render the RVT topography comparable to GS topography given the negative

correlation) RVT signal with the time series in each region with a 12-s lag. The RVT correlation

(RVTCORR) showed a weak (RVTCORR ranging from 0 to 0.05 while GSCORR ranging from

0 to 0.3, Figs 1B and 5B) but stable (spatial similarity between two days' rest: ICC = 0.9913, 95%

CI: [0.9901 0.9923], Fig 5C) spatial pattern in the resting state. In contrast, the spatial patterns of

RVTCORR varied (Fig 5B) and were less similar (ICC = 0.3983, 95% CI: [0.3712 0.4264], Fig

5C) across tasks, suggesting an unstable topography of RVTCORR during tasks. The spatial

pattern of rest-task modulation revealed significant reductions in somatosensory and visual

regions during tasks, and those regions were the ones with higher GSCORR in the resting state

(Fig 5D).

Finally, to quantify the contribution of respiration to the GS, we investigated the spatial

similarity between GSCORR and RVTCORR within conditions, i.e., resting state and tasks (Fig

5E). The spatial pattern of RVTCORR resembled the pattern of GSCORR in the resting state

(ICC = 0.8607, 95% CI: [0.8437 0.8760]) whereas that was not the case during the tasks (ICC =

0.57, 95% CI: [0.5271 0.6109]). These results confirmed the major contribution of respiration to

the GS in especially the resting state as observed in previous studies [3,16,17]. In contrast, the

contribution of respiration to GS topography was significantly reduced during the tasks,

suggesting a dissociation of GS and respiration.

Fig 5. Spatial topography of the respiration effect. (A) An illustration of cross-correlation

between the GS and RVT, and calculating a correlation map between the timeseries of

respiration and timeseries in each region of the gray matter. The respiration timeseries was first

transformed into the respiration volume per time (RVT), as the difference between the upper and

lower envelope (see Materials and Methods for details). Next the cross-correlation was

performed between the GS and RVT to determine the time lag with the strongest correlation, i.e.,

12 s (step 1). The RVT was flipped and correlated with the time series in gray matter with 12 s

lag (step 2) to obtain the RVTCORR map. The time interval showing significant rest-task

differences of the GS-RVT correlations was indicated by red line, thresholded at Bonferroni

corrected $\alpha < 0.01$. **(B)** Spatial patterns of RVTCORR in two days' resting states and seven tasks.

Top panel corresponded to the averaged spatial patterns for resting and task states, respectively.

Bottom panel corresponds to the spatial pattern in each condition. **(C)** The spatial similarity (ICC

± 95% CI) of RVTCORR between the two days' resting states, between the resting and task

states, and between the tasks. **(D)** Rest-task modulation of RVTCORR. **(E)** Spatial similarity

(ICC ± 95% CI) between GSCORR and RVTCORR for resting and task states, respectively.

Data available at DRYAD: https://doi.org/10.5061/dryad.xsj3tx9bw

Non-cortical GS components to rest-task modulation – Contributions from basal forebrain

The basal forebrain, reflecting the level of arousal, has been suggested as one of the neural origin

of the global signal [6,7]. We therefore investigate whether the rest-task modulation of GS

topography is driven by this subcortical region. We first replicated the previous findings in both

resting states and all tasks (Fig 6, top panel), by showing a negative correlation between the basal

forebrain and GS [6,7]. Extending these results, we demonstrated that their correlations were not

significantly differed during different tasks when compared to the resting state (see Fig 6 bottom

panel). This suggested that the level of arousal is stable across resting and task states, and the

rest-task modulation of GS on the cortical level, as described above, may not be driven by the

basal forebrain.

Fig 6. GSCORR at basal forebrain in rest and tasks. The maps are thresholded at Bonferroni

corrected α < 0.01. Top panel, group GSCORR comparisons with baseline (H0: r = 0). The

negative correlations are consistently seen across the two days' resting states and all task states.

Bottom panel, group GSCORR comparisons of task vs. rest show an absence of any significant

voxel in the basal forebrain. Data available at DRYAD: https://doi.org/10.5061/dryad.xsj3tx9bw

Non-cortical GS components to rest-task modulation – Contributions from physiological noise

As the GS may contain various types of noise [3], we checked whether the rest-task modulation

of GS topography was affected by head motion, respiration, cardiac, and signals from the

ventricle and white matter. We calculated the GSCORR under three procedures of noise

regression: 1) without any noise regression, 2) regressors including head motion and signals from

the ventricle and white matter as well as their first order derivatives, 3) regressors including

those in 2) plus the RVT and heart rate. If the task modulation of GSCORR (i.e., ΔGSCORR)

was driven merely by noise, then its spatial pattern, as calculated without noise, should show low

degrees of spatial similarity with the one after noise regression. We found a high spatial

similarity of the ΔGSCORR before and after nuisance noise regression (ICC > 0.95 in all cases)

(Fig 7). In addition, we observed that the inclusion of noise regression (i.e., head motion, white

matter and CSF) increased the effect size of ΔGSCORR in all seven tasks (increased percentage

in Cohen's d ranging from 22% to 39%), but was not further improved when adding respiration

and cardiac recording.

Together, these results suggested that GS topography during rest-task modulation cannot be

explained by non-neuronal components in the GS. Instead, regressing out non-neuronal

components increased the signal-to-noise ratio, and strengthened the presumed neuronal

component of GSCORR in the measured signal.

Fig 7. Rest-task modulation of GSCORR topography under different procedures of noise

regression. The maps were thresholded at Bonferroni corrected $\alpha < 0.01$. The spatial similarity of

rest-task modulation, i.e., ΔGSCORR, was measured based on the Cohen's d GSCORR

topography. The percentage of increased effect size were calculated as the increased overall |d|

value (relative to the d value from the previous step) divided by the |d| value from the previous

step. Data available at DRYAD: https://doi.org/10.5061/dryad.xsj3tx9bw

Reproducibility of the GS topography and its rest-task modulation

To examine the reliability of our findings, we calculated the GSCORR, GS-peak and the

occurrence rate of CAPs in an independent 7 Tesla dataset from HCP. This dataset included four

sessions of resting state, four sessions of movie watching, and six sessions of retinotopy stimuli

perception. Similar to the findings in the 3 Tesla dataset, we observed higher GSCORR and GS-

peak in somatosensory and visual cortex during the resting state (Fig 8A). Moreover, rest-task

modulation also showed (as seen in Figs 2B and 3C for the 3 Tesla dataset) a reduction of

GSCORR and GS-peak in the somatosensory cortex (Fig 8B). Finally, the CAPs showed a

comparable albeit smaller effect of rest-task modulation, with increased reoccurrence rate in

CAP1 and decreased reoccurrence rate in CAP2 (Fig 8C).

Fig 8. Reproducibility of the GS topography and its task modulation in HCP 7 Tesla

dataset. **(A)** Spatial topography measured by GSCORR and GS-peak in resting state and two

additional tasks (e.g., movie watching and retinotopy stimuli watching). **(B)** Grayordinate-based

group comparison for GS topography. The maps were tested with paired t-test between each task

and rest. The maps were thresholded at Bonferroni corrected α < 0.01, and illustrated by Cohen's

d. **(C)** Occurrence rates of CAPs and the difference between tasks and resting state. Data

available at DRYAD: https://doi.org/10.5061/dryad.xsj3tx9bw

Discussion

In this study, we demonstrated the informative nature of the GS through various lines of data on

rest-task modulation of its topography. We first showed that the spatial pattern of GS was

modulated during tasks with reduced contributions of somatosensory regions to GS topography.

This rest-task modulation was captured by the dynamics of GS topography in terms of transient

co-activation patterns that occur over brief epochs at the peak periods of the GS. By comparing

the spatial similarity between the GSCORR and the ones observed in relation to respiration, we

observed that the topography of respiration mimicked the topography of the GS mainly for the

resting state but significantly less so in the tasks; we therefore supposed that the GSCORR could

not equated with respiration. Finally, we ruled out the potential contributions from subcortical

areas or other sources of physiological noise on the rest-task modulation of GS topography.

GS topography – spatial distribution and rest-task modulation

We found that the GS exhibits a non-uniform topographical distribution across brain regions,

characterized by higher levels of GSCORR in primary sensory regions like auditory and visual

cortices. In contrast, we observed lower levels of GSCORR in higher-order cortical areas like the

prefrontal cortex. This replicated recent findings in both monkeys [7] and humans [3,6,11-13]. In

our study, we extended these resting state observations to the changes of GS topography in

different tasks, i.e., rest-task modulation. If the GS is subject to artifactual noise, i.e., nothing but

noise, one would expect no rest-task modulation of GS topography as the noise level may remain

the same during both resting and task states. If, in contrast, the GS reflects neuronally relevant

information, one would expect modulation effects of its topography during task states.

Our results support the latter. Specifically, we demonstrated that the GS topography was

modulated during different tasks in largely task-unspecific ways, that is, in more or less the same

manner during the seven different tasks. Moreover, we observed predominant task-unspecific

decreases of GSCORR in task irrelevant regions, e.g., somatosensory cortex, during the tasks. In

contrast, the task-related regions like the primary visual cortex (as almost of the tasks including

visual stimuli) exhibited unchanged GSCORR or, in a few regions, increase in GSCORR.

Together, we conclude that the GS does indeed provide relevant information about the brain's

neuronal activity in different states, i.e., rest and task. Therefore, we consider such rest-task

modulation as first novel line of support for the informative nature of GS and its GS topography.

Dynamics of GS topography – transient co-activation patterns

The GS topography is commonly measured by GS correlation [1,3,10], linear regression beta

weights between the GS and time series in each voxel [11,12], or whole brain functional

connectivity [14,15]. Recently, an analogous spatial pattern has also been observed in the

instantaneous co-activation pattern at the peak time points of the GS during the resting state [6].

Building on that, another study demonstrated that some co-activation patterns are phase-locked

to the stimulus onset during task states [32], Together, these findings suggest that the GS

topography is not static but rather dynamic consisting in a set of transient co-activation patterns

over brief epochs which can be modulated during task states.

In order to support the hypothesis of the dynamic nature of GS topography, we conducted the

same kind of analyses of co-activation patterns during the peak of the GS for both resting and

task states. We first found a high spatial similarity between the traditionally calculated static GS

topography and the co-activation pattern at the peaks of GS. Next, using a data driven clustering

approach, we demonstrated that the averaged co-activation pattern during the GS peak could be

decomposed into a set of finer grained co-activation patterns (see [20] for showing analogous

findings in animals). Most importantly, we showed that these dynamic co-activation patterns are

modulated during task states, in terms of a change in the frequency rates. For instance, the

occurrence rates of CAP1 increased while that of the CAP2 decreased. Summing up, our

observation of the dynamic nature of GS topography in terms of transient co-activation patterns

further support the assumption of the informative nature of GS.

Although we demonstrated that the rest-task modulation of global signal was associated with

alterations of the prevalence of CAPs, an alternative interpretation (or a potential confound) of

the results may be considered. More specifically, the 'resting state' has (trivially) no task and no

temporal structure, whereas each of the task states trivially does. This task structure may impose

temporal structure on both the GS (due to task-linked breathing and HR changes) and task-linked

BOLD responses (e.g. from visual or other stimuli associated with the task, responses such as

button presses, which are necessarily entrained to the task); both GS and task-linked BOLD

would have significant components along the vector defined by the response time course, making

them hard if not impossible to separate. Simply comparing estimates of GS during task vs rest

would be confounded by this difference in temporal structure of rest and task states. Due to the

fact that within the BOLD signal both entrainment to the temporal structure of the task and task-

linked neural activity are unavoidably correlated in time, it remains methodologically rather

challenging to decompose these two components in a meaningful way.

Subcortical and physiological substrates underlying GS topography

The GS has been shown to be closely related to physiological signals. While controlling the head

motion and scanning noise, the GS has been shown to exhibit high correlation with the

respiration rate [3,16,17,34]. We replicated the close relationship of respiration of GS by

demonstrating their spatial similarity during resting state. This suggests that the GS is indeed

coupled to respiration. Remarkably, we observed that the coupling of GS to respiration in terms

of their spatial similarity pattern was significantly decreased during all task states. However,

caveats still need to be taken for interpreting this effect. For example, breathing (and heart rate)

may be strongly entrained to task timing when the participants are engaged in a task. The GS due

to breathing (and heart rate) could thus have components that correlate or anticorrelate with the

neurally driven task related response in a relevant brain area. Therefore, the interplay among the

GS, task evoked brain activities and physiological signals may be an important question that

warrants more investigations."

Another source contributed to the GS, as observed by recent studies, is the basal forebrain as

supported by empirical data in both human [6] and macaque [7]. This region is functionally

involved with arousal or vigilance regulation [35-38]. We did not find statistically significant

difference of the rest-task modulation in the basal forebrain contributions to global signal.

However, caveats need to be taken for interpreting the null effect. That is, the absence of a rest-

task difference could not be evidence for a lack of involvement of the basal forebrain. First, the

basal forebrain is a very small structure with poor SNR [6], which may contribute to a Type II

error. Second, our analytical approach may be not sensitive enough to detect the basal forebrain's

contribution under the broad differences between resting and task states.

Nevertheless, from another angle, our results support the findings by Liu et al., 2018, in terms of

the strong negative correlation between the basal forebrain and global signal in general. We

found that the negative correlation is robust across both the resting and task states. The

seemingly invariant negative correlation may indicate that the arousal level is similar across the

resting and task states, and rest-task modulation most likely occurred on the cortical level that is

associated with cognitive processing.

Finally, the GS may also, in part, include widespread non-neuronal noise [3,21]. We therefore

checked the influence of noise regression on rest-task modulation of GS. Our data show highly

correlating and thus similar topographical distribution of rest-task GS modulation both with and

without physiological noise regression. If the GS topography were reflecting only the spatial

pattern of non-neuronal noise, one would have expected the opposite - namely, low or no

correlation in GS topography between analyses with and without physiological noise regression.

More importantly, we also demonstrated that regressing the noise can significantly improve the

signal-to-noise ratio in detecting rest-task modulation of GS topography; that further supports the

need for noise regression before calculating GS topography. Ideally, future investigations are

warranted to combine rest-task modulation in fMRI with electrophysiological investigation to

further corroborate its assumed neuronal basis.

Clinical relevance – GS topography in psychiatric disorders

The rest-task modulation of GS topography may provide a novel framework for interpreting

symptoms in psychiatric disorders. The GS topography during resting state has been investigated

in different psychiatric disorders like schizophrenia, bipolar disorders, and autism spectrum

disorders [10,11,14,15,39]. For example, the data showed altered GS topography with abnormal

GS increase in association regions in schizophrenia [11] as well as enhanced GS contribution

from motor regions in bipolar manic participants and from hippocampus/parahippocampus in

bipolar depression [10].

Albeit tentatively, GS increases and decreases in these regions may lead to corresponding

increases or decreases of the functions that are associated with the respective regions. For

instance, schizophrenia can be characterized by abnormal cognitive processing as in delusions

and formal thought disorders that are known to be related to abnormalities in that association

cortex that also shows GS increase [11]. While GS increase in motor cortex of manic subjects

may be related to increased motor function which symptomatically is manifest in psychomotor

agitation. Finally, depressed patients exhibit typically rumination with increased recollection of

autobiographical memories [10] which may be related to their increased GS in

hippocampus/parahippocampus.

Since these GS abnormalities were all observed during the resting state, one may assume that

these symptoms may neuronally be traced to virtual task states [40] in the respective regions

during resting state. If so, the psychiatric symptoms may mirror the 'normal' cognitive function

of these regions in an 'abnormal' neuronal context, that is, task-like state during the resting state.

One would then expect that, unlike in healthy subjects as in the present study, psychiatric

subjects may no longer show rest-task modulation of GS topography in specifically these regions

nor the task-related changes in the occurrence rates of their dynamic co-activation pattern.

Psychiatric symptoms may ultimately be traced to the confusion of resting and task states as

modulated by abnormalities in GS topography thus entailing what recently has been described as

"spatiotemporal psychopathology" [41-44].

Materials and Methods

Ethics statement

495   This paper utilised data collected for the HCP. The scanning protocol, participant recruitment

496   procedures, and informed written consent forms, including consent to share deidentified data,

497   were approved by the Washington University institutional review board [45].

Data Collection

We utilized the data set from the Human Connectome Project (HCP). The 3 Tesla dataset was

served for main analyses, and 7 Tesla served for replication [45,46]. All subjects from the HCP

1200-subject data release (Mar 01, 2017) having completed fMRI sessions were included (837

subjects in 3 Tesla and 145 subjects in7 Tesla, respectively). The age of the participants in the

project ranges from 22 to 35, and 54% were female.

Procedures of resting state and tasks in 3 Tesla

Data were collected over 2 days. On each day the data collection included 28 min of rest (eyes

open with fixation) fMRI data across two runs, and 30 min of task fMRI. Each of the seven tasks

was completed over two consecutive fMRI runs [47]. These tasks included seven distinct

domains: emotion, reward learning, language, motor, relational reasoning, social cognition and

working memory. Briefly, the emotion task involved matching fearful or angry faces to a target

face. The reward learning task involved a gambling task with monetary rewards and losses. The

language task involved auditory stimuli consisting of narrative stories and math problems, along

with questions to be answered regarding the prior auditory stimuli. The motor task involved

movement of the hands, tongue and feet. The relational reasoning task involved higher-order

cognitive reasoning regarding relations among features of presented shape stimuli. The social

cognition task used short video clips of moving shapes that interacted in some way or moved

randomly, with subjects making decisions about whether the shapes had social interactions. The

working memory (WM) task consisted of a visual N-back task, in which subjects indicate a

match of the current image to either a constant target image or two images previous. The details

of tasks can be found elsewhere [47].The whole-brain echo-planar imaging acquisitions were

acquired with a 32-channel head coil at a resolution of 2-mm isotropic and 0.72-s TR [48].

Procedures of resting state and tasks in 7 Tesla

Data collection in 7 Tesla contained a total of 14 fMRI runs, including 64 mins of rest, 64 mins

of movie watching, and 30 mins of retinotopy [48,49]. In each movie-watching run, subjects had

to watch a movie of approximately 15 minutes consisting several short clips separated by 20-

seconds rest periods. Different clips were used in different runs (Details are available at HCP

S1200 Release Reference). For retinotopy task, stimuli were constructed by creating slowly

moving apertures and placing a dynamic colorful texture within the apertures [50]. The data were

collected at a resolution of 1.6-mm isotropic and 1-s TR (multiband acceleration 5, in-plane

acceleration 2, 85 slices) [48].

Preprocessing

Preprocessing was carried out using Workbench [51] and custom code in MATLAB 2017b

(MathWorks). For 3 Tesla data, to match the steps of preprocessing across rest and task, the

version of HCP minimal preprocessing pipeline before FIX denoising was used (i.e., including

procedures of registration to MNI space, alignment for motion, fieldmap correction, and

MSMAll group registration[48]). Additional noise regression was applied by in-house code (see

details below). For 7 Tesla data, as the minimal preprocessed data in both rest and task had

performed FIX denoising [48], no additional noise regression was further applied.

For 3 Tesla data, the linear trend for each run were removed, and the nuisance time series

(ventricle, white matter, motions along with their first order derivatives) were regressed by using

linear regression [3,52]. The nuisance time series (ventricle, and white matter signals) were

extracted from volume-based minimal processing. No low-pass temporal filter was applied given

the possibility that frequency specificity might differ between resting and task state [52].

Preprocessing of physiological recordings

The cardiac and respiration recordings in 3 Tesla data set were analyzed. The preprocessing followed the previous studies [3,16,17,34]. More specifically, the respiratory signal was detrended linearly. The outliers were defined as the time points that deviated more than (approximately) 7 median absolute deviations (MAD) from the moving median value within a time window of 30 s. Subsequently, the respiratory signal was low-pass filtered at 5 Hz with a 2nd order Butterworth filter. The recording was then transformed into the time series of respiration volume per time (RVT), calculated as the difference between upper and lower envelope [16,17]. Outliers were replaced using linear interpolation. Finally, the RVT time series was resampled to the rate of fMRI recording (TR = 0.72 s).

The cardiac signal was initially band-pass filtered with a 2nd order Butterworth filter between 0.3 and 10 Hz. The peak was detected with minimum peak distance larger than 0.3 s. The cardiac signal was then computed in beats-per-minute (bpm) by multiplying the inverse of the time differences between pairs of adjacent peaks with 60. Outliers were corrected using the moving median method described above, and replaced using linear interpolation. Finally, the heart rate (HR) time series was resampled to the rate of fMRI (TR = 0.72 s).

Global signal correlation (GSCORR)

The global signal was calculated for each subject by averaging the standardized (z-score) fMRI signals across grayordinate. The GS topography was calculated by Pearson correlation (i.e., Global signal correlation/GSCORR) between the global signal and the time series in each

grayordinate [1,3]. The correlation r values were then transformed through fisher's z

transformation for statistical analyses [52,53].

Global signal co-activation pattern (GS-peak)

The method for calculating the GS-peak topography was adopted from previously studies [6,29].

Time points at top 17% global signal were selected, and the fMRI volumes at these time points

were averaged to generate the global signal co-activation pattern (GS-peak).

Whole brain co-activation pattern analysis (CAPs)

To investigate whether the global signal co-activation topography was a combination of several

specific recurring spatial pattern, we adopted an unsupervised machine-learning approach using

k-means clustering algorithm [30,33]. This procedure classified a set of fMRI volumes into

different categories (e.g. patterns) into k clusters based on their spatial similarity and thus

produced a set of co-activation patterns (CAP) or brain states. In such way, the original fMRI

(3D+time) data was transformed into a one-dimensional time-series of discrete CAP labels.

The analysis above was performed on the concatenated data of 3T data set, including resting

state and all tasks from all 837 subjects. To increase spatial signal to noise ratio (SNR) in local

brain regions, and make the computation achievable in this huge dataset, we performed the k-

means clustering at the region-of-interest (ROI) level [31], using a standard brain parcellation

with 1000 ROIs [27]. After clustering, the fMRI frames assigned to the same cluster were simply

averaged, resulting in k maps defined as CAPs. The CAP labels in 7T data set were assigned on

its maximal similarity to the pre-defined CAPs from 3T dataset, to make the results comparable

and generalizable across datasets [32].

Previous studies demonstrated a few recurring dominant network states explain the vast majority

of rs-fMRI temporal dynamics in both mouse [20] and human [31,33], we here inspected the

spatial patterns under 6 and 8 CAPs, and found that the 6 clusters leaded to the maximal number

of identifiable spatial patterns, and these patterns were conserved when the clusters increased to

8 (see S1 Fig).

Respiration topography (RVTCORR)

Cross-correlation over lags (-72 to 72 s) was first performed between global signal and RVT.

The cross-correlation demonstrates the global signal and RVT showed strongest negative

correlation with ~12 s lag. To make the topography of GSCORR and RVTCORR comparable,

the RVTCORR topography was calculated by Pearson correlation (i.e.,) between the flipped

RVT with 12 s lag and the time series in each grayordinate. The correlation r values were then

transformed through fisher's z transformation for statistical analyses [52,53].

Analysis of noise effect on GSCORR

To examine whether the global signal topography differed across states due to the difference of

nuisance signal contribution [3], we additionally calculated the GS topography with different

procedures of noise regression: 1) without any noise regression; 2) regressors including head

motion and signals from the ventricle and white matter as well as their first order derivatives; and

3) regressors including those in 2) plus the RVT and heart rate. The grayordinate-based

correlation for the task-rest difference was calculated to check the influence of nuisance signal

on rest-task difference [54].

Analysis of spatial similarity

To avoid the correlation induced by spatial adjacency in grayordinate-based correlation [55], the

analyses of absolute spatial similarities for GSCORR, GS-peak, and RVTCORR were performed

using intraclass correlation coefficient (ICC) [23], across the aforementioned 1000 ROIs at the

group level, with spatial normalization (z-score). In reporting these findings, the ICC was

categorized into five common intervals[26]: $0 < ICC \leq 0.2$ (slight); $0.2 < ICC \leq 0.4$ (fair); $0.4 <$

$ICC \leq 0.6$ (moderate); $0.6 < ICC \leq 0.8$ (substantial); and $0.8 < ICC \leq 1.0$ (almost perfect). The

significant difference between two ICCs was checked by fisher's z test [56].

Statistics

All statistical inferences for topography were based on two-tailed paired t-tests. In the current

study, we mainly report results with effect size using Cohen's d [57] instead of the t value, as

Cohen's d is independent of the sample size. The threshold for the differences in brain maps was

set at Bonferroni corrected $\alpha < 0.01$ across the whole brain.

Reference

1.   Fox MD, Zhang D, Snyder AZ, Raichle ME. The global signal and observed anticorrelated resting
state brain networks. J Neurophysiol. 2009;101(6):3270-83. Epub 2009/04/03. doi:
10.1152/jn.90777.2008. PubMed PMID: 19339462; PubMed Central PMCID: PMCPMC2694109.
2.   Murphy K, Birn RM, Handwerker DA, Jones TB, Bandettini PA. The impact of global signal
regression on resting state correlations: are anti-correlated networks introduced? Neuroimage.
2009;44(3):893-905.
3.   Power JD, Plitt M, Laumann TO, Martin A. Sources and implications of whole-brain fMRI signals
in humans. Neuroimage. 2017;146:609-25. Epub 2016/10/19. doi: 10.1016/j.neuroimage.2016.09.038.
PubMed PMID: 27751941; PubMed Central PMCID: PMCPMC5321814.
4.   Schölvinck ML, Maier A, Frank QY, Duyn JH, Leopold DA. Neural basis of global resting-state fMRI
activity. Proceedings of the National Academy of Sciences. 2010;107(22):10238-43.
5.   Lv Y, Margulies DS, Cameron Craddock R, Long X, Winter B, Gierhake D, et al. Identifying the
perfusion deficit in acute stroke with resting-state functional magnetic resonance imaging. Annals of
neurology. 2013;73(1):136-40.
6.   Liu X, de Zwart JA, Scholvinck ML, Chang C, Ye FQ, Leopold DA, et al. Subcortical evidence for a
contribution of arousal to fMRI studies of brain activity. Nat Commun. 2018;9(1):395. Epub 2018/01/28.
doi: 10.1038/s41467-017-02815-3. PubMed PMID: 29374172; PubMed Central PMCID:
PMCPMC5786066.
7.   Turchi J, Chang C, Ye FQ, Russ BE, Yu DK, Cortes CR, et al. The Basal Forebrain Regulates Global
Resting-State fMRI Fluctuations. Neuron. 2018;97(4):940-52 e4. Epub 2018/02/06. doi:
10.1016/j.neuron.2018.01.032. PubMed PMID: 29398365; PubMed Central PMCID: PMCPMC5823771.
8.   Orban C, Kong R, Li J, Chee MW, Yeo BT. Time of day is associated with paradoxical reductions in
global signal fluctuation and functional connectivity. PLoS Biology. 2020;18(2):e3000602.
9.   Yang GJ, Murray JD, Repovs G, Cole MW, Savic A, Glasser MF, et al. Altered global brain signal in
schizophrenia. Proc Natl Acad Sci U S A. 2014;111(20):7438-43. Epub 2014/05/07. doi:
10.1073/pnas.1405289111. PubMed PMID: 24799682; PubMed Central PMCID: PMCPMC4034208.
10.   Zhang J, Magioncalda P, Huang Z, Tan Z, Hu X, Hu Z, et al. Altered Global Signal Topography and
Its Different Regional Localization in Motor Cortex and Hippocampus in Mania and Depression.
Schizophr Bull. 2019;45(4):902-10. Epub 2018/10/05. doi: 10.1093/schbul/sby138. PubMed PMID:
30285255; PubMed Central PMCID: PMCPMC6581125.
11.   Yang GJ, Murray JD, Glasser M, Pearlson GD, Krystal JH, Schleifer C, et al. Altered Global Signal
Topography in Schizophrenia. Cereb Cortex. 2017;27(11):5156-69. Epub 2016/10/06. doi:
10.1093/cercor/bhw297. PubMed PMID: 27702810; PubMed Central PMCID: PMCPMC6075538.
12.   Li J, Bolt T, Bzdok D, Nomi JS, Yeo BTT, Spreng RN, et al. Topography and behavioral relevance of
the global signal in the human brain. Sci Rep. 2019;9(1):14286. Epub 2019/10/05. doi: 10.1038/s41598-
019-50750-8. PubMed PMID: 31582792; PubMed Central PMCID: PMCPMC6776616.
13.   Wen H, Liu Z. Broadband Electrophysiological Dynamics Contribute to Global Resting-State fMRI
Signal. J Neurosci. 2016;36(22):6030-40. Epub 2016/06/03. doi: 10.1523/JNEUROSCI.0187-16.2016.
PubMed PMID: 27251624; PubMed Central PMCID: PMCPMC4887567.
14.   Gotts SJ, Saad ZS, Jo HJ, Wallace GL, Cox RW, Martin A. The perils of global signal regression for
group comparisons: a case study of Autism Spectrum Disorders. Front Hum Neurosci. 2013;7:356. Epub
2013/07/23. doi: 10.3389/fnhum.2013.00356. PubMed PMID: 23874279; PubMed Central PMCID:
PMCPMC3709423.

15.   Gotts SJ, Simmons WK, Milbury LA, Wallace GL, Cox RW, Martin A. Fractionation of social brain
circuits in autism spectrum disorders. Brain. 2012;135(Pt 9):2711-25. Epub 2012/07/14. doi:
10.1093/brain/aws160. PubMed PMID: 22791801; PubMed Central PMCID: PMCPMC3437021.
16.   Birn RM, Diamond JB, Smith MA, Bandettini PA. Separating respiratory-variation-related
fluctuations from neuronal-activity-related fluctuations in fMRI. Neuroimage. 2006;31(4):1536-48. Epub
2006/04/25. doi: 10.1016/j.neuroimage.2006.02.048. PubMed PMID: 16632379.
17.   Birn RM, Smith MA, Jones TB, Bandettini PA. The respiration response function: the temporal
dynamics of fMRI signal fluctuations related to changes in respiration. Neuroimage. 2008;40(2):644-54.
Epub 2008/02/01. doi: 10.1016/j.neuroimage.2007.11.059. PubMed PMID: 18234517; PubMed Central
PMCID: PMCPMC2533266.
18.   Wise RG, Ide K, Poulin MJ, Tracey I. Resting fluctuations in arterial carbon dioxide induce
significant low frequency variations in BOLD signal. Neuroimage. 2004;21(4):1652-64. Epub 2004/03/31.
doi: 10.1016/j.neuroimage.2003.11.025. PubMed PMID: 15050588.
19.   Liu TT, Nalci A, Falahpour M. The global signal in fMRI: Nuisance or Information? Neuroimage.
2017;150:213-29. Epub 2017/02/19. doi: 10.1016/j.neuroimage.2017.02.036. PubMed PMID: 28213118;
PubMed Central PMCID: PMCPMC5406229.
20.   Gutierrez-Barragan D, Basson MA, Panzeri S, Gozzi A. Infraslow State Fluctuations Govern
Spontaneous fMRI Network Dynamics. Curr Biol. 2019;29(14):2295-306 e5. Epub 2019/07/16. doi:
10.1016/j.cub.2019.06.017. PubMed PMID: 31303490; PubMed Central PMCID: PMCPMC6657681.
21.   Murphy K, Fox MD. Towards a consensus regarding global signal regression for resting state
functional connectivity MRI. Neuroimage. 2017;154:169-73. Epub 2016/11/27. doi:
10.1016/j.neuroimage.2016.11.052. PubMed PMID: 27888059; PubMed Central PMCID:
PMCPMC5489207.
22.   Zuo XN, Xu T, Milham MP. Harnessing reliability for neuroscience research. Nat Hum Behav.
2019;3(8):768-71. Epub 2019/06/30. doi: 10.1038/s41562-019-0655-x. PubMed PMID: 31253883.
23.   Koo TK, Li MY. A guideline of selecting and reporting intraclass correlation coefficients for
reliability research. Journal of chiropractic medicine. 2016;15(2):155-63.
24.   Zuo X-N, Biswal BB, Poldrack RA. Reliability and Reproducibility in Functional Connectomics.
Frontiers in neuroscience. 2019;13:117.
25.   Chen B, Xu T, Zhou C, Wang L, Yang N, Wang Z, et al. Individual variability and test-retest
reliability revealed by ten repeated resting-state brain scans over one month. PLoS One. 2015;10(12).
26.   Landis JR, Koch GG. The measurement of observer agreement for categorical data. biometrics.
1977:159-74.
27.   Schaefer A, Kong R, Gordon EM, Laumann TO, Zuo XN, Holmes AJ, et al. Local-Global Parcellation
of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. Cereb Cortex.
2018;28(9):3095-114. Epub 2017/10/06. doi: 10.1093/cercor/bhx179. PubMed PMID: 28981612;
PubMed Central PMCID: PMCPMC6095216.
28.   Cohen J. Statistical power analysis. Current directions in psychological science. 1992;1(3):98-101.
29.   Liu X, Duyn JH. Time-varying functional network information extracted from brief instances of
spontaneous brain activity. Proc Natl Acad Sci U S A. 2013;110(11):4392-7. Epub 2013/02/27. doi:
10.1073/pnas.1216856110. PubMed PMID: 23440216; PubMed Central PMCID: PMCPMC3600481.
30.   Liu X, Zhang N, Chang C, Duyn JH. Co-activation patterns in resting-state fMRI signals.
Neuroimage. 2018;180(Pt B):485-94. Epub 2018/01/23. doi: 10.1016/j.neuroimage.2018.01.041.
PubMed PMID: 29355767; PubMed Central PMCID: PMCPMC6082734.
31.   Chen JE, Chang C, Greicius MD, Glover GH. Introducing co-activation pattern metrics to quantify
spontaneous brain network dynamics. Neuroimage. 2015;111:476-88. Epub 2015/02/11. doi:
10.1016/j.neuroimage.2015.01.057. PubMed PMID: 25662866; PubMed Central PMCID:
PMCPMC4386757.

32. Huang Z, Zhang J, Wu J, Mashour GA, Hudetz AG. Temporal circuit of macroscale dynamic brain
activity supports human consciousness. Science Advances. 2020;6(11):eaaz0087. doi:
10.1126/sciadv.aaz0087.
33. Huang Z, Zhang J, Wu J, Mashour G, Hudetz A. Temporal circuit of macroscale dynamic brain
activity supports human consciousness. Science advances. 2020.
34. Kassinopoulos M, Mitsis GD. Identification of physiological response functions to correct for
fluctuations in resting-state fMRI related to heart rate and respiration. Neuroimage. 2019;202:116150.
Epub 2019/09/06. doi: 10.1016/j.neuroimage.2019.116150. PubMed PMID: 31487547.
35. Wong CW, DeYoung PN, Liu TT. Differences in the resting-state fMRI global signal amplitude
between the eyes open and eyes closed states are related to changes in EEG vigilance. Neuroimage.
2016;124:24-31.
36. Wong CW, Olafsson V, Tal O, Liu TT. Anti-correlated networks, global signal regression, and the
effects of caffeine in resting-state functional MRI. Neuroimage. 2012;63(1):356-64. Epub 2012/06/30.
doi: 10.1016/j.neuroimage.2012.06.035. PubMed PMID: 22743194; PubMed Central PMCID:
PMCPMC3444518.
37. Wong CW, Olafsson V, Tal O, Liu TT. The amplitude of the resting-state fMRI global signal is
related to EEG vigilance measures. Neuroimage. 2013;83:983-90.
38. Chang C, Leopold DA, Scholvinck ML, Mandelkow H, Picchioni D, Liu X, et al. Tracking brain
arousal fluctuations with fMRI. Proc Natl Acad Sci U S A. 2016;113(16):4518-23. Epub 2016/04/07. doi:
10.1073/pnas.1520613113. PubMed PMID: 27051064; PubMed Central PMCID: PMCPMC4843437.
39. Wang X, Liao W, Han S, Li J, Zhang Y, Zhao J, et al. Altered dynamic global signal topography in
antipsychotic-naive adolescents with early-onset schizophrenia. Schizophrenia research. 2019;208:308-
16.
40. Northoff G, Qin P, Nakao T. Rest-stimulus interaction in the brain: a review. Trends Neurosci.
2010;33(6):277-84. Epub 2010/03/17. doi: 10.1016/j.tins.2010.02.006. PubMed PMID: 20226543.
41. Northoff G. Spatiotemporal psychopathology I: No rest for the brain's resting state activity in
depression? Spatiotemporal psychopathology of depressive symptoms. J Affect Disord. 2016;190:854-
66. Epub 2015/06/07. doi: 10.1016/j.jad.2015.05.007. PubMed PMID: 26048657.
42. Northoff G. Spatiotemporal Psychopathology II: How does a psychopathology of the brain's
resting state look like? Spatiotemporal approach and the history of psychopathology. J Affect Disord.
2016;190:867-79. Epub 2015/06/15. doi: 10.1016/j.jad.2015.05.008. PubMed PMID: 26071797.
43. Northoff G, Duncan NW. How do abnormalities in the brain's spontaneous activity translate into
symptoms in schizophrenia? From an overview of resting state activity findings to a proposed
spatiotemporal psychopathology. Prog Neurobiol. 2016;145-146:26-45. Epub 2016/08/18. doi:
10.1016/j.pneurobio.2016.08.003. PubMed PMID: 27531135.
44. Northoff G, Magioncalda P, Martino M, Lee HC, Tseng YC, Lane T. Too Fast or Too Slow? Time
and Neuronal Variability in Bipolar Disorder-A Combined Theoretical and Empirical Investigation.
Schizophr Bull. 2018;44(1):54-64. Epub 2017/05/20. doi: 10.1093/schbul/sbx050. PubMed PMID:
28525601; PubMed Central PMCID: PMCPMC5768053.
45. Van Essen DC, Smith SM, Barch DM, Behrens TE, Yacoub E, Ugurbil K, et al. The WU-Minn
Human Connectome Project: an overview. Neuroimage. 2013;80:62-79. Epub 2013/05/21. doi:
10.1016/j.neuroimage.2013.05.041. PubMed PMID: 23684880; PubMed Central PMCID:
PMCPMC3724347.
46. Glasser MF, Smith SM, Marcus DS, Andersson JL, Auerbach EJ, Behrens TE, et al. The Human
Connectome Project's neuroimaging approach. Nat Neurosci. 2016;19(9):1175-87. Epub 2016/08/30.
doi: 10.1038/nn.4361. PubMed PMID: 27571196; PubMed Central PMCID: PMCPMC6172654.
47. Barch DM, Burgess GC, Harms MP, Petersen SE, Schlaggar BL, Corbetta M, et al. Function in the
human connectome: task-fMRI and individual differences in behavior. Neuroimage. 2013;80:169-89.

Epub 2013/05/21. doi: 10.1016/j.neuroimage.2013.05.033. PubMed PMID: 23684877; PubMed Central
PMCID: PMCPMC4011498.
48.   Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, et al. The minimal
preprocessing pipelines for the Human Connectome Project. Neuroimage. 2013;80:105-24. Epub
2013/05/15. doi: 10.1016/j.neuroimage.2013.04.127. PubMed PMID: 23668970; PubMed Central
PMCID: PMCPMC3720813.
49.   Vu AT, Jamison K, Glasser MF, Smith SM, Coalson T, Moeller S, et al. Tradeoffs in pushing the
spatial resolution of fMRI for the 7T Human Connectome Project. Neuroimage. 2017;154:23-32.
50.   Benson NC, Jamison KW, Arcaro MJ, Vu AT, Glasser MF, Coalson TS, et al. The Human
Connectome Project 7 Tesla retinotopy dataset: Description and population receptive field analysis. J
Vis. 2018;18(13):23. Epub 2018/12/29. doi: 10.1167/18.13.23. PubMed PMID: 30593068; PubMed
Central PMCID: PMCPMC6314247.
51.   Marcus DS, Harwell J, Olsen T, Hodge M, Glasser MF, Prior F, et al. Informatics and data mining
tools and strategies for the human connectome project. Front Neuroinform. 2011;5:4. Epub
2011/07/12. doi: 10.3389/fninf.2011.00004. PubMed PMID: 21743807; PubMed Central PMCID:
PMCPMC3127103.
52.   Cole MW, Ito T, Bassett DS, Schultz DH. Activity flow over resting-state networks shapes
cognitive task activations. Nat Neurosci. 2016;19(12):1718-26. Epub 2016/11/01. doi: 10.1038/nn.4406.
PubMed PMID: 27723746; PubMed Central PMCID: PMCPMC5127712.
53.   Cole MW, Bassett DS, Power JD, Braver TS, Petersen SE. Intrinsic and task-evoked network
architectures of the human brain. Neuron. 2014;83(1):238-51. Epub 2014/07/06. doi:
10.1016/j.neuron.2014.05.014. PubMed PMID: 24991964; PubMed Central PMCID: PMCPMC4082806.
54.   Zhang J, Huang Z, Chen Y, Zhang J, Ghinda D, Nikolova Y, et al. Breakdown in the temporal and
spatial organization of spontaneous brain activity during general anesthesia. Hum Brain Mapp.
2018;39(5):2035-46. Epub 2018/01/30. doi: 10.1002/hbm.23984. PubMed PMID: 29377435.
55.   Jiang L, Zuo XN. Regional Homogeneity: A Multimodal, Multiscale Neuroimaging Marker of the
Human Connectome. Neuroscientist. 2016;22(5):486-505. Epub 2015/07/15. doi:
10.1177/1073858415595004. PubMed PMID: 26170004; PubMed Central PMCID: PMCPMC5021216.
56.   Fisher RA. On the'probable error'of a coefficient of correlation deduced from a small sample.
Metron. 1921;1:1-32.
57.   Cohen J. Statistical power analysis for the behavioral sciences 2nd edn. Erlbaum Associates,
Hillsdale; 1988.
802

Supporting information

S1 Fig. Transient co-activation patterns (CAPs) with 8 clusters. *A.* spatial topography of CAPs and their occurrence rates at the time points of GS-peak. Task modulation denoted the difference in the occurrence rate of the CAP's between task and resting state (Δrate). *B.* Spatial correlation between the CAPs. The eight CAPs were composed of four pairs of opposite co-activation patterns (i.e., CAP1 vs. CAP5, CAP2 vs. CAP6, CAP3 vs. CAP7, and CAP4 vs. CAP8), as denoted by their negative correlations. Data available at DRYAD: https://doi.org/10.5061/dryad.xsj3tx9bw

S2 Fig. Transient co-activation patterns (CAPs) across two days' rests and seven tasks. Top panel yielded the co-activation patterns in Fig 3. Bottom panel yielded the occurrence rate across resting state and seven tasks. Data available at DRYAD: https://doi.org/10.5061/dryad.xsj3tx9bw

A

Global signal → Pearson correlation → Time series in each region of gray matter → GSCORR map

B

Spatial Topography of GSCORR

REST   TASK

REST1   REST2   WM   GAMBLING   MOTOR   LANGUAGE   SOCIAL   RELATIONAL   EMOTION

C

Spatial Similarity of GSCORR

	REST1	REST2	WM	GAMBLING	MOTOR	LANGUAGE	SOCIAL	RELATIONAL	EMOTION
REST1	1.00	0.99	0.82	0.85	0.89	0.80	0.82	0.81	0.89
REST2	0.99	1.00	0.81	0.85	0.88	0.80	0.82	0.80	0.88
WM	0.82	0.81	0.99	0.97	0.94	0.94	0.90	0.96	0.97
GAMBLING	0.85	0.85	0.97	1.00	0.95	0.94	0.92	0.95	0.98
MOTOR	0.89	0.88	0.94	0.95	1.00	0.95	0.89	0.90	0.96
LANGUAGE	0.80	0.80	0.94	0.94	0.95	1.00	0.82	0.88	0.93
SOCIAL	0.82	0.82	0.90	0.92	0.89	0.82	0.99	0.91	0.93
RELATIONAL	0.81	0.80	0.96	0.95	0.90	0.88	0.91	1.00	0.96
EMOTION	0.89	0.88	0.97	0.98	0.96	0.93	0.93	0.96	1.00

ICC: 0 to 1

REST-REST: 0.99   REST-TASK: 0.86   TASK-TASK: 0.93

*** *** ***

A Mean of GSCORR

B ΔGSCORR (Task – Rest, Bonferroni corrected α < 0.01)

WM   GAMBLING   MOTOR

LANGUAGE   SOCIAL   RELATIONAL   EMOTION

C Overlapped Map Across 7 Tasks (Task – Rest)

Decrease   Unchanged   Increase

A

Averaging volumes at the peaks (e.g., top 17%) of global signal

GS-peak map

Global signal

B

Spatial Topography of GS-peak

REST   TASK

0.35

BOLD

REST1   WM   GAMBLING   MOTOR

REST2   LANGUAGE   SOCIAL   RELATIONAL   EMOTION

C

ΔGS-peak (Task – Rest, Bonferroni corrected α < 0.01)

WM   GAMBLING   MOTOR

Cohen's d

-1

LANGUAGE   SOCIAL   RELATIONAL   EMOTION

D

Spatial Similarity between GSCORR and GS-peak

GSCORR	REST1	REST2	WM	GAMBLING	MOTOR	LANGUAGE	SOCIAL	RELATIONAL	EMOTION
REST1	0.99	0.99	0.83	0.88	0.89	0.82	0.84	0.83	0.90
REST2	0.99	0.99	0.82	0.87	0.89	0.82	0.83	0.83	0.89
WM	0.79	0.78	0.99	0.97	0.93	0.93	0.92	0.96	0.96
GAMBLING	0.83	0.82	0.96	0.99	0.94	0.94	0.93	0.95	0.98
MOTOR	0.88	0.87	0.94	0.96	0.99	0.95	0.91	0.91	0.96
LANGUAGE	0.78	0.78	0.94	0.95	0.94	0.99	0.85	0.89	0.93
SOCIAL	0.81	0.80	0.88	0.92	0.88	0.82	0.99	0.91	0.92
RELATIONAL	0.79	0.78	0.94	0.94	0.90	0.87	0.92	0.99	0.95
EMOTION	0.87	0.86	0.95	0.98	0.95	0.92	0.94	0.96	0.99

GS-peak

0   ICC   1

A

Step 1: Clustering Frames into Co-activation Patterns (CAPs)

Time →

Subject s ↓

k-means →

Frame averaging →

CAP 1
CAP 2
⋮
CAP 5
CAP 6

Step 2: Decomposing GS-peak into A Combination of CAPs

Counting occurrence rates at GS-peak

CAP 1   CAP 2 ⋯ CAP 5   CAP 6

Global signal

B

CAP 1
Δrates: 15.6%

CAP 6
Δrates: -4.0%

CAP 2
Δrates: -13.4%

CAP 5
Δrates: 2.7%

CAP 3
Δrates: -3.5%

CAP 4
Δrates: 2.6%

40%
30%
20%
10%

— REST
— TASK

-0.15 ——— 0.15
BOLD

C   **Spatial Correlation between CAPs**

CAPs	1	2	3	4	5	6
1	1.00	-0.04	-0.42	-0.99	0.00	0.28
2	-0.04	1.00	-0.08	0.03	-0.96	0.12
3	-0.42	-0.08	1.00	0.32	0.18	-0.97
4	-0.99	0.03	0.32	1.00	-0.02	-0.16
5	0.00	-0.96	0.18	-0.02	1.00	-0.27
6	0.28	0.12	-0.97	-0.16	-0.27	1.00

D   **Spatial Correlation between CAPs and ΔGSCORR**

CAPs	1	2	3	4	5	6
WM	0.39	-0.74	0.12	-0.43	0.78	-0.24
GAMBLING	0.33	-0.72	0.05	-0.35	0.73	-0.15
MOTOR	0.16	-0.83	-0.08	-0.15	0.79	0.04
LANGUAGE	-0.06	-0.83	0.23	0.04	0.81	-0.25
SOCIAL	0.55	-0.35	-0.26	-0.57	0.40	0.12
RELATIONAL	0.68	-0.50	-0.15	-0.71	0.53	0.01
EMOTION	0.29	-0.67	0.06	-0.32	0.68	-0.15
AVERAGE	0.33	-0.66	-0.01	-0.36	0.67	-0.09

r = -1 ——— r = 1

A

Step 1: Cross-correlation between GS and RVT

Step 2: Calculating RVTCORR map

REST
TASK

12 sec

r

time (s)

-RVT with 12s lag

Time series in each region of gray matter

Pearson correlation

RVTCORR map

B **Spatial Topography of RVTCORR**

REST

TASK

RVTCORR (r)

REST1

REST2

WM   GAMBLING   MOTOR

LANGUAGE   SOCIAL   RELATIONAL   EMOTION

C **Spatial Similarity of RVTCORR**

ICC

0.99   0.55   0.40

REST-REST   REST-TASK   TASK-TASK

D **Δ RVTCORR (Task – Rest, Bonferroni corrected α < 0.01)**

WM   GAMBLING   MOTOR

LANGUAGE   SOCIAL   RELATIONAL   EMOTION

Cohen's d

E **Spatial Similarity between RVTCORR and GSCORR**

ICC

0.86   0.57

REST   TASK

GSCORR at Basal Forebrain

ΔGSCORR (Task − Rest , Bonferroni corrected α < 0.01)

A Spatial Topography

GSCORR

Rest   Movie   Retinotopy

r   0.3   0

GS-peak

BOLD   0.45   0

B Task – Rest (Bonferroni corrected α < 0.01)

GSCORR

Movie   Retinotopy

Cohen's d   1   -1

GS-peak

Cohen's d   1   -1

C

CAP 1

Δ rate
Movie: 5.3%
Retinotopy: 4.8%

CAP 6

Δ rate
Movie: -4.2%
Retinotopy: -1.2%

CAP 2

Δ rate
Movie: -4.4%
Retinotopy: -8.3%

CAP 5

Δ rate
Movie: 0.6%
Retinotopy: 7.9%

CAP 3

Δ rate
Movie: -0.37%
Retinotopy: -3.8%

40%
30%
20%
10%

- Rest
- Movie
- Retinotopy

Δ rate
Movie: 3.1%
Retinotopy: 0.58%

CAP 4

BOLD   -0.15   0.15