Running title (optional): **Choosing the optimal population for GWAS**

Core ideas (3-5 impact statements, 85 char max for each)

- Genome-wide association studies with mixture populations are expected to improve the

detection power of novel genes due to the increase of the sample size although the influence of

population structure is a concern.

- When a quantitative trait nucleotide (QTN) is polymorphic in a target population, a

combination of the target population and a population with higher diversity than the target

population improves the detection power of the QTN.

- We found that the fixation index (F_{ST}) and the expected heterozygosity (H_e) were strongly

related to the detection power of QTNs.

- Germplasm collections which have been already sequenced/genotyped are useful for improving

the detection power of GWAS without any addition of sequence costs by using a subset of them

with a target population.

Choosing the optimal population for a genome-wide association study: a simulation using whole-genome sequences from rice

Kosuke Hamazaki, Hiromi Kajiya-Kanegae, Masanori Yamasaki, Kaworu Ebana, Shiori Yabe, Hiroshi Nakagawa and Hiroyoshi Iwata*

Affiliations:

K. Hamazaki, H. Kajiya-Kanegae and H. Iwata, Department of Agricultural and Environmental Biology, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan; H. Kajiya-Kanegae, current address: Research Center for Agricultural Information Technology, National Agriculture and Food Research Organization, 3-1-1 Kannondai, Tsukuba, Ibaraki 305-8517, Japan; M. Yamasaki, Food Resources Education and Research Center, Graduate School of Agricultural Science, Kobe University, 1348 Uzurano, Kasai, Hyogo 675-2103, Japan; K. Ebana, Genetic Resources Center, National Agriculture and Food Research Organization, 2-1-2 Kannondai, Tsukuba, Ibaraki 305-8602, Japan; S. Yabe, Institute of Crop Science, National Agriculture and Food Research Organization, 2-1-2 Kannondai, Tsukuba, Ibaraki 305-8518, Japan; H. Nakagawa, Institute for Agro-Environmental Sciences, National Agriculture and Food Research Organization, 3-1-3 Kannondai, Tsukuba, Ibaraki 305-8604, Japan. *Corresponding author (aiwata@mail.ecc.u-tokyo.ac.jp)

Abbreviations:

AUC, area under the curve; CDR, correct detection rate; FDR, false discovery rate; FN, false

negative FP, false positive; F_{ST}, the fixation index; GWAS, genome-wide association study; H,

high; HM, higher-middle; H_e, the expected heterozygosity; L, low; LD, linkage disequilibrium;

LM, lower-middle; M, middle MAF, minor allele frequency QTL, quantitative trait loci; QTN,

quantitative trait nucleotide; ROC, receiver operating characteristic; SNP, single nucleotide

polymorphism; TN, true negative; TP, true positive.

44   **ABSTRACT**

A genome-wide association study (GWAS) needs to have a suitable population. The

factors that affect a GWAS, e.g. population structure, sample size, and sequence analysis and

field testing costs, need to be considered. Mixture populations containing subpopulations of

different genetic backgrounds may be suitable populations. We conducted simulation

experiments to see if a population with high genetic diversity, e.g., a diversity panel, should be

added to a target population, especially when the target population harbors small genetic

diversity. The target population was 112 accessions of *Oryza sativa* subsp. *japonica*, mainly

developed in Japan. We combined the target population with three populations that had higher

genetic diversities. These were 100 *indica* accessions, 100 *japonica* accessions, and 100

accessions with various genetic backgrounds. The results showed that the GWAS power with a

mixture population was generally higher than with a separate population. Also, the GWAS

optimal population varied depending on the fixation index F_{ST} of the quantitative trait nucleotide

(QTN) and its polymorphism of QTN in each population. When a QTN is polymorphic in a

target population, a target population combined with a higher diversity population improves the

QTN detection power. Investigating F_{ST} and the expected heterozygosity H_e as factors

influencing the detection power, we showed that SNPs with high F_{ST} or low H_e are less likely to

be detected by GWAS with mixture populations. Sequenced/genotyped germplasm collections

can improve the GWAS detection power by using a subset of them with a target population.

INTRODUCTION

Recently, as genome sequencing costs have continued to decrease (Metzker, 2010), the

whole-genome sequences of a large number of cultivars/lines have become available for major

crop species, such as rice (Li et al., 2014; Wang et al., 2018). A genome-wide association study

(GWAS) based on whole-genome sequences can more efficiently and accurately identify genes

that control important agronomic traits than previous methods (Koboldt et al., 2013; Ott et al.,

2015; Yano et al., 2016).

It is important to prepare an appropriate population to be analyzed when attempting to detect

candidate genes using GWAS techniques. For example, to avoid potential false positives caused

by population stratification/structure, a GWAS population should be selected that results in low

stratification (Begum et al., 2015; Yano et al., 2016). However, if such a population is selected as

an analytical population for a GWAS, the sample size may be limited and the detection power of

the GWAS will decrease (Korte and Farlow, 2013). Therefore, when designing an appropriate

GWAS population, one should be aware of the trade-off relationship between population

stratification and sample size.

When preparing the population to be analyzed, the factors that directly affect the GWAS

results, such as population structure, sample size, and the sequence analysis and cultivation

testing costs, need to be considered. In recent years, the whole-genome sequences of a large

number of cultivars/lines have become publicly available due to highly efficient sequencing

analyses and database enrichment. The publically available whole-genome sequence data will

improve GWASs and could enable researchers to avoid the costs of sequencing analyses. For

example, in rice, "The 3,000 Rice Genomes Project" (Li et al., 2014; Wang et al., 2018) by the

International Rice Research Institute (IRRI) is a well-known whole-genome sequence dataset

that is available in the "Rice SNP-Seek Database" (Alexandrov et al., 2015; Mansueto et al.,

2016; 2017). Therefore, an appropriate GWAS population could potentially utilize existing

public sequence data.

A mixture population obtained by mixing subpopulations with different genetic backgrounds

could also potentially be used in a GWAS. An advantage of using such a mixture population is

that it should improve the detection of causal variants by increasing the sample size. Conversely,

a GWAS with a mixture population may suffer from large numbers of false positives caused by

the population structure. Although a mixed effect model that suppresses the influence of the

population structure has been proposed (Yu et al., 2006), such a mixture population has rarely

been analyzed by a GWAS.

An actual data analysis of rice using whole-genome sequences showed that the detection

power of a GWAS improved when *Oryza sativa* subsp. *japonica* and *Oryza sativa* subsp. *indica*

populations were combined (Misra et al., 2017). Furthermore, the identification of new rice

genes using a GWAS and populations with extremely high genetic diversities has also been

previously reported (Zhao et al., 2011). Conversely, it has been reported that the genetic

differentiation between subpopulations in a population with high genetic diversity could cause a

reduction in the power of a GWAS (Huang et al., 2012). Therefore, real data studies have been

inconsistent about whether mixture populations or populations with high genetic diversities

should be used in a GWAS. However, these previous studies mostly analyzed actual data, and

there have been no theoretical simulation studies that have considered the possibility of using a

mixture population in a GWAS. Furthermore, no previous studies have discussed which kinds of

populations should be mixed to improve the GWAS detection power or which kinds of

populations are most appropriate for a GWAS. Therefore, in this study, we conducted simulation

experiments to see whether adding a population with a high genetic diversity compared to a

target population (e.g., adding a diversity panel to a target population) is appropriate, especially

when the genetic diversity of the target population is small.

MATERIALS AND METHODS

Materials (populations used in the GWAS)

In this study, 112 accessions of *Oryza sativa* subsp. *japonica* (referred to as "A"), which

were accessions that had mainly been developed in Japan, were used as a target population with

low genetic diversity (Yabe et al., 2016). We used the following three populations selected from

"The 3,000 Rice Genomes Project" (J. Y. Li et al., 2014), i.e., 100 accessions of *Oryza sativa*

subsp. *indica* (referred to as "B"), 100 accessions of *Oryza sativa* subsp. *japonica* (referred to as

"C" or temperate), and 100 accessions of *Oryza sativa* with various genetic backgrounds

(referred to as "D" or diverse), as populations with higher diversities than the target population

(Table S1 in Supplemental File 1). The process used to select each of the 100 accessions is

described in Supplemental File 9. One accession (IRIS ID: IRIS 313-11868) was duplicated in

populations B and D. Among populations B, C, and D, the B population was the most

differentiated from A, whereas C was the most similar to A. Population D contained subsp.

indica, subsp. *japonica*, and *aus*, and *aromatic* rice accessions, which meant that the D

population had the highest genetic diversity. Fig. 1 is an unrooted phylogenetic tree that shows

the genetic relationships among accessions belonging to populations A, B, C, and D.

Fig. 1. **Unrooted phylogenetic tree plot for four non-mixture populations.**

Unrooted phylogenetic tree plot for the four non-mixture populations, which consisted of 112

accessions of *japonica* (A), 100 accessions of *indica* (B), 100 accessions of temperate *japonica*

(C), and 100 diverse accessions (D) with neighbor-joining method.

The genetic relationships among the accessions were estimated by the neighbor-joining (NJ)

method (Saitou and Nei, 1987) using the R package "ape" version 5.3 (Paradis et al., 2004). The

genetic distances were estimated according to the Jukes and Cantor (1969) model. In addition to

these four populations, we synthesized three populations by combining population A with

populations B, C, or D. The mixture populations A + B, A + C, and A + D were named "E", "F",

and "G", respectively. We compared the QTN detection power the GWAS when the seven non-

mixture (A, B, C, and D) and mixture populations (E, F, and G) were used.

Genotype data

Whole genome sequencing data were available for the accessions (Jarquin et al., 2019).

Details about the DNA extraction and whole genome sequencing techniques are provided in a

previous report (Jarquin et al., 2019). The data sets deposited in the DDBJ Sequence Read

Archive (SRA106223, ERA358140, DRA000158, DRA000307, DRA000897, DRA000927,

DRA007273, DRA007256, and DRA008071) were reanalyzed. We processed the whole-genome

sequence data as follows so that they could be used in the GWAS. Adapters and low-quality

bases were removed from paired reads using the Trimmomatic v0.36 program (Bolger et al.,

2014). The preprocessed reads were aligned using Os-Nipponbare-Reference-IRGSP-1.0

(Kawahara et al., 2013) and the bwa-0.7.12 mem algorithm with the default options (H. Li,

2012). The binary alignment map (BAM) files deposited in the Rice SNP-Seek database were

also reanalyzed. Single nucleotide polymorphism (SNP) calling was based on alignments

determined using the Genome Analysis Toolkit (GATK), 3.7-0-gcfedb67 (McKenna et al., 2009;

Auwera et al., 2014) and Picard package V2.5.0 (http://broadinstitute.github.io/picard). The

mapped reads were realigned using RealignerTargetCreator and indelRealigner in the GATK

software. The SNPs and InDels were called at the population level using the UnifiedGenotyper

in GATK and the -glm BOTH option. We extracted bi-allelic sites in all the accessions from the

variants using VCFtools version 0.1.13 (Danecek et al., 2011). Then, imputations were imputed

using Beagle version 4.1 (Browning and Browning, 2016). Finally, we analyzed the SNPs with

minor allele frequencies (MAFs) $\geq$ 0.05 in each population. In the analysis, the genotypes were

represented as -1 (homozygous of the reference allele), 1 (homozygous of the alternative allele)

or 0 (heterozygous of the reference and alternative alleles). Out of all the whole-genome

sequence polymorphisms, only the SNPs on chromosome 1 were analyzed. The number of SNPs

on chromosome 1 in each population is shown in Table 1.

Table 1. Number of SNPs and the diversity level of non-mixture and mixture populations.

	Population name	Number of accessions	Number of SNPs	Diversity level†
A.	Japonica	112	72,110	263.095
B.	Indica	100	427,943	660.416
C.	Temperate japonica	100	135,665	362.649
D.	Diverse	100	647,731	798.646
E.	A + B	212	633,507	803.064
F.	A + C	212	151,675	334.606
G.	A + D	212	684,774	859.678

172   † Diversity level is the index that was used to indicate the degree of genetic diversity and is

173   described in the "Degree of genetic diversity index" section below.

Generating phenotype data

Phenotypic data were simulated using the following formula:

$$\mathbf{y} = \mathbf{X_1}\boldsymbol{\beta_1} + \mathbf{X_2}\boldsymbol{\beta_2} + \mathbf{X_3}\boldsymbol{\beta_3} + \mathbf{u} + \mathbf{e}, \qquad \#(\mathbf{Eq.\ 1})$$

where $\mathbf{y}$ is the vector that represents the simulated phenotypic values for all 411 accessions; $\mathbf{X}$ is the design matrix representing the genotypes of three quantitative trait nucleotides (QTNs) with scores -1, 0, or 1; $\boldsymbol{\beta} = [\beta_1 \quad \beta_2 \quad \beta_3]^T$ is the vector representing the effects of the three QTNs, $\mathbf{u}$ is the vector for polygenetic effects, and $\mathbf{e}$ is the residuals vector. Three QTN-SNPs whose MAF was equal to or larger than 0.05 in all 411 accessions (672,923 SNPs in total) were randomly selected from the SNPs on chromosome 1. The simulations were divided into five categories (low, lower-middle, middle, higher-middle, high) based on the fixation index (F_{ST}) between populations A and B for the first QTN (Fig. S1 in Supplemental File 2). We assumed that the first QTN had four times greater variance than the remaining two QTNs (referred to as "QTN1", "QTN2", and "QTN3" respectively). The remaining two QTNs were chosen randomly from SNPs where the F_{ST} between A and B were low (SNPs whose F_{ST} value was in the lower 20% category among the 672,923 SNPs). The F_{ST} for each marker was calculated according to Wright (1965) as follows:

$$F_{ST} = 1 - \frac{H_S}{H_T}, \qquad \#(\text{Eq. Error! Bookmark not defined.})$$

191   where H_S is the average of the expected heterozygosity based on the allele frequencies of

192   populations A and B, and H_T is the expected heterozygosity based on the average allele

193   frequency of populations A and B. H_S and H_T were calculated as follows:

$$H_S = \frac{N_A \cdot \{2p_A(1-p_A)\} + N_B \cdot \{2p_B(1-p_B)\}}{N_A + N_B}, \qquad \#(\text{Eq. 3})$$

$$H_T = 2\left(\frac{N_A p_A + N_B p_B}{N_A + N_B}\right)\left(1 - \frac{N_A p_A + N_B p_B}{N_A + N_B}\right), \qquad \#(\text{Eq. 4})$$

where p_A, p_B, N_A, and N_B are the allele frequencies and the sample sizes of populations A and B

respectively, and $N_A = 112$ and $N_B = 100$. The F_{ST} distribution between A and B is shown in

Fig. S1, which also shows the thresholds for the five F_{ST} categories.

The polygenetic effect in Eq. 5 was sampled from the multivariate normal distribution

whose variance-covariance matrix was proportional to the additive numerator relationship matrix

$\mathbf{A}$ and was normalized so that their variance was equal to that of the three QTN effects.

$$\mathbf{u} \sim \text{MVN}(0, \mathbf{G}), \qquad \#(\text{Eq. 5})$$

where $\mathbf{G} = \mathbf{A}\sigma_A^2$ is the genetic covariance matrix, and the additive genetic variance σ_A^2 was

automatically determined from the relationship with heritability. In this study, the additive

numerator relationship matrix $\mathbf{A}$ was estimated based on the marker genotype data for 402,509

SNPs, which consisted of the core SNPs (defined by the Rice SNP-Seek Database as the "404k

CoreSNP Dataset") in all 12 chromosomes (this marker genotype data was prepared separately

from the whole-genome sequence data), using the "A.mat" function in R package "rrBLUP"

version 4.5 (Endelman and Jannink, 2012; Endelman, 2011).

The residual **e** in Eq. 6 was sampled identically and independently from the normal distribution,

and was then normalized so that the narrow-sense heritability was equal to 0.6. Residual **e** was

calculated using the following formula:

$$\mathbf{e} \sim \mathrm{MVN}(0, \mathbf{I}\sigma_e^2), \qquad \#(\mathrm{Eq.}\ 6)$$

210   where **I** is an identity matrix, and the residual variance σ_e^2 was determined so that the heritability

211   was equal to 0.6.

214   **Genome-wide association study (GWAS) using simulated data**

215   We performed a GWAS on the seven non-mixture (A, B, C, D) and mixture populations

216   (E, F, and G) using the marker genotype data and the simulated phenotypic data. We fitted the

217   linear mixed model (Yu et al., 2006).

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{S}_i\alpha_i + \mathbf{Q}\mathbf{v} + \mathbf{Z}\mathbf{u} + \mathbf{e}, \qquad \#(\mathrm{Eq.}\ 7)$$

where **y** is the vector of phenotypic values, $\mathbf{X}\boldsymbol{\beta}$, $\mathbf{S}_i\alpha_i$, and $\mathbf{Q}\mathbf{v}$ are the fixed effects terms, $\mathbf{Z}\mathbf{u}$ is

the random effects term, and **e** is the residuals vector. $\boldsymbol{\beta}$ represents all of the fixed effects other

than $\mathbf{S}_i\boldsymbol{\alpha}_i$, and $\mathbf{Q}\mathbf{v}$, and X is the incidence design matrix corresponding to $\boldsymbol{\beta}$. In this study, $\mathbf{X}\boldsymbol{\beta}$

was an intercept. $\mathbf{S}_i\alpha_i$ is composed of $\mathbf{S}_i$, which is the i_{th} marker of the genotype data, and α_i,

which is the effect of that marker. $\mathbf{Q}\mathbf{v}$ is the term used to correct for the effect of population

structure, and in this study $\mathbf{Q}$ was the matrix of two eigenvectors corresponding to the upper two

eigenvalues of the additive numerator for relationship matrix **A**, Finally, **u** represents the

polygenetic effects, and **Z** is the incidence design matrix corresponding to **u**.

We used the EMMAX and P3D algorithms to reduce the computation time (Kennedy et al.,

1992; Kang et al., 2008; 2010; Zhang et al., 2010). The "GWAS" function in R package

"rrBLUP" version 4.5 (Endelman, 2011) was used to perform the GWAS described above.

231   **Evaluation of the simulation results**

The p-value (or $-\log_{10}(p)$) for each marker effect was estimated 100 times by the

GWAS in five patterns according to the size of the F_{ST} for the seven non-mixture/mixture

populations. In this study, the following summary statistics were mainly used to evaluate the

GWAS results.

In the 100 simulations, the QTNs were not always polymorphic in each population

(because the MAF of the whole population did not necessarily match the MAF of each individual

population). In such cases, the $-\log_{10}(p)$ value of a QTN that was not polymorphic within a

population could not be calculated. Therefore, when two SNPs were polymorphic within that

population and were adjacent to the QTN, then the statistic of the more significant SNP was used

as the QTN statistic. Since it was difficult to detect such QTNs using a GWAS, we calculated the

summary statistics by dividing two patterns depending on polymorphism patterns of QTN1, i.e.,

whether using all simulation results or using only results whose QTN1 was polymorphic in the

target population (referred to as "All" and "Polymorphic in the population", respectively).

Correct detection rate (CDR) and $-\log_{10}(p)$

The first summary statistic was whether the $-\log_{10}(p)$ rate for each QTN exceeded the

threshold in each GWAS (referred to as "CDR; correct detection rate"). We assumed that QTNs

would be successfully detected by the GWAS when the CDR was large. The $-\log_{10}(p)$ value

whose false discovery rate (FDR) was 0.05 was set as the threshold using the Benjamini-

Hochberg method (Benjamini and Hochberg, 1995; Storey and Tibshirani, 2003). As the second

summary statistic, we used the $-\log_{10}(p)$ for each QTN in each GWAS, and we also assumed

that QTNs were successfully detected by the GWAS when this statistic was large.

Area under the curve (AUC)

We also regarded the mean of the AUC as one summary statistic. The AUC refers to the

area under the receiver operating characteristic (ROC) curve (Fig. S2 in Supplemental File 3),

which was obtained by plotting the false positive rate on the horizontal axis and the true positive

rate on the vertical axis when the threshold was varied (Hanley and McNeil, 1982). The AUC

was calculated using the following formula:

$$\text{AUC} = \frac{1}{2}FPRs_1 TPRs_1 + \frac{1}{2}\sum_{i=2}^{m+1}(FPRs_i - FPRs_{i-1})(TPRs_i + TPRs_{i-1}), \#(\text{Eq.}8)$$

261   where m is the number of QTNs, and $m = 3$ in this study. The $FPRs$ and $TPRs$ are the $m+1$

262   vectors whose i_{th} elements are $FPRs_i$ and $TPRs_i$, respectively. $FPRs_i = TPRs_i = 1$ when

263     $i = m + 1$. When $1 \leqq i \leqq m$, the $FPRs_i$ and $TPRs_i$ represent the false positive rate and the true

264   positive rate at the time when i QTNs exceed the threshold, respectively. They were calculated

265   using the following formula:

$$FPRs_i = \frac{FP_i}{FP_i + TN_i} \qquad (1 \leqq i \leqq m), \#(\text{Eq}.\,9)$$

$$TPRs_i = \frac{TP_i}{TP_i + FN_i} \qquad (1 \leqq i \leqq m), \#(\text{Eq}.\,10)$$

where TP_i, FP_i, FN_i, and TN_i are the numbers of SNPs that are the true positives (where the SNP

is a QTN and exceeds the threshold), the false positives (where the SNP is not a QTN but

exceeds the threshold), the false negatives (where the SNP is a QTN but does not exceed the

threshold), and the true negatives (where the SNP is not a QTN and does not exceed the

threshold) at the time when i QTNs exceed the threshold respectively. When we evaluated the

true/false positive rate, we considered the existence of linkage disequilibrium (LD) by

investigating SNPs with LD as one set. In this study, we defined SNPs that satisfied the

conditions that they were within 300 kb from the focused SNP and the condition that their

squares of the correlation coefficients with the focused SNP were 0.35 or more as one set when

considering LD. When we counted TP_i, FP_i, FN_i, and TN_i, we counted the number of the sets

described above instead of the number of SNPs. The value for AUC calculated in this manner

takes a value between 0 and 1. The GWAS is more successful when the AUC is closer to 1.

Using the mean of the AUC as one of the summary statistics meant that it was possible to focus

on each QTN and evaluate the overall results of the GWAS.

Precision, recall, and F-measure

We calculated the mean of precision, the mean of recall, and the mean of the F-measure

as other summary statistics to evaluate the GWAS results. These summary statistics can be

calculated from the numbers of true positives, false positives, false negatives, and true negatives.

More specifically, the precision can be calculated using the following formula:

$$\text{Precision} = \frac{TP}{TP + FP}. \qquad \#(\text{Eq. 11})$$

287   We regarded an SNP as "positive" when the $-\log_{10}(p)$ of that SNP exceeded the

288   threshold described above. The precision represents the ratio of the detected SNPs that were

289   QTNs. The recall was defined using the following formula:

$$\text{Recall} = \frac{TP}{TP + FN}. \qquad \#(\text{Eq. 12})$$

290   The recall represents the proportion of QTNs detected by the GWAS. Finally, the *F*-

291   measure was calculated as the harmonic mean of the precision and the recall, and can be used to

292   comprehensively evaluate the GWAS results. The *F*-measure was calculated using the following

293   formula:

$$F = \frac{2 \cdot \text{Precision} \cdot \text{Recall}}{\text{Precision} + \text{Recall}}. \qquad \#(\text{Eq. 13})$$

The greater these summary statistics, the more accurate the GWAS results were.

Degree of genetic diversity index

In order to evaluate the relationship between genetic diversity and the CDR results, we

prepared an index that indicated the degree of genetic diversity in each population. The

Euclidean distance matrix between accessions for each population was calculated. The median

for the off-diagonal elements of the distance matrix was used to indicate the degree of genetic

diversity (referred to as the "diversity level", Table 1). The median was chosen as the diversity

level because the distribution of the distances between the accessions for E and G had a double

peak. This was because, for mixture populations such as E and G, the distance within the

subpopulations was small whereas the distance between subpopulations was large. Therefore, if

the mean of the distances (almost the same as Nei's gene diversity index (NEI, 1973)) is chosen

as the diversity level, then there is a risk of overestimating the diversity level.

311   **RESULTS**

312   **Comparisons between the CDR and AUC for the QTN1s in each population**

313   The CDRs of the QTN1s in each population were calculated under ten conditions: five levels

314 of F_{ST} between A and B and two patterns of QTN polymorphism, i.e., whether the QTN was

315 polymorphic or not in the target population (Fig. 2 and Table S2 in Supplemental File 5).

Fig. 2. **Correct detection rate for QTN1 in each population under ten conditions.**

The barplots of CDR of QTN1 in each population under ten conditions: five levels of F_{ST} of QTN1 and

two patterns of polymorphisms of QTN. Blue horizontal dashed lines indicate the CDR in the population

A for each population. A: *japonica*, B: *indica*, C: temperate *japonica*, D: diverse, E: A+ B, F: A + C, G:

A + D.

For almost all levels of F_{ST}, the CDRs for QTN1 in the mixture populations E, F, and G

were larger than in the corresponding non-mixture populations B, C, and D, regardless of the two

QTN polymorphism patterns (Fig. 2). The CDRs for QTN1 in the mixture populations E, F, and

G were always larger than in population A when all the simulation results were taken into

account (Fig. 2). When F_{ST} was low, and all simulation results were taken into account (Fig. 2a),

populations G and D, which were highly diverse populations, had a higher CDR than the other

populations. When F_{ST} was in the lower-middle or middle category, and all simulation results

were taken into account (Figs. 2b, c), population E had the highest CDR. The CDR of the highly

diverse populations G and D significantly decreased as F_{ST} increased. This result suggested that

the QTN1 effect could confound with the population structure at higher F_{ST} values, which meant

that it was difficult to detect QTN1 in a highly diverse population. When the F_{ST} value was in the

higher-middle or high level categories, and all simulation results were taken into account, (Figs.

2d, e), the CDR for QTN1 became quite low in all populations. In populations D, E, and G,

QTN1 was hardly detected because of the strong confounding effect of the population structure.

In the other populations, the expected heterozygosity (H_e) for QTN1 was extremely small (In A

and B, H_e was less than 0.1 in all 100 simulations). The small H_e may make the detection of

QTN1 difficult.

We excluded the simulations in which there were no polymorphisms in the population

so that the detection power of the GWAS when there were polymorphisms in an analyzed

population could be evaluated (Figs. 2f-j). When F_{ST} was low, population F had the highest CDR

and when F_{ST} was in the lower-middle or middle categories, population A had the highest CDR.

However, there were only 14 and 9 cases in which QTN1 was polymorphic in population A. In

general, the populations with low or moderate genetic diversities (A, C, and F) had higher CDRs

than the populations with high genetic diversities (D, E, and G). When F_{ST} was in the higher-

middle or high categories, the results were similar to when F_{ST} was in the lower-middle or

middle categories.

The CDRs of QTN2 and QTN3 were much lower than that of QTN1 because smaller

genetic variances were assigned to these QTLs than QTN1 (Table S2). As in the case of QTN1,

for almost all levels of F_{ST}, the CDRs of QTN2 and QTN3 were higher in the mixture

populations (E, F, and G) than their corresponding non-mixture populations (B, C, and D).

Furthermore, the CDRs for QTN2 and QTN3 in all the mixture populations were higher than for

population A. The CDRs for QTN2 and QTN3 were also larger when the F_{ST} for QTN1 was

higher.

Populations D and G had high AUC values in all cases (Table S2). Population F had a

smaller AUC than populations D and G, even when the CDR was highest in population F.

Comparisons of the $-\log_{10}(p)$ values for the GWAS on each mixture population

containing *japonica* (A)

We compared the $-\log_{10}(p)$ values for each QTN between populations mixed with the

japonica population (A) to see if QTN1 was polymorphic in A (Fig. 3). Comparing these values

allowed us to examine whether the detection power of the GWAS improved when genetic

resources with higher genetic diversities were added to target population A. There is no plot for

the high F_{ST} values because no QTN1 was polymorphic in population A over 100 simulations

when the F_{ST} of QTN1 was high.

Fig. 3. **Boxplots of $-\log_{10}(p)$ of each QTN when QTN1 was polymorphic in *japonica* (A).**

Boxplots of $-\log_{10}(p)$ of each QTN for each mixture population and *japonica* (A) when QTN1

was polymorphic in A. These plots are shown divided into four categories according to the F_{ST}

value for QTN1 (a: low, b: lower-middle, c: middle, d: higher-middle).

For all of the four F_{ST} levels, the detection power improved in all mixture populations

compared to A (Fig. 3). Population F showed the highest detectability, and this tendency was

conspicuous even when F_{ST} was in the middle or higher-middle categories (Figs. 3c, d,

respectively). This is because the QTN1 effect is less likely to be confounded with the population

structure in F than in the other mixture populations (E and G). Population G had the highest

$-\log_{10}(p)$ values for QTN2 and QTN3, although only slightly (Fig. 3).

384   **Factors affecting the detection power of QTNs in the mixture populations**

385   We considered the factors related to the detection power of QTNs in the mixture populations

386   by creating a figure that represented the relationship between F_{ST}, the expected heterozygosity

387   (H_e), and the QTN1 detection power (Fig. 4 and Fig. S4 in Supplemental File 6).

Fig. 4. **Relationship between F_{ST}, H_e, and the detection power of QTN1.**

391 The distribution of each marker is plotted thinly with between subpopulation F_{ST} on the

392 horizontal axis and H_e of each subpopulation on the vertical axis. The dark X marks on the plot

393 show the SNPs selected as QTN1s in this study. Red and purple marks were detected by GWAS,

394 and green and yellow ones were not detected by GWAS.

396    Detection of the QTNs by the GWAS was generally difficult when the between-

397 subpopulation F_{ST} value was high, or H_e was low (Fig. 4). There seemed to be a significant

398 difference between plots F and E or G (Fig. 4a, Fig. S4, and Fig. 4b). However, in population F,

399 because A and C are genetically close, the F_{ST} between the subpopulations was not high.

400 Therefore, the relationships between F_{ST}, H_e, and the GWAS detection power applied to all

401 mixture populations.

402  Some of the QTNs were detected by the GWAS when F_{ST} was in the medium category, and

403 H_e in one of the subpopulations was close to 0 (Fig. 4a and Fig. S4). This suggested that even if

404 the QTN was fixed in one subpopulation, the QTN may still be detected by the GWAS if another

405 subpopulation was added to the analysis.

410 **Comparisons among the precision, recall, and *F*-measure values for each population**

411  The three summary statistics (the mean of precision, the mean of recall, and the mean of the

412 *F*-measure) were also calculated under ten conditions (Fig. S5 in Supplemental File 7). The

413 precision of the mixture populations was better than the precision value for population A for

414 almost all F_{ST} categories when all simulation results were taken into account (Fig. S5a).

415 However, it is not necessarily true that the precision of the mixture populations outperformed

416 that of their original genetic resources (compare E with B, F with C, and G with D). The recall

417 values of the mixture populations were larger than for their original genetic resources under all

conditions. Finally, a comparison of the F-measure for each population showed that there

seemed to be no tendency associated with F_{ST}. Therefore, it was difficult to conclude which

population was suitable for a GWAS when the F-measure is used. These results indicated that

using mixture populations for a GWAS led to the detection of more SNPs, including QTNs.

Relationship between the CDR results and genetic diversity

The relationship between the CDR results for QTN1 and the degree of genetic diversity was

evaluated under the two QTN polymorphism patterns, i.e., whether or not QTN was polymorphic

in the population (Fig. S6 in Supplemental File 8). The CDRs for the mixture populations were

usually larger than for the non-mixture populations if their diversity levels were close (Fig. S6a,

b). A comparison of the results for the different F_{ST} categories showed that when F_{ST} was low,

the populations with the highest diversities, such as D or G, had the highest CDRs, and when F_{ST}

was in the lower-middle or middle categories, the populations with the second-highest

diversities, such as B or E, had the highest CDRs. Finally, when F_{ST} was in the higher-middle or

high categories, the populations with relatively low diversities, such as C or F, had the highest

CDRs (Fig. S6a). However, when the simulations in which there were no polymorphisms in the

population were excluded, the populations with relatively low diversities, such as A, C, or F, had

the highest CDRs in almost all the F_{ST} categories (Fig. S6b).

                                                DISCUSSION

                     Relationship between F_{ST} and QTN detection

One of the main results of this study was that the detection of QTNs was difficult in populations

with high genetic diversities, such as D, E, and G, when the F_{ST} for QTN1 between *japonica* (A)

and *indica* (B) was high. This was because the QTN effect confounds with the effect of

population structure in these populations. We also examined the reasons why the CDRs for

QTN2 and QTN3 were high when the QTN1 F_{ST} value was high.

In this study, phenotypic values were simulated using the following expression:

$$\mathbf{y} = \mathbf{X_1}\boldsymbol{\beta_1} + \mathbf{X_2}\boldsymbol{\beta_2} + \mathbf{X_3}\boldsymbol{\beta_3} + \mathbf{u} + \mathbf{e}, \qquad \#(\mathbf{Eq.\ 1})$$

where $\mathbf{u}$ is the polygenetic effect, and is the term that reflects differences between accessions and

thus differences between subpopulations. Therefore, if the degree of QTN1 genetic

differentiation between *japonica* (A) and *indica* (B) is high, it can be assumed that there is a high

correlation between $\mathbf{X_1}\beta_1$ and $\mathbf{u}$. In this study, we generated phenotypic values using a certain

variance ratio under the assumption that each term is independent. Therefore, if there is a

correlation between $\mathbf{X_1}\beta_1$ and $\mathbf{u}$, and the variance between these two terms is considered as one

unit, it can be assumed that the variance is smaller than the total value of the two variances under

the assumption of independence. Therefore, the variance of these two terms $(\mathbf{X_1}\beta_1 + \mathbf{u})$ in the

total phenotypic variance becomes smaller, whereas the variances caused by the terms $\mathbf{X_2}\beta_2$ and

$\mathbf{X_3}\beta_3$ become greater than those when it is assumed that each term is independent.

The GWAS model used in this study was

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{S}_i\alpha_i + \mathbf{Q}\mathbf{v} + \mathbf{Z}\mathbf{u} + \mathbf{e}, \#(\text{Eq.}\ 7)$$

where $\mathbf{Qv}$ is the term used to correct the effect of population structure, and Zu shows the

polygenetic effect. In this GWAS model, $\mathbf{S}_i\alpha_i$ and $\mathbf{Qv}$ or $\mathbf{Zu}$ have some correlation when

$\mathbf{S}_i = \mathbf{X}_1$. This correlation results in the underestimation of α_i by the terms originally used to

correct the effects of population structure or family relatedness, such as $\mathbf{Qv}$ and $\mathbf{Zu}$. Therefore,

QTN detection is quite difficult when a GWAS is performed on mixture populations. For QTN2

and QTN3, where $\mathbf{S}_i = \mathbf{X}_2$ or $\mathbf{S}_i = \mathbf{X}_3$, there is generally no correlation between $\mathbf{S}_i\alpha_i$ and $\mathbf{Qv}$ or

$\mathbf{Zu}$. Therefore, the detection of these QTNs is not related to these terms. Furthermore, the

variances represented by the terms $\mathbf{X}_2\beta_2$ and $\mathbf{X}_3\beta_3$ are considered to be higher when the QTN1

genetic differentiation is not high. Therefore, the CDRs of QTN2 and QTN3 were high when the

F_{ST} for QTN1 was high (Fig. 2 and Table S1). It has been suggested by Atwell et al. (2010) that a

bias may occur in the GWAS results when the QTN correlates with population structure or

family relatedness.

472   **Relationship between H_e and the QTN detection**

The detection of QTNs by a GWAS was difficult when the expected heterozygosity (H_e) in the

population was low. When H_e in the population was low, the MAF was low, and alleles and

mutations with low allele frequencies are known as "rare alleles" or "rare variants". In such

cases, the QTN effect when the H_e values are low may be concealed by the QTN effect when H_e

is not low or by the environmental effect because there are few accessions with one allele. For

this reason, it is generally challenging to detect QTNs in such cases, but a method to deal with

this problem has been developed (Wu et al., 2011).

One example of a "rare variant" that is common in plants is the haplotype condition. It has

been reported that haplotypes are difficult to detect using a GWAS (Stram, 2014). This is

because haplotypes are often "rare variants" and their H_e values in the population are low.

Another problem caused by "rare variants" is that the non-causal SNP whose LD is strong with

the "rare variant" may have a higher $-\log_{10}(p)$ value than the "rare variant". This occurrence,

known as "synthetic association", often happens when the minor allele frequency of the SNP is

higher than that of the "rare variant" (Dickson et al., 2010). These "synthetic associations" were

often detected in this simulation study.

490   **Summary and further discussion on each result**

Generally, the CDRs of the QTNs showed that the populations suitable for a GWAS were

different depending on whether all the QTNs were to be detected or only the polymorphic QTNs

in the target population. Specifically, if all QTNs are to be detected when the degree of genetic

differentiation between QTNs is low, then it is optimal to use a population with high genetic

diversity that has as many polymorphisms as possible. However, as the degree of genetic

differentiation becomes more extensive, a population with high genetic diversity is not suitable

for a GWAS because the QTN effect is more likely to confound with the population structure. In

contrast, a population with moderate genetic diversity, such as population F, was suitable for a

GWAS, regardless of the degree of genetic differentiation. This was partly because the QTN1

effect was less likely to confound with the population structure in F than in E or G, even when

F_{ST} was high. However, in either case, when the degree of genetic differentiation is extensive, it

is difficult to detect the QTNs in any population. Therefore a GWAS analysis is not suitable,

which means that another approach, such as biparental QTL mapping, must be used to identify

genes (Lander and Botstein, 1989).

Population F had a smaller AUC than populations D and G, even when the CDR for

population F was the highest. From its definition, AUC is more dependent on how low

$-\log_{10}(p)$ of the QTN with the lowest $-\log_{10}(p)$ value is than on how high the $-\log_{10}(p)$ of

the QTN with the highest $-\log_{10}(p)$ value is. Furthermore, in this study, the number of markers

for the GWAS differed (Table 1). When $-\log_{10}(p)$ values for the QTNs were similar among the

different populations, the larger number of markers meant that the true negative rate increased,

and the false positive rate decreased in a population, which resulted in an increase in the AUC of

a population with a larger number of markers, e.g. D and G.

A comparison of the mixture populations and *japonica* (A) using $-\log_{10}(p)$ showed

that when the QTNs are polymorphic in a target population with low genetic diversity, genetic

resources with higher genetic diversities should be added to the target population. However, in

order to avoid cases where the degree of genetic differentiation among the QTNs is extensive

between the target population and genetic resources, it is desirable to use populations that are

genetically close to the target population.

Finally, the results suggested that the F_{ST} differences between the subpopulations and

the expected heterozygosity (H_e) of each subpopulation greatly influenced QTN detection by the

GWAS in the mixture populations (Fig. 4 and Fig. S4). This result was in agreement with the

above finding that QTN detection using a GWAS was generally difficult when F_{ST} was high, or

H_e were low. However, these situations frequently happened when the F_{ST} between the

subpopulations was moderate. Therefore, even if a QTN is fixed in one subpopulation, it may be

possible to detect the QTN by adding another population to the analysis because when the H_e of

the QTN is low in one population and F_{ST} is moderate, it can be assumed that H_e is relatively

high in the other population. Therefore, the H_e of the mixture population as a whole becomes

larger and the detection of a QTN is possible unless the confounding of the effect of that QTN

with the population structure is extensive. Although this situation is not difficult to interpret, it is

extremely important that SNPs with high F_{ST} and low H_e values must exist in large numbers

among populations. After taking this fact into account, a GWAS with a mixture population can

be useful. Therefore, creating the proposed diagram shown in Fig. 4 and Fig. S4, will lead to a

quantitative understanding of what kind of SNPs can be detected by a GWAS in mixture

populations of interest.

537   **Relationships with using whole-genome sequences**

One of the major factors related to the QTN detection power was the fixation index F_{ST}

differences among subpopulations. When the F_{ST} difference between the *japonica* (A) and *indica*

(B) subpopulations was low, the CDR of the mixture populations was high. One example of such

markers is that mutations may have occurred at the same position in both populations after they

differentiated. Since such variants are relatively new variants, the LD relationship between these

variants and surrounding markers will be weak. Therefore, these variants cannot be detected

using marker genotype data with a small number of markers, such as an SNP array. However, the

use of whole-genome sequences will increase the marker density, which improves the possibility

of detecting such variants with a GWAS. In summary, using whole-genome sequences improves

the possibility of detecting QTNs with low F_{ST} values and the use of mixture populations should

further improve the QTN detection power. In this study, there were cases where SNPs in a low

LD region were selected as QTNs when F_{ST} was low.

553   **CONCLUSION**

In this study, we examined a way of selecting a population that was suitable for a GWAS by

conducting simulations using populations with various genetic backgrounds. We evaluated the

results of the simulations by dividing them into ten patterns according to two criteria: the degree

of genetic differentiation (F_{ST}) between two main subpopulations and QTN polymorphism in a

target population. When the QTNs are polymorphic in a target population, increasing the

population size by adding available genotypes to the target population improves the detection

power. We suggest that a population genetically similar to a target population is desirable. After

investigating F_{ST} and expected heterozygosity H_e as factors that may substantially influence the

detection power of a GWAS, the results showed that SNPs with high F_{ST} and low H_e values were

less likely to be detected by a GWAS that used mixture populations. These results indicated that

the detection power of a GWAS was improved by using mixture populations with different

genetic backgrounds. Furthermore, the use of publicly available whole-genome sequences meant

it was possible to increase the population size and to use polymorphic markers that were present

in high numbers, which should also improve the detection power of the GWAS.

ACKNOWLEDGMENTS

This study was supported by Grant-in-Aid for Scientific Research(A) (25252002), Grant-in-Aid for Scientific Research(B) (Grant number 15H04436), JST, PRESTO (Grant number JPMJPR15O6), the Cross-ministerial Strategic Innovation Promotion Program (SIP), the "Technologies for creating next-generation agriculture, forestry and fisheries" (funding agency: Bio-oriented Technology Research Advancement Institution, NARO), and JST CREST (Grant Number JPMJCR16O).

We are also grateful for the advice given by Dr. Ryokei Tanaka.

SUPPLEMENTAL MATERIAL

Supplemental File 1: **Table S1.** Information about the 299 rice accessions used in this study.

Supplemental File 2: **Fig. S1.** Histogram showing the F_{ST} differences between *japonica* (A) and *indica* (B).

Supplemental File 3: **Fig. S2.** Example of a ROC curve and the AUC.

Supplemental File 4: **Fig. S3.** Principal components analysis results for chromosome 1 and all the chromosomes.

Supplemental File 5: **Table S2.** Correct detection rate rates for all QTNs and the AUC in each population.

589   Supplemental File 6: **Fig. S4.** Relationship between F_{ST}, H_e, and the QTN1 detection power for the

590   population G.

591   Supplemental File 7: **Fig. S5.** Bar plots of the precision, the recall and the F-measure results.

592   Supplemental File 8: **Fig. S6.** Relationship between the diversity level and the CDR of QTN1.

593   Supplemental File 9: **Supplementary Note.** Additional information about the materials used in this study.

594   **OPTIONAL SECTIONS**

595   **Availability of data and material**

596   Whole genome sequencing data are available of 112 accessions of *Oryza sativa* subsp.

597   *japonica* in the DDBJ Sequence Read Archive (SRA106223, ERA358140, DRA000158,

598   DRA000307, DRA000897, DRA000927, DRA007273, DRA007256, and DRA008071). Whole

599   genome sequencing data for all the other accessions are available in the "Rice SNP-Seek

600   Database".

602   **Competing interests**

603   The authors declare that they have no competing interests.

605   **Author's contributions**

606   KH, HKK, and HI conceived and designed the study. KH and HI performed the

607   mathematical and statistical analysis. KH, HKK, MY, EK, SY and HN contributed to marker

608   genotyping. KH, HKK, and HI wrote the manuscript in consultation with MY, EK, SY, and HN.

609   All authors read and approved the final manuscript.

611   **REFERENCES**

Alexandrov, N., S. Tai, W. Wang, L. Mansueto, K. Palis, R.R. Fuentes, V.J. Ulat, et al. 2015.
SNP-Seek Database of SNPs Derived from 3000 Rice Genomes. Nucleic Acids Res.
43(D1):D1023–27. doi: 10.1093/nar/gku1039.

Atwell, S., Y.S. Huang, B.J. Vilhjálmsson, G. Willems, M. Horton, Y. Li, D. Meng, et al. 2010.
Genome-Wide Association Study of 107 Phenotypes in Arabidopsis Thaliana Inbred Lines.
Nature 465(7298):627–31. doi: 10.1038/nature08800.

Auwera, G.A. Van Der, M.O. Carneiro, C. Hartl, R. Poplin, A. Levy-moonshine, T. Jordan, K.
Shakir, et al. 2014. From FastQ Data to High Confidence Varant Calls: The Genonme
Analysis Toolkit Best Practices Pipeline. Curr Protoc Bioinformatics 11(10):1–33. doi:
10.1002/0471250953.bi1110s43.From.

Begum, H., J.E. Spindel, A. Lalusin, T. Borromeo, G. Gregorio, J. Hernandez, P. Virk, B.
Collard, and S.R. McCouch. 2015. Genome-Wide Association Mapping for Yield and Other
Agronomic Traits in an Elite Breeding Population of Tropical Rice (Oryza Sativa). PLoS
One 10(3):1–19. doi: 10.1371/journal.pone.0119873.

Benjamini, Y., and Y. Hochberg. 1995. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. J. R. Stat. Soc. 57(1):289–300. doi:
10.2307/2346101.

Bolger, A.M., M. Lohse, and B. Usadel. 2014. Trimmomatic: A Flexible Trimmer for Illumina
Sequence Data 30(15):2114–20. doi: 10.1093/bioinformatics/btu170.

Browning, B.L., and S.R. Browning. 2016. Genotype Imputation with Millions of Reference
Samples. Am. J. Hum. Genet. 98(1):116–26. doi: 10.1016/j.ajhg.2015.11.020.

Danecek, P., A. Auton, G. Abecasis, C.A. Albers, E. Banks, M.A. DePristo, R.E. Handsaker, et
al. 2011. The Variant Call Format and VCFtools 27(15):2156–58. doi:
10.1093/bioinformatics/btr330.

Dickson, S.P., K. Wang, I. Krantz, H. Hakonarson, and D.B. Goldstein. 2010. Rare Variants
Create Synthetic Genome-Wide Associations. PLoS Biol. 8(1):e1000294. doi:
10.1371/journal.pbio.1000294.

Endelman, J.B. 2011. Ridge Regression and Other Kernels for Genomic Selection with R
Package RrBLUP. Plant Genome J. 4(3):250. doi: 10.3835/plantgenome2011.08.0024.

Endelman, J.B., and J.L. Jannink. 2012. Shrinkage Estimation of the Realized Relationship
Matrix. G3 (Bethesda) 2(11):1405–13. doi: 10.1534/g3.112.004259.

Hanley, J.A., and B.J. McNeil. 1982. The Meaning and Use of the Area under a Receiver Operating Characteristic (ROC) Curve. Radiology 142(1):29–36. doi: 10.1080/02634938208400381.

Huang, X., Y. Zhao, X. Wei, C. Li, A. Wang, Q. Zhao, W. Li, et al. 2012. Genome-Wide Association Study of Flowering Time and Grain Yield Traits in a Worldwide Collection of Rice Germplasm. Nat. Genet. 44(1):32–39. doi: 10.1038/ng.1018.

Jarquin, D., H. Kajiya-Kanegae, C. Taishen, S. Yabe, R. Persa, J. Yu, H. Nakagawa, M. Yamazaki, and H. Iwata. 2019. Coupling Day Length Data and Genomic Prediction Tools for Predicting Time-Related Traits under Complex Scenarios, 703488.

Kang, H.M., J.H. Sul, S.K. Service, N.A. Zaitlen, S.Y. Kong, N.B. Freimer, C. Sabatti, and E. Eskin. 2010. Variance Component Model to Account for Sample Structure in Genome-Wide Association Studies. Nat. Genet. 42(4):348–54. doi: 10.1038/ng.548.

Kang, H.M., N.A. Zaitlen, C.M. Wade, A. Kirby, D. Heckerman, M.J. Daly, and E. Eskin. 2008. Efficient Control of Population Structure in Model Organism Association Mapping. Genetics 178(3):1709–23. doi: 10.1534/genetics.107.080101.

Kawahara, Y., M. de la Bastide, J.P. Hamilton, H. Kanamori, W.R. McCombie, S. Ouyang, D.C. Schwartz, et al. 2013. Improvement of the Oryza Sativa Nipponbare Reference Genome Using next Generation Sequence and Optical Map Data 6(1):1–10. doi: 10.1186/1939-8433-6-1.

Kennedy, B.W., M. Quinton, and J.A. van Arendonk. 1992. Estimation of Effects of Single Genes on Quantitative Traits. J. Anim. Sci. 70(7):2000–2012. doi: 10.1016/j.alcohol.2011.07.002.

Koboldt, D.C., K.M. Steinberg, D.E. Larson, R.K. Wilson, and E.R. Mardis. 2013. The Next-Generation Sequencing Revolution and Its Impact on Genomics. Cell 155(1):27–38. doi: 10.1016/j.cell.2013.09.006.

Korte, A., and A. Farlow. 2013. The Advantages and Limitations of Trait Analysis with GWAS: A Review. Plant Methods 9(1):29. doi: 10.1186/1746-4811-9-29.

Lander, E.S., and D. Botstein. 1989. Mapping Mendelian Factors Underlying Quantitative Traits Using RFLP Linkage Maps. Genetics 121:185–99. http://www.ncbi.nlm.nih.gov/pubmed/2563713.

Li, H. 2012. Exploring Single-Sample Snp and Indel Calling with Whole-Genome de Novo Assembly 28(14):1838–44. doi: 10.1093/bioinformatics/bts280.

Li, J.Y., J. Wang, and R.S. Zeigler. 2014. The 3,000 Rice Genomes Project: New Opportunities and Challenges for Future Rice Research. Gigascience 3(1):1–3. doi: 10.1186/2047-217X-3-8.

Mansueto, L., R.R. Fuentes, F.N. Borja, J. Detras, J.M. Abrio-Santos, D. Chebotarov, M. Sanciangco, et al. 2017. Rice SNP-Seek Database Update: New SNPs, Indels, and Queries. Nucleic Acids Res. 45(D1):D1075–81. doi: 10.1093/nar/gkw1135.

Mansueto, L., R.R. Fuentes, D. Chebotarov, F.N. Borja, J. Detras, J.M. Abriol-Santos, K. Palis, et al. 2016. SNP-Seek II: A Resource for Allele Mining and Analysis of Big Genomic Data in Oryza Sativa. Curr. Plant Biol. 7–8:16–25. doi: 10.1016/j.cpb.2016.12.003.

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K. Garimella, et al. 2009. The Genome Analysis Toolkit: A MapReduce Framework for Analyzing next-Generation DNA Sequencing Data. Proc. Int. Conf. Intellect. Capital, Knowl. Manag. Organ. Learn. 20:254–60. doi: 10.1101/gr.107524.110.20.

Metzker, M.L. 2010. Sequencing Technologies the next Generation. Nat. Rev. Genet. 11(1):31–46. doi: 10.1038/nrg2626.

Misra, G., S. Badoni, R. Anacleto, A. Graner, N. Alexandrov, and N. Sreenivasulu. 2017. Whole Genome Sequencing-Based Association Study to Unravel Genetic Architecture of Cooked Grain Width and Length Traits in Rice. Nat. Sci. Reports 7(1):12478. doi: 10.1038/s41598-017-12778-6.

NEI, M. 1973. Analysis of Gene Diversity in Subdivided Populations. Proc. Nat. Acad. Sci. USA 70(12):3321–23. doi: 10.1016/j.jasrep.2018.01.028.

Ott, J., J. Wang, and S.M. Leal. 2015. Genetic Linkage Analysis in the Age of Whole-Genome Sequencing. Nat. Rev. Genet. 16(5):275–84. doi: 10.1038/nrg3908.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: Analyses of Phylogenetics and Evolution in R Language 20(2):289–90. doi: 10.1093/bioinformatics/btg412.

Saitou, N., and M. Nei. 1987. The Neighbor-Joining Method - a New Method for Reconstructing Phylogenetic Trees. Mol. Biol. Evol. 4(4):406–25.

Storey, J.D., and R. Tibshirani. 2003. Statistical Significance for Genomewide Studies. Proc. Natl. Acad. Sci. 100(16):9440–45. doi: 10.1073/pnas.1530509100.

Stram, D. 2014. Design, Analysis, and Interpretation of Genome-Wide Association Scans. Heidelberg, New York: Springer Science+Business Media.

Wang, W., R. Mauleon, Z. Hu, D. Chebotarov, S. Tai, Z. Wu, M. Li, et al. 2018. Genomic Variation in 3,010 Diverse Accessions of Asian Cultivated Rice. Nature 557(7703):43–49. doi: 10.1038/s41586-018-0063-9.

Wright, S. 1965. THE INTERPRETATION OF POPULATION STRUCTURE BY F☐STATISTICS WITH SPECIAL REGARD TO SYSTEMS OF MATING. Evolution (N. Y). 19:395–420.

Wu, M.C., S. Lee, T. Cai, Y. Li, M. Boehnke, and X. Lin. 2011. Rare-Variant Association Testing for Sequencing Data with the Sequence Kernel Association Test. Am. J. Hum. Genet. 89(1):82–93. doi: 10.1016/j.ajhg.2011.05.029.

Yabe, S., M. Yamasaki, K. Ebana, T. Hayashi, and H. Iwata. 2016. Island-Model Genomic Selection for Long-Term Genetic Improvement of Autogamous Crops. PLoS One 11(4):1–21. doi: 10.1371/journal.pone.0153945.

Yano, K., E. Yamamoto, K. Aya, H. Takeuchi, P.C. Lo, L. Hu, M. Yamasaki, et al. 2016. Genome-Wide Association Study Using Whole-Genome Sequencing Rapidly Identifies New Genes Influencing Agronomic Traits in Rice. Nat. Genet. 48(8):927–34. doi: 10.1038/ng.3596.

Yu, J., G. Pressoir, W.H. Briggs, I. Vroh Bi, M. Yamasaki, J.F. Doebley, M.D. McMullen, et al.
2006. A Unified Mixed-Model Method for Association Mapping That Accounts for
Multiple Levels of Relatedness. Nat. Genet. 38(2):203–8. doi: 10.1038/ng1702.

Zhang, Z., E. Ersoz, C.Q. Lai, R.J. Todhunter, H.K. Tiwari, M.A. Gore, P.J. Bradbury, et al.
2010. Mixed Linear Model Approach Adapted for Genome-Wide Association Studies. Nat.
Genet. 42(4):355–60. doi: 10.1038/ng.546.

Zhao, K., C.-W. Tung, G.C. Eizenga, M.H. Wright, M.L. Ali, A.H. Price, G.J. Norton, et al.
2011. Genome-Wide Association Mapping Reveals a Rich Genetic Architecture of
Complex Traits in Oryza Sativa. Nat. Commun. 2:467. doi: 10.1038/ncomms1467.

FIGURES AND TABLES

Fig. 1. **Unrooted phylogenetic tree plot for four non-mixture populations.**

Unrooted phylogenetic tree plot for the four non-mixture populations, which consisted of 112

accessions of *japonica* (A), 100 accessions of *indica* (B), 100 accessions of temperate *japonica*

(C), and 100 diverse accessions (D) with neighbor-joining method.

Fig. 2. **Correct detection rate for QTN1 in each population under ten conditions.**

The barplots of CDR of QTN1 in each population under ten conditions: five levels of F_{ST} of QTN1 and

two patterns of polymorphisms of QTN. Blue horizontal dashed lines indicate the CDR in the population

A for each population. A: *japonica*, B: *indica*, C: temperate *japonica*, D: diverse, E: A+ B, F: A + C, G:

A + D.

Fig. 3. **Boxplots of $-\log_{10}(p)$ of each QTN when QTN1 was polymorphic in *japonica* (A).**

Boxplots of $-\log_{10}(p)$ of each QTN for each mixture population and *japonica* (A) when QTN1

was polymorphic in A. These plots are shown divided into four categories according to the F_{ST}

value for QTN1 (a: low, b: lower-middle, c: middle, d: higher-middle).

Fig. 4. **Relationship between F_{ST}, H_e, and the detection power of QTN1.**

The distribution of each marker is plotted thinly with between subpopulation F_{ST} on the

horizontal axis and H_e of each subpopulation on the vertical axis. The dark X marks on the plot

show the SNPs selected as QTN1s in this study. Red and purple marks were detected by GWAS,

and green and yellow ones were not detected by GWAS.

Table 1. Number of SNPs and the diversity level of non-mixture and mixture populations.

	Population name	Number of accessions	Number of SNPs	Diversity level†
A.	Japonica	112	72,110	263.095
B.	Indica	100	427,943	660.416
C.	Temperate japonica	100	135,665	362.649
D.	Diverse	100	647,731	798.646
E.	A + B	212	633,507	803.064
F.	A + C	212	151,675	334.606
G.	A + D	212	684,774	859.678

756　† Diversity level is the index to indicate the degree of genetic diversity, which is described in the

757　Materials and Method section.